Fine Motor Serious Game Training Improves Gait in Parkinson's Disease: A Pilot Study

Valentin Bégel, Ph.D. Institut des Sciences du Sport Santé de Paris, Paris Cité University, Paris, France; International Laboratory for Brain, Music and Sound Research (BRAMS), Montreal, Canada

Frédéric Puyjarinet, Ph.D. Montpellier Psychomotor Training Institute, UFR de Medicine Montpellier-Nîmes, University of Montpellier, Montpellier, France

Christian Gény, M.D., Ph.D. Department of Geriatrics, CHRU of Montpellier, Montpellier, France

Valérie Cochen De Cock, M.D., PhD. Department of Neurology, Beau Soleil Clinic, Montpellier, France; EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Ales, Montpellier, France

Serge Pinto, Ph.D. CNRS, Aix-Marseille Univ, LPL, Aix-en-Provence, France

Simone Dalla Bella, Ph.D, International Laboratory for Brain, Music and Sound Research (BRAMS), Montreal, Canada; Department of Psychology, University of Montreal, Montreal, Canada; Centre for Research on Brain, Language and Music (CRBLM), Montreal, Canada; University of Economics and Human Sciences in Warsaw, Warsaw, Poland

Word count: 4850

Number of figures: 3

Number of tables: 4

* Corresponding authors:

Valentin Bégel, Paris Cité University, valentin.begel@u-paris.fr, 1 rue Lacretelle, 75015 Paris,

France. https://orcid.org/0000-0003-4049-1126

Simone Dalla Bella, University of Montreal, simone.dalla.bella@umontreal.ca, Pavillon Marie-

Victorin, CP 6128 Succursale Centre-Ville, Montréal, QC, H3C 3J7, Canada

Abstract:

Crucial functions for human behavior such as gross motor skills (e.g., walking), cognitive,

rhythmic, and fine motor processes are mostly considered unrelated. However, these functions

may interact, but their relations are still poorly understood. Moreover, evidence of causal links

between them is scarce. Neurological disorders such as Parkinson's Disease affect all these

functions and thus provide a model to study the interplay between them. We tested the effect of a

training delivered using serious games on tablet, involving upper limb rhythmic and fine motor

functions, on walking capacities in patients with Parkinson's Disease (PwPD). PwPD gathered

into an Intervention group played either a rhythm game, or an adaptation of *Tetris*, four times a

week for six weeks. Before and after the training, gait was evaluated in spontaneous walking and

a dual task (counting backward while walking). A Control group of participants did not receive

any training. Gait speed, stride length and cadence improved in the Intervention group after the

training in comparison with the Control group. The improvement was observed in both

Intervention groups, in the spontaneous and in the dual-task conditions. These findings support

the hypothesis that gross (axial) motor functions can be trained by fine (lateralized) motor

training administered via serious games, possibly by stimulating the rhythmic, perceptual, and

cognitive resources sensory systems. This opening promising perspectives for telerehabilitation.

Patients with movement disorders could benefit from this promising low-cost and engaging

training method, fostering inclusivity and autonomy for people who have reduced access to the

clinic.

Keywords: Telerehabilitation, Serious Games, Motor functions, Rhythm, Gait, Parkinson's

Disease

Introduction

Theories such as embodied cognition[1,2] and active sensing,[3] as well empirical behavioral and neurofunctional evidence, emphasize the interplay between the cognitive, rhythmic, and fine motor domains and gross motor functions.[4–6] The interrelationship between fine and gross motor domains has been an object of study for decades.[7] Although fine and gross motor domains are theoretically and functionally separated, links between them are consistently reported.[8–10] For example, there is evidence that training gross motor functions (e.g., run, hop, jump) can improve fine motor functions (strike, dribble, kick, catch, throw) in children.[11] However, the opposite causal relationship, namely the effect of fine motor functions training on gross motor functions, remains largely unexplored.

Further evidence of the link between fine and gross motor domains comes from their relationship with cognitive abilities. For example, early locomotor experiences are essential for the development of cognitive skills, and gross and fine motor activities help foster language and cognitive development.[12,13] Cognition and gross motor functions (e.g., gait) are linked in aging, as cognitive skills such as executive function and attention play a pivotal role in motor control.[4,14]

An effective means to address the causal relationship between related cognitive or motor domains is to test patient populations who display deficits in these functions. Parkinson's

Disease (PD) is an interesting model to study the interaction between fine and gross motor domains. People with PD (PwPD) experience fine and gross motor difficulties both in axial (gait, speech, posture) and lateralized (limbs) movements due to the cardinal symptoms associated with the disease (bradykinesia, rigidity, tremor, and postural instability).[15] Fine motor functions are more impaired in PwPD with mild cognitive impairment, compared with patients without cognitive impairment.[16] Training gross motor functions via physical exercise improves cognition as well as gait in PD.[17] This suggests that fine and gross motor domains are related to cognition in PD, and transfer from one domain to another might be observed.

A key aspect of fine and gross motor functions is that sequences of motor events are organized in time. Specifically, activities such as walking or speech articulation, both typically altered in PwPD, rely on rhythmic sequences of movements which are precisely timed.[6,18] The capacities to produce motor rhythms are often assessed with tasks relying on fine motor skills, in tasks such as finger tapping.[19–21] PwPD typically display a more variable motor output in finger tapping tasks.[22,23] Increased variability in PD is also visible in rhythmic tasks involving gross motor functions (e.g., walking,[5]) or in speech.[6,24] Cognition and rhythmic skills are also related, as perceptual and rhythm processing correlate positively with executive functions (e.g., flexibility, inhibitory control), memory, and attention in various population, including PwPD.[25–27]

There is evidence that training gross motor functions can lead to transfer effects in the cognitive, fine and gross motor, and rhythmic domains. Training locomotor functions can improve fine motor skills in children,[11] and cognition in PD.[17] Gait training using rhythmic auditory stimulation improved fine motor and perceptual rhythmic skills (finger tapping with a periodic sound, duration discrimination and detection of misaligned beats in musical excerpts) in PD.[23] However, there is a paucity of studies in which fine motor functions are trained and transfer effects in gross motor functions are tested in PD. To our knowledge, the only exception is a recent study from our group in which fine motor functions were trained with a finger tapping game in PD, and the transfer effects on gait, tapping, and oromotor activity.[28] The focus of this study was narrow, though, as it took motor variability across different motor domains (manual, oromotor, and gait) as the main outcome measure. While the study showed a cross-domain transfer, by demonstrating a reduction in motor variability in the oromotor rhythmic task after the finger tapping rhythmic training, no effect on stride time variability was observed. However, this study did not test other gait spatio-temporal features (speed, stride length, and cadence).

The goal of this pilot study was to assess the effect of serious video game training protocols (one rhythmic game, and one non-rhythmic game) implemented on tablet and requiring finger movements, to improve gross, axial motor functions, namely gait spatio-temporal features, in PwPD. Serious games are games with pleasant and recreational aspects designed for purposes

beyond entertainment,[29] and have been proven efficient to train both fine and gross motor

domains in PD.[28,30]

We used two games that involve similar finger movements. One of them, *Rhythm

Workers*, is a rhythmic game that we developed and validated in PwPD.[21,28,31,32] Notably,

this form of rhythmic training differs from standard rhythmic auditory cueing, in which the

stimulation is typically delivered while patients walk.[22,23] The other game was an adaptation

of *Tetris*, a game that focuses on visuospatial ability and cognitive skills such as decision-

making.[33,34] One group of PwPD played *Rhythm Workers* and another played *Tetris* for two

months. A third group of PwPD did not receive any training during these two months. Before

and after the training period, we tested participants' gross motor skills (i.e., gait), as well as

rhythm perception and cognitive abilities. Some of the participants and the training protocol were

the same as in our previously published study;[28] here we focused on the effect of the training

on gait spatio-temporal parameters. We predicted that a telerehabilitation training targeting fine

motor skills, displayed via serious games (*Rhythm Workers* and/or *Tetris*) would improve

participants' gross motor control involved in gait, compared to a control group who did not

receive any training. We also expected that adding a rhythmic component to the training would

produce greater transfer to gait performance than non-rhythmic training.

Methods

Participants

Participants were recruited by a clinical research assistant, supervised by the team's

neurologist, at the Montpellier University Hospital. We recruited 45 participants with PD. Other

outcome measures collected for 33 of these participants were included in a previous publication,

in which we focused on speech and motor variability.[28] In this study, we present different gait

measures for them and 12 additional participants were tested who received the same training. All

participants who were contacted accepted to participate in the study, but three of them withdrew

their participation during the experiment, resulting in a final sample of 42 participants (12

women, mean age = 67.55, SD = 7.35, age range = 50-82). Patients' demographic and clinical

characteristics are presented in table 1. Patients' medication was stable for at least 4 weeks prior

to the start of the experiment, and they were under optimal medication for the evaluation sessions

(60–90 min after their usual dose). Participants were diagnosed with PD in accordance with the

UK Brain Bank criteria[15] and disease duration ranged from 2 to 25 years prior to the

experiment (mean duration: 10.05 years, SD = 5.61 years).

Table 1. Patients' demographic and clinical characteristics			
	Rhythm Workers (n = 19, 3 women)	Tetris (n = 11, 2 women)	No intervention (n = 12, 6 women)
	Mean (range)	Mean (range)	Mean (range)
Demographics			
Age	65.7 (50-77)	67.7 (56-82)	70.3 (63-79)
Disease duration (Years)	10.2 (2-25)	9.6 (2-23)	10.3 (6-18)
Clinical characteristics (MDS-UPDRS)			
Total score	64.8 (32-102)	57.4 (24-88)	56.7 (34-79)
Motor subscore (part 3)	30 (3-57)	26.3 (10-39)	29.45 (17-46)
Speech item (3.1)	1.5 (0-2)	1 (0-2)	1.2 (0-3)

Upper limb items (3.4a–3.6b)	7.1 (1-16)	4.7 (0-10)	8.2 (1-14)
Lower limb items (3.7a–3.8b)	6.5 (0-13)	5.2 (2-9)	6.36 (0-12)
Hoehn and Yahr score	2.23 (2-3)	2.2 (2-3)	2.27 (2-3)

Sample Size

Following recommendations by Whitehead et al. (2015),[35] the sample size was

estimated without formal power calculation, as it is not required to fulfil the objectives of a

small-scale, pilot study. We included a number of participants equivalent to the ones of

intervention studies using the same primary outcomes (gait parameters), which corresponds to 10

to 15 participants per group.[22,23]

Training protocol

Following a parallel-group design, participants were randomly assigned by a clinical

research assistant (using the free online application Group Maker -

https://livecloud.online/en/group-maker) to the Intervention (n = 30) or the Control group (n =

12). In the intervention group, participants were randomly assigned to the *Rhythm Workers* (*RW*

Group, n = 19) or *Tetris* Intervention group (*Tetris* Group, n = 11). Patients in the Intervention

group were instructed to play games on a tablet device (LG© G Pad 8.0 model) for at least 30

min, 4 times a week, during a period of 6 weeks, for a total of 12 hours.

Rhythm Workers is a music-based rhythm game consisting in finger tapping to the beat of

auditory stimuli varying in terms of rhythmic complexity (metronome, metrical sequences, and

musical excerpts).[21,32] The goal of the game is to synchronize the finger taps to the beat of

rhythmic stimuli. The greater the synchronization (i.e., alignment of the taps with the beats), the

higher the score. Feedback is provided by the score (maximum score = 100), and a minimum

score (= 70) is required to progress through difficulty levels. *Tetris* is a classical video game in

which pieces with different shapes move from the top to the bottom of the computer screen. In

the adapted version we developed, the player uses the tactile screen to move and/or rotate the

pieces by finger tapping anywhere on the screen, so each piece can fit in with previous pieces to

form complete rows (no gaps between pieces). Completed rows disappear, and the remaining

pieces drop down. Each completed row earns points, and the level stops when too many pieces

filled the screen.

Performance measures

Performance measures were collected before and after the training at the Montpellier University

Hospital by a team of researchers and clinical research assistants from March 2016 to January

2022.

Gait

Gait performance was quantified while participants walked over an eight-meter

GAITRite© 301 instrumented walkway system (CIR Systems Inc, Havertown, Pennsylvania)

composed of pressure-sensitive sensors that activated on foot contact. The walkway was disposed in an isolated hallway free from distraction. Participants started walking two meters away from the walkway to avoid variability (accelerations and decelerations) at the onset of the gait trial. They had to continue walking when they reached the walkway. In a first condition (Spontaneous gait), participants were instructed to walk naturally on the walkway. In a Dual task condition, participants had to verbally count backward from 100, subtracting in increments of seven. The Dual task condition was included to modulate the resource demands attributed to gait, introducing a cognitive load to and stimulate the interplay between gross motor functions and cognition. Three trials were performed in each condition. Spatiotemporal gait parameters, averaged from the three trials, were calculated for both Spontaneous and Dual tasks. Those included Speed (cm/sec), Cadence (steps/minute) and Stride length (cm).

Rhythm perception: Beat Alignment Test (BAT)

To control for rhythm perception skills of PwPD, the Beat Alignment Test (BAT) from the BAASTA was administered.[36,37] Participants listened to 72 twenty-beat excerpts from Bach's Badinerie and Rossini's William Tell Overture presented at three different tempi (inter-beat intervals, IBIs, of 450, 600 and 750 ms). Starting from the 7th beat, an isochronous sequence with a triangle timbre was superimposed to the music, either aligned (24 trials) to the musical beat or non-aligned (48 trials). Period changes (10% slower or faster than the IBIs, 24 trials) and phase changes (the tones preceded or followed the beats by 33 % of the IBIs, 24 trials)

were presented. Participants were asked to judge whether the metronome was aligned or not with the musical beat. The sensitivity index (d'), corresponding to the standardized difference between the hits (i.e., when a misaligned metronome was correctly detected) and false alarms (i.e., when a misalignment was erroneously reported) was calculated as an unbiased measure of detection performance.

Neuropsychological assessment

The Montreal Cognitive Assessment[38] was administered to assess global cognitive abilities. Further evaluation of memory and executive functions was performed using the Stroop Color Word Test (interference - time in seconds and number of errors),[39] the Digit Span subtest from Wechsler Adult Intelligence Scale version III (Forward and backward digit span),[40] and the Trail Making Test (TMT, part B, time in seconds and number of errors).[41]

Statistical analyses

All analyses were run in R Statistical Software.[42] Analyses of variance (ANOVA) were run with Afex package (v1.0-1)[43] and Tukey-corrected linear contrasts were run with the Emmeans package (1.7-2)[44]. First, we assessed the effect of the training by comparing the Intervention and Control groups. Gait spatiotemporal parameters (speed, stride length and cadence), results at the BAT and neuropsychological tests were entered as dependent variables in

a 2-way ANOVA with Group (Intervention vs. Control) as the between-subject factor and

Session (Pre vs. Post training) as the within-subject factor.

Z-scores of patients in the Interventions group were calculated relative to the

performance of the Control group [Z-score $= (Value - Mean_{Controls})/SD_{Controls}$] to ensure

that the effect observed in the Intervention group was not merely a test-retest effect. Z-scores

were computed across the two types of interventions for gait variables and the BAT scores

separately for pre and post-training measures. To quantify the effect of the interventions relative

to the performance of the Control group, Delta Z-scores ($\Delta = Z\text{-}score_{Post\text{-}Training} -$

$Z\text{-}score_{Pre\text{-}Training)}$) were calculated for each patient. Positive Delta scores reflect an

improvement of the performance. Delta Z-scores for gait spatiotemporal parameters (Speed,

Stride length and Cadence) for both the *Rhythm Workers* and the *Tetris* groups were entered as

dependent variables in a 2-way ANOVA with Group (*Rhythm Workers* vs. *Tetris*) as the

between-subject factor and Condition (Spontaneous vs. Dual task) as the within-subjects factor.

General Eta Squared are reported as a measure of effect sizes (.01: Small effect size, .06:

Medium effect size,14 or higher: Large effect size). In addition, the spontaneous and Dual-task

conditions were averaged, and t-tests were used to test if the mean of the Delta Z-scores at the

gait parameters differed from zero (absence of an effect of the training) in each group.

Finally, we used the clinically meaningful criteria in PD defined by Hass and

collaborators to quantify the benefits of the training at the individual level.[22,45,46] Following

Hass and collaborators criteria, a difference of 0.06 m/s in gait speed was considered as small, a

difference of 0.14 m/s was moderate, and a difference of 0.22 m/s was considered as large. We

then divided participants into two categories based on their response to the training. Participants

who showed a clinically significant improvement in gait speed (at least 0.6m/s in either

Spontaneous or Dual task conditions) were considered as participants with a Positive Response

(PR). Participants who did not respond positively to the training were considered as participants

with a Non-Positive Response (NPR, for a similar classification of patients' response to a

rhythmic training see Dalla Bella et al., 2017[22]). The difference between the PR and NPR

groups in total time spent playing the game (in hours) was assessed using Wilcoxon rank-sum

test. We computed t-tests on the Delta Z-scores to compare the effect of the training on gait

Speed, Stride length and Cadence in the PR and NPR groups. For each variable, significance

values were corrected for multiple comparisons.

Results

Intervention vs. Control

Gait spatio-temporal parameters (speed, stride length, and cadence) at Pre-training and

Post-training evaluation sessions are shown in Figure 1. Notably, the Intervention and Control

groups did not differ at baseline (Speed, Spontaneous: $p = .39$, Dual task: $p = .69$; Stride length,

Spontaneous: $p = .52$, Dual task: $p = 62$; Cadence, Spontaneous: $p = .49$, Dual task: $p = 46$).

Overall, participants in the Intervention group increased their gait speed by 5.01cm/second

(4.67%) in the Spontaneous task and 10.14cm/second (13.34%) in the Dual task, stride length by

3.93cm (3.24%) in the Spontaneous task and 7.26cm (6.61%) in the Dual task, and cadence by

1.25 step/min (1.18%) in the Spontaneous task and 5.95 step/min (7.33%) in the Dual task from

pre to post training. Participants in the Control group showed no significant change.

For gait Speed, the two-way ANOVA pointed to a Group x Session interaction, at the

threshold of statistical significance, $F(1,40) = 3.90$, $p = 0.055$, $\eta_G^2 = 0.07$ (Figure 1A, left panel).

The difference between Pre and Post was significant in the Intervention group, $t(40) = 2.25$, $p =$

0.03, $d = 0.58$, but not in the Control group, $t(40) = 0.91$, $p = .37$, $d = 0.37$. The Group X

Session interaction attained significance in the Dual task condition, $F(1,37) = 6.25$, $p = 0.017$, η_G^2

$= 0.006$ (Figure 1A, right panel). The difference between Pre and Post was significant in the

Intervention group, $t(37) = 3.37$, $p = 0.002$, $d = 0.92$, but not in the Control group, $t(40) = 0.75$,

$p = 0.45$, $d = 0.31$.

For Stride length, there was a Group x Session significant interaction in the Spontaneous

condition, $F(1,40) = 5.22$, $p = 0.028$, $\eta_G^2 = 0.006$ (Figure 1B, left panel). The difference between

Pre and Post was significant in the Intervention group, $t(40) = 3.04$, $p = 0.004$, $d = 0.79$, but not

in the Control group, $t(40) = 0.77$, $p = 0.44$, $d = 0.32$. In the Dual task condition, there was an

interaction between Group and Session on Stride length, at the threshold of statistical

significance, $F(1,37) = 4.01$, $p = 0.053$, partial $\eta_G^2 = 0.008$ (Figure 1B, right panel). The

difference between Pre and Post was significant in the Intervention group, $t(37) = 3.28$, $p =$

0.002, $d = 0.89$, but not in the Control group, $t(37) = 0.22$, $p = 0.83$, $d = 0.90$.

No main effects or interaction were found for Cadence (Spontaneous and Dual task

conditions), rhythm perception (The BAT) and in the neuropsychological tests (all ps > .05, see

supplementary table).

Figure 1

Gait spatiotemporal parameters (A = Speed, B = Stride length, C = Cadence) in the

Spontaneous task and Dual task for the Interventions and the Control groups. Errors bars

represent Standard Errors of the mean.

Rhythm Workers vs. Tetris

The Deltas, measured as the difference between the post and the pre-training at the averaged Spontaneous and Dual tasks Z-scores (relative to the Control group for the *Rhythm Workers* and *Tetris* subgroups) are shown in Figure 2. As can be seen, the Z-scores are generally larger than zero, indicating an improvement of the performance in both groups.

The ANOVA on gait Speed revealed no significant main effect or interactions.

The ANOVA on Stride length revealed a significant main effect of Group, $F(1,25) =$ 6.05, $p = 0.021$, $\eta_G^2 = 0.160$, suggesting that in general, the effect was more important in the *Tetris* group. The effect was also more important in the Dual task condition (Main effect of Condition, $F(1,25) = 8.00$, $p < 0.01$, $\eta_G^2 = 0.009$). Finally, there was an interaction between Group and Condition factors, $F(1,25) = 5.63$, $p < 0.026$, $\eta_G^2 = 0.46$. In the *Tetris* group, Delta Z-scores were significantly larger in the Dual task ($M = 1.79$, $SD = 1.31$) than in the Spontaneous ($M = 0.76$, $SD = 0.59$), $t(25) = 3.27$, $p = 0.003$, $d = 1.47$, whereas they did not differ in the RW Group (Spontaneous, $M = 0.32$, $SD = 0.64$; Dual task, $M = 0.45$, $SD = 1.28$), $t(25) = 0.37$, $p = 0.71$, $d = 0.13$.

For Cadence, there was no significant main effect or interactions.

We assessed whether differences in gait improvement were linked with the amount of time participants played the game in the two groups. Participants in the *Tetris* group played significantly more time than in the RW group (RW, $M = 5h$, $SE = 0.9h$; *Tetris*, $M = 29h$, $SE = 0.9h$), $t(10.23) = 2.81$, $p < 0.018$, $d = -1.40$).

Figure 2

Gait spatiotemporal results (Z-scores) at the gait parameters for the Rhythm Workers and the

*Tetris groups. Zero on the Y axis means that there is no improvement. * p < .05, ** p < .01, ****

p < .005, one-sample t-test against zero (zero = no effect of the interventions), corrected for the

four comparisons in each panel using Bonferroni correction. Errors bars represent Standard

Errors of the mean.

Although there were no effects of the interventions on the results at the BAT and the

neuropsychological tests, we conducted the Z-score analyses for these variables to highlight the

different effects of each training on rhythmic and cognitive skills. Participants in the RW group

had larger Delta Z-scores than participants in the *Tetris* group at the BAT (RW, *M* = 0.30, *SE* =

0.19; *Tetris*, *M* = -0.35, *SE* = 0.14), $F(1,25) = 4.86$, $p = 0.037$, $\eta_G^2 = 0.163$. We ran correlations

between the Delta Z-scores in the BAT and the gait variables to explore the relationship between

the improvement in rhythmic skills and in gait separately for each group. None of the

correlations were significant (all *p*s > .05). The two groups did not differ in the

neuropsychological tests (all *p*s > .05).

Interindividual differences

Overall, 20 patients out of 31 (67%) showed a clinically meaningful improvement (PR) in

the Intervention group. Eight of them showed a small effect (5 in the RW group, 3 in the Tetris

group), six a moderate effect (2 in the RW group, 4 in the Tetris group) and three a large effect

(2 in the RW group, 1 in the Tetris group) in Gait speed. The gait performance of PR and NPR in

the Intervention group (RW and Tetris groups taken together) are presented in Figure 3. The

improvement of gait parameters was significantly more important in PR than in NPR for Speed

(PR, *M* = .60, *SE* = .15; NPR, *M* = -.02, *SE* = .11), $t(16.50) = 3.47$, p = .001, $d = 1.39$, Stride

length (PR, *M* = .94, *SE* = .19; NPR, *M* = .12, *SE* = .30), $t(16.56) = 2.31$, p = .017, $d = 0.92$, and

Cadence (PR, *M* = .43, *SE* = .09; NPR, *M* = -.24, *SE* = .22), $t(12.24) = 2.85$, p = .007, $d = 1.31$.

Across the two groups, PR played significantly more than NPR (PR, *M* = 17h, *SE* = 5.5h; NPR,

M = 7h, *SE* = 1.5h; $t(21.52) = -1.77$, *p* = .045, $d = 0.49$).

Figure 3

Gait spatiotemporal results (Z-scores for the gait parameters) for PR and NPR. Average values

*for the Spontaneous and Dual task conditions are presented. * p < .05, ** p < .01*

Discussion

The aim of this pilot study was to test the effect of playing mobile video games involving

fine motor movement (finger movement) on one gross motor skill (gait). We provided first

evidence that spatio-temporal parameters in gait can be trained in patients in PD patients playing

a video game on tablet (a rhythm game or an adaptation of *Tetris)* training fine movement

abilities for six weeks. After the training period, patients showed improved walking, as revealed

mostly by faster speed, and increased stride length. The effect was visible in spontaneous

walking (4.67% for speed and 3.24% for stride length) as well as in a dual-task condition

consisting in counting backwards while stepping (12.34% for speed and 6.61% for stride length).

In comparison, a control group of PD participants without training did not show any

improvement.

The observed transfer effect from a fine motor control training, like finger tapping, to gait

spatio-temporal parameters is novel. Participants who trained fine motor control - in a rhythmic

or non-rhythmic game - displayed improvements of spontaneous gait performance just slightly

lower than observed after standard rhythmic auditory cueing therapies, and superior with regard

to dual-task performance. For example, in a previous study, PD patients improved their speed by

6% and their stride length by 4.5% in spontaneous walking after they underwent a one-month

rhythmic auditory cueing program.[22] .This is consistent with the average improvement

observed in recent meta-analyses of the effect of auditory cueing in PwPD.[47,48] Moreover, if

we focus on the gait effects of deep brain stimulation protocols, the intervention typically does

not show a superior general improvement. Patients who respond positively to deep brain

stimulation in terms of gait speed (53%) show an improvement of 5.72% on average.[49] In sum,

comparisons with previous studies suggest that fine motor training improves gait parameters in

spontaneous walking to a level slightly inferior to the benefits of rhythmic auditory cueing. This

is not surprising, as rhythmic auditory cueing directly trains gait, thus within-effector benefits are

expected.[22,50–52] Nevertheless, the effect was still quite high in spontaneous walking and

even more in dual task, compared for example with deep brain stimulation, given that gait was

not trained. Notably, more than two thirds of patients (67%) responded positively to the training

by showing a clinically meaningful improvement measured by gait speed, with almost one third

(29%) exhibiting average to large improvements.[45] This finding is promising, considering that

usually around 50% of the patients typically respond positively to rhythmic auditory cueing (see

recent studies that used the same criteria[45] to quantify the clinical benefits of the training

protocol[22,46]), which is similar to what is observed after deep brain stimulation.[49]

Moreover, there are important individual differences in the response to the training,

which may be partly explained an effect of dose, namely the true amount of training received.

Participants who responded positively to the training played on average more (17h) than the

instructed time (12h). In contrast, the ones who did not respond to the training played

significantly less (7h). In a future clinical trial, efforts should be made to increase participants'

adherence to the training, by improving the game design and add features that are more engaging

for PwPD.

The beneficial effects of gamified training of fine motor movements were visible for the

rhythmic game and Tetris alike. However, the mechanisms underlying these improvements are

likely to be different, as the two games engage different perceptual and cognitive functions. For

example, it has been shown in previous studies that that rhythmic auditory cueing is linked to

rhythmic sensorimotor skills, making rhythmic processes that are independent from an effector a

plausible mechanism underlying the improvement of gait parameters.[22,23,46] In line with this

idea, we reported in a recent study that rhythmic training in PD involving finger tapping reduces

motor variability in gait.[28] Different mechanisms may underlie the positive effect of *Tetris.*

The absence of improvement at a perceptual rhythmic task (the BAT) after the training in the

Tetris group confirms that *Tetris* does not influence rhythmic skills. The effect of *Tetris* on gross

motor skills could be attributed to selective perceptual and cognitive improvements. Balance and

gait are complex, biomechanically constrained skills that require perceptual (sensory strategies,

orientation in space) and cognitive resources.[53] Previous studies have shown that although

Tetris does not improve general cognitive skills,[54] it enhances specific skills such as selective

visual attention[55] and spatial skills.[56] These skills are impaired in PD and contribute to gait

and posture disorders.[57,58] Recently, Wielenski and colleagues conducted a pilot study that

showed that training visual attention skills improves gait in PD studies.[59] Sufficient visual

attentional and spatial orientation resources are necessary for postural control, and more so when

the task is complex, such as in dual-tasks.[60] The beneficial effect of the *Tetris* training on gait

is stronger in the dual-task condition (Stride length), further suggesting that *Tetris* training is

tapping into visual attention and spatial orientation skills. Although it is possible that the positive

effect of *Tetris* and *Rhythm Workers* are based on different mechanisms, we cannot rule out the

involvement of a common mechanism, possibly based on reward and motivation induced by

playing. Music and video games activate dopaminergic pathways associated with reward

processing and motivation.[61,62] Dysfunctions in these dopaminergic pathways are one of the main physiopathological features of PD.[63] Playing video games may therefore improve the activity of the dopaminergic system impaired in PD.

In conclusion, this short-scale study supports the hypothesis that one gross motor skill (gait) can be improved by fine motor, finger-tapping, training administered via video games. Findings in PwPD revealed a possible causal relationship between fine and gross motor skills. A future clinical trial with more participants will be needed to confirm these results, with an improved game design to increase participants' adherence to the training. Future studies with more exhaustive testing of motor, perceptual and cognitive skills before and after the training' to further characterize the mechanisms that mediate the transfer effect from fine motor to gross motor skills. This will also serve to rule out a placebo effect, as the effect of the training in the intervention group was shown in comparison with a passive control group. Nevertheless, the fact that positive responders played more than non-positive responders suggests that the training *per se* was efficient.

The results of this study have implications for clinical research and healthcare. Most methods to train gross motor skills require costly equipment as well as the presence of a clinician, and thereby cannot be carried out at home. The telerehabilitation technique that we used is low-cost, user-friendly, and fosters inclusivity and autonomy, as it could provide people

who have reduced access to clinics or patients who are temporarily immobile with an efficient training method.

Statements and Declarations

Ethics approval and consent to participate

This study was approved by the French Comité de Protection des Personnes (CPP N_2015-A01090-49). All participants gave their informed consent in accordance with the Helsinki Declaration (World Medical Association, 2024). This study was registered as a clinical trial (ClinicalTrials.gov, registration NCT02855710).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing Interests and Funding

Simone Dalla Bella and Valérie Cochen De Cock are board members of the BeatHealth startup dedicated to the design and commercialization of rhythm-based testing devices (including BAASTA) and cognitive therapies such as Rhythm Workers. For the remaining authors, no competing interests are to be declared.

This research was funded by a CIFRE grant (R153304C) to VB, the Junior Grant from the Institut Universitaire de France (IUF) to SDB, and the Maturation Technological Transfer grant (YouRhythm project) from SATTAxLR to SDB and the company NaturalPad.

Author contributions

V.B., S.P., and S.D.B conceived and planned the study. V.B, F.P., and C.G. collected the data.

V.B. and S.D.B analyzed, interpreted the data, and drafted the manuscript. All authors revised

the paper.

Acknowledgments

We thank all PwPDs for their collaboration in this study.

References

[1] Wilson M. Six views of embodied cognition. Psychon Bull Rev 2002;9:625–36. https://doi.org/10.3758/BF03196322.

[2] Wilson AD, Golonka S. Embodied Cognition is Not What you Think it is. Front Psychol 2013;4. https://doi.org/10.3389/fpsyg.2013.00058.

[3] Morillon B, Hackett TA, Kajikawa Y, et al. Predictive motor control of sensory dynamics in auditory active sensing. Curr Opin Neurobiol 2015;31:230–8. https://doi.org/10.1016/j.conb.2014.12.005.

[4] Amboni M, Barone P, Hausdorff JM. Cognitive contributions to gait and falls: Evidence and implications. Mov Disord 2013;28:1520–33. https://doi.org/10.1002/mds.25674.

[5] Cantiniaux S, Vaugoyeau M, Robert D, et al. Comparative analysis of gait and speech in Parkinson's disease: Hypokinetic or dysrhythmic disorders? J Neurol Neurosurg Psychiatry 2010;81:177–84. https://doi.org/10.1136/jnnp.2009.174375.

[6] Puyjarinet F, Bégel V, Gény C, et al. Heightened orofacial, manual, and gait variability in Parkinson's disease results from a general rhythmic impairment. Npj Park Dis 2019;5:19. https://doi.org/10.1038/s41531-019-0092-6.

[7] Seashore HG. Some relationships of fine and gross motor abilities. Res Q Am Assoc Health Phys Educ Recreat 1942;13:259–74. https://doi.org/10.1080/10671188.1942.10624754.

[8] Oberer N, Gashaj V, Roebers CM. Motor skills in kindergarten: Internal structure, cognitive correlates and relationships to background variables. Hum Mov Sci 2017;52:170–80. https://doi.org/10.1016/j.humov.2017.02.002.

[9] Cavallieri F, Di Rauso G, Gessani A, et al. A study on the correlations between acoustic speech variables and bradykinesia in advanced Parkinson's disease. Front Neurol 2023;14:1213772. https://doi.org/10.3389/fneur.2023.1213772.

[10] Cavallieri F, Gessani A, Merlo A, et al. Interplay between speech and gait variables in Parkinson's disease patients treated with subthalamic nucleus deep brain stimulation: A long□term instrumental assessment. Eur J Neurol 2023;30:1963–72. https://doi.org/10.1111/ene.15803.

[11] Sorgente V, Cohen EJ, Bravi R, et al. Crosstalk between gross and fine motor domains during late childhood: The influence of gross motor training on fine motor performances in primary school children. Int J Environ Res Public Health 2021;18. https://doi.org/10.3390/ijerph182111387.

[12] Gonzalez SL, Alvarez V, Nelson EL. Do Gross and Fine Motor Skills Differentially Contribute to Language Outcomes? A Systematic Review. Front Psychol 2019;10:2670–2670. https://doi.org/10.3389/fpsyg.2019.02670.

[13] Piek JP, Dawson L, Smith LM, et al. The role of early fine and gross motor development on later motor and cognitive ability. Hum Mov Sci 2008;27:668–81. https://doi.org/10.1016/j.humov.2007.11.002.

[14] Callisaya ML, Blizzard CL, Wood AG, et al. Longitudinal Relationships between Cognitive Decline and Gait Slowing: The Tasmanian Study of Cognition and Gait. J Gerontol - Ser Biol Sci Med Sci 2014;70:1226–32. https://doi.org/10.1093/gerona/glv066.

[15] Hughes a J, Daniel SE, Kilford L, et al. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. J Neurol Neurosurg Psychiatry 1992;55:181–4. https://doi.org/10.1136/jnnp.55.3.181.

[16] Dahdal P, Meyer A, Chaturvedi M, et al. Fine motor function skills in patients with Parkinson disease with and without mild cognitive impairment. Dement Geriatr Cogn Disord 2016;42:127–34. https://doi.org/10.1159/000448751.

[17] Intzandt B, Beck EN, Silveira CRA. The effects of exercise on cognition and gait in Parkinson's disease: A scoping review. Neurosci Biobehav Rev 2018;95:136–69. https://doi.org/10.1016/j.neubiorev.2018.09.018.

[18] Dalla Bella S, Falk S. Rhythmic processes in stuttering and Parkinson's disease. Rhythms Speech Lang. Cambridge University Press, Cambridge: n.d.

[19] Repp BH, Su Y-H. Sensorimotor synchronization: A review of recent research (2006–2012). Psychon Bull Rev 2013;20:403–52. https://doi.org/10.3758/s13423-012-0371-2.

[20] Dalla Bella S, Farrugia N, Benoit CE, et al. BAASTA: Battery for the Assessment of Auditory Sensorimotor and Timing Abilities. Behav Res Methods 2017;49:1128–45. https://doi.org/10.3758/s13428-016-0773-6.

[21] Dauvergne C, Bégel V, Gény C, et al. Home-based training of rhythmic skills with a serious game in Parkinson's disease: Usability and acceptability. Ann Phys Rehabil Med 2018;61:380–5. https://doi.org/10.1016/J.REHAB.2018.08.002.

[22] Dalla Bella S, Benoit C-E, Farrugia N, et al. Gait improvement via rhythmic stimulation in Parkinson's disease is linked to rhythmic skills. Nat Publ Group 2017;7:42005. https://doi.org/10.1038/srep42005.

[23] Benoit C-E, Dalla Bella S, Farrugia N, et al. Musically Cued Gait-Training Improves Both Perceptual and Motor Timing in Parkinson's Disease. Front Hum Neurosci 2014;8:494. https://doi.org/10.3389/fnhum.2014.00494.

[24] Skodda S, Flasskamp A, Schlegel U. Instability of syllable repetition as a model for impaired motor processing: is Parkinson's disease a "rhythm disorder"? J Neural Transm 2010;117:605–12. https://doi.org/10.1007/s00702-010-0390-y.

[25] Puyjarinet F, Bégel V, Lopez R, et al. Children and adults with Attention-Deficit/Hyperactivity Disorder cannot move to the beat. Sci Rep 2017 71 2017;7:1–11. https://doi.org/10.1038/s41598-017-11295-w.

[26] Tierney A, Kraus N. The ability to tap to a beat relates to cognitive, linguistic, and perceptual skills. Brain Lang 2013;124:225–31. https://doi.org/10.1016/j.bandl.2012.12.014.

[27] Biswas A, Hegde S, Jhunjhunwala K, et al. Two sides of the same coin: Impairment in perception of temporal components of rhythm and cognitive functions in Parkinson's disease. Basal Ganglia 2016;6:63–70. https://doi.org/10.1016/j.baga.2015.12.001.

[28] Puyjarinet F, Bégel V, Geny C, et al. At-Home Training With a Rhythmic Video Game for Improving Orofacial, Manual, and Gait Abilities in Parkinson's Disease: A Pilot Study. Front Neurosci 2022;16. https://doi.org/10.3389/fnins.2022.874032.

[29] Marsh T. Serious games continuum: Between games for purpose and experiential environments for purpose. Entertain Comput 2011;2:61–8. https://doi.org/10.1016/j.entcom.2010.12.004.

[30] de Oliveira LC, Mendes LC, de Lopes RA, et al. A systematic review of serious games used for rehabilitation of individuals with Parkinson's disease. Res Biomed Eng 2021;37:849–65. https://doi.org/10.1007/s42600-021-00162-3.

[31] Bégel V, Seilles A, Dalla Bella S. Rhythm Workers: A music-based serious game for training rhythm skills. Music Sci 2018;1. https://doi.org/10.1177/2059204318794369/ASSET/IMAGES/LARGE/10.1177_20592043 18794369-FIG6.JPEG.

[32] Dalla Bella S. Rhythmic serious games as an inclusive tool for music-based interventions. Ann N Y Acad Sci 2022;1517:15–24. https://doi.org/10.1111/nyas.14878.

[33] Haier RJ, Karama S, Leyba L, et al. MRI assessment of cortical thickness and functional activity changes in adolescent girls following three months of practice on a visual-spatial task. BMC Res Notes 2009;2:174. https://doi.org/10.1186/1756-0500-2-174.

[34] Lindstedt JK, Gray WD. Meta-T: Tetris⬚ as an experimental paradigm for cognitive skills research. Behav Res Methods 2015;47:945–65. https://doi.org/10.3758/s13428-014-0547-y.

[35] Whitehead AL, Julious SA, Cooper CL, et al. Estimating the sample size for a pilot randomised trial to minimise the overall trial sample size for the external pilot and main trial for a continuous outcome variable. Stat Methods Med Res 2016;25:1057–73. https://doi.org/10.1177/0962280215588241.

[36] Iversen JR, Patel AD. The Beat Alignment Test (BAT): Surveying beat processing abilities in the general population. In: Miyazaki K, Hiraga Y, Adachi M, et al., editors. Proc. 10th Int. Conf. Music Percept. Cogn. ICMPC10, Adelaide: Causal Productions; 2008, p. 465–8.

[37] Dalla Bella S, Foster N, Laflamme H, et al. Mobile version of the Battery for the Assessment of Auditory Sensorimotor and Timing Abilities (BAASTA): Implementation and adult norms. Behav Res Methods 2024. https://doi.org/10.3758/s13428-024-02363-x.

[38] Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. J Am Geriatr Soc 2005;53:695–9. https://doi.org/10.1111/j.1532-5415.2005.53221.x.

[39] Scarpina F, Tagini S. The stroop color and word test. Front Psychol 2017;8. https://doi.org/10.3389/fpsyg.2017.00557.

[40] Grégoire J. Factor structure of the French version of the Wechsler Adult Intelligence Scale-III. Educ Psychol Meas 2004;64:463–74. https://doi.org/10.1177/0013164403258452.

[41] Ashendorf L, Jefferson AL, O'Connor MK, et al. Trail Making Test errors in normal aging, mild cognitive impairment, and dementia. Arch Clin Neuropsychol 2008;23:129–37. https://doi.org/10.1016/j.acn.2007.11.005.

[42] R Core Team. R: A Language and Environment for Statistical computing. R Foundation for Statistical Computing 2023.

[43] Singmann H, Bolker B, Westfall J, et al. Afex: analysis of factorial experiments. R package version 1.3-0. 2023.

[44] Lenth R V. Emmeans: estimated marginal means, aka least-squares means. R package version 1.8.9 2023.

[45] Hass CJ, Bishop M, Moscovich M, et al. Defining the clinically meaningful difference in gait speed in persons with Parkinson disease. J Neurol Phys Ther 2014;38:233–8. https://doi.org/10.1097/NPT.0000000000000055.

[46] Cochen De Cock V, Dotov DG, Ihalainen P, et al. Rhythmic abilities and musical training in Parkinson's disease: do they help? Npj Park Dis 2018;4:8. https://doi.org/10.1038/s41531-018-0043-7.

[47] Wang L, Peng J, Ou-Yang J, et al. Effects of Rhythmic Auditory Stimulation on Gait and Motor Function in Parkinson's Disease: A Systematic Review and Meta-Analysis of Clinical Randomized Controlled Studies. Front Neurol 2022;13:818559. https://doi.org/10.3389/fneur.2022.818559.

[48] Ye X, Li L, He R, et al. Rhythmic auditory stimulation promotes gait recovery in Parkinson's patients: A systematic review and meta-analysis. Front Neurol 2022;13:940419. https://doi.org/10.3389/fneur.2022.940419.

[49] Wegrzyk J, Armand S, Chiuvé SC, et al. Education level affects dual-task gait after deep brain stimulation in Parkinson's disease. Parkinsonism Relat Disord 2019;68:65–8. https://doi.org/10.1016/j.parkreldis.2019.10.005.

[50] Ghai S, Ghai I, Schmitz G, et al. Effect of rhythmic auditory cueing on parkinsonian gait: A systematic review and meta-analysis. Sci Rep 2018;8:1–19. https://doi.org/10.1038/s41598-017-16232-5.

[51] Dalla Bella S. The use of rhythm in rehabilitation for patients with movement disorders. In: Cuddy LL, Belleville S, Moussard A, editors. Music Aging Brain. Academic Press, Elsevier; 2020, p. 383–406. https://doi.org/10.1016/B978-0-12-817422-7.00015-8.

[52] Bourdon A, Damm L, Dotov D, et al. Gait ecological assessment in persons with Parkinson's disease engaged in a synchronized musical rehabilitation program. Npj Park Dis 2025;11:12. https://doi.org/10.1038/s41531-024-00852-6.

[53] Sibley KM, Beauchamp MK, Van Ooteghem K, et al. Using the systems framework for postural control to analyze the components of balance evaluated in standardized balance measures: A scoping review. Arch. Phys. Med. Rehabil., vol. 96, W.B. Saunders; 2015, p. 122-132.e29. https://doi.org/10.1016/j.apmr.2014.06.021.

[54] Pilegard C, Mayer RE. Game over for Tetris as a platform for cognitive skill training. Contemp Educ Psychol 2018;54:29–41. https://doi.org/10.1016/j.cedpsych.2018.04.003.

[55] Belchior P, Marsiske M, Sisco SM, et al. Video game training to improve selective visual attention in older adults. Comput Hum Behav 2013;29:1318–24. https://doi.org/10.1016/j.chb.2013.01.034.

[56] Terlecki MS, Newcombe NS, Little M. Durable and generalized effects of spatial experience on mental rotation: Gender differences in growth patterns. Appl Cogn Psychol 2008;22:996–1013. https://doi.org/10.1002/acp.1420.

[57] Tommasi G, Fiorio M, Yelnik J, et al. Disentangling the role of cortico-basal ganglia loops in top–down and bottom–up visual attention: An investigation of attention deficits in parkinson disease. J Cogn Neurosci 2015;27:1215–37. https://doi.org/10.1162/jocn_a_00770.

[58] Possin KL. Visual spatial cognition in neurodegenerative disease. Neurocase 2010;16:466–87. https://doi.org/10.1080/13554791003730600.

[59] Wielinski CL, VandenDolder R, McManus L, et al. Attention training improves attention and gait in Parkinson disease: A pilot study. Parkinsonism Relat Disord 2019;60:188–9. https://doi.org/10.1016/j.parkreldis.2018.08.015.

[60] Woollacott M, Shumway-Cook A. Attention and the control of posture and gait: a review of an emerging area of research. Gait Posture 2002;16:1–14.

[61] Zatorre RJ. Musical pleasure and reward: mechanisms and dysfunction. Ann N Y Acad Sci 2015;1337:202–11. https://doi.org/10.1111/nyas.12677.

[62] Lorenz RC, Gleich T, Gallinat J, et al. Video game training and the reward system. Front Hum Neurosci 2015;9. https://doi.org/10.3389/fnhum.2015.00040.

[63] Cramb KML, Beccano-Kelly D, Cragg SJ, et al. Impaired dopamine release in Parkinson's disease. Brain 2023;146:3117–32. https://doi.org/10.1093/brain/awad064.

Supplementary table. Scores at the neuropsychological tests and the BAT				
	Intervention (N = 30)		No intervention (N = 12)	
	Mean (SD)		Mean (range)	
Total score	Pre	Post	Pre	Post
MoCA	25.30 (2.73)	25.42 (2.73)	25.73 (3.00)	25.73 (3.00)
Stroop (Errors)	2.76 (3.26)	3.22 (3.19)	2.60 (2.78)	2.95 (1.54)
Stroop (Time)	81.26 (42.76)	85.80 (53.62)	87.69 (29.57)	92.60 (27.13)
Digit Span (Forward)	8.11 (2.15)	8.00 (1.98)	8.91 (2.47)	7.91 (0.35)
Digit Span (Backward)	5.59 (1.80)	5.32 (1.83)	5.27 (2.05)	5.64 (2.42)
TMT B (Errors)	0.78 (0.90)	0.63 (1.05)	1.18 (1.15)	1.05 (1.44)
TMT B (Time)	67.00 (30.22)	72.46 (46.51)	85.14 (26.70)	81.86 (25.55)
BAT	1.79 (1.13)	1.96 (1.03)	1.84 (0.85)	1.87 (0.87)