AMF primes immune genes against *Puccinia hordei* (Brown rust) in *Hordeum vulgare*
but does not reduce pathogen burden

Claire Moulton-Brown[1]*, Karolina Brzezinska[2]*, Beatriz Orosa-Puente[2], Thorunn Helgason[1]†

[1]Institute of Ecology & Evolution, School of Biological Sciences, University of Edinburgh,
Edinburgh, EH9 3BF, UK; [2]Institute of Molecular Plant Sciences, School of Biological
Sciences, University of Edinburgh, Edinburgh, EH9 3BF, UK

* These authors contributed equally to this work
† Corresponding author: Thorunn Helgason,

Institute of Ecology & Evolution,
School of Biological Sciences,
University of Edinburgh,
Edinburgh, EH9 3BF, UK

thorunn.helgason@ed.ac.uk

Abstract

Barley (*Hordeum vulgare*) is susceptible to *Puccinia hordei* (leaf rust), a biotrophic foliar pathogen contributing to global yield losses. With rising food demand and increasing disease pressure, sustainable crop protection strategies are urgently needed to support UN Sustainable Development Goal 2: Zero Hunger. Arbuscular mycorrhizal (AM) fungi, including *Rhizophagus irregularis*, form symbioses with barley roots and can modulate host immunity through mycorrhiza-induced resistance (MIR). Here, we tested whether *R. irregularis* colonisation alters barley growth and defence responses during *P. hordei* infection.

AM fungal colonisation did not significantly reduce disease severity or mitigate pathogen-associated biomass loss at a single post-infection time point. However, co-infected plants showed enhanced expression of defence genes (*PR1*, *PR2*, *PR3* and *WRKY28*), which remained low in plants colonised by AM fungi alone, consistent with immune priming. RNA sequencing revealed AM fungal-associated reprogramming of the leaf transcriptome, including enrichment of defence, metabolism and ubiquitination-related processes.

These results indicate that *R. irregularis* reshapes barley immune regulatory networks at transcriptional and post-translational levels. Although these molecular changes did not translate into measurable phenotypic protection within the short experimental timeframe, they highlight the complexity and context dependence of MIR in cereal–rust interactions.

Introduction

Food security faces mounting pressure as global demand is projected to rise by 35–56 % by 2050 (van Dijk *et al.* 2021). Crops are fundamental to global food security, yet pathogens already reduce yields by 20–40 % (Ristaino *et al.* 2021). Barley (*Hordeum vulgare*), the world's fourth most cultivated cereal, is highly vulnerable to leaf rust (*Puccinia hordei),* which can reduce yield by up to 62 % in susceptible cultivars (Cotterill *et al.* 1992). Epidemics are driven by mild winters, wet springs and warm summers; conditions that are predicted to become more frequent with climate change (West *et al.* 2012). Current management relies on partially resistant cultivars, removal of volunteer hosts and fungicide application. However, these strategies are not fully effective as they are constrained by limited genetic resistance in UK barley (maximum Agriculture and Horticulture Development Board (AHDB) ratings of 6.2 for winter and 4.2 for spring varieties; AHDB 2025) and present sustainability concerns. Therefore, new approaches are needed to enhance crop resilience and reduce reliance on chemical inputs, in line with UN Sustainable Development Goal 2 (Zero Hunger and promotion of sustainable agriculture).

Arbuscular mycorrhizal (AM) symbiotic fungi represent an important strategy for promoting sustainable agriculture and enhancing food security. AM fungi (subphylum Glomeromycotina) colonise the roots of most vascular plants, including cereals (Wang and Qiu 2006). They facilitate nutrient and water uptake in exchange for sugars/lipids, leading to improved crop growth and yield, particularly under resource-limited conditions. Meta-analyses show that AM fungal inoculation increases cereal grain yields by up to 23 % under rainfed systems (Wu *et al.* 2022). Beyond nutrition, AM fungi also enhance tolerance to abiotic stressors such as drought and contribute to soil health via improved structure and microbial activity (Diagne *et al.* 2020).

At the molecular level, symbiosis with AM fungi has been shown to trigger extensive
transcriptional reprogramming in host plants, including the activation of genes involved in
nutrient transport, hormone signalling, and stress adaptation (Gao *et al.* 2023; Gong *et al.*
2023). Crucially, AM fungi can reduce susceptibility to pests and pathogens by modulating
host immunity in a phenomenon known as mycorrhiza-induced resistance (MIR) (Farhaoui *et*
al. 2025). The use of AM fungal symbioses is proposed as a sustainable strategy to improve
cereal crop productivity while reducing dependence on synthetic fertilisers and pesticides
(Bender, Wagg and van der Heijden 2016; Umer *et al.* 2025). To do this effectively, we must
understand the mechanisms that underpin these processes, so that either management
strategies that enhance AM function (Austen *et al.* 2022), or appropriate inoculants can be
identified that genuinely enhance productivity (Elliot *et al.* 2020).

Mycorrhiza-induced resistance (MIR) provides systemic protection against a broad range of
pathogens and shares similarities with both pathogen-triggered systemic acquired resistance
and rhizobacteria-induced systemic resistance (Cameron *et al.* 2013). Rather than triggering
constitutive defence activation, MIR predominantly functions through immune priming,
whereby defence responses are sensitised and more rapidly or strongly deployed upon
pathogen challenge, minimising the fitness costs associated with sustained immune
activation (Jung *et al.* 2012).

AM fungal colonisation remodels host immune signalling networks, enabling modified
defence responsiveness at sites distinct from the roots. Hormonal crosstalk plays a central
role in this process. Early models suggested that AM fungi suppress salicylic acid (SA)-
dependent defences while enhancing jasmonic acid (JA)-mediated responses, conferring
resistance to necrotrophs and herbivores (Cameron *et al.* 2013). However, more recent
findings indicate that MIR can amplify both SA- and JA-responsive genes upon pathogen
challenge, showing a more integrated modulation of multiple defence pathways (Fujita *et al.*
2022).

Importantly, MIR is systemic in nature, with AM fungal colonisation inducing transcriptional
reprogramming not only in roots but also in distal tissues such as leaves, where altered
defence signalling and priming responses to foliar pathogens are ultimately expressed (Pozo
et al. 2002). The MIR phenotype has been reported across diverse plant taxa. In tomatoes
(*Solanum lycopersicum*), *Funneliformis mosseae* colonisation enhanced resistance to
Alternaria blight through JA-dependent signalling, and protection was lost in JA-pathway
mutants (Song *et al.* 2015). In wheat (*Triticum aestivum*), AM fungal colonisation with *F.*
mosseae reduced susceptibility to powdery mildew (*Blumeria graminis*) via enhanced
accumulation of defence metabolites and upregulation of pathogenesis related (PR) and
phenylpropanoid biosynthetic genes (Mustafa *et al.* 2017). Similarly, *F. mosseae* was found
to prime wheat against *Zymoseptoria tritici* (Allario *et al.* 2025), and *Rhizophagus irregularis*
improves rice (*Oryza sativa*) resistance to *Magnaporthe oryzae,* which in turn increased
grain yield in field settings (Martín-Cardoso *et al.* 2025). These cases demonstrate the
relevance of MIR to both monocot and dicot systems and highlight its potential to enhance
disease resistance and yield concurrently, a central goal of sustainable crop protection.

Mechanistically, MIR involves transcriptional and post-transcriptional reprogramming of
immune signalling networks. Mycorrhizal colonisation alters the expression of transcription
factors, including WRKY proteins, which regulate both basal and inducible defences. In
apple (*Malus domestica*), AM fungal-induced *WRKY40* enhanced resistance to *Fusarium* by
activating *β-1,3-glucanase* expression (Wang *et al.* 2022). Priming also enhanced physical

defences, such as callose deposition, which restricts pathogen entry. Blocking callose
synthesis abolishes MIR protection in tomato (Sanmartín *et al.* 2021), indicating its functional
importance.

Beyond transcriptional changes, MIR includes modulation of the ubiquitin-proteasome
system (UPS), a major post-translational regulator of immunity. E3 ubiquitin ligases, which
confer substrate specificity in UPS, target immune regulators for proteasomal degradation,
balancing activation and repression of defences (de Vega, Newton and Sadanandom 2018).
Several *PUB* and *Arabidopsis Toxicos en Levadura* (*ATL*) family ligases are upregulated in
mycorrhizal roots and shoots (Wu *et al.* 2024), suggesting that AM fungi may fine-tune
immune signalling by modifying proteostasis. This regulation may allow plants to remain
immunologically alert without incurring the growth penalties of constitutive defence.

The magnitude and outcome of MIR can be highly context dependent. Factors including
plant nutrient status, AM fungal genotype, pathogen lifestyle, and environmental conditions
can influence whether mycorrhiza-induced changes translate into effective resistance
(Dejana *et al.* 2022; Weinberger *et al.* 2025). Understanding how these variables interact in
cereal systems is vital if MIR is to be harnessed as a reliable strategy for sustainable crop
protection. Despite compelling evidence from other pathosystems, the effectiveness and
molecular basis of MIR in barley, particularly against biotrophic foliar pathogens like *P.*
hordei, remains unstudied. Evidence from cereal-pathogen-symbiont tripartite systems is
needed to validate MIR as a practical tool for disease management. In this study, we
hypothesised that colonisation of barley by *Rhizophagus irregularis* (a widely distributed AM
fungus) primes the barley immune system, which could attenuate *P. hordei* infection or
disease development.

First, we quantified barley growth traits and rust disease severity to assess whether
arbuscular mycorrhizal fungal colonisation influences host fitness or pathogen susceptibility.
These measurements directly test the hypothesis that AM fungi enhance tolerance or
resistance to *Puccinia hordei* without incurring growth penalties, a hallmark of priming-based
immune responses. Second, we examined basal and pathogen-induced expression of key
defence-associated gene families, including pathogenesis-related (PR) proteins and WRKY
transcription factors. These gene groups were selected as central regulators of plant
immunity. Alterations in the basal expression of plant immune markers are indicative of a
primed state, whereas changes in induction kinetics or amplitude upon pathogen challenge
reflect enhanced defence activation. Third, we conducted transcriptomic analyses of leaf
tissue to capture systemic reprogramming of defence-related pathways in response to AM
fungal colonisation. This enabled assessment of broader immune signalling networks,
including transcriptional regulation and ubiquitin-mediated protein turnover, which are
increasingly recognised as key components of immune priming and defence fine-tuning.

By integrating physiological phenotyping with targeted and global transcriptional analyses,
this study aims to assess whether *R. irregularis* can induce mycorrhiza-induced resistance
(MIR) in the barley-*P. hordei* pathosystem and to characterise its potential mechanistic basis.
Given emerging evidence that MIR outcomes and underlying regulatory networks can vary
depending on host species, microbial symbionts, and pathogen lifestyles, dissecting MIR in
this economically relevant cereal-rust interaction is critical for distinguishing conserved
immune features from system-specific regulatory mechanisms. Such comparative resolution

is essential for assessing the translational potential of microbiome-based resistance

strategies across diverse crop-pathogen combinations.

Materials and Methods

Experimental design

A 2 × 2 factorial experiment was conducted to assess the impact of AM fungal colonisation

and foliar pathogen infection on barley. Treatments comprised: (i) **C**ontrol (no AM fungi, no

pathogen), (ii) **A**M fungi only (*Rhizophagus irregularis*), (iii) *Puccinia hordei* only, and (iv)

both AM fungi and *P. hordei* (**AP**). Each treatment group included six biological replicates,

with a subset of three replicates used for transcriptomic analysis and ubiquitination

quantification. Barley was grown for four weeks, then inoculated with *P. hordei* spores. The

experiment was terminated one week post-inoculation (Figure 1), a key timepoint for the

detection of differences in infection establishment and defence gene induction that are most

informative for elucidating MIR mechanisms (Arntzen and Parlevliet 1986).

Plant growth conditions and AM fungal inoculation

Seeds of *Hordeum vulgare* cv. Bowman, a rust-susceptible cultivar, were surface sterilised

by vortexing in 30 % (v/v) bleach supplemented with 0.1 % (v/v) Triton X-100 using three

sequential incubation steps (15 s vortex followed by 6 min at room temperature, repeated

twice, followed by a final 20 s vortex). Seeds were then washed three times with sterile

deionised water and sown in a 1:1 (v/v) mixture of autoclaved coarse silica sand and

Terragreen® clay (Oil-Dri, UK). Each 1 L pot was supplemented with 0.5 g L⁻¹ bonemeal and

watered twice weekly with phosphate-free Hoagland's solution to promote mycorrhization.

Plants were grown in a controlled-environment chamber (MLR-352H, PHC, Japan; 20°C day

/ 17°C night; 70 % relative humidity) under a 16 h light/8 h dark photoperiod, with fluorescent

lighting (FL40SSENW37) providing 150 µmol m⁻² s⁻¹.

AM fungi treated plants received commercial *Rhizophagus irregularis* inoculum (PlantWorks,

UK) at 20 g L⁻¹ of growth substrate, a rate previously shown to ensure consistent and robust

AM fungal colonisation in controlled experiments using sterilised media (Dejana *et al.* 2022;

Weinberger *et al.* 2025). Controls received autoclaved inoculum and microbial filtrate to

standardise non-AM fungal microbial communities.

Pathogen inoculation

Puccinia hordei isolate BBR 20-003 was obtained from the National Institute of Agricultural

Botany (NIAB). Urediniospores (~1.3 × 10☐ spores plant⁻¹) were applied to leaves using

talcum powder as an inert carrier (Roelfs, Singh and Saari 1992) at a spore: talcum ratio of

1:10 (w/w). Spores were mixed with talcum powder immediately prior to inoculation and

dusted onto the adaxial leaf surface using a fine brush to ensure uniform distribution. Control

plants received talcum powder only. To promote infection and prevent cross-infection, *P.*

hordei-inoculated plants were maintained at >95 % relative humidity for 12 hours post-

inoculation and transferred to a separate growth cabinet under identical growth conditions.

Confirmation of AM fungal colonisation

AM fungal colonisation was verified by both microscopy and qPCR on root samples. For

microscopy, roots were stained using the ink and vinegar method (Vierheilig *et al.* 1998).

Roots were stored in 40 % ethanol at 4°C, then cleared with 10 % KOH at 70°C for 45

minutes. They were stained with 5 % Brilliant Black ink (Pelikan) in 5 % acetic acid,
destained in 1 % acetic acid, and mounted on slides with 50 % (v/v) glycerol.

For AM fungal qPCR, DNA was extracted from roots using the DNeasy Plant Mini Kit
(Qiagen). Each 10 µL qPCR reaction included Fast SYBR™ Green Master Mix (Thermo
Fisher Scientific), 350 nM of each primer, and ~10 ng of root DNA. AM fungi were quantified
using the 18S rRNA gene with AM fungal-specific primers AMG1F and AM1 (Table S1;
Helgason et al. 1998; Hewins, Carrino-Kyker and Burke 2015), normalised to barley
Elongation Factor 1-α using primers HvEF1a-qF and HvEF1a-qR (Hua *et al.* 2015). Triplicate
reactions were run on a 7500 Fast Real-Time PCR System. The program comprised an
initial denaturation at 95°C for 20 s, followed by 40 cycles at 95°C for 10 s and 60°C for 30 s,
then a melt curve from 60-95°C. Data were analysed with Applied Biosystems 7500
Software (v2.3) using the Pfaffl method for relative quantification.

Rust infection quantification

Pathogen biomass in leaves was quantified by RT-qPCR (see Gene expression analysis
below for conditions). Leaf RNA was extracted and converted to cDNA (see below). Primers
targeting the *P. hordei* elongation factor 2 gene (Table S1) were used to measure pathogen
cDNA relative to barley cDNA (using a barley endogenous actin gene as reference). This
provided a relative infection index for each plant.

Barley resource allocation measurements

Root and shoot dry weights were recorded at harvest. Plants were divided into above- and
below-ground fractions and dried at 60°C until a constant weight was reached. Total biomass
and root:shoot ratio were calculated.

RNA extraction and cDNA synthesis

Total RNA was extracted from barley leaf tissue using a phenol-chloroform-isoamyl alcohol
method with LiCl precipitation, adapted from established protocols (Verwoerd, Dekker and
Hoekema 1989). Briefly, frozen plant material was ground under liquid nitrogen, extracted
with warmed phenol : chloroform : isoamyl alcohol (25 : 24 : 1), and subjected to sequential
organic washes. The aqueous phase was precipitated with 1/3 volume of 8 M LiCl overnight
at 4°C, and the resulting pellet was washed with 70 % ethanol. A secondary precipitation
with sodium acetate and 96 % ethanol was performed to ensure purity, and RNA pellets
were finally resuspended in RNase-free water. cDNA was synthesised using the
SuperScript™ III Reverse Transcriptase (Invitrogen) according to the manufacturer's
instructions.

Gene expression analysis

Quantitative RT-PCR was used to assess the expression of immune marker genes, *PR1*,
PR2, *PR3*, and *WRKY28*. Reactions used SYBR™ Green Universal Master Mix (Applied
Biosystems, Thermo Fisher Scientific). Reactions were conducted in technical triplicates,
each containing 2 µL of cDNA and 200 nM gene-specific primers (Table S1). The reactions
were run on a QuantStudio 5 system (Applied Biosystems, Thermo Fisher Scientific). The
cycling conditions were: 50°C for 2 minutes and 95°C for 10 minutes; 40 cycles of 95°C for
15 seconds, 60°C for 1 minute; and a melt curve at 95°C for 15 seconds, 60°C for 1 minute,
and 95°C for 15 seconds. Relative expression was calculated by the ΔCt method and
normalised to the barley housekeeping gene (Actin).

RNA sequencing and transcriptomics

RNA sequencing was conducted on a subset of samples (n = 3 per treatment). RNA sequencing was performed by Novogene Company Limited using the Illumina NovaSeq Platform (150 bp paired end). Reads were quality assessed using FastQC v0.11.9 (Andrews 2010). Adapter sequences were trimmed using Trimmomatic v0.39 (Bolger, Lohse and Usadel 2014). After adapter and quality trimming, we obtained on average 43.69, 42.83, 32.90, and 44.11 Mb clean reads from the Control, AM fungi, *P. hordei*, and AM fungi plus *P. hordei* treatments, respectively (Table S2), with a similar GC content across treatments (53.34 %). RNA-sequencing reads were aligned and annotated against the barley genome (MorexV3) using HISAT2 v2.1.0 (Kim *et al.* 2019).

Ubiquitination assay

Given the enrichment of ubiquitin-related genes in AM fungal-treated plants, global protein ubiquitination was assessed by Western blot. Total protein was extracted from frozen barley leaves using a RIPA-like buffer adapted from established plant protocols (Chico *et al.* 2020). Briefly, tissue was ground under liquid nitrogen and homogenised in extraction buffer (20 mM Tris pH 7.5, 50 mM NaCl, 1 mM EDTA, 0.5 % NP40, 0.5 % SDS, 0.5 % sodium deoxycholate) supplemented with protease inhibitors, 50 µM MG132, and 10 mM N-ethylmaleimide (NEM). Extracts were vortexed, incubated on ice for 10 min, and centrifuged at 10,000 rpm for 15 min at 4°C. The supernatant was collected for downstream analysis.

Protein extracts were heated in SDS sample buffer with 50 mM DTT at 80°C for 10 minutes and resolved by SDS-PAGE. Proteins were transferred to a nitrocellulose membrane via wet transfer at 20 V overnight. Transfer efficacy was verified with Ponceau staining. Membranes were blocked with 5 % (w/v) skim milk powder (Millipore) in PBST for 2 hours, then incubated with primary antibodies (Poly-ubiquitin P4D1 at 1:2500, Santa Cruz Biotechnology; Actin at 1:5000, Agrisera; actin was used as a loading control for normalisation) for 1.5 hours, followed by three 5-minute washes in PBST. Secondary antibody, anti-mouse 1:5000 (Cell Signaling Technology), incubation lasted 1 hour, followed by three additional washes. Detection was performed using a chemiluminescent substrate (SuperSignal™ West Pico PLUS, Thermo Fisher Scientific) and visualised with LI-COR® Odyssey Fc Imager with band intensity quantified using Image Studio 6.1 (LICORbio).

Statistical analysis

Statistical analyses were performed using R v4.5.1 (R Core Team 2025) implemented in RStudio version 2025.09.2 (RStudio Team 2025). Data processing, reshaping, and visualisation were performed using tidyverse v2.0.0 (Wickham *et al.* 2019), primarily dplyr v1.1.4, tidyr v1.3.1, and ggplot2 v4.0.1, with multi-panel figures assembled using cowplot (v1.2.0).

Physiological traits (root dry mass, shoot dry mass, and root:shoot ratio) and quantitative PCR (qPCR) data were analysed using two-way analysis of variance (ANOVA) to test for the main effects of arbuscular mycorrhizal fungal (AM fungal) inoculation (AM-, AM+) and *Puccinia hordei* infection (Puccinia-, Puccinia+), and their interaction. ANOVA models were fitted using the base R function aov() from the stats package, with F-tests and associated p-values obtained using anova(). qPCR data were log2-transformed prior to statistical analysis. Post-hoc comparisons among factorial treatment combinations were conducted using

estimated marginal means calculated with the emmeans() function from the emmeans
package v2.0.0 (Russell V. Lenth and Julia Piaskowski 2025). Pairwise comparisons were
adjusted for multiple testing using the Sidak method, and statistically distinguishable groups
were summarised using compact letter displays generated with the cld() function from the
multcomp package v1.4-29 (Hothorn, Bretz and Westfall 2008). Exact p-values are reported
in figures, with significance indicated by asterisks (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

RNA gene count data were quantified and standardised using HTseq2 v2.0.5 (Putri *et al.*
2022). Differential analysis was performed using edgeR v3.42.4 (Chen *et al.* 2025) by
filtering out CPM < 1 and normalising using a TMM method (Robinson and Oshlack 2010).
Differentially expressed genes (DEGs) were defined as: FDR < 0.05, $\log_2FC > 1$, logCPM >
2. Gene Ontology (GO) enrichment analysis of DEGs was done using g:Profiler (Raudvere
et al. 2019) with FDR correction. Hierarchical clustering (Ward's method) of DEGs was
employed to identify groups of genes with similar expression patterns across the four
conditions.

Ubiquitination data were normalised to actin and expressed as fold change relative to the
control treatment. Statistical analyses of ubiquitination data were performed on $\log_2$-
transformed fold-change values using the same two-way ANOVA framework described
above.

Assumptions of normality and homoscedasticity were assessed by visual inspection of
ANOVA residuals (residuals versus fitted values and normal Q-Q plots). Where necessary,
data were transformed prior to analysis to improve adherence to model assumptions and
stabilise variances. Independence of observations was ensured by the experimental design,
with each plant treated as an independent experimental unit.

Results

AM fungi alone did not reduce rust infection or improve biomass under pathogen
challenge

Barley plants were inoculated with *Rhizophagus irregularis*, and root colonisation was
confirmed 28 days post-inoculation. Microscopy revealed abundant arbuscules and hyphae
in the inoculated roots, whereas no mycorrhizal structures were observed in non-inoculated
control samples. qPCR further validated successful AM fungal colonisation in inoculated
plants (Figure 2 A-B). Low-level amplification detected in control samples is consistent with
previously reported non-specific amplification by the AM1 primer in the absence of AM fungal
templates (Lee, Lee and Young 2008). AM fungal presence did not significantly affect rust
severity one week after *P. hordei* infection; *P. hordei*-specific marker gene (PhEF2) showed
only a slight, non-significant increase in AM fungi-colonised plants (Figure 2 C).

Puccinia hordei infection resulted in a substantial reduction in plant biomass, with all infected
plants exhibiting approximately 20% lower total dry weight than uninfected controls,
regardless of AM fungal colonisation ($p = 0.000955$; Figure 2 D). No significant AM fungi ×
pathogen interaction was detected, and the root:shoot ratio remained unchanged (Figure
S1), indicating a proportional reduction in overall growth rather than altered biomass
allocation. In contrast, AM fungal colonisation had no detectable effect on barley growth
(Figure 2 D and S1), indicating that AM fungi did not impose a growth penalty under these
experimental conditions.

Overall, *P. hordei* caused substantial biomass losses regardless of AM fungal presence. We
next assessed whether AM fungal colonisation was associated with changes in molecular
immune responses that were not evident in infection or growth metrics in this timeframe.

AM fungi primed the expression of barley defence genes during *P. hordei* infection

Although AM fungi did not reduce rust severity in this experiment, we tested whether AM
colonisation altered defence gene expression patterns during infection. Transcript levels of
SA-responsive PR genes (*PR1*, *PR2*, *PR3*) and the transcription factor *WRKY28* were
measured. In the absence of *P. hordei*, *PR1*, *PR2* and *PR3* transcript levels remained low in
AM-colonised leaves and did not differ from non-colonised controls (Figure 3 A-C).

Following *P. hordei* infection, *PR1*, *PR2* and *PR3* were induced, with higher transcript levels
in AM-colonised infected plants than in infected plants without AM fungi (Figure 3 A-C).
WRKY28 showed a similar pattern, with low expression in uninfected plants and increased
expression following infection, including higher transcript levels in AM fungi-colonised
infected plants (Figure 3 D). This pattern is consistent with enhanced activation of higher-
order immune regulators.

Together, these results show that priming occurred at the transcriptional level. AM fungi
alone did not trigger immune gene expression in the absence of a pathogen but enhanced
the amplitude of the transcriptional response upon infection. The lack of immediate
protection despite strong PR induction suggests that *P. hordei* continues to suppress or
evade host defences, or that timing and physiological constraints limit the effectiveness of
the primed response.

AM fungi reprogrammed the barley leaf transcriptome

For a comprehensive view of the transcriptomic changes induced by AM fungi, we performed
RNA sequencing of barley leaves. AM fungal colonisation altered the expression in healthy
leaves of 526 genes (344 upregulated, 182 downregulated; Figure 4 A; Figure S2). A Gene
Ontology (GO) enrichment analysis revealed that upregulated genes were enriched for
defence, stress responses, and protein modification (Figure 4 B). Additionally, upregulated
transcripts included sugar transporters and amino acid biosynthesis genes, whereas
downregulated transcripts included those annotated for growth, photosynthesis and
gibberellin signalling (Table S3). This suggests AM fungi enhance the host's capacity to
regulate protein turnover. Induction of sugar transporters and amino acid synthesis genes
likely reflects increased nutrient exchange, while downregulated transcripts related to
growth, photosynthesis, and gibberellin signalling indicate a moderated growth programme.
These shifts did not affect overall biomass (Figure 2 D).

P. hordei infection in non-mycorrhizal plants triggered differential expression of 4,258 genes
(P/C: 2,317 upregulated, 1,941 downregulated), indicating a robust transcriptional
reprogramming in response to the pathogen. In comparison, AM fungal colonisation
attenuated the overall magnitude of rust-induced gene expression changes, reducing the
number of upregulated genes by ~15 % and downregulated genes by ~30 % (AP/A, 1,956
upregulated, 1,380 downregulated). Interestingly, during *P. hordei* infection, 231 genes were
differentially expressed in AM-colonised plants compared with non-mycorrhizal plants (AP/P,
163 upregulated, 68 downregulated), suggesting that AM fungi still shape a distinct
transcriptional response. Cluster analysis revealed that AM fungi amplified defence-related
genes, including PRs and glutathione-associated transcripts (Cluster 3), while dampening

broad kinase- and receptor-driven signalling cascades (Clusters 2 and 4; Figure 4 C-D).
Genes induced by AM fungi alone but suppressed during co-infection (Clusters 5 and 6) may
reflect negative feedback on pre-activated signalling modules. Together, these results
suggest that AM fungi modulate transcriptional changes by limiting broad stress responses
and enhancing defence-related gene expression.

AM fungi altered defence-associated signalling and transcriptional responses

To further investigate AM fungal-mediated modulation of plant immune response, we
examined key components of canonical immune signalling pathways. Transcriptional
reprogramming is a hallmark of immune activation and our targeted qPCR analyses had
already indicated that AM fungal colonisation influenced the expression of the immune-
associated transcription factor *WRKY28* (Figure 3 D), which showed enhanced induction in
AM fungi-colonised plants during *Puccinia* infection. This observation prompted a broader
investigation of transcription factor dynamics at the transcriptome scale.

We therefore focused on the WRKY transcription factor family, which comprises major
regulators of both biotic and abiotic stress responses (Javed and Gao 2023; Li *et al.* 2024)
and has previously been shown to be regulated during AM colonisation (Gallou, Declerck
and Cranenbrouck 2012). We found 74 annotated barley WRKY genes that were
differentially expressed across treatments (Table S4). In healthy plants, AM fungi generally
upregulated the expression of WRKY transcription factors, with 95 % of differentially
expressed WRKY genes showing increased expression. In contrast, during *P. hordei*
infection, AM fungi suppressed the majority of rust-induced WRKYs, downregulating 90 % of
those that were otherwise induced by infection alone (Figure 5 A). Notable exceptions
included *WRKY27*, *WRKY63*, and *WRKY1/38*, which were induced even more in co-infected
plants. Most differentially expressed WRKYs belonged to Classes II and III, consistent with a
role in biotic stress regulation.

To assess whether autoregulation and cross-regulation by WRKY transcription factors
contribute to the observed transcriptional changes, we then scanned the 500 bp upstream
promoter regions of all DEGs for W-box elements (TTGACY), the canonical WRKY binding
motif. Approximately 80 % of DEGs contained at least one W-box and gene ontology
enrichment analysis showed their involvement in phosphorylation, alongside terms related to
response to biotic stimulus, stress, and olefinic compound biosynthesis (Figure 5 B).

Since WRKYs are themselves regulated via phosphorylation by calcium-dependent protein
kinases (CDPKs), and AM fungal colonisation upregulated calcium-binding proteins, we also
examined calcium signalling components (Figure 5 C). Several CDPKs, calmodulin, and
calmodulin-like genes were upregulated in AM fungi-colonised healthy plants.

To investigate how AM fungi influence defence-related hormone signalling, we analysed
transcriptional changes in the ethylene, SA, and JA pathways. In healthy AM fungi-colonised
plants, ethylene-responsive genes such as *EIN3* were upregulated, while defence-
associated ethylene responses remained unchanged (Figure S3), indicating a shift towards
metabolic rather than immune regulation. In contrast, several JA synthesis pathway genes,
including *LOX* (lipoxygenase) genes, were expressed at lower levels in AM fungal-colonised
diseased plants (AP) than in plants infected with *P. hordei* alone (P). This did not, however,
affect downstream JA-responsive TIFY genes, which suggests that the effect on JA-
synthesis genes can be unlinked from JA-signalling in this context (Figure S4).

Similarly, for the SA-biosynthesis pathway, AM fungi caused downregulation of CM
(Chorismate mutase) and most PAL (Phenylalanine ammonia lyase) genes during *P. hordei*
infection (Figure S4), while enhancing SA-mediated signalling, as seen by the induction of
SA-responsive PR genes (Figure 3). Expression of *NPR1*, the central SA co-activator, was
also upregulated in AM fungal plants but suppressed by *P. hordei* regardless of AM fungal
status (Figure S4). In contrast, the strong induction of *PR* genes appears to be largely
independent of *NPR1* transcription, which is not markedly affected by pathogen challenge.
(Skelly, Frungillo and Spoel 2016; Withers and Dong 2016; Skelly *et al.* 2019). Overall, AM
fungi reshaped transcriptional profiles of hormone pathways.

AM fungi modulated expression of ubiquitin machinery genes without altering global
ubiquitin levels

Gene Ontology enrichment analysis of AM fungal-induced transcripts showed enrichment of
genes involved in protein modification processes (Figure 4 B), including ubiquitination-
related genes. To determine whether this transcriptional signal translated into changes at the
protein level, we assessed global protein ubiquitination across treatments (Figure S5).
Western blot analysis showed that ubiquitin signal was significantly increased by *Puccinia*
hordei infection ($F_{1,8}$ = 24.20, p = 0.001), with no significant effects of AM fungal inoculation
or AM fungi × *Puccinia* interaction (Figure 6 A), indicating that AM fungi do not induce broad
changes in total protein ubiquitination.

Despite the absence of a global effect, transcriptomic analysis revealed substantial AM
fungi-associated modulation of genes encoding ubiquitination enzymes. Many E3 ubiquitin
ligase genes were differentially expressed, as well as some ubiquitin E1 (activating
enzymes) and E2 (conjugating enzymes) (Figure 6 B). Expression patterns clustered into
five distinct groups with the top cluster upregulated by AM fungi in healthy and infected
plants (Figure 6 B). This small group included several RING/U-box-type E3 ligases from the
ATL family and PUB families, suggesting that AM fungi selectively modulate specific
components of the ubiquitin-proteasome system (UPS), rather than broadly altering the
cellular ubiquitome. In fact, the largest cluster of ubiquitination-related differentially
expressed genes (DEGs) was downregulated in AM-colonised plants during *P. hordei*
infection (Figure 6 B), supporting a targeted rather than global modulation of ubiquitin
signalling.

Discussion
AM fungi have been shown to induce resistance against various pathogens of crop plants
(Fiorilli *et al.* 2011, 2018; Campos-Soriano, García-Martínez and Segundo 2012), and
therefore their application as an alternative crop protection method has been widely explored
(Brito, Carvalho and Goss 2021; Lutz *et al.* 2023). In this study, we investigated whether
colonisation by the arbuscular mycorrhizal (AM) fungus *Rhizophagus irregularis* influences
barley fitness and disease outcomes during infection with the biotrophic foliar pathogen
Puccinia hordei, within the broader context of sustainable crop protection and UN
Sustainable Development Goal 2: Zero Hunger.

AM fungal colonisation induced pathogen-independent transcriptional reprogramming

While AM colonisation did not result in significant growth promotion, reduced infection load, or mitigated *P. hordei*-induced biomass loss, it triggered extensive transcriptional changes in the host. Notably, many of these changes were observed in the absence of pathogen challenge, indicating that AM fungi substantially reshape host regulatory networks independently of disease pressure.

A range of studies have shown that inoculation with AM fungi can promote plant growth, grain yield and vigour (Pellegrino *et al.* 2015; Fiorilli *et al.* 2018; Lutz *et al.* 2023). However, in our study, *Rhizophagus irregularis* colonisation did not result in measurable growth promotion under the tested conditions. This absence of a growth response may reflect limited compatibility between the specific host-symbiont combination, as plant responses to AM fungi are known to vary widely depending on both the fungal isolate and the host genotype (Munkvold *et al.* 2004; Hong *et al.* 2012; Koch *et al.* 2017). Barley cultivars exhibit varying degrees of mycorrhizal responsiveness, with differences in colonisation efficiency and functional outcomes (Peña Venegas *et al.* 2021; Gebremeskel *et al.* 2024; Marrassini *et al.* 2024). These genotype-dependent effects likely contribute to the observed lack of growth benefit in our system and suggest that the *Hordeum vulgare* cv. Bowman – *R. irregularis* pairing forms a transcriptionally active symbiosis with limited physiological output under the tested conditions.

In pathogen-free plants, AM colonisation has been shown to induce extensive transcriptional reprogramming associated with symbiosis establishment and maintenance. These changes include broad remodelling of primary metabolism and carbon allocation, reflecting a shift in host metabolic priorities rather than activation of defence responses (Smith and Smith 2011). Such transcriptional adjustments are thought to support symbiotic function and signalling while avoiding the fitness costs associated with constitutive immunity. Importantly, numerous studies have shown that these pathogen-independent molecular changes can occur without measurable penalties to host growth, highlighting that AM fungi can markedly reshape host regulatory and metabolic states while maintaining overall plant fitness (Smith and Smith 2011; Koch *et al.* 2017).

Pathogen-dependent immune priming in AM-colonised barley

In contrast to the pathogen-independent effects described above, a distinct set of transcriptional responses emerged specifically following *P. hordei* infection. Despite the absence of reduced pathogen burden, AM-colonised plants displayed enhanced induction of genes involved in broad-spectrum immune responses, including salicylic acid (SA)-associated pathways and cell wall defence mechanisms. Such patterns are consistent with immune priming, a hallmark of mycorrhiza-induced resistance (MIR), in which defence responses are sensitised but not constitutively activated (Jung *et al.* 2012; Huot *et al.* 2014).

Crucially, priming does not always confer full immunity. Instead, it can result in partial resistance, slower pathogen development, or reduced sporulation, outcomes that are often not captured by static disease assessments (Pieterse *et al.* 2014). This aligns with previous studies showing that MIR often manifests as subtle, ecologically relevant phenotypes such as delayed symptom onset or attenuated disease progression (Pozo *et al.* 2002; Pozo and

Azcón-Aguilar 2007; Jung *et al.* 2012). Future experiments quantifying disease dynamics
over time (e.g. latent period, spore production) could reveal benefits of MIR that static
measurements miss.

Whether broad-spectrum defence priming alone is sufficient for MIR establishment remains
debated. Several studies have demonstrated that AM fungi can induce pathogen-specific
immune reprogramming distinct from that of non-colonised plants (Campos-Soriano, García-
Martínez and Segundo 2012; Fiorilli *et al.* 2018). In our study, AM fungal colonisation
attenuated the overall magnitude of the *P. hordei*-induced transcriptional response and led to
the activation of a specific subset of defence-related genes. However, this reprogramming
alone was insufficient to confer effective resistance against a highly specialised biotrophic
pathogen such as *P. hordei*. As MIR is highly context-dependent and often biases JA- and
ethylene- mediated defences, it can antagonise SA pathway responses important against
biotrophic pathogens (Cameron *et al.* 2013; Liu *et al.* 2023). This could be reflected in the
elevated defence-related transcript levels, as seen here, without a corresponding increase in
downstream immune outputs (Caldo, Nettleton and Wise 2004; Zierold, Scholz and
Schweizer 2005; Polesani *et al.* 2010). Moreover, pathogen effectors can bypass the priming
by targeting specific immune proteins or disturbing host proteostasis (Bauters, Stojilković
and Gheysen 2021; Kainat *et al.* 2025). Therefore, our findings suggest that the outcome of
AM colonisation in the context of pathogen attack is not solely determined by transcriptional
priming but likely emerges from a complex tripartite interaction between host plant,
mycorrhizal symbiont, and pathogen. Disentangling these multilayered interactions remains
a significant challenge and will require more refined tools and approaches to fully understand
the context-dependence of mycorrhiza-induced resistance.

Transcriptional and post-translational regulators of the primed immune state

Our data identify WRKY transcription factors as potential regulators of the primed immune
state induced by *R. irregularis* in barley. Remarkably, nearly half of all annotated WRKY
genes were differentially expressed, representing a broader-scale alteration of regulatory
networks than previously reported in other systems such as potato, tomato, or apple (Gallou,
Declerck and Cranenbrouck 2012; Cervantes-Gámez *et al.* 2016; Wang *et al.* 2022). This
widespread regulation suggests that AM fungi fine-tune immune responses and
developmental trade-offs through WRKY-mediated transcriptional control. The co-induction
of calcium-dependent protein kinases and calmodulin-related genes further supports the
involvement of calcium-WRKY signalling modules in MIR, consistent with findings in other
host-AM fungal systems (Campos-Soriano, García-Martínez and Segundo 2012; Fiorilli *et al.*
2018).

In parallel, AM colonisation altered the expression of hormonal signalling genes in a manner
that deviates from canonical models of MIR. Suppression of jasmonic acid (JA) biosynthesis
genes and upregulation of TIFY repressors, which inhibit JA signalling, indicates a shift away
from typical JA/ethylene-mediated responses. Interestingly, while salicylic acid (SA)
biosynthesis and signalling genes were downregulated, SA-responsive defence markers
were upregulated during pathogen challenge. This pattern is consistent with an immune
priming mechanism previously described in rice, where AM fungi induced expression of
NPR1, the central coactivator of SA-responsive gene expression, in the leaves of healthy

colonised plants without activating the PR gene expression (Campos-Soriano and Segundo
2011).

This apparent disconnect between NPR1 transcript levels and downstream defence
activation is consistent with previous findings that *NPR1* activity is predominantly regulated
at the post-translational level through redox-dependent conformational changes,
phosphorylation and proteasomal turnover, rather than transcriptional control alone (Skelly,
Frungillo and Spoel 2016; Withers and Dong 2016; Skelly *et al.* 2019). These findings
suggest that AM fungi establish a preconditioned immune state that can be rapidly mobilised
upon pathogen challenge yet may be insufficient to overcome the virulence strategies of
highly specialised biotrophic pathogens such as *P. hordei*. Such priming is therefore likely to
depend not only on transcriptional readiness, but also on post-translational regulatory
mechanisms governing immune protein activation and stability.

In addition to transcriptional reprogramming, we observed significant modulation of the host
ubiquitin-proteasome system (UPS). Although global levels of ubiquitinated proteins
remained unchanged, AM colonisation led to the upregulation of numerous components of
the ubiquitination machinery, including E3 ligases from the PUB and ATL families (Figure 6
B), which are known to regulate diverse stress responses (Guzmán 2012; Trujillo 2018). This
seemingly paradoxical pattern may reflect increased protein turnover and dynamic post-
translational regulation of immune signalling, rather than broad immune activation. Notably,
PUB and ATL ligases have established roles in regulating stress and immune responses,
including chitin-triggered signalling and callose deposition (Berrocal-Lobo *et al.* 2010;
Maekawa *et al.* 2012). Together, our findings suggest that AM fungi-induced immune priming
operates through a multilayered regulatory framework, integrating transcriptional, hormonal,
and post-translational controls, to enhance defence readiness while maintaining metabolic
balance.

Conclusion
Achieving global food security under increasing biotic stress requires sustainable strategies
that reduce crop losses while minimising chemical inputs. While AM fungal colonisation did
not reduce pathogen burden at the phenotypic level in this study, the extensive
transcriptional reprogramming points to a robust immune priming effect that may confer
delayed or context-dependent protection not captured within the experimental timeframe of
this study. Importantly, this immune activation occurred without measurable growth penalties,
suggesting that *R. irregularis,* or other AM fungi, could, pending field validation, serve as a
sustainable tool to enhance crop resilience.

Realising this potential will require integrating AM fungal strategies into broader crop
protection frameworks and continued research into the ecological and mechanistic
dependencies that shape mycorrhiza-induced resistance. By leveraging the beneficial
properties of symbiotic soil fungi, we may unlock new ways to support plant health and
productivity in alignment with the goals of Zero Hunger and sustainable agriculture.

Conflict of Interest
The authors declare no competing interests.

Data availability

RNA-sequence data generated in this study have been deposited in the NCBI Sequence Read Archive (SRA) under accession number PRJNA1328502. The data and R code that support the findings of this study are openly available as a GitHub repository, archived at https://zenodo.org/records/18312009.

References

AHDB. *AHDB Recommended Lists for Cereals and Oilseeds 2025/26.*, 2025.

Allario T, Krzyzaniak Y, Magnin-Robert M *et al.* Defense responses related to mycorrhizal-induced resistance in wheat against Zymoseptoria tritici. *Biological Control* 2025;**203**, DOI: 10.1016/j.biocontrol.2025.105729.

Andrews S. FastQC - A quality control tool for high throughput sequence data. *Babraham Bioinformatics* 2010.

Arntzen FK, Parlevliet JE. Development of barley leaf rust, Puccinia hordei, infections in barley. II. Importance of early events at the site of penetration for partial resistance. *Euphytica* 1986;**35**:961–8.

Bauters L, Stojilković B, Gheysen G. Pathogens pulling the strings: Effectors manipulating salicylic acid and phenylpropanoid biosynthesis in plants. *Mol Plant Pathol* 2021;**22**, DOI: 10.1111/mpp.13123.

Bender SF, Wagg C, van der Heijden MGA. An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. *Trends Ecol Evol* 2016;**31**, DOI: 10.1016/j.tree.2016.02.016

Berrocal-Lobo M, Stone S, Yang X *et al.* ATL9, a RING zinc finger protein with E3 ubiquitin ligase activity implicated in chitin- and NADPH oxidase-mediated defense responses. *PLoS One* 2010;**5**, DOI: 10.1371/journal.pone.0014426.

Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* 2014;**30**, DOI: 10.1093/bioinformatics/btu170.

Brito I, Carvalho M, Goss MJ. Managing the functional diversity of arbuscular mycorrhizal fungi for the sustainable intensification of crop production. *Plants People Planet* 2021;**3**, DOI: 10.1002/ppp3.10212.

Caldo RA, Nettleton D, Wise RP. Interaction-dependent gene expression in Mla-specified response to barley powdery mildew. *Plant Cell* 2004;**16**, DOI: 10.1105/tpc.104.023382.

Cameron DD, Neal AL, van Wees SCM *et al.* Mycorrhiza-induced resistance: More than the sum of its parts? *Trends Plant Sci* 2013;**18**, DOI: 10.1016/j.tplants.2013.06.004.

Campos-Soriano L, García-Martínez J, Segundo BS. The arbuscular mycorrhizal symbiosis promotes the systemic induction of regulatory defence-related genes in rice leaves and confers resistance to pathogen infection. *Mol Plant Pathol* 2012;**13**, DOI: 10.1111/j.1364-3703.2011.00773.x.

Campos-Soriano L, Segundo BS. New insights into the signaling pathways controlling defense gene expression in rice roots during the arbuscular mycorrhizal symbiosis. *Plant Signal Behav* 2011;**6**, DOI: 10.4161/psb.6.4.14914.

Cervantes-Gámez RG, Bueno-Ibarra MA, Cruz-Mendívil A *et al.* Arbuscular Mycorrhizal Symbiosis-Induced Expression Changes in Solanum lycopersicum Leaves Revealed by RNA-seq Analysis. *Plant Mol Biol Report* 2016;**34**, DOI: 10.1007/s11105-015-0903-9.

Chen Y, Chen L, Lun ATL *et al.* edgeR v4: powerful differential analysis of sequencing data
with expanded functionality and improved support for small counts and larger datasets.
Nucleic Acids Res 2025;**53**, DOI: 10.1093/nar/gkaf018.

Chico JM, Lechner E, Fernandez-Barbero G *et al.* CUL3BPM E3 ubiquitin ligases regulate
MYC2, MYC3, and MYC4 stability and JA responses. *Proc Natl Acad Sci U S A*
2020;**117**, DOI: 10.1073/pnas.1912199117.

Cotterill PJ, Rees RG, Platz GJ *et al.* Effects of leaf rust on selected Australian Barleys. *Aust*
J Exp Agric 1992;**32**, DOI: 10.1071/EA9920747.

Dejana L, Ramírez-Serrano B, Rivero J *et al.* Phosphorus availability drives mycorrhiza
induced resistance in tomato. *Front Plant Sci* 2022;**13**, DOI:
10.3389/fpls.2022.1060926.

Diagne N, Ngom M, Djighaly PI *et al.* Roles of arbuscular mycorrhizal fungi on plant growth
and performance: importance in biotic and abiotic stressed regulation. *Diversity (Basel)*
2020;**12**, DOI: 10.3390/d12100370.

van Dijk M, Morley T, Rau ML *et al.* A meta-analysis of projected global food demand and
population at risk of hunger for the period 2010–2050. *Nat Food* 2021;**2**, DOI:
10.1038/s43016-021-00322-9.

Farhaoui A, Taoussi M, Laasli SE *et al.* Arbuscular mycorrhizal fungi and their role in plant
disease control: A state-of-the-art. *Microbe (Netherlands)* 2025;**8**, DOI:
10.1016/j.microb.2025.100438.

Fiorilli V, Catoni M, Francia D *et al.* The arbuscular mycorrhizal symbiosis reduces disease
severity in tomato plants infected by Botrytis cinerea. *Journal of Plant Pathology*
2011;**93**, DOI: 10.1400/16910

Fiorilli V, Vannini C, Ortolani F *et al.* Omics approaches revealed how arbuscular mycorrhizal
symbiosis enhances yield and resistance to leaf pathogen in wheat. *Sci Rep* 2018;**8**,
DOI: 10.1038/s41598-018-27622-8.

Fujita M, Kusajima M, Fukagawa M *et al.* Response of tomatoes primed by mycorrhizal
colonization to virulent and avirulent bacterial pathogens. *Sci Rep* 2022;**12**, DOI:
10.1038/s41598-022-08395-7.

Gallou A, Declerck S, Cranenbrouck S. Transcriptional regulation of defence genes and
involvement of the WRKY transcription factor in arbuscular mycorrhizal potato root
colonization. *Funct Integr Genomics* 2012;**12**, 10.1007/s10142-011-0241-4.

Gao Y, Huang S, Wang Y *et al.* Analysis of the molecular and biochemical mechanisms
involved in the symbiotic relationship between Arbuscular mycorrhiza fungi and Manihot
esculenta Crantz. *Front Plant Sci* 2023;**14**, DOI: 10.3389/fpls.2023.1130924.

Gebremeskel K, Birhane E, Habtu S *et al.* Arbuscular mycorrhizal fungi improve
morphological and yield performance of Eragrostis tef genotypes in Tigray, Ethiopia. *Sci*
Rep 2024;**14**, DOI: 10.1038/s41598-024-79628-0.

Gong M, Bai N, Wang P *et al.* Co-Inoculation with Arbuscular Mycorrhizal Fungi and Dark
Septate Endophytes under Drought Stress: Synergistic or Competitive Effects on Maize
Growth, Photosynthesis, Root Hydraulic Properties and Aquaporins? *Plants* 2023;**12**,
DOI: 10.3390/plants12142596.

Guzmán P. The prolific ATL family of RING-H2 ubiquitin ligases. *Plant Signal Behav* 2012;**7**,
DOI: 10.4161/psb.20851.

Helgason T, Daniell TJ, Husband R *et al.* Ploughing up the wood-wide web? *Nature*
1998;**394**, DOI: 10.1038/28764.

Hewins CR, Carrino-Kyker SR, Burke DJ. Seasonal variation in mycorrhizal fungi colonizing
roots of Allium tricoccum (wild leek) in a mature mixed hardwood forest. *Mycorrhiza*
2015;**25**, DOI: 10.1007/s00572-015-0628-5.

Hong JJ, Park YS, Bravo A *et al.* Diversity of morphology and function in arbuscular
mycorrhizal symbioses in Brachypodium distachyon. *Planta* 2012;**236**, DOI:
10.1007/s00425-012-1677-z.

Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models.
Biometrical Journal 2008;**50**, DOI: 10.1002/bimj.200810425.

Hua W, Zhu J, Shang Y *et al.* Identification of Suitable Reference Genes for Barley Gene
Expression Under Abiotic Stresses and Hormonal Treatments. *Plant Mol Biol Report*
2015;**33**, DOI: 10.1007/s11105-014-0807-0.

Huot B, Yao J, Montgomery BL *et al.* Growth-defense tradeoffs in plants: A balancing act to
optimize fitness. *Mol Plant* 2014;**7**, DOI: 10.1093/mp/ssu049.

Javed T, Gao SJ. WRKY transcription factors in plant defense. *Trends in Genetics* 2023;**39**,
DOI: 10.1016/j.tig.2023.07.001.

Jung SC, Martinez-Medina A, Lopez-Raez JA *et al.* Mycorrhiza-Induced Resistance and
Priming of Plant Defenses. *J Chem Ecol* 2012;**38**, DOI: 10.1007/s10886-012-0134-6.

Kainat, Mujtaba M, Wang Y *et al.* Effectors of plants pathogenic fungi and fungal like
microbes: a comprehensive review on mechanisms, roles, and host interactions. *Front
Plant Sci* 2025;**16**, DOI: 10.3389/fpls.2025.1626960.

Kim D, Paggi JM, Park C *et al.* Graph-based genome alignment and genotyping with HISAT2
and HISAT-genotype. *Nat Biotechnol* 2019;**37**, DOI: 10.1038/s41587-019-0201-4.

Koch AM, Antunes PM, Maherali H *et al.* Evolutionary asymmetry in the arbuscular
mycorrhizal symbiosis: conservatism in fungal morphology does not predict host plant
growth. *New Phytologist* 2017;**214**, DOI: 10.1111/nph.14465.

Lee J, Lee S, Young JPW. Improved PCR primers for the detection and identification of
arbuscular mycorrhizal fungi. *FEMS Microbiol Ecol* 2008;**65**, DOI: 10.1111/j.1574-
6941.2008.00531.x.

Li S, Khoso MA, Wu J *et al.* Exploring the mechanisms of WRKY transcription factors and
regulated pathways in response to abiotic stress. *Plant Stress* 2024;**12**, DOI:
10.1016/j.stress.2024.100429.

Liu WS, Wei YX, Deng PP *et al.* Conservation tillage increases surface soil organic carbon
stock by altering fungal communities and enzyme activity. *Environmental Science and
Pollution Research* 2023, DOI: 10.1007/s11356-023-28062-2.

Lutz S, Bodenhausen N, Hess J *et al.* Soil microbiome indicators can predict crop growth
response to large-scale inoculation with arbuscular mycorrhizal fungi. *Nat Microbiol*
2023;**8**, 10.1038/s41564-023-01520-w.

Maekawa S, Sato T, Asada Y *et al.* The Arabidopsis ubiquitin ligases ATL31 and ATL6 control
the defense response as well as the carbon/nitrogen response. *Plant Mol Biol* 2012;**79**,
DOI: 10.1007/s11103-012-9907-0.

Marrassini V, Ercoli L, Kuramae EE *et al.* Arbuscular mycorrhizal fungi originated from soils
with a fertility gradient highlight a strong intraspecies functional variability. *Applied Soil
Ecology* 2024;**197**, DOI: 10.1016/j.apsoil.2024.105344.

Martín-Cardoso H, Castillo L, Busturia I *et al.* Arbuscular Mycorrhizal Fungi Increase Blast
Resistance and Grain Yield in Japonica Rice Cultivars in Flooded Fields. *Rice* 2025;**18**,
DOI: 10.1186/s12284-025-00805-4.

Munkvold L, Kjøller R, Vestberg M *et al.* High functional diversity within species of arbuscular
mycorrhizal fungi. *New Phytologist* 2004;**164**, DOI: 10.1111/j.1469-8137.2004.01169.x.

Mustafa G, Khong NG, Tisserant B *et al.* Defence mechanisms associated with mycorrhiza-
induced resistance in wheat against powdery mildew. *Functional Plant Biology* 2017;**44**,
DOI: 10.1071/FP16206.

Pellegrino E, Öpik M, Bonari E *et al.* Responses of wheat to arbuscular mycorrhizal fungi: A
meta-analysis of field studies from 1975 to 2013. *Soil Biol Biochem* 2015;**84**, DOI:
10.1016/j.soilbio.2015.02.020.

Peña Venegas RA, Lee SJ, Thuita M *et al.* The Phosphate Inhibition Paradigm: Host and
Fungal Genotypes Determine Arbuscular Mycorrhizal Fungal Colonization and
Responsiveness to Inoculation in Cassava With Increasing Phosphorus Supply. *Front
Plant Sci* 2021;**12**, DOI: 10.3389/fpls.2021.693037.

Pieterse CMJ, Zamioudis C, Berendsen RL *et al.* Induced systemic resistance by beneficial
microbes. *Annu Rev Phytopathol* 2014;**52**, DOI: 10.1146/annurev-phyto-082712-
102340.

Polesani M, Bortesi L, Ferrarini A *et al.* General and species-specific transcriptional
responses to downy mildew infection in a susceptible (Vitis vinifera) and a resistant (V.
Riparia) grapevine species. *BMC Genomics* 2010;**11**, DOI: 10.1186/1471-2164-11-117.

Pozo MJ, Azcón-Aguilar C. Unraveling mycorrhiza-induced resistance. *Curr Opin Plant Biol*
2007;**10**, DOI: 10.1016/j.pbi.2007.05.004.

Pozo MJ, Cordier C, Dumas-Gaudot E *et al.* Localized versus systemic effect of arbuscular
mycorrhizal fungi on defence responses to Phytophthora infection in tomato plants. *J
Exp Bot* 2002;**53**, DOI: 10.1093/jexbot/53.368.525.

Putri GH, Anders S, Pyl PT *et al.* Analysing high-throughput sequencing data in Python with
HTSeq 2.0. *Bioinformatics* 2022;**38**, DOI: 10.1093/bioinformatics/btac166.

R Core Team. R: A Language and Environment for Statistical Computing. 2025.

Raudvere U, Kolberg L, Kuzmin I *et al.* G:Profiler: A web server for functional enrichment
analysis and conversions of gene lists (2019 update). *Nucleic Acids Res* 2019;**47**, DOI:
10.1093/nar/gkz369.

Ristaino JB, Anderson PK, Bebber DP *et al.* The persistent threat of emerging plant disease
pandemics to global food security. *Proc Natl Acad Sci U S A* 2021;**118**, DOI:
10.1073/pnas.2022239118.

Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis
of RNA-seq data. *Genome Biol* 2010;**11**, DOI: 10.1186/gb-2010-11-3-r25.

Roelfs AP, Singh RP, Saari EE. *Rust Diseases of Wheat: Concepts and Methods of Disease
Management.*, 1992.

RStudio Team. RStudio: Integrated Development Environment for R. 2025.

Russell V. Lenth and Julia Piaskowski. emmeans: Estimated Marginal Means, aka Least-
Squares Means. 2025, DOI: 10.32614/CRAN.package.emmeans.

Sanmartín N, Pastor V, Pastor-Fernández J *et al.* Role and mechanisms of callose priming in
mycorrhiza-induced resistance. *J Exp Bot* 2021;**71**, DOI: 10.1093/JXB/ERAA030.

Skelly MJ, Frungillo L, Spoel SH. Transcriptional regulation by complex interplay between
post-translational modifications. *Curr Opin Plant Biol* 2016;**33**, DOI:
10.1016/j.pbi.2016.07.004.

Skelly MJ, Furniss JJ, Grey HL *et al.* Dynamic ubiquitination determines transcriptional
activity of the plant immune coactivator NPR1. *Elife* 2019;**8**, DOI: 10.7554/eLife.47005.

Smith SE, Smith FA. Roles of arbuscular mycorrhizas in plant nutrition and growth: New paradigms from cellular to ecosystem scales. *Annu Rev Plant Biol* 2011;**62**, DOI: 10.1146/annurev-arplant-042110-103846.

Song Y, Chen D, Lu K *et al.* Enhanced tomato disease resistance primed by arbuscular mycorrhizal fungus. *Front Plant Sci* 2015;**6**, DOI: 10.3389/fpls.2015.00786.

Trujillo M. News from the PUB: Plant U-box type E3 ubiquitin ligases. *J Exp Bot* 2018;**69**, DOI: 10.1093/jxb/erx411.

Umer M, Anwar N, Mubeen M *et al.* Roles of arbuscular mycorrhizal fungi in plant growth and disease management for sustainable agriculture. *Front Microbiol* 2025;**16**, DOI: 10.3389/fmicb.2025.1616273.

de Vega D, Newton AC, Sadanandom A. Post-translational modifications in priming the plant immune system: ripe for exploitation? *FEBS Lett* 2018;**592**, DOI: 10.1002/1873-3468.13076.

Verwoerd TC, Dekker BMM, Hoekema A. A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Res* 1989;**17**, DOI: 10.1093/nar/17.6.2362.

Vierheilig H, Coughlan AP, Wyss U *et al.* Ink and vinegar, a simple staining technique for arbuscular-mycorrhizal fungi. *Appl Environ Microbiol* 1998;**64**:5004–7.

Wang B, Qiu YL. Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 2006;**16**, DOI: 10.1007/s00572-005-0033-6.

Wang M, Tang W, Xiang L *et al.* Involvement of MdWRKY40 in the defense of mycorrhizal apple against fusarium solani. *BMC Plant Biol* 2022;**22**, DOI: 10.1186/s12870-022-03753-z.

Weinberger N V., Cibils-Stewart X, Brien C *et al.* Plant phenotyping and root-associated metabolomics reveal insights into pathogen protection by diverse arbuscular mycorrhizal fungi. *Plants People Planet* 2025;**7**, DOI: 10.1002/ppp3.10627.

West JS, Townsend JA, Stevens M *et al.* Comparative biology of different plant pathogens to estimate effects of climate change on crop diseases in Europe. *Eur J Plant Pathol* 2012;**133**, DOI: 10.1007/s10658-011-9932-x.

Wickham H, Averick M, Bryan J *et al.* Welcome to the Tidyverse. *J Open Source Softw* 2019;**4**, DOI: 10.21105/joss.01686.

Withers J, Dong X. Posttranslational Modifications of NPR1: A Single Protein Playing Multiple Roles in Plant Immunity and Physiology. *PLoS Pathog* 2016;**12**, DOI: 10.1371/journal.ppat.1005707.

Wu S, Shi Z, Chen X *et al.* Arbuscular mycorrhizal fungi increase crop yields by improving biomass under rainfed condition: A meta-analysis. *PeerJ* 2022;**10**, DOI: 10.7717/peerj.12861.

Wu X, Lin T, Zhou X *et al.* Potato E3 ubiquitin ligase StRFP1 positively regulates late blight resistance by degrading sugar transporters StSWEET10c and StSWEET11. *New Phytologist* 2024;**243**, DOI: 10.1111/nph.19848.

Zierold U, Scholz U, Schweizer P. Transcriptome analysis of mlo-mediated resistance in the epidermis of barley. *Mol Plant Pathol* 2005;**6**, DOI: 10.1111/J.1364-3703.2005.00271.X.

Figure Legends

Figure 1 Overview of the experimental setup showing (A) the treatment setup, (B) the AMF
colonisation phase when pots were thinned to a single stem, (C) *Puccinia hordei* inoculation,
(D) the final harvest, including measurements of plant growth, AMF colonisation and qPCR-
based quantification of AMF, Puccinia and defence gene expression (n = 6), (E)
transcriptional and post-translational analyses performed on a subset of plants (n = 3) with
barley leaf transcriptomics and ubiquitination assay and (F) the experimental timeline. Figure
adapted from Weinberger et al. (2025).

Figure 2 Endpoint measurements of AMF colonisation, *Puccinia* infection, and plant biomass
after one week of *Puccinia hordei* infection. (A) Photograph of barley roots stained to show
AMF colonisation, highlighting vesicles – V, intraradical hyphae – IH, and arbuscules – A. (B)
Relative AMF DNA in barley roots, quantified as AMF copy number normalised to *HvEMF1α*,
five weeks post-inoculation. (C) Relative abundance of *Puccinia* in barley leaves, quantified
as *PhEF2* copy number normalized to *HvACT*, one-week post-infection. (D) Total dry
biomass of shoots and roots after five weeks of growth. Bars sharing different letters indicate
significant differences (p < 0.05). C, control; A, AMF addition; P, *Puccinia* addition; AP, AMF
plus *Puccinia* addition (mean ± SE, n = 6).

Figure 3 Relative expression of barley immune marker genes (A) *PR1*, (B) *PR2*, (C) *PR3*,
(D) *WRKY28*. Gene expression is relative to barley actin (HvACT); Bars sharing different
letters indicate significant differences (P < 0.05). C, control; A, AMF addition; P, *Puccinia*
addition; AP, AMF plus *Puccinia* addition (mean ± SE, n = 6).

Figure 4 Transcriptional responses to AMF colonisation and *Puccinia* infection in barley
leaves (n = 3). (A) Differentially expressed genes between control and AMF-colonised plants
(Control vs AMF; $\log_2$FC > 1, FDR < 0.05, logCPM > 2). (B) GO enrichment of AMF-
responsive genes shown in (A), using g:Profiler with Benjamini-Hochberg correction
(adjusted P < 0.05, term size > 2). (C) Heatmap of row-wise z-scored $\log_2$-transformed
expression [$\log_2(x + 1)$] for genes differentially expressed across treatments (C, control; A,
AMF addition; P, Puccinia addition; AP, AMF and Puccinia), clustered into seven co-
expression groups. (D) GO enrichment of gene clusters shown in (C), using g:Profiler with
g:SCS correction (adjusted P < 0.05).

Figure 4 AMF modulation of WRKY transcription factor expression and calcium signalling in
barley leaves (n = 3). (A) Row-scaled expression of differentially expressed WRKY genes
across control (C), AMF (A), Puccinia (P), and combined (AP) treatments ($\log_2$FC > 1, FDR
< 0.05, logCPM > 2), annotated by WRKY class and stress-response role. (B) GO
enrichment of differentially expressed genes containing predicted WRKY-binding motifs
within 500 bp upstream promoter regions (FIMO, P < 0.001). (C) Row-scaled expression of
differentially expressed calcium signalling genes across the same treatments ($\log_2$FC > 1,
FDR < 0.05, logCPM > 2). Calmodulin and calmodulin-like genes were annotated based on
(Cai et al. 2022); calcium-dependent protein kinase (*CDPK*) and *CDPK*-like genes were
annotated based on their closest Arabidopsis homologues.

Figure 5 Effects of AMF and *Puccinia hordei* on ubiquitination. (A) Ubiquitinated protein
levels normalised to actin and expressed relative to control (mean ± SE, n = 3); letters
denote Tukey HSD groupings (P < 0.05) and ANOVA P-values derive from a two-way
ANOVA on $\log_2$-transformed fold-change values. (B) Row-scaled expression of differentially

expressed ubiquitin-related genes across treatments, annotated by ubiquitination function
and pathway type. Genes are annotated by function and ubiquitin pathway type, with PUB
genes classified according to Ryu et al. (2019) and closest Arabidopsis homologues.
Treatments are ordered as C (control), P (*P. hordei*), A (AMF), and AP (AMF + *P. hordei*) for
consistency with the Western Blot.

(A)

V

IH

A

100 µm

Control
AMF
Puccinia
Both

(B)

AMF abundance

a   a

b   b

A   AP   C   P

A: p = 5.37e-05***
P: p = 0.545
AxP: p = 0.856

(C)

Puccinia abundance

30
20
10
0

a

a

b   b

A   AP   C   P

A: p = 0.275
P: p = 3.99e-05***
AxP: p = 0.275

(D)

Plant biomass (g)

3
2
1
0

ab   b

a

b

A   AP   C   P

A: p = 0.647
P: p = 0.000955***
AxP: p = 0.302

(A)

Puccinia — + — +
AMF — — + +

250
150
100
75

α Ubiquitin

α actin

A: p = 0.139
P: p = 0.001**
AxP: p = 0.897

Ubiquitin/Actin
(fold change vs Control)

C   P   A   AP

(B)

Type
ATG
RBR
RING
RING-H2
U-box
UBA
UBC
UEV

Function
E1
E2
E3

HvPUB36
HvPUB69
ATL31
ATL11
ARI7
ATL6-1
UBC26
ATL64
UBA1
HvPUB56
ATL57
HvPUB06
ATL2
HvPUB55
HvPUB68
RGLG2
HvPUB27
HvPUB29
ATL1-1
RBR-2
ATL6-2
ATL4
ATL60
HvPUB33
HvPUB07
HvPUB16
ATL5-1
ATL9
HvPUB48
HvPUB24
HvPUB23
BRIZ1
HvPUB74
ATL43
ATL74
RMA
ELC
HvPUB37
HvPUB18
ATL6-3
HvPUB72
RGLG1
ATL44
RHC1A
ATG3
ATL7
BB
ATL3
RHA1B
UBC28
HvPUB73
ATL6-3
HvPUB02
RDUF2
ATL66
ELC5
BHF2A
MBR2
HvPUB21
RMA3
ATL5
HvPUB50
HvPUB40
HvPUB71
AIP2
HvPUB28
RHC2A
ATL32
ATL4
MPSR1
MUP9

C   P   A   AP