Longitudinal Phylogenetic Inference of Copy Number Alterations and Single Nucleotide Variants from Single-Cell Sequencing

Ethan Kulman[1,2], Quaid Morris[2,*], and Rui Kuang[1,*]

[1] Department of Computer Science and Engineering, University of Minnesota, MN 55455

[2] Computational and Systems Biology Program, Sloan Kettering Institute, NY 10065

Abstract. Longitudinal phylogenetic reconstruction reveals how cancers evolve over time and respond to treatments. Advances in targeted single-cell sequencing, combined with longitudinal sampling, now enable detailed longitudinal tracking of single nucleotide variants (SNVs) and copy number alterations (CNAs) at single-cell resolution. While some phylogenetic inference methods can jointly reconstruct SNV and CNA evolution, none are designed for longitudinal targeted single-cell data. Here, we present LoPhy, a new sequential tree-building algorithm to reconstruct longitudinal phylogenies of SNVs and CNAs from targeted single-cell sequencing. LoPhy maximizes a novel factorized longitudinal tree reconstruction objective by incrementally growing the tree, adding SNVs and CNAs in the order they are observed across samples. Applied to a cohort of 15 acute myeloid leukemias (AMLs) and 4 TP53-mutated AMLs, LoPhy produced phylogenies that are biologically and temporally consistent with clinical observations, with many inferred CNAs validated by orthogonal bulk sequencing from the same patient. These reconstructions further reveal that AML clones selected after therapy are often defined by large-scale CNAs as well as SNVs, highlighting the role of CNAs in disease progression and resistance. More broadly, LoPhy can help uncover how SNVs and CNAs jointly shape the evolutionary trajectories of individual cancers at single-cell resolution. The LoPhy source code is available under a CC-BY-ND license at https://github.com/ethanumn/LoPhy.

Keywords: cancer phylogeny · longitudinal sequencing · cancer evolution.

---

[1] *These authors are co-senior authors

1   Introduction

Cancer evolves through the accumulation of somatic mutations, most commonly single nucleotide variants (SNVs) and copy number alterations (CNAs) [4, 12, 18]. This evolutionary process gives rise to genetically distinct populations of cancer cells, known as *clones*, each defined by a unique genome. The clonal composition of a cancer can change over time due to selective pressures such as immune response or therapy. Longitudinal sequencing, which involves sampling and sequencing cancer cells from the same patient at multiple time points, enables researchers to track these evolutionary dynamics and gain insight into how cancers adapt and progress. Recent advances in targeted single-cell DNA sequencing—particularly high-throughput microfluidic protocols known as *single-cell amplicon sequencing*—enable the detection of both SNVs and CNAs at single-cell resolution [10, 13]. Several studies have applied this technology to longitudinal cancer samples [10, 16], sequencing portions of oncogenes at ~40x per amplicon per cell. However, no computational methods currently exist to jointly reconstruct the longitudinal evolution of both SNVs and CNAs from this data—a gap this work addresses.

The evolutionary history of an individual cancer is typically represented as a phylogenetic tree, or *clone tree*, which depicts the ancestral relationships among cancer clones. Existing methods that reconstruct clone trees from longitudinal cancer samples focus exclusively on either SNVs [1, 6, 11] or CNAs [9], but not both. This largely reflects the limitations of previously available sequencing technologies. For example, single-cell whole-genome sequencing (scWGS) provides genome-wide coverage, enabling the detection of large-scale CNAs, but lacks the depth needed to reliably call SNVs. In contrast, older targeted single-cell sequencing technologies have sufficient depth to call SNVs, but too low of coverage to detect CNAs. Bulk DNA sequencing is useful for identifying SNVs and CNAs at the population level but lacks the resolution to detect changes in smaller subpopulations.

Two algorithms—BiTSC^2 [2] and COMPASS [14]—can jointly reconstruct clone trees containing both SNVs and CNAs from single-cell amplicon sequencing data, but neither is designed for longitudinal analysis. Applying them to longitudinal samples either requires reconstructing clone trees separately for each time point—which can yield disjoint evolutionary histories lacking temporal coherence—or pooling all samples together, which ignores sample-specific technical variation such as dropout rates and region-specific coverage. Failing to model sample-specific technical variation can lead to incorrect clone trees—clones may appear in samples where they were absent, or CNAs may be inferred from variations in coverage between samples, rather than genuine biological changes.

Here, we introduce LoPhy, an algorithm that reconstructs clone trees describing the joint evolution of SNVs and CNAs from longitudinal single-cell amplicon sequencing. The method relies on a factorized longitudinal tree reconstruction objective—a novel formulation derived in this work. To optimize this objective, LoPhy uses a sequential tree-building strategy in which mutations are added to the tree in the order they are first detected across longitudinal samples, ensuring that mutations detected earlier in time are placed earlier in evolution. Given S longitudinal samples, LoPhy generates a sequence of trees, $\mathcal{T}^1, \mathcal{T}^2, \ldots, \mathcal{T}^S$, where each tree $\mathcal{T}^s$ describes the evolutionary history up to time point s and forms a subtree of all subsequent trees. Each subtree is refined through hill climbing, which allows SNVs to be relocated and CNAs to be added or removed to maximize tree likelihood. This design naturally enforces longitudinal constraints while remaining computationally efficient, scaling to tens or even hundreds of thousands of cells. Structuring inference in this way simplifies the estimation of sample-specific technical parameters—such as dropout rates and region-specific coverage—making CNA inference more reliable. Using both simulations and real data, we show that LoPhy reconstructs temporally consistent and biologically plausible longitudinal clone trees describing the joint evolution of SNVs and CNAs.

We apply LoPhy to published longitudinal single-cell datasets comprising acute myeloid leukemia (AML) samples profiled with the Tapestri® platform (Mission Bio, Inc.), including a cohort of 15 AMLs undergoing diverse treatments and four TP53-mutated AMLs sampled before and after venetoclax therapy. The inferred clone trees reveal a coherent picture of how these cancers evolved over time, with many CNAs identified by LoPhy being orthogonally validated with bulk sequencing data from the same cancer. These longitudinal reconstructions reveal that AML clones emerging after therapy or at relapse are frequently defined by both SNVs and CNAs. While CNAs are well established drivers of oncogenesis, our results suggest that modeling them alongside SNVs is essential for fully characterizing key clonal populations.

Fig. 1. Overview of LoPhy. **a.** Four types of somatic events are modeled. **b.** Longitudinal reconstruction: LoPhy infers somatic events from longitudinal single-cell amplicon sequencing to build subtrees $\mathcal{T}^1, \mathcal{T}^2, \mathcal{T}^3$, representing evolution up to time of sampling. **c.** Inputs from longitudinal targeted single-cell sequencing include total coverage of targeted regions and locus-level alternative/reference read counts.

2   Materials and Methods

LoPhy reconstructs longitudinal clone trees from single-cell amplicon sequencing data collected at multiple time points from the same cancer. An overview of the method is provided in Figure 1. This section details the algorithm's main components: required sequencing inputs, clone tree representation, evolutionary model, likelihood model, and optimization strategy. Central to LoPhy is a novel factorized longitudinal likelihood, which extends concepts from our previous work [7].

2.1   Inputs from longitudinal single-cell amplicon sequencing

Single-cell amplicon sequencing focuses on small genomic regions (150-450 base pairs) within disease-related genes. Amplicons—generated by amplifying the reverse transcription products of these regions—are used to measure the frequency of the target regions in the genome and identify variants. For each cell j, the data from a single-cell amplicon sequencing experiment is summarized as follows: D_j is the total number of reads mapped across all targeted regions, D_{kj} is the number of reads mapped to region k, D_{ij} is the total number of reads mapped to locus i, and A_{ij} is the number of alternative reads observed at locus i. Each cell is associated with a specific longitudinal sample, allowing the data to be partitioned into sample-specific

subsets. We use $\mathbf{D}$ and $\mathbf{A}$ to denote the full dataset, containing read counts for all cells across all longitudinal samples. For a specific longitudinal sample s, we use $\mathbf{D}^s$ and $\mathbf{A}^s$ to represent the corresponding subset of data for cells originating from that sample.

2.2   Clone tree representation

Inferring a clone tree that includes both SNVs and CNAs involves reconstructing a rooted, directed tree $\mathcal{T}$, and clone assignment vector σ. Each node v in $\mathcal{T}$ represents a distinct clone, with the root corresponding to the germline (non-cancerous) clone, which may contain germline SNPs but no somatic mutations. The clone assignment vector σ specifies the clone membership of each cell, such that if $\sigma_j = v$, cell j is assigned to clone v. The copy number profile of cell j is determined by its assigned clone v, where $c_k(\sigma_j)$ denotes the total copy number for target region k, and $c_i^{(r)}(\sigma_j)$ and $c_i^{(a)}(\sigma_j)$ denote the number of copies of the reference and alternative alleles, respectively, at locus i.

2.3   Evolutionary model

The LoPhy algorithm models four types of somatic events: SNV acquisition, copy number gains, copy number losses, and copy-neutral loss of heterozygosity (CNLOH). CNAs may affect either individual target regions or entire chromosomes. Each variant locus lies within a target region, and its copy number can change if a CNA impacts the allele containing that region. Because targeted sequencing data are noisy and offer limited resolution for detecting multiple copy number changes along the same lineage, we restrict each region to be impacted by at most one CNA per lineage in the tree. SNV evolution in $\mathcal{T}$ follows the k-Dollo model [3], in which each SNV is acquired exactly once but may be lost at most once per lineage due to a deletion of the genomic region on which it resides. This combined evolutionary model for SNVs and CNAs is identical to that described in [14].

2.4   Longitudinal likelihood model

Given S longitudinal samples, LoPhy assumes the following factorized longitudinal likelihood:

$$\prod_{s=1}^{S} P(\mathbf{D}^s, \mathbf{A}^s \mid \mathcal{T}^s). \tag{1}$$

Each tree $\mathcal{T}^s$ represents the evolutionary history up to time point s and is a subtree of all subsequent trees, $\mathcal{T}^1 \subseteq \mathcal{T}^2 \subseteq \cdots \subseteq \mathcal{T}^S$, reflecting the accumulation of somatic mutations over time. The sample-specific likelihood, adopted from [14], is defined as:

$$
\begin{aligned}
P(\mathbf{D}^s, \mathbf{A}^s \mid \mathcal{T}^s) = \prod_{\text{cell } j} \sum_{\sigma_j} & \pi_{\sigma_j} \\
\times \prod_{\text{region } k} & \underbrace{P\big(D_{kj} \mid c_k(\sigma_j), \rho_k\big)}_{\text{Read depth likelihood}} \\
\times \prod_{\text{locus } i} & \underbrace{P\big(A_{ij} \mid D_{ij}, c_i^{(a)}(\sigma_j), c_i^{(r)}(\sigma_j), \mu_i^{(a)}, \mu_i^{(r)}\big)}_{\text{Allelic read count likelihood}}.
\end{aligned}
\tag{2}
$$

The sample-specific likelihood marginalizes over all possible clonal assignments for σ_j, with π_{σ_j} denoting the prior probability of assignment to clone σ_j. The read depth likelihood, which evaluates the inferred total copy numbers for each target region in each cell, and the allelic read count likelihood, which evaluates the inferred number of alternative and reference allele copies at each locus in each cell, are described below.

The factorized likelihood in Eq. 1 approximates the joint likelihood over the data from all S longitudinal samples:

$$\prod_{s=1}^{S} P(\mathbf{D}^s, \mathbf{A}^s \mid \mathcal{T}^s) \approx P(\mathbf{D}^1, \ldots, \mathbf{D}^S, \mathbf{A}^1, \ldots, \mathbf{A}^S \mid \mathcal{T}), \tag{3}$$

$$\approx P(\mathbf{D}, \mathbf{A} \mid \mathcal{T}).$$

The left and right hand sides of Eq. 3 are equivalent if and only if

$$\mathbf{D}^s, \mathbf{A}^s \perp\!\!\!\perp \mathbf{D}^{s^>}, \mathbf{A}^{s^>} \mid \mathbf{D}^{s^<}, \mathbf{A}^{s^<}, \tag{4}$$

which states that the read count data for sample s is conditionally independent of all data from future samples, $s^>$, given the data from all prior samples, $s^<$. This conditional independence assumption reflects

the idea that longitudinal sequencing captures cancer evolution over time. Genomic changes arising between two time points should be detectable, provided that cells carrying those mutations are sampled.

Read depth likelihood Current single-cell amplicon sequencing technologies have non-uniform coverage across amplicons [14]. Since each amplicon covers a portion of a disease-specific gene (i.e., a region), the probability a read falls into region k, denoted with ρ_k, is impacted by a variety of factors including the size of the region, its GC content, etc. We estimate ρ_k using the observed fraction of reads that fall into region k in a population of normal (i.e., absent of CNAs) cells. For a cell j with D_j total reads, assigned to clone σ_j, and copy number $c_l(\sigma_j)$ for region l, the expected read depth in region k is given by

$$\mathbb{E}(D_{kj}) = D_j \frac{c_k(\sigma_j)\rho_k}{\sum_l c_l(\sigma_j)\rho_l}. \tag{5}$$

Cells assigned to the same clone may have larger than expected variances in their regional read depths, a phenomenon known as *overdispersion*. To model this, we use a Negative Binomial distribution for each clone's regional read counts, parameterized by the mean $\mu = \mathbb{E}(D_{kj})$ and inverse-dispersion parameter θ:

$$P\left(D_{kj}\big|c_k(\sigma_j),\rho_k,\theta\right) = \frac{\Gamma(D_{kj}+\theta)}{\Gamma(D_{kj}+1)\Gamma(\theta)}\left(\frac{\theta}{\theta+\mu}\right)^{\theta}\left(\frac{\mu}{\theta+\mu}\right)^{D_{kj}}. \tag{6}$$

This parametrization, often referred to as the *overdispersed Poisson*, differs from the standard Poisson in that its variance is $\mu + \frac{\mu^2}{\theta}$, while the standard Poisson has variance equal to μ. Following [14], we set $\theta = 6.0$ to account for overdispersion; setting $\theta = \infty$ assumes no overdispersion and reduces it to the Poisson. We omit θ from the read depth likelihood in Eq. 2, since it is treated as fixed.

Allelic read count likelihood Although single-cell amplicon sequencing has low coverage, allelic imbalances resulting from CNAs can still be detected from shifts in the observed proportions of alternative and reference read counts. However, inferring the relative proportions of alternative and reference alleles from these read counts is complicated by overdispersion, which arises from allelic biases during amplification. To address this, the allelic read counts are modeled with a beta-binomial distribution. Using this model, the likelihood of observing A alternative reads out of D total reads at a position, given $c^{(a)}$ alternative alleles and $c^{(r)}$ reference alleles, is:

$$P(A \mid D, c^{(a)}, c^{(r)}, \omega, \epsilon) = \frac{\text{Beta}\left(A+\omega f, D-A+\omega(1-f)\right)}{\text{Beta}\left(\omega f, \omega(1-f)\right)}, \tag{7}$$

where $\omega > 0$ is the beta-binomial concentration parameter, $\text{Beta}(\cdot)$ denotes the Beta function, and f is the expected fraction of alternative reads assuming a fixed sequencing error rate of ϵ:

$$f = (1-\epsilon)\frac{c^{(a)}}{c^{(a)}+c^{(r)}} + \epsilon\frac{c^{(r)}}{c^{(a)}+c^{(r)}}.$$

Allelic read counts are also impacted by allele-specific dropout during sequencing, leading to false negatives. To account for this, we introduce global allele-specific dropout rates $\mu_i^{(a)}$ and $\mu_i^{(r)}$, representing the dropout probabilities of the alternative and reference alleles at locus i. The number of amplified alleles can be modeled using a binomial distribution. Specifically, the likelihood of amplifying l alternative and k reference alleles at locus i, given $c_i^{(a)}$ alternative and $c_i^{(r)}$ reference alleles copies, is:

$$P(l, k \mid c_i^{(a)}, c_i^{(r)}, \mu_i^{(a)}, \mu_i^{(r)}) = \text{Binom}(l \mid c_i^{(a)}, 1-\mu_i^{(a)}) \times \text{Binom}(k \mid c_i^{(r)}, 1-\mu_i^{(r)}), \tag{8}$$

where $\text{Binom}(k \mid n, p) = \binom{n}{k}p^k(1-p)^{n-k}$ is the binomial likelihood of observing k successes out of n trials with a success rate of p. To fully account for dropout uncertainty, the likelihood of the observed read count A is obtained by summing over all possible dropout scenarios:

$$P(A \mid D, c_i^{(a)}, c_i^{(r)}, \mu_i^{(a)}, \mu_i^{(r)}, \omega, \epsilon) = \sum_{\substack{0 \le l \le c_i^{(a)} \\ 0 \le k \le c_i^{(r)} \\ (l,k)\ne(0,0)}} P(l, k \mid c_i^{(a)}, c_i^{(r)}, \mu_i^{(a)}, \mu_i^{(r)})P(A \mid D, c_i^{(a)}, c_i^{(r)}, \omega, \epsilon). \tag{9}$$

For succinctness, we omit the fixed parameters ω, ϵ, defining the shorthand used in Eq. 2:

$$P(A \mid D, c_i^{(a)}, c_i^{(r)}, \mu_i^{(a)}, \mu_i^{(r)}, \omega, \epsilon) \equiv P(A \mid D, c_i^{(a)}, c_i^{(r)}, \mu_i^{(a)}, \mu_i^{(r)}). \tag{10}$$

2.5   Tree prior

Single-cell sequencing data are noisy, and complex evolutionary models can overfit the data by introducing spurious clones or CNAs. To mitigate this, we impose a prior on the number of clones and CNAs in the inferred tree.

Each CNA event incurs a penalty, with CNLOH assigned a stronger penalty than a gain or loss. CNAs affecting contiguous regions are counted as a single event, as large-scale alterations (e.g., whole-chromosome changes) are common. We also penalize the total number of clones, since incorporating more SNVs and CNAs into the tree increases the chance that multiple mutations originated in the same clone. Both penalties scale linearly with the number of cells, as larger datasets exert a proportionally greater influence on the likelihood. Let p_1 denote the penalty for the number of clones and p_2 the penalty for CNAs. The prior is defined as:

$$\log(P(\mathcal{T})) = -p_1 n_{\text{muts}} n_{\text{nodes}} - p_2 (n_{\text{gains}} + n_{\text{losses}} + 2 n_{\text{CNLOH}}) + \text{Constant}, \tag{11}$$

where n_{muts}, n_{nodes}, n_{gains}, n_{losses}, and n_{CNLOH} denote, respectively, the number of mutations, clones, non-contiguous gains, non-contiguous losses, and non-contiguous CNLOH events.

2.6 Algorithm

The LoPhy algorithm combines sequential tree-building with hill climbing to generate a sequence of clone trees $\mathcal{T}^1, \ldots, \mathcal{T}^S$ that maximize the factorized posterior:

$$\prod_{s=1}^{S} P(\mathbf{D}^s, \mathbf{A}^s \mid \mathcal{T}^s) P(\mathcal{T}^s). \tag{12}$$

Each factor in the posterior is the product of the prior (Eq. 11) and sample-specific likelihood (Eq. 2). The algorithm consists of three stages.

Hyperparameters Normal (non-cancerous) cell populations are first identified in each sample to estimate region-specific coverage probabilities ρ_k for each region k. LoPhy does this by independently reconstructing an SNV-only tree for each longitudinal sample, maximizing the sample-specific likelihood in Eq. 2, and designating cells assigned to the root node as the normal population. Finally, the sample in which each SNV is first detected is determined from the single-cell sequencing inputs.

Tree initialization The tree $\mathcal{T}^0$ is initialized with a root node that represents a normal cell population, which may have germline SNPs, but no somatic mutations.

Hill climbing For each sample s, SNVs first observed in that sample are incorporated into $\mathcal{T}^{s-1}$ by forming new clones that descend from existing ones. After introducing new SNVs, hill climbing proceeds for η epochs. At each iteration, either a newly added SNV or a genomic region is selected at random, and all valid modifications are evaluated. Three types of moves are permitted: (1) relocate an SNV, (2) remove a CNA, or (3) add a CNA. CNA moves are constrained so that only regions in lineages not previously altered by a CNA in $\mathcal{T}^{s-1}$ may be modified. All tree modifications respect longitudinal constraints: clones in $\mathcal{T}^{s-1}$ remain fixed, and new SNVs/CNAs are restricted to descendant clones. Tree modifications are accepted if they increase the posterior likelihood (Eq. 12).

2.7 Inferring clone assignment priors and dropout rates

Dropout rates and clone assignment priors are estimated for each sample s using its corresponding tree $\mathcal{T}^s$. We adopt the expectation–maximization (EM) algorithm from [14] to jointly estimate the prior probability of assignment to each node (clone) v, π_v, and the global dropout rates for the alternative and reference alleles at each locus i, $\mu_i^{(a)}$ and $\mu_i^{(r)}$. We place a Beta(α, β) prior on $\mu_i^{(a)}$ and $\mu_i^{(r)}$ centered at 0.05, taking advantage of its conjugacy with the binomial likelihood used to model dropout rates. For clone assignments, we use a flat Dirichlet$(1, \ldots, 1)$ prior, which is conjugate to the categorical distribution for clone assignment prior probabilities. The EM algorithm is run after every tree modification during hill climbing. There are two types of latent variables for each cell j: the clone assignment σ_j, and the number of amplified alternative and reference alleles for each locus i, $C_{ij}^{(a)}$ and $C_{ij}^{(r)}$. Given the current parameter estimates, the E-step is defined as:

$$Q(\sigma_j = v) = P(\sigma_j = v \mid \mathbf{D}_j, \mathbf{A}_j, \boldsymbol{\pi}, \boldsymbol{\mu}^{(a)}, \boldsymbol{\mu}^{(r)}) = \frac{\pi_v P(\mathbf{D}_j, \mathbf{A}_j \mid \sigma_j = v, \boldsymbol{\pi}, \boldsymbol{\mu}^{(a)}, \boldsymbol{\mu}^{(r)})}{\sum_{v'} \pi_{v'} P(\mathbf{D}_j, \mathbf{A}_j \mid \sigma_j = v', \boldsymbol{\pi}, \boldsymbol{\mu}^{(a)}, \boldsymbol{\mu}^{(r)})}, \tag{13}$$

$$Q(C_{ij}^{(a)} = l, C_{ij}^{(r)} = k \mid \sigma_j = v) = P(C_{ij}^{(a)} = l, C_{ij}^{(r)} = k \mid D_{ij}, A_{ij}, \sigma_j = v, \boldsymbol{\mu}^{(a)}, \boldsymbol{\mu}^{(r)})$$

$$= \frac{P(l, k \mid c_i^{(a)}(v), c_i^{(r)}(v), \mu_i^{(a)}, \mu_i^{(r)}) P(A_{ij} \mid D_{ij}, k, l, \omega, \epsilon)}{\sum_{k', l'} P(l', k' \mid c_i^{(a)}(v), c_i^{(r)}(v), \mu_i^{(a)}, \mu_i^{(r)}) P(A_{ij} \mid D_{ij}, k', l', \omega, \epsilon)}.$$

The update equations for the M-step are defined as:

$$\pi_v = \frac{1}{n_{\text{cells}}} \sum_{\text{cell } j} Q(\sigma_j = v) \tag{14}$$

$$\mu_i^{(r)} = \frac{\alpha - 1 + \sum_{\text{node } v} \sum_{\text{cell } j} Q(\sigma_j = v) \sum_{k,l} Q(C_{ij}^{(a)} = l, C_{ij}^{(r)} = k \mid \sigma_j = v)(c_i^{(r)}(v) - k)}{\alpha + \beta - 2 + \sum_{\text{node } v} \sum_{\text{cell } j} Q(\sigma_j = v) c_i^{(r)}(v)}. \tag{15}$$

$$\mu_i^{(a)} = \frac{\alpha - 1 + \sum_{\text{node } v} \sum_{\text{cell } j} Q(\sigma_j = v) \sum_{k,l} Q(C_{ij}^{(a)} = l, C_{ij}^{(r)} = k \mid \sigma_j = v)(c_i^{(a)}(v) - l)}{\alpha + \beta - 2 + \sum_{\text{node } v} \sum_{\text{cell } j} Q(\sigma_j = v) c_i^{(a)}(v)}, \tag{16}$$

3   Results

3.1   Evaluation on simulated longitudinal data

We benchmarked LoPhy on simulated longitudinal cancer data, comparing it to COMPASS [14], LACE [1], and SCITE [5]. COMPASS reconstructs clone trees incorporating both SNVs and CNAs but does not account for longitudinal sampling. LACE is designed for longitudinal data but models only SNVs. SCITE is a widely used method for SNV-based tree reconstruction, though it is not designed for longitudinal data. We also considered BiTSC^2[2], which supports joint reconstruction of SNVs and CNAs, but excluded it due to prohibitively long runtimes, even on small longitudinal datasets.

We simulated longitudinal cancer datasets to assess the benefits of jointly inferring SNVs and CNAs while explicitly modeling longitudinal sampling, compared to methods that either infer only SNVs or are not designed for longitudinal data. We simulated 40 longitudinal cancers in total: 10 for each setting of longitudinal samples (2, 3, 4, or 5). Each dataset spanned 2–5 time points, with 3,000 cells sampled per time point (6,000–15,000 cells per dataset). Sequencing data were generated to resemble output from the Tapestri $^{®}$ platform (see Appendix C), comprising 20 genomic regions, 20 SNVs, and 3 CNAs. Ground-truth longitudinal clone trees included 6–9 clones, with the number of clones increasing in datasets with more time points, reflecting the expectation that more longitudinal sampling reveals greater clonal diversity. The longitudinal dynamics for each simulated cancer were generated using a Moran process [8], a well-established model in population genetics.

Performance was evaluated using cell-to-clone assignment accuracy—comparing the inferred mutant (alternative) and total copy number profiles of each cell to its ground-truth profile. Specifically, we computed the mean absolute error (MAE) between inferred and ground-truth mutant copy numbers (MCN) and between inferred and ground-truth total copy numbers (TCN), denoted MCN-MAE and TCN-MAE, respectively. We also introduce a new metric, the false emergence rate (FER), that measures the fraction of cells assigned copy numbers found exclusively in clones from future samples and absent from all clones detected up to the cell's sampling time. FER captures a key pitfall of ignoring longitudinal structure: mutations may appear to emerge earlier in evolution than they actually did. Detailed descriptions of these metrics are provided in Appendix B.

Figure 2 shows the performance of LoPhy and competing methods on the 40 simulated longitudinal cancer datasets. LoPhy on average achieved the best performance (lowest error) across all evaluation metrics, regardless of the number of longitudinal samples, and showed consistently strong results—particularly for total copy number estimation. In contrast, COMPASS struggled with total copy number inference, with an average TCN-MAE exceeding 0.35 in all simulation settings. This is expected since it does not model coverage variation between longitudinal samples, which confounds CNA inference. For mutant copy numbers, LoPhy achieved the best average performance. While COMPASS and SCITE showed slightly lower MCN-MAE on a subset of datasets, the average difference was small—approximately 6×10^{-3}.

SCITE estimates whether loci are heterozygous or homozygous but does not infer total copy numbers; therefore, we report MCN-MAE for SCITE but exclude it from TCN-MAE comparisons. LACE outputs a binary matrix indicating the presence or absence of SNVs per cell, so we include it in the MCN-MAE comparisons while noting this limitation; it is likewise excluded from TCN-MAE results. SCITE performed well on MCN-MAE, showing lower error than COMPASS on average across all simulation settings. LACE exhibited poor MCN-MAE performance, likely due to its binary output, but performed well on the FER metric, as expected given its longitudinal modeling. LoPhy achieved perfect FER, whereas COMPASS frequently inferred mutant and/or total copy number changes earlier in evolution than they actually occurred.

Fig. 2. Evaluation of LoPhy, COMPASS, SCITE, and LACE on 40 simulated longitudinal cancer datasets. **a-c.** Distributions of evaluation metrics: MCN-MAE (*mutant copy number mean absolute error*), TCN-MAE (*total copy number mean absolute error*), and FER (*false emergence rate*). Box plots indicate the median (center line), first and third quartiles (box edges), range of data (whiskers), and outliers (unfilled circles). Lower values correspond to more accurate reconstructions. **d.** Examples of cell assignments which result in the three different error types.

Finally, LoPhy was substantially faster than COMPASS, completing datasets with 5 longitudinal samples in an average of about 2 minutes, compared to roughly 1 hour and 40 minutes for COMPASS. Runtime comparisons for all methods across the different simulation settings are provided in Appendix D.

3.2   Longitudinal phylogenetic analysis of acute myeloid leukemias

We applied LoPhy to a cohort of 15 longitudinally observed acute myeloid leukemias (AMLs) profiled with the Tapestri® platform [10], each with 2–5 time points sequenced using either a 50-amplicon panel (19 genes) or a 279-amplicon panel (37 genes). The authors in [10] provided clinical descriptions for all of these AMLs, which is summarized in Appendix A, detailing treatments and changes in the clonal composition of the AMLs over time. We demonstrate that our analyses with LoPhy are consistent with these clinical descriptions, and reveal that AML clones dominant after treatment or at relapse frequently harbor both SNVs and CNAs, illustrating the clinical relevance of jointly modeling SNVs and CNAs in a longitudinal framework. Below, we highlight several representative cases, including AML-99 and two AMLs where we directly compare reconstructions from LoPhy and COMPASS. Analyses of the remaining AMLs, including a second cohort of four TP53-mutated AMLs with samples collected before and after venetoclax treatment [16], are provided in Appendix E.

Dominant AML clones are characterized by both SNVs and CNAs AML-99 illustrates how incorporating longitudinal information and jointly modeling SNVs and CNAs yields more biologically meaningful clone trees. This patient was diagnosed with AML carrying an IDH2 R140Q mutation (AML-99-001) and had two additional samples collected during therapy (AML-99-002, AML-99-003). The patient remained in remission until relapse (AML-99-004), after which therapy was adjusted, though the disease remained refractory (AML-99-005).

Figure 3a shows the longitudinal clone tree reconstructed by LoPhy for AML-99, revealing that clones defined by CNAs dominated as the disease evolved. Across the first three samples (AML-99-001 through AML-99-003), the dominant clone carried a MYC/RAD21 gain and CNLOH of the RUNX1 reference allele, consistent with the bulk ASCAT [17] profile for AML-99-001 (Figure 3b) from [14]. At relapse, however, a new

Fig. 3. a. Longitudinal clone tree reconstructed by LoPhy for AML-99. Nodes containing CNAs are shown with bold outline. The time axis indicates sampling points and therapy regimens: Clad, cladribine; LDAC, low-dose cytarabine; Ena, enasidenib; VEN, venetoclax; DAC, decitabine. **b**. ASCAT profile for sample AML-99-001 from [14]. **c**. Fluctuations in size of key clonal populations in response to therapy.

clone emerged with the same MYC/RAD21 gain but CNLOH of the RUNX1 alternative allele, dominating AML-99-004 (Figure 3c). After therapy was adjusted, the relapse clone contracted, while a FLT3-positive subclone of the original MYC/RAD21 clone expanded to dominate AML-99-005 (Figure 3c).

A key insight from this reconstruction is that two independent CNLOH events occurred in the RUNX1 region on separate lineages in the tree: one deleting the reference allele early in the disease, and one deleting the alternative allele that appears only at relapse. At relapse, the lineage with CNLOH of the RUNX1 alternative allele expanded and became dominant, while the earlier lineage carrying CNLOH of the reference allele contracted (Figure 3c). The CNLOH at relapse is well supported by the data—cells in this lineage have

Fig. 4. Longitudinal reconstructions by LoPhy and COMPASS for AML-01 (**a**), and AML-83 (**b**).

median coverage ≥ 30 at the RUNX1 p.R162K locus, yet nearly all have zero (or near-zero) reads supporting the alternative allele. Bulk data for AML-99-004, if available, could independently confirm the CNLOH event at relapse.

We provide COMPASS's reconstruction for AML-99 in Appendix E, which agrees poorly with the clinical description for this patient and infers CNAs which cannot be validated. Across both cohorts—the 15 AMLs from [10] and the 4 TP53-mutated AMLs from [16]—LoPhy reconstructed clone trees where CNA-defined clones dominated in 17 of 19 AMLs. Bulk data exists for 5/17 of these CNA-dominated AMLs, supporting the majority of CNAs inferred by LoPhy.

Comparing and contrasting LoPhy and COMPASS longitudinal AML reconstructions Figure 4 shows longitudinal reconstructions by LoPhy and COMPASS for AML-01 and AML-83. These two cases highlight the importance of longitudinal constraints and estimating sample-specific coverage and dropout rates.

Figure 4a shows the longitudinal clone trees reconstructed by LoPhy and COMPASS for AML-01. This is a FLT3-ITD mutated AML with samples collected at diagnosis (AML-01-001) and relapse (AML-01-002). Treatment was given after diagnosis; however, the relapse sample remained positive for FLT3-ITD [10]. Although both LoPhy and COMPASS infer a CNLOH of the FLT3 reference allele, their reconstructions differ in its placement. LoPhy places the FLT3 CNLOH before the two WT1 mutations, identifying 85% of cells at diagnosis as positive for this event. COMPASS places the CNLOH event after the first WT1 mutation, assigning 86% of cells at diagnosis both WT1 mutations and the CNLOH event—even though no WT1 mutations were detected at diagnosis. This high rate of false emergences occurs because, without WT1 mutation data at diagnosis or longitudinal constraints, COMPASS relies solely on the strong CNLOH signal when assigning cells to clones. This also results in COMPASS estimating unrealistically high dropout rates of 50% for the WT1 loci, highlighting the importance of sample-specific dropout modeling for plausible longitudinal reconstructions.

Figure 4b shows the trees reconstructed by LoPhy and COMPASS for AML-83. This AML is characterized at diagnosis (AML-83-001) by an IDH2 p.R172K mutation. At relapse (AML-83-002), two additional NRAS mutations were detected that were absent at diagnosis [10]. Bulk copy number analysis using CNVkit [15], as reported in [14], supports a copy number gain on chromosome 8 at relapse (Figure 4b top left). LoPhy reconstructs an evolutionary history consistent with the case description in [10], correctly identify-

ing the IDH2 mutation at diagnosis, and the NRAS mutations and MYC/RAD21 gain on chromosome 8 at relapse. The MYC/RAD21 clone dominates at relapse, giving rise to three subclones distinguished by distinct FLT3/NRAS SNVs. In contrast, while COMPASS generally recovers the correct SNV evolution, it misses the MYC/RAD21 gain. This occurs because it does not model sample-specific coverage, so differences between the two sequencing experiments—particularly among the "normal" cells used to define baseline coverage—distort the signal needed to detect the gain. When run only on the relapse sample (AML-83-002), where sample-specific coverage does not need to be modeled, COMPASS reconstructs a tree nearly identical to LoPhy's, including the MYC/RAD21 gain. However, analyzing the two AML-83 samples independently with COMPASS produces inconsistent clone structures across the trees, obscuring the temporal ordering of genomic changes and the relationships between cells from the two samples.

Analyses with LoPhy of both AML cohorts—including the 15-AMLs from [10] and the 4 TP53-mutated AMLs from [16]—provide the first integrated view of how SNVs and CNAs jointly shaped clonal evolution in these AMLs over time. Analyses of the remaining AMLs, presented in Appendix E, mirror the findings for AML-01 and AML-83. COMPASS inferred trees that were inconsistent with longitudinal data or clinical descriptions in 7 of 19 AMLs, whereas LoPhy produced consistent reconstructions in all but one case, highlighting the robustness of its modeling framework. Note that previous analyses of these AMLs were limited either to SNV-only reconstructions or to reconstructing trees independently for each time point. In the original study [10], the 15 longitudinal AMLs were analyzed with SCITE [5] by pooling all time points and inferring clone trees based solely on SNVs; only AML-99 was later reanalyzed with COMPASS [14], which was used to reconstruct trees independently for each time point. The 4 TP53-mutated AMLs from [16] were likewise analyzed with COMPASS in [14], by reconstructing trees independently for each time point.

4   Discussion

Longitudinal single-cell sequencing is increasingly used to study cancer evolution, particularly in leukemias and lymphomas. Methods exist to reconstruct longitudinal clone trees from this data; however, they typically model the evolution of either single-nucleotide variants (SNVs) or copy number alterations (CNAs), but not both. New amplicon-based single-cell DNA sequencing technologies now enable reliable detection of both SNVs and CNAs, allowing detailed longitudinal tracking of these genomic alterations at single-cell resolution. Here, we show that reconstructing longitudinal clone trees from this new single-cell data reveals that dominant leukemia clones are often defined by both SNVs and CNAs, and that these clones frequently expand or contract in response to therapy. These insights are made possible by our algorithm, LoPhy, which optimizes a novel factorized objective to reconstruct clone trees from longitudinal single-cell amplicon sequencing. We show that LoPhy reconstructions are consistent with clinical observations and supported by orthogonal bulk sequencing data from the same cancers. The resulting trees provide continuous joint histories of how SNVs and CNAs emerge and evolve over time within individual cancers. The accuracy of our algorithm is further supported by simulations, where LoPhy consistently outperformed existing reconstruction methods across multiple evaluation metrics.

The current LoPhy algorithm has a few notable limitations. First, it relies on a greedy search strategy, which can become trapped in local maxima. While targeted single-cell sequencing experiments typically profile thousands of cells, making the reconstruction problem highly constrained, sparsely sampled subclones may be excluded from the inferred clone tree under this approach. This issue may be mitigated by increasing the number of cells sampled at each time point. Second, our copy number evolution model is potentially too simplistic: it only allows each region to be impacted by at most one CNA per lineage, models a restricted set of CNA types, and does not account for homozygous loss—a rare but possible event.

Future work could address these limitations. A more comprehensive copy number model that incorporates homozygous loss and multiple CNAs affecting the same region within a lineage is feasible, though these additions may become more relevant as single-cell technologies advance to better capture such events. The search strategy could be improved by replacing the greedy approach with a Markov Chain Monte Carlo framework, reducing the risk of becoming trapped in local maxima. However, this would require a carefully designed proposal distribution and acceptance criteria to avoid introducing spurious CNAs. Finally, while LoPhy was developed for single-cell amplicon sequencing, the framework could be extended to reconstruct longitudinal clone trees from single-cell whole-genome sequencing or bulk sequencing.

References

1. Ascolani, G., Angaroni, F., Maspero, D., Craighero, F., Bhavesh, N.L.S., Piazza, R., Damiani, C., Ramazzotti, D., Antoniotti, M., Graudenzi, A.: LACE 2.0: an interactive R tool for the inference and visualization of longitudinal cancer evolution. BMC Bioinformatics **24**(1), 99 (Mar 2023). https://doi.org/10.1186/s12859-023-05221-3, https://doi.org/10.1186/s12859-023-05221-3

2. Chen, Z., Gong, F., Wan, L., Ma, L.: BiTSC2: Bayesian inference of tumor clonal tree by joint analysis of single-cell SNV and CNA data. Briefings in Bioinformatics **23**(3), bbac092 (May 2022). https://doi.org/10.1093/bib/bbac092, https://doi.org/10.1093/bib/bbac092

3. Dollo, L.: THE LAWS OF EVOLUTION

4. Hanahan, D., Weinberg, R.A.: The Hallmarks of Cancer. Cell **100**(1), 57–70 (Jan 2000). https://doi.org/10.1016/S0092-8674(00)81683-9, https://www.cell.com/cell/abstract/S0092-8674(00)81683-9

5. Jahn, K., Kuipers, J., Beerenwinkel, N.: Tree inference for single-cell data. Genome Biology **17**(1), 86 (May 2016). https://doi.org/10.1186/s13059-016-0936-x, https://doi.org/10.1186/s13059-016-0936-x

6. Khan, R., Mallory, X.: scLongTree: an accurate computational tool to infer the longitudinal tree for scDNAseq data (Nov 2023). https://doi.org/10.1101/2023.11.11.566680, https://www.biorxiv.org/content/10.1101/2023.11.11.566680v1, pages: 2023.11.11.566680 Section: New Results

7. Kulman, E., Kuang, R., Morris, Q.: Orchard: building large cancer phylogenies using stochastic combinatorial search (Nov 2023). https://doi.org/10.48550/arXiv.2311.12917, http://arxiv.org/abs/2311.12917

8. Lacy, M.S., Jenner, A.L.: Impact of Resistance on Therapeutic Design: A Moran Model of Cancer Growth. Bulletin of Mathematical Biology **86**(4), 43 (2024). https://doi.org/10.1007/s11538-024-01272-6, https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10950993/

9. Lu, B., Curtius, K., Graham, T.A., Yang, Z., Barnes, C.P.: CNETML: maximum likelihood inference of phylogeny from copy number profiles of multiple samples. Genome Biology **24**(1), 144 (Jun 2023). https://doi.org/10.1186/s13059-023-02983-0, https://doi.org/10.1186/s13059-023-02983-0

10. Morita, K., Wang, F., Jahn, K., Hu, T., Tanaka, T., Sasaki, Y., Kuipers, J., Loghavi, S., Wang, S.A., Yan, Y., Furudate, K., Matthews, J., Little, L., Gumbs, C., Zhang, J., Song, X., Thompson, E., Patel, K.P., Bueso-Ramos, C.E., DiNardo, C.D., Ravandi, F., Jabbour, E., Andreeff, M., Cortes, J., Bhalla, K., Garcia-Manero, G., Kantarjian, H., Konopleva, M., Nakada, D., Navin, N., Beerenwinkel, N., Futreal, P.A., Takahashi, K.: Clonal evolution of acute myeloid leukemia revealed by high-throughput single-cell genomics. Nature Communications **11**(1), 5327 (Oct 2020). https://doi.org/10.1038/s41467-020-19119-8, https://www.nature.com/articles/s41467-020-19119-8, publisher: Nature Publishing Group

11. Myers, M.A., Satas, G., Raphael, B.J.: CALDER: Inferring Phylogenetic Trees from Longitudinal Tumor Samples. Cell Systems **8**(6), 514–522.e5 (Jun 2019). https://doi.org/10.1016/j.cels.2019.05.010, https://www.cell.com/cell-systems/abstract/S2405-4712(19)30191-7

12. Nowell, P.C.: The Clonal Evolution of Tumor Cell Populations. Science **194**(4260), 23–28 (Oct 1976). https://doi.org/10.1126/science.959840, https://www.science.org/doi/abs/10.1126/science.959840

13. Pellegrino, M., Sciambi, A., Treusch, S., Durruthy-Durruthy, R., Gokhale, K., Jacob, J., Chen, T.X., Geis, J.A., Oldham, W., Matthews, J., Kantarjian, H., Futreal, P.A., Patel, K., Jones, K.W., Takahashi, K., Eastburn, D.J.: High-throughput single-cell DNA sequencing of acute myeloid leukemia tumors with droplet microfluidics. Genome Research **28**(9), 1345–1352 (Sep 2018). https://doi.org/10.1101/gr.232272.117, https://genome.cshlp.org/content/28/9/1345, company: Cold Spring Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Harbor Laboratory Press Publisher: Cold Spring Harbor Lab

14. Sollier, E., Kuipers, J., Takahashi, K., Beerenwinkel, N., Jahn, K.: COMPASS: joint copy number and mutation phylogeny reconstruction from amplicon single-cell sequencing data. Nature Communications **14**(1), 4921 (Aug 2023). https://doi.org/10.1038/s41467-023-40378-8, https://www.nature.com/articles/s41467-023-40378-8, publisher: Nature Publishing Group

15. Talevich, E., Shain, A.H., Botton, T., Bastian, B.C.: CNVkit: Genome-Wide Copy Number Detection and Visualization from Targeted DNA Sequencing. PLOS Computational Biology **12**(4), e1004873 (Apr 2016). https://doi.org/10.1371/journal.pcbi.1004873, https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1004873, publisher: Public Library of Science

16. Thijssen, R., Diepstraten, S.T., Moujalled, D., Chew, E., Flensburg, C., Shi, M.X., Dengler, M.A., Litalien, V., MacRaild, S., Chen, M., Anstee, N.S., Reljić, B., Gabriel, S.S., Djajawi, T.M., Riffkin, C.D., Aubrey, B.J., Chang, C., Tai, L., Xu, Z., Morley, T., Pomilio, G., Bruedigam, C., Kallies, A., Stroud, D.A., Bajel, A., Kluck, R.M., Lane, S.W., Schoumacher, M., Banquet, S., Majewski, I.J., Strasser, A., Roberts, A.W., Huang, D.C.S.,

Brown, F.C., Kelly, G.L., Wei, A.H.: Intact TP-53 function is essential for sustaining durable responses to BH3-mimetic drugs in leukemias. Blood **137**(20), 2721–2735 (May 2021). https://doi.org/10.1182/blood.2020010167, https://doi.org/10.1182/blood.2020010167

17. Van Loo, P., Nordgard, S.H., Lingjærde, O.C., Russnes, H.G., Rye, I.H., Sun, W., Weigman, V.J., Marynen, P., Zetterberg, A., Naume, B., Perou, C.M., Børresen-Dale, A.L., Kristensen, V.N.: Allele-specific copy number analysis of tumors. Proceedings of the National Academy of Sciences **107**(39), 16910–16915 (Sep 2010). https://doi.org/10.1073/pnas.1009843107, https://www.pnas.org/doi/10.1073/pnas.1009843107, publisher: Proceedings of the National Academy of Sciences

18. Zack, T.I., Schumacher, S.E., Carter, S.L., Cherniack, A.D., Saksena, G., Tabak, B., Lawrence, M.S., Zhang, C.Z., Wala, J., Mermel, C.H., Sougnez, C., Gabriel, S.B., Hernandez, B., Shen, H., Laird, P.W., Getz, G., Meyerson, M., Beroukhim, R.: Pan-cancer patterns of somatic copy number alteration. Nature Genetics **45**(10), 1134–1140 (Oct 2013). https://doi.org/10.1038/ng.2760, https://www.nature.com/articles/ng.2760, publisher: Nature Publishing Group