Comparative Landscape of Small RNAs in Tissue and Liquid Biopsies for Liver

Transplant Outcomes

Monima Anam[1], Christine Watkins[2], Grace Rucker[2], Katie Marlow[1], Matthew Khalil[2],

Murat Dogan[2], James Eason [2,3,4], Jason Vanatta [2,3], Corey Eymard [2,3], Cem Kuscu[2],

Zhangli Su*[1], Canan Kuscu*[2]

[1] Department of Genetics, University of Alabama at Birmingham, Birmingham, AL, 35233,

United States

[2] Transplant Research Institute, Department of Surgery, University of Tennessee Health

Science Center, Memphis, TN, 38163, United States

[3] Methodist University Hospital, Memphis, TN, 38163, United States

[4] Current address: HCA Florida Hospitals, Largo, FL, 33770, United States

*Correspondence: Canan Kuscu (ckuscu@uthsc.edu) and Zhangli Su (zsu@uab.edu)

Abstract:

Background:

Ischemia-reperfusion injury (IRI) is an inevitable consequence of liver transplantation,

arising during donor organ procurement and reoxygenation. Severe IRI is a leading

contributor to early allograft dysfunction (EAD), a post-transplant complication associated

with reduced graft survival. Current postoperative biomarkers provide limited time for

intervention, highlighting a need to identify preoperative biomarkers of IRI. Meanwhile,

tRNA fragments (tRFs) have emerged as novel biomarkers in various diseases, but

remain unexplored in the context of liver transplant.

Results:

We performed small RNA sequencing on donor liver biopsies to investigate IRI-

associated transcript changes. In parallel, donor liver perfusates were analyzed as a non-

invasive surrogate for tissue profiling. Across samples, microRNAs (miRNAs) and tRFs

were the most abundant small RNAs. Perfusate expression strongly correlated with

paired biopsies, supporting its value as a non-invasive source. Comparison of post-

reperfusion versus pre-implantation biopsies revealed that IRI reprogrammed tRF

expression. Stratification by clinical outcome showed that patients who developed EAD

exhibited specific tRNA fragments signature in both biopsy and perfusate. Receiver

operating characteristic (ROC) analysis demonstrated that a tRF-based model achieved

an AUC of 0.7748, outperforming the donor risk index alone (AUC = 0.6804), representing

a 9.4% increase in discriminative capacity.

Conclusions:

These results are the first to establish tRFs as IRI-responsive biomolecules abundant in

both donor liver tissue and perfusate, a non-invasive biofluid. In particular, various tRF

species emerged as promising candidate biomarkers for early detection of EAD. These

results lay the foundation to further investigate the diagnostic and prognostic utility of tRF

species in clinical liver transplantation.

Keywords (3-10):

tRFs, non-invasive biomarkers, ischemia-reperfusion injury, early allograft dysfunction,

liver transplantation

Background

Ischemia-reperfusion injury (IRI) is an inevitable consequence of solid organ transplantation, initiated during donor organ procurement and cold storage, and exacerbated upon reperfusion in the recipient. The sudden reoxygenation of ischemic tissue induces a complex interplay of oxidative stress, metabolic disturbances, sterile inflammation, and cell death pathways that impair graft integrity (1). Severe IRI is a principal contributor to early allograft dysfunction (EAD), a clinical syndrome presenting within the first week post liver transplant, associated with increased morbidity, primary nonfunction, and diminished long-term graft survival (2). Current diagnostic criteria for EAD primarily rely on postoperative serum biochemical markers such as alanine aminotransferase (ALT) and aspartate aminotransferase (AST), which provide limited mechanistic insight and insufficient lead time for intervention (3). This underscores the need for high-resolution molecular biomarkers capable of detecting early transcriptional and post-transcriptional responses to IRI in graft tissue and accessible biofluids, thereby enabling predictive and mechanism-informed monitoring after liver transplantation.

Small RNAs are central regulators of post-transcriptional gene expression, modulating mRNA stability, translation, and chromatin organization across diverse biological contexts. Beyond well-characterized microRNAs and piRNAs, transfer RNA-derived fragments (tRFs) and stress-induced tRNA halves (tiRs) have recently emerged as abundant and functionally versatile small RNA species generated by precise cleavage of precursor or mature tRNAs (4). Their biogenesis is strongly induced by cellular stressors, including oxidative injury, nutrient deprivation, hypoxia, and immune activation (5,6). Functionally, tRFs repress global protein synthesis, promote stress granule

assembly, displace messenger RNAs from RNA-binding proteins such as YBX1, and

participate in RNA silencing pathways via Argonaute association (7–10). A recent

publication also demonstrated that hypoxia induced tRF promotes cellular homeostasis

and protects against kidney injury by enhancing autophagic flux (11). Through these

mechanisms, tRFs act as dynamic modulators of cellular adaptation to stress, influencing

survival, metabolism, and signaling, positioning them as promising molecular sentinels of

stress phenotypes in both experimental and clinical settings (12,13).

Recent advances in transcriptomic profiling have identified small RNA signatures

as a promising class of molecular biomarkers for disease diagnosis, prognosis, and

therapeutic monitoring. Their inherent stability in biofluids, resistance to RNase

degradation, and ability to reflect dynamic physiological states make them well-suited for

minimally invasive diagnostics (14,15). In liver transplantation, liver-specific miR-122

detected in recipient serum was shown to be correlated with ALT and AST levels, thus

IRI severity and EAD (16). Furthermore, donor organ preservation solution (perfusate)

has emerged as an attractive biospecimen for biomarker discovery, capturing molecules

released directly from the donor liver during storage and perfusion (17,18). While these

investigations provide important proof-of-concepts, they focused on targeted RNA

species rather than unbiased transcriptome-wide profiling. Consequently, the broader

RNA repertoire within perfusate, including stress-responsive small RNAs such as tRFs

and tiRs, remains largely unexplored. Here, we report the first transcriptome-wide small

RNA profiling of liver transplant biopsies alongside perfusate, aimed at uncovering

mechanistically linked early non-invasive biomarkers of graft function that complement

tissue-based assays.

Results

Small RNAs and tRFs are abundantly expressed in donor liver transplant biopsy samples.

To characterize small RNAs present in donor liver tissue and determine whether they change from pre- to post-transplantation, we collected donor liver biopsy samples at two timepoints: pre-implantation (L1) and post-reperfusion (L2), followed by small RNA sequencing analysis (**Fig. 1A**). Principal component analysis (PCA) revealed distinct separation between L1 and L2 small RNA expression profiles, with PC1 and PC2 accounting for 19% and 12% of the variance, respectively (**Fig. 1B).** A variety of small RNA species were detected through sequencing; however, miRNAs comprised the largest proportion of mapped reads, followed by tRFs (**Fig. 1C**), where the top fifty most highly expressed RNAs were exclusively miRNAs or tRFs (**Fig. S1A**). Four tRF subtypes dominated the tRF landscape: tiR5, tRF5, tRF3CCA, and tRF1 species (**Fig. 1D and E**). tiR5s are known as 5' tRNA halves, which are typically 30–35 nucleotides (nt) in length, while tRF5s are shorter 5' tRNA fragments that range from 15–29 nt long. tRF3CCAs are 3' tRNA fragments that retain 3' CCA tails, a feature of mature tRNAs. While tRF1s are the remnant 3' trailer sequences that are cleaved from precursor tRNAs to form mature tRNAs. These four tRF subtypes represented the majority of mapped tRF reads across samples. Additionally, tRFs derived from Glu- and Gly-carrying parental tRNAs were highly enriched relative to other amino acid isotypes (**Fig. S1B**). Furthermore, multiple tRF species rivaled the expression of the topmost expressed miRNAs, such as Glu CTC tiR5 and Gly GCC tiR5, which were the fourth- and sixth-most highly expressed small

RNAs across samples (**Fig. S1A**). Overall, we observe that tRFs are steadily abundant

across L1 and L2 samples, serving as promising candidates for further investigation.

Comparison of L2 vs. L1 expression profiles reveals IRI-associated tRF changes.

We next examined small RNAs differentially expressed in response to IRI by

comparing L2 and L1 samples, as L2 samples were collected post-reperfusion and

therefore serve as a proxy for transplant-associated IRI. DESeq2 analysis of L2 vs. L1

small RNA profiles revealed 182 differentially expressed small RNAs, including 116 tRFs

and 66 miRNAs (**Supplementary Table 1**). Of these, 102 (58 tRFs, 44 miRNAs) were

upregulated and 80 (58 tRFs, 22 miRNAs) were downregulated in L2 compared to L1

(**Fig. 2A and B**). Strikingly, tRFs of the same subtype displayed uniform directional

changes. Among the most abundant tRF classes, differentially expressed tiR5s were

consistently enriched in L2, while significant tRF3CCA species were consistently depleted

in L2. Similar trends were observed for differentially expressed tRF1 and tRF5 species,

with the majority showing upregulation following IRI (**Fig. 2B**). Boxplots of select tiR5 and

tRF3CCA species likewise exemplify and highlight this trend (**Fig. 2C-E**). Together, these

findings indicate that IRI is associated with subtype-specific changes in tRF expression,

suggesting that tRF dysregulation may be a characteristic feature of IRI.

IRI-associated tRF changes are stratified by transplant outcome.

Next, we stratified patients by early transplant outcome to assess whether IRI-

associated tRF changes were uniquely associated with graft dysfunction. Transplant

outcomes were based on day 1 AST (aspartate aminotransferase) values (2), where high

AST values (>2000) corresponded to early allograft dysfunction (EAD, n = 18) and

moderate/low AST values corresponded to normal function (NF, n = 30) **(Fig. 3A and**

S2A). DESeq2 analysis of L2 vs. L1 in each outcome group (EAD or NF) revealed group-

dependent regulation of tRFs. In the EAD group, L2 biopsies showed unique upregulation

of numerous tiR5 species not observed in NF samples **(Fig. 3B and C)**. In contrast, the

NF group exhibited unique upregulation of multiple tRF1 species in L2 vs. L1 comparisons

(Fig. 3B, S2C and D). Notably, no tiR5 species were downregulated in L2 in either group

(Fig. S2E). Downregulated RNAs in L2 were largely shared between groups and

consisted primarily of tRF3CCA species **(Fig. S2E and F)**. Overall, we observe that post-

IRI tRF changes are uniquely dysregulated in EAD and NF grafts, suggesting that tRFs

may be prognostically linked to graft outcome.

Comparison of EAD vs. NF small RNA profiles reveals transplant outcome-

associated tRF changes, independent of IRI.

To evaluate whether small RNA profiles differed between EAD and NF grafts, we

performed differential expression analysis (EAD vs. NF) separately within L1 and L2

samples. This approach allowed us to identify small RNAs associated with transplant

outcome independent of IRI-induced changes. DESeq2 analysis revealed multiple

differentially expressed small RNAs that were shared across EAD vs. NF comparisons.

Specifically, miR-452-5p and miR-224-5p were significantly upregulated in EAD samples

compared to NF samples, while Pro CGG tiR5 was downregulated (**Fig. 4A–E and Fig.**

S3A–B). Although IRI (L2 vs. L1) induced significant increases in Pro CGG tiR5

expression, levels remained consistently lower in EAD grafts compared to NF at both

timepoints. (**Fig. 4E**). Across both biopsy timepoints, ten miRNAs were commonly upregulated in EAD (**Fig. S3A**). In contrast, five small RNAs were consistently downregulated in EAD, with the majority being tRF3CCA species, alongside one tiR5 and one miRNA species (**Fig. S3B**). Together, these findings demonstrate that small RNA profiles differ between EAD and NF groups, regardless of IRI status, suggesting that pre-implantation profiles may be sufficiently informative to aid in predicting EAD.

Pre-transplant liver perfusate is enriched with tRFs and small RNA species.

To determine whether liver biopsy-associated tRF changes could be captured non-invasively, we analyzed small RNAs in donor liver preservation solution (now referred to as perfusate). Donor livers were submerged in preservation solution and transported on ice prior to transplantation. Immediately before implantation, perfusate was collected and analyzed based on EAD (n = 49) and NF (n = 112) presence (**Fig. S4A**). RNA was extracted from the perfusate and a robust signal corresponding to small RNA size was detected after library construction (**Fig. 5A and Fig. S4B**). The size distribution from the perfusate appears different from the one derived from tissue, highlighting specific enrichment of small RNAs in the perfusate. (**Fig. S4B**) As a readily accessible biofluid that does not require tissue biopsy, liver perfusate represents a promising non-invasive source for biomarker discovery. Small RNA sequencing revealed a high abundance of small RNAs in perfusate, with tRFs as the most prevalent class, followed by mitochondrial tRFs (MtRFs) (**Fig. 5B**). Comparisons between perfusates from EAD versus NF donors showed no significant differences in the overall distribution of small RNA classes; tRFs were similarly abundant across both groups (**Fig. 5C).** Despite the lack of global

differences, the consistently high levels of tRFs and other small RNAs in perfusate

highlight its potential as a platform for identifying novel RNA biomarkers predictive of

donor liver quality and transplant outcomes.

Pre-transplant perfusate small RNAs are differentially expressed in EAD and may

help predict transplant outcome.

We next investigated whether pre-transplant perfusate small RNAs reflect donor

liver quality and could serve as biomarkers for EAD. Due to the lower complexity of cDNA

libraries from perfusate samples, DESeq2 normalization was not suitable; therefore, we

quantified expression using reads per million (RPM) and identified differentially expressed

small RNAs using Student's T-tests. This analysis revealed multiple small RNAs

significantly dysregulated between EAD and NF samples (**Fig. 6A**). Perfusate samples

contained a higher proportion of tRFs compared to biopsy tissue, with the most abundant

species being tiR5 and tRF5 fragments (**Fig. 6B**). To assess the extent to which perfusate

reflects donor liver tissue small RNA profiles, we performed correlation analyses across

matched samples. Pearson correlations revealed moderate to strong relationships

between perfusate and both L1 (mean R = 0.70) and L2 (mean R = 0.72) biopsy profiles

(**Fig. S5A and S5B**), indicating that perfusate partially recapitulates the tissue molecular

landscape. We next compared differentially expressed RNAs across perfusate (EAD vs.

NF), L2 biopsies (EAD vs. NF), and L1 biopsies (EAD vs. NF) comparisons. Several

downregulated RNAs overlapped between perfusate and tissue datasets (**Fig. 6C and**

Fig. S5C). In particular, three highly expressed Proline-tiR5 species (Pro CGG tiR5, Pro

TGG tiR5, and Pro AGG tiR5) were consistently downregulated in EAD perfusates and

EAD biopsy tissues. Additionally, Ser TGA tRF1 was downregulated in both perfusate

and L2 biopsies (**Fig. 6A and 6C–E**). Given their consistent association with transplant

outcome and high expression in perfusate, we constructed a predictive model

incorporating these three Proline tiR5s together with Ser TGA tRF1. Receiver operating

characteristic (ROC) analysis demonstrated that the tRF-based model achieved an AUC

of 0.7582, outperforming the donor risk index alone (AUC = 0.6804). The donor risk index

is a quantitative metric developed to assess the risk of graft failure based on specific

donor characteristics including donor age, cold ischemia time, cause of death, and race

(19). A combined model integrating both perfusate tRFs and the donor risk index further

improved predictive capacity (AUC = 0.7748), representing a 9.4% increase over the

donor risk index alone (**Fig. 6F**). Finally, we assessed clinical correlations of the identified

perfusate tRFs. All four species showed a significant negative correlation with day 1 AST

and ALT values, further supporting their association with early graft injury (**Fig. S5D**).

Collectively, these findings demonstrate that pre-transplant liver perfusate is a non-

invasive source of small RNAs, particularly tRFs, that partially reflect donor liver tissue

profiles and are associated with transplant outcome. Perfusate-derived tRFs may thus

represent promising non-invasive biomarkers for predicting EAD in liver transplant.

Discussion

In this study, we performed small RNA sequencing on donor liver biopsies from 48

cases and reported abundant tRFs (**Fig. 1**). Analysis of paired post-reperfusion versus

pre-implantation biopsies revealed that ischemia-reperfusion injury increased tiR5 and

tRF1 species, alongside a decrease in tRF3CCA (**Fig. 2**). Notably, tiR5 was particularly

enriched in post-reperfusion L2 tissues from patients who developed EAD, whereas tRF1

upregulation was specific to those with NF graft outcomes (**Fig. 3**). In parallel, comparison

of EAD vs NF in L1 and L2 independently highlighted miRNAs and tRFs which are

important for EAD pathology (**Fig. 4**). As a proof of concept to use tissue preservation

solution perfusate for non-invasive biomarker discovery, we sequenced perfusate small

RNAs and observed abundant tRFs that are likely derived from liver tissues (**Fig. 5**).

Perfusate sampling further highlighted the abundance of tRFs, with Pro CGG tiR5, Pro

TGG tiR5, Pro AGG tiR5, and Ser TGA tRF1 repeatedly downregulated in EAD in both

tissues and perfusate (**Fig. 6**). Our integrated analysis of liver biopsies and perfusate

revealed that both compartments contain a shared repertoire of small RNAs specifically

tRFs, highlighting the utility of these small RNAs as non-invasive biomarkers for IRI and

post-transplant outcomes.

The observed increase of multiple tiR5 species in L2 (**Fig. 2B-D**) is likely due to

the hypoxia and oxidative stress incurred during static cold storage and reperfusion,

consistent with the well-established role of tiR5s as stress-induced RNAs, including in

hypoxic contexts (4,13). During cellular stress, these tiR5s have been shown to bind to

translation initiation factors such as eIF4G/A and displace the eIF4F complex from

mRNAs, resulting in global suppression of protein synthesis while permitting selective

translation of stress-adaptive transcripts (20). In addition, tiR5s facilitate the assembly of

stress granules, sequestering non-essential mRNAs to further conserve resources and

prioritize production of proteins critical for survival and recovery (8). Beyond stress

adaptation, infection-induced tiR5s have also been implicated in immune activation,

particularly through Toll-like receptor 7 interaction (21–23). Given that IRI is characterized

as a sterile inflammatory process in which innate immunity plays a central role, the

observed tiR5 induction may represent a mechanistic link between cellular stress and

immune activation. In particular, the greater tiR5 increase in patients who developed EAD

(**Fig. 3B-C**) could indicate more tissue damage or immune activation in those transplants.

Conversely, the specific increase in tRF1s upon reperfusion in patients with NF (**Fig. 3B**

and Fig. S2D) raises the possibility of a protective role for tRF1 that warrants further

mechanistic exploration.

The identification of tRFs as dominant small RNAs in perfusate is particularly

exciting and demonstrates their potential for use in liquid biopsy monitoring (**Fig. 5**). This

is particularly important for evaluating marginal organs and machine perfusion settings,

where clinically validated, predictive molecular markers are still lacking. Previously,

Verhoeven et al. identified mRNAs and microRNAs in graft preservation solution

(perfusate) correlating with early injury (17), while Matton et al. reported that miRNA

profiles in normothermic perfusion solution could inform liver viability assessments (18).

Both studies were limited to targeted approaches. Our application of unbiased, next-

generation sequencing to perfusate along with tissue small RNA profiling identified four

clinically relevant tRFs (**Fig. 6C**). Proline-derived tiR5s display diverse, context-

dependent functions, ranging from enhancing translation and myeloid differentiation in

the bone marrow niche (Pro CGG tiR5) (24) to driving oncogenesis through HPSE2

suppression in serrated pathway colorectal cancer (Pro TGG tiR5) (25), while a

constitutively expressed Pro tiR5 directly associates with ribosomes to inhibit protein

synthesis (26). Ser TGA tRF1 (also known as tRF-1001) suppresses proliferation by

inducing G2 arrest in colorectal cancer cells (27), but promotes proliferation and stemness

in breast cancer (28), with its levels responsive to nutrient stress. This loss of potential

protective tRF signaling in liver tissues may compromise hepatocellular stress adaptation

during ischemia–reperfusion, predisposing grafts to dysfunction. Moreover, the

expression levels of these four tRFs predicted EAD occurrence better than donor risk

index scores, highlighting their clinical potential. In summary, this unbiased study expands

the known repertoire of extracellular small RNAs and positions tRFs as strong candidates

for rapid, minimally invasive assessment of graft quality during ex situ organ preservation.

Conclusion

This study provides the first transcriptome-wide characterization of tRFs in both

human liver tissue and donor organ perfusate during transplantation. We demonstrate

that specific tRFs, particularly tiR5, are closely linked to reperfusion and EAD, whereas

tRF1 appears to be associated with favorable graft outcomes. The prevalence of tRFs in

perfusate highlights their potential as non-invasive biomarkers, bridging molecular

insights from tissue with clinically accessible diagnostics. By uncovering both shared and

compartment-specific signatures, our findings lay a foundation for future validation in

larger cohorts and under machine perfusion, advancing tRFs toward clinical application

for improved organ assessment and transplant precision.

Figures and Figure legends

Figure 1

Figure 1. Abundance of tRNA fragments (tRFs) in donor liver transplant biopsy samples.

(A) Liver biopsy samples were obtained from donor livers at two time points: pre-implantation (L1) and post-reperfusion (L2), followed by small RNA sequencing. **(B)** Principal component analysis (PCA) of small RNA sequencing data reveals distinct clustering between L1 and L2 small RNA profiles. **(C)** Stacked bar plot displaying the proportion of mapped reads for each small RNA type in L1 and L2 samples. For each patient (x-axis), the bars represent the relative proportion of total reads (y-axis) assigned to each small RNA type. Across samples, miRNAs comprised the largest majority of mapped reads, followed by tRFs. **(D)** Schematic illustrating four highlighted tRF subtypes. The tRF nomenclature echoes the region of tRNA from which tRFs are derived. The fragments shown are examples of each tRF subtype and do not represent all possible variants. tRF-1 species (green) are derived from precursor tRNAs (not pictured), while tiR5, tiR3, and tRF3CCA species (pink) are cleaved from mature tRNAs (blue). **(E)** Heatmap displaying the relative expression (Log2 RPM) of different tRF subtypes across L1 and L2 samples. Overall, biopsy samples showed abundant tRF expression.

Figure 2

Figure 2. Significant tRF alterations associated with ischemia-reperfusion injury (IRI) during liver transplantation.

(A) Volcano plot of differentially expressed small RNAs between L2 and L1 samples, identified using DESeq2. The x-axis shows the log2 fold change (L2 vs. L1), and the y-axis shows statistical significance (-log10 adjusted p-value). The top 10 most significantly altered small RNAs are labeled, many being tRF3CCA species downregulated in L2. **(B)** MA plot of differentially expressed small RNAs (L2 vs. L1) identified using DESeq2. The x-axis represents gene expression (log2 normalized counts), and the y-axis shows the

log2 fold change. Differentially expressed RNAs are color-coded by RNA type (tRF

subtype or miRNA). Select highly abundant, significantly altered tRFs are labeled,

including many tiR5 species that are upregulated in L2. **(C)** Primary and secondary

structure of parental tRNA-Glu-CTC-1-3, with its most abundant tiR5 fragment highlighted

in red (positions 1-35). Cleavage sites that generate the second and third most abundant

Glu CTC tiR5 isoforms from tRNA-Glu-CTC are marked in green and blue. **D)** Boxplot of

Glu CTC tiR5 expression in L1 vs. L2 samples, showing significant upregulation in L2,

consistent with IRI-associated induction. **(E)** Primary and secondary structure of parental

tRNA-Gln-CTG-1-4, with its corresponding tRF3CCA fragment highlighted in red

(positions 58–75). **(F)** Boxplot of Gln CTG tRF3CCA expression in L1 vs. L2 samples,

showing significant downregulation in L2, indicative of IRI-associated suppression.

Figure 3

A

L1/L2 Biopsy sample summary (n = 48)

Group	EAD	NF	p value
Number of patients	18	30	
Recipient mean age	57.3	56.6	N.S.
Recipient gender (F/M)	7/11	10/19	N.S.
Donor mean age	47.4	37.7	0.012 *
Donor gender (F/M)	9/9	16/14	N.S.
Mean CIT (hours)	4.89	4.34	N.S.
Mean AST (Day 1)	5472.8	1072.4	2.7e-04 ***
Mean ALT (Day 1)	1618.3	542.5	0.0015 **

C

Gly GCC tiR5

B

UP-regulated in L2 (post) vs L1 (pre)

EAD: **Gly GCC tiR5, Gly CCC tiR5, Glu TTC tiR5,** His GTG tiR5, Asp GTC tiR5, Val TAC tiR5, SeC TCA tiR5, Gly GCC tRF5, Glu TTC tRF5, Glu CTC tRF5, Gly CCC tRF5, Glu TTC tRF3, miR-27a-5p, miR-1246-5p, miR-3178-5p, miR-5195-3p, miR-8069-2-3p

17 | 14 | 4

NF: **Arg ACG tRF1,** Arg TCG tRF1, Val CAC tRF1, Val CAG tRF5

Tyr GTA tRF1, Thr TGT tRF1, Lys TTT tiR5, Lys CTT tiR5, Val CAC tiR5, Glu CTC tiR5, Pro CGG tiR5, Val TAC tiR3, Glu CTC tiR3, Gly CCC tiR3, miR-486-5p, miR-451a, miR-451a-5p, miR-4284-5p

D

Glu TTC tiR5

Figure 3. IRI-associated tRF alterations are transplant outcome specific.

(A) Clinical and demographic characteristics of the patient cohort, stratified by transplant outcome (EAD or NF), which was classified based on day 1 post-transplant aspartate aminotransferase (AST) levels. Cold ischemia time (CIT), the duration the donor liver remained on ice prior to transplantation, averaged 4.6 hours across groups. **(B)** Differential expression analysis of L2 vs. L1 samples was performed separately within EAD and NF groups. The Venn diagram displays small RNAs upregulated in L2 relative

to L1 for each group. In the EAD group, uniquely upregulated RNAs were predominantly

tiR5 species, whereas in the NF group, uniquely upregulated RNAs were mainly tRF1

species. A subset of tiR5 and tRF1 species was commonly upregulated in both groups,

alongside other tRF and miRNA species. **(C)** and **(D)** Boxplots showing normalized

expression (log10 transformed) of representative tiR5 species (Gly-GCC and Glu-TTC),

stratified by transplant outcome (EAD vs. NF) and biopsy timepoint (L2 vs. L1). Statistical

comparisons were performed using Wilcoxon rank-sum tests; p-values are indicated

above comparisons. Both panels exemplify the EAD-specific upregulation of tiR5s,

showing significant upregulation in L2 compared to L1 exclusively in EAD samples

(Student's t-test).

Figure 4. Transplant outcome-associated differences in liver biopsy small RNA

expression.

(A) Volcano plot of differentially expressed small RNAs between early EAD and NF in L1

samples, identified using DESeq2. The x-axis represents log2 fold change (EAD vs. NF),

and the y-axis indicates statistical significance (−log10 adjusted p-value). The top three

most significantly altered small RNAs are labeled. **(B)** Volcano plot of differentially

expressed small RNAs between EAD and NF in L2 biopsy samples. The top four most

significantly altered small RNAs are labeled. **(C)** and **(D)** Boxplots of normalized

expression (log10 transformed) of representative miRNAs (miR-452-5p and miR-224-5p),

stratified by transplant outcome (EAD vs. NF) and biopsy timepoint (L1 vs. L2). Both

miRNAs are consistently upregulated in EAD compared to NF, independent of time point.

Statistical comparisons were performed using Wilcoxon rank-sum tests; p-values are

shown above comparisons. **(E)** Boxplots of Pro CGG tiR5 expression, stratified by

transplant outcome and biopsy timepoint. While Pro GCC tiR5 expression increases after

reperfusion injury (L2 vs. L1), expression remains significantly lower in EAD samples

compared to NF at both timepoints (Student's t-test).

Figure 5

Figure 5. Pre-transplant liver perfusate solution is enriched with small RNAs.

(A) Donor livers were transported on ice in organ preservation solution prior to transplantation. Immediately before transplantation, the preservation solution (perfusate) was collected, and RNA was isolated for small RNA sequencing. **(B)** Composition of

mapped small RNAs in liver perfusate samples, stratified by EAD and NF groups. Across

both groups, tRFs represented the predominant RNA species, followed by mitochondrial

tRFs (MtRFs). **(C)** Relative abundance of mapped small RNA classes in liver perfusate

samples, stratified by NF and EAD groups. Each boxplot displays the percentage of total

mapped small RNAs in each class, with each point representing an individual sample.

RNA classes were similarly abundant in both EAD and NF samples, with no statistically

significant difference between groups (Wilcoxon test).

Figure 6

Figure 6. Pre-transplant perfusate small RNAs are differentially expressed based on transplant outcome.

(**A)** MA plot depicting differentially expressed small RNAs (EAD vs. NF) from pre-
transplant liver perfusate samples identified using Student's T-test. The x-axis represents
gene expression (log2 RPM), and the y-axis shows the log2 fold change. Significantly
differentially expressed RNAs are color-coded by RNA type (tRF subtype or miRNA).
Select highly abundant, significantly altered tRFs are labeled, including multiple proline-
tiR5 species that are downregulated in EAD**. (B)** Bar plot displaying the expression (mean
RPM) of the top fifty most highly expressed small RNAs in donor liver perfusate samples,
stratified by transplant outcome (EAD (orange) or NF (blue)). These top fifty genes consist
entirely of miRNAs and tRFs, which are outlined in red**. (C)** Venn diagram comparing
downregulated small RNAs in three comparisons: perfusate EAD vs. NF, L2 EAD vs. NF,
and L1 EAD vs. NF. One species, Pro CGG tiR5, was consistently downregulated across
all three comparisons, suggesting potential relevance in EAD. Additionally, two other
proline-tiR5 species (Pro TGG and Pro AGG) were commonly downregulated in EAD in
perfusate and L1 biopsy samples. **(D)** and **(E)** Boxplots showing normalized expression
(log2 RPM) of representative tRF species (Pro-CGG-tiR5 and Ser-TGA-tRF1) stratified
by transplant outcome (EAD vs. NF) in liver perfusate samples, where these tRFs were
significantly downregulated in grafts that developed EAD. Statistical comparisons were
performed using Student's T-tests; p-values are indicated above comparisons. **(F)** ROC
analysis evaluating the transplant-outcome predictive capacity of three models: donor risk
index (black), perfusate tRFs (blue), and a combined model incorporating both perfusate
tRFs and donor risk index (red). The donor risk index includes donor age, cold ischemia
time, cause of death, and race. The tRF model incorporates the expression of three Pro
tiR5s and Ser TGA tRF1 in pre-transplant perfusate. The resulting AUCs were 0.6804

(donor risk index), 0.7582 (tRFs), and 0.7748 (combined model), indicating that the

addition of perfusate tRFs improved the model's discriminatory capacity for EAD.

Methods

<u>Patient recruitment and specimen collection</u>

Liver tissue and perfusate samples were obtained from adult orthotopic liver transplant (OLT) recipients (≥18 years old) at Methodist University Hospital after obtaining informed consent. All procedures were conducted under protocols approved by the University of Tennessee Health Science Center Institutional Review Board (IRB#20-07742-XM and IRB#20-07640-XP).

Needle biopsy specimens were immediately immersed in RNA*later* (Thermo Fisher Scientific, Cat. #AM7021) and incubated at 4 °C overnight to ensure complete permeation. RNA*later* was then removed, and the samples were stored at –80 °C until RNA extraction. Perfusate samples were collected at the end of organ procurement concurrently with pre-implantation liver (L1) biopsies. The solutions were centrifuged at 2,400 rpm for 30 minutes at room temperature to remove debris. The resulting supernatant was aliquoted and stored at –80 °C until RNA isolation.

<u>RNA isolation and small RNA library preparation</u>

Total RNA was extracted from needle biopsies using the Direct-zol RNA MicroPrep Kit (Zymo Research, Cat. #R2062) according to the manufacturer's instructions, including on-column DNase I treatment to remove genomic DNA contamination. For DOPS samples, RNA was isolated from 500 µl of solution using the same kit, following the manufacturer's protocol for liquid samples, and eluted in 15 µl of RNase-free water. Purified RNA was stored at –80 °C, and repeated freeze–thaw cycles were avoided to minimize RNA degradation.

Small RNA libraries were prepared using the NEBNext Multiplex Small RNA Library Prep

Set for Illumina (New England Biolabs, Cat. #E7330) following the manufacturer's

instructions. For each sample, 100 ng of tissue RNA or 5 µl of DOPS RNA was used as

input. Indexing was performed using the NEBNext Multiplex Oligos for Illumina, with PCR

amplification carried out for 15 cycles for tissue RNA libraries and 18 cycles for perfusate

RNA libraries. Amplified products were resolved on an 8% polyacrylamide gel (Thermo

Fisher Scientific, Cat. #EC6215BOX), and target bands were excised and purified. Equal

amounts of libraries were pooled and sequenced on an Illumina NextSeq 500 platform,

generating approximately 10 million 75 bp single-end reads per sample.

<u>Small RNA-seq data analysis</u>

Raw small RNA sequencing reads were trimmed to remove adapter sequences using

Cutadapt (v5.1). Adapter-trimmed reads were then mapped and annotated using Unitas

(v1.7.3), a specialized tool for comprehensive small RNA annotation. For liver biopsy

samples, read count matrices generated by Unitas were imported into DESeq2 (v1.44.0)

in R for normalization and differential expression analysis. Prior to analysis, RNAs with

zero counts in more than 80% of samples were excluded to reduce low-abundance noise.

Data visualization was performed in R using ggplot2 (v3.5.1) and ggpubr (v0.6.0). For

perfusate samples, read count matrices generated by Unitas were imported into R and

normalized to reads per million (RPM). Given the lower complexity of perfusate libraries

relative to biopsy samples, differential expression analysis was performed using Student's

t-tests on RPM values. Data visualization was carried out in R with ggplot2 (v3.5.1) and

ggpubr (v0.6.0), and receiver operating characteristic (ROC) curves were generated

using the ROCR package (v1.0-11).

Declarations

<u>Ethics approval and consent to participate</u>

This study was conducted in accordance with the ethical standards of the University of

Tennessee Health Science Center Institutional Review Board, which approved all

protocols (IRB#20-07742-XM and IRB#20-07640-XP). Liver tissue and perfusate

samples were obtained from adult orthotopic liver transplant (OLT) recipients (≥18 years

old) at Methodist University Hospital after written informed consent was obtained from all

participants.

<u>Availability of data and materials</u>

The small RNA-seq data will be deposited to GEO.

<u>Competing interests</u>

The authors have no competing interests to declare.

<u>Funding</u>

This research was supported by NIH grants R00CA259526 (to Z.S.), T32NS121721 (to

M.A.), T32GM135028 (to K.M.), and Transplant Research Institute at UTHSC start up

fund (to C.K.).

Authors' contributions

C.W., G.R., and M.K. helped with patient sample processing, RNA isolation and small

RNA library preparation. M.D. and Cem K. coordinated sample collection, storage and

contributed to study design. J.E., J.V., and C.E. coordinated patient recruitment and

sample collection at the Methodist Hospital. M.A., K.M. and Z.S. performed data analysis

on RNA-seq. Canan K. and Z.S. supervised the study and wrote the manuscript with the

help of M.A and K.M. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank the Methodist University Hospital transplant team for their

assistance with patient recruitment, sample and clinical data collection. We are deeply

grateful to the patients and their families for their participation and generous donations to

this research. We also would like to acknowledge our colleagues at the Transplant

Research Institute for their helpful discussions and ongoing support. We would like to

thank IT Research Computing Team at the University of Alabama at Birmingham for NGS

support.

References

1. Zhai Y, Petrowsky H, Hong JC, Busuttil RW, Kupiec-Weglinski JW. Ischaemia-reperfusion injury in liver transplantation--from bench to bedside. Nat Rev Gastroenterol Hepatol. 2013 Feb;10(2):79–89.

2. Olthoff KM, Kulik L, Samstein B, Kaminski M, Abecassis M, Emond J, et al. Validation of a current definition of early allograft dysfunction in liver transplant recipients and analysis of risk factors. Liver Transpl. 2010 Aug;16(8):943–9.

3. Giraudi PJ, Codotto G, Baccarani U, Baralle F, Gondolesi G, Licastro D, et al. Ischemia/Reperfusion Injury and Outcomes in Liver Transplantation Assessed by Omics Technologies: Where Do We Stand? Transplantation. 2025 Aug 13;

4. Kumar P, Kuscu C, Dutta A. Biogenesis and Function of Transfer RNA-Related Fragments (tRFs). Trends Biochem Sci. 2016 Aug;41(8):679–89.

5. Li G, Manning AC, Bagi A, Yang X, Gokulnath P, Spanos M, et al. Distinct Stress-Dependent Signatures of Cellular and Extracellular tRNA-Derived Small RNAs. Adv Sci (Weinh). 2022 Jun;9(17):e2200829.

6. Su Z, Wilson B, Kumar P, Dutta A. Noncanonical Roles of tRNAs: tRNA Fragments and Beyond. Annu Rev Genet. 2020 Nov 23;54:47–69.

7. Gebetsberger J, Wyss L, Mleczko AM, Reuther J, Polacek N. A tRNA-derived fragment competes with mRNA for ribosome binding and regulates translation during stress. RNA Biol. 2016 Nov 28;14(10):1364–73.

8.   Emara MM, Ivanov P, Hickman T, Dawra N, Tisdale S, Kedersha N, et al. Angiogenin-

induced tRNA-derived Stress-induced RNAs Promote Stress-induced Stress Granule

Assembly. J Biol Chem. 2010 Apr 2;285(14):10959–68.

9.   Goodarzi H, Liu X, Nguyen HCB, Zhang S, Fish L, Tavazoie SF. Endogenous tRNA-

Derived Fragments Suppress Breast Cancer Progression via YBX1 Displacement.

Cell. 2015 May 7;161(4):790–802.

10. Kuscu C, Kumar P, Kiran M, Su Z, Malik A, Dutta A. tRNA fragments (tRFs) guide Ago

to regulate gene expression post-transcriptionally in a Dicer-independent manner.

RNA. 2018 Aug;24(8):1093–105.

11. Li G, Sun L, Xin C, Hao T, Kharel P, Manning AC, et al. A hypoxia-responsive tRNA-

derived small RNA confers renal protection through RNA autophagy. Science. 2025

Aug 28;389(6763):eadp5384.

12. Anderson P, Ivanov P. tRNA fragments in human health and disease. FEBS Lett.

2014 Nov 28;588(23):4297–304.

13. Pandey KK, Madhry D, Ravi Kumar YS, Malvankar S, Sapra L, Srivastava RK, et al.

Regulatory roles of tRNA-derived RNA fragments in human pathophysiology. Mol

Ther Nucleic Acids. 2021 Dec 3;26:161–73.

14. Costa B, Li Calzi M, Castellano M, Blanco V, Cuevasanta E, Litvan I, et al. Nicked

tRNAs are stable reservoirs of tRNA halves in cells and biofluids. Proc Natl Acad Sci

U S A. 2023 Jan 24;120(4):e2216330120.

15. Srinivasan S, Yeri A, Cheah PS, Chung A, Danielson K, De Hoff P, et al. Small RNA

Sequencing across Diverse Biofluids Identifies Optimal Methods for exRNA Isolation.

Cell. 2019 Apr 4;177(2):446-462.e16.

16. Farid WRR, Pan Q, van der Meer AJP, de Ruiter PE, Ramakrishnaiah V, de Jonge J,

et al. Hepatocyte-derived microRNAs as serum biomarkers of hepatic injury and

rejection after liver transplantation. Liver Transpl. 2012 Mar;18(3):290–7.

17. Khorsandi SE, Quaglia A, Salehi S, Jassem W, Vilca-Melendez H, Prachalias A, et al.

The microRNA Expression Profile in Donation after Cardiac Death (DCD) Livers and

Its Ability to Identify Primary Non Function. PLOS ONE. 2015 May 15;10(5):e0127073.

18. Matton APM, Selten JW, Roest HP, de Jonge J, IJzermans JNM, de Meijer VE, et al.

Cell-free microRNAs as early predictors of graft viability during ex vivo normothermic

machine perfusion of human donor livers. Clin Transplant. 2020 Mar;34(3):e13790.

19. Feng S, Goodrich NP, Bragg-Gresham JL, Dykstra DM, Punch JD, DebRoy MA, et al.

Characteristics associated with liver graft failure: the concept of a donor risk index.

Am J Transplant. 2006 Apr;6(4):783–90.

20. Ivanov P, Emara MM, Villen J, Gygi SP, Anderson P. Angiogenin-induced tRNA

fragments inhibit translation initiation. Mol Cell. 2011 Aug 19;43(4):613–23.

21. Pawar K, Kawamura T, Kirino Y. The tRNAVal half: A strong endogenous Toll-like

receptor 7 ligand with a 5'-terminal universal sequence signature. Proc Natl Acad Sci

U S A. 2024 May 7;121(19):e2319569121.

22. Pawar K, Shigematsu M, Sharbati S, Kirino Y. Infection-induced 5′-half molecules of

tRNAHisGUG activate Toll-like receptor 7. PLOS Biology. 2020 Dec

17;18(12):e3000982.

23. Yu J, Zhang X, Cai C, Zhou T, Chen Q. Small RNA and Toll-like receptor interactions:

origins and disease mechanisms. Trends Biochem Sci. 2025 May;50(5):385–401.

24. Kfoury YS, Ji F, Mazzola M, Sykes DB, Scherer AK, Anselmo A, et al. tiRNA signaling

via stress-regulated vesicle transfer in the hematopoietic niche. Cell Stem Cell. 2021

Dec 2;28(12):2090-2103.e9.

25. Wang XY, Zhou YJ, Chen HY, Chen JN, Chen SS, Chen HM, et al. 5'tiRNA-Pro-TGG,

a novel tRNA halve, promotes oncogenesis in sessile serrated lesions and serrated

pathway of colorectal cancer. World J Gastrointest Oncol. 2023 Jun 15;15(6):1005–

18.

26. Gonskikh Y, Gerstl M, Kos M, Borth N, Schosserer M, Grillari J, et al. Modulation of

mammalian translation by a ribosome-associated tRNA half. RNA Biol. 2020

Aug;17(8):1125–36.

27. Lee YS, Shibata Y, Malhotra A, Dutta A. A novel class of small RNAs: tRNA-derived

RNA fragments (tRFs). Genes Dev. 2009 Nov 15;23(22):2639–49.

28. Wan X, Shi W, Ma L, Wang L, Zheng R, He J, et al. A 3′‑pre‑tRNA‑derived small

RNA tRF‑1‑Ser regulated by 25(OH)D promotes proliferation and stemness by

inhibiting the function of MBNL1 in breast cancer. Clin Transl Med. 2024 May

9;14(5):e1681.