1 **Ecological drift drives microbiome variation in the red flour beetle**
2

3 Pratibha Sanjenbam, Harshith Koppa Guruswamy, Rittik Deb, Shivansh Singhal, Sneha
4 Garge, Shyamsunder Buddh, Shubha Govindarajan, and Deepa Agashe*
5
6 Address:
7 National Centre for Biological Sciences (NCBS–TIFR), GKVK Campus, Bellary Road,
8 Bengaluru, India, 560065
9
10 *Correspondence: dagashe@ncbs.res.in

11
12 **Short running title:** Sources of drift in the flour beetle microbiome

ABSTRACT

Many eukaryotes harbor diverse and complex microbial communities that provide significant benefits to the host. However, recent work demonstrates substantial variation in microbial communities across hosts and over time, and the causes of this instability are not always clear. We describe large temporal, among-individual and across-population variation in the microbiome of the red flour beetle *Tribolium castaneum*, a widespread pest of stored grain flour and a laboratory model organism. Across a 5-year period, the total load and composition of the bacterial community of laboratory beetle populations fluctuated dramatically, cycling between a few dominant taxa and hundreds of rare taxa. Our experiments show that this ecological drift is caused by a lack of maternal microbial transmission and large microbial load fluctuations in the primary reservoir of microbial inocula: the flour. Flour microbial load is important because newly hatched larvae (from eggs) and newly eclosed adults (from pupae) acquire microbes from ingested flour containing the fecal material of other individuals. The flour microbiome fluctuates depending on host population density and stage structure, with larval feces increasing microbial load, and antimicrobial secretions from mature adults decreasing it. Finally, periodic flour replacement during population maintenance also dramatically decreases flour bacterial load, contributing to stochasticity in microbial colonization. Our work thus represents a case study of significant and pervasive ecological drift even in a highly controlled laboratory environment. We suggest that such unstable microbial communities may not be uncommon, and can serve as good models to understand early stages of the evolution and ecology of host-microbial interactions.

Keywords
Microbiota, host-bacterial interaction, flour beetle, stochasticity, community assembly, microbiome

INTRODUCTION

Animal-microbe relationships are seemingly ubiquitous, and thus the focus of a large body of biological research, especially on obligately host-associated microbes (e.g., see (Holt et al., 2024; McFall-Ngai, 2024). However, several mammals, insects, and birds host transient or unstable microbiomes, and/or do not appear to acquire any specific benefits from the microbes that they harbour (e.g., Deb et al., 2019; Hammer et al., 2017, 2019; Łukasik & Kolasa, 2024; Phalnikar et al., 2018). Many host species also show high individual variation in their microbiomes (reviewed in, e.g., (Miller et al., 2018), even across individuals reared in identical conditions in the laboratory (Goulda et al., 2018; Obadia et al., 2017; Vega & Gore, 2017). What mechanisms and processes drive this variation?

Variability in the microbiota of a host species suggests a strong role for stochastic processes in driving microbial community assembly and maintenance. However, these processes may operate across different scales of organization, including at the level of the meta-community (Miller et al., 2018). Broadly, variation may arise from historical contingency or limited dispersal, or it may reflect ecological drift, whereby stochastic processes (birth, death, and dispersal) rather than deterministic processes govern the abundance of specific microbes

(Zhou & Ning, 2017). For instance, Vega and colleagues inoculated *Cenorhabditis* worms with similar numbers of two competitively equivalent *Escherichia coli* strains (Vega & Gore, 2017). At high inoculum densities, most worms were reliably colonized by a nearly equal mixture of both strains. But at low inoculum density, demographic stochasticity determined the outcome, with most worms colonized by only one of the two strains. Such ecological drift should be stronger when unopposed (or weakly opposed) by selection favouring obligate, consistent mutualism. On the other hand, determinism in community assembly does not necessarily reflect strong selection, because other processes — such as a strong environmental influence coupled with stable environments, or dietary specialization — may also drive deterministic colonization and assembly dynamics. Finally, an interaction between host- and microbe- specific factors may also lead to between-host microbiota variation; e.g., a combination of host control and microbial colonization may drive distinct routes of microbiome assembly in different individuals. Thus, several factors and processes may combine to drive microbiome variation across hosts. For example, strains of the symbiont *Lactobacillus plantarum* isolated from wild *Drosophila melanogaster* showed positive density-dependent colonization (i.e., were strong colonizers) — as observed in worms, discussed above — but strains isolated from laboratory flies or humans remained consistently poor colonizers regardless of density (Obadia et al., 2017). This study also uncovered additional effects of stochastic population bottlenecks, presence of competitor strains, and host genetics, highlighting the complexity of mechanisms underlying host-to-host microbiome variation. Even in a neutral model of community assembly (i.e., under no selection), weak dispersal from the environment to the host, coupled with a short host lifespan, led to different microbiome compositions in different hosts (Zapién-Campos et al., 2020). Thus, ecological drift may arise from several sources, driving large variation in host microbiomes. However, the relative contribution and consequences of these factors for ecological drift, especially for multi-species microbial communities, remains unclear.

Here, we focus on host-bacterial interactions in the red flour beetle *Tribolium castaneum*. These beetles are generalist, widespread pests of cereal grain flours, and complete their entire life cycle (egg to larva to pupa to adult) in flour. Hence, the beetles effectively live in a largely closed system, which can be efficiently reproduced in the laboratory under controlled environmental conditions such as temperature and humidity. Prior studies report somewhat diverse sets of flour beetle microbiomes, though generally dominated by bacteria from the family *Enterobacteriaceae*, with some variation across life stages (Agarwal & Agashe, 2020; Bi et al., 2024; Prakash et al., 2023; Zheng-Yan & Yang Meng-Ting LU Yu-Jie, 2020). Importantly, our previous work indicated a significant (but non-essential) beneficial effect of flour-associated microbes on beetle fitness parameters such as reproduction, survival, and immune function (Agarwal & Agashe, 2020; Prakash et al., 2023). However, across these two studies, we observed substantial variation in beetle microbiomes, which was also reflected in results from a different group finding variation in beetle microbiomes across sampling locations (Zheng-Yan & Yang Meng-Ting LU Yu-Jie, 2020). Given the observed fitness benefits of the microbiome for beetles, this variation is surprising, and indicated that the non-obligate host-bacterial interactions may be associated with significant ecological drift. Hence, this system presented a unique opportunity to measure and understand the factors responsible for ecological drift.

We first systematically quantified individual and population level as well as temporal
variation in the flour beetle microbiome across several years, using previously stored beetle
samples from populations that had been maintained in near-identical laboratory conditions.
Finding   large variation in all these cases, we conducted a series of experiments to
determine the causes of the apparent ecological drift. Specifically, we tested how the nature
and magnitude of variation in the beetle microbiome was affected by sources of bacterial
inocula entering the flour habitat (fresh flour, experimenter identity, and beetle feces) and
factors influencing bacterial survival and load in the habitat (beetle-secreted antimicrobials
and flour replacement during population maintenance). We show that ecological drift in the
flour beetle microbiome is driven by a combination of factors that alter the fecal bacterial
inoculum in the flour, as well as the timing of flour change implemented during population
maintenance (whereby used flour is replaced with fresh flour). We suggest that similar
factors at play in the beetles' natural habitat (flour and grain warehouses) may help explain
why a strong, obligate host-bacterial mutualism has not evolved in this system, despite
potential benefits for the host.
120
121   **METHODS**
122
123   **Population maintenance and sample storage**
124
For all our experiments, we used a large outbred stock population, described earlier
(Agarwal & Agashe, 2020). In brief, we allow 2000-3000 adults (~2-4 weeks old) to oviposit
on 750 g fresh wheat flour for 7 days in a large plastic box, and then remove the adults. After
allowing 2-3 weeks for offspring development, we conduct maintenance, replacing the used
flour with fresh wheat flour and removing dead individuals and larval molts. At this stage,
most individuals are either large larvae or pupae, and these are allowed to develop further
for another 2-3 weeks until we see a large number of adults. We then start a new stock
generation with a subset of these adults. Three independent replicate boxes of the outbred
line are maintained in this fashion, but we used one of them as our focal stock population,
and all experiments were done using individuals derived from this population, except where
noted otherwise.
136
137   Every packet of wheat flour used for our experiments is first frozen at -80°C for 4 hrs to
138   remove any prior pest infestation (Agarwal & Agashe, 2020) and is allowed to return to room
139   temperature before use. All beetle populations are maintained in plastic boxes with holes in
140   the lid for ventilation (except when noted otherwise), and kept in dark incubators at 33±1 °C.
141   Beetles or flour samples used for microbiome analysis were stored at -80°C.
142
143   **Collecting samples to estimate microbiome variation across time, replicate**
144   **populations, and generations**
145
146   *Experiment 1a:* To determine change in beetle microbiome composition in the same lineage
147   over time, in the year 2022, we created a new, parallel stock population. To obtain adults
148   that were exactly 2 weeks old, at each generation we allowed adults to lay eggs only for 48
149   hours (instead of 7 days in the usual protocol described above) (Fig S1A). Every 3 months,
150   we stored 2 week-old female beetles (n = 10) from the population.

Experiment 1b: To expand the above time series, we added samples from the same outbred stock population, which were stored for other projects from 2015–2021. These represent opportunistic samples and are therefore not spaced out systematically across the years. Since these were sampled directly from the stock boxes, precise adult age was not known; however, given the beetle life cycle, we estimated that the females were between 2-4 weeks old, comparable to our 2022 samples. When possible (July 2020, February 2021, and March 2022), we also analysed males from the same population (n=10) to assess sex-specific variation.

Experiment 1c: To determine microbiome variation across replicate populations with similar original ancestry, we sampled and stored individuals from the three independent stock boxes described above, in February 2016 and again in March 2022. As described for Experiment 1a, samples for March 2022 were collected following 48 hours of oviposition, and therefore, females were 2 weeks old. In contrast, samples stored in February 2016 were 2-4 week old females (Fig S1B). When available, we also analysed males from the same boxes.

Experiment 2: To test for differences between parent and offspring microbiomes, we divided ~600 2-week-old adults from the stock population into groups of 200, and allowed each group to oviposit in 200 g fresh wheat flour for 7 days, creating three replicate boxes (Fig S1C). This protocol mimics our stock population maintenance. From each replicate box, we sampled 1-week old offspring (n = 10 females and 10 males) for microbiome analysis. These data allowed us to test for (a) variation in parent vs. offspring microbiomes and (b) variation across replicate boxes of offspring of parents from the same stock box.

Testing the effect of external factors that may influence the beetle microbiome

Experiment 3: To test the effect of factors such as experimenter identity, exposure to sterile vs. normal air, and flour sterilization, we allowed 1500 ~4 week-old adults from the stock population to oviposit for 48 hrs in 600 g double sifted fresh wheat flour (Fig S2A). To measure the effects of air exposure, we sifted out 100 eggs per treatment and placed each set of eggs in 100 g fresh wheat flour, with normal air supply (box lid had holes) vs. sterile air (box without holes, but opened under sterile air supply in a laminar airflow twice a week). Similarly, we generated two more batches of eggs and placed them in 100 g of either normal or dry-heated wheat flour (90°C for 45 mins). In all cases, we allowed eggs to develop for 5 weeks, i.e., until they were 2-week-old adults, and stored for microbiome analysis (n = 5 males and 5 females/treatment). The sterilized wheat flour served as our positive control, wherein we expected a distinct microbiome and lower total bacterial load due to the heat treatment. To test the effect of experimenter identity, we sampled 1500 adults from the stock population and split them equally between two experimenters. Each experimenter allowed adults to oviposit for 7 days in independent boxes containing 600 g double sifted fresh wheat flour. After 5 weeks, the resulting adult offspring (1000 each) served as each experimenter's independent set of parents. These were allowed to oviposit for 48 hrs in 300 g of double sifted fresh wheat flour, and offspring were sampled when they turned 2-week-old adults.

Testing for maternal transmission of the beetle microbiome

Experiment 4a: To test for maternal microbiome transmission, we allowed ~50 2-week-old
adult beetles to oviposit for 48 hrs on 5 g fresh wheat flour in each of three replicate boxes.
We sampled the parents (n = 5 females/replicate) and eggs (n = 3 replicates, with 50 pooled
eggs/replicate), and stored them for microbiome analysis (Fig S3A).

Experiment 4b: In experiment 4a, contamination of the egg surface from flour or adult feces
was a concern. To address this, we directly dissected ovaries and eggs from ~2 week old
stock females (n=3) that were previously stored at -80°C. We rinsed each female in 70%
ethanol and washed with sterile saline (0.9% sodium chloride). We placed the beetle on a
5% agar plate and cut the posterior tip of the abdomen using fine scissors (Fine Science
Tools) under a stereomicroscope (Leica V4.12). We applied pressure to the sternum to force
the abdominal contents outside, then teased apart the ovary with needles and forceps and
collected eggs and ovaries from each female for DNA extraction (Fig S3B). We ensured that
we did not disrupt the gut during dissection.

Experiment 4c: In the above experiment, we used ~2 week old females. In subsequent
experiments (see below), we observed that the adult microbiome took ~4 weeks to stabilize,
and it was possible that vertical transmission could occur at that stage. Hence, we analysed
the microbiomes of dissected eggs from females collected from stock populations at 2, 4,
and 6 weeks post eclosion (Fig S3C). We dissected them as described above, and collected
3 groups of 10 eggs per age category, with each group containing eggs from 2-5 females.

Testing the effect of fecal inoculum on the beetle microbiome

Experiment 5: To test whether beetle feces serve as a source of bacterial inoculum, we
conducted a series of experiments (Experiments 5-8). First, if adult feces are a key source of
inoculum, we expected that bacterial load would be positively correlated with adult density.
To test this, we took 2-4 week old adults, and created two densities by placing 200 (low) or
2000 adults (high) in 300 g wheat flour. We allowed adults to oviposit for 7 days and then
removed them. We split the flour (including eggs) into three replicate lines (100 g each), and
allowed offspring to develop for 5 weeks, i.e., until they were 2-week-old adults. We stored
offspring at -80°C for microbiome analysis (n = 5 females/replicate/treatment). For one
replicate per treatment, we also stored 5 males (Fig S2B).

Experiment 6: Next, determined whether the larval and adult fecal microbiomes were
different, and changed with age. To obtain individuals of known age, we allowed ~200 2-4-
week-old adults from the stock population to oviposit in 200 g of fresh wheat flour, and
removed the adults after 48 hrs. Once we started observing adult offspring (after 4 weeks),
we sampled adults every week for 6 weeks, placing them in empty 1.5 ml tubes (Eppendorf)
for 24 hrs to collect their feces (n = 30 adults/tube, with 3 replicate tubes per time point).
After this, we placed the adults back in their box to maintain the same population density
until the next sampling point. Thus, we obtained adult fecal microbiomes from 1-6 weeks of
age. Similarly, we obtained larval fecal microbiomes, sampling every 3 days for 21 days,
until the larvae began pupating (n = 30 larvae/tube, with 3 replicate tubes per time point). We

242   immediately processed all fecal samples for microbiome analysis, as described in later
243   sections (Fig S4A).
244
Experiment 7: To determine the effect of larval vs. adult fecal inoculum on recipient
microbiomes, we added feces to fresh wheat flour and monitored the microbiome of recipient
pupae until they were 6-week-old adults (i.e., for 7 weeks total). We first prepared 'enriched
flour' containing either larval or adult feces, by adding either 1000 larvae (10-14 days old) or
200 adults (2-4 weeks old) to 50 g of wheat flour for 3 days. Then, we mixed 80% of this
enriched flour with 20% of fresh wheat flour to ensure that nutrient depletion of the enriched
flour would not lead to starvation in the recipients. We placed single recipient pupae
collected from our stock population in 1.5 mL tubes containing either 0.5 g of fresh wheat
flour (control) or flour enriched with larval or adult feces. Every week, we sampled and
stored individuals (n = 10 females/week/treatment) for further processing (Fig S4B).
255
Experiment 8: Finally, we tested whether the amount of larval fecal inoculum would alter
recipient beetle microbiomes. We generated larval-enriched wheat flour as described above
(Experiment 7), and then prepared a series of flour mixes with different proportions of
enriched flour (0%, 25%, 50%, 75%, 100%), adding fresh wheat flour as required. We
placed single pupae in each treatment as described above for 3 weeks, and sampled at the
end point (i.e., 2-week-old adults) for microbiome analysis (Fig S4C).
262
263   **Measuring the impact of quinones on bacteria and the beetle microbiome**
264
Experiment 9: Adult flour beetles secrete antimicrobial quinones from specialized stink
glands (Pedrini et al., 2015; Prendeville & Stevens, 2002), and we suspected that this may
explain why adult presence reduced flour bacterial loads and increased individual level
microbiome variation (see Results). To test whether beetle-associated bacteria are
differentially sensitive to quinones, we first isolated bacteria from beetle life stages and used
flour, using multiple solid growth media (Table S1). To prepare glycerol stocks, we scraped
colonies from the plate, resuspended them in 800 µL of autoclaved 60% glycerol solution,
and stored at -80°C. We extracted genomic DNA for each isolate, and identified it by Sanger
sequencing the 16S rRNA gene (Table S1). To prepare quinone extracts, we took 100 6-
week-old adult beetles from the stock population and placed them in an empty Petri plate
(90 mm diameter). We stimulated the release of quinones from stink glands using a cold
shock, placing the plate on ice until beetles were immobile. We removed the plate from the
ice, allowed beetles to recover, and repeated the process for a total of 30 mins. We removed
the adults and collected the released quinones by adding 1 mL of hexane to the plate. After
mixing with a pipette, we transferred the quinone extract to a 1.5 mL tube. We revived
bacterial stocks by streaking on Luria-Bertani (LB) agar (Difco) and M-Enterococcus agar
(HiMedia) plates, incubated at 28°C for 24 hrs, picked up multiple colonies, suspended in
0.9% autoclaved saline and normalized all cultures to an OD_{600} of 0.1 using sterile saline.
We spread the culture on 1.5% LB agar plates using sterile swabs to make a bacterial lawn,
and dried the plates inside a laminar hood for a few minutes. We placed 4 sterile filter paper
discs (6 mm diameter) onto the prepared bacterial lawn plates, spaced evenly apart. We
added 20 µL of the prepared quinone extract onto 3 of the discs and the solvent (hexane) as
our negative control for the 4[th] disc (1 Petriplate per isolate, with 3 technical replicates of the

test and 1 control). We incubated plates at 28°C for 14-16 hrs (~40 hrs for *Enterococcus*
isolates because of slow growth), and measured the zone of inhibition around each disc as a
measure of sensitivity to quinones.
291
Experiment 10: We tested whether focal bacterial strains are also sensitive to quinones
secreted in wheat flour. We took fresh wheat flour and added feces (which are rich in the
beetle-associated *Izhakiella* strain that we are unable to culture), or 0.3% lyophilized
bacterial cells (*Acinetobacter radioresistens* or *Enterococcus faecium*) (Knorr Eileen et al.,
2015), either with or without quinone extract. The treatments without quinones allowed us to
measure bacterial death in the absence of quinones. We prepared quinone extract (as
described for Experiment 9) and added 50 µL extract to 0.5 g of flour in 1.5 ml tubes with
mixing, every 4th day. We periodically stored a subset of tubes at -80°C to measure the
bacterial death rate (n = 8 replicate tubes/treatment/time point) (Fig S5A).
301
Next, we tested whether quinones perturb microbial load in the flour in the presence of
beetles, which the flour bacteria could colonize. We prepared flour enriched with larval or
adult feces, as described for Experiment 7. In another treatment, we added both larval feces
and quinone discs. However, here the quinone preparation differed slightly: we placed 100
beetles directly in hexane at -80°C (without a cold shock), and added quinone extract to the
flour only at the beginning of the experiment. We placed single pupae isolated from the stock
population in a 1.5 mL tube containing 0.5 g of the appropriate flour for each treatment. We
sampled a subset of pupae every week for 6 weeks, storing 10 females/week/treatment at -
80°C for further processing (Fig S4B).
311
Experiment 11: To test the impact of quinones on the microbiome, we used RNA-mediated
interference (RNAi, using double-stranded RNA, dsRNA) to knock down expression of the
gene *GT39*, which is involved in quinone synthesis (Sawada et al., 2020). As negative
controls, we included beetles that were left untreated, injected with TE buffer (10mM Tris,
1mM EDTA buffer to control for effects of injection), or with GFP (green fluorescent protein)
dsRNA (which should not influence quinone synthesis). To synthesize *GT39* dsRNA, we
extracted DNA from female beetles (as described below), and amplified the gene of interest
using primers with the T7-promoter, described in (Sawada et al., 2020).
GT39_Forward- 5'-TAATACGACTCACTATAGGG-GGAGGTCACCCAGAACAACT-3';
GT39_Reverse-5'-TAATACGACTCACTATAGGG-TGACATCCCTTGGCACATATTC-3';
To obtain the dsRNA for GFP, we used a strain of *Escherichia coli* MG1655 carrying GFP to
get template DNA, and used the following primers that we designed.
GFP_Forward-5'-TAATACGACTCACTATAGGG-GCAGCCGGATCCTTTGTATAG-3';
GFP_Reverse-5'-TAATACGACTCACTATAGGG-TGCTGAAGTCAAGTTTGAAGGTG-3'.
We used 500 ng of the amplified DNA as a template for in-vitro transcription in a 20 µL
reaction using the Thermo MEGAscript RNAi Kit. We followed the kit instructions, with
overnight incubation at 37°C followed by elution of the dsRNA in TE buffer. We injected 0.2
µL of the dsRNA solution (~200 ng/pupae) in early-stage (~2 day old) pupae using a
microinjector. We handled control pupae in an identical fashion, except untreated controls
which were not injected. We isolated pupae in 1.5 mL tubes containing 0.5 g of fresh wheat
flour for one week, during which time the pupae eclosed into adults. We grouped these
adults into 10 pairs (10 individuals per sex) in 3 g of flour in a small plastic box (35 mm

diameter) for 3 weeks (i.e., until we obtained ~4 week old adults) (n = 7 replicate
boxes/treatment). One week after grouping, we sampled and dissected a few females to
check the stink gland phenotype (n = 2 females from each of 3 replicate boxes per
treatment). At the final time point, we stored all remaining beetles until further processing
(Fig S5B).
339
340 **Measuring the impact of periodic flour replacement on the beetle microbiome**
341
Experiment 12: Our results suggested that fresh flour has very little bacterial inoculum, that
fecal inoculum from larvae is critical for the colonization of the beetle gut, and that adult-
secreted quinones selectively suppress the growth of specific bacteria. This led us to predict
that in our stock populations, substantial ecological drift may be generated due to a
combination of these factors and our stock maintenance protocol, which involves periodic
replacement of the flour. To test this, we sampled and stored flour (0.5 g) and beetles (n =
10 for each available life stage) every 5 days from our main stock population. We sampled
and analysed microbiomes across two successive discrete-generation cycles. We also
similarly sampled an independent experimental population that was maintained under
continuous generation cycles, in which the number of larvae does not fluctuate as much as
under discrete generations.
353
354 **DNA extraction and sequencing for flour and beetle samples**
355
We extracted DNA from stored (–80°C) or fresh beetles (larvae, pupae, or adults) using the
Qiagen DNeasy Blood and Tissue kit, with the following modifications. We first surface-
sterilized each sample by dipping it in 70% ethanol, and then washed it with sterile water.
For adult beetles, we removed the wings (elytra) and legs. The beetles were again surface
sterilized, washed, and kept in individual 1.5 mL tubes. We flash froze each beetle in liquid
nitrogen and crushed it using a micro-pestle. Next, we added 180 μl of lysis buffer and 20 μl
Proteinase K solution from the kit, vortexed the sample, and kept it at 56ºC for ~18 hours.
Then, we followed the rest of the kit protocol, eluting DNA in 30 μl of nuclease-free water.
364
To obtain the wheat flour microbiome, we modified the protocol to increase the yield of
bacterial DNA and minimize plant (flour) DNA. We took 100 mg of wheat flour, added 300 μL
of Redford buffer (1 M tris HCl; 0.5 M EDTA; 1.24% Triton), and placed the tube horizontally
with periodic gentle shaking on a vortex for 2 hrs, as described in (Sanjenbam & Agashe,
2024). We pelleted the flour debris by centrifuging at 10,000 rpm for 5 mins, and used the
supernatant for further processing as described above.
371
We quantified the DNA concentration in each sample using the Qubit dsDNA HS assay kit.
We amplified the V3-V4 hypervariable region of the 16S rRNA gene using a modified
Illumina 10N primer to facilitate detection of rare taxa (Naik et al., 2023). We followed the
standard Illumina 16S metagenome amplification protocol, with the following modification.
For wheat flour and larval samples, which have high chloroplast and mitochondrial
contamination, we added 25 and 15 μM PNAs, respectively (peptide nucleic acids, to
specifically block chloroplast and mitochondrial DNA amplification; (Lundberg et al., 2013)).
We prepared libraries from the amplified product as per standard Illumina protocols and

sequenced on the Illumina Miseq platform (300 x 2 paired-end reads). In each MiSeq flow
cell, we also included one negative control, where we did not add any DNA template, but
carried out all amplification steps.
383
384 **Measuring total bacterial load using quantitative PCR**
385
We measured absolute bacterial load in the same beetle and flour samples used for
microbiome analysis (n = 6-10 for beetles and 2-4 for wheat flour). We used 1 μL of
extracted DNA solution from each sample (described above) and obtained 200 bp amplicons
for quantification, using primers designed to amplify only bacterial 16S rRNA (forward: 5'-
CGGTAATACGGAGGGTGCAA-3'; reverse: 5'- TCCTCCAGATCTCTACGCAT -3'), using
the qPCR protocol described in (Sanjenbam & Agashe, 2024). For beetle samples, we
normalized the amount of bacterial DNA to the amount of beetle DNA, using primers for the
host gene *rps3* (not found in bacteria or plants) (forward 5'-CAGGGGTCTGTGTGCCATTG-
3'; reverse 5'-TCACACCCTGCAAAATAACCACT-3'). Similarly, we normalized flour samples
to the actin gene *ACT2* (forward: 5'-CAAATCATGTTTGAGACCTTCAATG-3'; reverse 5'-
ACCAGAATCCAACACGATACCTG-3') (Gordon et al., 2016) that represents the amount of
plant DNA. In this manner, we effectively estimated the total bacterial load per gram host
tissue (or flour), as $2^{-\Delta Ct}$, where ΔCt = Ct(16S rRNA) – Ct(rps3 or actin).
399
400 **Data analysis**
401
We used R for all statistical analysis (CoreTeam, 2016). To determine the microbiome
composition from the amplicon sequencing output from MiSeq, we used the DADA2 pipeline
(Callahan et al., 2016) to generate a taxonomic table using 100% identity with the Silva
reference database (training set v138.1) (ref). We first filtered out chloroplast and
mitochondrial reads, obtaining an average of 30,000 reads (range 20,000-60,000) in beetle
samples, and 5,000 reads (range 5,000-15,000) in flour samples. We obtained less than 100
reads on average in our negative controls, suggesting no significant contribution of
contaminants from reagents. We used a subset of samples to conduct a rarefaction analysis,
finding that ~10,000 bacterial reads for the beetle samples, and ~5,000 bacterial reads for
the flour samples, led to saturation of community richness respectively (Fig S6). For samples
with fewer reads than these thresholds, we repeated the sequencing. Samples that
repeatedly failed to generate sufficient reads were excluded from further analysis.
414
We used the R packages 'phyloseq' and 'vegan' to estimate Shannon diversity in all
samples (McMurdie & Holmes, 2015; Oksanen et al., 2019). To quantify individual variation in
microbiome composition, we used the package 'NST' (Ning et al., 2019) to calculate the
taxonomic beta diversity (β_{rc} - Raup-Crick Dissimilarity) (Chase et al., 2011). We carried out
PERMANOVA using the 'adonis2' function in the package 'vegan' in R to quantify the impact
of various factors on the beetle microbiome. Finally, we used the R package 'ggplot2' for
visualization.
422
423
424 **RESULTS**
425

Beetle microbiomes vary across time, replicate populations, generations, and individuals

We first analyzed the microbiomes of female beetles sampled from the same outbred stock population across years (**Experiments 1a and 1b, Table 1**). The bacterial community composition varied significantly across time (**Fig 1A**; PERMANOVA, timepoint effect (month/year combination): $R^2 = 0.25$, P = 0.0009; year effect: $R^2 = 0.13$, P = 0.0009), as did bacterial richness and diversity (**Fig S7**; Kruskal-Wallis tests, $P_{diversity} = 0.01$; $P_{richness} = 6.5$ e-05). We observed similar patterns across three sampled time points for males, indicating that underlying processes were not sex-specific (**Fig S8**). Next, we tested whether three independently maintained stock populations initiated from the same founders also acquired divergent microbiomes (**Experiment 1c, Table 1**). In February 2016, the populations had indeed diverged, although we could not identify the dominant bacterial taxon (**Fig 1B**; PERMANOVA, lineage effect: $R^2 = 0.13$, P = 0.003). In March 2022, the three populations appeared to harbour more similar microbiomes dominated by *Izhakiella*, though the community composition still differed significantly (**Fig 1B**; lineage effect: $R^2 = 0.10$, P = 0.004). Males at this time point showed similar patterns (PERMANOVA, lineage effect: $R^2 = 0.11$, P = 0.0009, **Fig S8).** Thus, in addition to temporal changes observed in a given population over several years, the microbiomes of independently maintained stock populations also diverged significantly from each other.

Next, we tested for shorter-term variation in microbiome composition (**Experiment 2**). For three replicate "offspring" boxes set up from the same set of stock-derived parents, we again observed significant variation in microbiomes (PERMANOVA, replicate effect, females: $R^2 = 0.23$, P=0.0009; males: $R^2 = 0.22$, P=0.0009). Thus, beetle microbiomes varied dramatically across time and across independent lineages maintained under the same laboratory conditions. Despite this variation, however, a few bacterial genera such as *Izhakiella*, *Enterococcus*, *Pseudomonas*, and *Acinetobacter* often dominated the community (**Fig 1A**), and were identified as members of the core beetle microbiome (**Fig S9**). For instance, *Izhakiella* and *Enterococcus* were consistently present in ~80% of beetle samples with a moderate relative abundance of 10%, whereas *Pseudomonas* and *Acinetobacter* were found in ~50% of the samples at a low relative abundance of 1% (**Fig S9**).

We then analysed the degree of individual-level variation in microbiome composition, which also appeared to be high (**Fig 1A**). Using taxonomic beta diversity (β_{rc}) as a metric of microbiome diversity, we found that the degree of individual variation in a population also changed across time (**Fig 11D**; Kruskal-Wallis test, time effect: P < 2.2e-16). For instance, females had highly similar microbiomes in September 2015, July 2020, and June 2022 (median β_{rc} approaching -1), whereas females sampled from the same stock population had highly dissimilar microbiomes in March 2021 and December 2022 (median β_{rc} approaching +1) (**Fig 1D**). We also tested whether individual variation in microbiome composition was associated with total bacterial load, measured using qPCR (normalized with a beetle-specific primer to obtain an estimate of total bacterial DNA per unit host DNA for each sample). The bacterial load per beetle varied significantly over time (ANOVA, load ~ time: P = 4.64e-06 **Fig 1E**), but was not correlated with β_{rc} (Spearman's rho = -0.14, P = 0.16). Both results were robust to removing a potential outlier data point from August 2020 (ANOVA, load ~

time: P = 0.012; Spearman's rho = -0.15, P = 0.17). In contrast, the total bacterial load was
positively correlated with the relative abundance of the genus *Izhakiella*, suggesting that this
genus may colonize beetles consistently and well (**Fig 1F**; Spearman's rho =0.34, P =
0.0002). We did not observe such correlations for the relative abundance of *Enterococcus* or
for rare taxa that sometimes had a relatively high combined relative abundance (e.g., **Fig
1A**, March 2022 samples) (**Fig S10**).
478
Together, these results showed that the beetle microbiome composition and load vary a lot
across individuals, generations, and populations; but dominant taxa recur, including the
genus *Izhakiella,* which appears to be a strong colonizer. Next, we considered factors that
could drive this variation across individuals and over time. The flour beetle system is a
relatively closed environment, where the flour serves as both habitat and food for all life
stages. All individuals consume flour and deposit fecal material in the flour (except pupae,
which do not feed), such that microbes continuously circulate through beetles and flour,
enabling dispersal and exchange across host individuals. Hence, there are broadly three
sources of bacterial inocula in this system: the environment, flour, and beetles. We
considered each of these in turn.
489
490   **Sources of bacterial inoculum: external environmental factors and fresh flour**
491
We first tested whether variation in external sources of bacterial inocula such as air,
experimenter, and fresh flour could contribute to the variation in beetle microbiomes. We set
up independent experiments to test these factors, with the following treatments: exposure to
normal vs. sterile air; experimenter 1 vs experimenter 2; and fresh flour vs. heat-treated
fresh flour (**Experiment 3**). In each case, we found relatively weak effects of the treatment.
Air sterility had no significant effect, experimenter identity explained ~15% of the variation in
bacterial community composition (PERMANOVA, R^2 = 0.15, P = 0.02), and flour treatment
explained ~13% of the variation (PERMANOVA, R^2 = 0.13, P = 0.012) (**Fig 2A**) in adult
female microbiomes. All treatments had a significant effect on adult male microbiomes but
only ~13%-16% of variation was explained by these factors (**Fig S11**). All factors also
influenced the degree of individual variation (**Fig 2B**; Wilcoxon tests, $P_{air\ sterility}$ = 0.0001;
$P_{experimenter}$ = 0.003; P_{flour} = 0.0004). For instance, experimenter 1 generated individuals with
more similar microbiomes compared to experimenter 2, and individuals reared on dry heated
wheat flour had highly dissimilar microbiomes compared to those fed fresh wheat flour.
Although air sterility and flour affected the relative abundance of *Izhakiella* (Pearson chi-
square test, $P_{air\ sterility}$ = 0.005546, P_{flour} = 4.531e-09; **Fig 2A**), none of the factors influenced
total bacterial load (**Fig 2C**; ANOVA, load ~ factor, $P_{experimenter}$= 0.3; P_{air} = 0.6; P_{flour} = 0.3),
which tended to be lower than that observed in our temporal sampling experiment (compare
Fig 2C with **Fig 1E**). The experimenter identity effect was potentially confounded by an
unexpected difference in population density between the boxes handled by the two
experimenters (e.g., experimenter 1 had ~700 larvae whereas experimenter 2 had ~1200
larvae on the 18th day). This density effect was specifically tested later. Together, our results
suggested that the tested external environmental factors are unlikely to explain the observed
large variation in beetle microbiomes, though it is possible that these factors may be
important in combination and/or over several generations.
517

Sources of bacterial inoculum: host-associated factors

Next, we investigated the role of possible host-associated sources of bacterial inoculum on the beetle microbiome, which may include maternal transmission through the egg, and horizontal transmission via larval or adult beetle feces. In a preliminary experiment, we observed that microbiomes of groups of freshly laid eggs were different from those of their mothers (**Experiment 4a**, PERMANOVA, R^2 = 0.09, P = 0.002, **Fig 3A**) with lower abundance of *Izhakiella*, in contrast to expectations under maternal transmission. For eggs and ovaries dissected from females before oviposition (**Experiment 4b**, **Fig 3B**), we again observed very low relative abundance of *Izhakiella* (~3%, **Fig 3C**). Finally, we found that the egg microbiome varied significantly with maternal age (**Experiment 4c;** PERMANOVA, age, R^2 = 0.35, P = 0.014; **Fig 3D**), indicating unstable egg microbiomes. Together, these results suggest a lack of substantial and consistent bacterial transmission from mother to offspring via eggs.

Therefore, we turned to the possibility that feces could serve as an important source of bacteria that colonize beetle guts. If true, higher population densities should reduce the stochasticity in colonization by increasing fecal inoculum in the flour. Recall that our previous results from Experiment 3 (different experimenters happened to generate populations with distinct densities, which also had different microbiomes) are consistent with this prediction. Hence, we set up a new experiment where a single experimenter set up beetle populations at either low or high adult density. Here, we observed that compared to low density populations, individuals reared in high density populations had similar total bacterial loads (ANOVA, load ~ density, P = 0.06) but higher relative abundance of *Izhakiella* (average 47% vs. 18% in high vs. low density respectively), lower individual variation (Wilcoxon test, P = 3.10e-14), and distinct microbiomes (**Experiment 5**, PERMANOVA, density effect combining all replicates, R^2 = 0.06, P = 0.0009; **Fig 4A, Fig S12**). Replicate populations within each treatment also varied significantly (PERMANOVA, replicate effect: low density, R^2 = 0.17, P = 0.0009; high density, R^2 = 0.13, P = 0.0009), potenitialy reducing the effect of population density. Nonetheless, these results pointed to fecal inoculum as an important factor affecting microbiome variability.

In the above experiment, we focused on adult fecal material, but larvae are also a potentially important source of fecal microbial inoculum. On one hand, the microbiomes of larvae and adults could be similar since all life stages consume the same flour, which includes fecal material from all individuals in the population. On the other hand, given the distinct feeding patterns and physiology of larvae and adults, distinct microbiomes could be associated with each life stage. Indeed, we found significant differences in both the microbiome composition and load of larvae vs. adults sampled from the same populations (**Experiment 3**, PERMANOVA comparing larvae vs. parents; R^2 = 0.21 , P = 0.0009; **Fig 4E**). Larvae had significantly higher abundance of *Izhakiella* (two experimental blocks: 30% vs. 14% in larvae and adults respectively, Chi-square test, P = 0.01; and 53% vs. 6%, P = 9.42e-10). Given that the larval stage is the primary growing stage and larvae consume a large amount of flour to support the rapid increase in body size, we speculated that larval feces may serve as a major source of bacterial inoculum in the flour.

We tested this by sampling feces of larvae vs. adults across development. The larval fecal
microbiome composition was very stable and was primarily composed of *Izhakiella* (~80%)
with the highest bacterial load at days 9-10, which is approximately halfway through the total
larval period of ~20 days (**Experiment 6**, **Fig 5A**, **Fig S13A**). In contrast, the adult fecal
microbiome was highly variable until 2 weeks post-eclosion, and then stabilized with a large
increase in total bacterial load (**Fig 5B**, **Fig S13B**). Overall, the bacterial load in larval feces
was higher than in adult feces (ANOVA, load ~ age x stage, P_{age} = 3.52e-6; P_{stage} = 0.0007;
$P_{age \, x \, stage}$ = 0.0007). Given that most of our stored adult beetle samples (analysed in the first
section of the Results) were ~2 weeks old, these results may help explain why we observed
high variability in beetles sampled across time (**Fig 1A**). Further, exposing pupae to larval
feces led to higher bacterial loads, more stable microbiomes, and lower individual variation
in the eclosed adults, compared to pupae exposed to flour enriched with adult feces
(**Experiment 7**, **Fig 5C-D**, statistics reported in the figure legend, **Fig S13C**). The stabilizing
effect of larval feces also increased with the proportion of fecal inoculum provided in the flour
(**Experiment 8, Fig 5F-G**, statistics reported in the figure legend, **Fig S13D-E**). Together,
these results support a strong role of fecal-oral transmission in microbiome assembly and
stability, with larval feces as the main source of inoculum for the beetle microbiome.
581
582   **The role of adult-secreted quionones**
583
The results above indicated that fecal inoculum is important, and that it is life-stage specific.
However, beyond the effect of fecal volume and turnover in the gut (which is expected to be
higher for larvae), we suspected that adults also have an inhibitory effect on bacterial load in
the flour, given prior work showing antimicrobial effects of quinone compounds secreted
from adult stink glands in the flour (Pedrini et al., 2015; Prendeville & Stevens, 2002). The shift in
adult microbiomes between 2-3 weeks of age coincides with the stink glands becoming fully
functional (Lisa M. Unruh et al., 1998; Markarian et al., 1978). Further, the effect of adult
density observed above (higher density reduces individual microbiome variation) could be
mediated by density-dependent upregulation of quinones (Khan et al., 2017). Hence, we
tested whether bacteria in the beetle microbiome were sensitive to adult-secreted quinones
(**Experiment 9**).
595
We found differential sensitivity of beetle-associated culturable bacteria to stink gland
extracts from 6-week old females; the solvent negative control did not inhibit growth (**Fig 6A**,
ANOVA, zone of inhibition ~ isolates, P = 0.001). Different isolates of the same genus also
had distinct sensitivity (e.g., *Enterococcus* and *Acinetobacter*), demonstrating large
functional variation even in closely related bacteria. Notably, opportunistic pathogens such
as *Serratia* were among the most resistant to quinones. Unfortunately, we were unable to
culture *Izhakiella* from beetles or larvae; hence we used *I. capsodis* that was previously
isolated from mirid bugs (Aizenberg-Gershtein et al., 2016). This species was very sensitive
to quinones, but given the variation observed among other congeners, it is unclear whether
our beetle-associated *Izhakiella* strain(s) have similar quinone resistance as *I. capsodis*.
606
We next tested whether quinones influence bacterial persistence in the flour, in the absence
of live adult beetles (or larvae) (**Experiment 10**). We added large quantities of lyophilized
bacteria or adult fecal material to flour, and monitored bacterial loads over time, with or

without added quinone extracts. Quinones had differential effects on the load of different
bacteria in flour (**Fig 6B, Table S2**). The total bacterial load in fresh flour as well as flour
enriched with adult feces generally declined over time (likely due to dessication in the dry
flour environment), and the decline was exacerbated on adding quinones (**Fig 6B,** Fig
S14A). However, lyophilized *Enterococcus* survived for at least ~2 weeks in flour, and
declined only when we added quinones. Surprisingly, the load of *Acinetobacter* increased
significantly over time (**Fig 6B,** Fig S14A). We were unable to identify the cause of this
counterintuitive bacterial growth, which is not expected to happen in dry flour. Overall, the
decline in bacterial load due to quinones was robust to a different method of quinone
extraction (**Fig S14B**). Thus, adult secreted quinones could play an important role in
modulating bacterial loads in flour, potentially influencing microbiome assembly and
dynamics.
622
To test this hypothesis, we suppressed quinone production in adult beetles via RNA-
mediated interference of expression of the gene *GT39*, which is involved in quinone
synthesis (Sawada et al., 2020) (**Experiment 11**). The dsGT39 treated beetles had missing
or underdeveloped stink glands, confirming that the RNAi was effective (**Fig S5C**).
Compared to controls, the bacterial load in these beetles was significantly higher, as
expected (**Fig 6C**, t test, dsGFP vs. dsGT39: P = 0.014). The quinone knockdown beetles
also had significantly lower individual variation in microbiomes (**Fig 6D,** dsGFP vs. dsGT39,
Wilcoxon test, P = 0.02). However, the microbiome composition of beetles was similar in
both cases (PERMANOVA, dsGFP vs. dsGT39: $R^2 = 0.05$, P = 0.4), and we did not observe
high *Izhakiella* abundance in any of the beetles (**Fig S14C**). Thus, the effect of quinone
production was weaker than we expected, perhaps due to the lack of initial larval fecal
inoculum in the fresh flour supplied to pupae in this experiment. Nevertheless, these results
demonstrate an important role for adult-secreted quinones in regulating flour beetle
microbiomes.
637
638 **Population maintenance protocol affects flour microbiomes and bacterial load**
639
Our experiments thus far highlighted two important factors driving beetle microbiome
assembly and stability: larval fecal inoculum, which increases bacterial loads in the flour, and
adult-secreted quinones, which suppress it. However, the periodic replacement and
replenishment of flour — a critical part of flour beetle population maintenance — could
potentially dilute the effect of these two factors, and therefore also influence microbiomes.
As individuals grow and fecal matter accumulates, the flour quality declines, flour can get
moldy, and populations become more susceptible to infection and nutrient deficiencies.
Therefore, periodically replacing conditioned flour with fresh flour is essential for maintaining
healthy laboratory populations of flour beetles.
649
In our maintenance protocol (**Fig 7A**), flour replacement typically coincides with low
abundance of larvae, and high abundance of pupae or young adults in the population.
Hence, most of the larval inoculum might be effectively removed during the flour change,
with very few remaining larvae that could contribute new fecal inoculum in the flour. Instead,
as new adults eclose and mature, the flour accumulates increasing concentrations of
quinones (Pedrini et al., 2015; Prendeville & Stevens, 2002), which should suppress bacterial

loads until the eventual increase in the number of larvae can counteract the effect of quinones. This plausible effect of flour replacement is supported by changes in the microbiome composition and total bacterial load in flour sampled across two generations in the same stock population (**Fig 7B**). The disruptive effect of flour replacement is also seen in populations maintained under continuous generation cycles, where larval abundance does not fluctuate as much as under discrete generation cycles (**Fig 7C, Fig S15C**). In both sets of populations, bacterial loads dropped precipitously after a flour change, and gradually increased over time — coincident with increasing number of larvae — along with *Izhakiella* abundance.

Supporting our earlier results (**Fig 1F**), we again found that the total bacterial load strongly correlates with the relative abundance of *Izhakiella* in the flour from stock populations (**Fig 7D**, Spearman's rank correlation rho = 0.5, P = 0.0001), reinforcing the conclusion that this genus is a strong colonizer of flour beetles. This is further evidenced by the mismatch in flour and beetle microbiomes (**Fig 7B–C**, beetle microbiomes shown under the x-axis; also see **Fig S15** for individual level composition): at each sampling time, beetle microbiomes are often more enriched in *Izhakiella* than the flour microbiome.

DISCUSSION

Here, we identify and quantify the effect of multiple sources of stochasticity in community assembly and maintenance of the flour beetle microbiota. Our results show that several aspects of beetle life history and ecology explain the observed ecological drift in the microbiome (i.e., apparently idiosyncratic changes in the composition of the microbial community). (1) Bacteria are not maternally transmitted through the eggs or ovaries. (2) Larvae host large and consistent bacterial communities. Feces of larvae thus represent a key source of bacterial inoculum in the flour, likely because high larval feeding rates allow rapid proliferation and movement of bacteria between flour and gut, and between individuals. (3) In comparison, adult beetles harbor smaller and more variable microbiota. Hence, adult feces represent a smaller inoculum source, with the effect of lower feeding rates in adults compounded by the secretion of antimicrobial quinones. (4) Larval microbiomes stabilize rapidly compared to adult microbiomes, such that even a brief period of larval fecal enrichment can increase and homogenize flour and beetle microbial communities. (5) Stock maintenance protocols exacerbate ecological drift for two related reasons. First, the periodic replacement of used flour with fresh flour dramatically reduces flour microbial loads, mimicking frequent migration of adults in grain warehouses to colonize new resource patches. Second, at the time of such flour change, most larvae are either in the last instar (the non-feeding stage) or have already pupated, while newly eclosed adults begin secreting quinones. As a result, bacterial load in the flour does not increase rapidly. Drawing upon these results, we propose a model for bacterial transmission, persistence and host colonization in the flour beetle system (**Fig 8**). Together, the periodic flour replacement and absence of larval fecal inoculum, lack of vertical bacterial transmission, and quinone secretion by adults cause substantial stochasticity in community assembly, leading to the observed ecological drift in adult flour beetle microbiomes. Flour beetles may thus serve as a good example of neutral community assembly, with a short host lifespan and weak microbial dispersal into hosts: two factors that are predicted to drive large individual variation

702 in microbiome composition (Zapién-Campos et al., 2020). Our results contribute to the
703 growing body of research on the role of stochastic colonization on insect-microbe
704 associations, which is important to more broadly understand insect populations and
705 adaptation to environmental change (Łukasik & Kolasa, 2024).
706
Our work allows us to speculate about the mechanisms that may prevent or facilitate the
establishment of obligate host-microbial interactions, which involve both host and bacterial
control (Schmidt & Engel, 2021). A recent mathematical model showed that in the absence of
strict vertical transmission, horizontal or environmental transmission of microbiota is critical
for the establishment of a host-microbial relationship (Leftwich et al., 2020). In *T. castaneum,*
we did not observe evidence of vertical transmission, and we find that horizontal
transmission is context-dependent. The lack of vertical transmission may be interpreted in
different ways. It may arise due to constraints placed by ecological drift in adult
microbiomes, e.g., because young ovipositing females do not have a stable microbiome to
transmit. In turn, the absence of such transmission may also contribute towards preventing
the establishment of a stable microbiome, by enhancing microbiome stochasticity at the
beginning of an individual's development. Thus, the lack of vertical transmission may explain
why a strict host-microbe relationship has not evolved in flour beetles, despite evidence for
host fitness benefits (Agarwal & Agashe, 2020). On the other hand, weak (rather than strong)
vertical transmission could also be selectively favoured, e.g., because it can drive among-
host variation that is adaptive in fluctuating environments (Bruijning et al., 2022). It is
plausible that the frequent colonization of new habitat patches by *T. castaneum* — a feature
of its life history as a human-associated pest — may generate such selection for weak
vertical transmission of microbiota. Yet another possibility is that strong selection on
quinones as signals of population density — important for negative density dependent
population regulation (Khan et al., 2017) — may also indirectly impede the establishment of
a stable microbiome. Finally, it is possible that all of these mechanisms contribute to the lack
of stable host-microbiome associations in flour beetles, and that similar processes may be at
play in several other species that lack such relationships (Hammer et al., 2019). Although
distinguishing cause and effect in our case is difficult, it may be possible via evolution
experiments where each contributing factor can be independently manipulated and its effect
on the emergence of vertical transmission quantified. Prior laboratory experiments with stink
bugs suggest that such evolution is plausible within relatively short timescales (Koga et al.,
2022). More generally, experimental evolution approaches may be fruitful to understand the
relative contribution of several factors driving stochastic vs. deterministic dynamics, and the
emergence of strong host-microbial associations or lack thereof.
738
Our results also complement prior work on microbiome variation in insects that undergo
large ecological shifts across developmental stages (Girard et al., 2023). Dramatic internal
reorganization during metamorphosis in holometabolous insects (including beetles) is
associated with high beta diversity and microbiome turnover, compared to hemimetabolous
insects (Manthey et al., 2023). Microbiome shifts across life stages are also common in
insects (Engel & Moran, 2013), including dragonflies ((Nobles & Jackson, 2020), butterflies
(Phalnikar et al., 2018; Sanaei et al., 2024), and mosquitoes (Moll et al., 2001). For instance,
gut microbiome assembly in monarch butterflies mirrors the patterns we observed in flour

beetles, whereby the microbiome load is high in larvae and decreases in pupae and adults
(Sanaei et al., 2024). However, in most cases, the microbiome shift across insect life stages
is typically associated with a niche shift that can directly influence the microbiome by altering
the influx of bacteria and/or available nutrients. The case of *T. castaneum* is remarkable
because the microbiome shift is not associated with a diet or habitat change; in fact all life
stages share the same habitat that serves as the route for rapid fecal-oral transmission. We
speculate that in flour beetles, the life stage-specific variation may derive from differential
feeding rates (allowing, e.g., faster bacterial proliferation in larval guts as reported in (Mason
et al., 2023; Wang et al., 2020), distinct gut morphology (e.g., the larval gut may be more
suitable for bacterial growth(Engel & Moran, 2013)), or different immune responses (e.g.,
adults may have more effective immune function, suppressing bacterial growth in the gut
(Tate et al., 2021). Note that quinone secretion by adults probably does not contribute to
stage-specific microbiome variation, because quinones suppress overall and taxon-specific
microbial load in the flour, which should therefore influence available inocula for all life
stages. We hope that future work can distinguish between these causes of life stage-specific
variation in flour beetle microbiomes.
763
Here, we analysed broad patterns of variation in microbiota composition, demonstrating
significant ecological drift. However, this does not imply that community assembly is entirely
stochastic: indeed, we observed that a relatively small number of genera (such as *Izhakiella*
and *Enterococcus*) repeatedly dominated, indicating a role for deterministic factors. Thus,
several questions remain about factors that guide community assembly and maintenance —
which specific taxa are more vs. less likely to colonize, and under what conditions? In the
fruit fly, successful colonization by a focal species depended on stochasticity as well as the
identity and presence of other species, likely due to deterministic outcomes of inter-species
interactions (Jones et al., 2022). Similarly, processes and phenomena such as priority
effects, differential survival, differential competitive ability, and niche use may determine
precisely which taxa will dominate the beetle microbiome at a given time (**Fig 8**). In our case,
it remains unclear whether, why, and when *Izhakiella* is a strong colonizer; this requires
further work, which is currently challenging due to our inability to culture the beetle-
associated strain of *Izhakiella*. We also acknowledge that our current analysis is coarse-
grained, and further interesting patterns may emerge at the level of variation and dynamics
across bacterial strains, or at higher level functional classes. We hope to address these
issues in future work. Finally, in closing we note that even non-essential bacterial symbionts
may significantly influence host ecology and life history, e.g., by providing transient benefits
such as protection against parasites, predators, and pathogens (Łukasik & Kolasa, 2024). It
remains to be seen whether any of the observed colonizers of the flour beetle gut provide
the host with fitness benefits, or whether the community truly represents neutral assembly.
785

786 **ACKNOWLEDGEMENTS**
787
We thank Basabi Bagchi and other members of the Agashe lab for discussion and critical
comments on the manuscript; Dhanush A, Prapti Satpathy and Manjunath Reddy for
laboratory assistance, and Sampath Kumar and the insectary facility at the Tata Institute for
Genetics and Society for training on and access to their microinjection platform. We
acknowledge funding and support from the National Centre for Biological Sciences and the

Department of Atomic Energy, Government of India (Project Identification No. RTI 4006); the
DBT/Wellcome Trust India Alliance (award number IA/I/17/1/503091 to DA); and an SERB
National Postdoctoral Fellowship (PDF/2018/001441 to RD).
796
797 **AUTHOR CONTRIBUTIONS**
798
799 DA, PS, RD, SG, SS, SB designed the project; PS, HKG, SG, ShG, SS, SB and RD
800 conducted experiments; PS analysed data; DA and PS wrote the manuscript with input from
801 all authors; DA acquired funding.
802
803 **DATA AVAILABILITY**
804
805 All the raw sequencing data are available in the Sequence Read Archive (SRA) of the
806 National Centre for Biotechnology Information (NCBI) with SRA accession numbers (X)
807 under Bioproject (X). 16S rRNA Sanger sequenced data are available in the GenBank of the
808 National Centre for Biotechnology Information (NCBI) with accession numbers (X). All
809 processed data used for analysis are available on Figshare (X).
810
811
812 **TABLES**
813
814 **Table 1: Summary of experiments conducted, and corresponding figures.**
815

Experiment	Description	Figures
1a,b	Quantify temporal variation in microbiomes: compare microbiomes of adults from a single stock population stored at -80° C from 2015–2022	Fig 1A,D-F
1c	Quantify microbiome variation across independently evolving lineages: compare microbiomes of adults from independent stock population boxes founded by same parental stock, sampled at the same time	Fig 1B
2	Quantify microbiome variation across replicate population boxes: compare microbiomes of adult offspring across 3 replicate boxes set up using the same stock parents	Fig 1C, Fig 4D-F
3	Determine microbiomes as a function of external factors such as experimenter identity, air sterility, and fresh vs. heat-treated wheat flour	Fig 2
4a	Test for maternal microbiome transmission, comparing microbiomes of females and eggs	Fig 3A-B
4b	Test for maternal microbiome transmission, analysing microbiomes of dissected female eggs and ovaries	Fig 3D-E
4c	Test for consistent maternal microbiome transmission, comparing microbiomes of eggs dissected from ageing females	Fig 3F
5	Quantify effect of population density: comparing microbiomes of adult offspring in parental populations housed at two	Fig 4A-C

	different densities	
6	Quantify fecal microbiomes of larvae and adults across development	Fig 5A-B
7	Quantify effect of fecal microbial inoculum on adult microbiomes, comparing adults provided with fresh flour, flour enriched with larval vs. adult feces, or flour enriched with larval feces and added quinones	Fig 5C-D
8	Quantify effect of amount of larval feces on beetle microbiome: comparing microbiomes of adults fed with different proportions of larval feces in fresh flour	Fig 5E-F
9	Testing the sensitivity of beetle-associated bacterial strains to adult-secreted antimicrobial quinones	Fig 6A
10	Testing the effect of quinones on bacterial load in flour: comparing temporal change in flour microbial loads, after inoculating with lyophilized focal bacteria or larval or adult feces, with vs. without quinones	Fig 6B
11	Testing the effect of adult quinones on beetle microbiomes: comparing microbiomes of beetles with vs. without RNAi-mediated knockdown of quinone pathway	Fig 6C-D
12	Testing the effect of flour replacement on the flour microbiome: comparing flour microbiomes across two cycles (generations) for stock populations maintained under discrete or continuous generations.	Fig 7

    FIGURES

Fig 1: Beetle microbiome and bacterial load varies across time, individuals,
populations, and generations. Results from Experiments 1 and 2 are shown here. (A–C)
The relative abundance of bacterial ASVs (identified to genus level) for each female beetle
sampled from a population as a function of (A) time (B) replicate populations (three different
lineages) (C) replicate populations from the same stock population. (D) Taxonomic beta
diversity (β_{rc}) of bacterial communities across time, for samples shown in panel A. Positive
values indicate that samples are more dissimilar; negative values indicate that samples are
taxonomically more similar to each other. (E) Mean total bacterial load per beetle
(normalized to total amount of host DNA per sample) for a subset of samples shown in panel
A. The dotted grey line represents the overall average bacterial load across all samples,
excluding August 2020 samples. Error bars represent standard error. (F) Correlation
between the mean relative abundance of *Izhakiella* and the mean total bacterial load per
population, for samples shown in panel E. The trendline shown is a linear model fit with
R^2=0.22. Sample size (number of female beetles) is indicated in parentheses.

Fig 2: External environmental factors explain very little variation in the beetle
microbiome. Results of Experiment 3, testing the effects of external factors. (A) Bacterial
community composition in female beetles as a function of experimenter identity, air sterility,
and flour treatment; each factor was tested in separate experiments (see Fig S11 for results
for male beetles). (B) Taxonomic beta diversity (β_{rc}) of bacterial communities for samples
shown in panel A. (C) Mean total bacterial load (normalized to total host DNA) for a subset
of samples shown in panel A (n = 3 females/treatment). Error bars represent standard error.
For reference, the dotted grey line indicates the mean total bacterial load per beetle
observed for the samples shown in Fig 1E.
843

844
845

Fig 3: No evidence for consistent maternal microbiome transmission to eggs. (A)
Results from Experiment 4a, determining bacterial community composition in 2-week-old
mothers and their eggs. (B) Image showing an example of dissected ovaries and eggs from
a 2-week-old female. (C) Results of Experiment 4b, determining bacterial community
composition in ovaries dissected from 2-week-old females (O) and their eggs (E). (D)
Results of Experiment 4c determining the bacterial community composition in eggs
dissected from ovaries of females of different ages. Sample sizes are indicated in
parentheses below bars.

Fig 4: Feces are a major source of beetle microbiome inoculum. (A–C) Results of
Experiment 5 determining the bacterial community characteristics of the offspring of 2-week-
old females ovipositing at different adult densities (LD = low density; HD = high density; R1,
R2, R3 = replicates). (A) Mean total bacterial load (±se) (B) Bacterial community composition
(C) Taxonomic beta diversity (β_{rc}). (D–F) Results from Experiment 2 comparing bacterial
communities of females and their 14-day old larval offspring. (D) Mean total bacterial load
(±se). (E–F) Bacterial community composition (E) and Taxonomic beta diversity (β_{rc}) (F) of
adult females (M = mother) and 2 week old larvae (O = offspring). Sample sizes are
indicated in parentheses.
864

865
866

Fig 5: Source and amount of host fecal inoculum in flour alter the beetle microbiome.
(A–B) Results of Experiment 6 to determine the time course of mean total bacterial load (±se) and average relative abundance of bacterial taxa (shown as pie charts) in the feces of (A) larvae and (B) adults (n = 3 replicate tubes, each containing 30 larvae or adults) (see Fig S13A-B for microbiome composition). (C–D) Results of Experiment 7 to determine microbiome establishment in pupae exposed to flour enriched with larval or adult feces, or nothing (control), as a function of the age of eclosed adult females. (C) Median taxonomic beta diversity (β_{rc}) (see Fig S13C for microbiome composition) (ANOVA, P_{treat} = 0.79, P_{time} = 0.009, $P_{treat \times time}$ = 1.19e-06) (D) Mean total bacterial load and (E) relative abundance of *Izhakiella* (n = 6 females/treatment/age) (ANOVA, P_{treat} = 0.006, P_{time} = 0.003, $P_{treat \times time}$ = 0.0002). (F–G) Results of Experiment 8 to test the effect of amount of fecal inoculum on beetle microbiome. (F) Mean total bacterial load in the flour (n = 2 technical replicates/treatment) (ANOVA, P_{treat} = 0.03) and in beetles (n = 4 females/treatment) (ANOVA, P_{treat} = 0.003). Pie charts show the average bacterial community composition in 25% and 100% enriched flour (see Fig S13D for the full dataset). (G) Taxonomic beta diversity (β_{rc}) (see Fig S13E for full dataset) (Kruskal-Wallis tests, $P_{diversity}$ = 9.586e-06). Error bars represent standard error.

Fig 6: Effect of quinones on the adult microbiome. (A) Results of Experiment 9 to measure the antimicrobial effect of quinone extracts on several bacterial strains isolated from beetles (n = 3 technical replicates/strain). A larger zone of inhibition indicates greater sensitivity of the strain to quinones. The images show a dissected beetle with paired abdominal stink glands (blue arrows), and examples of agar plates with bacterial growth inhibition due to quinones added to filter paper discs (the bottom right disc in each plate serves as a solvent-only negative control). (B) Results of Experiment 10 showing the effect of quinones on mean (±se) total bacterial load in the flour (n = 8 technical replicates/treatment/timepoint). Results of ANOVA are shown for each panel. Pie charts show the average bacterial community composition; outer circles indicate treatment (solid line: without quinone; dotted line: with quinones) (see Table S2 and Fig S14A for data for each replicate). (C–D) Results of Experiment 11, testing the effect of suppressing quinones using RNAi for genes GT39 (quinone production) and GFP (negative control); with untreated beetles and injection with buffer serving as additional controls. (C) Mean (±se) total bacterial load in beetles (n = 4 females/treatment) (t test). (D) Taxonomic beta diversity (β_{rc}) of experimental beetles (see Fig S14C for microbiomes of individual females). Asterisks indicate significant differences between dsGT39 and dsGFP beetles (Wilcox test).

904
905
Fig 7: Effect of population maintenance protocol on the beetle microbiome. (A)
schematic showing the stock maintenance protocol. Blue arrows mark timepoints when the
flour is replaced with fresh flour. (B) Results of Experiment 12 showing the time course of
mean (±se) total bacterial load over time, in flour sampled every 5 days from stock
populations maintained under discrete generation cycles (n = 2 technical replicates per
timepoint). Pie charts show the average relative abundance of bacterial taxa in flour samples
(within the plot area) and in sampled beetles (under the x-axis). Schematics along the x-axis
indicate typical stage structure of the population at each timepoint. (C) Results of
Experiment 12, showing data as described for panel B, for populations maintained under
continuous generations. Fig S15 shows microbiome composition for each replicate in the
flour as well as for beetles sampled for populations in both panels B and C. (D) The
relationship between mean total bacterial load and relative abundance of *Izhakiella* in flour
samples for samples shown in panels B and C (ANOVA, load ~ relative abundance of
Izhakiella (rel) x population (pop); $P_{rel} < 0.0001$, $P_{pop} = 0.7$, $P_{rel\ x\ pop} = 0.9$).
920

Fig 8: Proposed model for key processes shaping the red flour beetle microbiome. We
consider three different aspects of host-bacterial interactions: bacterial inoculum and survival
in the flour, and host colonization. Life cycle transitions are shown with curved arrows. Blue
arrows indicate increased bacterial transmission; red indicates inhibitory effects; increasing
arrow thickness indicates stronger effects. Hypothesized factors with unknown effects (not
tested in this study) are marked by a box with dashed borders.
929

REFERENCES

Agarwal, A., & Agashe, D. (2020). The red flour beetle Tribolium castaneum: A model for host-microbiome interactions. *PLoS ONE*, *15*(10 October). https://doi.org/10.1371/journal.pone.0239051

Aizenberg-Gershtein, Y., Laviad, S., Samuni-Blank, M., & Halpern, M. (2016). Izhakiella capsodis gen. nov., sp. nov., in the family Enterobacteriaceae, isolated from the mirid bug Capsodes infuscatus. *International Journal of Systematic and Evolutionary Microbiology*, *66*(3), 1364–1370. https://doi.org/10.1099/ijsem.0.000886

Bi, J., Zhang, L., & He, Y. (2024). Metabolomics and microbiome provide new insights into mechanisms of the variations between the larva and adult stages of the Model Beetle Tribolium castaneum (Herbst). *Journal of Asia-Pacific Entomology*, *27*(2). https://doi.org/10.1016/j.aspen.2024.102241

Bruijning, M., Henry, L. P., Forsberg, S. K. G., Metcalf, C. J. E., & Ayroles, J. F. (2022). Natural selection for imprecise vertical transmission in host–microbiota systems. *Nature Ecology and Evolution*, *6*(1), 77–87. https://doi.org/10.1038/s41559-021-01593-y

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Chase, J. M., Kraft, N. J. B., Smith, K. G., Vellend, M., & Inouye, B. D. (2011). Using null models to disentangle variation in community dissimilarity from variation in α-diversity. *Ecosphere*, *2*(2). https://doi.org/10.1890/ES10-00117.1

CoreTeam, R. (2016). *R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.*

Deb, R., Nair, A., & Agashe, D. (2019). Host dietary specialization and neutral assembly shape gut bacterial communities of wild dragonflies. *PeerJ*, *2019*(11), 1–30. https://doi.org/10.7717/peerj.8058

Engel, P., & Moran, N. A. (2013). The gut microbiota of insects - diversity in structure and function. In *FEMS Microbiology Reviews* (Vol. 37, Issue 5, pp. 699–735). https://doi.org/10.1111/1574-6976.12025

Girard, M., Luis, P., Valiente Moro, C., & Minard, G. (2023). Crosstalk between the microbiota and insect postembryonic development. In *Trends in Microbiology* (Vol. 31, Issue 2, pp. 181–196). Elsevier Ltd. https://doi.org/10.1016/j.tim.2022.08.013

Gordon, C. S., Rajagopalan, N., Risseeuw, E. P., Surpin, M., Ball, F. J., Barber, C. J., Buhrow, L. M., Clark, S. M., Page, J. E., Todd, C. D., Abrams, S. R., & Loewen, M. C. (2016). Characterization of triticum aestivum abscisic acid receptors and a possible role for these in mediating fusairum head blight susceptibility in wheat. *PLoS ONE*, *11*(10). https://doi.org/10.1371/journal.pone.0164996

Goulda, A. L., Zhang Vivian, Lamberti Lisa, Jones Eric W., Obadia Benjamin, Korasidis Nikolaos, Gavryushkin Alex, Carlson Jean M., Ludington William B., & Beerenwinkel Niko. (2018). Microbiome interactions shape host fitness. *Proceedings of the National Academy of Sciences*, *115*(51), 11951–11960. https://doi.org/10.5061/dryad.2sr6316

Hammer, T. J., Janzen, D. H., Hallwachs, W., Jaffe, S. P., & Fierer, N. (2017). Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(36), 9641–9646. https://doi.org/10.1073/pnas.1707186114

Hammer, T. J., Sanders, J. G., & Fierer, N. (2019). Not all animals need a microbiome. In *FEMS Microbiology Letters* (Vol. 366, Issue 10). Oxford University Press. https://doi.org/10.1093/femsle/fnz117

Holt, J. R., Cavichiolli de Oliveira, N., Medina, R. F., Malacrinò, A., & Lindsey, A. R. I. (2024). Insect–microbe interactions and their influence on organisms and ecosystems. In *Ecology and Evolution* (Vol. 14, Issue 7). John Wiley and Sons Ltd. https://doi.org/10.1002/ece3.11699

Jones, E. W., Carlson, J. M., Sivak, D. A., & Ludington, W. B. (2022). Stochastic microbiome assembly depends on context. https://doi.org/10.1073/pnas.2115877119/-/DCSupplemental

Khan, I., Prakash, A., Umarani, M., Sasidharan, R., Masagalli, J. N., Lama, P., Venkatesan, R., & Agashe, D. (2017). *Female Density-Dependent Chemical Warfare Underlies Fitness Effects of Group Sex Ratio in Flour Beetles.* https://doi.org/10.5061/dryad.p9v3q

Koga, R., Moriyama, M., Onodera-Tanifuji, N., Ishii, Y., Takai, H., Mizutani, M., Oguchi, K., Okura, R., Suzuki, S., Gotoh, Y., Hayashi, T., Seki, M., Suzuki, Y., Nishide, Y., Hosokawa, T., Wakamoto, Y., Furusawa, C., & Fukatsu, T. (2022). Single mutation makes Escherichia coli an insect mutualist. *Nature Microbiology*, *7*(8), 1141–1150. https://doi.org/10.1038/s41564-022-01179-9

Leftwich, P. T., Edgington, M. P., & Chapman, T. (2020). Transmission efficiency drives host-microbe associations. In *Proceedings of the Royal Society B: Biological Sciences* (Vol. 287, Issue 1934). Royal Society Publishing. https://doi.org/10.1098/rspb.2020.0820

Lisa M. Unruh, Rongda Xu, & Karl J. Kramer. (1998). Benzoquinone levels as a function of age and gender of the red flour beetle, Tribolium castaneum. *Insect Biochemistry and Molecular Biology*.

Łukasik, P., & Kolasa, M. R. (2024). With a little help from my friends: The roles of microbial symbionts in insect populations and communities. In *Philosophical Transactions of the Royal Society B: Biological Sciences* (Vol. 379, Issue 1904). Royal Society Publishing. https://doi.org/10.1098/rstb.2023.0122

Lundberg, D. S., Yourstone, S., Mieczkowski, P., Jones, C. D., & Dangl, J. L. (2013). Practical innovations for high-throughput amplicon sequencing. *Nature Methods*, *10*(10), 999–1002. https://doi.org/10.1038/nmeth.2634

Manthey, C., Johnston, P. R., Nakagawa, S., & Rolff, J. (2023). Complete metamorphosis and microbiota turnover in insects. *Molecular Ecology*, *32*(23), 6543–6551. https://doi.org/10.1111/mec.16673

Markarian, H., Florentine, G. J., & Pratt, J. J. (1978). Quinone production of some species of tribolium. In *J. Insecr Phxsiol* (Vol. 24). Pergamon Press Ltd.

Mason, C. J., Auth, J., & Geib, S. M. (2023). Gut bacterial population and community dynamics following adult emergence in pest tephritid fruit flies. *Scientific Reports*, *13*(1). https://doi.org/10.1038/s41598-023-40562-2

McFall-Ngai, M. (2024). Symbiosis takes a front and center role in biology. *PLoS Biology*, *22*(4). https://doi.org/10.1371/journal.pbio.3002571

McMurdie, P. J., & Holmes, S. (2015). Shiny-phyloseq: Web application for interactive microbiome analysis with provenance tracking. *Bioinformatics*, *31*(2), 282–283. https://doi.org/10.1093/bioinformatics/btu616

Miller, E. T., Svanbäck, R., & Bohannan, B. J. M. (2018). Microbiomes as Metacommunities: Understanding Host-Associated Microbes through Metacommunity Ecology. In *Trends in Ecology and Evolution* (Vol. 33, Issue 12, pp. 926–935). Elsevier Ltd. https://doi.org/10.1016/j.tree.2018.09.002

Moll, R. M., Romoser, W. S., Modrzakowski, M. C., Moncayo, A. C., & Lerdthusnee, K. (2001). Meconial Peritrophic Membranes and the Fate of Midgut Bacteria During Mosquito (Diptera: Culicidae) Metamorphosis. In *J. Med. Entomol* (Vol. 38, Issue 1). https://academic.oup.com/jme/article/38/1/29/1005225

Naik, T., Sharda, M., Lakshminarayanan, C. P., Virbhadra, K., & Pandit, A. (2023). High-quality single amplicon sequencing method for illumina MiSeq platform using pool of 'N' (0–10) spacer-linked target specific primers without PhiX spike-in. *BMC Genomics*, *24*(1). https://doi.org/10.1186/s12864-023-09233-4

Ning, D., Deng, Y., Tiedje, J. M., & Zhou, J. (2019). A general framework for quantitatively assessing ecological stochasticity. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(34), 16892–16898. https://doi.org/10.1073/pnas.1904623116

Nobles, S., & Jackson, C. R. (2020). Effects of life stage, site, and species on the dragonfly gut microbiome. *Microorganisms*, *8*(2). https://doi.org/10.3390/microorganisms8020183

Obadia, B., Güvener, Z. T., Zhang, V., Ceja-Navarro, J. A., Brodie, E. L., Ja, W. W., & Ludington, W. B. (2017). Probabilistic Invasion Underlies Natural Gut Microbiome Stability. *Current Biology*, *27*(13), 1999-2006.e8. https://doi.org/10.1016/j.cub.2017.05.034

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Mcglinn, D., Minchin, P. R., Hara, R. B. O., & Simpson, G. L. (2019). *vegan: Community Ecology Package.R package version 2.5-4*. https://CRAN.R-project.org/package=vegan.

Pedrini, N., Ortiz-Urquiza, A., Huarte-Bonnet, C., Fan, Y., Juárez, M. P., & Keyhani, N. O. (2015). Tenebrionid secretions and a fungal benzoquinone oxidoreductase form competing components of an arms race between a host and pathogen. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(28), E3651–E3660. https://doi.org/10.1073/pnas.1504552112

Phalnikar, K., Kunte, K., & Agashe, D. (2018). Dietary and developmental shifts in Subject Category⬜: *Royal Society Open Science*, *5*, 171559.

Prakash, A., Agashe, D., & Khan, I. (2023). Alteration of diet microbiota limits the experimentally evolved immune priming response in flour beetles, but not pathogen resistance. *Journal of Evolutionary Biology*, *36*(12), 1745–1752. https://doi.org/10.1111/jeb.14213

Prendeville, H. R., & Stevens, L. (2002). MICROBE INHIBITION BY Tribolium FLOUR BEETLES VARIES WITH BEETLE SPECIES, STRAIN, SEX, AND MICROBE GROUP. In *Journal of Chemical Ecology* (Vol. 28, Issue 6). http://www.ars.usda.gov/is/np/systematics/

Sanaei, E., Chavez, J., Harris, E. V., Alcaide, T. Y., Baffour-Addo, K., Bugay, M. J., Adams, K. L., Zelaya, A., De Roode, J. C., & Gerardo, N. M. (2024). Microbiome analysis of monarch butterflies reveals effects of development and diet. *FEMS Microbiology Ecology*, *100*(12). https://doi.org/10.1093/femsec/fiae143

Sanjenbam, P., & Agashe, D. (2024). Divergence and convergence in epiphytic and endophytic phyllosphere bacterial communities of rice landraces. *MSphere*, *9*(11). https://doi.org/10.1128/msphere.00765-24

Sawada, M., Sano, T., Hanakawa, K., Sirasoonthorn, P., Oi, T., & Miura, K. (2020). Benzoquinone synthesis-related genes of Tribolium castaneum confer the robust antifungal host defense to the adult beetles through the inhibition of conidial germination on the body surface. *Journal of Invertebrate Pathology*, *169*. https://doi.org/10.1016/j.jip.2019.107298

Schmidt, K., & Engel, P. (2021). Mechanisms underlying gut microbiota-host interactions in insects. In *Journal of Experimental Biology* (Vol. 224, Issue 2). Company of Biologists Ltd. https://doi.org/10.1242/jeb.207696

Tate, A. T., Perry, A., & Jent, D. (2021). Larvae and adults exhibit contrasting patterns of immune gene expression and infection resistance in wild flour beetle populations. *Ecological Entomology*, *46*(5), 1230–1235. https://doi.org/10.1111/een.13066

Knorr Eileen, Schmidtberg Henrike, Arslan Derya, Bingsohn Linda, & Vilcinskas Andreas. (2015). Translocation of bacteria from the gut to the eggs triggers maternal transgenerational immune priming in Tribolium castaneum _ Enhanced Reader. *Biology Letters*, *11*.

Vega, N. M., & Gore, J. (2017). Stochastic assembly produces heterogeneous communities in the Caenorhabditis elegans intestine. *PLoS Biology*, *15*(3). https://doi.org/10.1371/journal.pbio.2000633

Wang, M., Xiang, X., & Wan, X. (2020). Divergence in gut bacterial community among life stages of the rainbow stag beetle Phalacrognathus muelleri (Coleptera: Lucanidae). *Insects*, *11*(10), 1–13. https://doi.org/10.3390/insects11100719

Zapién-Campos, R., Sieber, M., & Traulsen, A. (2020). Stochastic colonization of hosts with a finite lifespan can drive individual host microbes out of equilibrium. *PLoS Computational Biology*, *16*(11). https://doi.org/10.1371/journal.pcbi.1008392

Zheng-Yan, W., & Yang Meng-Ting LU Yu-Jie, W. H. (2020). Diversity of intestinal bacteria
in different geographic populations of Tribolium castaneum (Coleoptera:
Tenebrionidae). *Chinese Journal of Applied Entomology*, *2020*(3), 617–622.
https://doi.org/10.7679/j.issn.2095–1353.2020.062
Zhou, J., & Ning, D. (2017). *Stochastic Community Assembly: Does It Matter in Microbial*
Ecology? https://doi.org/10.1128/MMBR
1100
1101