Mapping prefrontal afferents along development

Anne Günther[a*], Malte Stelzer[a*], Andreas Franzelin[b], Annette Marquardt[a], Thomas Oertner[b],
Ileana L. Hanganu-Opatz[a#], Jastyn A. Pöpplau[a#]

[a]Institute of Developmental Neurosciences, Center for Molecular Neurobiology Hamburg,
Hamburg Center of Neuroscience, University Medical Center Hamburg-Eppendorf,
Falkenried 94, 20251, Hamburg, Germany

[b]Institute of Synaptic Physiology, Center for Molecular Neurobiology Hamburg, Hamburg
Center of Neuroscience, University Medical Center Hamburg-Eppendorf, Falkenried 94,
20251, Hamburg, Germany

[*]These authors share first authorship
[#]These authors share last authorship
12

Corresponding authors: Jastyn A. Pöpplau
                             jastyn.poepplau@zmnh.uni-hamburg.de
                             Falkenried 94, 20251 Hamburg, Germany
                             Lead contact

                             Ileana L. Hanganu-Opatz
                             hangop@zmnh.uni-hamburg.de
                             Falkenried 94, 20251 Hamburg, Germany

13
Running title: Development of prefrontal afferents

Keywords: Retrograde labeling, development, neonatal, adolescence, mPFC,corpus
callosum, contralateral, myelin, prefrontal afferents

6 Figures
1 Supplementary figure
2 Supplementary tables
20

Abstract

The medial prefrontal cortex (mPFC) acts as a hub for cognitive, emotional, and social
processes, integrating neuronal inputs from numerous brain regions. Knowledge on the
processes underlying formation and myelination of prefrontal afferents is essential for a
comprehensive understanding of the emergence, maturation, and refinement of mPFC-
dependent behaviors. However, despite detailed investigation of long-range projections to the
mPFC in the adult brain, little is known about their maturation. Here, we use retrograde
labeling, light-sheet imaging, and automatic cell segmentation to quantify afferent projections
to the mPFC from neonatal to adult age. We show that densities of ipsi- and contralateral
prefrontal afferents change along development from a widespread bilateral distribution at
neonatal age to a predominantly ipsilateral organization. Furthermore, myelination of
interhemispheric prefrontal afferents starts only after the decline of neonatal contralateral
projections and is followed by temporally elevated projection densities from limbic brain
regions. Overall, these distinct developmental dynamics of prefrontal afferents might have
major implications for the maturation of mPFC-dependent functional and behavioral outputs.

1. Introduction

The mPFC, a central hub of integrative processing in the brain, is involved in a wide range of
cognitive, emotional, and social functions. It integrates direct axonal inputs from numerous
cortical and subcortical regions through local and long-range circuits to guide behavior
(Hoover and Vertes, 2007; Ährlund-Richter et al., 2019; Hanganu-Opatz et al., 2023). Within
the PFC, functions are anatomically and physiologically compartmentalized, with medial
subdivisions (*i.e.*, anterior cingulate (ACA), prelimbic (PL), and infralimbic (ILA) areas)
primarily contributing to sensorimotor functions, action-planning, and decision making, while
ventral regions (*i.e.*, orbitofrontal cortex (ORB)) as well as adjacent cortices (*i.e.*, secondary
motor cortices (MOs)) control emotional and motivational behaviors, respectively
(Heidbreder and Groenewegen, 2003; Laubach et al., 2018; Anastasiades and Carter, 2021).
In contrast, long-range inputs to the mPFC provide essential information on environmental
context and internal state from other regions (Spellman et al., 2015; Burgos-Robles et al.,
2017). For instance, inputs from the ventromedial thalamus (VM) to the mPFC regulate
arousal states (Honjoh et al., 2018), while hippocampal projections contribute to working,
recognition, and episodic memory (MacDonald et al., 2011; Takehara-Nishiuchi, 2014; Chao
et al., 2017, 2020).

However, not only ipsilateral projections shape prefrontal circuits. Interhemispheric
connectivity has been shown to increase the complexity of prefrontal networks (Floresco and
Ghods-Sharifi, 2007; Chao et al., 2016; Wang et al., 2023). Numerous studies have
investigated the projections crossing the corpus callosum (CC) in rodents and humans (Yorke
Jr. and Caviness Jr., 1975; van der Knaap and van der Ham, 2011; Innocenti et al., 2022).
While callosal connectivity has been traditionally assumed to be predominantly homotopic
(*i.e.,* mPFC to mPFC) (Yorke Jr. and Caviness Jr., 1975; De Benedictis et al., 2016), more
recent studies have revealed substantial heterotopy of these projections (Szczupak et al.,
2023). Notably, specific functions of homotopic and heterotopic callosal connections have
been described in rodents. On the homotopic level, sensory processing has been shown to be
influenced by contra- and ipsilateral ACA projections (Wang et al., 2023), whereas
disruptions of heterotopic pathways between basolateral amygdala (BLA) and ACA lead to
impaired decision making (Floresco and Ghods-Sharifi, 2007).

In contrast to this detailed dissection of adult prefrontal afferents, little is known about
their development. Projections from other brain regions reach the mPFC during development
and are thought to undergo dynamic reorganization during early life to promote maturation of
higher-order cognitive (*e.g.*, working memory) and emotional functions (*e.g.*, fear learning)
(Chini and Hanganu-Opatz, 2021). For example, projections from the mediodorsal thalamus
(MD) and from the BLA progressively increase in density during development and modulate
the maturation of local prefrontal circuits (Cunningham et al., 2002; Maren and Quirk, 2004;
Rios and Villalobos, 2004; Parnaudeau et al., 2018; Klune et al., 2021). However, prior
studies along development have focused on isolated pathways and time points, precluding a
large-scale age-dependent mapping of afferents.

In this study, we filled this knowledge gap and systematically monitored the
development of axonal projections to the mPFC from birth to adulthood. Using retrograde
viral tracing in combination with light-sheet microscopy, we generated a comprehensive atlas

of distinct regional afferent projection patterns to the mPFC during early development. Based
on these results, we used the same retrograde viral tracing approach combined with confocal
microscopy to apply automatic cell segmentation for quantification of projecting cells from
different regions of the brain along the entire development. Additionally, we assessed the
degree of myelination of retrogradely labeled inputs using immunohistochemical staining of
myelin. Our data provide a comprehensive, brain-wide (*i.e.*, ACA, PL, IL, ORB, MOs,
agranular insular cortex (AI), piriform cortex (PIRI), BLA, basomedial amygdala (BMA),
retrosplenial cortex (RSP), entorhinal cortex (ENT), ventral and intermediate CA1 (CA1v,
CA1i), ventral and dorsal subiculum (SUBv, SUBd), claustrum (CLA), anterior nuclei
(ATN), VM, MD, supramammillary nucleus (SUM), periaqueductal gray (PAG), and ventral
tegmental area (VTA)) developmental overview, revealing distinct, region- and age-specific
dynamics of prefrontal afferent and giving crucial insights into the evolving architecture of
prefrontal circuits.

2. STAR Methods

2.1. Key resource table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Myelin Basic Protein antibody	Abcam	Cat# ab40390, RRID: AB_114152
Goat-Anti-Rabbit IgG (H&L) DyLight 633	Novus	Cat# NBP1-76057, RRID: AB_11059121
Bacterial and virus strains		
rAAV9_CAG-eGFP	Addgene	Addgene viral prep #37825
rAAV9rg_mCaMKII-mCherry	Addgene	Addgene viral prep #114469-AAVrg
Experimental models: Organisms/strains		
Mouse: C57Bl/6J	Universitätsklinikum Hamburg-Eppendorf – Animal facility	Mouse: C57Bl/6J
Software and algorithms		
Matlab 2017b, 2021b,	MathWorks	RRID:SCR_001622
ImageJ 1.52b	ImageJ	RRID:SCR_003070
R Statistical Software 4.2	R Project for Statistical Computing	RRID:SCR_001905
Anaconda 1.9.6	Conda	RRID:SCR_018317
Cellpose2	https://www.cellpose.org/	https://www.cellpose.org/
Custom analysis code	This paper	Available upon request
Other		

Confocal microscope	Zeiss	LSN700
Confocal microscope	Olympus	F1000

2.2. Animals

96
All experiments were performed in compliance with German law and guidelines of the
European Union for the use of animals in research (European Union Directive 2010/63/EU)
and were approved by the local ethical committee (Behörde für Gesundheit und
Verbraucherschutz Hamburg, N19/121, N23/110). Experiments were carried out in C57BL/6J
mice of both sexes. Animals were housed in a 12 h light/12 h dark cycle and were given
access to water and food *ad libitum*. The day of birth was considered postnatal day (P) 0.

2.3. Virus injections

Viral vectors were injected unilaterally into the PFC of the right hemisphere of C57BL/6J
mice. Two viral vectors were applied simultaneously: rAAV9rg_mCaMKII-mCherry
(#114469-AAVrg, Addgene, Watertown, MA, United States) for retrograde labeling of
excitatory projection-neurons and rAAV9_CAG-eGFP (#37825 Addgene, Watertown, MA,
United States) for local labeling of cells at the injection site. Viral vectors were prepared in a
genome copy ratio of 1:2.5 and injected at the same time using a stereotax (Kostka and
Bitzenhofer, 2022) (Figure 1A). Viral vectors were injected at four different time points, for
which volumes and coordinates were adjusted to compensate for brain volume to result in a
comparable virus spread across development, *i.e.* P0 (neonatal, 130 nl at a rate of 120 nl/min,
0.3 mm posterior to the inferior cerebral vein, 0.2 mm lateral to the midline and 0.4-0.8 mm
ventral from the brain surface), P10 (pre-juvenile, 170 nl at a rate of 120 nl/min, 1,6 mm
anterior to Bregma, 0.2 mm lateral to midline and 0,8-1,2 mm ventral from the brain surface),
P20 (juvenile, 190 nl at a rate of 120 nl/min, 1,8 mm anterior to Bregma, 0.2 mm lateral to
midline and 1-1,4 mm ventral from the brain surface), and P50 (adult, 210 nl at a rate of 120
nl/min, 1,9 mm anterior to Bregma, 0.2 mm lateral to midline and 1,1-1,5 mm ventral from
the brain surface). Mice were transcardially perfused with 4% paraformaldehyde (PFA) 10
days after virus injection, *i.e.* at P10 (neonatal, n=6), P20 (pre-juvenile, n=6), P30 (juvenile,
n=6), and P60 (adult, n=6). To further account for differences of viral spread at the injection
site, brains were manually categorized as low spread, normal spread, or high spread. Low
spread was defined as expression in only one mPFC subdivision or a very low degree of
labeled cell bodies in more than one subdivision (neonatal n=1, pre-juvenile n=1). High
spread was defined as labeled cell bodies in all three subdivisions of the mPFC, while also
slightly exceeding mPFC boundaries on the dorsal-ventral or anterior-posterior axis (neonatal
n=1, pre-juvenile n=1, juvenile n=1, adult n=1). Accordingly, normal spread was defined as
expression in at least two subdivisions without exceeding mPFC boundaries (neonatal n=4,
pre-juvenile n=4, juvenile n=5, adult n=5). Virus spread was included in statistical analysis
(see Statistical analysis, Supplementary discussion, and Statistics table S1&2).

2.4. Tissue clearing and light-sheet imaging

Light-sheet imaging on an open source mesoSPIM (Voigt et al., 2019) was used to generate
brain-wide overviews of mPFC-projecting brain regions at neonatal (injected at P0, perfused
at P10) and pre-juvenile (injected at P10, perfused at P20) age. Perfusion-fixed brains were

cleared with SHIELD reagent (Park et al., 2019) to preserve endogenous fluorescence and
imaged in EasyIndex solution (RI = 1.465, Life Canvas Technologies) with an Orca Flash 4.0
camera (Hamamatsu).

2.5. Cell quantification

For confocal microscopy, brains were prepared in coronal 100 µm sections and mounted on
slides using Vectashield with DAPI (Vector Laboratories, Burlingame, CA, United States).
For cell quantification, images were obtained using a confocal microscope (LSN700, Zeiss,
Germany) with a 20x objective (2,048 x 2,048 pixels; 319.5 x 319.5 µm). Areas were
demarcated according to the coronal P56 mouse reference atlas from the Allen Brain Atlas
(mouse.brain-map.org). Additionally, images of mice injected at P0 were also compared to a
developing mouse brain atlas (Paxinos et al. 2007). Images of three consecutive coronal brain
slices were taken for each brain region in the ipsi- and the contralateral hemispheres. Images
were segmented with Cellpose 2 in Anaconda (Python 3.8.) using both the standard model
algorithm 'cyto2' for the images of cells transduced by rAAV9rg_CaMKII-mCherry and a
self-trained algorithm as described for the human-in-the-loop approach (Pachitariu and
Stringer, 2022) for rAAV9_CAG-eGFP-transduced cells. After automatic cell segmentation,
all images were validated by visual inspection. Confocal images from 22 prefrontal and long-
range projecting brain sections with mCherry-labeled pyramidal neurons were quantified as a
measure of projection strength to the mPFC across development. Images of eGFP-labeled
cells were acquired from prefrontal areas at the injection site (ACA, PL, ILA, ORB, MOs) for
normalization of quantification of retrogradely labeled projections. Normalized cell count
(NCC) was calculated per brain area using mCherry-labeled cells of a single image from each
brain area of an animal (r) and all eGFP-labeled cells of an animal (g☐) as: $NCC = \frac{r}{r+\bar{g}}$.
Additionally, the mean cell count (x) was calculated as the mean of mCherry-labeled cells
from all sections of one region for all animals from all age groups.

2.6. Immunohistochemistry

Immunohistochemical staining of myelin basic protein (MBP) at the CC was performed on
the corresponding sections of injected brains after perfusion at P10 (n=4), P20 (n=3), P30
(n=3), and P60 (n=3) as previously described (Parbst et al., 2025). Briefly, brain slices were
permeabilized using 0.8% Triton X-100 (Sigma-Aldrich, MO, USA) and blocked with 5%
normal goat serum and 5% normal donkey serum (Jackson Immuno Research, PA, USA) for
1 h. Samples were incubated overnight at 4°C with an MBP antibody (1:500, Abcam
ab40390). As a secondary antibody Goat-Anti-Rabbit IgG (H&L) DyLight 633 (1:1000,
Novus, NBP1-76057) was used.

2.7. Myelin quantification

For myelin quantification, images were obtained using a confocal microscope (F1000,
Olympus, Tokyo, Japan) with a 60x objective and 2x zoom (1024 x 1024 pixels; 105.47 x
105.47 mm). Immunohistochemically stained samples from injected brains were used to
image axonal fibers and myelin at the mid-section of the CC. For every animal, 3-4 stacks
from different brain slices (10-30 single images, 150 nm interval) were acquired. After
preprocessing and thresholding in ImageJ, colocalization of MBP-positive and mCherry-

labeled structures expression was calculated in Matlab R2023a using a custom-written
algorithm similar to (Pöpplau et al., 2024). Values were normalized for stack depth.

2.8. Statistical analysis

Statistical analysis was carried out in R using linear mixed effect models (LME) of the 'lme4'
package in R 4.4.1. We assessed the impact of *age* (fixed effect) on NCCs, while accounting
for random effects, including *sex*, *viral spread,* and *image number* to consider for multiple
image-based measurements per *individual animal*. Analyzed image stacks of myelinated
structures, labeled axonal fibers, and their colocalization were similarly analyzed with LME
models to determine the effect of fixed and random effects on myelinization of axons.
Briefly, likelihood ratio tests were performed to compare log-likelihoods between the full and
the reduced model. To test for a significant effect of age-group (fixed effect), sex, or viral
spread (random effects), these parameters were removed from the full model to calculate
statistical significance using the 'anova' function. Post-hoc pairwise comparisons between
age-groups were conducted using the Tukey's Honest Significant Difference test using the
'multcomp' package. P-values were adjusted using the Holm method 'holm' to account for
multiple comparisons (see Statistics table S1).

3. Results

3.1. mPFC afferents show age-dependent trajectories

Resolving the developmental dynamics of long-range projections to the mPFC from different
brain regions and between hemispheres is the basis for understanding how prefrontal circuits
gain and develop their functions. In a first step, we assessed the maturation of prefrontal
afferents during early development using light-sheet microscopy on cleared brain tissue. For
this, we injected rAAV9rg_mCaMKII-mCherry and rAAV9_CAG-eGFP unilaterally into the
mPFC at two different time points (*i.e.,* neonatal P0 and pre-juvenile P10). Tissue was
perfused and cleared 10 days after injection (*i.e.,* neonatal P10, pre-juvenile P20) (Figure 1A,
B). Whole-brain imaging revealed retrogradely labeled cells in multiple cortical and
subcortical brain regions of both hemispheres at both investigated time points (Figure 1B).
Based on this qualitative overview, we chose 22 brain regions with direct axonal projections
to the mPFC along development to analyze in detail. We also included the PAG and VTA
(Figure 1B), that showed prominent glutamatergic afferent projections during development
which is not reported for the adult brain (Hoover and Vertes, 2007; Ährlund-Richter et al.,
2019). Next, we used confocal microscopy in slices of injected brains across the entire
development (*i.e.,* neonatal P0, pre-juvenile P10, juvenile P20, adult P50). Subsequent cell
segmentation (Figure 1C-F) to assess projection strength by calculating the NCC (normalized
cell count: calculated as ratio of mCherry-labeled cells from each brain area *vs*. all eGFP- and
mCherry-labeled cells, see Materials and Methods section) in prefrontal subdivisions (ACA,
PL, ILA, ORB, MOs) as well as in cortical (AI, PIRI, BLA, BMA, RSP, ENT, CA1v, CA1i,
SUBv, SUBd) and subcortical areas (CLA, ATN, VM, MD, SUM, PAG, VTA), projecting
ipsi- and contralaterally to the mPFC.

3.2. Medial but not adjacent local prefrontal connectivity increases along development

In order to assess local prefrontal connectivity, we quantified retrogradely labeled cells in prefrontal subdivisions of the ipsilateral hemisphere (marked as [i] for all regions). The density of labeled cells in mPFC and adjacent subdivisions changed with age but was not influenced by sex or viral spread (Figure 2A-E). For the ACA[i] and PL[i] the NCC increased from neonatal to pre-juvenile age and remained stable afterwards (Figure 2A,B). Solely ILA[i] local connectivity peaked at juvenile age and decreased in adults (Figure 2C). In contrast, the NCC in adjacent MOs[i] and ORB[i] did not significantly change with age (Figure 2D,E). Taken together, these data indicate that especially PL and ILA become strongly integrated into local prefrontal connectivity during development.

3.3. Cortical long-range afferents to the mPFC peak at (pre-)juvenile age

Next, we quantified the densities of long-range afferents from ipsilateral cortical (RSP[i], PIRI[i], AI[i], ENT[i], SUBv[i], SUBd[i], CA1i[i], and CA1v[i]) as well as cortical adjacent regions (BLA[i], BMA[i], CLA[i]). Sex of investigated mice or viral spread had no confounding effects on the results. In most regions, with exception of AI[i] that showed a consistently high density of afferents, and CA1i[i] (Figure 3A,B), prefrontal afferents revealed distinct age-dependent patterns. The densities of projections from BMA[i], CLA[i], and BLA[i] strongly increased from neonatal to pre-juvenile age and showed only slight changes later on (Figure 3C-E). Notably, BLA[i]-to-mPFC afferents were nearly absent at neonatal age, emerging towards pre-juvenile age. A similar pattern, though less striking, was also detected for RSP[i], ENT[i], and SUBv[i] (Figure 3F-I,K). However, the densities of these prefrontal afferents decreased during later development, resulting in similar values when comparing neonatal and adult mice. A rather unusual developmental dynamic was detected for SUBd[i] with dense projections to the mPFC at neonatal age that almost disappeared at adult age (Figure 3J).

Thus, numerous ipsilateral projections target the mPFC during development and reveal distinct dynamics. Overall, long-range connectivity seems to be particularly dense during the pre-juvenile / juvenile period.

3.4. Prefrontal projections from thalamic and hypothalamic nuclei develop in an inverse manner

Next, we analyzed long-range prefrontal afferents arising from ipsilateral subcortical brain structures (ATN[i], VM[i], MD[i], SUM[i], PAG[i], VTA[i]). For the ATN[i], we quantified the interanterodorsal nucleus as well as ventral and dorsal sections of the AM. VTA[i]-to-mPFC projections remained unchanged throughout development (Figure 4A). In contrast, the density of VM[i] and ATN[i] afferents increased from neonatal to pre-juvenile age and remained stable afterwards, even though the density of projections at neonatal age was rather low in VM[i] and high in ATN[i] (Figure 4B,C). Notably, ATN[i] was the only region for which we observed an effect of viral spread which decreased with age (see Supplementary discussion, Statistics table S1&S2). The density of labeled cells in the MD[i] decreased with age, indicating a sparsification of prefrontal afferents (Figure 4D). Similarly, the density of SUM[i]-to-mPFC projections decreased from pre-juvenile to juvenile age with a significant effect of sex, revealing higher projection densities for male mice (Figure 4E, Statistics table S1&S2).

260 For the PAG[i], the density of retrogradely labeled cells was generally low but followed a
261 similar trend as most of the cortical areas and showed a peak at pre-juvenile age (Figure 4F).

262   Thus, most prefrontal afferents from subcortical regions reach high densities at pre-
263 juvenile age, yet their dynamics, either increasing or decreasing with age, are region-specific.
264

3.5. Contralateral prefrontal afferents undergo pruning along development

Adult mPFC has been reported to receive afferents from numerous contralateral regions
(marked as [c] for all regions) (Anastasiades and Carter, 2022). To clarify how these
projections evolve along development, we monitored contralateral prefrontal afferents from
birth to adulthood alongside ipsilateral projections. While no age-dependent changes of
prefrontal afferent density from contralateral ACA[c], ILA[c] and MOs[c] were detected (Figure
S1A), the number of labeled cells in the PL[c], similarly to the PL[i], increased from neonatal to
pre-juvenile age and remained stable afterwards (Figure 5A). In contrast, the number of
labeled cells in the ORB[c] decreased from neonatal to juvenile and adult age (Figure 5B).
Whereas projection density from ORB[i] showed no sex-dependent differences, an effect of sex
was detected for the ORB[c], the number of labeled cells being higher in male compared to
female mice (see Statistics table S2). Other contralateral regions were affected neither by sex
nor viral spread.

Based on their observed developmental dynamics, we classified contralateral long-
range afferents into four distinct groups. First, afferents that were present at birth and their
density remaining constant throughout development. The projections from ENT[c] and CLA[c]
are characteristic for this pattern (Figure 5C; S1A). Second, afferents that were present early
during development and strongly decrease later on. This patterns was detected for
contralateral afferents from MD[c], RSP[c], AI[c], ATN[c], SUBd[c], PIRI[c], SUM[c], and VTA[c] (Figure
5D, E; S1B, C). The RSP[c] was the only contralateral brain region for which we detected a
density increase of retrogradely-labeled cells from neonatal to pre-juvenile age followed by a
substantial decrease at older age (Figure 5E). Similar to the ORB[c], VTA[c] was one of the few
regions that showed an age-dependent change in contra- but not ipsilateral projections to
mPFC (Figure 5B; S1C). Third, the number of retrogradely labeled cells increased with age
in a similar manner as their corresponding ipsilateral sections. Afferents from BLA[c] and
BMA[c] belong to this group, despite the density differences between the neonatal and adult
group being minor (Figure 5F; S1E). Fourth, almost no prefrontal afferents were detected
during the entire development. This pattern was characteristic for CA1v[c], CA1i[c], SUBv[c],
VM[c], and PAG[c] (Figure S1D).

Taken together, these data reveal that the pruning of contralateral afferents along
development follow age- and region-specific patterns, resulting in denser ipsilateral
projections from most brain regions to the mPFC at adult age than those from the
contralateral hemisphere.
298

3.6. Myelination of contralateral prefrontal afferents emerges after neonatal age

In addition to monitoring the density of contralateral prefrontal afferents, we also assessed
their myelination. For this, we focused on the cortico-cortical projections crossing the CC.

The CC represents the largest white matter tract in the mammalian brain (Fitsiori et al., 2011;
De León Reyes et al., 2020; Szczupak et al., 2021) and it has been extensively studied at
adult age (Yorke Jr. and Caviness Jr., 1975; van der Knaap and van der Ham, 2011; De León
Reyes et al., 2020; Innocenti et al., 2022). We characterized the developmental dynamics of
myelination of contralateral prefrontal afferents in the previously defined age groups
(neonatal, pre-juvenile, juvenile, and adult) (Figure 6A). Overall myelin expression increased
with age (Figure 6B). Retrogradely labeled axons that colocalized with myelin were sparse in
neonatal mice, but their density strongly increased at pre-juvenile age and remained elevated
until adulthood (Figure 6C). Similarly, the portion of myelin colocalizing with labeled axons
increased from neonatal to adult age (Figure 6D).

4. Discussion

The PFC is a higher-order cortical region that is essential for complex cognitive and affective
processes such as decision making, working memory, and emotional regulation. Its
maturation is protracted when compared to primary sensory cortices (Chini and Hanganu-
Opatz, 2021), suggesting a prolonged time window for structural and functional refinement of
prefrontal circuits. In this study, we unraveled the developmental dynamics of local and long-
range ipsilateral and contralateral projections to the mPFC, with a specific focus on
developmental changes in projection strength from different regions (*i.e.*, ACA, PL, IL, ORB,
MOs, AI, PIRI, BLA, BMA, RSP, ENT, CA1v, CA1i, SUBv, SUBd, CLA, ATN, VM, MD,
SUM, PAG, and VTA) as well as the onset of myelination. Our findings reveal regionally
distinct maturational dynamics of prefrontal afferents that might underlie functional and
behavioral development.

4.1. Developmental shift from bilateral to unilateral projections to the mPFC

At birth, afferent projections to the mPFC are largely bilateral. However, this pattern shifts
progressively towards ipsilateral projections during maturation, with only a subset of cortical
and limbic areas retaining contralateral connectivity in adulthood (*i.e.*, ACA[c], PL[c], ILA[c],
MOs[c], ORB[c], AI[c], BLA[c], BMA[c], and ENT[c]). The most commonly observed dynamic of
contralateral projections was a decline of projection density after neonatal age (*i.e.*, ORB[c],
AI[c], PIRI[c], SUBd[c], ATN[c], MD[c], SUM[c]). While postnatal pruning of perinatal contralateral
projections for the ATN and MD has been reported (Minciacchi and Granato, 1989; De León
Reyes et al., 2019), our results imply that this unilaterality of prefrontal projections might
even be initiated before birth, as contralateral projections at neonatal age are already less
dense than ipsilateral ones.

Our data show that myelination of CC-crossing prefrontal afferents begins only after
the loss of most contralateral projections. In adult animals, the CC, is the largest white matter
tract in the mammalian brain (Fitsiori et al., 2011; De León Reyes et al., 2020) and contains
mostly cortico-cortical fibers (Szczupak et al., 2021), although evidence also suggests the
presence of cortico-subcortical connections (Milardi et al., 2015; De Benedictis et al., 2016).
In line with previous studies (Sturrock, 1980), we detected low amounts of myelin in the CC
and found that mPFC axonal fibers are largely unwrapped at neonatal age. As has been
previously shown (Paus et al., 1999; Fornari et al., 2007), myelination of prefrontal afferents
increases strikingly during the second postnatal week, co-occurring with weaning and the

onset of more complex behavioral abilities (Klune et al., 2021). As oligodendrocyte
engagement and axonal wrapping are promoted by activity-dependent mechanisms (de Faria
et al., 2019), the absence of myelin in early-stage interhemispheric fibers supports the idea
that these projections might lack the necessary activity for functional embedding into mature
circuits. In line with this hypothesis, the emergence of activity follows a protracted
development in the mPFC (Brockmann et al., 2011). This transient bilateral connectivity may
serve as a scaffolding mechanism for the maturation of local prefrontal microcircuits,
facilitating early synchronization across hemispheres. The question of whether early
contralateral prefrontal projections are functionally active remains to be addressed.
Alternatively, these projections could reflect developmental remnants of cerebral ontogenesis
that, in the absence of a specific function, are pruned with ongoing maturation.

In line with this, most ipsilaterally projecting regions show an overall increase of
input strength from neonatal to adult age, some areas (*e.g.*, MD[i] and SUBd[i]) demonstrate an
overall reduction in mPFC afferents, similar to their contralateral counterparts. This result
partially contradicts earlier qualitative, not statistically validated reports, suggesting a
rebound of MD input in adulthood (Rios and Villalobos, 2004). Given the well-established
role of the MD in working memory and cognitive functions (Parnaudeau et al., 2018),
developmental pruning of its projections to the mPFC appears paradoxical. However, pruning
of excessive, non-informative, or redundant projections might improve signal fidelity. Our
data suggest that this refinement of projections to the mPFC continues into adulthood,
potentially facilitating the emergence of precise, goal-directed behavior.

4.2.   Connectivity to the mPFC peaks during early adolescence

One of the most salient observations in our data is the pronounced increase of projections to
the mPFC during early adolescence (*i.e.*, pre-juvenile and juvenile group), coinciding with
the emergence of higher-order cognitive functions (Klune et al., 2021; Thies et al., 2025). To
this point, we could show that afferents from the limbic regions ENT[i], SUBv[i], CA1v[i], BLA[i],
BLA[c], and BMA[c] peak in projection strength during early adolescence. For example, the
SUBv is known to be involved in the reward circuit of the brain (Ding et al., 2020) and a role
of the CA1 in anticipation of consequences has been proposed (Fyhn et al., 2002; Numan,
2015). Thus, this temporal alignment of projections to the mPFC supports behavioral studies
that implicate adolescence as a sensitive window for risk taking and cognitive flexibility in
humans (van Duijvenvoorde et al., 2016; McCormick and Telzer, 2018) and in rodents
(Johnson and Wilbrecht, 2011; Klune et al., 2021; Thies et al., 2025). Similarly, projections
from the amygdala (*i.e.*, BLA and BMA of both hemispheres) are nearly absent in neonatal
mice but strongly increase towards pre-juvenile age. This developmental trajectory has been
previously described (Cunningham et al., 2002) and might be essential for the consolidation
of fear memories. Notably, the BLA has also been shown to play a role in decision-making
paradigms, in which effort and reward values have to be evaluated (Floresco and Ghods-
Sharifi, 2007). Moreover, increased projection strength from the AM (as part of the ATN[i]) to
the mPFC could facilitate dopamine-dependent reward processing, as its role in reinforcing
goal-directed behaviors has been shown in adult mice (Yang et al., 2022). Since the density
of dopaminergic receptors in the mPFC peaks during adolescence (Larsen and Luna, 2018),

the maturation of these inputs might contribute to the anatomical foundation for adaptive
decision-making and behavioral flexibility.

While some brain areas maintained their ipsilateral prefrontal innervations strength
after initial increase, other regions showed a decline of projections after early adolescence
(*i.e.*, PIRI[i], RSP[i], and SUM[i]). These brain regions are associated with olfactory processing,
spatial navigation, and episodic memory, all of which become particularly relevant during the
transition from maternal dependency to independent exploration (Vann et al., 2009; Ito et al.,
2018; Loureiro et al., 2019). The PIRI projects to the ORB as well as to the mPFC (Ährlund-
Richter et al., 2019; Bhattarai et al., 2022). These pathways have been reported to be relevant
for assessment of food safety (Loureiro et al., 2019). The SUM and RSP are involved in both
prefrontal and hippocampal networks and play a role in spatial navigation as well as episodic
memory (Vann et al., 2009; Ito et al., 2018). We hypothesize that transient innervation of the
early adolescent mPFC by these areas facilitates critical behavioral adaptations at the time of
weaning, including food selection and environmental orientation.

4.3.   Concluding remarks

In summary, our data reveal that the development of mPFC afferents is neither linear
nor uniform but characterized by region-specific dynamics of input formation, stabilization,
and pruning (see Supplementary discussion). These anatomical changes closely parallel
known milestones of behavioral and functional maturation, suggesting structural development
as a framework for functional circuit integration and adult-like cognitive functions. These
findings help to understand how early afferent maturation shapes the integrative functions of
the mPFC and how disruptions may result in lifelong lasting phenotypes, such as those
associated with neurodevelopmental disorders.

5. Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

6. CRediT authorship contribution

Conceptualization, supervision, and project administration: JAP, AG, and ILH-O.
Investigation: MS, AG, JAP, AF, and AM. Methodology and validation: AG and JAP. Data
curation, formal analysis, and visualization: MS, JAP, and AG. Writing – original draft: JAP,
AG, and MS. ILH-O, JAP, AG, and TO: Writing – review and editing. All authors have
approved the final version of the manuscript.

7. Funding

German Research Foundation FOR5159 TP1 (437610067) to ILH-O.
German Research Foundation FOR5159 start-up fund (437610067) to JAP
German Research Foundation (563625846) to I.L.H.-O.

8. Acknowledgments

We thank P. Putthoff and A. Dahlmann for excellent technical assistance.

9. Resource availability

Lead contact

Jastyn A. Pöpplau, jastyn.poepplau@zmnh.uni-hamburg.de

Materials availability

The study did not generate new unique reagents.

Data and code availability

All data needed to evaluate the conclusions are available in the figures, main text, or the
supplementary materials. Raw images are available from the lead contact upon request.
Custom image analysis code is available from the lead contact upon request.

10. Glossary

mPFC   medial prefrontal cortex
ACA   anterior cingulate cortex
PL   prelimbic cortex
IL   infralimbic cortex
ORB   orbitofrontal cortex
MO   secondary motor cortex
AI   agranular insular cortex
PIRI   piriform cortex (),
BLA   basolateral amygdala,
BMA   basomedial amygdala

453	RSP	retrosplenial cortex
454	ENT	entorhinal cortex
455	CA1v	ventral CA1
456	CA1i	intermediate CA1
457	SUBv	ventral subiculum
458	SUBd	dorsal subiculum
459	CLA	claustrum
460	ATN	anterior nuclei
461	VM	ventromedial thalamus
462	MD	mediodorsal thalamus
463	SUM	supramammillary nucleus
464	PAG	periaqueductal gray
465	VTA	ventral tegmental area
466	NCC	normalized cell count
467	rAAV	recombinant adeno-associated virus
468	P	postnatal day
469		

11. References.

Ährlund-Richter, S., Xuan, Y., van Lunteren, J. A., Kim, H., Ortiz, C., Pollak Dorocic, I., et
al. (2019). A whole-brain atlas of monosynaptic input targeting four different cell
types in the medial prefrontal cortex of the mouse. *Nat. Neurosci.* 22, 657–668. doi:
10.1038/s41593-019-0354-y

Anastasiades, P. G., and Carter, A. G. (2021). Circuit organization of the rodent medial
prefrontal cortex. *Trends Neurosci.* 44, 550–563. doi: 10.1016/j.tins.2021.03.006

Bhattarai, J. P., Etyemez, S., Jaaro-Peled, H., Janke, E., Leon Tolosa, U. D., Kamiya, A., et
al. (2022). Olfactory modulation of the medial prefrontal cortex circuitry:
Implications for social cognition. *Semin. Cell Dev. Biol.* 129, 31–39. doi:
10.1016/j.semcdb.2021.03.022

Brockmann, M. D., Pöschel, B., Cichon, N., and Hanganu-Opatz, I. L. (2011). Coupled
oscillations mediate directed interactions between prefrontal cortex and hippocampus
of the neonatal rat. *Neuron* 71, 332–347. doi: 10.1016/j.neuron.2011.05.041

Burgos-Robles, A., Kimchi, E. Y., Izadmehr, E. M., Porzenheim, M. J., Ramos-Guasp, W.
A., Nieh, E. H., et al. (2017). Amygdala inputs to prefrontal cortex guide behavior
amid conflicting cues of reward and punishment. *Nat. Neurosci.* 20, 824–835. doi:
10.1038/nn.4553

Chao, O. Y., de Souza Silva, M. A., Yang, Y.-M., and Huston, J. P. (2020). The medial
prefrontal cortex - hippocampus circuit that integrates information of object, place and
time to construct episodic memory in rodents: Behavioral, anatomical and
neurochemical properties. *Neurosci. Biobehav. Rev.* 113, 373–407. doi:
10.1016/j.neubiorev.2020.04.007

Chao, O. Y., Huston, J. P., Li, J.-S., Wang, A.-L., and de Souza Silva, M. A. (2016). The
medial prefrontal cortex-lateral entorhinal cortex circuit is essential for episodic-like
memory and associative object-recognition. *Hippocampus* 26, 633–645. doi:
10.1002/hipo.22547

Chao, O. Y., Nikolaus, S., Lira Brandão, M., Huston, J. P., and de Souza Silva, M. A. (2017).
Interaction between the medial prefrontal cortex and hippocampal CA1 area is
essential for episodic-like memory in rats. *Neurobiol. Learn. Mem.* 141, 72–77. doi:
10.1016/j.nlm.2017.03.019

Chini, M., and Hanganu-Opatz, I. L. (2021). Prefrontal Cortex Development in Health and
Disease: Lessons from Rodents and Humans. *Trends Neurosci.* 44, 227–240. doi:
10.1016/j.tins.2020.10.017

Cunningham, M. G., Bhattacharyya, S., and Benes, F. M. (2002). Amygdalo-cortical
sprouting continues into early adulthood: implications for the development of normal
and abnormal function during adolescence. *J. Comp. Neurol.* 453, 116–130. doi:
10.1002/cne.10376

De Benedictis, A., Petit, L., Descoteaux, M., Marras, C. E., Barbareschi, M., Corsini, F., et al.
(2016). New insights in the homotopic and heterotopic connectivity of the frontal

portion of the human corpus callosum revealed by microdissection and diffusion
tractography. *Hum. Brain Mapp.* 37, 4718–4735. doi: 10.1002/hbm.23339

de Faria, O., Gonsalvez, D. G., Nicholson, M., and Xiao, J. (2019). Activity-dependent
central nervous system myelination throughout life. *J. Neurochem.* 148, 447–461. doi:
10.1111/jnc.14592

De León Reyes, N. S., Bragg-Gonzalo, L., and Nieto, M. (2020). Development and plasticity
of the corpus callosum. *Dev. Camb. Engl.* 147, dev189738. doi: 10.1242/dev.189738

De León Reyes, N. S., Mederos, S., Varela, I., Weiss, L. A., Perea, G., Galazo, M. J., et al.
(2019). Transient callosal projections of L4 neurons are eliminated for the acquisition
of local connectivity. *Nat. Commun.* 10, 4549. doi: 10.1038/s41467-019-12495-w

Ding, S.-L., Yao, Z., Hirokawa, K. E., Nguyen, T. N., Graybuck, L. T., Fong, O., et al.
(2020). Distinct Transcriptomic Cell Types and Neural Circuits of the Subiculum and
Prosubiculum along the Dorsal-Ventral Axis. *Cell Rep.* 31, 107648. doi:
10.1016/j.celrep.2020.107648

Fitsiori, A., Nguyen, D., Karentzos, A., Delavelle, J., and Vargas, M. I. (2011). The corpus
callosum: white matter or terra incognita. *Br. J. Radiol.* 84, 5–18. doi:
10.1259/bjr/21946513

Floresco, S. B., and Ghods-Sharifi, S. (2007). Amygdala-Prefrontal Cortical Circuitry
Regulates Effort-Based Decision Making. *Cereb. Cortex* 17, 251–260. doi:
10.1093/cercor/bhj143

Fornari, E., Knyazeva, M. G., Meuli, R., and Maeder, P. (2007). Myelination shapes
functional activity in the developing brain. *NeuroImage* 38, 511–518. doi:
10.1016/j.neuroimage.2007.07.010

Fyhn, M., Molden, S., Hollup, S., Moser, M.-B., and Moser, E. (2002). Hippocampal neurons
responding to first-time dislocation of a target object. *Neuron* 35, 555–566. doi:
10.1016/s0896-6273(02)00784-5

Hanganu-Opatz, I. L., Klausberger, T., Sigurdsson, T., Nieder, A., Jacob, S. N., Bartos, M., et
al. (2023). Resolving the prefrontal mechanisms of adaptive cognitive behaviors: A
cross-species   perspective.   *Neuron*   111,   1020–1036.   doi:
10.1016/j.neuron.2023.03.017

Heidbreder, C. A., and Groenewegen, H. J. (2003). The medial prefrontal cortex in the rat:
evidence for a dorso-ventral distinction based upon functional and anatomical
characteristics.   *Neurosci.   Biobehav.   Rev.*   27,   555–579.   doi:
10.1016/j.neubiorev.2003.09.003

Honjoh, S., Sasai, S., Schiereck, S. S., Nagai, H., Tononi, G., and Cirelli, C. (2018).
Regulation of cortical activity and arousal by the matrix cells of the ventromedial
thalamic nucleus. *Nat. Commun.* 9, 2100. doi: 10.1038/s41467-018-04497-x

Hoover, W. B., and Vertes, R. P. (2007). Anatomical analysis of afferent projections to the
medial prefrontal cortex in the rat. *Brain Struct. Funct.* 212, 149–179. doi:
10.1007/s00429-007-0150-4

Innocenti, G. M., Schmidt, K., Milleret, C., Fabri, M., Knyazeva, M. G., Battaglia-Mayer, A.,
et al. (2022). The functional characterization of callosal connections. *Prog. Neurobiol.*
208, 102186. doi: 10.1016/j.pneurobio.2021.102186

Ito, H. T., Moser, E. I., and Moser, M.-B. (2018). Supramammillary Nucleus Modulates
Spike-Time Coordination in the Prefrontal-Thalamo-Hippocampal Circuit during
Navigation. *Neuron* 99, 576-587.e5. doi: 10.1016/j.neuron.2018.07.021

Johnson, C., and Wilbrecht, L. (2011). Juvenile mice show greater flexibility in multiple
choice reversal learning than adults. *Dev. Cogn. Neurosci.* 1, 540–551. doi:
10.1016/j.dcn.2011.05.008

Klune, C. B., Jin, B., and DeNardo, L. A. (2021). Linking mPFC circuit maturation to the
developmental regulation of emotional memory and cognitive flexibility. *eLife* 10,
e64567. doi: 10.7554/eLife.64567

Kostka, J. K., and Bitzenhofer, S. H. (2022). Postnatal Development of Centrifugal Inputs to
the Olfactory Bulb. *Front. Neurosci.* 16. doi: 10.3389/fnins.2022.815282

Larsen, B., and Luna, B. (2018). Adolescence as a neurobiological critical period for the
development of higher-order cognition. *Neurosci. Biobehav. Rev.* 94, 179–195. doi:
10.1016/j.neubiorev.2018.09.005

Laubach, M., Amarante, L. M., Swanson, K., and White, S. R. (2018). What, If Anything, Is
Rodent Prefrontal Cortex? *eNeuro* 5. doi: 10.1523/ENEURO.0315-18.2018

Loureiro, M., Achargui, R., Flakowski, J., Van Zessen, R., Stefanelli, T., Pascoli, V., et al.
(2019). Social transmission of food safety depends on synaptic plasticity in the
prefrontal cortex. *Science* 364, 991–995. doi: 10.1126/science.aaw5842

MacDonald, C. J., Lepage, K. Q., Eden, U. T., and Eichenbaum, H. (2011). Hippocampal
"Time Cells" Bridge the Gap in Memory for Discontiguous Events. *Neuron* 71, 737–
749. doi: 10.1016/j.neuron.2011.07.012

Maren, S., and Quirk, G. J. (2004). Neuronal signalling of fear memory. *Nat. Rev. Neurosci.*
5, 844–852. doi: 10.1038/nrn1535

McCormick, E. M., and Telzer, E. H. (2018). Not Doomed to Repeat: Enhanced Medial
Prefrontal Cortex Tracking of Errors Promotes Adaptive Behavior during
Adolescence. *J. Cogn. Neurosci.* 30, 281–289. doi: 10.1162/jocn_a_01206

Milardi, D., Bramanti, P., Milazzo, C., Finocchio, G., Arrigo, A., Santoro, G., et al. (2015).
Cortical and Subcortical Connections of the Human Claustrum Revealed In Vivo by
Constrained Spherical Deconvolution Tractography. *Cereb. Cortex* 25, 406–414. doi:
10.1093/cercor/bht231

Minciacchi, D., and Granato, A. (1989). Development of the thalamocortical system:
Transient-crossed projections to the frontal cortex in neonatal rats. *J. Comp. Neurol.*
281, 1–12. doi: 10.1002/cne.902810102

Numan, R. (2015). A Prefrontal-Hippocampal Comparator for Goal-Directed Behavior: The
Intentional Self and Episodic Memory. *Front. Behav. Neurosci.* 9. doi:
10.3389/fnbeh.2015.00323

Pachitariu, M., and Stringer, C. (2022). Cellpose 2.0: how to train your own model. *Nat.*
Methods 19, 1634–1641. doi: 10.1038/s41592-022-01663-4

Parbst, F., Kostka, J. K., Günther, A., Chen, Y.-N., Hanganu-Opatz, I. L., and Bitzenhofer, S.
H. (2025). Developmental Olfactory Dysfunction and Abnormal Odor Memory in
Immune-Challenged   Disc1+/−   Mice.   *J.   Neurosci.*   45.   doi:
10.1523/JNEUROSCI.1007-24.2025

Park, Y.-G., Sohn, C. H., Chen, R., McCue, M., Yun, D. H., Drummond, G. T., et al. (2019).
Protection of tissue physicochemical properties using polyfunctional crosslinkers.
Nat. Biotechnol. 37, 73–83. doi: 10.1038/nbt.4281

Parnaudeau, S., Bolkan, S. S., and Kellendonk, C. (2018). The Mediodorsal Thalamus: An
Essential Partner of the Prefrontal Cortex for Cognition. *Biol. Psychiatry* 83, 648–
656. doi: 10.1016/j.biopsych.2017.11.008

Paus, T., Zijdenbos, A., Worsley, K., Collins, D. L., Blumenthal, J., Giedd, J. N., et al.
(1999). Structural Maturation of Neural Pathways in Children and Adolescents: In
Vivo Study. *Science* 283, 1908–1911. doi: 10.1126/science.283.5409.1908

Pöpplau, J. A., Schwarze, T., Dorofeikova, M., Pochinok, I., Günther, A., Marquardt, A., et
al. (2024). Reorganization of adolescent prefrontal cortex circuitry is required for
mouse   cognitive   maturation.   *Neuron*   112,   421-440.e7.   doi:
10.1016/j.neuron.2023.10.024

Rios, O., and Villalobos, J. (2004). Postnatal development of the afferent projections from the
dorsomedial thalamic nucleus to the frontal cortex in mice. *Brain Res. Dev. Brain*
Res. 150, 47–50. doi: 10.1016/j.devbrainres.2004.02.005

Spellman, T., Rigotti, M., Ahmari, S. E., Fusi, S., Gogos, J. A., and Gordon, J. A. (2015).
Hippocampal–prefrontal input supports spatial encoding in working memory. *Nature*
522, 309–314. doi: 10.1038/nature14445

Sturrock, R. R. (1980). Myelination of the Mouse Corpus Callosum. *Neuropathol. Appl.*
Neurobiol. 6, 415–420. doi: 10.1111/j.1365-2990.1980.tb00219.x

Szczupak, D., Iack, P. M., Liu, C., IRC5 Consortium, Tovar-Moll, F., Lent, R., et al. (2021).
Direct Interhemispheric Cortical Communication via Thalamic Commissures: A New
White-Matter Pathway in the Rodent Brain. *Cereb. Cortex* 31, 4642–4651. doi:
10.1093/cercor/bhab112

Szczupak, D., Iack, P. M., Rayêe, D., Liu, C., Lent, R., Tovar-Moll, F., et al. (2023). The
relevance of heterotopic callosal fibers to interhemispheric connectivity of the
mammalian brain. *Cereb. Cortex* 33, 4752–4760. doi: 10.1093/cercor/bhac377

Takehara-Nishiuchi, K. (2014). Entorhinal cortex and consolidated memory. *Neurosci. Res.*
84, 27–33. doi: 10.1016/j.neures.2014.02.012

Thies, A. M., Pochinok, I., Marquardt, A., Dorofeikova, M., Hanganu-Opatz, I. L., and
Pöpplau, J. A. (2025). Trajectories of working memory and decision making abilities
along juvenile development in mice. *Front. Neurosci.* 19. doi:
10.3389/fnins.2025.1524931

van der Knaap, L. J., and van der Ham, I. J. M. (2011). How does the corpus callosum
mediate interhemispheric transfer? A review. *Behav. Brain Res.* 223, 211–221. doi:
10.1016/j.bbr.2011.04.018

van Duijvenvoorde, A. C. K., Peters, S., Braams, B. R., and Crone, E. A. (2016). What
motivates adolescents? Neural responses to rewards and their influence on
adolescents' risk taking, learning, and cognitive control. *Neurosci. Biobehav. Rev.* 70,
135–147. doi: 10.1016/j.neubiorev.2016.06.037

Vann, S. D., Aggleton, J. P., and Maguire, E. A. (2009). What does the retrosplenial cortex
do? *Nat. Rev. Neurosci.* 10, 792–802. doi: 10.1038/nrn2733

Voigt, F. F., Kirschenbaum, D., Platonova, E., Pagès, S., Campbell, R. A. A., Kastli, R., et al.
(2019). The mesoSPIM initiative: open-source light-sheet microscopes for imaging
cleared tissue. *Nat. Methods* 16, 1105–1108. doi: 10.1038/s41592-019-0554-0

Wang, Y., Chen, Z., Ma, G., Wang, L., Liu, Y., Qin, M., et al. (2023). A frontal transcallosal
inhibition loop mediates interhemispheric balance in visuospatial processing. *Nat.*
Commun. 14, 5213. doi: 10.1038/s41467-023-40985-5

Yang, C., Hu, Y., Talishinsky, A. D., Potter, C. T., Calva, C. B., Ramsey, L. A., et al. (2022).
Medial prefrontal cortex and anteromedial thalamus interaction regulates goal-
directed behavior and dopaminergic neuron activity. *Nat. Commun.* 13, 1386. doi:
10.1038/s41467-022-28892-7

Yorke Jr., C. H., and Caviness Jr., V. S. (1975). Interhemispheric neocortical connections of
the corpus callosum in the normal mouse: A study based on anterograde and
retrograde methods. *J. Comp. Neurol.* 164, 233–245. doi: 10.1002/cne.901640206

A Injection site Long-range projection

C Age groups at injection and perfusion

Neo Pre Juv Adu

P0 P10 P20 P50

P0 → P10 P10 → P20 P20 → P30 P50 → P60

B mCherry P0 P10

2mm

D eGFP 100µm

mCherry 1mm 100µm

E Neo Pre Juv Adu

1mm 1mm 1mm 1mm

F Ipsi Neo Pre Juv Adu

100µm

Contra

100µm

Figure 1

Figure 2

Figure 3

Figure 5

Figure 6