Title: A deep-time landscape of plant *cis*-regulatory sequence evolution

Authors: Kirk R. Amundson[1] [†], Anat Hendelman[2,3] [†], Danielle Ciren[4], Hailong Yang[1], Amber E. de Neve[1], Shai Tal[5], Adar Sulema[5], David Jackson[2], Madelaine E. Bartlett[1,6] *, Zachary B. Lippman[2,3,4] *, Idan Efroni[5] *

Affiliations:
[1]Department of Biology, University of Massachusetts; Amherst, MA, USA.
[2]Cold Spring Harbor Laboratory; Cold Spring Harbor, NY, USA.
[3]Howard Hughes Medical Institute, Cold Spring Harbor Laboratory; Cold Spring Harbor, NY, 4 USA.
[4]Cold Spring Harbor Laboratory, School of Biological Sciences; Cold Spring Harbor, NY, USA.
[5]Institute of Plant Science, The Hebrew University; Rehovot, Israel.
[6]Sainsbury Laboratory Cambridge University (SLCU); Cambridge, UK.

[†] These authors contributed equally to this work.
* Co-corresponding authors

Abstract: Developmental gene function is conserved over deep time, but *cis*-regulatory sequence conservation is rarely found. Rapid sequence turnover, paleopolyploidy, structural variation, and limited phylogenomic sampling have impeded conserved non-coding sequence (CNS) discovery. Using Conservatory, an algorithm that leverages microsynteny and iterative alignments to map CNS-gene associations over evolution, we uncovered ~2.3M CNSs, including over 3,000 predating angiosperms, from 284 plant species spanning 300 million years of diversification. Ancient CNSs were enriched near developmental regulators, and mutating these CNSs near *HOMEOBOX* genes produced strong phenotypes. Tracing CNS evolution uncovered key principles: CNS spacing varies, but order is conserved; genomic rearrangements form new CNS-gene associations; and ancient CNSs are preferentially retained among paralogs, but are often lost as cohorts or evolve into lineage-specific CNSs.

One Sentence Summary:
Conservatory maps ancient regulatory elements and uncovers regulatory evolution dynamics.

5   **Main Text:** *Cis*-regulatory sequence conservation and divergence likely underlie phenotypic evolution (*1–3*). Defining principles that govern *cis*-regulatory evolution is key to deciphering the genetic basis of natural and artificial selection, and essential for targeted manipulation of gene regulation (*4–6*). A long-standing paradox in the evolution of regulatory sequences is that homologous genes in distantly related species apparently lack *cis*-regulatory sequence

10   conservation, despite similar expression patterns and functions (*7*). However, identifying *cis*-regulatory elements (CREs) and establishing their orthology across species is often technically challenging, especially over deep time.

Evolutionary conservation can identify CREs as conservation indicates selection for preserved regulatory function (*8–14*). Conserved non-coding sequences (CNSs) are often identified through

15   whole-genome alignment approaches (*15, 16*), but these methods fail in comparing distantly related species, because regulatory sequence evolves rapidly. Additionally, frequent macro- and micro-genomic rearrangements, gene duplications, and whole-genome duplications obscure orthology relationships. Thus, homologous CREs often cannot be directly aligned in distantly related species (*17, 18*). Iterative alignment through intermediate "bridge" genomes improves

20   CNS discovery across broader evolutionary distances (*18, 19*). However, these strategies must still contend with pervasive structural variation and the challenges of complex, highly rearranged, paleopolyploid, or repetitive genomes typical of many eukaryotes (*20*).

The challenges of identifying CNSs are magnified in plant genomes. Repeated cycles of whole-genome duplication, subsequent fractionation to diploidization, and the unequal, often biased

25   retention of paralogous genes obscure gene and CNS orthology relationships. As a result, most identified plant CNSs are limited in phylogenetic scope, with only a handful extending beyond taxonomic families (*21–25*).

Here, we present Conservatory (www.conservatorycns.com), an orthogroup-centric CNS discovery algorithm developed for detecting noncoding sequence conservation that accounts for

30   complex gene orthology relationships, extreme structural variation, and rapid sequence divergence. Using Conservatory, we annotated and resolved CNS orthologies and associated genes across 314 plant genomes representing 284 species spanning green plant diversity. Our analysis uncovered widespread conservation of noncoding sequences at unprecedented evolutionary depths. We use this comprehensive atlas to trace ~300 million years of *cis*-

35   regulatory sequence evolution in plants (*26*), uncovering fundamental principles shaping the dynamic formation, conservation, and turnover of plant regulatory sequences.

Ancient *cis*-regulatory elements are common in plant genomes

To address challenges in identifying CNSs, we developed 'Conservatory', a large-scale comparative genomics algorithm that captures sequence conservation across deep time, while

40   accounting for gene duplication and rapid sequence divergence. Conservatory identifies orthologs within a particular taxonomic family, then aligns up to 120 kb of intergenic sequence surrounding each ortholog to a common reference genome. To account for frequent paleopolyploidy, fractionation, and structural rearrangements, multiple co-orthologs for each reference gene are retained, based on microsynteny and sequence conservation data. CNSs

45   surrounding candidate orthologs were then identified using *phyloP* (*27*), and ancestral sequence reconstruction (*FastML*) (*28*) was used to infer CNS sequences at the crown nodes of each taxonomic family. Ancestral sequences were then used to search the regulatory regions of all putative orthologs outside of the taxonomic family using high-sensitivity alignment. This two-step alignment, first within and then between families, allows for CNS discovery despite

5   divergence of flanking sequence. To account for reference genome bias, independent CNS
       datasets were generated using multiple reference genomes. The algorithm then uses 'bridge
       genomes' to infer the homology of CNSs that cannot be directly aligned, but can be linked by
       intermediate sequences (*19*). After filtering, Conservatory generates a catalog of CNSs and their
       orthologous relations (**fig. S1**).

10   We applied Conservatory to 314 genomes, representing 284 species spanning green plant
       diversity, using 10 reference genomes from the Solanaceae, Asteraceae, Brassicaceae, Fabaceae,
       Poaceae, and Lemnaceae families (**Fig. 1A; table S1**). We discovered 32.8 million CNSs across
       all genomes, and grouped them into 2.3 million unique CNSs based on sequence orthology.
       Conservatory CNSs were short (median length 12-40 bp; **Fig. 1B**), consistent with previous
15   reports (*21–24*, *29*, *30*). They were enriched for transcription factor binding sites and activating
       histone modifications, and depleted for DNA methylation and repressive histone marks (**Fig.
       1C**), as in other plant regulatory sequences (*22*, *31*). The Conservatory CNSs were also
       significantly enriched in accessible chromatin regions in arabidopsis, tomato, and maize ($p < $ 1e-
       4, permutation test; **table S2**). Moreover, the CNSs were preferentially located at the centers of
20   open chromatin regions, likely pinpointing specific functional sequences in cis-regulatory space
       (**Fig. 1D-F**). Thus, Conservatory CNSs have the hallmarks of functional *cis*-regulatory elements.

       We estimated the age of CNSs by mapping them to a phylogenetic tree with nodes dated
       according to published best estimates (see methods). Most (2.15 million; 92%) were conserved
       within plant families. Remarkably, we identified many ancient CNSs that emerged prior to the
25   diversification of the eudicots (n=31,314; 1.34%), angiosperms (n=3,321; 0.14%), or even seed
       plants (n=633; 0.02%) (**Fig. 1G; table S3**). To assess the functional categories of genes
       harboring the most ancient seed plant CNSs, we performed gene ontology (GO) term analysis.
       Similar to ancient metazoan CNSs, genes associated with angiosperm and seed plant CNSs were
       significantly enriched for developmental and transcriptional biological processes (**Fig. 1H**),
30   suggesting critical roles in conserved developmental programs.

       Mutations in deeply conserved sequences disrupt development

       To determine the functional roles of deeply conserved CNS-gene associations in development,
       we focused on genes with canonical roles in embryogenesis (*32*). The ancient homeobox gene
       WUSCHEL HOMEOBOX 9 (*WOX9*) has conserved functions in development, and null mutants
35   in tomato are embryonic lethal (*9*). We re-evaluated our *SlWOX9* promoter allelic series (*9*) and
       found that mutations disrupting a seed-plant-level CNS caused partially penetrant vegetative
       growth defects, likely stemming from defective shoot apical meristem development. We
       generated two additional alleles disrupting this ancient CNS (S217), and they also replicated
       these phenotypes (**Fig. 2A-C**).

40   To extend our functional analyses, we analyzed *WOX2,* which is expressed in the arabidopsis
       zygote and acts in embryogenesis (*32*). To ask if *WOX2* function is conserved in tomato, we
       generated two independent coding null alleles in *SlWOX2*, neither of which could be recovered
       as homozygotes, indicating fully penetrant embryonic lethality. This is in contrast to arabidopsis,
       where *WOX2* is redundant with some of its paralogs (*32*), likely reflecting the distant
45   relationship between tomato and arabidopsis. Conservatory identified three angiosperm-level and
       three eudicot-level CNSs in the *SlWOX2* promoter (**Fig. 2D**). We used CRISPR-Cas9
       mutagenesis to generate a *SlWOX2* promoter allelic series (*SlWOX2^{Pro}*). Deletions overlapping
       the angiosperm CNSs resulted in strong segregation distortion, indicating partially penetrant
       embryonic lethality. Rare homozygotes of these promoter alleles had vegetative defects, similar

5   to the *SlWOX9^{Pro}* alleles that disrupted a seed plant CNS (**Fig. 2B- F**) (*9*). Taken together, these results demonstrate key roles for ancient CNSs in embryogenesis, with partially penetrant phenotypes indicating roles for additional CREs.

We next asked if ancient CNSs play similar roles in other developmental genes, by analyzing Conservatory CNSs in known CREs of eight transcription factors that regulate meristem, leaf,

10   flower, or epidermal development (*13*, *33–38*). In all cases, functional CREs overlapped with ancient CNSs found by Conservatory (**fig. S2**). In particular, the *WOX* gene *WUSCHEL (WUS)*, a conserved regulator of stem cell proliferation in meristems (*39*), has two defined regulatory modules. One module drives *WUS* expression in ovules, and the other in meristems (**Fig. 3A**) (*40*). We discovered five ancient CNSs in the meristem module, including conserved binding

15   sites for Type-B RESPONSE REGULATORS (RRs). These transcription factors act downstream of cytokinin to induce *WUS* expression in multiple species (**Fig. 3A**) (*41*, *42*). This correspondence between classical CRE studies and Conservatory CNSs is consistent with an important role for ancient CNSs in development.

Two core meristem regulators uncover CNS evolutionary dynamics

20   We utilized Conservatory identification of CNS homologous relationships across species to investigate how CNSs evolve, first focusing on the deeply conserved *WUS* meristem regulatory module. In arabidopsis, Conservatory found five partly overlapping CNSs in this module. The S115 CNS sequence, deeply conserved even in the gymnosperm *Cycas panzhiuaensis*, was duplicated and found in distal and proximal positions (**Fig. 3A**). Cross-species analysis suggested

25   that the core spatial configuration of this meristem module emerged at the base of the angiosperms, while the S115 duplication likely occurred at the base of the eudicots. Strikingly, while most of the CNSs in this module were lost in the grasses, the deeply conserved S115, which contained a Type-B RR binding site important for meristem expression (*41*, *42*), was retained (**Fig. 3B**). Consistently, the *WUS* meristem expression pattern is conserved in the

30   grasses, despite divergent meristem morphology and developmental dynamics (*43*, *44*).

WUS underwent several independent duplications, and CNS analysis for the duplication that occurred at the base of the Asteraceae showed that associated *cis*-regulatory sequences were duplicated with the coding regions. However, these sequences then exhibited species-specific patterns of divergence and loss (**Fig. 3B**). In lettuce (*Lactuca sativa)*, only one angiosperm-level

35   CNS was retained in both *WUS* paralogs, while the other ancient CNSs were differentially retained and partitioned between the two paralogs. The products of the same ancestral duplication diverged differently in sunflower (*Helianthus annuus)*, with one *WUS* paralog maintaining all ancient CNSs, and the other maintaining only some of them (**Fig. 3B**). Both species appeared to have lost the distal copy of S115, but closer examination revealed an

40   Asteraceae-specific CNS similar to the ancestral S115 sequence in the collinear location (**fig. S3A-C**), suggesting sequence divergence followed by purifying selection in the Asteraceae. Analysis of an additional *WUS* duplication that occurred at the base of a large clade of grasses revealed similar evolutionary patterns, with S115 and Poaceae-specific CNS shared between the paralogs, but each paralog also had lineage-specific CNSs (**fig. S3D**). Notably, despite

45   divergence in the regulatory sequences of many *WUS* paralogs, the ancient meristem module CNSs were maintained across angiosperms (**Fig. 3B**).

Conservatory associates CNSs and genes based on conserved syntenic relationships, and a CNS can be associated with more than one nearby gene. Analysis of the ancient S115 CNS found that it was associated with multiple orthogroups at different phylogenetic levels. Throughout

5 angiosperms, the S115 CNS was commonly associated with genes encoding the chromatin remodeler TRA1A, and in eudicots with a gene encoding YAF9, which is in the TRA1A protein complex (*45*). Other associations were also identified, including a grass-specific one with a gene encoding an uncharacterized CCCH zinc-finger protein (**fig. S4A**). Remarkably, these associations were conserved despite frequent genomic rearrangements and rapid decay of plant

10 genome synteny (*46*), suggesting that the S115 CNS may indeed regulate the expression of multiple genes (**fig. S4B**).

To characterize CNS evolution on shorter time scales, we next analyzed the maize *TEOSINTE BRANCHED1* (*TB1*) gene. *TB1* encodes a TCP transcription factor with conserved roles in axillary branching (*47–49*). Regions upstream of *TB1* orthologs are associated with maize (*Zea*

15 *mays*), sorghum (*Sorghum bicolor*), and pearl millet (*Pennisetum glaucum*) domestication (*50–52*). Analysis of the maize domestication locus, located ~65 kb upstream of *TB1,* uncovered multiple CNSs separated into distal and proximal collinear clusters (P1-P6 and P7-P12, respectively; **Fig. 3C**). Tracing the evolution of *TB1* CNSs revealed that the proximal cluster emerged at the base of the grasses (~70 mya), while the distal cluster emerged later, before a

20 major grass radiation that includes maize (PACMAD clade, ~33 mya, **Fig. 3D**). Notably, a gain-of-function *Hopscotch* transposon insertion associated with domestication separated the proximal and distal CNS clusters in maize, leaving the CNSs intact. Individual fragments of the regulatory region containing these CNS act as repressors (*53*), suggesting that the gain-of-function TB1 phenotype in maize could be caused by the *Hopscotch* insertion disrupting CNS spacing in a

25 conserved repressor element (**Fig. 3A, F**).

Comparison of CNS spacing and order in *TB1* and *WUS* CNSs revealed that CNS colinearity was remarkably conserved among species, with the notable exception of an inversion in several grasses (**Fig. 3F**). In contrast, the distances between CNSs and their associated genes varied and were CNS-specific. Thus, the *WUS* meristem CNS module was within 1.5 kb of the start codon

30 in 224 of 240 species, and only in 8 cases was more than 10 kb upstream. The position of the *TB1* enhancer was much more variable (**Fig. 3E**), and in some cases, such as sorghum, it was closer to a different gene than to *TB1* itself. In other grasses, such as rice (*Oryza sativa*), a different gene was located between the CNS cluster and the *TB1* ortholog coding region (**Fig. 3F**).

35 Overall, our analysis of *WUS* and *TB1* revealed that CNSs maintained after duplication diverge in particular lineages, change position relative to their associated genes, or even get lost as cohorts in specific lineages. Interestingly, gene duplication was associated with duplication of associated CNSs, which remained as multiple copies in the genome. We next asked whether such dynamics characterized CNS evolution on a global scale.

40 **Genome rearrangements alter CNS position and gene associations**

To study the evolutionary dynamics of CNS position, we first analyzed the positions of all CNSs relative to their associated genes, grouped by CNS age. The distribution of CNS positions was genome-specific, but overall, most CNSs were located upstream of associated genes, and were often close to coding sequences, with older CNSs tending to be closer to start codons (**Fig. 4A**).

45 However, many CNSs across all age categories were more than 25 kb from their associated genes. CNSs were often closer to a different gene than to their associated gene, and 25% to 50% of CNSs-gene associations skipped over intervening, non-associated genes (**Fig. 4B; table S5**).

Long-range CNS-gene interactions and interactions skipping over other genes are common in metazoans (*54*), but are only rarely reported in plants. While they may be common in the large

maize genome (*55*), we were surprised to find so many distal or non-adjacent associations, including in compact genomes. Most genes intervening between CNSs and their associated genes were associated with CNSs of their own, and most of the closest non-associated genes were not orthologous to associated genes. This suggests that CNS-gene associations that skip over non-associated genes were unlikely to be caused by annotation artifacts or by tandem gene duplications (**fig. S5A-B**). To provide empirical support for such long-range CNS-gene associations, we investigated the concordance of predicted CNS-gene associations with predicted chromatin loops, as *cis*-regulatory elements and gene interactions might be maintained through chromatin looping (*55*). We carried out chromatin-conformation capture sequencing on tomato meristem tissue, and reanalyzed published datasets for arabidopsis and maize, focusing on loops connecting intergenic regions to genes (*55–57*). Compared to loops that did not overlap with CNSs, loops with CNSs at their intergenic edges were more likely to be connected to genes ($p < 0.001$, chi-square test) (**Fig. 4C**). This suggests that intergenic DNA sequences looping to genes are conserved, consistent with Conservatory CNSs acting as regulatory elements. To directly assess the concordance of CNS-gene associations with chromatin looping, we focused on loops connecting intergenic regions to genes that harbored CNSs, and quantified CNS-gene associations supported by loops. Across all three species, 52-67% of loops showed at least one concordant CNS-gene association (**Fig. 4D-E**), indicating that CNS-gene associations can pinpoint physical chromatin interactions and are likely retained over large genomic distances, despite evolutionary variability in their positions.

Our analysis uncovered an unexpected relationship between the depth of CNS conservation and relative position to their associated genes, with ancient CNSs occurring closer to transcription start sites. One explanation for this observation is that the functions of these CNSs are position-dependent. In support of this, analysis of CNS mobility (defined as the standard variation of its relative position) over short evolutionary distances (i.e., within taxonomic families) revealed that mobility of ancient CNSs was still significantly lower than that of comparable family-level CNSs ($p<2.2e-16$; Wilcoxon test; **Fig. 4F**).

Evolution of new *cis*-regulatory sequences from ancestral CNSs

Analysis of the *WUS* meristem module revealed that even ancient CNSs can diverge or be lost in specific lineages. To assess the extent of this phenomenon, we examined the presence or absence of ancient CNSs across species. Remarkably, the number of angiosperm-level CNSs detected in each species varied widely, yet was highly consistent within families, suggesting that most losses or divergences occurred during early events at the base of each family (**Fig. 5A**). The consistency within families suggests that this CNS loss is a biological phenomenon, rather than an artifact of genome annotation or assembly quality. Using a maximum parsimony approach, we mapped the most likely phylogenetic nodes where angiosperm CNS loss or divergence occurred. While there was a gradual overall trend of CNS loss or divergence, we also detected punctuated, large-scale losses in families such as Brassicaceae, Asteraceae, Poaceae, and Lemnaceae (**Fig. 5B**). In the grasses, this large-scale CNS loss is consistent with their rapid evolutionary diversification (*58*). Indeed, lost CNSs were associated with genes such as *WOX4* and *KANADI*, which play key roles in vasculature and leaf development, processes in which grasses display unique morphological innovations (**table S6**).

The apparent loss of ancient CNSs may reflect degradation and loss of function, or, as for *WUS* in the Asteraceae, divergence accompanied by the emergence of lineage-specific CNSs (**fig. S3**). Because sparse phylogenetic sampling limits detection of early divergence events across species,

5   we analyzed paralog evolution within genomes as a proxy for early stages of divergence, as in these cases, ancestral CNS complements can be reconstructed and regulatory turnover directly traced. We identified duplicate pairs derived from whole-genome duplications in our 10 reference genomes (82). Following duplication, the number of shared CNSs declined with age, but retention patterns differed between duplicates: one paralog generally maintained a stable

10   CNS complement, whereas the other experienced an initial drop before stabilizing. Meanwhile, paralog-specific CNSs steadily accumulated, revealing an early phase of divergence marked by CNS loss that was subsequently balanced by the gain of new CNSs (**Fig. 5C**). Notably, as observed for *WUS* paralogs, ancient CNSs were preferentially maintained in both paralogs compared to younger, family-level ones. Consistently, these old CNSs accumulated in higher

15   copy numbers in genomes (**Fig. 5D-E**).

Finally, to determine whether new paralog-specific CNSs arise *de novo* or are derived from ancestral sequences, we examined paralog pairs in which an old CNS was retained in one paralog but replaced by a different CNS at the same position in the other paralog. We reasoned that if these sequences are derived, they should still retain some sequence similarity to the old CNS.

20   Using CNSs retained in both paralogs as alignment anchors, we compared the sequence similarity between old CNSs and new paralog-specific CNSs at syntenic sites versus those in non-syntenic sites. Syntenic replaced–new CNS pairs showed significantly higher identity than non-syntenic controls (**Fig. 5F**; $p < 0.001$; Dunn test), suggesting that these new, paralog-specific CNSs originate from ancestral sequences rather than from entirely new sequences.

Discussion

Conservation and divergence of cis-regulatory sequences are key forces driving morphological evolution, yet their evolutionary dynamics remain poorly understood. Using Conservatory as a platform for CNS discovery, we identified and traced the evolution of over 2 million CNSs, including many that predate the emergence of angiosperms.

30   Why are these sequences so deeply conserved? Their enrichment near developmental regulators suggests they govern critical regulatory programs. This is supported by the severe developmental phenotypes observed when such sequences are disrupted (**Fig. 3**) (*9–12, 14*). Moreover, their retention in high copy numbers following repeated gene duplications suggests that their function may be less dependent on the surrounding sequence environment. Thus, similar to ancient

35   metazoan CNSs, ancient plant CNSs may encode functionally-enriched core regulatory sequences (*59*). The observed gradient of CRE selective constraints, with slowly evolving deeply conserved sequences flanked by faster evolving lineage-specific sequences, suggests a model whereby deeply conserved sequences form the foundation of core expression programs, which are modified by peripheral sequences, whose modifications contribute to lineage-specific trait

40   variation and innovation.

Following gene duplication, the regulatory landscapes of the resulting paralogs rapidly diverge (*60, 61*). In our analyses, we observed clear asymmetry: one paralog largely retained ancestral CNSs, whereas the other underwent rapid turnover, first losing conserved elements and then acquiring paralog-specific CNSs. Although new regulatory sequences are often thought to arise

45   *de novo* from neutrally evolving noncoding DNA, many may originate from older sequences (*5, 62*). By tracing CNS dynamics in paralogs, we provide direct evidence that new CNSs often emerge through divergence of ancestral elements rather than *de novo* formation. Furthermore,

5   this suggests that certain sequences possess intrinsic properties that predispose them to regulatory function and long-term retention.

One of the paradoxes in CNS evolution is that deeply conserved sequences can nevertheless be lost in cohorts within specific lineages. Such loss may result from extreme developmental systems drift, in which, after gene duplication or divergence of other regulatory sequences, a

10   deeply conserved element becomes redundant and gradually degenerates through mutation (*63*, *64*). Alternatively, it may reflect technical limitations: orthologous CREs are difficult to align and identify across large evolutionary distances (*17*, *18*). While leveraging hundreds of genomes helped bridge these gaps, sparse phylogenomic coverage remains a major constraint for reliably identifying CNS orthology. For example, the apparent loss of many deeply conserved CNSs in

15   grasses and duckweeds is consistent with large-scale *cis*-regulatory rewiring during their early divergence, yet poor sampling of monocots outside these families means that many lineage-specific CNSs may still be orthologous to ancestral angiosperm elements. Here, we provide a unified map of CNS evolution in plants, based on 314 genomes. As pan-genomic resources grow, improved phylogenomic resolution will further clarify orthologous relationships between

20   CNSs and guide deeper computational analyses and functional assays to refine our understanding of regulatory sequence evolution and provide even higher resolution targets for trait engineering (*5*, *65*).

25   **References and Notes**

1. S. B. Carroll, Evo-devo and an expanding evolutionary synthesis: a genetic theory of morphological evolution. *Cell* **134**, 25–36 (2008).

2. P. J. Wittkopp, G. Kalay, Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat. Rev. Genet.* **13**, 59–69 (2011).

3. D. L. Stern, V. Orgogozo, The loci of evolution: how predictable is genetic evolution? *Evolution* **62**, 2155–2177 (2008).

4. D. Villar, P. Flicek, D. T. Odom, Evolution of transcription factor binding in metazoans - mechanisms and functional implications. *Nat. Rev. Genet.* **15**, 221–233 (2014).

5. X. Li, R. J. Schmitz, Cis-regulatory dynamics in plant domestication. *Trends Genet.*, doi: 10.1016/j.tig.2025.02.005 (2025).

6. A. P. Marand, L. Jiang, F. Gomez-Cano, M. A. A. Minow, X. Zhang, J. P. Mendieta, Z. Luo, S. Bang, H. Yan, C. Meyer, L. Schlegel, F. Johannes, R. J. Schmitz, The genetic architecture of cell type-specific cis regulation in maize. *Science* **388**, eads6601 (2025).

7. D. Villar, C. Berthelot, S. Aldridge, T. F. Rayner, M. Lukk, M. Pignatelli, T. J. Park, R. Deaville, J. T. Erichsen, A. J. Jasinska, J. M. A. Turner, M. F. Bertelsen, E. P. Murchison, P. Flicek, D. T. Odom, Enhancer evolution across 20 mammalian species. *Cell* **160**, 554–566 (2015).

8. E. Z. Kvon, O. K. Kamneva, U. S. Melo, I. Barozzi, M. Osterwalder, B. J. Mannion, V. Tissières, C. S. Pickle, I. Plajzer-Frick, E. A. Lee, M. Kato, T. H. Garvin, J. A. Akiyama, V. Afzal, J. Lopez-Rios, E. M. Rubin, D. E. Dickel, L. A. Pennacchio, A. Visel, Progressive loss of function in a limb enhancer during snake evolution. *Cell* **167**, 633–642.e11 (2016).

9. A. Hendelman, S. Zebell, D. Rodriguez-Leal, N. Dukler, G. Robitaille, X. Wu, J. Kostyun, L.

Tal, P. Wang, M. E. Bartlett, Y. Eshed, I. Efroni, Z. B. Lippman, Conserved pleiotropy of an ancient plant homeobox gene uncovered by cis-regulatory dissection. *Cell* **184**, 1724–1739.e16 (2021).

10. A. Lanctot, A. Hendelman, P. Udilovich, G. M. Robitaille, Z. B. Lippman, Antagonizing cis-regulatory elements of a conserved flowering gene mediate developmental robustness. *Proc. Natl. Acad. Sci. U. S. A.* **122**, e2421990122 (2025).

11. D. Ciren, S. Zebell, Z. B. Lippman, Extreme restructuring of cis-regulatory regions controlling a deeply conserved plant stem cell regulator. *PLoS Genet.* **20**, e1011174 (2024).

12. X. Wang, L. Aguirre, D. Rodríguez-Leal, A. Hendelman, M. Benoit, Z. B. Lippman, Dissecting cis-regulatory control of quantitative trait variation in a plant stem cell circuit. *Nat. Plants* **7**, 419–427 (2021).

13. T. C. Tran, K. Mähl, C. Kappel, Y. Dakhiya, A. Sampathkumar, A. Sicard, M. Lenhard, Altered interactions between cis-regulatory elements partially resolve BLADE-ON-PETIOLE genetic redundancy in Capsella rubella. *Plant Cell* **36**, 4637–4657 (2024).

14. S. G. Zebell, C. Martí-Gómez, B. Fitzgerald, C. P. Cunha, M. Lach, B. M. Seman, A. Hendelman, S. Sretenovic, Y. Qi, M. Bartlett, Y. Eshed, D. M. McCandlish, Z. B. Lippman, Cryptic variation fuels plant phenotypic change through hierarchical epistasis. *Nature* **644**, 984–992 (2025).

15. M. J. Christmas, I. M. Kaplow, D. P. Genereux, M. X. Dong, G. M. Hughes, X. Li, P. F. Sullivan, A. G. Hindle, G. Andrews, J. C. Armstrong, M. Bianchi, A. M. Breit, M. Diekhans, C. Fanter, N. M. Foley, D. B. Goodman, L. Goodman, K. C. Keough, B. Kirilenko, A. Kowalczyk, C. Lawless, A. L. Lind, J. R. S. Meadows, L. R. Moreira, R. W. Redlich, L. Ryan, R. Swofford, A. Valenzuela, F. Wagner, O. Wallerman, A. R. Brown, J. Damas, K. Fan, J. Gatesy, J. Grimshaw, J. Johnson, S. V. Kozyrev, A. J. Lawler, V. D. Marinescu, K. M. Morrill, A. Osmanski, N. S. Paulat, B. N. Phan, S. K. Reilly, D. E. Schäffer, C. Steiner, M. A. Supple, A. P. Wilder, M. E. Wirthlin, J. R. Xue, Zoonomia Consortium§, B. W. Birren, S. Gazal, R. M. Hubley, K.-P. Koepfli, T. Marques-Bonet, W. K. Meyer, M. Nweeia, P. C. Sabeti, B. Shapiro, A. F. A. Smit, M. S. Springer, E. C. Teeling, Z. Weng, M. Hiller, D. L. Levesque, H. A. Lewin, W. J. Murphy, A. Navarro, B. Paten, K. S. Pollard, D. A. Ray, I. Ruf, O. A. Ryder, A. R. Pfenning, K. Lindblad-Toh, E. K. Karlsson, Evolutionary constraint and innovation across hundreds of placental mammals. *Science* **380**, eabn3943 (2023).

16. J. Stiller, S. Feng, A.-A. Chowdhury, I. Rivas-González, D. A. Duchêne, Q. Fang, Y. Deng, A. Kozlov, A. Stamatakis, S. Claramunt, J. M. T. Nguyen, S. Y. W. Ho, B. C. Faircloth, J. Haag, P. Houde, J. Cracraft, M. Balaban, U. Mai, G. Chen, R. Gao, C. Zhou, Y. Xie, Z. Huang, Z. Cao, Z. Yan, H. A. Ogilvie, L. Nakhleh, B. Lindow, B. Morel, J. Fjeldså, P. A. Hosner, R. R. da Fonseca, B. Petersen, J. A. Tobias, T. Székely, J. D. Kennedy, A. H. Reeve, A. Liker, M. Stervander, A. Antunes, D. T. Tietze, M. F. Bertelsen, F. Lei, C. Rahbek, G. R. Graves, M. H. Schierup, T. Warnow, E. L. Braun, M. T. P. Gilbert, E. D. Jarvis, S. Mirarab, G. Zhang, Complexity of avian evolution revealed by family-level genomes. *Nature* **629**, 851–860 (2024).

17. E. S. Wong, D. Zheng, S. Z. Tan, N. L. Bower, V. Garside, G. Vanwalleghem, F. Gaiti, E. Scott, B. M. Hogan, K. Kikuchi, E. McGlinn, M. Francois, B. M. Degnan, Deep conservation of the enhancer regulatory code in animals. *Science* **370**, eaax8137 (2020).

18. M. H. Q. Phan, T. Zehnder, F. Puntieri, A. Magg, B. Majchrzycka, M. Antonović, H.

Wieler, B.-W. Lo, D. Baranasic, B. Lenhard, F. Müller, M. Vingron, D. M. Ibrahim, Conservation of regulatory elements with highly diverged sequences across large evolutionary distances. *Nat. Genet.* **57**, 1524–1534 (2025).

19.   I. Braasch, A. R. Gehrke, J. J. Smith, K. Kawasaki, T. Manousaki, J. Pasquier, A. Amores, T. Desvignes, P. Batzel, J. Catchen, A. M. Berlin, M. S. Campbell, D. Barrell, K. J. Martin, J. F. Mulley, V. Ravi, A. P. Lee, T. Nakamura, D. Chalopin, S. Fan, D. Wcisel, C. Cañestro, J. Sydes, F. E. G. Beaudry, Y. Sun, J. Hertel, M. J. Beam, M. Fasold, M. Ishiyama, J. Johnson, S. Kehr, M. Lara, J. H. Letaw, G. W. Litman, R. T. Litman, M. Mikami, T. Ota, N. R. Saha, L. Williams, P. F. Stadler, H. Wang, J. S. Taylor, Q. Fontenot, A. Ferrara, S. M. J. Searle, B. Aken, M. Yandell, I. Schneider, J. A. Yoder, J.-N. Volff, A. Meyer, C. T. Amemiya, B. Venkatesh, P. W. H. Holland, Y. Guiguen, J. Bobe, N. H. Shubin, F. Di Palma, J. Alföldi, K. Lindblad-Toh, J. H. Postlethwait, The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. *Nat. Genet.* **48**, 427–437 (2016).

20.   B. Song, E. S. Buckler, M. C. Stitzer, New whole-genome alignment tools are needed for tapping into plant diversity. *Trends Plant Sci.* **29**, 355–369 (2024).

21.   A. Haudry, A. E. Platts, E. Vello, D. R. Hoen, M. Leclercq, R. J. Williamson, E. Forczek, Z. Joly-Lopez, J. G. Steffen, K. M. Hazzouri, K. Dewar, J. R. Stinchcombe, D. J. Schoen, X. Wang, J. Schmutz, C. D. Town, P. P. Edger, J. C. Pires, K. S. Schumaker, D. E. Jarvis, T. Mandáková, M. A. Lysak, E. van den Bergh, M. E. Schranz, P. M. Harrison, A. M. Moses, T. E. Bureau, S. I. Wright, M. Blanchette, An atlas of over 90,000 conserved noncoding sequences provides insight into crucifer regulatory regions. *Nat. Genet.* **45**, 891–898 (2013).

22.   B. Song, E. S. Buckler, H. Wang, Y. Wu, E. Rees, E. A. Kellogg, D. J. Gates, M. Khaipho-Burch, P. J. Bradbury, J. Ross-Ibarra, M. B. Hufford, M. C. Romay, Conserved noncoding sequences provide insights into regulatory sequence and loss of gene expression in maize. *Genome Res.* **31**, 1245–1257 (2021).

23.   M. C. Stitzer, A. S. Seetharam, A. Scheben, S.-K. Hsu, A. J. Schulz, T. AuBuchon-Elder, M. El-Walid, T. H. Ferebee, C. O. Hale, T. La, Z.-Y. Liu, S. J. McMorrow, P. Minx, A. Phillips, M. Syring, T. Wrightsman, J. Zhai, R. Pasquet, C. McAllister, S. Malcomber, P. Traiperm, D. Layton, J. Zhong, D. E. Costich, R. K. Dawe, K. Fengler, C. Harris, Z. Irelan, V. Llaca, P. Parakkal, G. Zastrow-Hayes, M. R. Woodhouse, E. K. S. Cannon, J. Portwood II, C. M. Andorf, P. S. Albert, J. A. Birchler, A. Siepel, J. Ross-Ibarra, M. C. Romay, E. Kellogg, E. S. Buckler IV, M. Hufford, Extensive genome evolution distinguishes maize within a stable tribe of grasses, *bioRxiv* (2025). https://doi.org/10.1101/2025.01.22.633974.

24.   H. Xin, X. Liu, S. Chai, X. Yang, H. Li, B. Wang, Y. Xu, S. Lin, X. Zhong, B. Liu, Z. Lu, Z. Zhang, Identification and functional characterization of conserved cis-regulatory elements responsible for early fruit development in cucurbit crops. *Plant Cell* **36**, 2272–2288 (2024).

25.   X. Lai, S. Behera, Z. Liang, Y. Lu, J. S. Deogun, J. C. Schnable, STAG-CNS: An order-aware conserved noncoding sequences discovery tool for arbitrary numbers of species. *Mol. Plant* **10**, 990–999 (2017).

26.   S. Magallón, S. Gómez-Acevedo, L. L. Sánchez-Reyes, T. Hernández-Hernández, A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytol.* **207**, 437–453 (2015).

27. K. S. Pollard, M. J. Hubisz, K. R. Rosenbloom, A. Siepel, Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res.* **20**, 110–121 (2010).

28. H. Ashkenazy, O. Penn, A. Doron-Faigenboim, O. Cohen, G. Cannarozzi, O. Zomer, T. Pupko, FastML: a web server for probabilistic reconstruction of ancestral sequences. *Nucleic Acids Res.* **40**, W580–4 (2012).

29. J. Van de Velde, M. Van Bel, D. Vaneechoutte, K. Vandepoele, A collection of conserved noncoding sequences to study gene regulation in flowering plants. *Plant Physiol.* **171**, 2586–2598 (2016).

30. D. Burgess, M. Freeling, The most deeply conserved noncoding sequences in plants serve similar functions to those in vertebrates despite large differences in evolutionary rates. *Plant Cell* **26**, 946–961 (2014).

31. Z. Lu, A. P. Marand, W. A. Ricci, C. L. Ethridge, X. Zhang, R. J. Schmitz, The prevalence, evolution and chromatin signatures of plant regulatory elements. *Nat. Plants* **5**, 1250–1259 (2019).

32. H. Breuninger, E. Rikirsch, M. Hermann, M. Ueda, T. Laux, Differential expression of WOX genes mediates apical-basal axis formation in the Arabidopsis embryo. *Dev. Cell* **14**, 867–876 (2008).

33. J. A. Aguilar-Martínez, N. Uchida, B. Townsley, D. A. West, A. Yanez, N. Lynn, S. Kimura, N. Sinha, Transcriptional, posttranscriptional, and posttranslational regulation of SHOOT MERISTEMLESS gene expression in Arabidopsis determines gene function in the shoot apex. *Plant Physiol.* **167**, 424–442 (2015).

34. B. Raatz, A. Eicker, G. Schmitz, E. Fuss, D. Müller, S. Rossmann, K. Theres, Specific expression of LATERAL SUPPRESSOR is controlled by an evolutionarily conserved 3' enhancer: LAS/Ls is regulated by a downstream enhancer. *Plant J.* **68**, 400–412 (2011).

35. K. Watanabe, K. Okada, Two discrete cis elements control the Abaxial side-specific expression of the FILAMENTOUS FLOWER gene in Arabidopsis. *Plant Cell* **15**, 2592–2602 (2003).

36. L. Ye, B. Wang, W. Zhang, H. Shan, H. Kong, Gains and losses of Cis-regulatory elements led to divergence of the Arabidopsis APETALA1 and CAULIFLOWER duplicate genes in the time, space, and level of expression and regulation of one paralog by the other. *Plant Physiol.* **171**, 1055–1069 (2016).

37. M. Abe, T. Takahashi, Y. Komeda, Identification of a cis-regulatory element for L1 layer-specific gene expression, which is targeted by an L1-specific homeodomain protein: Identification of the L1 box. *Plant J.* **26**, 487–494 (2001).

38. M. Kerstens, Y. Boele, A. Moralez-Cruz, C. Roelofsen, P. Wang, L. A. Baumgart, R. O'Malley, G. Sanchez-Perez, B. Scheres, V. Willemsen, Two deeply conserved non-coding sequences control PLETHORA1/2 expression and coordinate embryo and root development. *Plant Commun.* **0**, 101466 (2025).

39. M. Somssich, B. I. Je, R. Simon, D. Jackson, CLAVATA-WUSCHEL signaling in the shoot meristem. *Development* **143**, 3238–3248 (2016).

40. I. Bäurle, T. Laux, Regulation of WUSCHEL transcription in the stem cell niche of the Arabidopsis shoot meristem. *Plant Cell* **17**, 2271–2280 (2005).

41.   T.-Q. Zhang, H. Lian, C.-M. Zhou, L. Xu, Y. Jiao, J.-W. Wang, A two-step model for de Novo activation of WUSCHEL during plant shoot regeneration. *Plant Cell* **29**, 1073–1087 (2017).

42.   Z. Chen, W. Li, C. Gaines, A. Buck, M. Galli, A. Gallavotti, Structural variation at the maize WUSCHEL1 locus alters stem cell organization in inflorescences. *Nat. Commun.* **12**, 2378 (2021).

43.   J. Nardmann, W. Werr, The shoot stem cell niche in angiosperms: expression patterns of WUS orthologues in rice and maize imply major modifications in the course of mono- and dicot evolution. *Mol. Biol. Evol.* **23**, 2492–2504 (2006).

44.   B. I. Je, J. Gruel, Y. K. Lee, P. Bommert, E. D. Arevalo, A. L. Eveland, Q. Wu, A. Goldshmidt, R. Meeley, M. Bartlett, M. Komatsu, H. Sakai, H. Jönsson, D. Jackson, Signaling from maize organ primordia via FASCIATED EAR3 regulates stem cell proliferation and yield traits. *Nat. Genet.* **48**, 785–791 (2016).

45.   M. E. Potok, Y. Wang, L. Xu, Z. Zhong, W. Liu, S. Feng, B. Naranbaatar, S. Rayatpisheh, Z. Wang, J. A. Wohlschlegel, I. Ausin, S. E. Jacobsen, Arabidopsis SWR1-associated protein methyl-CpG-binding domain 9 is required for histone H2A.Z deposition. *Nat. Commun.* **10**, 3352 (2019).

46.   T. Zhao, M. E. Schranz, Network-based microsynteny analysis identifies major differences and genomic outliers in mammalian and angiosperm genomes. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 2165–2174 (2019).

47.   T. H. Kebrom, B. L. Burson, S. A. Finlayson, Phytochrome B represses Teosinte Branched1 expression and induces sorghum axillary bud outgrowth in response to light signals. *Plant Physiol.* **140**, 1109–1117 (2006).

48.   J. A. Aguilar-Martínez, C. Poza-Carrión, P. Cubas, Arabidopsis BRANCHED1 acts as an integrator of branching signals within axillary buds. *Plant Cell* **19**, 458–472 (2007).

49.   J. Doebley, A. Stec, L. Hubbard, The evolution of apical dominance in maize. *Nature* **386**, 485–488 (1997).

50.   R. M. Clark, E. Linton, J. Messing, J. F. Doebley, Pattern of diversity in the genomic region near the maize domestication gene tb1. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 700–707 (2004).

51.   M.-S. Remigereau, G. Lakis, S. Rekima, M. Leveugle, M. C. Fontaine, T. Langin, A. Sarr, T. Robert, Cereal domestication and evolution of branching: evidence for soft selection in the Tb1 orthologue of pearl millet (Pennisetum glaucum [L.] R. Br.). *PLoS One* **6**, e22404 (2011).

52.   X. Wu, Y. Liu, H. Luo, L. Shang, C. Leng, Z. Liu, Z. Li, X. Lu, H. Cai, H. Hao, H.-C. Jing, Genomic footprints of sorghum domestication and breeding selection for multiple end uses. *Mol. Plant* **15**, 537–551 (2022).

53.   A. Studer, Q. Zhao, J. Ross-Ibarra, J. Doebley, Identification of a functional transposon insertion in the maize domestication gene tb1. *Nat. Genet.* **43**, 1160–1163 (2011).

54.   A. Sanyal, B. R. Lajoie, G. Jain, J. Dekker, The long-range interaction landscape of gene promoters. *Nature* **489**, 109–113 (2012).

55. W. A. Ricci, Z. Lu, L. Ji, A. P. Marand, C. L. Ethridge, N. G. Murphy, J. M. Noshay, M. Galli, M. K. Mejía-Guerra, M. Colomé-Tatché, F. Johannes, M. J. Rowley, V. G. Corces, J. Zhai, M. J. Scanlon, E. S. Buckler, A. Gallavotti, N. M. Springer, R. J. Schmitz, X. Zhang, Widespread long-range cis-regulatory elements in the maize genome. *Nat. Plants* **5**, 1237–1249 (2019).

56. C. Liu, C. Wang, G. Wang, C. Becker, M. Zaidem, D. Weigel, Genome-wide analysis of chromatin packing in Arabidopsis thaliana at single-gene resolution. *Genome Res.* **26**, 1057–1068 (2016).

57. Y. Sun, L. Dong, Y. Zhang, D. Lin, W. Xu, C. Ke, L. Han, L. Deng, G. Li, D. Jackson, X. Li, F. Yang, 3D genome architecture coordinates trans and cis regulation of differentially expressed ear and tassel genes in maize. *Genome Biol.* **21**, 143 (2020).

58. E. A. Kellogg, Evolutionary history of the grasses. *Plant Physiol.* **125**, 1198–1205 (2001).

59. D. Singh, T. Hoyt, S. V. Yi, Evolution of Enhancers through Duplication, *Evolutionary Biology* (2025). https://www.biorxiv.org/content/10.1101/2025.04.09.648062v1.abstract.

60. M. Benoit, K. M. Jenike, J. W. Satterlee, S. Ramakrishnan, I. Gentile, A. Hendelman, M. Passalacqua, H. Suresh, H. Shohat, G. M. Robitaille, B. Fitzgerald, M. Alonge, X. Wang, R. Santos, J. He, S. Ou, H. Golan, Y. Green, K. Swartwood, N. G. Karavolias, G. P. Sierra, A. Orejuela, F. Roda, S. Goodwin, W. R. McCombie, E. Kizito, E. Gagnon, S. Knapp, T. Särkinen, A. Frary, J. Gillis, J. Van Eck, M. C. Schatz, Z. Lippman, Solanum pan-genetics reveals paralogues as contingencies in crop engineering. *Nature* **640**, 135–145 (2025).

61. X. Li, X. Zhang, R. J. Schmitz, Using single-cell genomics to explore transcriptional divergence and cis -regulatory dynamics of duplicated genes, *Evolutionary Biology* (2025). https://www.biorxiv.org/content/10.1101/2025.05.07.652760v2.full.pdf.

62. J. M. C. McDonald, R. D. Reed, Beyond modular enhancers: new questions in cis-regulatory evolution. *Trends Ecol. Evol.* **39**, 1035–1046 (2024).

63. P. L. van der Jagt, S. Oud, R. M. A. Vroomans, Ubiquitous systems drift in the evolution of development, *bioRxiv* (2025). https://doi.org/10.1101/2025.04.23.650280.

64. Á. McColgan, J. DiFrisco, Understanding developmental system drift. *Development* **151**, dev203054 (2024).

65. G. Li, L. An, W. Yang, L. Yang, T. Wei, J. Shi, J. Wang, J. H. Doonan, K. Xie, A. R. Fernie, E. S. Lagudah, R. A. Wing, C. Gao, Integrated biotechnological and AI innovations for crop improvement. *Nature* **643**, 925–937 (2025).

66. S. Ou, W. Su, Y. Liao, K. Chougule, J. R. A. Agda, A. J. Hellinga, C. S. B. Lugo, T. A. Elliott, D. Ware, T. Peterson, N. Jiang, C. N. Hirsch, M. B. Hufford, Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol.* **20**, 275 (2019).

67. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

68. J. L. Steenwyk, T. J. Buida 3rd, Y. Li, X.-X. Shen, A. Rokas, ClipKIT: A multiple sequence alignment trimming software for accurate phylogenomic inference. *PLoS Biol.* **18**, e3001007 (2020).

69. A. M. Kozlov, D. Darriba, T. Flouri, B. Morel, A. Stamatakis, RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**, 4453–4455 (2019).

70. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol.* **20**, 238 (2019).

71. One Thousand Plant Transcriptomes Initiative, One thousand plant transcriptomes and the phylogenomics of green plants. *Nat. Commun.*

72. B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von Haeseler, R. Lanfear, IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

73. J. Kim, M. J. Sanderson, Penalized likelihood phylogenetic inference: bridging the parsimony-likelihood gap. *Syst. Biol.* **57**, 665–674 (2008).

74. M. J. Sanderson, Estimating absolute rates of molecular evolution and divergence times: a penalized likelihood approach. *Mol. Biol. Evol.* **19**, 101–109 (2002).

75. E. Paradis, Molecular dating of phylogenies by likelihood methods: a comparison of models and a new information criterion. *Mol. Phylogenet. Evol.* **67**, 436–444 (2013).

76. S. Kumar, M. Suleski, J. M. Craig, A. E. Kasprowicz, M. Sanderford, M. Li, G. Stecher, S. B. Hedges, TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol. Biol. Evol.* **39** (2022).

77. J. P. Rose, T. J. Kleist, S. D. Löfstrand, B. T. Drew, J. Schönenberger, K. J. Sytsma, Phylogeny, historical biogeography, and diversification of angiosperm order Ericales suggest ancient Neotropical and East Asian connections. *Mol. Phylogenet. Evol.* **122**, 59–79 (2018).

78. M. A. Koch, B. Haubold, T. Mitchell-Olds, Comparative evolutionary analysis of chalcone synthase and alcohol dehydrogenase loci in Arabidopsis, Arabis, and related genera (Brassicaceae). *Mol. Biol. Evol.* **17**, 1483–1498 (2000).

79. A. Franzke, D. German, I. A. Al-Shehbaz, K. Mummenhoff, *Arabidopsis* family ties: molecular phylogeny and age estimates in Brassicaceae. *Taxon* **58**, 425–437 (2009).

80. M. A. Beilstein, N. S. Nagalingum, M. D. Clements, S. R. Manchester, S. Mathews, Dated molecular phylogenies indicate a Miocene origin for Arabidopsis thaliana. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 18724–18728 (2010).

81. T. L. P. Couvreur, A. Franzke, I. A. Al-Shehbaz, F. T. Bakker, M. A. Koch, K. Mummenhoff, Molecular phylogenetics, temporal diversification, and principles of evolution in the mustard family (Brassicaceae). *Mol. Biol. Evol.* **27**, 55–71 (2010).

82. S. Kagale, S. J. Robinson, J. Nixon, R. Xiao, T. Huebert, J. Condie, D. Kessler, W. E. Clarke, P. P. Edger, M. G. Links, A. G. Sharpe, I. A. P. Parkin, Polyploid evolution of the Brassicaceae during the Cenozoic era. *Plant Cell* **26**, 2777–2791 (2014).

83. N. Hohmann, E. M. Wolf, M. A. Lysak, M. A. Koch, A time-calibrated road map of Brassicaceae species radiation and evolutionary history. *Plant Cell* **27**, 2770–2784 (2015).

84. C.-H. Huang, R. Sun, Y. Hu, L. Zeng, N. Zhang, L. Cai, Q. Zhang, M. A. Koch, I. Al-Shehbaz, P. P. Edger, J. C. Pires, D.-Y. Tan, Y. Zhong, H. Ma, Resolution of Brassicaceae phylogeny using nuclear genes uncovers nested radiations and supports convergent

morphological evolution. *Mol. Biol. Evol.* **33**, 394–412 (2016).

85. S. Mohammadin, K. Peterse, S. J. van de Kerke, L. W. Chatrou, A. A. Dönmez, K. Mummenhoff, J. C. Pires, P. P. Edger, I. A. Al-Shehbaz, M. E. Schranz, Anatolian origins and diversification of Aethionema, the sister lineage of the core Brassicaceae. *Am. J. Bot.* **104**, 1042–1054 (2017).

86. X. Guo, J. Liu, G. Hao, L. Zhang, K. Mao, X. Wang, D. Zhang, T. Ma, Q. Hu, I. A. Al-Shehbaz, M. A. Koch, Plastome phylogeny and early diversification of Brassicaceae. *BMC Genomics* **18**, 176 (2017).

87. T. Mandáková, Z. Li, M. S. Barker, M. A. Lysak, Diverse genome organization following 13 independent mesopolyploid events in Brassicaceae contrasts with convergent patterns of gene retention. *Plant J.* **91**, 3–21 (2017).

88. X.-C. Huang, D. A. German, M. A. Koch, Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. *Ann. Bot.* **125**, 29–47 (2020).

89. N. Walden, D. A. German, E. M. Wolf, M. Kiefer, P. Rigault, X.-C. Huang, C. Kiefer, R. Schmickl, A. Franzke, B. Neuffer, K. Mummenhoff, M. A. Koch, Nested whole-genome duplications coincide with diversification and high morphological disparity in Brassicaceae. *Nat. Commun.* **11**, 3795 (2020).

90. K. P. Hendriks, C. Kiefer, I. A. Al-Shehbaz, C. D. Bailey, A. Hooft van Huysduynen, L. A. Nikolov, L. Nauheimer, A. R. Zuntini, D. A. German, A. Franzke, M. A. Koch, M. A. Lysak, Ó. Toro-Núñez, B. Özüdoğru, V. R. Invernón, N. Walden, O. Maurin, N. M. Hay, P. Shushkov, T. Mandáková, M. E. Schranz, M. Thulin, M. D. Windham, I. Rešetnik, S. Španiel, E. Ly, J. C. Pires, A. Harkess, B. Neuffer, R. Vogt, C. Bräuchler, H. Rainer, S. B. Janssens, M. Schmull, A. Forrest, A. Guggisberg, S. Zmarzty, B. J. Lepschi, N. Scarlett, F. W. Stauffer, I. Schönberger, P. Heenan, W. J. Baker, F. Forest, K. Mummenhoff, F. Lens, Global Brassicaceae phylogeny based on filtering of 1,000-gene dataset. *Curr. Biol.* **33**, 4052–4068.e6 (2023).

91. Y. Zhao, R. Zhang, K.-W. Jiang, J. Qi, Y. Hu, J. Guo, R. Zhu, T. Zhang, A. N. Egan, T.-S. Yi, C.-H. Huang, H. Ma, Nuclear phylotranscriptomics and phylogenomics support numerous polyploidization events and hypotheses for the evolution of rhizobial nitrogen-fixing symbiosis in Fabaceae. *Mol. Plant* **14**, 748–773 (2021).

92. E. J. M. Koenen, D. I. Ojeda, F. T. Bakker, J. J. Wieringa, C. Kidner, O. J. Hardy, R. T. Pennington, P. S. Herendeen, A. Bruneau, C. E. Hughes, The origin of the legumes is a complex paleopolyploid phylogenomic tangle closely associated with the Cretaceous-Paleogene (K-Pg) mass extinction event. *Syst. Biol.* **70**, 508–526 (2021).

93. Y.-H. Li, G. Zhou, J. Ma, W. Jiang, L.-G. Jin, Z. Zhang, Y. Guo, J. Zhang, Y. Sui, L. Zheng, S.-S. Zhang, Q. Zuo, X.-H. Shi, Y.-F. Li, W.-K. Zhang, Y. Hu, G. Kong, H.-L. Hong, B. Tan, J. Song, Z.-X. Liu, Y. Wang, H. Ruan, C. K. L. Yeung, J. Liu, H. Wang, L.-J. Zhang, R.-X. Guan, K.-J. Wang, W.-B. Li, S.-Y. Chen, R.-Z. Chang, Z. Jiang, S. A. Jackson, R. Li, L.-J. Qiu, De novo assembly of soybean wild relatives for pan-genome analysis of diversity and agronomic traits. *Nat. Biotechnol.* **32**, 1045–1052 (2014).

94. M. Y. Kim, S. Lee, K. Van, T.-H. Kim, S.-C. Jeong, I.-Y. Choi, D.-S. Kim, Y.-S. Lee, D. Park, J. Ma, W.-Y. Kim, B.-C. Kim, S. Park, K.-A. Lee, D. H. Kim, K. H. Kim, J. H. Shin,

5   Y. E. Jang, K. D. Kim, W. X. Liu, T. Chaisan, Y. J. Kang, Y.-H. Lee, K.-H. Kim, J.-K. Moon, J. Schmutz, S. A. Jackson, J. Bhak, S.-H. Lee, Whole-genome sequencing and intensive analysis of the undomesticated soybean (Glycine soja Sieb. and Zucc.) genome. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 22032–22037 (2010).

95.   Y. Zhuang, X. Wang, X. Li, J. Hu, L. Fan, J. B. Landis, S. B. Cannon, J. Grimwood, J. Schmutz, S. A. Jackson, J. J. Doyle, X. S. Zhang, D. Zhang, J. Ma, Phylogenomics of the genus Glycine sheds light on polyploid evolution and life-strategy transition. *Nat. Plants* **8**, 233–244 (2022).

96.   J. C. Villarreal A, B. J. Crandall-Stotler, M. L. Hart, D. G. Long, L. L. Forrest, Divergence times and the evolution of morphological complexity in an early land plant lineage (Marchantiopsida) with a slow molecular rate. *New Phytol.* **209**, 1734–1746 (2016).

97.   C.-H. Huang, C. Zhang, M. Liu, Y. Hu, T. Gao, J. Qi, H. Ma, Multiple polyploidization events across Asteraceae with two nested events in the early history revealed by nuclear phylogenomics. *Mol. Biol. Evol.* **33**, 2820–2835 (2016).

98.   J. R. Mandel, R. B. Dikow, C. M. Siniscalchi, R. Thapa, L. E. Watson, V. A. Funk, A fully resolved backbone phylogeny reveals numerous dispersals and explosive diversifications throughout the history of Asteraceae. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 14083–14088 (2019).

99.   C. Zhang, C.-H. Huang, M. Liu, Y. Hu, J. L. Panero, F. Luebert, T. Gao, H. Ma, Phylotranscriptomic insights into Asteraceae diversity, polyploidy, and morphological innovation. *J. Integr. Plant Biol.* **63**, 1273–1293 (2021).

100.   J. L. Panero, B. S. Crozier, Macroevolutionary dynamics in the early diversification of Asteraceae. *Mol. Phylogenet. Evol.* **99**, 116–132 (2016).

101.   R. Chu, X. Xu, Z. Lu, Y. Ma, H. Cheng, S. Zhu, F. T. Bakker, M. E. Schranz, Z. Wei, Plastome-based phylogeny and biogeography of Lactuca L. (Asteraceae) support revised lettuce gene pool categories. *Front. Plant Sci.* **13**, 978417 (2022).

102.   N. Kilian, A. Sennikov, Z.-H. Wang, B. Gemeinholzer, J.-W. Zhang, Sub-Paratethyan origin and Middle to Late Miocene principal diversification of the Lactucinae (Compositae: Cichorieae) inferred from molecular phylogenetics, divergence-dating and biogeographic analysis. *Taxon* **66**, 675–703 (2017).

103.   K. Tremetsberger, B. Gemeinholzer, H. Zetzsche, S. Blackmore, N. Kilian, S. Talavera, Divergence time estimation in Cichorieae (Asteraceae) using a fossil-calibrated relaxed molecular clock. *Org. Divers. Evol.* **13**, 1–13 (2013).

104.   J. Bechteler, G. Peñaloza-Bojacá, D. Bell, J. Gordon Burleigh, S. F. McDaniel, E. Christine Davis, E. B. Sessa, A. Bippus, D. Christine Cargill, S. Chantanoarrapint, I. Draper, L. Endara, L. L. Forrest, R. Garilleti, S. W. Graham, S. Huttunen, J. J. Lazo, F. Lara, J. Larraín, L. R. Lewis, D. G. Long, D. Quandt, K. Renzaglia, A. Schäfer-Verwimp, G. E. Lee, A. M. Sierra, M. von Konrat, C. E. Zartman, M. R. Pereira, B. Goffinet, J. C. Villarreal A, Comprehensive phylogenomic time tree of bryophytes reveals deep relationships and uncovers gene incongruences in the last 500 million years of diversification. *Am. J. Bot.* **110**, e16249 (2023).

105.   B. J. Harris, J. W. Clark, D. Schrempf, G. J. Szöllősi, P. C. J. Donoghue, A. M. Hetherington, T. A. Williams, Divergent evolutionary trajectories of bryophytes and

5   tracheophytes from a complex common ancestor of land plants. *Nat. Ecol. Evol.* **6**, 1634–1643 (2022).

106.   J. D. Buenrostro, B. Wu, U. M. Litzenburger, D. Ruff, M. L. Gonzales, M. P. Snyder, H. Y. Chang, W. J. Greenleaf, Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**, 486–490 (2015).

107.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

108.   P. S. Hosmani, M. Flores-Gonzalez, H. van de Geest, F. Maumus, L. V. Bakker, E. Schijlen, J. van Haarst, J. Cordewener, G. Sanchez-Perez, S. Peters, Z. Fei, J. J. Giovannoni, L. A. Mueller, S. Saha, An improved de novo assembly and annotation of the tomato reference genome using single-molecule sequencing, Hi-C proximity ligation and optical maps, *Genomics* (2019). https://www.biorxiv.org/content/10.1101/767764v1.abstract.

109.   H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM, *arXiv [q-bio.GN]* (2013). http://arxiv.org/abs/1303.3997.

110.   H. Zhao, Z. Sun, J. Wang, H. Huang, J.-P. Kocher, L. Wang, CrossMap: a versatile tool for coordinate conversion between genome assemblies. *Bioinformatics* **30**, 1006–1007 (2014).

111.   X. Tu, M. K. Mejía-Guerra, J. A. Valdes Franco, D. Tzeng, P.-Y. Chu, W. Shen, Y. Wei, X. Dai, P. Li, E. S. Buckler, S. Zhong, Reconstructing the maize leaf regulatory network using ChIP-seq data of 104 transcription factors. *Nat. Commun.* **11**, 5089 (2020).

112.   M. Galli, A. Khakhar, Z. Lu, Z. Chen, S. Sen, T. Joshi, J. L. Nemhauser, R. J. Schmitz, A. Gallavotti, The DNA binding landscape of the maize AUXIN RESPONSE FACTOR family. *Nat. Commun.* **9**, 4526 (2018).

113.   M. Galli, Z. Chen, T. Ghandour, A. Chaudhry, J. Gregory, F. Feng, M. Li, N. Schleif, X. Zhang, Y. Dong, G. Song, J. W. Walley, G. Chuck, C. Whipple, H. F. Kaeppler, S.-S. C. Huang, A. Gallavotti, Transcription factor binding divergence drives transcriptional and phenotypic variation in maize. *Nat. Plants* **11**, 1205–1219 (2025).

114.   W. J. Kent, A. S. Zweig, G. Barber, A. S. Hinrichs, D. Karolchik, BigWig and BigBed: enabling browsing of large distributed datasets. *Bioinformatics* **26**, 2204–2207 (2010).

115.   G. Ambrosini, R. Groux, P. Bucher, PWMScan: a fast tool for scanning entire genomes with a position-specific weight matrix. *Bioinformatics* **34**, 2483–2484 (2018).

116.   K. N. Chang, S. Zhong, M. T. Weirauch, G. Hon, M. Pelizzola, H. Li, S.-S. C. Huang, R. J. Schmitz, M. A. Urich, D. Kuo, J. R. Nery, H. Qiao, A. Yang, A. Jamali, H. Chen, T. Ideker, B. Ren, Z. Bar-Joseph, T. R. Hughes, J. R. Ecker, Temporal transcriptional response to ethylene gas drives growth hormone cross-regulation in Arabidopsis. *Elife* **2**, e00675 (2013).

117.   Y. Zhang, O. Mayba, A. Pfeiffer, H. Shi, J. M. Tepperman, T. P. Speed, P. H. Quail, A quartet of PIF bHLH factors provides a transcriptionally centered signaling hub that regulates seedling morphogenesis through differential expression-patterning of shared target genes in Arabidopsis. *PLoS Genet.* **9**, e1003244 (2013).

118.   W. Deng, H. Ying, C. A. Helliwell, J. M. Taylor, W. J. Peacock, E. S. Dennis, FLOWERING LOCUS C (FLC) regulates development pathways throughout the life cycle

of Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 6680–6685 (2011).

119.   N. Gonzalez, L. Pauwels, A. Baekelandt, L. De Milde, J. Van Leene, N. Besbrugge, K. S. Heyndrickx, A. Cuéllar Pérez, A. N. Durand, R. De Clercq, E. Van De Slijke, R. Vanden Bossche, D. Eeckhout, K. Gevaert, K. Vandepoele, G. De Jaeger, A. Goossens, D. Inzé, A repressor protein complex regulates leaf growth in Arabidopsis. *Plant Cell* **27**, 2273–2287 (2015).

120.   A. Ritter, S. Iñigo, P. Fernández-Calvo, K. S. Heyndrickx, S. Dhondt, H. Shi, L. De Milde, R. Vanden Bossche, R. De Clercq, D. Eeckhout, M. Ron, D. E. Somers, D. Inzé, K. Gevaert, G. De Jaeger, K. Vandepoele, L. Pauwels, A. Goossens, The transcriptional repressor complex FRS7-FRS12 regulates flowering time and growth in Arabidopsis. *Nat. Commun.* **8**, 15235 (2017).

121.   D. Ezer, S. J. K. Shepherd, A. Brestovitsky, P. Dickinson, S. Cortijo, V. Charoensawan, M. S. Box, S. Biswas, K. E. Jaeger, P. A. Wigge, The G-box transcriptional regulatory code in Arabidopsis. *Plant Physiol.* **175**, 628–640 (2017).

122.   P. Hornitschek, M. V. Kohnen, S. Lorrain, J. Rougemont, K. Ljung, I. López-Vidriero, J. M. Franco-Zorrilla, R. Solano, M. Trevisan, S. Pradervand, I. Xenarios, C. Fankhauser, Phytochrome interacting factors 4 and 5 control seedling growth in changing light conditions by directly controlling auxin signaling: PIF4 and PIF5 control auxin signaling. *Plant J.* **71**, 699–711 (2012).

123.   A. Verkest, T. Abeel, K. S. Heyndrickx, J. Van Leene, C. Lanz, E. Van De Slijke, N. De Winne, D. Eeckhout, G. Persiau, F. Van Breusegem, D. Inzé, K. Vandepoele, G. De Jaeger, A generic tool for transcription factor target gene discovery in Arabidopsis cell suspension cultures based on tandem chromatin affinity purification. *Plant Physiol.* **164**, 1122–1133 (2014).

124.   F. Chen, B. Li, G. Li, J.-B. Charron, M. Dai, X. Shi, X. W. Deng, Arabidopsis phytochrome A directly targets numerous promoters for individualized modulation of genes in a wide range of pathways. *Plant Cell* **26**, 1949–1966 (2014).

125.   Y. Kurihara, Y. Makita, M. Kawashima, H. Hamasaki, Y. Y. Yamamoto, M. Matsui, Next-generation sequencing of genomic DNA fragments bound to a transcription factor in vitro reveals its regulatory potential. *Genes (Basel)* **5**, 1115–1131 (2014).

126.   U. V. Pedmale, S.-S. C. Huang, M. Zander, B. J. Cole, J. Hetzel, K. Ljung, P. A. B. Reis, P. Sridevi, K. Nito, J. R. Nery, J. R. Ecker, J. Chory, Cryptochromes interact directly with PIFs to control plant growth in limiting blue light. *Cell* **164**, 233–245 (2016).

127.   A. Pfeiffer, H. Shi, J. M. Tepperman, Y. Zhang, P. H. Quail, Combinatorial complexity in a transcriptionally centered signaling hub in Arabidopsis. *Mol. Plant* **7**, 1598–1618 (2014).

128.   K. Kaufmann, F. Wellmer, J. M. Muiño, T. Ferrier, S. E. Wuest, V. Kumar, A. Serrano-Mislata, F. Madueño, P. Krajewski, E. M. Meyerowitz, G. C. Angenent, J. L. Riechmann, Orchestration of floral initiation by APETALA1. *Science* **328**, 85–89 (2010).

129.   L. Yant, J. Mathieu, T. T. Dinh, F. Ott, C. Lanz, H. Wollmann, X. Chen, M. Schmid, Orchestration of the floral transition and floral development in Arabidopsis by the bifunctional transcription factor APETALA2. *Plant Cell* **22**, 2156–2170 (2010).

130.   E. Moyroud, E. G. Minguet, F. Ott, L. Yant, D. Posé, M. Monniaux, S. Blanchet, O.

Bastien, E. Thévenon, D. Weigel, M. Schmid, F. Parcy, Prediction of regulatory interactions from genome sequences using a biophysical model for the Arabidopsis LEAFY transcription factor. *Plant Cell* **23**, 1293–1306 (2011).

131. W. Huang, P. Pérez-García, A. Pokhilko, A. J. Millar, I. Antoshechkin, J. L. Riechmann, P. Mas, Mapping the core of the Arabidopsis circadian clock defines the network structure of the oscillator. *Science* **336**, 75–79 (2012).

132. R. G. H. Immink, D. Posé, S. Ferrario, F. Ott, K. Kaufmann, F. L. Valentim, S. de Folter, F. van der Wal, A. D. J. van Dijk, M. Schmid, G. C. Angenent, Characterization of SOC1's central role in flowering by the identification of its upstream and downstream regulators. *Plant Physiol.* **160**, 433–449 (2012).

133. S. E. Wuest, D. S. O'Maoileidigh, L. Rae, K. Kwasniewska, A. Raganelli, K. Hanczaryk, A. J. Lohan, B. Loftus, E. Graciet, F. Wellmer, Molecular basis for the specification of floral organs by APETALA3 and PISTILLATA. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 13452–13457 (2012).

134. N. Nakamichi, T. Kiba, M. Kamioka, T. Suzuki, T. Yamashino, T. Higashiyama, H. Sakakibara, T. Mizuno, Transcriptional repressor PRR5 directly regulates clock-output pathways. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 17123–17128 (2012).

135. T. Liu, J. Carlsson, T. Takeuchi, L. Newton, E. M. Farré, Direct regulation of abiotic responses by the Arabidopsis circadian clock component PRR7. *Plant J.* **76**, 101–114 (2013).

136. P. Merelo, Y. Xie, L. Brand, F. Ott, D. Weigel, J. L. Bowman, M. G. Heisler, S. Wenkel, Genome-wide identification of KANADI1 target genes. *PLoS One* **8**, e77341 (2013).

137. J. Heyman, T. Cools, F. Vandenbussche, K. S. Heyndrickx, J. Van Leene, I. Vercauteren, S. Vanderauwera, K. Vandepoele, G. De Jaeger, D. Van Der Straeten, L. De Veylder, ERF115 controls root quiescent center cell division and stem cell replenishment. *Science* **342**, 860–863 (2013).

138. M. Fan, M.-Y. Bai, J.-G. Kim, T. Wang, E. Oh, L. Chen, C. H. Park, S.-H. Son, S.-K. Kim, M. B. Mudgett, Z.-Y. Wang, The bHLH transcription factor HBI1 mediates the trade-off between growth and pathogen-associated molecular pattern-triggered immunity in Arabidopsis. *Plant Cell* **26**, 828–841 (2014).

139. E. Oh, J.-Y. Zhu, M.-Y. Bai, R. A. Arenhart, Y. Sun, Z.-Y. Wang, Cell elongation is regulated through a central circuit of interacting transcription factors in the Arabidopsis hypocotyl. *Elife* **3** (2014).

140. F. Chen, B. Li, J. Demone, J.-B. Charron, X. Shi, X. W. Deng, Photoreceptor partner FHY1 has an independent role in gene modulation and plant development under far-red light. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 11888–11893 (2014).

141. O. S. Lau, K. A. Davies, J. Chang, J. Adrian, M. H. Rowe, C. E. Ballenger, D. C. Bergmann, Direct roles of SPEECHLESS in the specification of stomatal self-renewing cells. *Science* **345**, 1605–1609 (2014).

142. H. Zhang, X. Zhao, J. Li, H. Cai, X. W. Deng, L. Li, MicroRNA408 is critical for the HY5-SPL7 gene network that mediates the coordinated response to light and copper. *Plant Cell* **26**, 4933–4953 (2014).

143. S. Liu, B. Kracher, J. Ziegler, R. P. Birkenbihl, I. E. Somssich, Negative regulation of

ABA signaling by WRKY33 is critical for Arabidopsis immunity towards Botrytis cinerea 2100. *Elife* **4**, e07295 (2015).

144.   N. Marín-de la Rosa, A. Pfeiffer, K. Hill, A. Locascio, R. P. Bhalerao, P. Miskolczi, A. L. Grønlund, A. Wanchoo-Kohli, S. G. Thomas, M. J. Bennett, J. U. Lohmann, M. A. Blázquez, D. Alabadí, Genome wide binding site analysis reveals transcriptional coactivation of cytokinin-responsive genes by DELLA proteins. *PLoS Genet.* **11**, e1005337 (2015).

145.   M. Kamioka, S. Takao, T. Suzuki, K. Taki, T. Higashiyama, T. Kinoshita, N. Nakamichi, Direct repression of evening genes by CIRCADIAN CLOCK-ASSOCIATED1 in the Arabidopsis circadian clock. *Plant Cell* **28**, 696–711 (2016).

146.   C. Sayou, M. H. Nanao, M. Jamin, D. Posé, E. Thévenon, L. Grégoire, G. Tichtinsky, G. Denay, F. Ott, M. Peirats Llobet, M. Schmid, R. Dumas, F. Parcy, A SAM oligomerization domain shapes the genomic binding landscape of the LEAFY transcription factor. *Nat. Commun.* **7**, 11222 (2016).

147.   R. C. O'Malley, S.-S. C. Huang, L. Song, M. G. Lewsey, A. Bartlett, J. R. Nery, M. Galli, A. Gallavotti, J. R. Ecker, Cistrome and epicistrome features shape the regulatory DNA landscape. *Cell* **165**, 1280–1292 (2016).

148.   S. Bencivenga, A. Serrano-Mislata, M. Bush, S. Fox, R. Sablowski, Control of oriented tissue growth through repression of organ boundary genes promotes stem morphogenesis. *Dev. Cell* **39**, 198–208 (2016).

149.   R. P. Birkenbihl, B. Kracher, M. Roccaro, I. E. Somssich, Induced genome-wide binding of three Arabidopsis WRKY transcription factors during early MAMP-triggered immunity. *Plant Cell* **29**, 20–38 (2017).

150.   Y. O. Zubo, I. C. Blakley, M. V. Yamburenko, J. M. Worthen, I. H. Street, J. M. Franco-Zorrilla, W. Zhang, K. Hill, T. Raines, R. Solano, J. J. Kieber, A. E. Loraine, G. E. Schaller, Cytokinin induces genome-wide binding of the type-B response regulator ARR10 to regulate growth and development in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5995–E6004 (2017).

151.   Z. Lu, B. T. Hofmeister, C. Vollmers, R. M. DuBois, R. J. Schmitz, Combining ATAC-seq with nuclei sorting for discovery of cis-regulatory regions in plant genomes. *Nucleic Acids Res.* **45**, e41 (2017).

152.   C. M. Alexandre, J. R. Urton, K. Jean-Baptiste, J. Huddleston, M. W. Dorrity, J. T. Cuperus, A. M. Sullivan, F. Bemm, D. Jolic, A. A. Arsovski, A. Thompson, J. L. Nemhauser, S. Fields, D. Weigel, K. L. Bubb, C. Queitsch, Complex relationships between chromatin accessibility, sequence divergence, and gene expression in Arabidopsis thaliana. *Mol. Biol. Evol.* **35**, 837–854 (2018).

153.   Y. Liu, T. Tian, K. Zhang, Q. You, H. Yan, N. Zhao, X. Yi, W. Xu, Z. Su, PCSD: a plant chromatin state database. *Nucleic Acids Res.* **46**, D1157–D1167 (2018).

154.   Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nusbaum, R. M. Myers, M. Brown, W. Li, X. S. Liu, Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

155.   H. Stroud, M. V. C. Greenberg, S. Feng, Y. V. Bernatavichute, S. E. Jacobsen, Comprehensive analysis of silencing mutants reveals complex regulation of the Arabidopsis

methylome. *Cell* **152**, 352–364 (2013).

156. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* **16**, 284–287 (2012).

157. AnnotationHub, *Bioconductor*. https://bioconductor.org/packages/AnnotationHub.

158. J. Van Eck, P. Keen, M. Tjahjadi, Agrobacterium tumefaciens-mediated transformation of tomato. *Methods Mol. Biol.* **1864**, 225–234 (2019).

159. D. Rodríguez-Leal, Z. H. Lemmon, J. Man, M. E. Bartlett, Z. B. Lippman, Engineering quantitative trait variation for crop improvement by genome editing. *Cell* **171**, 470–480.e8 (2017).

160. S. Soyk, Z. H. Lemmon, M. Oved, J. Fisher, K. L. Liberatore, S. J. Park, A. Goren, K. Jiang, A. Ramos, E. van der Knaap, J. Van Eck, D. Zamir, Y. Eshed, Z. B. Lippman, Bypassing negative epistasis on yield in tomato imposed by a domestication gene. *Cell* **169**, 1142–1155.e12 (2017).

161. S. Werner, C. Engler, E. Weber, R. Gruetzner, S. Marillonnet, Fast track assembly of multigene constructs using Golden Gate cloning and the MoClo system. *Bioeng. Bugs* **3**, 38–43 (2012).

162. D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2: Improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

163. G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, Ggtree: An r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

164. L.-G. Wang, T. T.-Y. Lam, S. Xu, Z. Dai, L. Zhou, T. Feng, P. Guo, C. W. Dunn, B. R. Jones, T. Bradley, H. Zhu, Y. Guan, Y. Jiang, G. Yu, Treeio: An R package for phylogenetic tree input and output with richly annotated and associated data. *Mol. Biol. Evol.* **37**, 599–603 (2020).

165. T. Hackl, M. Ankenbrand, B. van Adrichem, D. Wilkins, K. Haslinger, gggenomes: effective and versatile visualizations for comparative genomics, *arXiv [q-bio.GN]* (2024). http://arxiv.org/abs/2411.13556.

166. P. Danecek, J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T. Keane, S. A. McCarthy, R. M. Davies, H. Li, Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).

167. A. M. Waterhouse, J. B. Procter, D. M. A. Martin, M. Clamp, G. J. Barton, Jalview Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009).

168. Z. B. Lippman, O. Cohen, J. P. Alvarez, M. Abu-Abied, I. Pekker, I. Paran, Y. Eshed, D. Zamir, The making of a compound inflorescence in tomato and related nightshades. *PLoS Biol.* **6**, e288 (2008).

169. N. Servant, N. Varoquaux, B. R. Lajoie, E. Viara, C.-J. Chen, J.-P. Vert, E. Heard, J. Dekker, E. Barillot, HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol.* **16**, 259 (2015).

170. M. Alonge, L. Lebeigle, M. Kirsche, K. Jenike, S. Ou, S. Aganezov, X. Wang, Z. B.

Lippman, M. C. Schatz, S. Soyk, Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. *Genome Biol.* **23**, 258 (2022).

171.   T. Yang, F. Zhang, G. G. Yardımcı, F. Song, R. C. Hardison, W. S. Noble, F. Yue, Q. Li, HiCRep: assessing the reproducibility of Hi-C data using a stratum-adjusted correlation coefficient. *Genome Res.* **27**, 1939–1949 (2017).

172.   N. Abdennur, L. A. Mirny, Cooler: scalable storage for Hi-C data and other genomically labeled arrays. *Bioinformatics* **36**, 311–316 (2020).

173.   A. Kaul, S. Bhattacharyya, F. Ay, Identifying statistically significant chromatin contacts from Hi-C data with FitHiC2. *Nat. Protoc.* **15**, 991–1012 (2020).

174.   H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).

175.   K. A. Riemondy, R. M. Sheridan, A. Gillen, Y. Yu, C. G. Bennett, J. R. Hesselberth, valr: Reproducible genome interval analysis in R. *F1000Res.* **6**, 1025 (2017).

176.   X. Qiao, Q. Li, H. Yin, K. Qi, L. Li, R. Wang, S. Zhang, A. H. Paterson, Gene duplication and evolution in recurring polyploidization-diploidization cycles in plants. *Genome Biol.* **20**, 38 (2019).

177.   N. Uchida, B. Townsley, K.-H. Chung, N. Sinha, Regulation of SHOOT MERISTEMLESS genes via an upstream-conserved noncoding sequence coordinates leaf development. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 15953–15958 (2007).

Acknowledgments: We thank the members of the Bartlett, Lippman, and Efroni laboratories for their support and feedback; T. Harrington, G. Robitaille, B. Seman for technical assistance; T. Mulligan, K. Schlecht, S. Qiao, and B. Fitzgerald for assistance with plant care; H. Golan for website design.

Funding:

Binational Science Foundation 2021660 (IE)

Israeli Science Foundation grant 928/22 (IE)

Howard Hughes Medical Institute International Research Scholar grant 55008730 (IE)

National Science Foundation Plant Genome Research Program grant IOS-2129189 (DJ, MEB, ZBL)

USDA-NIFA Postdoctoral Fellowship (KRA)

Gatsby Foundation (MEB)

Howard Hughes Medical Institute (ZBL)

Author contributions:

Conceptualization: DJ, MEB, ZBL, IE

Investigation: KRA, AH, DC, HY, AEdN, ST, AS, MEB, ZBL, IE

Software: AH, AS, IE

5   Funding acquisition: DJ, MEB, ZBL, IE

Supervision: MEB, ZBL, IE

Writing – original draft: KRA, AH, MEB, ZBL, IE

Writing – reviewing and editing: KRA, AH, DJ, MEB, ZBL, IE

10   **Competing interests:** Z.B.L and D.J. are consultants for Inari Agriculture, and Z.B.L is a member of their scientific strategy board. All other authors declare that they have no competing interests.

Data and materials availability: Raw and processed data for Conservatory are available from
15   www.conservatorycns.com;   Conservatory   code   is   available   at https://github.com/idanefroni/Conservatory; Tomato Hi-C and ATAC-Seq raw data are available from NCBI Gene Expression Omnibus record GSE307325.

Supplementary Materials
Materials and Methods
20   Figs. S1 to S5
Tables S1 to S6
References (66-177)

Fig. 1. Identification of deeply conserved plant *cis*-regulatory elements. (**A**) Phylogeny of the
species used in this study. The six families with reference species are colored. (**B**) Median CNS
length per genome. (**C**) Enrichment of DNA methylation, histone modification, open chromatin,
10   and transcription factor binding sites in CNSs, compared to intergenic background regions in
arabidopsis and maize. (**D**) Distribution of the percentages of accessible chromatin regions
(ACRs) of arabidopsis, tomato, and maize that overlap CNSs (pink) versus control intergenic
regions (gray). (**E**) Position distribution within overlapping ACRs for CNSs (pink) or control
intergenic regions (gray). CNS are enriched at ACR midpoints. (**F**) Percentage of CNSs vs
15   control background (bkg) sequences that overlap ACRs. (**G**) Frequency histograms of CNS age
estimations. (**H**) Functional enrichment of arabidopsis genes associated with angiosperm and
seed plant CNSs.

Fig. 2. CRISPR-Cas9 mutagenesis of ancient CNSs near *HOMEOBOX* genes disrupts embryonic and vegetative development. (**A**) CNS annotations of the *SlWOX9* promoter. Distinct modules controlling expression in the embryo and shoot apical meristem (pro-Reg1) or inflorescences (pro-Reg2) (*9*) are outlined in gray. Multiple sequence alignment of seed plant

10   CNS (S217) within pro-Reg1 for ten representative angiosperm species, with a conserved WRKY/NAC motif in pink. See table S4 for gene names. (**B**) Schematic of five *slwox9^Pro-Reg1-TFBS^* alleles targeting CNS S217. (**C**) Percentages of offspring with vegetative defects in wild type (WT) and in five F2 families segregating *slwox9^Pro-Reg1-TFBS^* alleles. (**D**) CNS annotations of the *SlWOX2* promoter and a *slwox2^Pro^* allelic series generated by CRISPR-Cas9. Dashed red lines

15   indicate deletions. Black and gray skulls mark fully or partially expressive embryonic lethal

alleles, respectively. The right panel shows genotypic proportions of F2 populations and chi-square statistics under a 1:2:1 wild type:heterozygous:homozygous expectation. Note the top 2 rows are *Slwox2* CRISPR null alleles. (**E**) Detailed view of deeply conserved *SlWOX2^{Pro}* region, showing deletions in the different *Slwox2Pro* alleles. (**F**) Representative vegetative phenotypes of *slwox2^{Pro-6}* homozygotes, with multiple cotyledons (red arrowheads) and leaf fusion (white asterisk), and *slwox2^{Pro-3}* homozygotes with a terminated SAM (white arrowhead).

Fig. 3. CNSs associated with homologs of *WUSCHEL* and *TEOSINTE BRANCHED1* illustrate principles of plant *cis-regulatory* evolution. (**A**) CNS annotation of the *AtWUS* promoter. The conserved regulatory module specifying *WUS* expression in the SAM (*40*) is shown at high detail, with multiple sequence alignment of the seed plant CNS S115 for twelve representative angiosperm species. The red line marks the position of a *B. distachyon*-specific insertion that was omitted for visualization purposes. (**B**) Presence-absence matrix (left) and microsynteny plot (right) of deeply conserved CNSs in the *WUS* meristem module. Asterisks indicate intragenic duplication of a CNS. (**C**) CNS annotation of the *ZmTB1* promoter. ACRs on either side of a *Hopscotch* transposon each contain six CNSs conserved in the grass family. Shown below are multiple sequence alignments of CNSs P10 and P11 in twelve selected grass

5   species, and the presence-absence matrix of these CNSs in (**D**). P1-P6 are specific to one clade, while P7-P12 are broadly conserved among grasses. (**E**) Distributions of CNS distances from the translation start sites of *WUS* and *TB1* orthologs. (**F**) Microsynteny plot of *TB1* orthologs. The P1-12 block is highlighted in purple. Note the varied position of this CNS group in different grasses. See table S4 for gene names.

Fig. 4. Conservation of CNS position and gene associations. (**A**) Distributions of CNS-gene distances, stratified by CNS conservation level. (**B**) Proportions of CNSs associated with the closest genes, with other adjacent genes, or that skip adjacent genes, for each of our ten reference genomes. (**C**) Proportions of intergenic chromatin loop edges in arabidopsis, tomato, and maize, stratified by overlap with at least one CNS and whether the other edge of the loop connects to a gene. Chromatin loops connecting CNSs with genes are more prevalent than those that do not overlap CNSs ($p < 0.001$; chi-square test). (**D**) Proportions of CNS-containing intergenic-genic chromatin loops with one or more concordant CNS-gene associations. (**E**) Example of long-range CNS-gene association supported by chromatin looping. (**F**) Distribution of CNS mobility (standard deviation of relative position in orthogroups of taxonomic family) of ancient (angiosperm and seed plant) vs family level CNS ($p < 2.2e\text{-}16$; Wilcoxon test).

Fig. 5. Diversification of CNSs throughout plant evolution. (**A**) Presence-absence matrix of angiosperm CNSs among all 314 genomes in this study. Rows correspond to genomes, columns to angiosperm CNSs, and tiles are color-coded by taxonomic family. The lower right panel highlights outlier monocot species. (**B**) Lineage-specific loss of angiosperm CNSs. Each line

10   tracks the number of angiosperm CNSs in a single genome from 200 Mya to the present day. Lines are color-coded according to the taxonomic family. (**C**) The normalized CNS number in diverging paralogs resulting from whole-genome duplications in 10 reference genomes. Shown is the number of shared CNS amongst the paralogs, paralog-specific CNS, and the number in each of the paralogs. To control for variation in CNS number between genomes, the CNS count per

15   gene was normalized to the mean CNS count per gene in the relevant genome. (**D**) Enrichment of angiosperm-level CNS retention in duplicate gene pairs from the reference genomes. Two genomes were used from each family. As paralogs diverged, the preference to maintain the angiosperm-level CNS increased. (**E**) Distribution of CNS copy number stratified by CNS conservation level. (**F**) Distribution of pairwise identity for nonsyntenic CNSs, for syntenic

20   CNSs with different names, and syntenic orthologous CNSs. Syntenic CNS were more similar than Non-syntenic ones (p<0.001; Wilcoxon test).

Materials and Methods

Conservatory algorithm

The Conservatory algorithm was written in perl and R. All scripts, installation instructions and related data files are available at https://github.com/idanefroni/Conservatory/. All references to scripts below are to those in the github scripts/ directory. Output files, including raw data, calibrated trees, and GFF3 tracks for all 314 genomes used in this study, are provided at www.conservatorycns.com. The conservatory algorithm was run on the Hebrew University High Performance Computing cluster, using up to 1,000 nodes at a time. Overall run time for producing and merging the entire dataset was approximately 7 days. Jobs used to run the algorithm are provided in the jobs folder of the GitHub directory.

1. Genome resources

Genome assemblies and annotations were downloaded from public repositories, including NCBI, Ensembl, CoGe, Phytozome, and project-specific websites as described in table S1. We analyzed 314 genomes from 68 plant families. For each genome, we obtained FASTA format files containing genome and protein sequences, and GFF3 gene annotation files. Ten genome assemblies were designated as family-specific reference genomes: *Arabidopsis thaliana* and *Arabis alpina* for Brassicaceae, *Medicago truncatula* and *Glycine max* for Fabaceae, *Lemna minor* strain 7210 for Lemnaceae, *Brachypodium distachyon* and *Zea mays* for Poaceae, *Lactuca saligna* for the Asteraceae, and *Solanum lycopersicum* and *Physalis grisea* for the Solanaceae. For these ten reference genomes, we generated repeat annotation using EDTA *(66)* with default settings.

2. Phylogenetic inference

Conservatory requires two types of phylogenetic trees. First, family-specific rate-calibrated trees are required for each reference species. Second, a single all-species tree with age estimation for each node. To generate phylogenies for each of the reference families, we used our initial orthogroup assignments to identify low and single-copy nuclear genes in each family. The predicted peptide sequences for each of these orthogroups were aligned with mafft (v7.313) *(67)*. Alignments were trimmed with clipkit *(68)* and concatenated. Constraint trees were generated for all families, using published phylogenies. Maximum likelihood tree inference was then carried out with RAxML-NG version 1.1.0 *(69)*, providing the concatenated orthogroup alignments, a constraint tree, and partitions, and specifying a JTT+G substitution model.

To generate the all-species tree, we clustered the predicted peptide sequences from each genome into orthogroups using OrthoFinder *(70)*. To identify low-copy orthogroups suitable for species tree construction, we aligned tomato peptide sequences from the unmasked multiple sequence alignments of the 1KP project *(71)*, to tomato SL4.0 predicted peptides with blastp at default settings, retaining the top-scoring hit for each 1KP peptide. This recovered 386 of the 410 1KP orthogroups. As a quality control, we inspected the distribution of these orthogroups present in each genome, and the number of genomes with each orthogroup. Among 160 orthogroups present in 300-500 total copies among all genomes, we selected 84 orthogroups at random. Another 16 orthogroups were specifically chosen for *Apium graveolens* and *Nelumbo nucifera*

genes, for a total of 100 orthogroups. Alignments and phylogenetic inference were conducted as for the family trees, except phylogenetic inference was performed using IQTree2 version 2.0.3 *(72)*. Node ages were estimated with the penalized likelihood approach *(73, 74)* implemented in the R package ape *(75)*. We used a relaxed model of substitution rate variation among branches and up to 80 alternative iterations between estimating rates and dates. Calibration points were obtained from TimeTree *(76)* and from the literature *(77–105)* and are available from the authors upon request.

3. Genome and mapping database construction

Whole-genome sequences and gene annotations (in GFF3 format) were preprocessed using the processGenomes script, which standardized FASTA headers, indexed sequences for rapid retrieval, and parsed gene names to generate standardized gene coordinate files. Repetitive sequences for all reference species were masked based on the EDTA annotation (see above). Regulatory sequences were extracted using the buildRegulatorySeqDB script. For each gene, non-coding regions were extracted from the genome. The length of upstream regulatory regions was determined as the 97th quantile of intergenic region lengths, with a maximum of 100 kb. The downstream regulatory region was one-fifth the length of the upstream region. Upstream sequences were taken from the translation start site, and downstream from the translation end site. To prevent contamination by coding sequence, CDS were masked based on genome annotation.

Homolog databases were created using the buildOrthologDB script. First, reciprocal all-vs-all BLAST against all other genomes in the database was performed. Each gene pair was accepted as orthologous if each gene was the top BLAST hit of the other in the corresponding search direction. In cases where multiple high-scoring hits existed, a cut-off E-value was calculated at 0.66 of the maximum E-value score. Up to 16 genes with an E-value higher than the cutoff were defined as a group of homologs. Sequences shorter than 120 amino acids were treated with higher sensitivity, using relaxed E-value cutoffs. The final output was a tab-delimited table containing homolog groups with associated alignment scores.

CNS identification

Identification of CNS for a given gene was performed using buildConservation script. For each gene in the reference species, candidate homolog genes were identified in the target species using the homolog mapping database. The regulatory regions of the gene were filtered for low complexity regions and then aligned using LastZ (--format=maf --gap=200,100 --nochain --noytrim --seed=match4 --gapped --strand=both --step=1 --ambiguous=iupac --identity=70 --ydrop=1000 --hspthreshold=2000 --gappedthresh=2000) and alignment scores calculated. The highest-scoring homolog was selected as the putative ortholog; however, if multiple candidates had comparable high scores, multiple orthologs were retained per species to account for recent duplications or lineage-specific expansions. For species that had multiple available genomes, the ortholog from the genome with the best alignment score for the regulatory regions was selected.

Alignments were refined by trimming poorly aligned ends and merging nearby high-scoring alignment blocks. Conservation within the aligned sequences was assessed using *phyloP (27)*, using the family tree, with rates estimated from the background substitution rates, as

detailed above. Conserved blocks were defined as regions with phyloP p-values lower than a species-specific p-value cut-off and over stretches of at least 15 bp. The species-specific values were selected so that, on average, the number of conserved bp per gene is 250. Candidate conserved blocks were filtered for a minimum sequence identity of over 70% and a minimum length threshold of 8, and regions overlapping masked or low-quality genomic regions were removed.

For each CNS, ancestral sequences were reconstructed. First, sequences were aligned using mafft (--auto --thread -1 --ep 0.3), followed by *FastML* (-b -qf -mh) using the family tree. Then, each reconstructed ancestral sequence was aligned with homologous gene regulatory sequences of all genomes from outside the family using sensitive local alignment with relaxed parameters (--HSPThreshold=1600). Hits were retained if they met identity (> 70%) and length (> 12 bp) thresholds.

To obtain a unified CNS map for a reference genome, the mergeCNS script was used. CNS entries were compared based on genomic coordinates, and overlapping CNSs were collapsed into a single entry, prioritizing longer sequences, higher conservation scores, and broader conservation across species. A final filtering step was used to remove low-quality and spurious CNS associations.

Following the merge of the CNSs, sequences that result from unannotated coding sequences were filtered. CNS sequences were used for a tblastn against a combined protein dataset of all annotated proteins in the 314 genomes. CNSs that had more than 5 hits at Evalue lower than 0.0001, or had longer than 200 aligned basepair were considered as unannotated open reading frames and were removed from the dataset using the scripts blast_filter.sh.

To obtain a single CNS map for multiple reference species, the CNS dataset files were concatenated and then merged using the mergeCNS script with the --merge-deep parameter. This algorithm searches for overlapping CNSs from different reference species. If a sufficient number of CNSs overlap, they are merged to form a single CNS. The combined sequences are then realigned using mafft and then split or merged based on the depth of conservation. The CNS-gene association is filtered based on phylogenetic relationships. Finally, ancestral sequences are reconstructed, as detailed above, and CNS age is determined based on the deepest node in the all-species tree common to all species with the CNS.

Assay for transposase-accessible chromatin using sequencing (ATAC-Seq) libraries

ATAC-seq was performed as previously described (9). Briefly, six biological replicates of 1g of freshly collected terminal leaflet of young M82 leaves, were ground in ice-cold isolation buffer (300 mM sucrose, 20 mM Tris pH 8.0, 5 mM $MgCl_2$, 5 mM KCl, 5 mM beta-mercaptoethanol, 35% glycerol, 0.2% Triton X-100). After grinding, the suspension was filtered through a series of cell strainers (the smallest being 70 mm). Samples were further disrupted using a Dounce homogenizer, and then washed four times with ice-cold isolation buffer. The enriched nuclei samples were resuspended in Resuspension Buffer (50 mM Tris pH 8.0, 5 mM $MgCl_2$, 5 mM beta-mercaptoethanol, 20% glycerol). In order to continue with 100,000 nuclei, a small portion of the nuclei-enriched sample was stained with 4,6-Diamidino-2-Phenylindole (DAPI), and counted under a microscope using a hematocytometer. For the transposase reaction

100,000 nuclei were resuspended in 22.5 ml freezing buffer (5 mM $MgCl_2$, 0.1 mM EDTA, 50 mM TRIS pH 8.0, 40% glycerol), and were mixed with 2X DMF Buffer (66 mM Tris-acetate (pH 7.8), 132 K-Acetate, 20 mM Mg-Acetate, 32% DMF) and 2.5ul Transposase enzyme from Illumina Nextera Kit (Catalog No. FC-121-1031), and incubated at 37°C for 30 min. Tagmented DNA was isolated using NEB Monarch PCR Purification kit (Catalog No. NEB #T1030). Next-generation sequencing (NGS) libraries were amplified in two steps [106] using KAPA HiFi HotStart ReadyMix PCR kit (Catalog No. KK2601). NGS libraries were PE50 sequenced on an Illumina MiSeq SY-410-1003 instrument.

ATAC-seq reads were trimmed with trimmomatic [107] and aligned to Heinz SL4.0 reference genome [108] with BWA mem [109]. Reads that mapped to either chloroplast or mitochondria were filtered out together with PCR duplicate reads (picard MarkDuplicates) and multiple-position aligned reads. Finally, only high-quality properly paired reads were retained for the analysis (samtools view -b -h -f 3 -F 4 -F 8 -F 256 -F 1024 -F 2048 -q 30). ATAC-seq peak calling was done by Genrich V0.6 (https://github.com/jsh58/Genrich) using all six libraries with the following parameters: -j -r -v -q 0.5 -g 30.

Definition of background regions

To calculate CNS enrichment for functional genomic marks, we defined an appropriate background space as the genomic regions that Conservatory searches for CNSs. This is a genome-specific length upstream and downstream of all protein-coding genes, excluding annotated transposons and genes within this space. For each gene, we identified the regions spanning the start and stop codons of the longest splice isoform, then used bedtools slop to expand these regions upstream of start codons by the 97.5th percentile of intergenic distances in the genome at hand, up to a maximum distance 100 kb, and downstream of the stop codon by 20% of the upstream distance, rounded to the nearest 1 kb. Genome-specific distances are provided in table S1. Next, from each expanded window, we used bedtools subtract to remove regions corresponding to protein-coding genes, transposable elements and assembly gaps. The resulting gene-specific intervals were merged to delineate each genome's background space.

Functional genomic marks

Coordinate files in GFF3, BED, bigwig or narrowPeak format were retrieved from original publications or public databases and lifted over to different versions of genome assemblies as needed using CrossMap [110], as specified in table S2.

Maize ChIP-seq peaks [111] in B73 version 4 coordinates were retrieved from NCBI Gene Expression Omnibus (GEO) and converted to B73 version 5 coordinates, using the B73 v4 to v5 chain file available from Ensembl release 61. Maize DAP-seq in v5 coordinates [112, 113] was obtained from GEO. Other maize epigenomic marks (histone modifications, ATAC-seq, bisulfite-seq) in B73v5 coordinates were obtained from MaizeGDB. Maize DNA methylation in bigwig-formatted coverage and methylation tracks were converted to bedgraph with bigWigToBedGraph [114]. Transcription factor binding sites were predicted for the maize genome with PWMscan [115] for all nonredundant position weight matrices in the JASPAR core 2024 set, using a cutoff of 1e-5. Predicted binding sites were subsequently filtered to retain only those sites with the highest observed score for each PWM.

Arabidopsis ChIP-seq *(116–150)* and ATAC-seq *(151, 152)* datasets were downloaded from the Plant Chromatin State Database *(153)* as bigwig files and converted to bedgraph format for peak calling with MACS3 bdgpeakcall *(154)*. Arabidopsis DNA methylation data *(155)* was downloaded from NCBI GEO in wig format. Histone modification data were downloaded from PCSD *(153)* and broad peaks identified with MACS3.

Published tomato meristem ATAC-seq *(9)* were obtained from NCBI GEO.

For each functional genomic mark, we counted how many unique CNS base pairs overlapped with that feature and with the background regions described above. Then, enrichment values were calculated as:

$$\log2((\text{mark} \cap \text{CNS} / \text{CNS}) / (\text{mark} \cap \text{background} / \text{background}))$$

Gene ontology enrichment

Gene ontology enrichment was performed using the clusterProfiler R package *(156)*. For enrichments in each species, the corresponding annotation was retrieved from AnnotationHub *(157)*. Enrichments were carried out using genes with at least one associated CNS as the background genes, compared to genes that were associated with angiosperm or tracheophyte CNSs.

Plant phenotyping

We defined vegetative trait phenotypes as shoot apical meristem (SAM) termination, cotyledon defects (multiple cotyledons, lobed cotyledons, and fused cotyledons), leaf fusion, and phyllotaxis defects. Vegetative trait phenotyping was done on young seedlings at various stages of vegetative growth. In order to determine if a mutant allele was embryonic lethal, an F2 population of at least 125 seedlings was screened and genotyped.

CRISPR-Cas9 mutagenesis, plant transformation, and selection of mutant alleles

CRISPR-Cas9 mutagenesis and generation of transgenic tomato plants were performed as previously described *(158)*. Briefly, guide RNAs (gRNAs) were designed using the Geneious Prime software (https://www.geneious.com) (table S6). The Golden Gate cloning system was used to assemble the binary vector containing the Cas9 and the specific gRNAs *(159–161)*. Final binary vectors were then transformed into the tomato cultivar M82 or groundcherry by Agrobacterium tumefaciens-mediated transformation through tissue culture *(158)*. First-generation transgenic plants (T_0) were genotyped with specific primers surrounding the target sites. To generate new alleles and purify them from potential spontaneous mutations or CRISPR-Cas9 off-target effects following plant transformation, all CRISPR-Cas9 T_0 transgenic lines were backcrossed to parental wild type M82 cultivar plants. These F1 populations were then screened for plants lacking the Cas9 transgene, and PCR products of the targeted regions were sequenced to confirm inheritance of alleles. Selected F1 plants were self-fertilized to generate F2 populations, and these segregating populations were used to validate the phenotypic effects of each allele by co-segregation. F2 or F3 homozygous mutant plants were then used for quantitative phenotypic analyses. Screening for the embryonic lethal phenotype and validation of expected segregation ratios was performed on F2 families.

Gene tree inference

To reconstruct the evolutionary history of CNSs associated with *TB1* or *WUS*, orthologs, the predicted peptide sequences of each genome were clustered into orthogroups with OrthoFinder *(70)*. Genes in the same orthogroup as the gene of interest (either *TB1* or *WUS*) that shared at least one CNS with the focal gene were retained for further analysis. For each filtered orthogroup, amino acid translations were aligned using MAFFT (version 7.526) *(67)* and trimmed with clipkit *(68)*. For the *WUS* tree, only homologs from the Asteraceae and outgroup taxa were retained. Maximum likelihood trees were inferred with IQtree2 *(72)* with 1000 ultrafast bootstrap replicates *(162)*. CNS presence-absence tip states were visualized with the R packages ggtree *(163)* and treeio *(164)*, and CNS microsynteny was visualized with the R package gggenomes *(165)*. CNSs were extracted using genome-specific GFF3 annotations with samtools (version 1.19) *(166)*, aligned with MAFFT (v7.508) and visualized in Jalview (v2.11.5.0) *(167)*.

Chromatin interactions

Maize Hi-C and Hi-ChIP loop edges in B73 version 4 coordinates were obtained from previous studies *(55, 57)* and converted to B73v5 coordinates using CrossMap v0.7.3 *(110)*. *Solanum lycopersicum* cv. M82 *anantha (an)* mutants were grown in pots under greenhouse conditions *(168)*. The cauliflower-like tissue from young *an* inflorescences was harvested 10 weeks after sowing, with tissue collected from two different plants to obtain two biological replicates (3.3 g and 3.1 g, respectively). Hi-C data was generated using the Arima Hi-C 2.0 Kit (PN A410110) according to the manufacturer's user guide for plant tissues (PN A160163 v00). Nuclei extraction was done using the CelLytic Plant Nuclei Isolation/Extraction Kit according to the manufacturer's recommendations for cell lysis and semi-pure preparation of nuclei, with the addition of a Dounce homogenizer to increase nuclei yield (Sigma, CELLYTPN1). DNA fragmentation was performed with the Bioruptor Pico (Diagenode) using 4 cycles of 15 s sonication followed by 90 s rest. Library preparation was performed using the Swift Biosciences Accel-NGS 2S Plus DNA Library Kit, Swift Biosciences Indexing Kit, KAPA Library Quantification Kit for Illumina Platforms, and KAPA Library Amplification Kit according to the Arima Hi-C 2.0 library preparation guide (PN A160164 v00). Each library was sequenced in paired-end mode (2 × 250 bp) on an Illumina NovaSeq 6000 SP flowcell, yielding 463,915,437 reads for replicate 1 and 461,940,842 reads for replicate 2. Raw sequencing reads were processed using HiC-Pro v3.1.0, mapping reads to the M82 genome (SollycM82_v1.0) with a MAPQ threshold >30 *(169, 170)*. Valid read pairs were identified after filtering out invalid pairs and PCR duplicates, and both raw and normalized contact maps were generated at multiple bin sizes (5 kb, 10 kb, 20 kb, 40 kb, 150 kb, 500 kb, and 1 Mb). Following filtering, 167,352,522 and 175,494,920 valid reads were retained for replicates 1 and 2, respectively. Contact matrices were converted to .mcool format with Cooler, and the reproducibility between replicates was evaluated using HiCRep *(171, 172)*. HiC-Pro v3.1.0 was then used to merge valid read pairs from both replicates and generate contact matrices. FitHiC2 was used to identify significant chromatin looping interactions at 5 kb resolution on the merged data (q-value 0.01, maximum loop distance of 1 Mb, minimum loop distance of 10 kb), identifying a total of 9,573 loops *(173)*. Loop edge sequences were then aligned to the SL4.0 genome using minimap2 with asm10 parameters *(174)* and filtered to retain cases where both edges mapped to the same SL4.0 chromosome exactly once, with primary alignment tags (tp:A:P) for both edges.

To test for CNS enrichment in loops that connected intergenic regions to genes and proximal promoters, we intersected loop edges with genes including 500 bp of upstream sequence and with CNSs. We then identified loops for which one or both edges did not intersect a gene and its 500 bp proximal promoter. For loops that connected intergenic regions, one edge was chosen at random. We then constructed a 2x2 contingency table, stratifying loops by their connectivity (intergenic-genic or intergenic-intergenic) and whether the intergenic edge contained a CNS, and tested for statistical significance with a chi-square test. To evaluate the concordance of CNS-gene associations with chromatin loops in Arabidopsis, tomato and maize, we filtered to retain intergenic-genic loops that overlapped with CNSs at intergenic edges and that spanned less than the maximum upstream distance searched by Conservatory in each genome (100 kb in maize and tomato and 11 kb in Arabidopsis). For each of these loops, we then examined the gene associations of each CNS within the intergenic loop edge. If a CNS within a loop was associated with a gene, and the genic edge of that loop overlapped with the same gene, or with 500 bp of upstream sequence, it was counted as supported. All intersection operations were carried out in R using the valr package *(175)*.

Comparison of CNS space between paralog pairs

Paralog pairs were identified in each genome using DupGenFinder *(176)* with *Amborella trichopoda* as the outgroup. For each paralog pair, the normalized number of CNSs specific to one pair was compared to the total CNS associated with both pairs.

Fig. S1. Flow chart of the Conservatory algorithm. (**A**) Preprocessing of genome sequence and annotation data (**B**) Construction of the homologs database (**C**) Identification of CNS per gene. (**D**) Merging of CNS for all genes in a given reference genome. (**E**) Merging CNS databases from different reference genomes.

Fig. S2. CNSs identified within experimentally validated CREs. (A-F) Microsynteny maps and multiple sequence alignments for CNSs located within experimentally validated CREs of *FILAMENTOUS FLOWER (FIL)* (*35*) **(A)**, *LATERAL SUPPRESSOR (LAS)* (*34*) (B), *BLADE-ON-PETIOLE (BOP)* (*13*) (C), *PROTODERMAL FACTOR 1 (PDF1)* (*37*) (D), *APETALA1 (AP1)* (*36*) (E), and *SHOOT MERISTEMLESS (STM)* (*33, 177*) (F).

Fig. S3. Gene duplication results in paralog-specific divergence of ancient CNSs. (A) Gene tree of WUS showing the duplication at the base of the Asteraceae that gave rise to *WUSa* and *WUS*b. **(B)** Microsynteny map of *WUSa* and *WUSb* in the Asteraceae**.** Note that *WUSa* maintains greater conservation of the meristem CNS module, while *WUSa* exhibits increased divergence with Asteraceae-specific CNSs (shown in orange). **(C)** Multiple sequence alignment of the ancient S115 from representative angiosperms and the collinear Asteraceae-specific CNS in *WUSa*. **(D)** Microsynteny map of *WUS* homologs in the grasses showing conservation amongst paralogs in the PACMAD clade.

Fig. S4. Conserved association of a CNS with multiple genes. (A-B) A phylogeny (A) and microsynteny map (B) of the association of the seed plant CNS S115 with WUS and other genes across evolution. Note the particularly strong synteny conservation with TRA1A, despite many genomic rearrangements.

Fig. S5. Association of CNS with non-adjacent genes. (**A**) Proportion of genes located between a CNS and an associated gene that have CNS of their own. (**B**) Proportion of genes located between CNS and an associated gene that belong to either the same or different orthogroup. The vast majority belong to a different orthogroup, indicating that the gene skipping is not due to tandem duplications.

Table S1. A list of all species and genomes used in the Conservatory analyses.

Table S2. Functional genomic datasets reanalyzed for CNS overlap enrichment analyses.

Table S3. Number of CNSs of different conservation levels in all genomes.

Table S4. Gene list indicator shown in figures and supplementary figures.

Table S5. Fraction of CNSs classified as closest to associated gene, not associated with closest gene, or as skip associations per species.

Table S6. Arabidopsis genes associated with CNS lost in the Poaceae.

Table S7. gRNA and primer sequences used in this study.