Structural Basis for Sarbecovirus Rc-o319 Spike Adaptation to *Rhinolophus cornutus* Bat ACE2 and Constraints on Switching to Human ACE2

Jingjing Wang[1,2,†], Zexuan Li[1,3,†], Yong Ma[1,2,†], Zimu Li[1,2,4,5], Hang Yuan[1,2,3], Chuanying Niu[1,5], Yifeng Teng[1,3], Banghui Liu[1,2], Mei Li[4], Min Zhou[4], Wenxiu Liu[4], Huimin Feng[1], Jing Chen[4], Jun He[1,2], Xinwen Chen[4,*], Xiaoli Xiong[1,2,*]

1. State Key Laboratory of Respiratory Disease, Guangdong Provincial Key Laboratory of Stem Cell and Regenerative Medicine, Guangdong-Hong Kong Joint Laboratory for Stem Cell and Regenerative Medicine, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China.

2. Guangzhou Medical University-Guangzhou Institutes of Biomedicine and Health Joint School of Life Sciences, Guangzhou Medical University, Guangzhou 511436, China.

3. University of Chinese Academy of Sciences, Beijing 100864, China.

4. Guangzhou National Laboratory, Guangzhou 510005, China.

5. University of Science and Technology of China, Hefei 230026, Anhui, China

†These authors contributed equally to this work.

*Corresponding author. Email: chen_xinwen@gzlab.ac.cn (X.C.); xiong_xiaoli@gibh.ac.cn (X.X.)

Abstract

Bat sarbecoviruses often exhibit species-dependent ACE2 specificity. Understanding the determinants of receptor specificity enables better assessment of the cross-species transmission potential of sarbecoviruses. Here, we characterize the S-protein of Rc-o319, a sarbecovirus identified in Japanese *Rhinolophus cornutus* bats. Featuring an unusual 9-amino-acid deletion within its receptor binding motif (RBM), Rc-o319 S-protein utilizes its cognate *R. cornutus* ACE2 (bACE2$_{R.cor}$) but not human ACE2 (hACE2), demonstrating highly restricted receptor specificity. Cryo-EM structures reveal two locked prefusion conformations of the Rc-o319 S-trimer and define a novel type-1b receptor-binding domain (RBD), featuring a distinct beta-loop (BL) within the RBM due to the RBM-deletion. The Rc-o319-RBD:bACE2$_{R.cor}$ complex structure reveals unique interactions mediated by the specialized BL and RBM-loop of Rc-o319-RBD and by a bACE2$_{R.cor}$ glycan. Structure-guided mutagenesis demonstrates that changes in BL and RBM-loop within the Rc-o319 S-RBD must occur simultaneously to allow medium-to-high-affinity hACE2 binding. Comparative assays further show that the bACE2$_{R.cor}$ receptor supports only a subset of sarbecoviruses, highlighting its restricted sarbecovirus compatibility. Our findings establish the Rc-o319 S-protein as a structurally and functionally specialized adaptation to *R. cornutus* ACE2 and identify the structural constraints limiting its cross-species transmission potential.

Author Summary

SARS-related viruses are widely found in horseshoe bats. Some can bind the human ACE2 receptor with variable affinities, whereas others bind only bat ACE2 receptors. Understanding the basis for this difference is critical for assessing spillover risk. We studied the spike protein of a bat sarbecovirus, Rc-o319, isolated from Japanese *Rhinolophus cornutus* horseshoe bats. Although Rc-o319 is genetically related to SARS-CoV-2, it is unable to bind the human ACE2 receptor. Structural analyses and functional experiments revealed that the Rc-o319 spike protein possesses a distinct receptor-binding motif (RBM). This RBM features a specialized "beta-loop" that replaces the "large-loop" found in human-infecting sarbecoviruses. The beta-loop enables high-affinity binding to its cognate *Rhinolophus cornutus* bat ACE2 receptor while simultaneously preventing optimal interaction with human ACE2. We further found that acquisition of high affinity for the human ACE2 receptor would require coordinated changes across multiple regions of the RBM, including changing the beta-loop into a large-loop structure. Together, our findings demonstrate that the Rc-o319 spike protein is highly adapted to its cognate bat ACE2 receptor and faces substantial structural constraints that limit its ability to switch to binding human ACE2.

Introduction

The SARS-CoV-1 (also known as SARS-CoV) outbreak (1-4) and the SARS-CoV-2 pandemic have profoundly impacted global public health. Surveillance efforts have since detected multiple SARS-related coronaviruses (SARSr-CoVs) primarily in horseshoe bats (genus *Rhinolophus*) worldwide (5-9), suggesting they are the likely natural reservoir for SARSr-CoVs (10-12). Along with SARS-CoV-1 and SARS-CoV-2, SARSr-CoVs have been classified into the *Sarbecovirus* subgenus in the *Betacoronavirus* family (https://ictv.global/taxonomy). Sarbecoviruses share considerable similarity in their spike (S) proteins with amino-acid (AA) sequence identity > 63% (13-18). Like other coronaviruses (CoVs), sarbecovirus S-protein assembles into a large, glycosylated homo-trimer on the virion surface. Each S-protein protomer can be divided into two parts: the S1 part, responsible for cell attachment, and the S2 part, facilitating virus-cell membrane fusion (19-21). For a sarbecovirus S-protein, its S1 parts can be further divided as the N-terminal domain (NTD), receptor binding domain (RBD), Domain C (also called subdomain-1, SD-1) and Domain D (also called subdomain-2, SD-2) (17, 19, 22-25). The S-RBD can be further divided into a core domain, onto which a receptor binding motif (RBM) region is attached (17, 26). The RBM primarily contacts the receptor, forming the interaction interface with the receptor (24, 26-28).

For both SARS-CoV-1 (26, 29) and SARS-CoV-2 (30, 31) their S-RBDs bind with the human ACE2 (hACE) receptor with high affinities in the nano-molar range, suggesting that high-affinity hACE2 binding is likely a key prerequisite for spill-over into the human population. Of note, not all bat sarbecoviruses can bind hACE2 with high-affinity (5, 7, 12-16, 32, 33). Phylogenetic classification of SARSr-CoVs based on S-RBD sequences has defined multiple clades (34). Clade 1 sarbecoviruses utilize ACE2 and can be further subdivided into clade 1a (SARS-CoV-1 lineage) and clade 1b (SARS-CoV-2 lineage), both of which possess the most complete RBMs, lacking deletions of more than one amino acid. In contrast, no cellular receptor has been identified for clade 2 sarbecoviruses, which harbor deletions at two distinct regions within the RBM. Finally, clade 3 viruses contain a single loop deletion within the RBM and exhibit variable capacities for ACE2

utilization. Based on these distinct RBM deletion patterns and their associated

structural features among clades 1-3 viruses, we previously classified their RBDs

as type-1, type-3, and type-2, respectively (17).

Among the three types of sarbecovirus RBDs with distinct RBM structural features,

four featured RBM regions have been defined, namely the large-loop (LL), lamella

(LM), small-loop (SL), and finally the anchor-loop (AL) (17). Type-1 RBD S-

proteins, including those from clade 1a and 1b sarbecoviruses such as SARS-CoV-

1, SARS-CoV-2 and WIV1, possess intact LL and SL regions. Type-1 RBD S-

proteins often bind hACE2, albeit with varying affinities: SARS-CoV-1 and

SARS-CoV-2 bind hACE2 with high affinities, whereas WIV1 binds hACE2 with

a moderate affinity (35). We found high-affinity hACE2 binding requires

synergistic favourable hACE2 interactions across the four featured RBM regions.

Type-2 RBD S-proteins, including those from clade 3 sarbecoviruses such as

BM48-31 and BtKY72, have an intact LL but a deleted SL. They generally lack

hACE binding but can bind selected bat ACE2 (bACE2) orthologs with

intermediate affinities. Type-3 RBD S-proteins, including those from clade 2

sarbecoviruses such as HeB2013 and RmYN02, have both their LL and SL regions

deleted. To date, type-3 RBD S-proteins have not been reported to bind either

hACE2 or bACE2.

Ongoing surveillance in Japan has recently discovered several new SARSr-CoVs

in the little Japanese horseshoe bat - *Rhinolophus cornutus* (*R.cor*) detected in

2020 using samples collected in 2013 (7). The presentative virus, Rc-o319, is

distantly related to SARS-CoV-2, sharing 81.5% whole-genome identity. The

nucleotide and amino acid sequences of Rc-o319 S-protein are more homologous

to those of viruses belonging to the SARS-CoV-2 lineage (clade 1b) than to those

of the SARS-CoV-1 lineage (clade 1a) within the *Sarbcecovirus* subgenus (**Fig.**

S2). Interestingly, although its S-protein has 76.7% AA identity to SARS-CoV-2

S-protein, its RBM contains a unique nine-amino-acid deletion relative to SARS-

CoV-2 that removes residues identified critical for ACE2 binding in other SARSr-

CoVs and likely alters its RBM structure (7). Deletions in the RBMs of SARSr-

CoVs have been shown to greatly affect ACE2 binding (28, 36). In this study, we

surveyed binding of Rc-o319 S-protein to various ACE2 orthologs to understand

its specificity. Further, we determined the structures of the Rc-o319 S-trimers and

of the Rc-o319-RBD bound to *R. cornutus* ACE2 (bACE2$_{R.cor}$), thereby

establishing the Rc-o319 RBD as a novel structurally-defined type-1b RBD. By

analysing the molecular determinants of receptor specificity revealed by our

structures, we identify the changes necessary to switch the specificity of the Rc-

o319 S-protein to hACE2. These findings provide insights into Rc-o319's

adaptations to various ACE2 orthologs and provide an assessment of its cross-

species transmission potential.

Results

Rc-o319 S utilizes a narrow range of *Rhinolophus* bat ACE2 orthologs as receptors

We used a S:ACE2 interaction mediated cell-cell fusion assay to identify ACE2 orthologs that can interact with Rc-o319 S-protein (37). Effector cells expressing Rc-o319 S-proteins were tested against receptor cells expressing a panel of ACE2 orthologs for fusion activity (**Fig 1A**). In this assay, several *Rhinolophus affinis* (*Ra*) and *Rhinolophus sinicus* (*Rs*) ACE2 molecules, including *Ra*5538, *Ra*9479, *Rs*1446, *Rs*3359, and *Rs*3366, were found to mediate varying levels of cell-cell fusion, with *Ra*9479 showing the highest activity and *Rs*3366 the lowest. However, all were substantially weaker than the cognate $bACE2_{R.cor}$ receptor. Rc-o319 S-protein is unable to utilize pangolin and human ACE2 (hACE2) orthologs for fusion. Receptor binding assays using biolayer interferometry (BLI) show that Rc-o319-RBD-Fc and Rc-o319-S-trimer bind $bACE2_{R.cor}$ with moderate-to-high affinities (**Fig 1B** left panels, 76 nM for RBD-Fc and 27 nM for S-trimer). Although Rc-o319 appears to somewhat use $bACE2_{Ra9479}$, showing weak fusion activity in the cell-cell fusion assay. BLI assays were unable to detect $bACE2_{Ra9479}$ binding by the Rc-o319 S-trimer and dimeric Rc-o319-RBD-Fc (**Fig 1B**, middle panels). Rc-o319-S-trimer binds $bACE2_{R.cor}$ with relatively weak responses and an estimated affinity of 27 nM, tighter than that observed for the dimeric Rc-o319-RBD-Fc, indicating avidity-amplified receptor binding (**Fig 1B,** left panels). Consistent with the cell-cell fusion assay results, neither Rc-o319-RBD-Fc nor Rc-o319-S-trimer was able to bind hACE2 (**Fig 1B**, right panels). These results demonstrate that Rc-o319-S-protein has a restricted ACE2 specificity, with a clear preference towards its cognate $bACE2_{R.cor}$ receptor (**Fig 1A** and **B**).

Structures of the Rc-o319 S-protein identify a new type of RBM

To understand the unusual restricted receptor specificity of Rc-o319 S-protein, purified Rc-o319 S-protein ectodomain was imaged by cryo-electron microscopy (Cryo-EM) (**S1 Fig**), revealing that Rc-o319 S-trimers adopt two distinct conformations (**Fig 1D-I, S1 Fig, S3 Fig, and S4A, F and G Fig**). In both conformations, the 3 RBDs in each S-trimer are all in "down" positions (**Fig 1D**

and **G**). Detailed structural analysis reveals tight trimer packing for both
conformations. Based on their structural features, the two conformations differ
primarily at domain D (**Fig 2A** and **B, S3** and **S4F** and **S4G Figs**) and have been
identified as "locked-1" and "locked-2" conformations previously identified for
both SARS-CoV-1 and SARS-CoV-2 S-trimers (22, 25, 38-40).

In the locked-1 Rc-o319 S-trimer, each RBD contains a density that can be
modelled as a fatty acid with 18-carbon atoms, previous mass-spectrometry
analysis identified it being either linoleic acid (41, 42) or oleic acid (43) (**Fig 1E**
and **S4C Fig**). Consistent with previous observations, the aliphatic tail of the fatty
acid is bound within the hydrophobic fatty-acid binding pocket of one RBD, while
its carboxylate group forms salt-bridges with $R382_{Rc-o319}$ of a neighbouring RBD
(**Fig 1E** and **S4C Fig**); these fatty acids mediate inter-protomer interactions likely
stabilizing the S-trimer in the observed locked-1 conformation. Each NTD also
contains a density corresponding to a biliverdin molecule. The binding of
biliverdin has been correlated with evasion of antibody immunity (44, 45) (**Fig 1D**
and **1G** and **S4D Fig**). Consistent with being in a locked conformation, the fusion
peptide proximal region (FPPR) is ordered (**Fig 1F**). Finally, consistent with the
locked-1 conformation observed for the S-proteins of SARS-CoV-1 (25) and
SARS-CoV-2 (40), residues $580\text{-}605_{Rc-o319}$ form a large disordered loop within
Domain D of the Rc-o319 S-trimer (**Fig 1D** and **F and S4F Fig**).

In the locked-2 conformation, no fatty acid density was observed within the RBD
fatty-acid binding pocket (**Fig 1H**). This is accompanied with a restructuring of the
fatty-acid binding pocket (**Fig 1H** and **S4C Fig**) as observed when the fatty-acid
binding pockets were vacated in other sarbecovirus S-trimers (41, 46). Each NTD
still contains a density modelled as a biliverdin (**Fig 1D** and **G**). The FPPR is
ordered (**Fig. 1I**). Differently, the disordered Domain D loop in the locked-1
conformation refolds and forms a large ordered helical structure, packing against
the core of the Domain D and the junction between Domains C and D with non-
interupted cryo-EM density (**Fig 1F and S4G Fig**) (25, 40).

A structural change between locked-1 and locked-2 conformations in the Domain
C-D junction region can be observed similar to previous observations on the S-
proteins of SARS-CoV-1 and SARS-CoV-2 (25, 40) (**S4A, 4F** and **4G Fig**). Near
the 3-fold axis, to the top of the S-trimer, $D468_{Rc-o319}$ forms an intra-protomer
hydrogen bond with $Y473_{Rc-o319}$ in the locked-1 conformation (**Fig 2A**). Whereas in
locked-2 conformation, $D468_{Rc-o319}$ mediates two inter-protomer hydrogen bonds
with $S469_{Rc-o319}$ of a neighbouring S-protomer (**Fig 2B**). Of Note, in both
conformations, $K458_{Rc-o319}$, equivalent to $Q493_{SARS2}$, previously identified as an
important determinant for ACE2 specificity (17, 47, 48), is found to form an inter-
protomer hydrogen bond to the glycan attaching to $N344_{Rc-o319}$ of a neighbouring
S-protomer (**Fig 2A** and **B, the second panels**). Inter-protomer hydrogen bonds
likely contribute to S-trimer stability.

In both Rc-o319 S-trimer conformations, their NTDs are well resolved, revealing
five loops that are markedly shorter than those in SARS-CoV-2 (**Fig 2C and D** and
S4E Fig). In addition, a four-amino-acid insertion (residues $189-192_{Rc-o319}$) is
identified in the Rc-o319 NTD, forming an "inserted loop" exposed to the exterior
of the trimer (**S4B Fig**). It's well-established that the SARS-CoV-2 S-protein NTD
is under strong immune pressure (49, 50). The truncated and elongated loops
observed in the Rc-o319 S-protein NTD may therefore confer a distinct antigenic
profile, although whether these features reflect immune-driven selection in bats
remains to be determined.

Finally, Rc-o319 S-trimer structures show that their RBDs form structures distinct
from previous characterized sarbecovirus RBDs (**Fig 2E-G** and **S5 Fig**). In
contrast to type-3 RBDs, where 13- to 14-amino-acid deletions relative to SARS-
CoV-2 eliminate the large-loop (LL) (**S5C** and **S5F-G Fig**), the Rc-o319 RBD
retains an extension in a similar region as the LL of Type-1 RBDs, despite a nine-
amino-acid deletion (**Fig 2E and 2G**). In the type-1 RBD, the LL is formed by two
half-loops stabilized by an intra-loop disulfide bond (**Fig 2F**). By contrast, the
equivalent extension in the Rc-o319 RBD comprises two small antiparallel β-
sheets connected by a short loop (**Fig 2E** and **Fig 3**). We designate this distinct
extension a beta-loop (BL) structure. Due to the nine-amino-acid deletion, the
stabilizing disulfide bond found in the LL is lost. The shorter BL occupies a

position roughly corresponding to the C-terminal half-loop of the LL in type-1
RBDs (**Figs 2E-G** and **3D)**, albeit with notable conformational and tilting angle
differences (**Figs 2G,** also see **3D**). This distinct beta-loop (BL) structure, by
comparison with the LL structure in type-1 RBDs (**Fig 2F**), suggests that the Rc-
o319-RBD likely binds ACE2 with distinct interactions.

Structure of the Rc-o319-RBD:bACE2$_{R.cor}$ complex

To further understand receptor binding by the Rc-o319 S-protein, we determined a
structure of Rc-o319-RBD in complex with its cognate bACE2$_{R.cor}$ receptor (**Fig 3**
and **S1B Fig**). This structure uncovers several previously unrecognized interactions
at the Rc-o319-RBD:bACE2$_{R.cor}$ interface, even though the molecular surfaces
employed for binding largely mirror those in other sarbecovirus RBD:ACE2
complexes (**Fig 3A and B**). These interactions are mediated by the specialized
RBD BL and RBM-loop and a glycan linked to Asn38 of bACE2$_{R.cor}$ (**Fig 3A, E**
and **H-I**) (interfacial residues in the RBD and ACE2 are denoted using single-letter
and three-letter amino acid codes, respectively). The buried surface area (BSA) of
the complex was calculated as 1005 Å^2, which includes a contribution of 252 Å^2
from the binding of the Asn38-linked glycan. In line with previous interface
analysis, the binding interface can be divided into four regions, namely the beta-
loop (BL), lamella (LM), anchor loop (AL) and small loop (SL) regions (**Fig 3F-I**).

The beta-loop (BL) of Rc-o319 RBD engages highly distinctive interactions (**Fig**
3A and **F**). Due to its distinct structure and protrusion from the RBM, the BL
contacts the bACE2$_{R.cor}$ near Lys27, forming a hydrogen bond with the side chain
of Lys27 (**Fig 3A** and **F**). In contrast, LL in type-1 RBDs contacts ACE2 near
residue 24, more towards the ACE2 N-terminus (**Fig 3J**). Distinctively, BL residue
Y447$_{Rc-o319}$ forms a charged hydrogen bond with Asp31 of bACE2$_{R.cor}$ (**Fig 3A** and
F). Of note, the positively charged residue K454$_{Rc-o319}$, unique to the Rc-o319
RBM, extends from the BL to form a salt-bridge with Glu75$_{Rc-o319}$ of bACE2$_{R.cor}$
(**Fig 3A and F**) and is positioned in close proximity to the negatively charged
bACE2$_{R.cor}$ residues Asp31 and Glu35 (**Fig 3F-G**). Changing this residue

(K454Y$_{Rc-o319}$) to the corresponding residue Y in SARS-CoV-2 substantially

reduced bACE2$_{R.cor}$ binding **(S9B Fig)**.

Distinct from SARS-CoV-2, the large Rc-o319 RBM-loop (**Figs 2E, 3C and 3F**

and **S7 Fig**) adjacent and N-terminal to the BL contributes to bACE2$_{R.cor}$ binding

and contacts the BL structure with specialized interactions (compare **Figs 3F** and

3J): W430$_{Rc-o319}$ in the RBM-loop forms a cation-π interaction with Lys27 of

bACE2$_{R.cor}$, and likely stabilizes the BL via a π-π interaction with the BL residue

Y447$_{Rc-o319}$ (**Fig 3F** and **S7B Fig** and **see below**).

In the lamella (LM) region, K458$_{Rc-o319}$ forms a salt-bridge with bACE2$_{R.cor}$ Glu35

(**Fig 3G**). Distinctively, in this region, the Asn38-glycan is part of the ACE2-RBD

interface, forming hydrogen bonds with N459$_{Rc-o319}$ (**Fig 3G**). Also in this region,

Lys353$_{Rc-o319}$ forms a hydrogen bond with the G461$_{Rc-o319}$ backbone (**Fig 3G**). In

the anchor-loop (AL) region, a hydrogen bond network is observed among

bACE2$_{R.cor}$ residues Tyr41, Asn330, Glu335, Arg357 and the RBD residue S465

(**Fig 3H**). Further, the glycan attached to Asn38 of bACE2$_{R.cor}$ forms an inter-

molecular hydrogen bond with bACE2$_{R.cor}$ residue Glu42 and also hydrogen bonds

with RBD residue Y463$_{Rc-o319}$ (**Fig 3H**). Finally, in the small-loop (SL) region,

F423$_{Rc-o319}$ forms a sugar-π contact with the glycan attached to Asn38 (**Fig 3I**).

These interactions, by comparison with those used by SARS-CoV-2 to engage

ACE2 (**Figs 3J-M**), show that Rc-o319 RBD engages its cognate bACE2$_{R.cor}$

receptor with unique characteristics.

Molecular determinates underlying Rc-o319 specificity for bACE2$_{R.cor}$

To understand the specificity of the Rc-o319 S-protein towards the bACE2$_{R.cor}$

receptor. We employed a mutagenesis approach to systematically change RBM

regions to match those of SARS-CoV-2 (BL change = swapping of 443-456$_{Rc-o319}$

"AHYDYQVGTQFKSS" to 469-491$_{SARS2}$ "STEIYQAGSTPCNGVEGFNCYFP";

SL change = F423Y$_{Rc-o319}$; LM change = K458Q$_{Rc-o319}$; AL change = S465T$_{Rc-}$

$_{o319}$+A466N$_{Rc-o319}$+H470Y$_{Rc-o319}$; RBM-loop change = W430F$_{Rc-o319}$+S432K$_{Rc-o319}$).

These variants were used to assess hACE2 and bACE2$_{R.cor}$ binding by BLI assays
using RBD-Fc proteins, as well as receptor utilization by cell-cell fusion and
pseudovirus entry assays using full-length S proteins. A similar approach has been
previously used to elucidate how high-affinity hACE2 binding may be acquired by
S-proteins with type-2 RBDs, as we defined previously (17), from clade 3
sarbecoviruses.

We found that changing AL and SL regions individually had very little effect in
improving hACE2 binding (**Fig 4D** and **E, the second panels**). The SL change
retained very strong bACE2$_{R.cor}$ utilization (**Fig 4E, the third** and **fifth panels**),
whereas the AL change reduced bACE2$_{R.cor}$ utilization somewhat in cell-cell fusion
(**Fig 4D, the third** and **fifth panels**) and pseudovirus entry assays (**Fig 4M**).
Interestingly, the AL change very slightly increased bACE2$_{R.cor}$ binding when
assayed using both Rc-o319-RBD-Fc and the Rc-o319 S-trimer (**Fig 4D, the first**
panels and **S9E** and **S14F Figs**). The discrepancy between the increased BLI
affinity and the reduced cellular activity may reflect differences in assay conditions.
In the cellular context, phenotypes may also be influenced by S-protein-related
factors beyond receptor binding, such as S-protein stability, efficiency of surface
expression, incorporation into pseudovirus particles, and activation by cellular
proteases, which could contribute to the observed discrepancy (see the Limitations
section). Changing the LM region alone (the LM change) reduced bACE2$_{R.cor}$
binding and utilization (**Fig 4C, the first, third** and **fifth panels**) but still failed to
confer hACE2-mediated fusion activity and pseudovirus entry (**Fig 4C, the fourth**
and **fifth panels**).

Swapping of the Rc-o319 BL region to the LL sequence of SARS-CoV-2-RBD
(the BL change), appears to confer Rc-o319 with the ability to utilize hACE2 for
cell fusion activity at the expense of substantially reduced fusion activity mediated
by its cognate bACE2$_{R.cor}$ receptor (**Fig 4B**). Consistently, the BL change conferred
hACE2 mediated pseudovirus entry, while pseudovirus entry mediated by the
cognate bACE2$_{R.cor}$ reduced by ~52-fold (**Fig 4M**). Despite conferring detectable
cell-cell fusion activity and pseudovirus entry, hACE2 binding by the dimeric BL
variant of the Rc-o319 RBD-Fc protein remained below the detection limit of the
BLI assay, while bACE2$_{R.cor}$ binding was markedly reduced and fell below the
detection limit of the BLI assay (**Fig 4B, the first panels**). A subsequent BLI assay
using the BL variant of the Rc-o319 S-trimer detected weak hACE2 binding (**S10B**

Fig), likely made detectable by amplification through trivalent binding-mediated
avidity, underscoring a dependence of assay sensitivity on binding valency. As
cell-based assays involve highly multivalent interactions mediated by multiple S-
trimers and receptors, avidity-enhanced binding renders them substantially more
sensitive to weak interactions (also see the Limitations section).

Therefore, the above results show that replacing each RBM region singly with its
SARS-CoV-2 counterpart in the Rc-o319 RBD-Fc failed to confer high-affinity
binding to hACE2. This finding is consistent with our previous observations that
high-affinity hACE2 binding requires a sarbecovirus RBD to establish optimal
contacts across its all four RBM regions, including LL/BL, LM, AL, and SL(17).
A similar requirement for synergistic interactions across multiple RBM regions to
achieve high-affinity receptor binding has also been observed in merbecoviruses
(51).

However, unexpectedly, the BL+LM+AL+SL variant, designed to introduce
optimal interactions across all four RBM regions, exhibited enhanced hACE2
utilization in cell-cell fusion and pseudovirus entry assays, yet its hACE2 binding
remained too weak to be detected by BLI (**Fig 4F, the second** and **fourth panels**).
The BL+LM+AL+SL variant of Rc-o319-RBD-Fc protein also showed no
bACE2$_{R.cor}$ binding by BLI, and consequently exhibited very little bACE2$_{R.cor}$
mediated fusion activity (**Fig 4F, the first** and **third panels**). Therefore, in the
case of Rc-o319 S-protein, determinants outside of the BL, LM, AL, and SL
regions likely hinder high-affinity hACE binding.

Structural comparison shows that W430$_{Rc-o319}$ and S432$_{Rc-o319}$ in the large RBM-
loop (amino acid (AA) 429-442$_{Rc-o319}$, corresponding to AA 455-468$_{SARS2}$), located
N-terminal to the BL, are unique to the Rc-o319 RBD (**Fig 3C** and **F**, **S5G** and **S7**
Figs). In SARS-CoV-2-RBD, the corresponding residues F456$_{SARS2}$ and K458$_{SARS2}$,
respectively interact with the LL residues Y473$_{SARS2}$ and E471$_{SARS2}$ (**Fig 3J** and
S7A and **S7D Fig**). These interactions likely stabilize the SARS-CoV-2 LL
structure. We introduced these residues into the Rc-o319 BL+LM+AL+SL variant

as the W430F$_{Rc\text{-}o319}$+S432K$_{Rc\text{-}o319}$ change (collectively named as the "RBM-loop"

change). BLI binding results show that the dimeric BL+SL+LM+AL+RBM-loop

variant of Rc-o319-RBD-Fc is able to bind hACE2 with a much increased affinity

of 31.7 nM (**Fig 4G**, **the second panel**). We also attempted the swap of Rc-o319

RBM for the SARS-CoV-2 RBM (Rc-o319-RBD-RBM$_{SARS2}$), as expected, the Rc-

o319-RBD-Fc variant with the swapped RBM also showed high-affinity binding to

hACE2 with only slightly higher affinity (12.3 nM) than the

BL+LM+AL+SL+RBM-loop variant (**compare Fig 4G** and **H**, **the second panels**).

Consistently, both BL+LM+AL+SL+RBM-loop and RBM swapped Rc-o319-

RBD-RBM$_{SARS2}$ show effective hACE2 mediated cell-fusion activities (**Fig 4G** and

H, **the fourth** and **fifth panels**). Therefore, our mutagenesis results identify

residues in the RBM-loop of Rc-o319, N-terminal to the BL, that are incompatible

with the LL structure of type-1 RBDs, thereby preventing the LL from mediating

optimal ACE2 interactions (**S6C, S7C** and **S7D Figs**).

To further clarify the role of the RBM-loop in hACE2 recognition, we introduced

the RBM-loop change alone into the Rc-o319 S-protein. This change did not

confer hACE2-dependent cell-cell fusion (**S8B Fig**), consistent with structural

analyses indicating that the RBM-loop does not make direct contacts with ACE2

(**Fig 3**). No detectable hACE2 binding was observed for the Rc-o319 BL-variant

RBD-Fc (**Fig 4B**), and a similar lack of hACE2 binding was observed for the Rc-

o319 BL+LM-variant RBD-Fc in BLI assays (**Fig 4I**), despite the corresponding S-

proteins mediating clear hACE2-dependent cell-cell fusion and pseudovirus entry

(**Fig 4B, I** and **M**). When the RBM-loop change was introduced into the Rc-o319

BL- or BL+LM-variant RBD-Fc (i.e., BL+RBM-loop and BL+LM+RBM-loop

variants), weak but detectable hACE2 binding could be detected in BLI assays (**Fig**

4B, I-K). These results confirm that the RBM-loop plays a modulatory role,

essential for mediating high-affintiy ACE2 interaction.

Utilization of bACE2$_{R.cor}$ by different types of sarbecoviruses

Our structural and biochemical analyses show that the Rc-o319 S-protein has a

highly restrictive specificity towards its cognate bACE2$_{R.cor}$ receptor, we further

characterized the potential of bACE2$_{R.cor}$ to support diverse sarbecoviruses (**Fig 5**).
A panel of sarbecovirus S-proteins was surface-expressed on effector cells and
tested for cell-cell fusion activity against receptor cells expressing bACE2$_{R.cor}$ (**Fig
5A**). The panel included S-proteins with type-1 RBDs (17) from clade 1a and 1b
sarbeviruses, including SARS-CoV-1, LYRa11, SHC014, WIV1, RaTG13,
BANAL-20-52, BANAL-20-236, and the SARS-CoV-2 ancestral strain; S-proteins
with type-2 RBDs (17) from clade 3 sarbeviruses, including BtKY72, BM48-31,
RfGB02, and RhGB07; and S-proteins with type-3 RBDs (17) from clade 2
sarbeviruses, including GX2013, HeB2013, YN2013, and RmYN02. The results
show that bACE2$_{R.cor}$ supported receptor binding and mediated cell fusion
activities of several clade 1 sarbecovirus S-proteins with type-1 RBDs (**Fig 5B** and
C). However, potent fusion activity was only observed for several S-proteins of
viruses from the SARS-CoV-2 lineage (clade 1b). Although the SARS-CoV-1 S-
protein exhibits only intermediate bACE2$_{R.cor}$-mediated cell-cell fusion activity, it
is nevertheless the most potent among S-proteins from viruses within the SARS-
CoV-1 lineage (clade 1a) (**Fig 5A-C**). Among bACE2$_{R.cor}$ utilizing S-proteins, only
the BANAL-20-236 S-trimer was found to bind bACE2$_{R.cor}$ in BLI assays, whereas
S-trimers from BANAL-20-52, SARS-CoV-2, and SARS-CoV-1 exhibited no
detetctable binding to bACE2$_{R.cor}$ despite displaying robust cell-cell fusion activity
(**Fig 5C and S11B, C, E, and F Fig**). These observations indicate that, in addition
to receptor binding, complex cell-associated factors, potentially including S-
protein expression level, protein stability, and susceptibility to activation by
cellular proteases, can critically influence S-protein-mediated membrane fusion
(also see the Limitations section).

The Asn38-linked glycan in bACE2$_{R.cor}$ is involved in S-protein binding, a feature
observed for the first time between an ACE2 and a sarbecovirus S-protein. To
understand the effect of this glycan on S-protein function, we removed the glycan
by introducing a Thr40Ala change (designated as 'bACE2$_{R.cor}$-ΔGLC$_{38}$'). In cell-
cell fusion assays, the bACE2$_{R.cor}$-ΔGLC$_{38}$ variant receptor showed a tendency to
mediate enhanced fusion activities for all the sarbecovirus S-proteins capable of
utilizing the WT bACE2$_{R.cor}$ (**Fig 5B and C**). BLI assay results show that the
removal of the Asn38-glycan was able to enhance bACE2$_{R.cor}$ binding not only for
Rc-o319 but also for the BANAL-20-52 and BANAL-20-236 S-trimers (**S11A**

Fig). In particular, although unable to utilize WT bACE2$_{R.cor}$, BtKY72 S-protein,
with a type-2 RBM, of a clade 3 sarbevirus, was able to utilize bACE2$_{R.cor}$-ΔGLC$_{38}$,
albeit exhibiting only very weak cell fusion activity. Consistently, BLI binding
results show that the BtKY72 S-trimer was unable to bind bACE2$_{R.cor}$, but gained
bACE2$_{R.cor}$-ΔGLC$_{38}$ binding with an estimated affinity of 11.8 nM (**S11H Fig, the**
second panel).

To further assess the functional role of the Asn38-glycan, an N-linked
glycosylation site was introduced into a *Rhinolophus affinis* bat ACE2
(bACE2$_{Ra9479}$), by substituting Asp38 with Asn, generating bACE2$_{Ra9479}$-GLC$_{38}$. In
both cell-cell fusion and BLI assays, BtKY72 and SARS-CoV-1 S-proteins
exhibited substantially reduced fusion activity utilizing bACE2$_{Ra9479}$-GLC$_{38}$
compared with the WT bACE2$_{Ra9479}$ receptor (**S16B** and **F Fig; S11J** and **K Fig**).
In parallel, we swapped the RBMs of SARS-CoV-1 and BtKY72 S-proteins with
that of Rc-o319, generating chimeric SARS1$_{Rc-o319RBM}$ and BtKY72$_{Rc-o319RBM}$ S-
proteins. Although being expressed by the cell (**S16I** and **J Fig**), the RBM
swapping appeared to render the SARS1$_{Rc-o319RBM}$ S-protein non-functional, unable
to mediate cell-cell fusion mediated by either bACE2$_{Ra9479}$ or bACE2$_{R.cor}$ receptor
and their Asn38-glycan variants (**S16C** and **D Fig**). The BtKY72$_{Rc-o319RBM}$ S-
protein showed substantially reduced utilization of the bACE2$_{Ra9479}$ and
bACE2$_{Ra9479}$-GLC$_{38}$ receptors (**S16F** and **H Fig**). In contrast, its utilization of both
bACE2$_{R.cor}$ and bACE2$_{R.cor}$-ΔGLC$_{38}$ was only marginally enhanced compared with
the WT BtKY72 S protein, yet remained substantially weaker than that of the Rc-
o319 S protein (**S16E** and **G Fig** and **Fig 4A**). These results highlight compatibility
constraints between different RBMs and S-protein backbones, indicating that RBM
changes can adversely affect overall S-protein function.

Discussion

In this study, our receptor binding and structural analyses show that Rc-o319 S-
protein exhibits a clear preference for its cognate bACE2$_{R.cor}$ receptor with marked
reduced utilization of *R. sinicus* and *R. affinis* ACE2 receptors. Rc-o319 S-protein
showed no utilization of human and pangolin ACE2s. A closer relationship among

Asian *Rhinolophus* bats (52) probably allowed Rc-o319 S-protein to cross-bind *R.*
sinicus and *R. affinis* ACE2s. Previously, it has been shown that weak binding of
SARS-CoV-2 S-protein to mouse ACE2 (mACE2) was unable to facilitate
effective mouse infection, and increased mACE2 binding, thorugh the $N501Y_{SARS2}$
change, was key to effective mouse infection (53). Therefore, weak *R. sinicus* and
R. affinis ACE2 binding may not allow effective infection of *R. sinicus* and *R.*
affinis bats by the Rc-o319 virus.

Our structural analyses reveal that the Rc-o319 S-trimer adopts two distinct
conformations: a locked-1 conformation with fatty-acid-bound RBDs and a locked-
2 conformation in which those fatty-acid binding pockets are unoccupied. Previous
cryo-EM studies of bat sarbecovirus S-trimers have nearly always captured them in
locked-2 conformations with and without fatty acid bound (17). Locked-1
conformation has only been reported for the bat sarbecovirus WIV1 S-trimer (54).
Both locked-1 and locked-2 conformations have been observed for SARS-CoV-1
and SARS-CoV-2 S-trimers. The locked-1 conformation appears to be always
identified with fatty acid bound, whereas the locked-2 conformation has been
resolved with or without fatty acid bound. Interestingly, these two conformations
appear to be more difficult to capture for SARS-CoV-1 (25) and SARS-CoV-2 (22,
40, 55) S-trimers and their capture often requires deliberate stabilization of the S-
trimers. The predominance of locked conformations in bat sarbecovirus S-trimers,
versus them being rare conformations in human-infecting sarbecoviruses, has been
proposed to reflect likely differences in S-trimer stability (17). We have also
proposed that tightly packed S-trimers in locked conformations facilitate viral
egress (40). As the preferred conformation for bat sarbecovirus S-proteins, these
locked conformations have also been postulated to represent adaptations for fecal-
oral transmission and immune evasion in their natural hosts (54, 56).

Due to residue-31 of bat ACE2 molecules often being a negative charged Asp or
Glu, including in $bACE2_{R.cor}$, a positively charged LM region, such as $K458_{Rc-o319}$
(corresponding to $Q493_{SARS2}$), has been noted as an important determinant for
specificity towards bACE2 (47). Notably, type-1 RBDs often feature neutral LM
regions, whereas type-2 RBDs commonly contain positively charged LM regions

(17). Although BANAL-20-236 possesses a type-1 RBD, it unusually contains a
lysine in the LM region, which may confer an enhanced ability to bind a broader
range of bACE2 molecules (56). Monomeric BANAL-20-236-RBD was
previously measured to bind $bACE2_{Ra5538}$ with a moderate affinity of 104 nM (17);
it also binds $bACE2_{R.cor}$ with a reduced affinity of 981 nM (**S11L Fig**). SARS-
CoV-2-RBD is highly similar to BANAL-20-236, however, with a neutral Q in the
LM region, SARS-CoV-2-RBD-Fc binding to $bACE2_{R.cor}$ was too weak to be
measured by the BLI assay (**S9H Fig**). With a $Q493K_{SARS2}$ point mutation in LM,
SARS-CoV-2-RBD-Fc binds $bACE2_{R.cor}$ in BLI assays (compare **S9H** and **S9L**
Fig). In addition, SARS-CoV-1-RBD-Fc with an LM change ($N479K_{SARS1}$) gained
weak $bACE2_{R.cor}$ binding (**S8K Fig**). These results are further consistent with the
effect of the LM change ($K458Q_{Rc-o319}$) in the Rc-o319 S-protein, which
considerably reduced binding to $bACE2_{R.cor}$ (**Fig 4C**). Finally, although the SARS-
CoV-2, BANAL-20-236, and BANAL-20-52 S-proteins from clade1b
sarbecoviruses all exhibited robust fusion activities, only BANAL-20-236 S-trimer,
harbouring a lysine in the LM region, showed detectable binding to $bACE2_{R.cor}$ in
BLI assays (**S11E Fig**). These observations collectively suggest that a positive-
charged LM region contributes to $bACE2_{R.cor}$ receptor affinity and ultilization.

The Rc-o319-RBD:$bACE2_{R.cor}$ complex structure reveals that Rc-o319-RBD
employs a featured beta-loop (BL) structure to bind to the $bACE2_{R.cor}$ molecule
using a previous unobserved interaction mode that involves interactions with
$bACE2_{R.cor}$ residues Lys27, Asp31, and Glu75 (**Fig 3A** and **F**). BL change in Rc-
o319-RBD-Fc is even more detrimental than the LM change, rendering $bACE2_{R.cor}$
binding completely undetectable (compare **Fig 4B** and **C**). Notably, Lys27 is
uniquely present in $bACE2_{R.cor}$ and $bACE2_{R.ferr}$ among related ACE2 molecules.
However, whereas most ACE2 molecules contain a Glu75 residue, $bACE2_{R.ferr}$
instead contains a Lys75, which would abolish the $K454_{Rc-o319}$-Glu75 salt bridge
observed in our Rc-o319-RBD:$bACE2_{R.cor}$ complex structure (**Fig 3A** and **F**) and
likely introduce electrostatic repulsion with the BL residue $K454_{Rc-o319}$. The
specific interaction pattern between Rc-o319-RBD-BL and $bACE2_{R.cor}$
demonstrates an adaptation of the Rc-o319 RBM region to its cognate $bACE2_{R.cor}$
receptor.

While this manuscript was in preparation, Kosugi et al. reported a cryo-EM

structure of the Rc-o319-RBD:bACE2$_{R.cor}$ complex (57), corroborating the

interactions we observed and identifying Y463$_{Rc-o319}$ and S465$_{Rc-o319}$ in the AL

region as critical for bACE2$_{R.cor}$ binding. Importantly, beyond these AL region

interactions, we show that the BL and LM regions also govern bACE2$_{R.cor}$

engagement, and that replacing the Rc-o319 BL with the SARS-CoV-2 LL

sequence has a substantially greater impact on bACE2$_{R.cor}$ binding than multiple

LM mutations (**Fig 4B** vs **Fig 4C** and see **S9D Fig** for the K458A$_{Rc-o319}$

substitution), establishing the BL as a principal determinant of high-affinity

binding in Rc-o319 S-protein.

Consistently, swapping the BL with the LL of SARS-CoV-2 (the BL change) alone

enabled detectable hACE2-mediated cell-cell fusion activity, even though hACE2

binding by BL variants remained undetectable using RBD-Fc (**Fig 4B**) or only

very weak using S-trimer (**S10B Fig**) in BLI assays. Combined BL and RBM-loop

substitutions enabled Rc-o319 RBD-Fc binding to both hACE2 and bACE2$_{R.cor}$ to

be detected by BLI, albeit with very weak affinities (**Fig 4B** and **J**). These results

highlight the importance of the BL/LL region together with its corresponding

RBM-loop structure in ACE2 recognition, as well as the adaptation of the Rc-o319

BL/RBM-loop structures to bACE2$_{R.cor}$. Cell-cell fusion assays showed that the BL

change alone was sufficient to confer hACE2-mediated fusion, whereas the RBM-

loop change alone was insufficient (**S8B** and **S8C Fig**), indicating that changes

within the RBM BL region constitute the primary determinants enabling hACE2

utilization (**Fig 4** and **S8 Fig**).

Our data further show that the specialized Rc-o319 BL loop co-adapts with the Rc-

o319 RBM-loop to confer high-affinity binding to its cognate bACE2$_{R.cor}$ receptor.

This co-adaptation renders the RBM-loop incompatible with the SARS-CoV-2 LL

region (**S7C** and **F Fig**), preventing an introduced LL region from mediating high-

affinity hACE2 binding in BLI assays (compare **Fig 4F** and **G,** compare **Fig 4I** and

K). Conversely, changing LL to the Rc-o319 BL in SARS-CoV-2-RBD

completely abolishes binding to hACE2 and bACE2$_{R.cor}$ (**S12A** and **S12E Fig**),
likely due to incompatibility between the introduced Rc-o319 BL and the SARS-
CoV-2 RBM-loop. These observations further suggest that the Rc-o319 BL and
RBM-loop are coevolved structures adapted for bACE2$_{R.cor}$ binding, and their
unique structural features and interactions engaged support classifying the Rc-
o319-RBD as a novel type. We designated this RBD as type-1b because, like type-
1 RBDs, it lacks a deletion in the SL region, yet it features a specialized BL
structure in place of the LL characteristic of previously studied type-1 RBDs.
Notably, our data show the specialized BL and RBM-loop of the Rc-o319 RBM
severely limit its capacity for high-affinity hACE2 binding, with extensive
mutations and insertions in these regions likely required to achieve the binding
affinity necessary for cross-species transmission.

Finally, our results show that, in contrast to the highly efficient utilization of
bACE2$_{R.cor}$ by the Rc-o319 S-protein, bACE2$_{R.cor}$ supports only limited usage by
selected clade1 sarbecovirus S-proteins with type-1 RBDs. Notably, most of these
S-proteins are from viruses that belong to the SARS-CoV-2 lineage (clade 1b),
highlighting this lineage's extraordinary ability to utilize a broad range of ACE2
orthologs. Although multiple S-proteins were capable of mediating bACE2$_{R.cor}$-
dependent cell-cell fusion, most exhibited little or no detectable bACE2$_{R.cor}$
binding in BLI assays, suggesting that the Rc-o319 S-protein has evolved a
uniquely specialized specificity and affinity for bACE2$_{R.cor}$. Further, although
structural analysis shows that the Asn38-linked glycan of bACE2$_{R.cor}$ forms
specific interactions with Rc-o319 RBD residues F423$_{Rc-o319}$, Y426$_{Rc-o319}$, H444$_{Rc-o319}$,
$_{o319}$, and N459$_{Rc-o319}$, our data indicate that this glycan generally dampens
bACE2$_{R.cor}$ utilization by sarbecovirus S-proteins, including for Rc-o319, likely
through steric effects that reduce S-protein binding. Introduction of an Asn38-
linked glycan into bACE2$_{Ra9479}$ also dampened its usage by SARS-CoV-1 and
BtKY72 S-proteins. We noted that, in cell-cell fusion assays, the Asn38-linked
glycan had a greater impact on S-proteins mediating weak to intermediate fusion
activity than on those with strong fusion activity (**Fig 5**). We speculate that this
glycan may exert a more pronounced restrictive effect on viruses with limited entry
capacities. Interestingly, the Asn38-linked glycan is also positioned near the

binding interface of HKU5-like, ACE2-utilizing merbecovirus S-proteins and may
therefore modulate their receptor binding (58, 59).

In summary, the complementary interactions between the specialized RBM,
particularly the BL region, of the Rc-o319 S-protein and the corresponding
interacting residues in $bACE2_{R.cor}$ appears to reflect a highly specific virus-host co-
adaptation (**S17 Fig** and **Table S10**). To enable such adaptation, the RBM of Rc-
o319 has diverged substantially from that of the SARS-CoV-2 lineage, to which it
is most closely related, resulting in a highly restricted ACE2 specificity. Notably,
the RBM is also a major target of immune pressure, as demonstrated by the
ongoing evolution of SARS-CoV-2 (60-64). Even though certain SARS-CoV-2
lineage viruses can use $bACE2_{R.cor}$, we still do not know what evolutionary forces,
perhaps immune pressure, pushed Rc-o319 S toward specialization for its host
receptor.

Limitations of the Study:

A limitation of this study is that the biophysical BLI assays employed relatively
low-valency analytes (monomeric RBD, dimeric ACE2, or trimeric S-protein),
thereby limiting avidity-mediated binding amplification and sensitivity to weak
interactions. In contrast, cell-cell fusion and pseudovirus entry assays (**Figs 4, 5,**
and S16 Fig) involve highly multivalent interactions between multiple surface-
displayed S-trimers and ACE2 molecules, resulting in substantially enhanced
avidity and sensitivity. Accordingly, several interactions that were undetectable by
BLI nonetheless supported detectable fusion or entry in cell-based assays (**Figs. 4B,**
4F, 4G, 4I, 4K, 4L, and 5). A further limitation is that, in the cellular context,
phenotypes can be influenced by S-protein-related factors beyond receptor binding,
potentially including S-protein stability, efficiency of surface expression,
incorporation into pseudovirus particles, and activation by cellular proteases. This
is exemplified in Fig. 5, where the SARS-CoV-2 S-protein exhibited the strongest
fusion activity mediated by both WT $bACE2_{R.cor}$ and the glycan mutant
$bACE2_{R.cor}$-ΔGLC_{38}, despite no detectable binding to either receptor in BLI assays.
An additional example is shown in **S16 Fig**, where a SARS-CoV-1 S-protein

variant carrying the Rc-o319 RBM (SARS1$_{\text{Rc-o319RBM}}$) failed to mediate
bACE2$_{R.cor}$-dependent fusion despite detectable expression, suggesting
incompatibility between the swapped RBM and the S-protein backbone. While
cell-based assay readouts often scale with large differences in receptor-binding
affinity, they may not scale linearly when affinity differences are modest and the
above-discussed S-protein-related factors are involved. Several RBM-modified S-
proteins exhibited modestly increased ACE2 binding in BLI assays but reduced
fusion or entry activity in cell-based assays (**Fig 4A vs 4D and Fig 4G vs 4H**). We
speculate that RBM changes may affect S-protein stability, surface presentation, or
interactions with host factors, leading to nonlinear relationships between receptor-
binding affinity and cellular phenotype. Finally, future studies using animal models
and authentic viruses would be needed to further assess the relevance of receptor-
affinity changes in the context of infection.

Materials and Methods

Protein constructs

S gene of Rc-o319 (Taxonomy ID: 694009) was codon-optimized and synthesized by Sangon Biotech (Shanghai, China). The transmembrane domain and C-terminal end of Rc-o319 (AA 1176-1235) were removed before the C-terminus of the coding sequence was fused to a T4 trimerization foldon, an HRV3C cleavage site, an Octo-His tag, and a double strep-tag to allow secretory expression and purification.

Rhinolophus cornutus (*R.cor*, Genebank: BCG67443.1), human (hACE2, Genebank: BAB40370.1), *Rhinolophus ferrumequinum* (*R.ferr*, Genbank: BAH02663.1) and greater mouse-eared bat *Myotis lucifugus* (*M.luci*, Genbank: XP_023609438.1) ACE genes were synthesized. Pangolin (Pangolin, Genbank: XP_017505752.1), *Rhinolophus affinis* 9479 (*Ra9479*, Genbank: QMQ39227), *Rhinolophus affinis* 787 (*Ra5538*, Genbank: QMQ39222), *Rhinolophus sinicus* WJ2 (*Rs*-WJ2, Genbank: QMQ39202.1), *Rhinolophus sinicus* 1434 (*Rs*1434, Genbank: QMQ39216.1), *Rhinolophus sinicus* 1438 (*Rs*1438, Genbank: QMQ39203.1), *Rhinolophus sinicus* 1446 (*Rs*1446 Genbank: MT394194.1), *Rhinolophus sinicus* 3357 (*Rs*3357 Genbank: AGZ48803.1), *Rhinolophus sinicus* 3359 (*Rs*3359 Genbank: QMQ39211.1), *Rhinolophus sinicus* 3366 (*Rs*3366, Genbank: QMQ39215.1) and *Rhinolophus sinicus* 5720 (*Rs*5720, Genbank: MT394182.1) ACE2 gene sequences were from a previous study (65). The peptidase domain and the C-terminal collectrin-like domain of ACE2, bearing a carboxy-terminal Hexa-His tag, was generated for hACE2. Due to the presence of the C-terminal collectrin-like domain, these his-tagged ACE2 (AA 1-731) proteins form dimers. Additionally, the same length of ACE2s were fused to the Fc domain of human IgG with one thrombin site and a GGGG linker in between, resulting in Fc-tagged ACE2.

Rc-o319 RBD and variants (AA 293-506), SARS-CoV-2 RBD (AA 319-541), were fused with human IgG Fc domains to generate dimeric RBD-Fc proteins.

Protein expression and purification

Proteins were expressed by transient transfection in Expi293F cells, 1 mg of DNA was transfected into 1 L of cells using Linear Polyethylenimine (PEI) transfection reagent. After transfection, cells were cultured at 33 °C for 4 days for S-proteins, 5 days for RBD and ACE2 proteins. For his-tagged S-proteins and RBD proteins, the supernatants were harvested by centrifugation and 25 mM phosphate pH 8.0, 3 mM imidazole, 300 mM NaCl, and 0.5 mM PMSF were supplemented. Supernatant was recirculated onto a 5-ml Talon Cobalt column for about 3 times and the column was washed by 100 ml buffer A (25 mM phosphate pH 8.0, 5 mM imidazole, and 300 mM NaCl). Proteins were eluted using a linear gradient with increasing concentration of Buffer B (25 mM phosphate pH 8.0, 500 mM imidazole, and 300 mM NaCl). Eluted protein samples were quality checked by SDS-PAGE, concentrated and buffer-exchanged into PBS using a 100 kDa (for S-proteins) or 10 kDa (for RBD proteins) MWCO Amicon Ultra filtration device (Merck Millipore). The concentrated proteins were frozen in liquid nitrogen and stored at -80 °C.

For RBD-Fc and ACE2-Fc proteins, 5 days post-transfection, each culture supernatant was cleared by filtering through a 0.45 μm filter membrane before being loaded onto a 5 mL HiTrap Protein A column (Cytiva). The column was washed by PBS (10 mM Na_2HPO_4 and 1.8 mM KH_2PO_4 pH 7.4, 137 mM NaCl, and 2.7mM KCl). Proteins were eluted using a linear gradient with increasing concentration of a citrate buffer (0.1 M citric acid, pH 3.0). Eluted protein fractions were immediately neutralized with equal volumes of 1.0 M Tris-HCl pH 8.0 buffer. RBD-Fc and ACE2-Fc proteins were concentrated, buffer-exchanged into PBS before being frozen and stored at -80 °C.

Biolayer Interferometry (BLI) binding analysis

Binding of Rc-o319 S-proteins or RBDs to ACE2-Fc proteins was assessed using biolayer interferometry (BLI) on an Octet RED 96 instrument (FortéBio, USA). All steps were performed at 25 °C with an orbital shaking speed of 1000 rpm. All reagents were formulated in PBS-TB buffer (PBS with 0.02% v/v Tween-20 and 0.1% w/v BSA). Before the experiments, all protein A biosensors were pre-equilibrated in the PBS-TB buffer for 10 min. ACE2-Fc proteins (11 μg/ml) were immobilized onto Protein A biosensors (FortéBio, USA) to a response value of

~1.2 nm. After a 60 s baseline step in PBS-TB, the biosensors immobilized with
ACE2-Fc proteins were submerged in the analyte (S-proteins or RBDs) solutions
for 300 s to measure association, before being submerged in PBS-TB for 600 s to
measure dissociation. Data were reference subtracted and analyzed in the FortéBio
Data Analysis Software (FortéBio) by fitting to 1:1 or 2:1 binding models to
determine k_{on}, k_{off}, and K_D values as previously described (66). Raw data and fits
were plotted using GraphPad Prism 8 (GraphPad Software).

Binding of dimeric his-tagged hACE2, pangolin ACE2, and bACE2s to Rc-o319
RBD-Fc protein was assessed using the BLI assay. The RBD-Fc proteins (11
µg/ml) were loaded onto Protein A biosensors (FortéBio, USA) to a level of ~1.2
nm. After a 60 s baseline step in PBS-TB buffer, the biosensors immobilized with
RBD-Fc proteins were submerged in the ACE2 protein solutions for 300 s to
measure association, before being submerged in PBS-TB for 600 s to measure
dissociation. Data were reference subtracted and analyzed in the FortéBio Data
Analysis Software.

Cryo-EM grid preparation and data acquisition
Cryo-EM grids were prepared with Rc-o319 S-trimer samples at concentration of
1.65 mg/ml. All grids were prepared by applying 3.0 µl of S-trimer solutions
directly to glow discharged (15 mA, 30 s) holey carbon grids (Quantifoil Cu mesh
300 R1.2/R1.3) before blotting. Cryo-EM grids were blotted for 2.5 s with a blot
force of 4, before being plunge-frozen into liquid ethane using a Vitrobot Mark IV
(ThermoFisher) at 4 °C and 100% humidity.

For Rc-o319 S-proteins, micrographs were collected using the EPU software
(Thermo Fisher) on a Titan Krios G3i electron microscope (Thermo Fisher)
operated at 300 keV. Micrographs were recorded at a nominal magnification of
130,000× on a Falcon 4 direct electron detector coupled to a Selectris-X energy
filter set with a slit width of 10 eV and with a defocus range between -0.8 to -2.0
µm. Each movie was collected with a dose rate of 25 e⁻/pixel/second, and exposed
for 2.28 seconds, resulting in a total dose of 50 e⁻/Å² with a calibrated pixel size of
0.93 Å.

For complex formation, wild-type dimeric Rc-o319 RBD was mixed with dimeric
Fc-tagged bACE2$_{R.cor}$ at a 1.2:1 molar ratio, followed by incubation at 4 °C in the

presence of 0.01% glutaraldehyde for 20 min. The glutaraldehyde was quenched
by adding Tris-HCl to a final concentration of 50 mM. Gel filtration was
performed to remove excess RBD on a Superose 6 10/300 GL column (Cytiva)
equilibrated in PBS buffer. Complex formation was confirmed by SDS-PAGE, and
the Rc-o319-RBD:bACE2$_{R.cor}$ complex peak was collected and concentrated to
prepare cryo-EM grids. 3 μL of 6.8mg/mL the RBD-ACE2 complex was mixed
with 8 mM 3-[(3-Cholamidopropyl)dimethylammonio]-2-hydroxy-1-
propanesulfonate (CHAPSO) before being applied onto glow discharged (15 mA,
30 s) Ni-Au-300 mesh R1.2/1.3 grids. Grids were blotted for 2.5 s with a blot force
of 5 before being plunge frozen into liquid ethane using a Vitrobot Mark IV
(Thermo Fisher Scientific) at 4 °C and 100% humidity. Micrographs were
collected on a Titan Krios electron microscope (Thermo Fisher) operated at 300
keV using the EPU software (Thermo Fisher). Micrographs were recorded at a
nominal magnification of 130,000× on a K3 direct electron detector (Gatan)
coupled to a Gatan BioQuantum energy filter set with a slit width of 20 eV
(Thermo Fisher) and with a defocus range between -0.8 to -2.0 μm. Each movie
was collected with a dose rate of 18.7 e^{-}/pixel/second, exposed for 1.78 seconds,
resulting in a total dose of 50 e^{-}/Å^2 with a calibrated pixel size of 0.65 Å.

Image processing
For dataset processed by the CryoSPARC package (67), particles were picked
before 2D classification was performed. Well featured particles from the 2D
classification were subjected to *ab initio* 3D classification. Homogeneous
Refinement and Local Refinement were applied to the set of particles with good
features identified by the *ab initio* 3D classification. For Rc-o319 S-trimer, *ab*
initio 3D classification was performed with the resolution limit restricted to 10.0 Å
to minimize overfitting and emphasize large-scale conformational differences.
Eight initial classes were obtained from this procedure. Three well-resolved classes
were subsequently subjected to independent non-uniform refinement. The resulting
refined maps were then rigid-body aligned and compared using pairwise map-map
correlation analysis to quantitatively assess structural similarity among the classes.
Classes yielding similar structures were combined and subjected to final
refinement, resulting in two Rc-o319 S-trimer structures adopting distinct
conformations.

For the Rc-o319-RBD:bACE2$_{R.cor}$ complex dataset, data processing was performed
mainly using cryoSPARC v4.6.2 (67). After patch-motion correction and CTF
estimation, micrographs with CTF fit resolution above 6Å were picked by the
curate exposures job. Particles were then picked using a blob picker to generate
templates for the subsequent template picker. Picked particles were extracted at 3x
binning and subjected to 2D classification, *ab initio* reconstruction, and
heterogeneous refinement. Particles belongs to classes with high-resolution details
were selected for a final round of Topaz picker by the Topaz cross validation job.
Picked particles were extracted at 3x binning and subjected to 2D classification, *ab*
initio reconstruction, and heterogeneous refinement again. Highly similar classes in
heterogeneous refinement showing high-resolution details were combined and re-
extracted at 1.5x binning and subjected to non-uniform refinement with
optimization of per-group CTF parameter analysis. The CTF-refined particles and
volumes were subjected to reference-based motion correction and an extra round of
non-uniform refinement to get the final dimeric structure.

Due to conformational dynamics in the dimeric RBD-ACE2 complex structure,
several local regions were resolved to lower resolution. Local refinement was
applied to improve the local resolution at the RBD-ACE2 binding interface.
Particles were symmetry expanded, the dynamic parts were masked, before the
particles were subtracted through masking. The subtracted particles were imported
to perform NU-Refine and local refinement jobs.

Model building and structure refinement
A previous determined SARS-CoV-2 S-trimer structure (PDB ID: 6ZP2) (22) was
used as the starting models for the Rc-o319 S-trimer structures. Model building
and adjustment were performed manually in Coot 0.9 (68). Steric clash and
sidechain rotamer conformations were improved using the Namdinator web server
(69). Final structures after additional manual adjustments were refined and
validated in PHENIX-1.18.261 (70). The SARS-CoV-2-RBD:hACE2-B^{0}AT1
complex structure (PDB ID: 6M17, EMD-30039) (27, 71) was used as the starting
model for the Rc-o319 RBD:bACE2$_{R.cor}$ complex structure. The starting model was
fitted in the cryo-EM density in Chimera 1.18.0 before manual adjustments were
carried out in Coot 0.9. The model was refined by multiple rounds of refinement in

Namdinator and PHENIX. Data collection and refinement statistics for all the
reported structures are provided in Table S1.

GFP-split fusion assay
The split-GFP system based cell-cell fusion assay mediated by the interaction
between the S-protein and the ACE2 receptor was previously described (37).
Briefly, HEK293T cells transfected with full length S-proteins and pQCXIP-
GFP1-10 were prepared as effector cells, while HEK293T cells transfected with
bat ACE2 (*R.cor*, *Rs*-WJ2, *Rs*1446, *Rs*1434, *Rs*1438, *Rs*3357, *Rs*3366, *Rs*3359,
*Rs*5720, *Ra*9479, *Ra*5538, *R.ferr*, and *M.luci*) or pangolin ACE2 or human ACE2
(hACE2) and pQCXIP-BSR-GFP11 served as receptor cells. 24 h after transfection,
effector and receptor cells were washed and resuspended in DMEM containing 10%
FBS without trypsin treatment, mixed at a 1:1 ratio, and plated at ~2×10^5 cells per
well in a 96-well plate. Fluorescence images were recorded at the indicated time
points using a Nikon TS2-S-SM fluorescence microscope. The GFP area was
quantified with ImageJ. S-protein expression was analyzed via flow cytometry
using a SARS-CoV-2 S2 antibody (1:2500 dilution, Sino Biological, 40590-T62)
and western blot using an anti-FLAG antibody (1:1000 dilution, Sino Biological,
109143-MM13). The GFP-positive area was normalized to the mean fluorescence
intensity of surface S-proteins, and resulting values used to calculate the area under
the curve (AUC) for statistical analysis.

VSV-Pseudovirus entry assay
VSV pseudotypes were generated according to a published protocol (72). Briefly,
HEK293T cells were transfected with 50 μg Rc-o319 WT or variant S plasmid into
a 15 cm cell culture dish. After 24h, the cells were washed twice and inoculated
with VSV*△G-Luc at a multiplicity of infection of 0.1 for 1h. After the inoculum
was removed, the cells were washed 5 times using PBS with 2% FPS and further
cultured with a DMEM culture medium for 24h. The supernatant was harvested
and centrifuged at 3000 rpm to free cellular debris, filtered through a 0.45 μm
syringe filter, 10-fold concentration and stored at– 80°C in small aliquots.

To quantify entry of pseudovirons into cells, HEK293T cells stably co-expressing
TMPRSS2 and ACE2 (2×10^5 cells in 100 μL DMEM per well) were seeded in 96-
well plates and inoculated with an equal amount of pseudotype particles. 24 h post-

transduction, culture supernatants were aspirated and cells lysed in 1x cell culture
lysis reagent (prepared from 5x stock, Promega) for 20 min at room temperature.
The lysates were then transferred to white 96-well plates and luciferase activity
was analyzed by the luciferase assay system (Promega).

<u>Western Blot</u>
To analyze S proteins and ACE2 expression in cells, HEK293T cells were
transfected with expression vectors and then incubated for an additional 24h. Then,
cells were washed with ice-cold PBS, and soluble proteins were extracted with cell
lysis buffer (100 mM Tris-HCl pH 8, 150 mM NaCl, 1% NP-40) containing 1 mM
PMSF (Beyotime, ST505) and 2.5 U/mL BeyoZonase Super Nuclease (Beyotime,
D7121). Non-reducing SDS–PAGE loading buffer (4×) was added to each sample
immediately before boiling at 98 °C for 10 minutes and subjected to SDS-PAGE
and immunoblotting.

To analyze of S protein incorporation into pseudotyped particles, 1 mL of the each
VSV pseudotype preparation was subjected to ultracentrifugation (25 000 × g, 120
min, 4°C). The supernatant (1 mL) was carefully removed, and the remaining
pellet was resuspended in 20 μL of 5x SDS-sample buffer, heated at 98 °C for 10
min and then analyzed by SDS-PAGE and immunoblotting.

Equal amounts of protein samples were separated by 4-10% SDS-PAGE and
transferred to PVDF membranes (Millipore). After blocking with 5% (w/v)
skimmed milk in TBST buffer, the membranes were incubated with primary
antibodies, such as rabbit anti-SARS-CoV-2 S2 polyclonal antibody (1:2,500
dilution, Sino Biological, 40590-T62), mouse monoclonal antibody targeting
FLAG tag (1:2,500 dilution, TransGen Biotech, HT201), beta-actin (1:5,000
dilution, TransGen Biotech, HC201) and mouse anti-VSV matrix protein (Kerafast,
EB0011, 1:2,500). Horseradish peroxidase-conjugated (HRP) goat anti-mouse IgG
antibody (1:10,000 dilution, Beyotime, A0216) and goat anti-rabbit IgG antibody
(1:10,000 dilution, Beyotime, A0208) were used as secondary antibodies. The
quantitative results of VSV-M were analyzed using ImageJ software.

Flow cytometry

To validate surface expression of S protein and its binding affinity for ACE2
following transfection, we performed flow cytometry using a BD Accuri C6 Plus
flow cytometer, and data were analyzed with FlowJo software. HEK293T cells
transfected with S protein expression plasmids were harvested 24 h post-
transfection by centrifugation (3,000 rpm, 5 min, 4 °C), resuspended in ice-
cold PBS containing 1% BSA, and incubated on ice for 1 h with rabbit anti–
SARS-CoV-2 S2 polyclonal antibody (Sino Biological, Cat. #40590-T62, 1:500)
and/or recombinant human ACE2-Fc fusion protein (10 μg/mL). Cells were then
washed twice with PBS and incubated on ice for 45 min with Alexa Fluor 488–
conjugated goat anti-rabbit IgG (H+L) secondary antibody (Beyotime, A0423,
1:500) and/or APC-conjugated anti-human IgG Fc antibody (BioLegend, 410711,
1:1000).

Statistical analysis

880
All statistical analyses were performed using GraphPad Prism software (Graphpad
Version 10.6.1). The significance of differences between the two groups was
determined with an unpaired, two-tailed Student's *t*-test. For all analyses, only a
probability (*p*) value of 0.05 or lower were considered statistically significant ($p >$
0.05 [n.s, not significant], $p \leq 0.05$ [*], $p \leq 0.01$ [**], $p \leq 0.001$ [***], $p \leq 0.0001$
[****]).

Data availability

Cryo-EM density maps and coordinates generated in this study have been deposited in the Electron Microscopy Data Bank (EMDB; www.ebi.ac.uk/emdb/) and the Protein Data Bank (PDB; www.rcsb.org/ with the following accession numbers: Rc-o319 locked-1 S-trimer: EMD-65717 and PDB:9W75; Rc-o319 locked-2 S-trimer: EMD-65718 and PDB: 9W76; Dimeric Rc-o319-RBD:bACE2$_{R.cor}$ complex: EMD-65719 and PDB: 9W77, monomeric Rc-o319-RBD:bACE2$_{R.cor}$ complex: EMD65720 and PDB: 9W78. All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Acknowledgments

We thank the staff of the GIBH-CAS Cryo-EM Facility for their help with cryo-EM sample preparation and data collection. We thank the staff of the Guangzhou Laboratory Cryo-EM Facility for their help with cryo-EM sample data collection. This work was supported by the National Natural Science Foundation of China (32570199 to X.X.; 82341085 to X.X.); the National Key R&D Program of China (2021YFA1300903 to X.X.); the Emergency Key Program of Guangzhou Laboratory (EKPG21-06 to X.X.); the R&D Program of Guangzhou National Laboratory (SRPG22-002 to X.X.; SRPG22-003 to J.H); the Guangdong Basic and Applied Basic Research Foundation (2021A1515011289 to X.X.); the Science and Technology Planning Project of Guangdong Province, China (2023B1212060050 to X.X.; 2023B1212120009 to X.X.); the Basic Research Project of Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences (GIBHBRP24-02 to X.X.); the Natural Science Fund of Guangdong Province (2022A1515110211 to J.W.; 2614050002966 to J.W.); the Young Doctoral Starting Sail Project of the Guangzhou Municipal Science and Technology Bureau (2024A04J4197 to J.W.; 2024A04J4357 to Y.M.); and the Grants from State Key Laboratory of Respiratory Diseases (SKLRD-Z-202515 to Y.M.). X.X. acknowledges Start-up grants from the Chinese Academy of Sciences.

Author contributions

J.W., Z.L., and Y.M. conceived the study under the direction of X.C. and X.X. J.W. and Z.L. performed the clone design, protein purification and EM-structure

determination study; Y.M. performed the cell-cell fusion and FCAS assays; Z. L.,
Y.T., H.Y. performed BLI assays; J.W., B.L., C.N. and X.X. analysied the BLI
results; J.W., Z.L., M.L., H.F. and M.Z. helped collect cryo-EM data under the
supervision of J.H.and X.X.; Z.L., Zimu Li and J.W. processed cryo-EM data
under the supervision of X.X.; J.W. Z.L. and X.X. analyzed cryo-EM structures;
J.W., Z.L. and Y.M. prepared the figures under the supervision of X.X.; J.W., Z.L.,
Y.M. and X.X. wrote the paper with input from all authors; J.C. and J.H. reviewed
and edited the manuscript; X.C. and X.X. supervised the research.
932

933   **Conflict of interest**
934   The authors declare no competing interests.

References

1. Peiris JS, Chu CM, Cheng VC, Chan KS, Hung IF, Poon LL, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. Lancet. 2003;361(9371):1767-72.
2. Drosten C, Gunther S, Preiser W, van der Werf S, Brodt HR, Becker S, et al. Identification of a novel coronavirus in patients with severe acute respiratory syndrome. N Engl J Med. 2003;348(20):1967-76.
3. Fouchier RA, Kuiken T, Schutten M, van Amerongen G, van Doornum GJ, van den Hoogen BG, et al. Aetiology: Koch's postulates fulfilled for SARS virus. Nature. 2003;423(6937):240.
4. Peiris JS, Guan Y, Yuen KY. Severe acute respiratory syndrome. Nat Med. 2004;10(12 Suppl):S88-97.
5. Temmam S, Vongphayloth K, Baquero E, Munier S, Bonomi M, Regnault B, et al. Bat coronaviruses related to SARS-CoV-2 and infectious for human cells. Nature. 2022;604(7905):330-6.
6. Wacharapluesadee S, Tan CW, Maneeorn P, Duengkae P, Zhu F, Joyjinda Y, et al. Evidence for SARS-CoV-2 related coronaviruses circulating in bats and pangolins in Southeast Asia. Nat Commun. 2021;12(1):972.
7. Murakami S, Kitamura T, Suzuki J, Sato R, Aoi T, Fujii M, et al. Detection and Characterization of Bat Sarbecovirus Phylogenetically Related to SARS-CoV-2, Japan. Emerg Infect Dis. 2020;26(12):3025-9.
8. Drexler JF, Gloza-Rausch F, Glende J, Corman VM, Muth D, Goettsche M, et al. Genomic characterization of severe acute respiratory syndrome-related coronavirus in European bats and classification of coronaviruses based on partial RNA-dependent RNA polymerase gene sequences. J Virol. 2010;84(21):11336-49.
9. Tao Y, Tong S. Complete Genome Sequence of a Severe Acute Respiratory Syndrome-Related Coronavirus from Kenyan Bats. Microbiol Resour Announc. 2019;8(28).
10. Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, et al. Bats are natural reservoirs of SARS-like coronaviruses. Science. 2005;310(5748):676-9.
11. Ge XY, Li JL, Yang XL, Chmura AA, Zhu G, Epstein JH, et al. Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. Nature. 2013;503(7477):535-8.
12. Hu B, Ge X, Wang LF, Shi Z. Bat origin of human coronaviruses. Virol J. 2015;12:221.
13. Zhou H, Ji J, Chen X, Bi Y, Li J, Wang Q, et al. Identification of novel bat coronaviruses sheds light on the evolutionary origins of SARS-CoV-2 and related viruses. Cell. 2021;184(17):4380-91 e14.
14. Delaune D, Hul V, Karlsson EA, Hassanin A, Ou TP, Baidaliuk A, et al. A novel SARS-CoV-2 related coronavirus in bats from Cambodia. Nat Commun. 2021;12(1):6563.
15. Alkhovsky S, Lenshin S, Romashin A, Vishnevskaya T, Vyshemirsky O, Bulycheva Y, et al. SARS-like Coronaviruses in Horseshoe Bats (Rhinolophus spp.) in Russia, 2020. Viruses. 2022;14(1):113.

16.   Guo H, Li A, Dong TY, Si HR, Hu B, Li B, et al. Isolation of ACE2-dependent and
-independent sarbecoviruses from Chinese horseshoe bats. J Virol.
2023;97(9):e0039523.
17.   Wang J, Ma Y, Li Z, Yuan H, Liu B, Li Z, et al. SARS-related coronavirus S-
protein structures reveal synergistic RBM interactions underpinning high-affinity human
ACE2 binding. Sci Adv. 2025;11(11):eadr8772.
18.   Wells HL, Letko M, Lasso G, Ssebide B, Nziza J, Byarugaba DK, et al. The
evolutionary history of ACE2 usage within the coronavirus subgenus Sarbecovirus.
Virus Evol. 2021;7(1):veab007.
19.   Zhang J, Xiao T, Cai Y, Chen B. Structure of SARS-CoV-2 spike protein. Curr
Opin Virol. 2021;50:173-82.
20.   Bosch BJ, van der Zee R, de Haan CA, Rottier PJ. The coronavirus spike protein
is a class I virus fusion protein: structural and functional characterization of the fusion
core complex. J Virol. 2003;77(16):8801-11.
21.   Harrison SC. Viral membrane fusion. Nat Struct Mol Biol. 2008;15(7):690-8.
22.   Xiong X, Qu K, Ciazynska KA, Hosmillo M, Carter AP, Ebrahimi S, et al. A
thermostable, closed SARS-CoV-2 spike protein trimer. Nat Struct Mol Biol.
2020;27(10):934-41.
23.   Gui M, Song W, Zhou H, Xu J, Chen S, Xiang Y, et al. Cryo-electron microscopy
structures of the SARS-CoV spike glycoprotein reveal a prerequisite conformational
state for receptor binding. Cell Res. 2017;27(1):119-29.
24.   Song W, Gui M, Wang X, Xiang Y. Cryo-EM structure of the SARS coronavirus
spike glycoprotein in complex with its host cell receptor ACE2. PLoS Pathog.
2018;14(8):e1007236.
25.   Zhang X, Li Z, Zhang Y, Liu Y, Wang J, Liu B, et al. Disulfide stabilization reveals
conserved dynamic features between SARS-CoV-1 and SARS-CoV-2 spikes. Life Sci
Alliance. 2023;6(9):e202201796.
26.   Li F, Li W, Farzan M, Harrison SC. Structure of SARS coronavirus spike
receptor-binding domain complexed with receptor. Science. 2005;309(5742):1864-8.
27.   Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition
of SARS-CoV-2 by full-length human ACE2. Science. 2020;367(6485):1444-8.
28.   Si JY, Chen YM, Sun YH, Gu MX, Huang ML, Shi LL, et al. Sarbecovirus RBD
indels and specific residues dictating multi-species ACE2 adaptiveness. Nat Commun.
2024;15(1):8869.
29.   Li W, Zhang C, Sui J, Kuhn JH, Moore MJ, Luo S, et al. Receptor and viral
determinants of SARS-coronavirus adaptation to human ACE2. EMBO J.
2005;24(8):1634-43.
30.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia
outbreak associated with a new coronavirus of probable bat origin. Nature.
2020;579(7798):270-3.
31.   Han P, Su C, Zhang Y, Bai C, Zheng A, Qiao C, et al. Molecular insights into
receptor binding of recent emerging SARS-CoV-2 variants. Nat Commun.
2021;12(1):6103.
32.   Crook JM, Murphy I, Carter DP, Pullan ST, Carroll M, Vipond R, et al.
Metagenomic identification of a new sarbecovirus from horseshoe bats in Europe.
Scientific Reports. 2021;11(1):14723.

33.   Markotter W, Geldenhuys M, Jansen van Vuren P, Kemp A, Mortlock M,
Mudakikwa A, et al. Paramyxo- and Coronaviruses in Rwandan Bats. Trop Med Infect
Dis. 2019;4(3):99.
34.   Guo H, Hu B, Si HR, Zhu Y, Zhang W, Li B, et al. Identification of a novel lineage
bat SARS-related coronaviruses that use bat ACE2 receptor. Emerg Microbes Infect.
2021;10(1):1507-14.
35.   Qiao S, Wang X. Structural determinants of spike infectivity in bat SARS-like
coronaviruses RsSHC014 and WIV1. J Virol. 2024;98(8):e0034224.
36.   Hu Y, Liu K, Han P, Xu Z, Zheng A, Pan X, et al. Host range and structural
analysis of bat-origin RshSTT182/200 coronavirus binding to human ACE2 and its
animal orthologs. EMBO J. 2022:e111737.
37.   Ma Y, Li P, Hu Y, Qiu T, Wang L, Lu H, et al. Spike substitution T813S increases
Sarbecovirus fusogenicity by enhancing the usage of TMPRSS2. PLOS Pathogens.
2023;19(5):e1011123.
38.   Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, Jr., et al. Distinct
conformational states of SARS-CoV-2 spike protein. Science. 2020;369(6511):1586-92.
39.   Ke Z, Oton J, Qu K, Cortese M, Zila V, McKeane L, et al. Structures and
distributions of SARS-CoV-2 spike proteins on intact virions. Nature.
2020;588(7838):498-502.
40.   Qu K, Chen Q, Ciazynska KA, Liu B, Zhang X, Wang J, et al. Engineered
disulfide reveals structural dynamics of locked SARS-CoV-2 spike. PLoS Pathog.
2022;18(7):e1010583.
41.   Toelzer C, Gupta K, Yadav SKN, Borucu U, Davidson AD, Kavanagh Williamson
M, et al. Free fatty acid binding pocket in the locked structure of SARS-CoV-2 spike
protein. Science. 2020;370(6517):725-30.
42.   Zhang S, Qiao S, Yu J, Zeng J, Shan S, Tian L, et al. Bat and pangolin
coronavirus spike glycoprotein structures provide insights into SARS-CoV-2 evolution.
Nat Commun. 2021;12(1):1607.
43.   Ma J, Su D, Sun Y, Huang X, Liang Y, Fang L, et al. Cryo-electron Microscopy
Structure of S-Trimer, a Subunit Vaccine Candidate for COVID-19. J Virol.
2021;95(11):10.1128/jvi.00194-21.
44.   Rosa A, Pye VE, Graham C, Muir L, Seow J, Ng KW, et al. SARS-CoV-2 can
recruit a heme metabolite to evade antibody immunity. Sci Adv. 2021;7(22):eabg7607.
45.   Freeman SL, Oliveira ASF, Gallio AE, Rosa A, Simitakou MK, Arthur CJ, et al.
Heme binding to the SARS-CoV-2 spike glycoprotein. J Biol Chem. 2023;299(8):105014.
46.   Lee J, Zepeda SK, Park YJ, Taylor AL, Quispe J, Stewart C, et al. Broad receptor
tropism and immunogenicity of a clade 3 sarbecovirus. Cell Host Microbe.
2023;31(12):1961-73 e11.
47.   Li F. Structural analysis of major species barriers between humans and palm
civets for severe acute respiratory syndrome coronavirus infections. J Virol.
2008;82(14):6984-91.
48.   Dadonaite B, Brown J, McMahon TE, Farrell AG, Figgins MD, Asarnow D, et al.
Spike deep mutational scanning helps predict success of SARS-CoV-2 clades. Nature.
2024;631(8021):617-26.

49.   Harvey WT, Carabelli AM, Jackson B, Gupta RK, Thomson EC, Harrison EM, et
al. SARS-CoV-2 variants, spike mutations and immune escape. Nat Rev Microbiol.
2021;19(7):409-24.
50.   McCallum M, De Marco A, Lempp FA, Tortorici MA, Pinto D, Walls AC, et al. N-
terminal domain antigenic mapping reveals a site of vulnerability for SARS-CoV-2. Cell.
2021;184(9):2332-47 e16.
51.   Yuan H, Wang J, Ma Y, Li Z, Gao X, Habib G, et al. Structures and receptor
binding activities of merbecovirus spike proteins reveal key signatures for human DPP4
adaptation. Sci Adv. 2025;11(28):eadv7296.
52.   Chornelia A, Hughes AC. The evolutionary history and ancestral biogeographic
range estimation of old-world Rhinolophidae and Hipposideridae (Chiroptera). BMC
Ecol Evol. 2022;22(1):112.
53.   Shuai H, Chan JF, Yuen TT, Yoon C, Hu JC, Wen L, et al. Emerging SARS-CoV-
2 variants expand species tropism to murines. EBioMedicine. 2021;73:103643.
54.   Hills FR, Eruera AR, Hodgkinson-Bean J, Jorge F, Easingwood R, Brown SHJ, et
al. Variation in structural motifs within SARS-related coronavirus spike proteins. PLoS
Pathog. 2024;20(5):e1012158.
55.   Zhou T, Tsybovsky Y, Gorman J, Rapp M, Cerutti G, Chuang GY, et al. Cryo-EM
Structures of SARS-CoV-2 Spike without and with ACE2 Reveal a pH-Dependent
Switch to Mediate Endosomal Positioning of Receptor-Binding Domains. Cell Host
Microbe. 2020;28(6):867-79 e5.
56.   Ou X, Xu G, Li P, Liu Y, Zan F, Liu P, et al. Host susceptibility and structural and
immunological insight of S proteins of two SARS-CoV-2 closely related bat
coronaviruses. Cell Discov. 2023;9(1):78.
57.   Kosugi Y, Matsumoto K, Lytras S, Plianchaisuk A, Tolentino JE, Fujita S, et al.
Molecular basis of sarbecovirus evolution and receptor tropism in natural hosts,
potential intermediate hosts, and humans. Cell Rep. 2025;44(9):116220.
58.   Park YJ, Liu C, Lee J, Brown JT, Ma CB, Liu P, et al. Molecular basis of
convergent evolution of ACE2 receptor utilization among HKU5 coronaviruses. Cell.
2025;188(6):1711-28 e21.
59.   Chen J, Zhang W, Li Y, Liu C, Dong T, Chen H, et al. Bat-infecting merbecovirus
HKU5-CoV lineage 2 can use human ACE2 as a cell entry receptor. Cell.
2025;188(6):1729-42 e16.
60.   Liu B, Niu X, Deng Y, Zhang Z, Wang Y, Gao X, et al. An unconventional VH1-2
antibody tolerates escape mutations and shows an antigenic hotspot on SARS-CoV-2
spike. Cell Rep. 2024;43(6):114265.
61.   Yan Q, Gao X, Liu B, Hou R, He P, Ma Y, et al. Antibodies utilizing VL6-57 light
chains target a convergent cryptic epitope on SARS-CoV-2 spike protein and potentially
drive the genesis of Omicron variants. Nat Commun. 2024;15(1):7585.
62.   He P, Liu B, Gao X, Yan Q, Pei R, Sun J, et al. SARS-CoV-2 Delta and Omicron
variants evade population antibody response by mutations in a single spike epitope. Nat
Microbiol. 2022;7(10):1635-49.
63.   Yu H, Liu B, Zhang Y, Gao X, Wang Q, Xiang H, et al. Somatically hypermutated
antibodies isolated from SARS-CoV-2 Delta infected patients cross-neutralize
heterologous variants. Nat Commun. 2023;14(1):1058.

64.   Chen B, Farzan M, Choe H. SARS-CoV-2 spike protein: structure, viral entry and
variants. Nat Rev Microbiol. 2025;23(7):455-68.
65.   Guo H, Hu BJ, Yang XL, Zeng LP, Li B, Ouyang SY, et al. Evolutionary Arms
Race between Virus and Host Drives Genetic Diversity in Bat Severe Acute Respiratory
Syndrome-Related Coronavirus Spike Genes. J Virol. 2020;94(20).
66.   Carnell GW, Ciazynska KA, Wells DA, Xiong X, Aguinam ET, McLaughlin SH, et
al. SARS-CoV-2 Spike Protein Stabilized in the Closed State Induces Potent
Neutralizing Responses. J Virol. 2021;95(15):e0020321.
67.   Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. Nat Methods. 2017;14(3):290-6.
68.   Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr. 2010;66(Pt 4):486-501.
69.   Kidmose RT, Juhl J, Nissen P, Boesen T, Karlsen JL, Pedersen BP. Namdinator
- automatic molecular dynamics flexible fitting of structural models into cryo-EM and
crystallography experimental maps. IUCrJ. 2019;6(Pt 4):526-31.
70.   Afonine PV, Poon BK, Read RJ, Sobolev OV, Terwilliger TC, Urzhumtsev A, et al.
Real-space refinement in PHENIX for cryo-EM and crystallography. Acta Crystallogr D
Struct Biol. 2018;74(Pt 6):531-44.
71.   Buchrieser J, Dufloo J, Hubert M, Monel B, Planas D, Rajah MM, et al. Syncytia
formation by SARS-CoV-2-infected cells. EMBO J. 2020;39(23):e106267.
72.   Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et
al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a
Clinically Proven Protease Inhibitor. Cell. 2020;181(2):271-80 e8.

1139

1140

Fig 1. Rc-o319 S-protein utilizes a narrow range of *Rhinolophus* bat ACE2

orthologs as receptors and adopts two locked, 3-RBD-down conformations.

(A) Cell-cell fusion assays using receptor cells expressing bat, pangolin or human

ACE2 molecules and effector cells expressing the Rc-o319 S-proteins. Fused cells

show GFP signals. Representative images of cell-cell fusion captured at 6 hours

post-transfection are shown. Scale bar, 500 μm. **(B)** Binding of Rc-o319 S-trimer

to immobilized, dimeric $bACE2_{R.cor}$, hACE2 and $bACE2_{Ra9479}$ as assessed by BLI

assays. **(C)** A schematic representation showing the eight functional domains or

regions of Rc-o319 S-protein. **(D)** Cryo-EM density and associated molecular

model showing the overall architecture of a Rc-o319 S-trimer in a locked-1

conformation. **(E)** Lipid binding pocket in the RBD of locked-1 Rc-o319 S-trimer.

Modeled linoleic acid (magenta) interacts with the shown hydrophobic residues of

the lipid binding pocket. Salt bridge and hydrogen bond are shown as dashed lines.

(F) Structure of the Domain D region in the locked-1 Rc-o319 S-trimer. Consistent

with a locked conformation, the fusion peptide proximal region (FPPR) is ordered.

Residues $580\text{-}605_{Rc\text{-}o319}$ form a large disordered loop in the locked-1 conformation.

(G) Cryo-EM density and associated molecular model showing the overall

architecture of a locked-2 Rc-o319 S-trimer. **(H)** The RBD lipid binding pocket is

empty with $Y339_{Rc\text{-}o319}$ side-chain reorientated in the locked-2 conformation. **(I)**

Residues $580\text{-}605_{Rc\text{-}o319}$ refold and form an ordered structure. A stabilizing cation-

π interaction (red dashed line) is formed between Domain D residue $W598_{Rc\text{-}o319}$

and Domain C-D junction residue $R293_{Rc\text{-}o319}$.

1163

1164

Fig 2. Rc-o319 S-trimer structures reveal an RBD featuring a novel RBM and an NTD with altered loops.

(A-B) Topviews of locked-1 and locked-2 Rc-o319 S-trimer structures. N-linked glycans are shown as light gray spheres. NTDs are colored blue, RBDs are colored orange (locked-1, A) or pink (locked-2, B). The lower panels show intra- or inter-protomer interactions involving RBD residues. $K458_{Rc-o319}$ forms an inter-protomer hydrogen bond with $N344_{Rc-o319}$ of an RBD of a neighboring protomer in both locked-1 and locked-2 S-trimers. In the locked-1 conformation, $D468_{Rc-o319}$ forms an intra-protomer hydrogen bond with $Y473_{Rc-o319}$; in the locked-2 conformation, $D468_{Rc-o319}$ instead mediates two inter-protomer hydrogen bonds with $S469_{Rc-o319}$ of an adjacent protomer. **(C-D)** Comparison of Rc-o319 (C) and SARS-CoV-2 (D) N-terminal domain (NTD) structures. The five highly variable NTD loops (N1-N5) are shown in the Rc-o319 and SARS-CoV-2 (PDB: 7XU4) NTD structures. A four-amino-acid residue insertion is highlighted in the Rc-o319 NTD in red. **(E)** The structure of the Rc-o319 RBD. The receptor binding motif (RBM) of the RBD is colored pink and the core of the RBD is colored gray. Disulfide bonds within the RBD are indicated. Featured structural elements, beta-Loop (BL), small loop (SL), Lamella (LM) sheets, β5 and β6, and the large RBM loop are indicated. **(E-G)** For comparison, the SARS-CoV-2 RBD (PDB:7XU4) is shown (F) and superposed on the Rc-o319 RBD structure (E).

1186

Fig 3. Cryo-EM structure of the Rc-o319-RBD:bACE2$_{R.cor}$ complex with a detailed comparison of RBD:ACE2 interactions versus those in the SARS-CoV-2-RBD:hACE2 complex.

(A) Molecular structure of a monomeric Rc-o319-RBD:bACE2$_{R.cor}$ complex
extracted from the dimeric Rc-o319-RBD:bACE2$_{R.cor}$ complex structure resolved
by cryo-EM (see Figure S1). Detailed interactions formed between the Rc-o319-
RBD and bACE2$_{R.cor}$ are shown with hydrogen bonds and salt bridges shown as
dashed lines. **(B)** Superposition of the Rc-o319-RBD:bACE2$_{R.cor}$ and SARS-CoV-
2-RBD:hACE2 complex structures. The interface areas between the RBD and
ACE2 in the two structures are divided into four regions: beta-loop (BL in Rc-
o319-RBD)/large-loop (LL in SARS-CoV-2-RBD) (purple box), lamella (LM,
blue box), anchor-loop (AL, green box), and small-loop (SL, orange box). **(C-D)**
Comparison of the Rc-o319 RBM-loop and beta-loop (BL) with the RBM-loop and
large-loop (LL) of SARS-CoV-2. **(E)** An N-linked glycan (shown in grey stick
representation overlaid with white cryo-EM density) attached to Asn38 of
bACE2$_{R.cor}$ is involved in the binding interface with Rc-o319-RBD, representing a
unique structural feature. **(F-I)** Detailed interactions identified in the beta-
loop+RBM-loop (BL, purple box), lamella (LM, blue box), anchor-loop (AL,
green box), and small-loop (SL, orange box) regions of the interface between Rc-
o319-RBD and bACE2$_{R.cor}$. **(J-M)** For comparison, detailed interactions are shown
for the corresponding regions in the interface between SARS-CoV-2-RBD and
hACE2. Hydrogen bonds, pi-pi (sugar-pi) interactions and cation-pi interactions
are indicated by black, yellow (green) and red dashed lines, respectively.

1210

1211

Fig 4. ACE2 utilization by wild-type and variant Rc-o319 RBD-Fc in BLI and wild-type and variant Rc-o319 S-proteins in cell-cell fusion and Pseudovirus entry assays.

(A-L) Top two panels, binding of bACE2$_{R.cor}$ and hACE2 by wild-type (WT) and variant Rc-o319-RBD-Fc proteins as assessed by BLI assays. Dimeric Rc-o319-RBD-Fc proteins were immobilized on the BLI sensors and tested binding against dimeric *Rhinolophus cornutus* bat ACE2 (bACE2$_{R.cor}$) or hACE2 protein as the analyte in solution. bACE2$_{R.cor}$ and hACE2 were three-fold serial diluted from 3000 to 4.1 nM. K_D values estimated from each binding experiment are shown alongside the corresponding binding curves. Lower two panels, Representative images of cell-cell fusion captured at 12 hours post-transfection are shown. Scale

bar, 500 μm. Effector cells expressing WT or variant Rc-o319 S-proteins were
tested against receptor cells expressing either bACE2$_{R.cor}$ or hACE2. The bottom
panels show the quantification of cell-cell fusion measuring the GFP+ area at 2, 6,
12 hours (h) post-transfection. Data are presented as mean ± s.d., n = 2. The *P*
values were determined by unpaired t-test comparing wild type (WT) with each
Rc-o319 variant. **P* < 0.05, ***P* < 0.01, ****P* < 0.001; NS, not significant.
(M) Pseudovirus assay. Entry of VSV particles pseudotyped with Rc-o319 S-
protein variants into HEK293T cells transiently expressing hACE2 or bACE2$_{R.cor}$
and co-expressing TMPRSS2. Luciferase activity was measured 24 h post-
transduction. Data are presented as mean ± s.d. (n = 3). *P* values were calculated
using unpaired t tests comparing wild-type (WT) with each Rc-o319 variant.
**P* < 0.05, ***P* < 0.01, ****P* < 0.001 and *****P* < 0.0001.

1235

1236

Fig 5. bACE2$_{R.cor}$ utilization by different sarbecoviruses S-proteins. (A)
Different sarbecovirus S-proteins are sorted based on S-protein phylogenetics
(clade 1-3) and grouped by RBD types defined structurally. Type-1 RBD strains
are shown in red (SARS-CoV-2 clade, Clade 1b) and blue (SARS-CoV-1 clade,
clade 1a), type-2 strains in orange, type-3 strains in black and type-1b strains in
green. Quantification of S-protein expression on the cell surface of effector cells by
flow cytometry using an S2 antibody as the probe. Flow cytometry was repeated
three times with representative data shown. **(B)** Effector cells shown in (A) were
tested against receptor cells expressing hACE2 (gray line), bACE2$_{R.cor}$ (red line)
and bACE2$_{R.cor}$-ΔGLC$_{38}$ (green line, bACE2$_{R.cor}$ with the Thr40Ala mutation to
remove the Asn38-glycan). Cell-cell fusion was quantified by assessing GFP+
areas (y-axis) at 4, 18, and 24 hours (h, x-axis) post-transfection (pt). Data are
presented as mean ± s.d., n = 2. *P* values were calculated using unpaired t-tests
comparing fusion activities mediated by bACE2$_{R.cor}$ and bACE2$_{R.cor}$-ΔGLC$_{38}$.
*$P\square<\square0.05$, **$P\square<\square0.01$, ***$P\square<\square0.001$; NS, not significant. **(C)**
Representative images of cell-cell fusion captured at 24 hours post-transfection are
shown against receptor cells expressing different ACE2. Scale bar, 500 μm.

1254

1255