Effects of Simulated Microgravity on Human Hematological and Anemia-Related Biomarkers: A Systematic Review and Meta-Analysis

Ahmed Ayman Mohamed Abolyazed[1], Wafaa Ahmed Elbehery[1], Hanan Mahmoud Elsayed[2], Yousef Hamdy ElSheeta[1], Abdalla Yasser Elnemr[1]

[1] Faculty of Medicine, Tanta University, Tanta, Egypt

[2] Department of Pediatrics and Neonatology, Tanta General Hospital, Tanta, Egypt

Correspondence:

Ahmed Ayman Mohamed Abolyazed

Ahmed.31010459@med.tanta.edu.eg

Ahmedabolyazed21@gmail.com

Keywords: Space anemia, microgravity, spaceflight, ground-based analogs.

Abstract

Background: Space anemia has been observed in astronauts after spaceflights for a long time, but its exact mechanism remains unclear. Because of the practical and ethical challenges of performing human studies in space, researchers use ground-based analogs to study the hematological changes that occur upon exposure to microgravity. Understanding these changes is important for protecting astronauts during long spaceflights and for guiding medical treatment on Earth in cases of prolonged inactivity or bed rest.

Methods: The literature search was done in April 2025 in PubMed, Web of Science, Scopus, and Cochrane. The inclusion criteria are randomized or non-randomized controlled trials and experimental repeated-measures studies that enrolled humans exposed to simulated microgravity for at least 24 hours. This study followed PRISMA (2020) guidelines and was registered in PROSPERO under CRD420251104161. Data was analyzed by Comprehensive Meta-Analysis software.

Results: Ten studies, involving 152 participants, met the inclusion criteria. During exposure, the first effect was a decreased plasma volume, leading to hemoconcentration. Erythropoietin concentration decreased, while parameters dependent on plasma volume, such as hemoglobin, hematocrit, and RBC count, falsely increased. Total hemoglobin mass and red cell volume showed a marked decrease, suggesting suppressed erythropoiesis. Iron status markers increased significantly, while classic hemolysis markers did not confirm increased hemolysis. Post-reambulation, erythropoietin rebounded, while the true decreases in hemoglobin, hematocrit, and RBC count were revealed as hemoconcentration resolved.

Conclusion: Simulated microgravity induces hemoconcentration, erythropoietin
suppression, and altered iron metabolism. Further investigation is needed to clarify the
role of hemolysis and to develop efficient countermeasures.

1. Introduction

Prolonged exposure to spaceflight causes multiple physiological changes in the
human body. Microgravity, exposure to radiation, physical and emotional stress,
nutritional changes, and disrupted circadian cycles affect many of the body's
functions, such as vision, the health of muscles and bones, and the immune system.
[1]. An important physiological change that occurs is fluid redistribution as a result of
microgravity. This redistribution causes various changes in different hematological
parameters, and these changes are referred to as space anemia.

The history of space anemia started in the 1960s following missions in NASA's
Gemini program when reductions in red blood cell mass were documented after
returning to the earth [2]. These early findings were the core for recognition of space
anemia as a physiological response to spaceflight.

Even though space anemia has been known about for a long time, the exact
mechanism is still not fully understood. Different hypotheses were proposed to
explain the reduction in red blood cell mass following exposure to microgravity.
Suggested mechanisms include decreased erythropoietin (EPO) production,
suppression of erythropoiesis, enhanced hemolysis, sequestration, and RBC
dysfunction, but the precise physiological basis of space anemia remains unclear and
not fully established [3,4].

As humanity prepares for longer space missions, understanding the exact
physiological responses of exposure to the space environment becomes essential, and
among these responses, space anemia is considered a key concern [5]. Deeper
understanding of this condition is vital for developing suitable countermeasures to
support the travelers during these prolonged missions.

Studying the effect of microgravity on different physiological aspects in space faced
logistical, financial, and ethical challenges, so to overcome these limitations, different
ground-based models have been developed to simulate the effects of microgravity
under controlled conditions. Among the most commonly used are head-down tilt bed
rest and dry immersion, which not only mimic the physiological effects of
microgravity but also offer a high degree of experimental control, which helps the
researchers isolate specific variables and study them with greater precision [6–8].

In addition to the great benefits in the field of space medicine, understanding the
effect of models like head-down tilt bed rest and dry immersion on the hematological
parameters can also have important clinical applications on earth. These models are
used to mimic the effects of prolonged physical inactivity, which is a very common
problem among various groups of patients, like the elderly and those with chronic

immobilization, providing us with more knowledge about hematological and systemic changes associated with extended bed rest. Improved understanding of these changes can help us develop more strategies for preventing or managing anemia and support overall health in these vulnerable groups. [9,10].

Despite the increasing number of studies about the effects of simulated microgravity on anemia-related biomarkers, the findings were inconsistent with variability in study designs, participant characteristics, and, most importantly, reported outcomes. Previous reviews discussed the available literature, but this study quantitatively synthesizes the evidence, determines the overall effect size, and assesses changes in hematological biomarkers both during and after exposure to simulated microgravity, aiming to advance current understanding of microgravity-induced anemia and support the development of both space mission countermeasures and Earth-based clinical applications.

2. Methods

This study was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines [11]. We followed the Cochrane Handbook guidelines in doing all the steps [12]. The study was registered in the International Prospective Register of Systematic Reviews (PROSPERO) under the registration number CRD420251104161.

2.1 Inclusion criteria:

Randomized or non-randomized controlled trials and experimental repeated-measures studies that enrolled human participants exposed to simulated microgravity for at least 24 hours were eligible. Included studies had to provide quantitative data on anemia-related hematological markers. Only full-text articles published in English were considered, with no restrictions on publication date, due to the limited number of studies available in this field.

2.2 Exclusion criteria:

Studies that used animals, in vitro or computational models, didn't have any microgravity exposure, were observational, case reports/series, conference abstracts, editorials, commentaries, or didn't report extractable numerical data for the outcomes we were looking for were excluded.

2.3 Information Sources and Search Strategy

Two authors conducted a comprehensive literature search in four major electronic databases: PubMed, Cochrane Library, Scopus, and Web of Science. The search was performed in April 2025. No filters were applied during the search to ensure broad coverage. Reference lists of included studies were also screened manually to identify additional eligible records. The complete search strategy is provided in the supplementary material.

2.4 Study selection:

The results of the literature search were collected in an Excel sheet and screened through the title and abstract by two independent authors. Then, the retrieved studies were further screened through full text to determine the final included studies. A third author was consulted to resolve any conflict.

2.5 Data extraction:

Data extraction was performed by two reviewers separately and revised by a third reviewer. The reviewers collected general study information (study design, type of simulated microgravity, exposure duration, sample size, participant gender, presence or absence of a control group for microgravity or for other interventions, and the nature of such interventions if applicable). Baseline participant characteristics were also extracted, including age, body mass index (BMI), height, weight, and VO_2 max.

Regarding outcomes, the reviewers extracted quantitative data for hematological markers, which are hemoglobin, hematocrit, erythropoietin (EPO), plasma volume (PV), red cell volume (RCV), red blood cell (RBC) count, reticulocytes, total hemoglobin mass (tHb mass), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), serum iron, transferrin saturation, total and indirect bilirubin, and haptoglobin.

Each outcome was extracted at two primary time points: (1) baseline during the ambulatory period and (2) during or immediately after exposure to microgravity to assess the impact of the intervention. When available, a third post-recovery time point was also extracted, which is defined as 24 hours or more after reambulation to normal gravity. When multiple values were reported for the same outcome on different days within the same time window, we selected the most representative value of the response during that phase. The same approach was applied uniformly across all included studies. All extracted data were organized and presented in tables.

2.6 Data Processing and Transformation

All outcome data were standardized to mean ± standard deviation (SD) before analysis. For the studies that reported standard errors (SE) [13,14] or 95% confidence intervals (CI) [15] instead of SDs, the SD was derived using Cochrane-recommended formulas [12]. Where results were presented separately for males and females [16,17], means and SDs were pooled using the participant-weighted method described in the Cochrane Handbook [12].

In some studies, both groups were exposed to simulated microgravity, but one group also received an additional intervention (e.g., nutrition, artificial gravity, or thigh cuffs). Since all included studies directly reported that these interventions had no significant effect on hematological outcomes, if the data weren't presented pooled [18], we pooled them into a single group. Means and standard deviations were combined using Cochrane-recommended formulas [12].

Numerical values that weren't available in text or tables were digitized from figures using WebPlotDigitizer. All hematological outcomes were converted to uniform units (e.g., hemoglobin in g/dL, serum iron in μmol/L), allowing for the use of mean differences (MD) rather than standardized mean differences (SMD), which provided more clinically interpretable effect estimates.

In the study by Dunn et al. (1984), outcome data were extracted only from the group that underwent 7 days of continuous horizontal or 6-degree head-down tilt (n = 18). The other group that completed 28 days of strict horizontal bed rest (n = 6) was excluded, as horizontal bed rest is no longer considered a valid analogue for simulated microgravity [6,7,19]. The authors reported results for the 18 participants as a single group without differentiating between horizontal and HDT positions, so we used the aggregated means and standard deviations provided for this combined group [13].

Also, the study by Dunn et al. (1984) was excluded from the quantitative synthesis of erythropoietin (EPO) outcomes because of problems with the methods used. In that study, researchers used a fetal mouse liver cell (FMLC) bioassay to check the levels of EPO, which is known to give values that are much higher and not standardized compared to newer immunoassays like ELISA or radioimmunoassay (RIA) [20]. Since there is no reliable method to calibrate the outcomes, including this study would have resulted in less accurate pooled estimates and made comparisons between studies more difficult [13].

In the study by Gunga et al. (1996), participants were studied under multiple conditions, including actual spaceflight and several ground-based microgravity analogues, such as −6° head-down tilt (HDT) and horizontal bed rest. Only the HDT data were used in this meta-analysis, as the authors reported that HDT was the analogue that most closely matched the response seen during real spaceflight [21].

In the study by Trudel et al. (2017), there was no clear report of indirect (unconjugated) bilirubin. However, there were values for both total and direct (conjugated) bilirubin. So, following standard clinical practice, indirect bilirubin was calculated by subtracting direct bilirubin from total bilirubin [14]. This method is widely used to estimate unconjugated bilirubin when it is not directly given, and it is described in major clinical chemistry references such as Tietz Textbook of Clinical Chemistry [22].

2.7 Risk of Bias Assessment

Risk of bias was evaluated by two reviewers using tools selected to match each study's design and analytic structure. Because the main goal of this review was to measure changes in hematological markers after being exposed to simulated microgravity, most studies were evaluated using the NIH Quality Assessment Tool for Before−After (Pre-Post) Studies with No Control Group [**23**]. This choice was appropriate because, in many randomized trials, both groups underwent simulated

microgravity, and the randomized co-intervention (e.g., nutritional modifications, thigh cuffs) was reported to have no measurable effect on the outcomes of interest.

One study included a non-randomized control group that did not experience simulated microgravity; it was assessed with the Joanna Briggs Institute (JBI) Checklist for Quasi-Experimental Studies [24], which is better for comparing groups without randomization.

Each item in the selected tools was evaluated as Yes, No, Cannot Determine, Not Reported, or Not Applicable, based on the extent to which the study met the requirement.

2.8 Statistical Analysis

One author has performed data analysis, and it was revised by another author. It was done by Comprehensive Meta-Analysis software, version 3.0.

Pooled effect sizes were calculated as mean differences (MD) with corresponding 95% confidence intervals (CI), using the inverse-variance method for weighting. The pre-post correlation coefficient (r), required for within-subject comparisons, was neither reported in any of the included studies nor could it be calculated from the available data. Accordingly, we assumed a value of r = 0.5, as recommended by the Cochrane Handbook for Systematic Reviews of Interventions [12]. To evaluate the validity of this assumption, sensitivity analyses were conducted using alternative r values of 0.2, 0.7, and 0.9. All of these values gave us consistent estimates, with no significant changes in direction or magnitude.

Heterogeneity was evaluated using Cochran's Q test and quantified by the I^2 statistic. When heterogeneity was present (p < 0.1 for Q or I^2 > 50%), a random-effects model was used. Without it, a fixed-effect model was used.

2.9 Reporting Bias Assessment

Assessment of reporting bias (e.g., by funnel plots or Egger's test) was not performed due to the small number of studies contributing to each outcome (range: 3 to 8). The Cochrane Handbook for Systematic Reviews of Interventions says that these methods are usually not reliable and don't have enough power when fewer than 10 studies are included in a meta-analysis [12]. For this reason, the risk of reporting bias due to missing results could not be formally assessed.

3. Results

3.1 Study Selection

The initial literature search through the previously mentioned databases has revealed 1274 articles, where 193 of them were duplicates. The number of articles identified through each database was as follows: 347 in PubMed, 19 in Cochrane, 328 in WOS, and 580 in Scopus. Title and abstract screening were done on 1081 articles, and 52

articles were retrieved, which were then screened by the full text. The authors have excluded 42 articles for the following reasons: narrative reviews (n = 5), non-human studies (n = 11), different intervention (n = 4), different outcomes (n = 7), incompatible study design (n = 5), lack of quantitative data (n = 6), insufficient duration of microgravity exposure (n = 1), mathematical modeling (n = 1), and language restrictions (n = 2).

Finally, 10 studies were eligible for inclusion in our study [13–18,21,25–27]. The study flow diagram is shown in **Fig. 1**.

3.2 Study Characteristics

Ten studies were included in this systematic review and meta-analysis, with a total of 152 participants.

Two ground-based models were used to simulate microgravity conditions. Eight studies used the 6-degree head-down tilt (HDT) model, with exposure times ranging from 4 to 60 days. Two studies used dry immersion (DI) protocols for 5 days each. Study designs included randomized clinical trials, non-randomized controlled studies, and pre-post experimental designs. Four studies included additional interventions like nutritional supplements, anti-gravity (AG) training, or thigh cuffs. The other studies did not include any additional interventions. **Table 1** displays the characteristics of the included studies. Details of the demographics of the participants are shown in **Table 2**. Reported ages ranged from the early twenties to late forties, and body mass index (BMI) values were generally within normal limits. Most participants were male, although two studies included female participants. In general, the studies were very heterogeneous in terms of methodology (design, type of intervention, and characteristics of participants), which provided a broad dataset to assess the physiological impact of simulated microgravity. Baseline values for the measured physiological outcomes are presented in **Table 3**.

3.3 Risk of Bias Assessment

Risk of bias was assessed using two tools based on study design. Nine studies were assessed by the NIH Quality Assessment Tool for Before-After (Pre-Post) Studies without Control Groups, and one study was assessed by the JBI Checklist for Quasi-Experimental Studies. Low to moderate risk of bias was found in the majority of the included studies. For the studies that were evaluated by the NIH Tool, the majority met the criteria for clearly defined research questions, valid outcome measurements, and appropriate statistical analyses. However, due to the specialized nature of this field of study, sample sizes were generally small, and several studies lacked clear details regarding blinding procedures. **Table 4** provides a detailed overview of the risk of bias evaluations for every study.

For the study that was evaluated by the JBI Tool, most criteria were met, including clarity of cause-effect relationships, similarity between groups, and appropriate

statistical methods, but the study did not include multiple pre- and post-intervention
measurements except for the erythropoietin, which might be considered a drawback.
Details of the assessment are shown in **Table 5**.

3.4 Outcomes

3.4.1 Core Anemia and Circulatory Volume Indicators

Hemoglobin

Eight studies were analyzed to assess the changes in hemoglobin (Hb) levels during
the exposure to microgravity. The meta-analysis showed a statistically significant
increase in Hb concentrations with a consistent elevation across studies, which was
indicated by mean difference (MD). The pooled value was +1.21 g/dL (95% CI: 0.86
to 1.57, p < 0.001). The random-effects model was used because of the significant
heterogeneity (I^2 = 71.85%, p = 0.001) **(Fig. 2a)**.

Five studies were analyzed to assess the changes after reambulation to normal gravity.
In contrast, the meta-analysis showed a significant reduction in Hb levels. The pooled
MD value was –0.60 g/dL (95% CI: –0.75 to –0.44, p < 0.001). As no heterogeneity
was detected (I^2 = 0%, p = 0.676), the fixed-effects model was used **(Fig. 2b)**.

Hematocrit

Eight studies were analyzed to assess the changes in hematocrit (Hct) levels during
the exposure to microgravity. The meta-analysis showed a statistically significant
increase in Hct concentrations, which was reflected by the mean difference (MD). The
pooled value was +3.10% (95% CI: 1.85 to 4.34, p < 0.001). The random-effects
model was used because of the significant heterogeneity (I^2 = 82.27%, p < 0.001)
(Fig. 3a).

After reambulation to normal gravity, Four studies were analyzed to assess the
changes in Hct levels. The meta-analysis showed a significant reduction in Hct levels.
The pooled MD value was –1.93% (95% CI: –3.44 to –0.43, p = 0.012). Because of
the significant heterogeneity (I^2 = 76.22%, p = 0.006), a random-effects model was
used **(Fig. 3b)**.

Total Hemoglobin Mass

Five studies were used to assess the changes in total hemoglobin mass (tHb mass)
during the exposure to microgravity. The meta-analysis showed a statistically
significant reduction in tHb mass, which was reflected by mean difference (MD). The
pooled value was –58.06 g (95% CI: –88.94 to –27.17, p < 0.001). The random-
effects model was used because of the moderate heterogeneity (I^2 = 51.81%, p =
0.081) **(Fig. 4a)**.

Red Cell Volume

Four studies were analyzed to assess the changes in red cell volume (RCV) during the exposure to microgravity. The meta-analysis showed a statistically significant decrease in RCV, which was demonstrated by mean difference (MD). The pooled value was –196.38 mL (95% CI: –350.91 to –41.85, p = 0.013). The random-effects model was used because of the moderate heterogeneity (I^2 = 62.22%, p = 0.047) **(Fig. 4b)**.

Plasma Volume

Five studies were used to assess the changes in plasma volume (PV) during the exposure to microgravity. The analysis showed a statistically significant reduction in PV, which was reflected by mean difference (MD). The pooled value was –533.10 mL (95% CI: –722.08 to –344.13, p < 0.001). The random-effects model was used because of the high heterogeneity (I^2 = 69.87%, p = 0.010) **(Fig. 5)**.

3.4.2 Erythropoiesis and Red Cell Indices

Erythropoietin

Six studies were analyzed to assess the changes in erythropoietin (EPO) levels during the exposure to microgravity. The meta-analysis showed a statistically significant decline in EPO concentrations, which was demonstrated by mean difference (MD). The pooled value was –1.96 mIU/mL (95% CI: –2.65 to –1.26, p < 0.001). As low heterogeneity was detected (I^2 = 22.3%, p = 0.266), the fixed-effects model was used. **(Fig. 6a)**.

After reambulation to normal gravity, Four studies were analyzed to assess the changes in EPO levels. The meta-analysis showed a significant rebound in EPO levels. The pooled MD value was +2.68 mIU/mL (95% CI: 1.57 to 3.79, p < 0.001). The fixed-effects model was used as there was low heterogeneity (I^2 = 33.39%, p = 0.212) **(Fig. 6b)**.

Reticulocyte Count

Six studies were used to assess the changes in reticulocyte percentage during the exposure to microgravity. The meta-analysis indicated no statistically significant change, which was demonstrated by mean difference (MD). The pooled value was –0.05% (95% CI: –0.15 to 0.04, p = 0.279). The random-effects model was used because of the significant heterogeneity (I^2 = 71.37%, p = 0.004) **(Fig. 7a)**.

After returning to normal gravity, three studies were analyzed to assess the changes in reticulocyte percentage. Although a slight increase was observed (MD = +0.16%, 95% CI: –0.13 to 0.44), this difference was not statistically significant (p = 0.291). A random-effects model was used as there was high heterogeneity (I^2 = 89.92%, p < 0.001) **(Fig. 7b)**.

Red Blood Cell Count

Six studies were analyzed to assess the changes in red blood cell (RBC) count during the exposure to microgravity. The meta-analysis showed a statistically significant increase in RBC levels, which was reflected by the mean difference (MD). The pooled value was $+0.41 \times 10^6/\mu L$ (95% CI: 0.24 to 0.58, p < 0.001). The random-effects model was used because of the significant heterogeneity ($I^2 = 88.16\%$, p < 0.001) **(Fig. 8a)**.

After returning to normal gravity, three studies were analyzed to assess the changes in RBC levels. The meta-analysis showed a significant decrease in RBC levels. The pooled MD value was $-0.19 \times 10^6/\mu L$ (95% CI: –0.27 to –0.11, p < 0.001). No heterogeneity was detected ($I^2 = 0\%$, p = 0.787), so a fixed-effects model was applied **(Fig. 8b)**.

3.4.3 Red Cell Morphological Indices

Mean Corpuscular Volume

Four studies were analyzed to assess the changes in mean corpuscular volume (MCV) during the exposure to microgravity. The meta-analysis indicated no statistically significant change, which was demonstrated by mean difference (MD). The pooled value was +0.13 fL (95% CI: –0.48 to 0.74, p = 0.673). The fixed-effects model was used because of the moderate heterogeneity ($I^2 = 46.24\%$, p = 0.134) **(Supplementary Fig. 1)**.

Mean Corpuscular Hemoglobin

Four studies were used to assess the changes in mean corpuscular hemoglobin (MCH) during the exposure to microgravity. The meta-analysis indicated no statistically significant change, which was evidenced by mean difference (MD). The pooled value was +0.08 pg (95% CI: –0.14 to 0.30, p = 0.482). A fixed-effects model was used because of the absence of heterogeneity ($I^2 = 0\%$, p = 0.970) **(Supplementary Fig. 2)**.

Mean Corpuscular Hemoglobin Concentration

Five studies were used to assess the changes in mean corpuscular hemoglobin concentration (MCHC) during the exposure to microgravity. The pooled mean difference was +0.25 g/dL (95% CI: –0.03 to 0.53, p = 0.084), which reflects no statistically significant change. The random-effects model was used because of the high heterogeneity ($I^2 = 77.37\%$, p = 0.001) **(Supplementary Fig. 3)**.

3.4.4 Iron Metabolism and Hemolysis-Associated Biomarkers

Serum Iron

Four studies were analyzed to assess the changes in serum iron concentrations during the exposure to microgravity. The meta-analysis showed a statistically significant increase in iron levels, which was reflected by the mean difference (MD). The pooled

value was +4.59 µmol/L (95% CI: 0.97 to 8.20, p = 0.013). The random-effects model was used because of the significant heterogeneity (I^2 = 96.2%, p < 0.001) **(Fig. 9a)**.

Transferrin Saturation

Four studies were analyzed to assess the changes in transferrin saturation during the exposure to microgravity. Meta-analysis showed a statistically significant increase, which was reflected by mean difference (MD). The pooled value was +8.97% (95% CI: 6.63 to 11.31, p < 0.001). The random-effects model was used because of the moderate heterogeneity (I^2 = 69.17%, p = 0.021) **(Fig. 9b)**.

Total Bilirubin

Five studies were used to assess the changes in total bilirubin levels during the exposure to microgravity. The meta-analysis indicated no statistically significant change, which was demonstrated by mean difference (MD). The pooled value was +0.10 µmol/L (95% CI: −1.36 to 1.56, p = 0.890). The random-effects model was used because of the moderate heterogeneity (I^2 = 56.78%, p = 0.055) **(Fig. 10a)**.

Indirect Bilirubin

Three studies were used to assess the changes in indirect (unconjugated) bilirubin levels during the exposure to microgravity. The meta-analysis demonstrated no statistically significant change, which was demonstrated by mean difference (MD). The pooled value was +0.68 µmol/L (95% CI: −0.03 to 1.40, p = 0.062), which did not reach statistical significance. No heterogeneity was detected (I^2 = 0%, p = 0.472), so a fixed-effects model was applied **(Fig. 10b)**.

Haptoglobin

Five studies were used to assess the changes in haptoglobin concentrations during the exposure to microgravity. The meta-analysis indicated no statistically significant change, which was demonstrated by mean difference (MD). The pooled value was +0.063 g/L (95% CI: −0.007 to 0.133, p = 0.077). No heterogeneity was detected (I^2 = 0%, p = 0.500), so a fixed-effects model was applied **(Supplementary Fig. 4)**.

4. Discussion

This systematic review and meta-analysis dealt with data from ten studies to try to give a comprehensive overview of the hematological effects of exposure to short- to medium-duration simulated microgravity in humans. Because of the relevance of models like head-down tilt and dry immersion to both spaceflight and prolonged physical inactivity on earth, studying the effect of these models can help us understand the physiological changes that happen and then help us develop countermeasures to overcome them, whether in space or in clinical strategies for patients on earth [28].

Analysis of various physiological biomarkers showed several effects upon exposure to microgravity, such as hemoconcentration due to plasma volume contraction, decreased erythropoiesis, iron metabolism changes, and some degree of quantitatively unsupported hemolysis based on available biomarker data. These changes appeared to occur without significant changes to the red blood cell morphology, which supports the idea that those are physiological adaptations rather than a pathological response [29].

The early redistribution of body fluids and the resulting decrease in plasma volume are the most important physiological changes that explain many of the observed results. This causes hemoconcentration, which makes the blood look more concentrated not because an actual increase in red blood cells happened but because of the decrease in the plasma volume [30–32]. This particular change occurs quickly after exposure to simulated microgravity, which greatly affects how several hematological markers should be interpreted.

That is why the increase in hemoglobin and hematocrit should not be seen as a true elevation of red cell mass but as a result of the plasma volume reduction [33,34]. The first observation that supports this interpretation is that after moving back to normal gravity, both hemoglobin and hematocrit levels drop significantly below baseline levels [35]. The second observation is that total hemoglobin mass (tHb) and red cell volume (RCV), which are independent of blood volume, both reflect real physiological loss and decline upon exposure.

The same pattern is seen with red blood cell count, which initially seems to be elevated during exposure because of its dependence on plasma volume for concentration. However, after reambulation, this increased RBC count is reversibly decreased beyond baseline levels. This further supports that early changes are hemoconcentration, not enhanced erythropoiesis [32].

The actual decrease in red blood cell volume and associated indices can be linked to the fact that erythropoietic activity was low during simulated microgravity exposure. This is best reflected by the constant reduction of erythropoietin (EPO), which is an important hormone regulating red blood cell production [36]. The reduction in EPO levels is probably central in mediating the decrease in total hemoglobin mass and red cell volume because erythropoietic drive is actively reduced.

Notably, the EPO suppression seems to be a secondary effect of reduced plasma volume. The central fluid shift caused by microgravity raises renal perfusion and oxygenation, which are well-known causes of EPO suppression in microgravity analogs. This further shows the importance of fluid dynamics in determining how the body responds to microgravity [33,37]. It also supports the idea that these changes are a coordinated physiological adaptation rather than a pathological process.

After reambulation, EPO levels rise again, likely because the plasma volume has been restored. This hormonal recovery allows other blood parameters to gradually return to

normal over time. This supports the idea that spaceflight-associated anemia is
predominantly a self-resolving condition recovering gradually after landing, assuming
no permanent organ damage has developed [4].

Because of erythropoietin suppression and general downregulation of erythropoiesis,
there are no significant changes in reticulocyte percentage during or post exposure to
simulated microgravity. This is consistent with the assumption that there is a
hypoproliferative state where there is diminished bone marrow output [38–41].
Moreover, the short follow-up period after reambulation in most cases probably
limited the ability to observe any delayed reticulocyte rebound that may occur
following the EPO recovery post-reambulation.

This suppressed erythropoietic response also makes interpreting potential hemolysis
more difficult. Physiologically, reticulocyte production usually increases in response
to hemolysis [42], but in our condition that response may be muted or masked due to
the existing hypoproliferative state. Therefore, not observing reticulocytosis cannot be
taken as conclusive proof that there is no hemolysis, which represents a major
limitation concerning the interpretation of hematological dynamics under suppressed
erythropoiesis.

In all studies evaluating mean corpuscular volume (MCV), mean corpuscular
hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC), no
significant changes were noted during exposure to simulated microgravity. These
indices reflect the dimensions and hemoglobin content of red blood cells [43] and,
given their constancy, suggest that erythropoiesis, although decreased in quantity, still
maintained some structural order, as there were no macrocytic nor microcytic changes
observed.

Simulated microgravity exposure was associated with an increase in serum iron
concentration, alongside a rise in transferrin saturation. Collectively, these results
indicate that there is a measurable elevation in the circulating iron levels [44].
Increased circulating iron levels may result from microgravity's hypoproliferative
effects slowing the utilization of iron for erythropoiesis. It may also be due to the
release of iron from red blood cells as a consequence of hemolysis. Likely, both
processes, with different contributions, led to the changes observed.

In assessing the possible occurrence of hemolysis, we planned to assess endogenous
carbon monoxide concentration since it is a more modern and sensitive biomarker for
the breakdown of red blood cells [45], but due to the limited number of studies
addressing this outcome, it was not possible to conduct a meta-analysis. For this
reason, we relied on more traditional but widely available markers such as total
bilirubin as well as indirect (unconjugated) bilirubin. The pooled analysis showed no
statistically significant changes in either marker with exposure to simulated
microgravity. Although these findings do not provide strong evidence to support
hemolysis, they do not provide sufficient evidence to confidently rule it out either.

The analysis of haptoglobin, another marker of hemolysis [46], also showed no significant change during simulated microgravity exposure. This adds more complexity in interpreting hemolytic activity in this situation, especially in the consideration of whether any hemolysis occurring is primarily intravascular or extravascular in nature. In addition, it should be noted that haptoglobin is an acute phase reactant, which means it is liable to changes in inflammatory conditions like those that may be encountered in spaceflight [47,48]. This form of systemic inflammation could increase haptoglobin levels, masking the decreases associated with hemolysis. These aspects limit its reliability as a hemolysis marker in this context.

Several studies that include human subjects exposed to microgravity during spaceflights have reported results that align with the current analysis, like the reductions in hemoglobin concentration, hematocrit levels, and red blood cell mass (RBCM), which have all been documented and are consistent with the results of our study [39,49,50]. The agreement between spaceflight data and ground-based analogues increases the confidence in the observed patterns and their applicability in different conditions.

Several studies conducted during spaceflight missions have described an early and persistent suppression of erythropoietin (EPO) during microgravity exposure [39,51]. This initial decline indicates disruption of the erythropoietic response in advance of any detectable decreases in red blood cell mass. The reduction in EPO concentration has been demonstrated to inversely correlate with central venous pressure, which is a relationship seen in residents of high altitude too, indicating a shared physiological mechanism [33,34,37]. The severity of spaceflight-associated anemia and recovery time are linked to mission duration, showing the impact of extended erythropoietic suppression, as indicated by the available epidemiological data [49,50].

New papers of less than five years have supported the idea that hemolysis occurs during exposure to microgravity, and they are supported by using recently developed methods to precisely quantify carbon monoxide (CO) elimination as an indicator. One study on actual microgravity showed sustained increases in alveolar CO elimination over the duration of a six-month space mission in 14 astronauts [4]. In addition, another study based on ground simulation of microgravity using head-down tilt bed rest also reported increases in CO elimination and bilirubin levels [15], which support the assumption of the occurrence of hemolysis in microgravity, whether in spaceflights or in the simulation models on earth. In contrast, a study which used 21 days of head-down tilt (HDT) bed rest used the same indicator showed no significant increase in CO levels or other hemolytic markers challenging the assumption that hemolysis is a mechanism for red cell loss under simulated microgravity conditions [14].

These contradictory results show the need for more studies to evaluate the nature and magnitude of hemolysis during exposure to microgravity. The use of sensitive and

non-invasive indicators as CO elimination should be expanded to detect minor changes in hemolysis that might be missed by traditional hematological parameters [52,53].

It is also important to discuss that the lack of significant changes in the MCV, MCH, and MCHC does not mean that there are no changes in the structure of the red blood cell or its function. Microgravity has been linked with a number of changes in the cytosol of erythrocytes, such as diminished total antioxidant capacity, increased levels of reactive oxygen species (ROS), and lower glutathione concentration, which threaten cell integrity. Moreover, elevated concentrations of phosphatidylcholine and nitric oxide have been reported, which may impair RBC-mediated vasoregulation as well as ATP release and may also contribute to alterations in membrane deformability [54]. Research conducted using blood samples from spaceflights indicates that an increase in phosphatidylcholine levels may lead to increased rigidity of the membrane of red blood cells, which is known to limit the functional flexibility of the cells [55,56].

These findings show that space medicine still has much potential to advance, as every physiological change caused by microgravity leads to important clinical effects. Even plasma volume loss, which could be underestimated, has been shown to cause orthostatic intolerance (OI) in astronauts, especially after short-duration spaceflights [57,58]. This shows the need for the development of integrated solutions that should ensure fluid replacement, erythropoiesis stimulation, and erythrocyte destruction prevention. Enhancing these countermeasures may contribute not only to the safety of long-duration space missions but also help in the development of treatments for similar conditions on Earth.

This study has multiple strengths, such as the systematic method used to collect and analyze data from several microgravity analog studies. It also provided a detailed and comprehensive overview of the hematological response to simulated microgravity. It included appropriate consideration of multiple outcomes across different domains like blood volume status, erythropoietic activity, red cell morphology, iron metabolism, and hemolysis. This broad coverage improves the explanation of the results and enables a more coherent understanding of space anemia.

Also, there are some limitations that need to be acknowledged. The total number of studies that were included in our analysis was small because of the limited number of studies in this field. Also, the sample size in each study was relatively small, which might decrease their statistical power. In addition, the follow-up periods after reambulation were relatively short for many studies, which limited the ability to give a broader picture for some outcomes.

As for our recommendations for future studies, further research is required to better evaluate the nature and extent of hemolysis during exposure to microgravity. At the same time, there is a need for well-designed studies to develop specific

countermeasures intended to maintain red blood cell mass and function. Such
measures might include protocols for artificial gravity, exercise programs,
pharmacological interventions, or any other methods designed to help the human body
adjust to this extreme environment.

Conclusion
This research provides a comprehensive overview of the hematological changes
during simulated microgravity exposure. The findings support the existence of
hemoconcentration, erythropoietin suppression, and altered iron metabolism. Many of
these changes may be considered a coordinated physiological adaptation rather than a
pathological process. However, the possible contributions of low-grade hemolysis
require further investigation. Further investigation is also necessary to understand
these mechanisms and develop strategies that can aid humans during spaceflights and
in relevant clinical situations on Earth.

Statements and Declarations

Conflict of interest

The authors have no relevant financial or non-financial interests to disclose.

Funding

No funding was received for conducting this study.

Author contributions

600   **Availability of data and material**

601   All data generated or analyzed during this study are included in this published article
602   and its supplementary information files.

603   **Acknowledgments**

References

References

1.   Williams D, Kuipers A, Mukai C, Thirsk R. Acclimation during space flight: effects on
human physiology. Can Med Assoc J. 2009;180: 1317–1323.
doi:10.1503/cmaj.090628

2.   Fischer CL, Johnson PC, Berry CA. Red blood cell mass and plasma volume changes in
manned space flight. JAMA. 1967;200: 579–83.

3.   Alfrey CP, Rice L, Udden MM, Driscoll TB. Neocytolysis: physiological down-regulator
of red-cell mass. The Lancet. 1997;349: 1389–1390. doi:10.1016/S0140-
6736(96)09208-2

4.   Trudel G, Shahin N, Ramsay T, Laneuville O, Louati H. Hemolysis contributes to
anemia during long-duration space flight. Nat Med. 2022;28: 59–62.
doi:10.1038/s41591-021-01637-7

5.   Lansiaux E, Jain N, Yatin Chodnekar S, Siddiq A, Ibrahim M, Yèche M, et al.
Understanding the complexities of space anaemia in extended space missions:
revelations from microgravitational odyssey. Front Physiol. 2024;15.
doi:10.3389/fphys.2024.1321468

6.   Watenpaugh DE. Analogs of microgravity: head-down tilt and water immersion. J
Appl Physiol. 2016;120: 904–914. doi:10.1152/japplphysiol.00986.2015

7.   Hargens AR, Vico L. Long-duration bed rest as an analog to microgravity. J Appl
Physiol. 2016;120: 891–903. doi:10.1152/japplphysiol.00935.2015

8.   Tomilovskaya E, Shigueva T, Sayenko D, Rukavishnikov I, Kozlovskaya I. Dry Immersion
as a Ground-Based Model of Microgravity Physiological Effects. Front Physiol.
2019;10. doi:10.3389/fphys.2019.00284

9.   Rizzo AM, Corsetto PA, Montorfano G, Milani S, Zava S, Tavella S, et al. Effects of
Long-Term Space Flight on Erythrocytes and Oxidative Stress of Rodents. PLoS One.
2012;7: e32361. doi:10.1371/journal.pone.0032361

10.   Strollo F, Gentile S, Strollo G, Mambro A, Vernikos J. Recent Progress in Space
Physiology and Aging. Front Physiol. 2018;9. doi:10.3389/fphys.2018.01551

11.   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al.
Updating guidance for reporting systematic reviews: development of the PRISMA
2020 statement. J Clin Epidemiol. 2021;134: 103–112.
doi:10.1016/j.jclinepi.2021.02.003

12.   Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors.
Cochrane Handbook for Systematic Reviews of Interventions. Wiley; 2019.
doi:10.1002/9781119536604

13.   Dunn CDR, Lange RD, Kimzey SL, Johnson PC, Leach CS. Serum erythropoietin titers
644   during prolonged bedrest; relevance to the ?anaemia? of space flight. Eur J Appl
645   Physiol Occup Physiol. 1984;52: 178–182. doi:10.1007/BF00433389

646   14.   Trudel G, Uhthoff HK, Laneuville O. Hemolysis during and after 21 days of head-
647   down-tilt bed rest. Physiol Rep. 2017;5: e13469. doi:10.14814/phy2.13469

648   15.   Culliton K, Louati H, Laneuville O, Ramsay T, Trudel G. Six degrees head-down tilt bed
649   rest caused low-grade hemolysis: a prospective randomized clinical trial. NPJ
650   Microgravity. 2021;7: 4. doi:10.1038/s41526-021-00132-0

651   16.   Horeau M, Ropert M, Mulder E, Tank J, Frings-Meuthen P, Armbrecht G, et al. Iron
652   metabolism regulation in females and males exposed to simulated microgravity:
653   results from the randomized trial Artificial Gravity Bed Rest—European Space Agency
654   (AGBRESA). Am J Clin Nutr. 2022;116: 1430–1440. doi:10.1093/ajcn/nqac205

655   17.   Horeau M, Navasiolava N, Van Ombergen A, Custaud M-A, Robin A, Ropert M, et al.
656   Dry immersion rapidly disturbs iron metabolism in men and women: results from the
657   VIVALDI studies. NPJ Microgravity. 2024;10: 68. doi:10.1038/s41526-024-00399-z

658   18.   Nay K, Koechlin-Ramonatxo C, Rochdi S, Island M, Orfila L, Treffel L, et al. Simulated
659   microgravity disturbs iron metabolism and distribution in humans: Lessons from dry
660   immersion, an innovative ground-based human model. The FASEB Journal. 2020;34:
661   14920–14929. doi:10.1096/fj.202001199RR

662   19.   Pavy-Le Traon A, Heer M, Narici M V., Rittweger J, Vernikos J. From space to Earth:
663   advances in human physiology from 20 years of bed rest studies (1986–2006). Eur J
664   Appl Physiol. 2007;101: 143–194. doi:10.1007/s00421-007-0474-z

665   20.   Sakata S, Nakatani A, Jimaru D, Ueda M, Kohzuki H, Ohga Y, et al. Measurement of
666   Mouse Plasma Erythropoietin by an Improved ELISA. Comparative Haematology
667   International. 1999;9: 110–116. doi:10.1007/s005800050019

668   21.   Gunga HC, Kirsch K, Baartz F, Maillet A, Gharib C, Nalishiti W, et al. Erythropoietin
669   under real and simulated microgravity conditions in humans. J Appl Physiol. 1996;81:
670   761–773. doi:10.1152/jappl.1996.81.2.761

671   22.   Jung K. Tietz Fundamentals of Clinical Chemistry, 6th edition. Carl A. Burtis, Edward R.
672   Ashwood, and David E. Bruns, editors. St Louis, MO: Saunders/Elsevier, 2008, 976 pp,
673           $96.95. ISBN 978-0-7216-3865-2. Clin Chem. 2008;54: 1933–1933.
674   doi:10.1373/clinchem.2007.101378

675   23.   National Heart Lung, Institute B. Study quality assessment tools. 2014.

676   24.   Joanna Briggs Institute. Checklist for quasi-experimental studies (non-randomized
677   experimental studies). 2020.

678   25.   Ryan BJ, Goodrich JA, Schmidt WF, Stothard ER, Wright KP, Byrnes WC. Haemoglobin
679   mass alterations in healthy humans following four-day head-down tilt bed rest. Exp
680   Physiol. 2016;101: 628–640. doi:10.1113/EP085665

681   26.   Lampe L, Wienhold K, Meyer G, Baisch F, Maass H, Hollmann W, et al. Effects of
682   simulated microgravity (HDT) on blood fluidity. J Appl Physiol. 1992;73: 1366–1369.
683   doi:10.1152/jappl.1992.73.4.1366

684   27.   Branch JD, Pate RR, Bodary PF, Convertino VA. Red cell volume and [erythropoietin]
685   responses during exposure to simulated microgravity. Aviat Space Environ Med.
686   1998;69: 347–51.

687   28.   Hargens AR, Bhattacharya R, Schneider SM. Space physiology VI: exercise, artificial
688   gravity, and countermeasure development for prolonged space flight. Eur J Appl
689   Physiol. 2013;113: 2183–2192. doi:10.1007/s00421-012-2523-5

690   29.   Smith SM. Red blood cell and iron metabolism during space flight. Nutrition. 2002;18:
691   864–866. doi:10.1016/S0899-9007(02)00912-7

692   30.   Otto JM, Plumb JOM, Clissold E, Kumar SB, Wakeham DJ, Schmidt W, et al.
693   Hemoglobin concentration, total hemoglobin mass and plasma volume in patients:
694   implications for anemia. Haematologica. 2017;102: 1477–1485.
695   doi:10.3324/haematol.2017.169680

696   31.   Bilancio G, Lombardi C, Pisot R, De Santo NG, Cavallo P, Cirillo M. Effects of Bed-Rest
697   on Urea and Creatinine: Correlation with Changes in Fat-Free Mass. PLoS One.
698   2014;9: e108805. doi:10.1371/journal.pone.0108805

699   32.   Leach CS, Alfrey CP, Suki WN, Leonard JI, Rambaut PC, Inners LD, et al. Regulation of
700   body fluid compartments during short-term spaceflight. J Appl Physiol. 1996;81: 105–
701   116. doi:10.1152/jappl.1996.81.1.105

702   33.   Garrett-Bakelman FE, Darshi M, Green SJ, Gur RC, Lin L, Macias BR, et al. The NASA
703   Twins Study: A multidimensional analysis of a year-long human spaceflight. Science
704   (1979). 2019;364. doi:10.1126/science.aau8650

705   34.   Leach CS, Johnson PC. Influence of Spaceflight on Erythrokinetics in Man. Science
706   (1979). 1984;225: 216–218. doi:10.1126/science.6729477

707   35.   Smith SM, Davis-Street JE, Fontenot TB, Lane HW. Assessment of a portable clinical
708   blood analyzer during space flight. Clin Chem. 1997;43: 1056–65.

709   36.   Jelkmann W. Regulation of erythropoietin production. J Physiol. 2011;589: 1251–
710   1258. doi:10.1113/jphysiol.2010.195057

711   37.   De Santo NG, Cirillo M, Kirsch KA, Correale G, Drummer C, Frassl W, et al. Anemia and
712   Erythropoietin in Space Flights. Semin Nephrol. 2005;25: 379–387.
713   doi:10.1016/j.semnephrol.2005.05.006

714   38.   Blaber EA, Finkelstein H, Dvorochkin N, Sato KY, Yousuf R, Burns BP, et al.
715   Microgravity Reduces the Differentiation and Regenerative Potential of Embryonic
716   Stem Cells. Stem Cells Dev. 2015;24: 2605–2621. doi:10.1089/scd.2015.0218

717   39.   Udden MM, Driscoll TB, Pickett MH, Leach-Huntoon CS, Alfrey CP. Decreased
718   production of red blood cells in human subjects exposed to microgravity. J Lab Clin
719   Med. 1995;125: 442–9.

720   40.   Alfrey CP, Udden MM, Leach-Huntoon C, Driscoll T, Pickett MH. Control of red blood
721   cell mass in spaceflight. J Appl Physiol. 1996;81: 98–104.
722   doi:10.1152/jappl.1996.81.1.98

723   41.   Tavassoli M. Anemia of spaceflight. Blood. 1982;60: 1059–67.

724   42.   Rai D, Wilson AM, Moosavi L. Histology, Reticulocytes. StatPearls [Internet]. Treasure
725   Island (FL): StatPearls Publishing; 2025. Available:
726   https://www.ncbi.nlm.nih.gov/books/NBK542172/

727   43.   Walker HK, Hall WD, Hurst JW, editors. Clinical Methods: The History, Physical, and
728   Laboratory Examinations. 3rd ed. Boston: Butterworths; 1990. Available:
729   https://www.ncbi.nlm.nih.gov/books/NBK201/

730   44.   Zwart SR, Morgan JL, Smith SM. Iron status and its relations with oxidative damage
731   and bone loss during long-duration space flight on the International Space Station.
732   Am J Clin Nutr. 2013;98: 217–223. doi:10.3945/ajcn.112.056465

733   45.   Shahin N, Louati H, Trudel G. Measuring Human Hemolysis Clinically and in Extreme
734   Environments Using Endogenous Carbon Monoxide Elimination. Ann Biomed Eng.
735   2020;48: 1540–1550. doi:10.1007/s10439-020-02473-5

736   46.   Alayash AI, Andersen CBF, Moestrup SK, Bülow L. Haptoglobin: the hemoglobin
737   detoxifier in plasma. Trends Biotechnol. 2013;31: 2–3.
738   doi:10.1016/j.tibtech.2012.10.003

739   47.   Tocci D, Ducai T, Stoute CAB, Hopkins G, Sabbir MG, Beheshti A, et al. "Monitoring
740   inflammatory, immune system mediators, and mitochondrial changes related to brain
741   metabolism during space flight." Front Immunol. 2024;15.
742   doi:10.3389/fimmu.2024.1422864

743   48.   Davis CM, Allen AR, Bowles DE. Consequences of space radiation on the brain and
744   cardiovascular system. Journal of Environmental Science and Health, Part C. 2021;39:
745   180–218. doi:10.1080/26896583.2021.1891825

746   49.   Kunz H, Quiriarte H, Simpson RJ, Ploutz-Snyder R, McMonigal K, Sams C, et al.
747   Alterations in hematologic indices during long-duration spaceflight. BMC Hematol.
748   2017;17: 12. doi:10.1186/s12878-017-0083-y

50.   Trudel G, Shafer J, Laneuville O, Ramsay T. Characterizing the effect of exposure to
microgravity on anemia: more space is worse. Am J Hematol. 2020;95: 267–273.
doi:10.1002/ajh.25699

51.   Leach CS, Johnson PC, Cintron NM. The endocrine system in space flight. Acta
Astronaut. 1988;17: 161–166. doi:10.1016/0094-5765(88)90017-3

52.   Levitt D, Levitt M. Carbon monoxide: a critical quantitative analysis and review of the
extent and limitations of its second messenger function. Clin Pharmacol. 2015; 37.
doi:10.2147/CPAA.S79626

53.   Coburn RF. The measurement of endogenous carbon monoxide production. J Appl
Physiol. 2012;112: 1949–1955. doi:10.1152/japplphysiol.00174.2012

54.   Bennett-Guerrero E, Veldman TH, Doctor A, Telen MJ, Ortel TL, Reid TS, et al.
Evolution of adverse changes in stored RBCs. Proceedings of the National Academy of
Sciences. 2007;104: 17063–17068. doi:10.1073/pnas.0708160104

55.   Boonstra J. Growth factor-induced signal transduction in adherent mammalian cells is
sensitive to gravity. The FASEB Journal. 1999;13. doi:10.1096/fasebj.13.9001.s35

56.   Ivanova SM, Morukov B V, Labetskaia OI, Iarlykova I V, Levina AA, Kozinets GI.
[Morphobiochemical assay of the red blood system in members of the prime crews of
the International Space Station]. Aviakosm Ekolog Med. 2006;40: 9–15.

57.   Fritsch-Yelle JM, Whitson PA, Bondar RL, Brown TE. Subnormal norepinephrine
release relates to presyncope in astronauts after spaceflight. J Appl Physiol. 1996;81:
2134–2141. doi:10.1152/jappl.1996.81.5.2134

58.   Buckey JC, Lane LD, Levine BD, Watenpaugh DE, Wright SJ, Moore WE, et al.
Orthostatic intolerance after spaceflight. J Appl Physiol. 1996;81: 7–18.
doi:10.1152/jappl.1996.81.1.7

Figure captions

Fig. 1 PRISMA 2020 flow diagram illustrating the study selection process

Fig. 2 Forest plots of changes in Hemoglobin concentrations (a) Hemoglobin during
exposure to simulated microgravity (b) Hemoglobin after reambulation

Fig. 3 Forest plots of changes in Hematocrit concentrations (a) Hematocrit during
exposure to simulated microgravity (b) Hematocrit after reambulation

Fig. 4 Forest plot of changes in (a) total hemoglobin mass and (b) red cell volume
during simulated microgravity exposure

Fig. 5 Forest plot of changes in plasma volume during simulated microgravity
exposure

Fig. 6 Forest plots of changes in erythropoietin concentration (a) during simulated
microgravity exposure and (b) after reambulation

Fig. 7 Forest plots of changes in reticulocyte percentage (a) during simulated
microgravity exposure and (b) after reambulation

Fig. 8 Forest plots of changes in red blood cell count (a) during simulated
microgravity exposure and (b) after reambulation

Fig. 9 Forest plots of changes in (a) serum iron and (b) transferrin saturation during
simulated microgravity exposure

Fig. 10 Forest plots of changes in (a) total bilirubin concentration, (b) indirect
bilirubin concentration during simulated microgravity exposure

Tables:

Table 1. Characteristics of included studies.

Study Name	Study Design	Microgravity Type	Exposure Duration	Sample Size (n)	Participant Gender	Microgravity Control Group Present?	Other Intervention Control Group Present?	Intervention Type
Branch1998	Non-randomized Controlled Experimental Study	6-degree head-down tilt (HDT)	16 days	6	6 males	Yes	No	
Culliton2021	Randomized Clinical Trial	6-degree head-down tilt (HDT)	60 days	20	20 males	No	Yes	Nutrition
dunn1984	Experimental Study - Pre-post Design	Continuous horizontal or 6-degree head-down tilt (HDT)	7 days	18	18 males	No	No	
gunga1996	Experimental Study - Pre-post Design	6-degree head-down tilt (HDT)	42 days	8	8 males	No	No	
Horeau2022	Randomized Clinical Trial	6-degree head-down tilt (HDT)	60 days	23	7 females, 16 males	No	Yes	AG training
Horeau2024	Experimental Study - Pre-post Design	Dry Immersion (DI)	5 days	37	18 females, 19 males	No	No	
lampe1992	Experimental Study - Pre-post Design	6-degree head-down tilt (HDT)	10 days	6	6 males	No	No	
Nay 2020	Randomized Clinical Trial	Dry Immersion (DI)	5 days	17	18 males	No	Yes	Thigh cuffs
ryan2016	Experimental Study - Pre-post Design	6-degree head-down tilt (HDT)	4 days	7	7 males	No	No	
trudel2017	Cross-over Experimental Study	6-degree head-down tilt (HDT)	21 days	10	10 males	No	Yes	Nutrition

Table 2. Baseline demographic characteristics of participants in the included studies.

Study Name	Age (years)	BMI (kg/m²)	Height (cm)	Weight (kg)	VO₂ Max (ml/min/kg)
Branch1998	39 ± 7		182.0 ± 6.5	81.7 ± 12.0	32.7 ± 8.1
Culliton2021	34.2 ± 7.8	22–27			
dunn1984				68.8 ± 2.12	
gunga1996	27.9 ± 2.3		175 ± 4	74.5 ± 8.3	
Horeau2022	33.27 ± 9.30	24.30 ± 1.99	174.03 ± 5.91	74.57 ± 7.83	
Horeau2024	28.38 ± 4.51	22.46 ± 1.83	170.83 ± 5.06	65.82 ± 6.49	
lampe1992	26				42.5
Nay 2020	33.6 ± 5.5	23.5 ± 2.5	178 ± 6	74.4 ± 8.0	46.1 ± 6.9
ryan2016	21 ± 3		180 ± 11	73 ± 10	50 ± 6
trudel2017	32	23.2			

Age, body mass index (BMI), height, weight, and VO₂ max are reported as mean ± standard deviation.

Table 3. Baseline values for the measured physiological outcomes across included studies.

Study Name	Hemoglobin (g/dL)	Hematocrit (%)	Erythropoietin (mIU/ml)	Plasma Volume (ml)	RCV (ml)	RBC Count (×10⁶/μL)	Reticulocytes (%)	Total Hemoglobin Mass (g)	MCV (fL)	MCH (pg)	MCHC (g/dL)	Iron (μmol/L)	Transferrin Saturation (%)	Bilirubin (μmol/L)	Indirect Bilirubin (μmol/L)	Haptoglobin (g/L)
Branch1998	14.5 ± 0.9	43.0 ± 3.4	11.6 ± 2.9	3769 ± 512	2845 ± 410											
Culliton2021	13.7 ± 0.64		12.2 ± 4.3			4.529 ± 0.29	1.05 ± 0.39	818.2 ± 104.6				14.57 ± 3.47	29.0 ± 9.1	11.5 ± 4.8		1.03 ± 0.29
dunn1984		41.4 ± 2.12	120 ± 84.9				0.60 ± 0.3									
gunga1996	14.2 ± 1.4	44.2 ± 2.2	11.4 ± 3.3													
Horeau2022	14.96 ± 1.01	39.75 ± 4.82		3030.61 ± 480.49	2382.35 ± 374.77	4.69 ± 0.35	0.95 ± 0.19	813.59 ± 122.45	86.39 ± 4.38	29.7 ± 1.66	34.39 ± 0.64	16.35 ± 4.05	26.33 ± 6.45	11.74 ± 6.53	11.40 ± 6.61	1.01 ± 0.40
Horeau2024				3275.76 ± 571.05		4.64 ± 0.29	1.02 ± 0.35	710.46 ± 103.14	91.49 ± 3.52	31.11 ± 1.1	34.11 ± 0.40	18.08 ± 1.82	30.49 ± 2.96			
lampe1992	14.5 ± 1.10	42.3 ± 3.46				4.73 ± 0.39			90.5 ± 2.06	31.4 ± 1.57	34.6 ± 1.07					
Nay 2020	14.55 ± 0.88	41.96 ± 2.33	7.70 ± 2.70			4.85 ± 0.37	1.15 ± 0.20					19.06 ± 2.74	32.69 ± 4.90	8.00 ± 2.27	4.65 ± 1.57	0.85 ± 0.23
ryan2016	15.7 ± 0.4	45.3 ± 1.8	7.1 ± 1.8	3369 ± 569	2356 ± 371		1.33 ± 0.28	817 ± 135			34.7 ± 1.0			8.55 ± 4.1		0.54 ± 0.32
trudel2017	14.0 ± 0.32	41.1 ± 1.3	10.8 ± 2.85	4004 ± 351	2459 ± 196.1	4.83 ± 0.16	1.10 ± 0.14	838 ± 66.4	83.9 ± 1.3	28.9 ± 0.3	34.1 ± 0.3			14.5 ± 4.3	8.9 ± 4.55	1.11 ± 0.44

Values are presented as mean ± standard deviation

Table 4. Risk of bias assessment using NIH Tool for Before-After (Pre-Post) Studies with No Control Group

Study	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12
Culliton et al., 2021	Y	Y	Y	NR	Y	Y	Y	NR	Y	Y	Y	NA
Dunn et al., 1984	Y	N	CD	NR	Y	Y	Y	NR	Y	Y	Y	NA
Gunga et al., 1996	Y	Y	Y	NR	CD	Y	Y	NR	Y	Y	Y	NA
Horeau et al., 2022	Y	Y	Y	NR	Y	Y	Y	NR	Y	Y	Y	NA
Horeau et al., 2024	Y	Y	Y	NR	Y	Y	Y	NR	Y	Y	N	NA
Lampe et al., 1992	Y	Y	CD	NR	CD	Y	Y	NR	Y	Y	Y	NA
Nay et al., 2020	Y	Y	Y	NR	Y	Y	Y	NR	Y	Y	N	NA
Ryan et al., 2016	Y	Y	Y	NR	CD	Y	Y	NR	Y	Y	Y	NA
Trudel et al., 2017	Y	Y	Y	NR	CD	Y	Y	NR	Y	Y	Y	NA

*Y, yes; N, No; CD, cannot determine; NA, not applicable; NR, not reported

Table 5. Risk of bias assessment using JBI Checklist for Quasi-Experiments for the study Branch et al., 1998

JBI Checklist Item	Assessment
1. Is it clear what is the 'cause' and what is the 'effect'?	Yes
2. Were the participants in any comparisons similar?	Yes
3. Were the participants receiving similar care other than the exposure?	Yes
4. Was there a control group?	Yes
5. Were there multiple measurements of the outcome pre/post intervention?	No
6. Was follow-up complete and adequately described?	Yes
7. Were outcomes measured the same way in both groups?	Yes
8. Were outcomes measured in a reliable way?	Yes
9. Was appropriate statistical analysis used?	Yes

Figure

a

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Branch1998	0.800	0.534	0.285	-0.246	1.846	1.499	0.134
Culliton2021	1.300	0.189	0.036	0.929	1.671	6.867	0.000
Gunga1996	1.600	0.434	0.189	0.748	2.452	3.683	0.000
Horeau2022	0.420	0.209	0.043	0.011	0.829	2.014	0.044
Lampe1992	1.200	0.449	0.202	0.320	2.080	2.672	0.008
Nay2020	1.720	0.228	0.052	1.273	2.167	7.546	0.000
Ryan2016	1.700	0.295	0.087	1.121	2.279	5.759	0.000
Trudel2017	1.000	0.173	0.030	0.662	1.338	5.796	0.000
	1.212	0.180	0.033	0.858	1.565	6.717	0.000

Random-effects model used. Heterogeneity: I² = 71.85%, p = 0.001

b

	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Culliton2021	-0.600	0.171	0.029	-0.936	-0.264	-3.499	0.000
Gunga1996	-1.000	0.430	0.185	-1.843	-0.157	-2.325	0.020
Lampe1992	-0.100	0.413	0.171	-0.909	0.709	-0.242	0.809
Ryan2016	-0.600	0.262	0.069	-1.113	-0.087	-2.291	0.022
Trudel2017	-0.600	0.101	0.010	-0.798	-0.402	-5.929	0.000
	-0.595	0.080	0.006	-0.751	-0.439	-7.469	0.000

Fixed-effects model used. No heterogeneity was detected across studies (I² = 0%, p = 0.676)

Figure

a

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Branch1998	0.900	1.283	1.647	-1.615	3.415	0.701	0.483
Dunn1984	1.400	0.458	0.210	0.502	2.298	3.055	0.002
Gunga1996	3.200	0.796	0.634	1.640	4.760	4.020	0.000
Horeau2022	0.580	1.067	1.139	-1.511	2.671	0.544	0.587
Lampe1992	5.400	1.299	1.687	2.854	7.946	4.157	0.000
Nay2020	5.260	0.687	0.471	3.914	6.606	7.662	0.000
Ryan2016	5.100	0.900	0.810	3.336	6.864	5.667	0.000
Trudel2017	2.900	0.466	0.217	1.987	3.813	6.225	0.000
	3.095	0.637	0.406	1.847	4.344	4.859	0.000

Random-effects model used. Heterogeneity: $I^2 = 82.27\%$, $p < 0.001$

b

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Gunga1996	-3.900	0.696	0.485	-5.265	-2.535	-5.600	0.000
Lampe1992	0.900	1.254	1.571	-1.557	3.357	0.718	0.473
Ryan2016	-1.900	0.818	0.669	-3.503	-0.297	-2.324	0.020
Trudel2017	-1.900	0.466	0.217	-2.813	-0.987	-4.079	0.000
	-1.934	0.768	0.590	-3.439	-0.429	-2.518	0.012

Random-effects model used. Heterogeneity: $I^2 = 76.22\%$, $p = 0.006$

Figure

a

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Culliton2021	-83.200	27.241	742.049	-136.591	-29.809	-3.054	0.002
Horeau2022	-42.670	24.822	616.117	-91.320	5.980	-1.719	0.086
Horeau2024	-44.700	15.739	247.708	-75.547	-13.853	-2.840	0.005
Ryan2016	32.000	52.196	2724.429	-70.302	134.302	0.613	0.540
Trudel2017	-95.000	20.076	403.033	-134.348	-55.652	-4.732	0.000
	-58.058	15.757	248.290	-88.942	-27.174	-3.685	0.000

Random-effects model used. Heterogeneity: $I^2 = 51.81\%$, $p = 0.081$

b

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Branch1998	-354.000	151.396	22920.667	-650.730	-57.270	-2.338	0.019
Horeau2022	-154.310	72.254	5220.703	-295.926	-12.694	-2.136	0.033
Ryan2016	99.000	150.790	22737.571	-196.543	394.543	0.657	0.511
Trudel2017	-302.000	60.099	3611.939	-419.793	-184.207	-5.025	0.000
	-196.380	78.845	6216.543	-350.914	-41.847	-2.491	0.013

Random-effects model used. Heterogeneity: $I^2 = 62.22\%$, $p = 0.047$

Figure

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Branch1998	-580.000	181.718	33021.500	-936.161	-223.839	-3.192	0.001
Horeau2022	-314.350	87.080	7582.860	-485.023	-143.677	-3.610	0.000
Horeau2024	-756.170	85.547	7318.300	-923.839	-588.501	-8.839	0.000
Ryan2016	-464.000	211.742	44834.714	-879.007	-48.993	-2.191	0.028
Trudel2017	-534.000	99.714	9942.929	-729.436	-338.564	-5.355	0.000
	-533.102	96.417	9296.252	-722.076	-344.128	-5.529	0.000

Random-effects model used. Heterogeneity: I² = 69.87%, p = 0.010

Figure

a

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Branch1998	-0.400	1.128	1.272	-2.610	1.810	-0.355	0.723	
Culliton2021	-3.300	0.839	0.704	-4.944	-1.656	-3.934	0.000	
Gunga1996	-2.300	1.504	2.261	-5.247	0.647	-1.530	0.126	
Nay2020	-2.250	0.606	0.367	-3.438	-1.062	-3.713	0.000	
Ryan2016	-1.800	0.844	0.713	-3.455	-0.145	-2.132	0.033	
Trudel2017	-0.800	0.901	0.812	-2.566	0.966	-0.888	0.375	
	-1.956	0.353	0.125	-2.648	-1.263	-5.534	0.000	

Lower EPO   Higher EPO

Fixed-effects model used. Heterogeneity: $I^2 = 22.3\%$, $p = 0.266$

b

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	4.100	1.476	2.179	1.207	6.993	2.778	0.005	
Gunga1996	3.300	1.452	2.109	0.454	6.146	2.272	0.023	
Ryan2016	1.300	0.872	0.760	-0.409	3.009	1.491	0.136	
Trudel2017	3.700	1.081	1.169	1.581	5.819	3.423	0.001	
	2.681	0.568	0.322	1.569	3.793	4.724	0.000	

Lower EPO   Higher EPO

Fixed-effects model used. Heterogeneity: $I^2 = 33.39\%$, $p = 0.212$

Figure

a

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	-0.130	0.079	0.006	-0.285	0.025	-1.644	0.100	
Dunn1984	-0.110	0.061	0.004	-0.230	0.010	-1.791	0.073	
Horeau2022	0.010	0.041	0.002	-0.070	0.090	0.246	0.806	
Horeau2024	0.090	0.061	0.004	-0.030	0.210	1.473	0.141	
Nay2020	-0.200	0.056	0.003	-0.309	-0.091	-3.599	0.000	
Ryan2016	0.060	0.124	0.015	-0.183	0.303	0.485	0.628	
	-0.053	0.049	0.002	-0.148	0.043	-1.082	0.279	

Lower Reticulocyte count   Higher Reticulocyte count

Random-effects model used. Heterogeneity: I² = 71.37%, p = 0.004

b

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	0.420	0.087	0.008	0.249	0.591	4.816	0.000	
Ryan2016	0.060	0.124	0.015	-0.183	0.303	0.485	0.628	
Trudel2017	-0.010	0.041	0.002	-0.091	0.071	-0.241	0.809	
	0.155	0.146	0.021	-0.132	0.441	1.057	0.291	

Lower Reticulocyte count   Higher Reticulocyte count

Random-effects model used. Heterogeneity: I² = 89.92, p < 0.001

Figure

a

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	0.478	0.075	0.006	0.330	0.626	6.340	0.000	
Horeau2022	0.240	0.075	0.006	0.093	0.387	3.194	0.001	
Horeau2024	0.620	0.048	0.002	0.527	0.713	13.005	0.000	
Lampe1992	0.380	0.146	0.021	0.095	0.665	2.609	0.009	
Nay2020	0.550	0.112	0.013	0.330	0.770	4.892	0.000	
Trudel2017	0.190	0.056	0.003	0.080	0.300	3.396	0.001	
	0.408	0.089	0.008	0.235	0.582	4.609	0.000	

Lower RBC Count   Higher RBC Count

Random-effects model used. Heterogeneity: I^2 = 88.16%, p < 0.001

b

	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	-0.181	0.074	0.005	-0.326	-0.036	-2.448	0.014	
Lampe1992	-0.110	0.138	0.019	-0.381	0.161	-0.797	0.425	
Trudel2017	-0.210	0.056	0.003	-0.320	-0.100	-3.754	0.000	
	-0.191	0.042	0.002	-0.274	-0.108	-4.499	0.000	

Lower RBC Count   Higher RBC Count

Fixed-effects model used. Heterogeneity: I^2 = 0.00%, p = 0.787

Figure

a

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Culliton2021	1.870	0.936	0.876	0.035	3.705	1.998	0.046
Horeau2022	4.110	1.118	1.250	1.919	6.301	3.676	0.000
Horeau2024	8.570	0.341	0.116	7.901	9.239	25.119	0.000
Nay2020	3.570	0.765	0.585	2.071	5.069	4.669	0.000
	4.586	1.843	3.397	0.974	8.198	2.488	0.013

Lower Iron   Higher Iron

Random-effects model used. Heterogeneity: I² = 96.2, p < 0.001

b

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Culliton2021	4.100	2.081	4.331	0.021	8.179	1.970	0.049
Horeau2022	8.320	2.008	4.032	4.384	12.256	4.143	0.000
Horeau2024	9.810	0.502	0.252	8.826	10.794	19.539	0.000
Nay2020	11.500	1.282	1.644	8.987	14.013	8.969	0.000
	8.968	1.195	1.427	6.626	11.309	7.507	0.000

Lower Transferrin Saturation   Higher Transferrin Saturation

Random-effects model used. Heterogeneity: I² = 69.17%, p = 0.021

Figure

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Culliton2021	1.800	1.173	1.376	-0.499	4.099	1.534	0.125	
Horeau2022	0.450	1.342	1.802	-2.181	3.081	0.335	0.737	
Nay2020	0.900	0.557	0.310	-0.191	1.991	1.617	0.106	
Ryan2016	-3.050	1.343	1.804	-5.683	-0.417	-2.271	0.023	
Trudel2017	-0.500	1.360	1.849	-3.165	2.165	-0.368	0.713	
	0.103	0.745	0.556	-1.358	1.564	0.138	0.890	

Random-effects model used. Heterogeneity: I² = 56.78%, p = 0.055

b

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Horeau2022	0.260	1.387	1.923	-2.458	2.978	0.187	0.851	
Nay2020	0.840	0.393	0.155	0.069	1.611	2.135	0.033	
Trudel2017	-0.900	1.417	2.009	-3.678	1.878	-0.635	0.525	
	0.684	0.366	0.134	-0.033	1.401	1.870	0.062	

Fixed-effects model used. Heterogeneity: I² = 0.00%, p = 0.472

Figure

	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Horeau2022	-1.390	0.855	0.731	-3.066	0.286	-1.625	0.104
Horeau2024	0.000	0.541	0.293	-1.060	1.060	0.000	1.000
Lampe1992	-0.400	0.976	0.952	-2.312	1.512	-0.410	0.682
Trudel2017	0.800	0.466	0.217	-0.113	1.713	1.717	0.086
	0.131	0.309	0.096	-0.476	0.737	0.423	0.673

Fixed-effects model used. Heterogeneity: I² = 46.24%, p = 0.134

Figure

Study name	Statistics for each study							Difference in means and 95% CI
	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Horeau2022	0.010	0.327	0.107	-0.631	0.651	0.031	0.976	
Horeau2024	0.050	0.177	0.031	-0.297	0.397	0.283	0.777	
Lampe1992	0.300	0.560	0.313	-0.797	1.397	0.536	0.592	
Trudel2017	0.100	0.164	0.027	-0.222	0.422	0.609	0.543	
	0.078	0.111	0.012	-0.139	0.295	0.704	0.482	

Lower MCH   Higher MCH

Fixed-effects model used. Heterogeneity: I^2 = 0.00%, p = 0.970

Figure

Study name	Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value
Horeau2022	0.580	0.132	0.018	0.320	0.840	4.380	0.000
Horeau2024	0.000	0.072	0.005	-0.142	0.142	0.000	1.000
Lampe1992	0.800	0.413	0.170	-0.009	1.609	1.938	0.053
Ryan2016	-0.100	0.594	0.353	-1.264	1.064	-0.168	0.866
Trudel2017	0.100	0.095	0.009	-0.086	0.286	1.054	0.292
	0.246	0.143	0.020	-0.033	0.525	1.726	0.084

Random-effects model used. Heterogeneity: I² = 77.37%, p = 0.001

Figure

Fixed-effects model used. Heterogeneity: I² = 0.00%, p = 0.500

Figure