Principles of protein abundance regulation across single cells in a mammalian tissue

Andrew Leduc[1], Gergana Shipkovenska[2], Yanxin Xu[2], Alexander Franks[3], & Nikolai Slavov[1,4,✉]

[1]Departments of Bioengineering, Biology, Chemistry and Chemical Biology, Single Cell Proteomics Center, and Barnett Institute, Northeastern University, Boston, MA 02115, USA
[2]Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA, USA
[3]University of California, Santa Barbra, Santa Barbara, CA, USA
[4]Parallel Squared Technology Institute, Watertown, MA 02472, USA

✉ Correspondence: leduc.an@northeastern.edu or nslavov@northeastern.edu
∈ Code & Data: github.com/SlavovLab/SC-metabolic_labeling

Protein synthesis and clearance are major regulatory steps of gene expression, but their in vivo regulatory roles across the cells comprising complex tissues remains unexplored. Here, we systematically quantify protein synthesis and clearance across over 4,200 cells from a primary tissue. Through integration with single-cell transcriptomics, we report the first quantitative analysis of how individual cell types regulate their proteomes across the continuum of gene expression. Our analysis quantifies the relative contributions of RNA abundance, translation, and protein clearance to the abundance variation of thousands of proteins. These results reveal an putative organizing principle: The contributions of both translation and protein clearance are linearly dependent on the cell growth rate. Further, we find that some proteins are primarily regulated by one mechanism (RNA abundance, translation, or clearance) across all cell types while the abundances of other proteins is dominated by different regulatory mechanisms across cell types. Our reliable multimodal measurements enabled quantifying and functionally interpreting molecular variation across single cells from the same cell type. The protein-protein correlations are substantially stronger than the mRNA-mRNA ones, which is mediated by protein clearance regulation. The protein-protein correlations are stronger not only for directly interacting proteins but also between functional sets of proteins. Further, these protein correlations allow identifying cell-type specific functional clusters. These clusters vary across cell types, revealing differences in metabolic processes coordination, partially mediated by protein clearance regulation. Our approach provides a scalable multiplexed framework for quantifying the regulatory processes shaping mammalian tissues and reveals organizing principles determining the relative contributions of translation and protein clearance to the proteomes of primary mammalian cells.

Main

Metabolic pulse labeling proteomics has suggested that protein synthesis and clearance (the joint contribution of degradation and dilution) significantly regulate cellular proteomes[1–4]. Further, these studies have revealed how reliance on synthesis and clearance is shaped by biological contexts such as tissue type[4,5] and cellular growth rate[4,6]. However, studies of gene expression regulation within primary tissues have mostly focused on regulation of RNA abundance as they can be measured by scalable single-cell DNA sequencing technologies. Thus, the regulation of protein synthesis and degradation across different cell types within complex tissues remain unexplored.

Regulation of protein synthesis and degradation may also contribute to functional differences across individual single cells within a cell type. Due to the sparse nature of the data generated by single-cell DNA sequencing approaches, analysis is usually restricted to comparing clusters of cells or in vitro grown cell lines[7,8]. Analyzing protein covariation across single cells with a cell type may identify regulatory mechanisms[9], which motivates us to explore this potential within a mammalian tissue.

Using scalable single-cell mass spectrometry (MS), we sought to elucidate principles of protein abundance regulation in a complex mammalian tissue across the full continuum of gene expression. We chose murine trachea as a model and quantified the the contributions of RNA abundance, translation and protein clearance across diverse cell types, Fig. 1a. Our analysis uncovers the ways in which the primary mode of gene regulation differs for individual proteins and functional groups of proteins, as well as across the same proteins depending on the specific cell type. We also reveal substantial functional protein covariation across single cells of the same type and its underlying regulatory factors. Protein clearance substantially shapes both protein abundance and protein-protein covariation in individual single cells, substantially contributing to discordance between mRNA and protein covariation patterns.

To enable this analysis, we introduced combined multiplexing of isotopic peaks from metabolic pulse labeling with isotopic chemical barcoding for sample multiplexing via the plexDIA framework[10], Fig. 1b. Previously, plexDIA has been used for sample multiplexing (enabling measuring thousands of cells from a human tissue[11]) and for metabolic pulse labeling for protein turnover measurements[12,13], but not for both simultaneously. We reasoned that combining these differ-

ent approaches could enable quantifying protein synthesis and clearance rates while increasing the throughput of single cell analysis, Fig. 1b. We achieved this by a metabolic pulse with +6 Da heavy lysine combined by multiplexing single cells with mTRAQ $\Delta 0$ and $\Delta 8$ tags creating a 16 Da offset between lysine containing peptides from different single cells. To prepare single cell samples, we leveraged the nPOP glass-slide sample preparation[14,15]. This approach facilitated preparing 1,300 single cells per batch with multiplexing two cells per set.

This design empowered the analysis of over 4,200 single cells from 6 mice fed with heavy Lys food for either 5 or 10 days, Fig. 1a,c. We aligned this data set with over 36,000 single cells from mice in the space of 2,201 shared gene products quantified at the mRNA and protein level[16]. We first clustered and annotated each dataset separately based on abundance of common marker proteins and then aligned the cell types, Extended Data Fig. 1a. To validate cell type assignment across data sets, we correlated relative levels of protein and mRNA abundance across cell types. We observed remarkably strong agreement for matched cell types. The strong correlations among independently annotated cell type clusters indicate that our data support robust cell type assignment without the need for more complex integration approaches, Fig. 1d and Extended Data Fig. 1b.

To improve the quantification, we devised strategies to accounting for artifacts such as variability in liquid chromatography (LC) and missing values. These approaches were added as functionality to our QuantQC package used for downstream processing of searched single cell data[15]. First, due to the large number of LC-MS/MS runs, we noticed the intensities of certain peptides drifting as a function of the run number, Fig. 2a. Similar to previous approaches applied in metabolics[17], we regressed out unwanted variation for each peptide using a spline fit to the LC run order, see methods. To account for the influence of missing data, we devised a missingness-aware approach to improve fold change estimates across cell types. For each protein, we identified the percentage of cells with missing values caused by either biological or technical factors[18], allowing us to scale fold changes proportionally to biological missingness, see methods. We identified cell size as a leading factor for technical missingness with LC batch and label explaining missingness to a lesser extent Extended Data Fig. 2b-e. The approach improved agreement between relative protein and RNA abundance estimates (Extended Data Fig. 2f-h), suggesting improved quantification.

To achieve accurate estimates of protein synthesis and clearance rates, we explicitly modeled the amino acid recycling (Extended Data Fig. 3a), which is significant for *in vivo* metabolic pulse

Figure 1 | Single-cell metabolic pulse proteomics and transcriptomics in murine trachea. a, Mice for sc-proteomics were fed a diet of +6 Da heavy lysine for 5 days ($n = 4$) or 10 days ($n = 2$). Single cells prepared via nPOP multiplexed sample preparation were labeled with either mTRAQ $\Delta0$ or $\Delta8$. **b**, Mass spectra from precursor ions for peptide LYSEFLGK are plotted showing the 4 co-eluting isotopic states, $\Delta0$ labeled peptides from the first labeled cell with lysine +0 Da or +6 Da and the $\Delta8$ labeled peptides from the second cell, resulting in a 16 Da mass shifted precursors for the +0 and +6 lysine. Single cells from the trachea of 4 different mice were processed for sc-mRNA seq by 10x Genomics. **c**, The resulting dataset contains 4,200 single cells with proteomic measurements of peptides containing light and heavy lys and 36,000 cells with RNA data. UMAP visualization shows the corresponding cell types clustered in the space of either protein or RNA abundance. **d**, The relative abundance of protein and mRNA levels in corresponding cell types are strongly correlated (diagonal). **e**, Number of proteins quantified per cell. Across all cells, protein abundances was quantified for 2,215 unique proteins and synthesis and degradation rates for 1,506 unique proteins. Details shown in Extended Data Fig. 4 **f**, Protein abundance at steady-state for a given protein can be modeled as the rate of protein synthesis (which equals mRNA concentration multiplied by translation efficiency, the number of proteins transcribed per mRNA),and the rate of proteins cleared via degradation, secretion, or dilution due to cell growth. **g**, The reproducibility of estimates for protein and mRNA abundance, protein clearance, and translation per mRNA is visualized in basal cells by comparing technical replicates of averaged non-overlapping subsets of single cell.

labeling experiments[19]. First, we computed the exchange of free lysine amino acids in each cell type utilizing miss-cleaved peptides[2], see methods. The rate of exchange differed depending on cell type, generally higher for epithelial and immune cell types compared to mesenchymal and smooth muscle, Extended Data Fig. 3b. Cell-type specific lysine exchange rates were then used to correct half lives for all single cells across each respective cell type. Correcting for recycling reduced half life of protein clearance from roughly 7 days on average to 3 days, Extended Data Fig. 3c. Further, applying this correction harmonized clearance rates computed from 5 and 10 day cells, Extended Data Fig. 3d. On average, clearance rates were lowest for chondrocytes and smooth muscle cells while being highest for basal cells, Extended Data Fig. 3f. Overall, we quantified clearance rates for 1,506 proteins with an average of 514 proteins per cell, Fig. 1e.

Clearance rates can vary across cell types cultured in vitro[20], and we focus on quantifying this variation across the cell types of a tissue. To examine this, we correlated clearance rates across cell types. The average similarity across cell types is $\rho = 0.50$ Extended Data Fig. 3g, much lower than the reliability estimated from independent subsets of cells of the same type, as shown on the diagonal. For example, a correlation of $\rho = 0.51$ between basal cells and chondrocytes while internal estimates for each cell type are above $\rho = 0.75$. This difference strongly suggests cell-type specific rates of protein clearance. Thus, we sought to quantify and understand how such regulated protein clearance plays into the larger picture of gene expression.

To this end, we inferred a range of parameters spanning the gene expression process. Under steady state, protein synthesis equals the product of protein clearance rate and abundance[4,21], Fig. 1f. Protein synthesis itself is the joint contribution of the amount of mRNA molecules and the rate at which each molecule is translated into proteins. Our data enabled us to estimate each of these quantities and their influence at the cell type level.

We next sought to examine the reliability of the inferred quantities. For the scRNA-seq, we achieved depth of at least 3,000 unique transcripts per with at least 10k unique UMIs per single cell, comparable to the best published 10x genomics data sets, Extended Data Fig. 4a. For the single-cell proteomics data, we quantified abundances for 2,205 proteins with an average of 805 per single cell, Fig. 1e. Variation on the number of proteins quantified can be attributed to cell size variation, Extended Data Fig. 4b. Further, the strong correlation between cell size and total protein highlights the consistency and precision of the sample preparation. To evaluate the accuracy of

protein quantification, we examined the consistency of relative abundances for peptides mapping to the same protein[22], Extended Data Fig. 4c. For example, the 5 most abundant peptides from the protein Tgm2 show high correlation across single cells, $\rho = 0.88$. On average, peptides originating from a protein correlated with $\rho = 0.50$. As expected, proteins that varied more significantly from cell to cell displayed higher correlations, Extended Data Fig. 4d.

To evaluate the consistency of the estimates within cell types, we partitioned cells into two populations and computed the average mRNA and protein abundance, protein clearance rate and translation rate, Fig. 1g and Extended Data Fig. 4e. The reliability of agreement between the estimates was above $\rho = 0.85$ for protein abundance, mRNA abundance, and translation. Lower correlation, on average $\rho = 0.69$, for clearance rates might reflect the smaller dynamic range of the clearance rates. These estimates allowed us to account for the influence of measurement noise when determining how different regulatory modes impact protein abundance across cell types.

Growth rate and absolute abundance analysis

One property that has been suggested to influence the mode of gene expression regulation is growth rate[4,6,21]. Some reports have emphasized the compensation of protein dilution by changes in synthesis rates[6,21] while others have emphasized its impact on altering protein abundance[4]. No estimates exist for primary cell types directly from mammalian tissues. To investigate this influence, we first estimated cell growth rate based on the incorporation of histone proteins[12]. Here, we explicitly computed the effective doubling time of each cell type by counting the number of cell divisions that occurred since the feeding started, see methods Fig. 2a. Cell types displayed a variety of growth rates ranging from 9.9 day doubling time to no recorded division events within the 10 day pulse.

These growth differences are important since the growth related dilution contributes to protein clearance. At faster growth rates, the rate of dilution assumes a larger fraction of protein clearance, reducing the protein to protein variability in clearance rates[4,21], Fig. 2b,c. As a result, the dynamic range of clearance rates is reduced (the smallest rate is the rate of cell growth), thus directly affecting the scaling of clearance rates between cell types, as visualized via the shallow slope when comparing clearance rates between cell types with growing at different rates, Fig. 2b. The dependence of this slope scaling with growth rate is strongly confirmed by the fact that the slope between

degradation rates for these cell types is close to 1, Extended Data Fig. 5a.

Independent from the slope scaling, the data reveal significant variation (scatter) of the clearance rates of basal cells and chondrocytes, with a correlation of only 0.50, despite higher reliability of the clearance rate estimates (0.7 - 0.75). This trend is further observed between most cell types, Extended Data Fig. 3g. This variation reflect cell-type specific differences in growth rates, as indeed confirmed by the fact that the correlation between the corresponding protein degradation rates is the same.

In addition to cell-type specific degradation, clearance rates have a wider dynamic range at low growth (Fig. 2b), and thus the potential for stronger impact on protein abundance. To evaluate this potential across all cell types, we compare the dynamic range of the clearance rates variation across proteins to the corresponding cell growth rate. We found a strong inverse relationship with growth rate, $\rho = -0.82$, Fig. 2c, consistent with the expectation. To more directly quantify if the potential is realized, we examines the joint distribution of clearance rates and protein abundance in chondrocytes. It shows a protein clearance explains roughly 45 % of the variation in protein abundance estimated by square of the Pearson correlation, Fig. 2d. Conversely, only 29% of variance is explained for faster growing basal cells, Extended Data Fig. 5b. Overall, the extent of abundance variation explained by clearance attenuates as growth rate increases, following a clear linear dependence Fig. 2e. This result extends our previous observation made across tissues to cell types within a tissue[4].

Accurately estimating this influence required accounting for reliability of the measurements. To do so, we applied the Spearman correction to the fraction of variance explained based on reliability estimates derived in Extended Data Fig. 4e, see methods. We also applied this method to estimating the fraction of variance explained by mRNA abundance, Fig. 2d. While both mRNA abundance and clearance are estimated relatively independently of protein abundance, translation shares peptide ionization bias with protein abundance estimates. Thus, estimates for fraction of variation explained by translation were taken to be the remaining variance. These estimates are in agreement with another estimate derived from the square of the Pearson correlation, Extended Data Fig. 5c. Corrected and uncorrected estimates for each mode of regulation are displayed in Extended Data Fig. 5d. The contribution of mRNA abundance did not correlate strongly to growth rate. In contrast, translational regulation increased in direct proportion to growth rate, Fig. 2d.

Figure 2 | The regulatory contributions of translation and clearance to absolute protein abundance scales linearly with growth rate a, Plotting the recycling adjusted heavy to light ratios of histone peptides reveals multimodal distributions, with modes corresponding to the number of times cells have divided and partitioned histones. These distributions were used to infer cell doubling time, which is proportional to growth rate, $ln(2)/k_g$. **b**, Comparisons of clearance rates between different cell types, ordered by growth-rate differences between the cell types. For cell types with more similar growth rates, the slope of the clearance rates between the cell types converges to 1. **c**, The variability in degradation rates across proteins is estimated by the standard deviation of the log transformed clearance rates and plotted against growth rate, $\rho = -0.82$. **d**, For the most slowly growing chondrocyte cells (with no recorded cell divisions), protein clearance explains 0.45 % of protein abundance variation. **e**, Growth rate strongly predicts the influence of degradation and translation on absolute protein abundance. The influence of translation is proportional to the growth rate while the influence of clearance is inversely proportional.

Relative regulation across cell types

Having analyzed how different regulatory modes influence absolute abundance variation across protein , we next used our our measurements to characterize how individual proteins are regulated across cell types. Similarly, we aim to quantify the impact of all regulatory modes. The simplest

possibility is that the relative abundances of a protein is regulated primarily by single mechanism. This is lustrated with Sult1d1, which primarily co-vary with mRNA abundance, $\rho = 0.98$, Fig. 3a. Other proteins such has Sec14I2 primary co-vary with translation, $\rho = 0.98$. To estimate measurement reliability, we leverage the quantification of distinct peptide sequences for a given protein along with sample preparation replicates within a Bayesian framework. This allowed us to reflect higher uncertainty in the translation quantities, which share uncertainty from the mRNA abundance, protein abundance, and protein clarence measurements, see methods. Our uncertainty estimates control for factors such as sample preparation variability and potential MS interferences.

To explore the full range of regulation for all quantified proteins, we correlated protein abundance to mRNA abundance, translation rate, and clearance half life, Fig. 3b and Fig. 6a. The correlations for all regulatory modes were broadly spread and positively biased, indicating that proteins are regulated by different modes to different degrees. The average correlation was the highest for mRNA abundance, suggesting it significantly influences the abundance of many proteins. For some protein groups, such as actin filaments, RNA abundance appears the dominant regulatory factor. For other protein groups, translation and protein clearance appear the dominant regulatory factors, Fig. 3b. Specifically, different pathways controlling metabolic functions, translation, localization, and degradation of proteins were primarily controlled by translation and clearance. We further extended this analysis to important functional groups of proteins such as chromatin binding proteins, mRNA binding proteins, and E3 ligase, Fig. 3c. Clearance played a strong role in regulating relative levels of mitochondria and the proteosome, while translation had a prominent regulatory role for E3 ligases. The relative influence of translation and clearance has only previously been estimated across a single cell type cultured in vitro and responding to stimulus[2]. Our results suggest stronger overall regulatory impact when extending measurements to the new context of how diverse cell types from a primary tissue regulate their proteomes.

The univariate analysis in Fig. 3, suggests that most proteins are regulated by multiple mechanisms, and we further investigated this with bivariate analysis, Extended Data Fig. 6. We observed that as the correlations between proteins and mRNA abundance decreases, the correlation between protein and translation or clearance rates increase, Extended Data Fig. 6a. Interestingly, proteins with the strongest correlation to mRNA abundance had a strong inverse correlation to clearance half life. This suggests cell type specific proteins with fast turnover require even tighter transcrip-

Figure 3 | Regulation of relative protein abundance across cell types a, Examples of proteins regulated by a primary mechanism across all cell types, mRNA abundance for Sutl1d1 and translation control for Sec12I2. **b**, Correlations of cell-type average protein abundance to mRNA abundance, translation rate, and protein clearance half lives for all proteins. Example proteins predominantly controlled by mRNA abundance, translation, or clearance are plotted, displaying correlations for individual proteins grouped by GO terms. **c**, Average correlations for each mode of regulation to proteins from functional groups. **d**, Clearance contributes significantly to protein fold changes between basal cells and chondrocytes, but less for other cell-ype pairs growing faster, as shown in the heatmap. **e**, Examining protein S100a1 demonstrates how regulation methods can be cell type specific, with mRNA abundance and clearance explaining high abundance in chondrocytes, but low translation explaining low abundance in basal cells. **f**, Heatmaps are displayed for proteins organized by the primarily regulatory mode, mRNA abundance, translation, or clearance. For proteins primarily regulated by translation in chondrocytes, there is substantial diversity in regulation patterns in fibroblasts. **g**, GO terms for proteins enriched to be regulated primarily through transcription, translation and clearance show cell type specific trends.

tional control. However, these mechanisms may not only act independently, but synergistically. Indeed, proteins such as St13 are co-regulated by both transcription and translation, Extended Data Fig. 6b. This can also be observed for groups of proteins, such as certain keratin filaments synergistically regulated by both transcription and translation in basal cells Extended Data Fig. 6c. The contributions of all modes to the regulation of proteins abundance is listed in supplementary table 1, both for individual proteins and GO terms. Collectively, our results indicate that the dominant mode of regulation across cell types varies across proteins.

The primary mode of regulation also varies across cell types. For example, while protein clearance explains 38% of variation in relative protein abundance between basal cells and chondrocytes, it explains little variation for the same set of proteins between different cell types, such as basal and immune cells, Fig. 3d. Generally, we see clearance explain the most significant amount of variance between cell types with slower growth rates such as chondrocytes and smooth muscle cells. This is because cell-type specific differences in degradation exert greater effect on protein clearance when clearance rates are less influenced by protein dilution.

Next, we increase our resolution and consider the variation in the regulatory mode both across cell types and proteins. As an example, S100a1, is up-regulated in chondrocytes because of increased RNA abundance and decreased clearance while in basal cells, it reduced abundance is primarily achieved by translationally suppression, Fig. 3e. To visualize such trends, we plotted the subsets of proteins whose relative abundance in chondrocytes is primarily regulated by a single mode of regulation, Fig. 3f, see methods. To explore of the mode of regulation is protein-specific and similar across cell types, we focused on the subset of proteins primarily regulated by translation in chondrocytes. Visualizing their regulatory modes fibroblasts shows a mix of regulatory strategies, including significant control by transcription and clearance, Fig. 3f. These results generalize the observations from S100a1 and indicate that the mode of regulation is both protein and cell-type specific.

To visualize this result more globally and representatively, we systematically identified functional groups of proteins whose regulation is dominated by a single regulatory mode across all cell types, Fig. 3g. Some terms such as cytoplasmic translation were regulated by different modes across cell types such as transcription in basal cells and translation in secretory and immune cells. Many structural process were regulated by clearance across cell types such as microtublue poly-

merization in ciliated cells, lamellipodium organization in smooth muscle cells and intermediate filament organization in immune cells.

Comparing mRNA and protein covariation patterns within cell type

Although grouping cells by type is convenient and explains much of the protein abundance variability, cells vary functionally even within a cell type. These functional variations are mediated in part by regulated protein abundance changes. Indeed, when examining the most numerous basal cell population, we found that some proteins such as Cavin1 varied at least 30 fold, Fig. 4a. Further, the abundance of Cavin1 varied uni-modally across single cells, suggesting that further sub-clustering would incompletely explain its variation. Interestingly, mRNA levels for Cavin1 varied about 4 fold less. This difference in dynamic range for the protein and RNA products of a gene generalize to all analyzed proteins: the protein abundances vary to a greater degree across single cells than mRNA for over 90 % of quantified genes products, Extended Data Fig. 7a. To better understand the implications of this variation and distinguish between biological and technical origins, we sought to analyze covariation patterns to identify potential functions associated. Further, as we did not measure mRNA and protein abundance in the same single cells, comparing single-cell covariation patterns at the protein and mRNA level enabled us to determine the extent to which protein level covariation is regulated transcriptionally.

The large variation of Cavin1 within basal cells is likely reflecting the variable abundance of caveolae structures, as supported by the strong correlation ($\rho = 0.81$) to the other subunit, Cav1, Fig. 4a. At the mRNA level, we observed a weaker correlation between Cavin1 and Cav1 of $\rho = 0.34$. Extending this analysis to all protein pairs indicates an overall similarity (Pearson correlation) between pairwise correlations of RNA and proteins, $\Psi = 0.31$. Beyond the similarity of covariation at the mRNA and protein level, this result emphasizes significantly stronger correlations among protein pairs. For example, over 5,000 pairwise protein correlations exceed a correlation of 0.3, compared to just 340 at the mRNA level.

To determine whether these high correlations reflected cellular functions, we compared groups of related gene products. Proteins annotated to interact in either the CORUM or STRING database showed significantly higher correlation at the protein level as comapred to mRNA, Fig. 4c and Extended Data Fig. 7b, see methods. These results indicate that protein correlations capture func-

tional covariation and motivated us to examine differences in clustered correlation modules. We applied hierarchical clustering on a subset of proteins and mRNAs, see methods. First, this subset consisted of proteins and mRNAs for which the proteins had at least one pairwise correlation above 0.3. This resulted in a group of roughly 400 unique proteins. We then plotted the correlation patterns against each other as the lower and upper triangle of a correlation heatmap[7], Fig. 4d and Extended Data Fig. 7c. We observed several distinct correlation modules primarily present at the protein level, Extended Data Fig. 7c, each found to be enriched for different functional processes, FDR < 0.01. For example, a module of proteins enriched for mesenchymal transition had an average correlation of 0.43 at the protein level and 0.02 at the mRNA level. Applying the same procedure, but on a subset of gene products where the mRNAs contained at least one pairwise correlation above 0.2, we found fewer confident clusters and many had no clear functional enrichment, Extended Data Fig. 7c.

We then set out to understand why certain covariation patterns would be present primarily at the protein level. One potential reason is that active regulation of protein translation or clearance contribute to the functional protein covariation not observed at the level of transcripts. As we measured protein clearance rate with single-cell resolution, we investigated its contribution. To gain intuition, we first inspected the chromatin bound proteins Hmga1 and Hmgn2 whose correlation at the protein level is 0.56, while only 0.01 at the level of transcripts Fig. 4d. This difference may stem from the regulation of Hmga1 protein abundance by clearance, reflected in a correlation of $\rho = -0.48$ between Hmga1 clearance rates and abundance across cells, Fig. 4d. To explore if this example is representative and clearance regulation contributes globally, we plotted the difference in correlations between mRNA and protein abundance, versus the extent to which each protein was regulated by clearance (the average of the correlations of clearance and abundance for each protein across single cells). We observed that pairs of gene products more highly correlated at the protein level are significantly more likely to be controlled by protein clearance, Fig. 4f.

We then expanded our analysis to test whether these trends generalize to other cell types and depend on factors such as growth rate. We constrained our analysis to populations with at least 500 single cells to ensure a sufficient number of data points when computing protein-protein correlations. To estimate the contribution of clearance to protein-protein correlations, we compared the pairwise correlations between protein abundance and clearance and quantified the similarity

Figure 4 | Protein covariation within cell type is only partially reflected in mRNA covariation and significantly shaped by protein clearance regulation a, The dynamic range of measured protein concentrations for Cavin1 is about 4 fold larger than at the mRNA level. Pairwise scatter plots of relative fold changes across single basal cells for Cav1 and Cavin1 transcripts and proteins. **b**, A scatter plot of all pairwise protein-protein versus corresponding mRNA-mRNA correlations reveals a correlation $\Psi = 0.31$. **c**, Pairwise correlations for all proteins and transcripts are centered around 0, but correlations for interacting proteins are positive and larger than the corresponding mRNA correlations on average 0.12. **d**, Protein correlation modules detected via hierarchical clustering are plotted with correlations for the corresponding transcripts. **e**, Proteins Hmga1 and Hmgn2 are correlated but their mRNAs are not. This discrepancy is partly explained by the clearance regulation of these proteins by clearance reflected in the correlations between abundance and clearance rates. **f**, The extent of protein clearance regulation for protein pairs can explain the difference between their correlation and the correlations among corresponding mRNA pairs, $\rho = -0.44$. **g**, Pairwise protein clearance rate correlations are similar to the corresponding protein abundance correlations, $\Psi = 0.24$. This agreement Ψ depends linearly on the cell growth rate. The agreement between pairwise protein and mRNA correlations exhibits the inverse linear dependence. **h**, GO terms whose members covary significantly, FDR of 1 % for proteins and mRNA. Proteins show considerable differences across cell types while differences in covariation of mRNA levels are less present.

by the correlation Ψ, analogous to Fig. 4b; for chondrocytes, clearance $\Psi = 0.24$, suggesting that protein clearance contributes significantly to shaping protein-protein correlations, Fig. 4g. Analogously, we computed for other cell types clearance Ψ and RNA Ψ (correlation between RNA-RNA and protein-protein correlations), and plotted them as a function of the growth rate, Fig. 4g. The results indicate strong linear dependencies: For more slowly growing cell types such as chondrocytes, clearance covariation more strongly reflected protein abundance covariation. This trend was reversed for when comparing protein and mRNA covariation patterns to cell growth rate. These results were consistent with our previous observations about more slowly growing cells relying more strongly on regulation by protein degradation. However, the extent of regulation by transcription within cell type depends more on growth rate than observed at the absolute abundance level.

Having established differences in correlation patterns for proteins and RNAs across single cell, we next examined the differences in covariation patterns within functional groups of proteins and mRNAs. For each cell type, we computed the protein-protein and mRNA-mRNA correlations within a functional group and compared them to the corresponding distribution of all correlations within each modality. We observed 315 GO terms with significantly (at 1% FDR) higher protein correlations in at least on cell type. In contrast, only 185 GO terms reached significance at the mRNA level (supplemental table 2, Fig. 4h), despite the larger number of single cells. These differences suggest that protein abundance variation is better at capturing functional state variation across individual cells. These functional states show significant degrees of cell-type specificity, as demonstrated by the series of different functional groups most strongly co-regulated within individual cell types, Fig. 4h. Notably, terms such as ER-Golgi intermediate compartment and other terms involved in protein secretion were significant correlated in the secretory cell population, while proteins participating in glycolysis strongly co-varied in the chondrocyte cells inhabiting low oxygen environments,[23].

Within cell type protein correlation

Having observed strong co-varation patterns at the protein level, we sought to further leverage this signal to infer differences in the coordination of proteins functions across cell types. We originally observed large dynamic range spanning 50 fold for the concentration of intermediate filaments (IF) in both basal cells (Keratin 5) and fibroblasts (Vimentin). Curious as to why these cells would have

such varied concentrations, we explored which proteins were correlated to high IF concentration by computing the correlations of all proteins to Krt5 and Vim in both fibroblast and basal cells, i.e., correlation vectors. Interestingly, the correlation vectors for either Krt5 and Vim showed no similarity between basal cells and fibroblasts, Fig. 5a. However, the correlation vectors of the primary IF in each cell type (Krt5 in basal cells and Vim in fibroblasts) are strongly correlated, $\rho = 0.80$. This correlation pattern illustrates how correlations can reveal functional changes.

To understand the functional variability associated with high or low IF concentration, we next inspected the subset of proteins with high positive and negative correlations to the primary IFs, Fig. 5b. We observed a gradient of protein abundance present in both cell types. It reflects a shift from cells with high concentration of cytoskeletal proteins (suggesting high mechanical force) to cells with high concentration of proteins participating in cellular redox response to stress. These results suggest a common phenotypic axis of variation, with unique proteins participating from each cell type based on the similarity of their their primary functional role contingent on cellular context.

We next sought to generalize this analysis to identify additional proteins that potentially change their functional co-ordination across cell types. The first step in our procedure was to compute vectors of pairwise correlations for the ith protein, denoted as $\mathbf{b}_i$ in basal cells and $\mathbf{f}_i$ for fibroblasts. We then quantified the similarity between these correlation vectors by their correlations, $\langle \mathbf{b}_i, \mathbf{f}_i \rangle$ as previously described[24,25], Fig. 5c, supplemental table 3. The distribution of correlations was on average positive, mean $\rho = 0.21$, suggesting shared covariation patterns between cell types. However, many proteins showed minimal correlation vector similarity across cell types. We then partitioned the proteins into two sets of highly similar and dissimilar proteins with a threshold for high ($\rho > 0.50$) and low ($\rho > 0.10$) similarity. High similarity proteins from set 1 constituted an axis of common covariation across cell types, including groups proteins participating in cell metabolism, cytoskeleton, and translation, Fig. 5c.

We then utilized this common axis of covariation to compare the covariation for proteins from set 2 across cell types. By clustering correlations between set 1 and set 2 proteins for each cell type, we identified opposing covariation patterns, Fig. 5d. In basal cells, set 2 proteins associated with intermediate filament and fatty acid metabolism (purple cluster) positively correlated with set 1 proteins related to the TCA cycle and cellular stress, while proteins associated with glycoprotein

Figure 5 | **Cell-type specific covariation identifies functional exchanges between proteins a**, Vimentin is the primary intermediate filament (IF) in fibroblasts while Krt5/15 are the primary IFs in basal cells. The correlation vectors (to all other proteins) of each IF are cell-type specific and similar between basal and fibroblasts cells. **b**, Protein abundance across single cells shows gradients of cells defined by the co-expression of either cytoskeletal proteins (suggesting cells are under higher mechanical strain) or redox proteins (suggesting high oxidative stress). **c**, The correlations between correlation vectors for all proteins across basal and fibroblast cells are broadly distributed with many proteins showing high covariation (set 1) while fewer proteins showing low covariation agreement (set 2). A clustergram of protein proteins from set 1 defines functional clusters. **d**, Clustergrams of pairwise correlations between proteins from set 1 and set 2 identify clusters (marked by purple and green) with changing correlations between basal and fibroblast cells.

metabolism and cell adhesion (green cluster) correlated positively with set 1 proteins involved in translation and glycolysis. These trends were reversed in fibroblasts, Fig. 5d, suggesting that these cell types rely on different strategies to provide energy for the same processes.

To understand how these differences in protein-protein correlations are achieved, we examined

the potential for regulation by transcription and protein clearance. Applying the same correlation vector analysis to the corresponding mRNA and clearance rate data resulted in much weaker correlation structuresExtended Data Fig. 8a. To reduce the potential contribution of measurement noise, we compared the co-variances of the averaged mRNA and clearance rates across proteins from the two clusters of set 1 and set 2 proteins, Extended Data Fig. 8b. This revealed correlation structures in the mRNA abundance and protein clearance similar to the structure of to protein-protein correlations This similarity suggests that both RNA abundance and protein clearance contribute to the protein clusters in Fig. 5d.

Discussion

Our data and analysis allowed us extend the analysis of pioneering studies of gene expression regulation in vitro[1,2] to the single cells comprising a complex tissue. This expansion revealed new regulatory trends: Across cell types, differences in cell growth rate proved a strong driver of gene regulatory mechanisms. Our results suggest substantial influence of translation regulation at fast growth rate. This is consistent with previous observations made in rapidly proliferating cancer cells[1]. Our results extend to slower growth rates, were cells gradually shift from translational to clearance regulation of protein abundance.

We observe remarkably simple linear scaling of the contributions of translation and protein clearance to protein abundance variation. This linear scaling is similar to scaling laws that have been observed as a function of cell growth rate in bacteria[26] and yeast[27]. We can mechanistically explain the increasing role of protein clearance with the decreasing dilution rate at slower growth rates, if (and only if) this dilution effect is not compensated by adjustment of synthesis rates as previously posited[4,6,21]. In contrast, the scaling of the translation contribution remains phenomenological but consistent with previous estimates from rapidly growing cells in vitro[1,2]. The discovery of these simple relationships in the growth-rate dependence of gene expression control will stimulate further research into the regulatory mechanisms in primary mammalian cells. This research will be facilitated by the multiplexing framework introduced here; it enables scaling the analysis of protein synthesis and clearance rates by mass tags, which can further increase the plex and thus the throughput[28,29]. The number of multiplexed samples can also be increased with existing mass

tags by modeling the overlap of isotopic envelopes, as done by JMod[30].

The accuracy and scale of our dataset enabled functional analysis based on covariation of proteins and RNA within a cell type. Advances in single-cell proteomics[31–34] enable analyzing thousands of single cells[11,35–37] and performing protein covariation analysis[9,38,39]. Here, we analyze differences between mRNA and protein correlations and their origins. We find that regulated protein clearance strongly contributes to the differences in mRNA and protein covariation patterns (Fig. 5), but it does not account for all differences. In addition to measurement noise, additional contributors include translational regulation and and the longer half lives of proteins; thus, transcriptional bursts are integrated into steadier protein levels that reflect the transcriptional history of the cell, not just a momentary point estimate of mRNA abundance[40].

Our approach opens the door to characterizing gene expression regulation in mammalian tissues and identified a potential organizing principle in the role of growth rate. It also demonstrates the a feasibility of functionally interpreting protein covariation within individual cell types, that will be amplified by technological developments[41].

Resource availability:

Lead contact: Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Andrew Leduc (leduc.an@northeastern.edu).

Materials availability: This study did not generate new unique reagents.

Data availability: All raw data and search engine outputs can be found on MassIVE with ID: MSV000098940. Single cell mRNA sequencing data was collected as part of Lin et al.[16] and was previously deposited to the GEO:GSE244215. Additional processed data needed for reproducing the analysis can be found on Zenodo under DOI 10.5281/zenodo.14902833.

Supplemental data table 1

Supplemental data table 2

Supplemental data table 3

Code availability: The code used for data analysis and figures is at: github.com/Slavovlab/sc_metabolic_pulse

Acknowledgments: We thank Georg Wallmann for assistance generating the custom spectral library used for assessing amino acid recycling, Zhixun Dou and Jayaraj Rajagopal for early discussions, and Vladamir Ondruska and Stefan Binder for help operating and maintaining mass spectrometry instruments. The work was funded by an Allen Distinguished Investigator award through The Paul G. Allen Frontiers Group to N.S., a Bits to Bytes award from MLSC to N.S., an NIGMS award R01GM144967 to N.S., and a MIRA award from the NIGMS of the NIH (R35GM148218) to N.S, and a UH3CA268117 award from NIH to N.S.

Competing Interests: N.S. is a founding director and CEO of Parallel Squared Technology Institute, which is a nonprofit research institute. The authors declare that they have no other competing interests.

Correspondence: Correspondence and requests should be addressed to nslavov@northeastern.edu, nslavov@alum.mit.edu

Author Contributions

Experimental design: A.L. and N.S.

Sample preparation: A.L, G.S., Y.X.

Raising funding & supervision: N.S., A.F.

Data analysis: A.L. and N.S.

Writing & editing: A.L. and N.S.

Methods

Mouse model, handling and metabolic pulse

All mice experiments were performed in compliance with the Institutional Animal Care and Use Committee at Massachusetts General Hospital. Both male and female C57BL/6 mice, either 24-month-old or 4-month-old, were ordered from the NIA. Murine proteomes were metabolically labeled with 13C6 Lysine. Mouse Express L-LYSINE [13C6, 99%] MOUSE FEED kit was purchased from Cambridge Isotopes Laboratories. Mice were first acclimated to the Lys+0 feed for a period of 10 days before switching to the Lys+6 diet for either 5 or 10 days. Mice were fed 5g per day for either 5 days or 10 days before euthanasia. Mice were euthanized with CO2 followed by cervical dislocation. Tissues were harvested post-euthanasia and perfusion with PBS.

Tracheal single cell suspension generation

Immediately after euthanasia, murine tracheas were dissected and dissociated to single cells in a two-step protocol. Tracheas were first incubated in 20 u/ml papain and 7.5 u/ml DNase I for 30 min at 37 degrees C with rotation, then triturated to release cells. Cells, debris and undissociated tracheal husks were pelleted and resuspended for a second enzymatic digestion in a mix of 350 u/ml collagenase I, 250 u/ml hyaluronidase, 60 u/ml dispase, 3.2 u/ml papain and 75 u/ml DNase I for 20min at 37°C with rotation. After the second digestion, the husks including the cartilaginous scaffold had been dissociated fully. Single cells were filtered through a 100μm filter and red blood cells were lysed using standard RBC lysis buffer. Single cells were then prepared fresh for either single cell proteomics or mRNA sequencing.

Descriptions of the age, gender and feeding time for each mice can be found in the metadata file on Zenodo.

Single-cell and bulk sample preparation

Mouse trachea single-cell suspensions were collected on 4 different days, resulting in 4 different single-cell sample preparations following the procedure outlined below.

Samples were prepared using the nPOP sample preparation method for multiplexed single cell proteomics[14] using cell permeability staining to select only intact cells for MS analysis[37]. Cells were concentrated at 1000 cells per μL in 1x PBS and were incubated on ice and in the dark for 20 min with Sytox Green Dead Cell Stain at a concentration of 1 μM (Thermo Fisher S34860). Cells were then washed to remove dye and resuspended in 1X PBS at a concentration of 300 cells per μL for eventual cell sorting. Cells were then sorted in a volume of 300 pL into 9 nL of 100% DMSO droplets on the surface of a fluorocarbon-coated glass slide for cell lysis using the CellenONE cell sorter and liquid handler. Cells sorting gates excluded all cells with measurable green fluorescent intensity (Sytox Green positive) and selected cells between 11 and 27 μm in diameter with elongation below 2. The single cells were incubated for overnight in digestion buffer of 100 ng/μL Trypsin Gold (Promega V5280) and 20mM HEPES. Droplet stability was maintained with the aid of the CellenONE's humidifier and slide cooling system, set to 75% relative humidity and one degree below the due point.

The remaining single-cell suspension was used to prepare bulk digest for spectral library generation. Briefly, the remaining cells were pelted and resuspended in mass spectrometry grade water at a concentration of 1000 cells per μL. The cells were then lysed by freezing to -80 degrees C and then heating to 90 degrees C for 10 minutes[42]. The lysate was digested overnight at 37 degrees C by adding trypsin and 8.5 pH TEAB buffer to a concentration of 20 ng/μL and 100mM, respectively.

The next day, the nPOP single-cell samples were labeled with 20 nL of mTRAQ reagents dissolved in 100% DMSO at a concentration of 1/60th unit per μL. Samples were pooled using the CellenONE in a 50%/50% solution of acetonitrile and water and dispensed into a 384 well PCR plate, dried down in a speed vac and stored at -20C for later injection for LC/MS analysis. Bulk samples were labeled by adding 20 μL of mTRAQ d0 label at a concentration of 1/20th unit per μL to 40 μL of digested bulk peptide at a concentration of 1000 cells of digested peptides per μL. Samples were incubated for 1 hour and then dried down and resuspended in 40 μL of 0.1 % Formic

acid and 0.02 % DDM for LC-MS/MS injection.

LC-MS/MS data acquisition

All samples were separated via online nLC on a Vanquish Neo UHPLC using a 25cm x 75 μl
IonOpticks Aurora Series UHPLC column (AUR2-25075C18A). Single-cell sets resting in the
wells of a 384-multiwell plate were re-suspended in 1 μl of 0.1 % Formic acid, 0.02 % DDM for
injection. All samples were run on a 30 minute total length run to run method with 15 minutes
of active gradient. 0.1 % formic acid in water was used for Buffer A and a 20% solution of 0.1%
formic acid mixed in 80 % acetonitrile was used for buffer B. A constant flow rate of 200nl/min
was used throughout sample loading and separation. Samples were loaded onto the column for 20
minutes. The gradient started at 6 % buffer B and ramped to 38% B buffer over 20 minutes. The
gradient was then ramped to 95% B buffer over 2 minutes and stayed at that level for 3 minutes
to remove excess peptide and protein residue from column. The gradient then dropped to 1% B
buffer over 0.1 minutes and stayed at that level for 4.9 minutes for the re-equilibration.

All samples were analyzed on a timsTOF SCP mass spectrometer. Instrument control and
data acquisition were performed via timsControl 3.1. The data acquisition method consisted of 8
PASEF frames with 26 Th MS2 windows (1 Th overlaps). An MS1 scan was taken every 2 PASEF
frames to increase the frequency of precursor sampling[43], resulting in 4 MS1 scans per duty cycle.
The MS1 scan range was 100-1700 m/z, while MS2 scan range was 300-1000 m/z. The 1/K0
range was between 0.64 and 1.20 and collision energy was set at 20eV at 1/K0 = 0.60 and 59eV
at 1/K0 = 1.60, collision RF was set to 2000 Vpp. The ramp and accumulation times were 100
milliseconds and estimated duty cycle time is 1.28 seconds.

sc-mRNA sample preparation, sequencing, and read alignment

The single-cell suspensions were used for barcoding and library preparation with a Chromium
Next GEM Single Cell 3' Kit v3.1 according to the manufacturer's protocol. Libraries were se-
quenced on NovaSeq 6000 S2 with paired-end reads at a depth of 50,000 reads per cell. Reads
were processed and aligned to the mm10 mouse genome reference using CellRanger. We collected

these data and reported them as part of Lin et al.[16]. The data was previously deposited to the GEO:GSE244215.

Interpreting raw mass spectra

All MS runs were searched with DIA-NN v1.9.0[44]. The two bulk runs were first searched using an *in silico* generated spectral library from a murine protein sequence fasta downloaded from Uniprot in September 2024 with mTRAQ d0 specified as a fixed modification with mass of 140.0949630177 on all n-termini and lysine. Additional parameters specified were scan window of 5, Mass accuracy of 15 ppm, and MS1 accuracy of 5 ppm. All other parameters were set to defaults. The resulting search generated a empirical spectral library with 4,056 proteins from 56,321 precursors that was used to analyze the single-cells samples.

Single-cell raw files were searched with match between runs in batches of roughly 600 raw files grouped by the days in which the samples were prepared. DIA-NN settings were the same as for the bulk samples with the addition of the peak-translation and channels arguments for plexDIA. Four channels were specified for mTRAQ d0 and light lysine, mTRAQ d0 and heavy lysine, mTRAQ d8 and light lysine, mTRAQ d8 and heavy lysine as follows:

1. mTRAQ,0,nK,0:0

2. mTRAQ,1,nK,0:6.020129

3. mTRAQ,2,nK,8.0141988132:8.0141988132

4. mTRAQ,3,nK,8.0141988132:14.0343278132

We also performed an additional search with the intention of identifying missed cleaved peptides that were generated with both heavy and light lysines to assess amino acid recycling. To do so, we customized to our empirical library to duplicate all entries so we could manually specify light and heavy lysines. For peptides with two lysines, 4 different precursors were created, one with both light, one with both heavy, and the two combinations of one light and one heavy. Runs were then searched with mTRAQ d0 as a fixed modification and two channels for peptides labeled with d0 and labeled with d8. The downside to this approach is that the peak translation would not

apply across heavy and light precursors with the same label. Thus the results were only used for comparing the heavy and light distribution for peptides with multiple lysines, subsequently used for determining the rate of amino acid recycling.

Plotting raw mass spectra

To plot the raw mass spectra, we utilized the alphaTims GUI[45] to locate precursor ions at the retention time, M/Z, and ion mobility range for peptide LYSEFLGK as reported by DIA-NN.

SCP pre-processing and batch corrections of abundances

Each single cell protein data set defined by individual nPOP sample preparations was pre-processed using the QuantQC R package[15]. The package first maps relevant meta data to the DIA-NN report file by integrating the dispensing log files from the cellenONE. This includes information such as cell size, label, and run order.

For processing of protein abundances, first, light and heavy intensities for each lysine containing peptides from a given single cell were summed to reflect the total abundance of the peptide. To remove failed cells, total MS intensity summed from all peptides was plotted for all single cells including negative control droplets that received all the reagents but no single cell. An intensity threshold was then set that removed all negative control droplets along with all single cells that had intensities smaller than the negative control with the highest signal. The peptide by single cell matrix was then generated, allowing maximum 5 peptides mapping to the same protein. For proteins with greater than 5 peptides, the most abundant 5 peptides, computed from average intensity across all cells, were chosen. Data was normalized to relative log_2 fold changes via the method described in Khan et al.[11]. Briefly, sample loading normalization was performed using the intersected median across all cells as opposed to that of each cell. Specifically, a reference vector was created from the median value for each peptide across all cells. The median difference between the non missing peptide intensities from each cell and the reference vector were then used to scale all peptide intensities for that cell. Each peptide was then divided by the mean expression across all single cells and the data was log transformed.

Next, to control for variation that occurs across LC runs, a spline was fit to the vector of peptide fold changes and the order to which the MS samples were run. The degrees of freedom for the spline fit were set to the number of single cells with present values divided by 10 with a maximum value of 20. For all spline fits with R^2 greater than 0.1, the variation due to the spline fit was regressed out by subtracting the fit line.

Peptide data was then collapsed to the protein level by taking the median value for all peptides mapping to the same protein within each cell type, resulting in a protein by cell type matrix. To correct for bias due to mTRAQ label used, data was first imputed using KNN with K equal to 3, and then batch corrected using the ComBat R package with the label (d0 or d8) specified as a co-variate.

To generate the final data matrix for all data sets, the data for each sample prep was concatenated together and combat was applied with the sample preparation as a co-variate.

Clustering, dimensionality reduction and cell type assignment

Cell types for mRNA data were annotated previously as described in Lin et al.[16] Louvain clustering and UMAP dimensionality reduction was performed on the final protein by cell data matrix via the Seurat R package. Clustering resolution was set to a value of 0.1. Cell types for each cluster were annotated using established cell type markers as shown in Extended Data Fig. 1a. To validate clustering, the average protein and mRNA log_2 fold changes were correlated for all cell types and showed strong agreement for corresponding cell types, Fig. 1c.

Accounting for missing values in SCP data

To effectively estimate protein abundances at the cell type level, we sought to account for missing data in individual single cells. To do this we first needed to distinguish between missing values due to 1) technical factors such as run to run variability, labeling bias, and variable quantification depth due to cell size variation and 2) reduced abundance due to type specific regulation. This functionality, as described below, is incorporated in the QuantQC package.

To this end, we first formulated a technical effect abundance model that predicted the abun-

dance of each peptide data point based on LC-MS/MS run order, sample preparation batch, and mass tag label. First, we started with the peptide level raw abundance data across single cells, and normalized the data matrix to log_2 relative fold changes based on the reference vector approach described above. We then fit a spline model to each peptide fold change using the run order as the independent variable. The spline was fit separately for samples with either mTRAQ d0 or d8 labels and for each sample preparation batch. Again, the degrees of freedom for the spline was set to floor of the number of data points divided by 10, with a max value of 20. We then used this model to predict abundances for all present and missing peptide data points.

Next, we formulated two different missingness models. Each model was trained to predict whether a peptide data point is present or missing by fitting a logistic regression based on a number of features. The first model predicted missingness due to technical factors by utilizing 1) the abundance prediction from the technical effect abundance model and 2) the cell size. The second model included 3) the assigned cell type as a feature to capture missingness due to cell type specific regulation.

The difference in number predicted missing values between the technical only and cell type logistic regression models was taken as the number of single cells missing due to biological reasons, i.e., the peptide abundance is below the detection limit of the measurements. To better estimate the average protein abundance in a cell type in the presence of biological missingness, the values below the limited of detection were assumed to equal the minimum value measured in this cell type. The cell type averages were then computed including the included minimum values. This correction resulted in improved agreement of fold changes estimated across cell types when compared to scRNA-seq, Extended Data Fig. 2f-h.

Estimating clearance rates in vivo

To estimate the change in the fraction of light lysine in the unbound lysine pool over time, $\gamma(t)$, we first calculated the fraction of heavy lysine in the free pool at our two time points. To do this, we utilized the average ratio of partially labeled to fully labeled peptides as done by Jovanovic et al.[2]. This ratio was computed as the average across all miscleaved peptide from all single cells within each cell type. We then fit an equation of the form $\gamma(t) = 1 - 0.5e^{-at} - 0.5e^{-bt}$ to estimate the

change in available light lysine over time as previously developed and used in ref.[19] for each cell type individually. Cell types were also fit separately within each age group. Because of the loss of old mice from the 10 day experiment, we estimated the recycling ratio for the old mice at 10 days to be equivalent to the difference between the old and young mice at day 5, Extended Data Fig. 3b.

Having solved for the change in the free lysine pool over time, we then formulated the change in any given light peptide over time $\frac{dL_i}{dt}$ via the equation:

$$\frac{dL_i}{dt} = k_{s,i}\gamma(t) - k_{r,i}L(t)$$

We then integrate this equation to solve for $L_i(t)$ for the initial condition $L(0) = L_0$ where L_0 is defined as the sum of heavy and light intensities under the steady state assumption that overall protein concentration is not changing over time. Integrating and further substituting $k_{s,i}$ via the steady-state equality $L_0 = \frac{k_{s,i}}{k_{r,i}}$. This results in the following equation:

$$L(t) = k_{r,i}L_0 \left(\frac{0.5}{k_{r,i}-a}e^{-at} + \frac{0.5}{k_{r,i}-b}e^{-bt} \right) + \left(L_0 - k_{r,i}L_0 \left(\frac{0.5}{k_{r,i}-a} + \frac{0.5}{k_{r,i}-b} \right) \right) e^{-k_{r,i}t}$$

The arithmetic was performed in Mathematica to avoid manual integration errors. Lastly, this equation was then solved for via a non-linear solver to find $k_{d,i}$ for each peptide from each single cell, with equation parameters for $\gamma(t)$ depending on the cell type and mouse age.

Computing gene regulatory parameters

The following derivations are more comprehensively derived in Baum *et al.*[21]. Briefly, in cells that are not growing, under the steady state assumption protein concentrations can be formulated as a balance of rate of synthesis and degradation.

$$p_i = \frac{k_{s,i}}{k_{d_i}}$$

In cells that are growing, the steady state model can be maintained under the assumption that protein concentrations remain constant as cells double the amount of all proteins at an equal rate.

However, under this regime, we need to introduce an additional term to reflect protein clearance due not only to degradation, but to dilution as well.

$$p_i = \frac{k_{s,i}}{k_{d,i} + k_g}$$

The dilution rate k_g is constant for all proteins and is inversely proportional to the rate of cell division, T_{CC}, $k_g = log(2)/T_{CC}$. For both models, we define the overall amount of protein clearance from degradation and dilution is $k_{r_i} = k_{d_i} + k_g$, with $k_g = 0$ in the absence of growth.

To determine the average cell division time across a population of single cells, we leverage the fact that histones degrade very slowly, but double and partition themselves roughly evenly upon cell division. Thus, the distribution of heavy over light histone abundances takes on a discrete nature where it is possible to asses the number of times a cell had divided. We then use the following equation to determine the doubling time, $N(t) = N_0 2^{t/T_{CC}}$ where t is the pulse duration, $N(t)$ is the number of single cells in the population at the pulse duration t, and N_0 is the initial number of cells. N_0 can be computed as the following:

$N_0 = \sum_{d=0}^{D} \frac{n_d}{2^d}$

where n_d are the number of cells that have divided d times. Error bars are the standard deviation of T_{CC} estimates across the 4 sample preparations.

Lastly, the synthesis rate can be further decomposed into the number of mRNA copies for the *ith* gene in the cell, $mRNA_i$, multiplied by the average rate the mRNA is translated.

$$k_{s,i} = k_{t,i} \cdot mRNA_i$$

Resulting in the final expression defining the regulation of the *ith* gene:

$$p_i = \frac{k_{t,i} \cdot mRNA_i}{k_{d,i} + k_g}$$

Handling measurement noise for absolute abundance control

To handle measurement noise when estimating the extent that mRNA abundance, translation, and protein clearance influence absolute protein abundance, we utilized reliability measurements from

average estimates from correlating random subsets of single cells. For two random subsets of cells, we required at least 5 observed values to contribute to each average before computing the reliability. From this approach, we are underpowered to estimate the influence of systematic biases contributing to absolute copy number of proteins and mRNAs. Phenomena such as ionization and digestion bias for protein measurements, and cell lysis, transcript capturing and PCR biases for mRNA measurements might bias absolute abundance estimates across different proteins and transcripts. Thus, estimates of reliability for absolute protein and mRNA abundances were multiplied by 0.7 to reflect these systematic biases,[4,46]. This adjustment reflects comparison of different mRNA sequencing technologies and peptides generated from different protease digests. No further correction was applied to the protein clearance estimates as these biases should cancel out when computing the ratios between heavy and light precursors.

We then used the Spearman correction to correct the percent variance explained by both clearance and mRNA abundance[47]. In the corrected estimates, the translation contributes the remaining variance. In the uncorrected estimates, the raw R^2 of each modality is displayed in Extended Data Fig. 5e. Estimates made from the remaining variance correlate well with those made by regressing the translation rates against protein abundance, $\rho = 0.75$, Extended Data Fig. 5d.

To estimate uncertainty in the variance-explained statistics, we applied a nonparametric bootstrap procedure at the level of genes averages. For each cell type, paired values of the regulatory variable (e.g., clearance rate) and protein abundance were resampled with replacement to the same size as the original dataset. The variance-explained statistic was recalculated for each bootstrap sample, yielding a distribution of bootstrap estimates. We repeated this procedure 1,000 times per cell type and summarized the dispersion of the bootstrap distribution by its standard deviation, which we report as the bootstrap error for the variance-explained measure.

Estimating relative levels by Bayesian inference

To estimate mRNA abundance, translation rate, protein clearance and relative protein abundance of each protein in each cell type, we relied on replicate estimates from peptides mapping to the same protein, individual single cells, and measurements made from sample preparation replicates. We formulated the problem using Bayesian inference to estimate measurement noise for individual

quantities in individual cell types.

Specifically, we modeled the log transformed parameters, including relative fold changes compared to the average across cells for protein abundance ($logP$), mRNA abundance ($logM$), translation rate ($logT$), and clearance rate ($logC$). These values were each inferred for for gene g in cell type c via the log transformation of the previously established relationship between the quantities:

$$\log p_{g,c} = \log mRNA_{g,c} + \log \mathrm{k_t}_{g,c} - \log \mathrm{k_r}_{g,c}$$

In our model, the protein abundance, mRNA abundance, and clearance rates were the only observed values. Translation rate is inferred from the remaining difference in protein levels not explained by mRNA abundance and protein clearance.

The observed quantities were modeled using Gaussian likelihoods with standard errors derived from the number of replicate observations. For protein abundance ($\bar{p}_m$) and clearance rates ($\bar{c}_m$), observed quantities were the peptide level cell type average of the relative log_2 fold changes compared to the average value across cell types for one of the four sample preparation replicates.

Protein abundance observations were modeled as:

$$\bar{p}_m \sim \mathcal{N}(logP_{g,c}, \ \sigma = 1/\sqrt{n_{\mathrm{pro}}[m]})$$

where g and c denote the gene and cell type indices for observation m, and $n_{\mathrm{pro}}[m]$ is the number of cells where the peptide was observed in the given replicate. Similarly, observed clearance rates were modeled as:

$$\bar{c}_m \sim \mathcal{N}(logC_{g,c}, \ \sigma = 1/\sqrt{n_{\mathrm{clear}}[m]})$$

For observed mRNA ($\bar{M}_m$) quantities were similarly the cell type average of the relative log_2 fold changes compared to the average fold change within replicate.

$$\bar{M}_m \sim \mathcal{N}(logM_{g,c}, \ \sigma = 1/\sqrt{n_{\mathrm{rna}}[m]})$$

Here $n_{\mathrm{rna}}[m]$ was the number of non-zero observations of the transcript in the given cell type and replicate.

We placed zero-centered Gaussian priors with standard deviation that reflected our prior certainty for each quantity:

$$logM_{g,c} \sim \mathcal{N}(0,\ 5)$$

$$logC_{g,c} \sim \mathcal{N}(0,\ 3)$$

$$logT_{g,c} \sim \mathcal{N}(0,\ 1)$$

This model enabled us to propagate measurement noise and quantify uncertainty in the estimates of gene expression regulation for each gene-cell type pair, allowing robust decomposition of protein abundance into transcriptional, translational, and degradative components.

Correlation analysis across cell types

To determine the extent to which each protein is regulated by mRNA abundance, translation, and clearance, we correlated proteins abundances to the extent of regulation by each mode across cell types. So that a positive correlations would reflect strong regulation for each mode, we utilized clearance half life, $log(2)/K_r$, instead of clearance rate. To determine functional groups of proteins primarily regulated by a single mode of regulation, we performed a gene set enrichment analysis on the correlation distribution for each mode of regulation. Within each mode, we compared the distribution correlation for genes in each gene set with at least 4 proteins present, to the remaining distribution of correlations using a t-test. We then corrected the distribution of p-values within each mode of regulation for multiple hypothesis testing with the Benjamini-Hochberg (BH) method.

Cell type specific gene regulation analysis

To plot the regulatory influences for gene products that were predominantly regulated by a single mode of gene expression in individual cell type, we plotted in Fig. 3 all genes where the contribution from a single mode was greater than the contribution of the other two. We clustered data individually for mRNA, translation, and clearance and plotted the genes passing this threshold stacked for each modality, Fig. 3f.

To identify functional sets of proteins that were over represented in the genes primarily reg-

ulated by a single mode of gene expression, we utilized the compareCluster function from the R package clusterprofiler. Briefly, this utilizes a hyper-geometric test to compare the genes that passed the threshold to the background set of all genes where mRNA, translation, and clearance were quantified. All plotted terms met a significance threshold of FDR of 5 % across at least one of the 6 cell types.

Single cell protein-mRNA-clearance covariation analysis

We plotted all pairwise correlations for intersected sets of proteins and mRNAs where the proteins had over 100 pairwise observations to robustly support the computed correlation. Curate a list of protein pairs that that directly interact we took all pairwise protein combinations from complexes in the Corum protein data base (release 5.0) as well as including all reported interactions in the STRINGdb with confidence score above 0.7.

To identify protein or mRNA specific correlation modules, we filtered for proteins that contained at least one pairwise correlation above a certain threshold. We first set the threshold at 0.3 for proteins and then plotted the heatmap via hierarchical clusters. We then extracted the clusters at a resolution of the top 7 clusters. Lastly we looked for GO term over enrichment for proteins in these clusters compared to the background set using the clusterProfiler r package (Release 3.21)[48]. Several terms were enriched for each of the 7 clusters and the most significant GO term was displayed below the heatmap. The analogous procedure was applied to the mRNA data but few transcripts displayed pairwise correlations above 0.3. Reducing the threshold to 0.20, we identified 5 clusters with greater than 10 transcripts. However, only 3 of the 5 clusters contained GO terms with significant over enrichment.

To quantify the contribution of protein clearance to the differences between the corresponding protein-protein and mRNA-mRNA correlations, we started by estimating protein regulation by clearance, r_c; r_c was estimated by correlating protein abundances across single cells to the paired clearance rates across the same single cells. We then plotted the difference between pairwise correlations for the same pair of gene products at the mRNA and protein level versus the average r_c for the protein pair. The resulting correlation, $\rho = -0.44$, suggests significant contribution of clearance to the difference in correlations.

We then created a linear model including as additional explanatory variables the average absolute half life of the two proteins as well as the average raw counts of each transcript. This increased the explanatory power of the model, R = 0.55.

Single-cell functional covariation analysis

We test for enriched correlations in functional groups of genes within cell type at the protein and mRNA levels. We performed this analysis in the space of intersected proteins and mRNA quantified in each cell type. For each GO term with at least 4 genes, we performed a t-test between the distribution of all pairwise genes from the go term and the background distributions of pairwise correlations for all other proteins or transcripts. We then corrected the distribution of p-values for multiple hypothesis testing with the Benjamini-Hochberg (BH) method. We plotted all terms which were significant at 1% FDR in at least one cell type at the protein or mRNA level.

Fibroblast vs basal covariation analysis

To compare within cell type covariation patterns across fibroblast and basal cells, we first computed pairwise correlations for all proteins across single cells within a cell type. For each cell type, we defined correlation vector for each protein as the vector of correlations to all other proteins. To compare covariation patterns across cell types, we correlated these correlation vectors for shared proteins between basal and fibroblast cells. A null distribution was computed by randomizing abundances across cells and proteins for each cell type and computing the same procedure.

We then took a threshold of all proteins with a correlation vector similarity of Pearson correlation above 0.5 and denoted this the set of proteins with common covariation patterns as set 1. The similar correlation structure of pairwise proteins can be observed via the heatmap, Fig. 5c. Clusters were annotated for functional groups with the procedure described for the protein-mRNA comparison, but was only applied on the basal cells due to the existing high correlation similarity between the two cell types. All proteins with a correlation of correlation vector below 0.1 were denoted as set 2 proteins. Differences in the covariation between set 1 and set 2 proteins across cell types were identified by clustering the correlation matrix defined by correlations between all

set 1 and set 2 proteins in basal cells, Fig. 5d.

References

1. Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473,** 337–342 (2011).

2. Jovanovic, M. *et al.* Dynamic profiling of the protein life cycle in response to pathogens. *Science* **347,** 1259038 (2015).

3. Buccitelli, C. & Selbach, M. mRNAs, proteins and the emerging principles of gene expression control. *Nature Reviews Genetics* **21,** 630–644 (2020).

4. Leduc, A. & Slavov, N. Protein degradation and growth dependent dilution substantially shape mammalian proteomes. *bioRxiv,* 2025–02. https://doi.org/10.1101/2025.02.10.637566 (2025).

5. Li, W. *et al.* Turnover atlas of proteome and phosphoproteome across mouse tissues and brain regions. *Cell* **188,** 2267–2287 (2025).

6. Zhang, T. *et al.* Proteome-wide modulation of degradation dynamics in response to growth arrest. *Proceedings of the National Academy of Sciences* **114,** E10329–E10338 (2017).

7. Chapman, A. R. *et al.* Correlated gene modules uncovered by high-precision single-cell transcriptomics. *Proceedings of the National Academy of Sciences* **119,** e2206938119 (2022).

8. Tarbier, M. *et al.* Nuclear gene proximity and protein interactions shape transcript covariations in mammalian single cells. *Nature communications* **11,** 5445 (2020).

9. Slavov, N. Learning from natural variation across the proteomes of single cells. *PLOS Biology* **20,** 1–4. https://doi.org/10.1371/journal.pbio.3001512 (Jan. 2022).

10. Derks, J. *et al.* Increasing the throughput of sensitive proteomics by plexDIA. *Nature Biotechnology.* https://doi.org/10.1038/s41587-022-01389-w (2022).

11. Khan, S. *et al.* Inferring post-transcriptional regulation within and across cell types in human testis. *bioRxiv.* https://doi.org/10.1101/2024.10.08.617313 (2024).

12. Sabatier, P. *et al.* Global analysis of protein turnover dynamics in single cells. *Cell* **188,** 2433–2450 (2025).

13. Borteçen, T., Müller, T. & Krijgsveld, J. An integrated workflow for quantitative analysis of the newly synthesized proteome. *Nature Communications* **14,** 8237 (2023).

14. Leduc, A., Huffman, R. G., Cantlon, J., Khan, S. & Slavov, N. Exploring functional protein covariation across single cells using nPOP. *Genome Biology* **23,** 261. `https://doi.org/10.1186/s13059-022-02817-5` (2022).

15. Leduc, A., Khoury, L., Cantlon, J., Khan, S. & Slavov, N. Massively parallel sample preparation for multiplexed single-cell proteomics using nPOP. *Nature Protocols,* 1–27. `https://doi.org/10.1038/s41596-024-01033-8` (2024).

16. Lin, B. *et al.* Airway hillocks are injury-resistant reservoirs of unique plastic stem cells. *Nature* **629,** 869–877 (2024).

17. Märtens, A. *et al.* Instrumental drift in untargeted metabolomics: Optimizing data quality with intrastudy QC samples. *Metabolites* **13,** 665 (2023).

18. Vanderaa, C. & Gatto, L. scplainer: using linear models to understand mass spectrometry-based single-cell proteomics data. *Genome Biology* **26,** 237 (2025).

19. Fornasiero, E. F. *et al.* Precisely measured protein lifetimes in the mouse brain reveal differences across tissues and subcellular fractions. *Nature communications* **9,** 4230 (2018).

20. Mathieson, T. *et al.* Systematic analysis of protein turnover in primary cells. *Nature communications* **9,** 689 (2018).

21. Baum, K., Schuchhardt, J., Wolf, J. & Busse, D. Of gene expression and cell division time: a mathematical framework for advanced differential gene expression and data analysis. *Cell Systems* **9,** 569–579 (2019).

22. Gatto, L. *et al.* Initial recommendations for performing, benchmarking, and reporting single-cell proteomics experiments. *Nat. Methods* **20,** 375–386. `https://doi.org/10.1038/s41592-023-01785-3` (2023).

23. Shi, J. & Du, G. Metabolic reprogramming of glycolysis favors cartilage progenitor cells rejuvenation. *Joint Bone Spine* **91,** 105634 (2024).

24. Slavov, N. & Dawson, K. A. Correlation signature of the macroscopic states of the gene regulatory network in cancer. *Proceedings of the National Academy of Sciences* **106,** 4079–4084. eprint: http://www.pnas.org/content/106/11/4079.full.pdf+html (2009).

25. Specht, H. *et al.* Single-cell proteomic and transcriptomic analysis of macrophage heterogeneity using SCoPE2. *Genome Biology* **22** (2021).

26. Iyer-Biswas, S. *et al.* Scaling laws governing stochastic growth and division of single bacterial cells. *Proceedings of the National Academy of Sciences* **111,** 15912–15917 (2014).

27. Slavov, N. & Botstein, D. Coupling among growth rate response, metabolic cycle, and cell division cycle in yeast. *Molecular Biology of the Cell* **22,** 1997–2009 (2011).

28. Derks, J. & Slavov, N. Strategies for increasing the depth and throughput of protein analysis by plexDIA. *Journal of Proteome Research* **22,** 697–705. https://doi.org/10.1021/acs.jproteome.2c00721 (2023).

29. Specht, H. *et al.* PSMtags improve peptide sequencing and throughput in sensitive proteomics. *bioRxiv,* 2025–05 (2025).

30. McDonnell, K. *et al.* JMod: Joint modeling of mass spectra for empowering multiplexed DIA proteomics. *bioRxiv.* https://www.biorxiv.org/content/early/2025/05/27/2025.05.22.655512 (2025).

31. Budnik, B., Levy, E., Harmange, G. & Slavov, N. SCoPE-MS: mass-spectrometry of single mammalian cells quantifies proteome heterogeneity during cell differentiation. *Genome Biology* **19,** 161. https://doi.org/10.1186/s13059-018-1547-5 (2018).

32. Johnston, S. M. *et al.* Rapid, One-Step Sample Processing for Label-Free Single-Cell Proteomics. en. *J. Am. Soc. Mass Spectrom.* ISSN: 1044-0305, 1879-1123. https://pubs.acs.org/doi/10.1021/jasms.3c00159 (July 2023).

33. Huffman, R. G. *et al.* Prioritized mass spectrometry increases the depth, sensitivity and data completeness of single-cell proteomics. *Nat. Methods.* http://dx.doi.org/10.1038/s41592-023-01830-1 (2023).

34. Sinn, L. R. *et al.* Slice-PASEF: Maximising Ion Utilisation in LC-MS Proteomics. *bioRxiv* (2025).

35. Furtwängler, B. *et al.* Mapping early human blood cell differentiation using single-cell proteomics and transcriptomics. *Science,* eadr8785 (2025).

36. Karagach, S., Smollich, J., Atrakchi, O., Mohan, V. & Geiger, T. High throughput single-cell proteomics of in vivo cells. *Molecular & Cellular Proteomics,* 101018 (2025).

37. Leduc, A., Xu, Y., Shipkovenska, G., Dou, Z. & Slavov, N. Limiting the impact of protein leakage in single-cell proteomics. *Nature Communications* **16,** 4169 (2025).

38. Khan, S., Conover, R., Asthagiri, A. R. & Slavov, N. Dynamics of single-cell protein co-variation during epithelial-mesenchymal transition. *Journal of Proteome Research.* https://pubs.acs.org/doi/10.1021/acs.jproteome.4c00277 (2024).

39. Derks, J. *et al.* Single-nucleus proteomics identifies regulators of protein transport. *bioRxiv.* https://doi.org/10.1101/2024.06.17.599449 (2024).

40. Raj, A. & van Oudenaarden, A. Single-molecule approaches to stochastic gene expression. *Annual review of biophysics* **38,** 255–270 (2009).

41. Slavov, N. Single-Cell Proteomic Technologies: Tools in the quest for principles. arXiv: 2506.18198 [q-bio.QM]. https://arxiv.org/abs/2506.18198 (2025).

42. Specht, H. *et al.* Automated sample preparation for high-throughput single-cell proteomics. *bioRxiv 10.1101/399774.* https://doi.org/10.1101/399774 (2018).

43. Wallmann, G., Leduc, A. & Slavov, N. Data-Driven Optimization of DIA Mass-Spectrometry by DO-MS. *Journal of Proteome Research.* https://doi.org/10.1101/2023.02.02.526809 (2023).

44. Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. *Nature methods* **17,** 41–44 (2020).

45. Willems, S., Voytik, E., Skowronek, P., Strauss, M. T. & Mann, M. AlphaTims: indexing trapped ion mobility spectrometry–TOF data for fast and easy accession and visualization. *Molecular & Cellular Proteomics* **20,** 100149 (2021).

46. Hornung, B. V. *et al.* Comparison of single cell transcriptome sequencing methods: of mice and men. *Genes* **14,** 2226 (2023).

47. Franks, A., Airoldi, E. & Slavov, N. Post-transcriptional regulation across human tissues. *PLoS computational biology* **13,** e1005535. https://doi.org/10.1371/journal.pcbi.1007082 (2017).

48. Yu, G. Thirteen years of clusterProfiler. *The Innovation* **5** (2024).

Extended Data Figures

Extended Data Fig. 1 | Validating the alignment of the mRNA and protein cell-type clusters a,, Abundance of cell-type specific markers for each of the 7 cell types. The protein levels are plotted as log_2 concentrations, and the mRNA levels as raw counts. **b**, Scatter plots for relative protein levels in each cell type. The levels are plotted as log_2 ratios relative to the mean across all cells types.

Extended Data Fig. 2 | Processing and normalization of the single-cell proteomics data a, Visualization of bias in peptide abundance based on deviations that occur systematically across runs and spline fit to data. R^2 of the spline fit for all measured peptides. Regression of trend was applied to peptides with an $R^2 > 0.1$. The abundance of the peptide VISLSGEHSIIGR2 does not vary with run order after correction. **b**, Fraction of missing peptide data points in a cell is a function of total MS signal measured or similarly the cells and the cell size. **c**, Results of a logistic regression model fit to predict intensity based on retention time and label of the single cell show moderate predictive power on determining whether a peptide data point is missing or observed. **d**, The ROC curve for a logistic regression model fit to predict missingness of peptide data points based on technical factors (cell size, retention time and label). Including the cell-type identity improves the predictions. **e**, For the least abundant proteins in chondrocytes, including cell-type information increased the prediction accuracy for missing values. **f**, Log_2 protein and mRNA fold changes for chondrocytes plotted before and after missing data correction. **g**, Slopes of mRNA protein fold changes across gene products for all cell types plotted before and after correction. **h**, Similar to panel g, but the slopes are estimated across cell types.

Extended Data Fig. 3 | Adjusting clearance for amino acid recycling a, At each time point, three data points are sampled, the heavy amino acid intensity, the light amino acid intensity, and the fraction of heavy lysine available as predicted from miscleaved peptides. **b**, Free heavy lysine pool computed from peptides with missed cleavages as the ratio (HL/HH)/(HL/HH+2). Data points represent average estimates across all peptides for a given single cell. Fit line to the average data points within cell type for 5 and 10 days for the sum of two exponential for basal cells. Due to the loss of old mice at the 10 day time point, the 10 day time point was estimated using the same difference between old and young at the 5 day time point. **c**, Adjusted clearance half life distribution computed before and after recycling correction. **d**, Slope between 5 and 10 day cells computed before and after the adjustment for amino acid recycling. **e**, Basal cell clearance rate for all proteins plotted for 24 month and 2 month mice shows slope of 1 and lower average clearance rate of 0.04 in older mice. **f**, Clearance rate half life distributions for all cell types. **g**, Correlation between average clearance rate distributions for all cell types. The diagonal represents the correlation of average clarence rates for the same cell type computed from non-overlapping subsets of single cells.

Extended Data Fig. 4 | Quantitative accuracy and reliability of mRNA abundance, protein abundance, and clearance rate estimates a, Quality plots for single-cell mRNA data show the number of unique transcripts detected on average per cell and the number of unique UMIs detected. **b**, The log transformed sum of all protein intensity per cell plotted against the cell volume show that total protein is highly proportional to cell size. **a**, Relative protein log_2 fold changes for two peptides from protein Tgm2 across single cells. Correlations of protein fold changes for the to 5 most abundant peptides have an average Pearson correlation of 0.88. **d**, The average correlation for all proteins is plotted against the average protein fold change across single cells for each data set. More variable proteins with higher average fold change correlate more highly. **e**, The average reliability for each modality calculated by taking averages from two random subsets of single cells and plotting the correlation between subsets.

Extended Data Fig. 5 | Cell growth modulates regulation of absolute abundance a, Despite the difference in the slope of clearance rates between basal cells and chondrocytes (shown in Fig. 2), the degradation rate slope is 0.95. This suggests that the shallow slope is due to the influence of cell growth. **b**, Scatter plot of clearance rates and absolute protein abundance in basal cells shows R^2 of 0.26. **c**, Scatter plot of mRNA copies and and absolute protein abundance secretory cells show R^2 of 0.51. **d**, Scatter plot translation contribution estimates computed from the squared Pearson correlation between translation rates and protein abundance, and cell growth rate still shows strong positive trend.

Extended Data Fig. 6 | The modes controlling relative protein levels can be shared across cell types or cell-type specific. a, Joint distributions of correlations. The top panel shows distributions of correlations between protein and mRNA abundance versus bins of the correlations between protein abundance and protein clearance. The bottom panel shows distributions of correlations between protein and mRNA abundance versus bins of the correlations between protein abundance and translation rate. **c**, Examples of synergistic regulation of St13 across cell types. **c**, A heatmap showing synergistic regulation by mRNA abundance and translation in basal cells, and translation and clearance in chondrocytes.

Extended Data Fig. 7 | Single-cell mRNA and protein covariation a, A comparison of the dynamic range of mRNA and protein abundance within basal cells indicates larger variation for proteins. The dynamic range is estimated by the standard deviation of log_2 fold changes relative to the mean for protein and mRNA levels. **b**, Correlations between directly interacting proteins within each cell type for proteins that have over 100 pairwise observations and the corresponding mRNAs. **c**, mRNA dominant correlation modules lack enriched GO terms and exhibit low correspondence to protein data. **d**, Prediction of the difference between protein-protein and mRNA-mRNA correlations by a linear model that incorporates the protein clearance rates, the correlation between the relative clearance rates and proteins levels, and the average mRNA counts. **e**, A heat map displaying average correlations for the full list of GO terms that were significant in at least one cell type.

Extended Data Fig. 8 │ Analyzing the differences between mRNA and protein correlation vectors a, mRNA-mRNA correlations corresponding to the protein-protein correlations displayed in Fig. 5d. **b**, The protein abundance, protein clearance, or mRNA abundance for gene products from the clusters in Fig. 5d were averaged for each cluster, and the cluster averages correlated. The protein clusters have the largest correlations that exhibit qualitative similarity with both clearances and mRNA correlations, suggesting contributions of these processes to the observed cell-type specific changes in protein covariation.