Sexual dimorphism in pollen foraging and sensory traits in *Heliconius* butterflies

José Borrero[1], David F. Rivas-Sánchez[2], Daniel Shane Wright[1], Caroline Nicole Bacquet[3], Stephen H. Montgomery[2], Alexander Keller[4] & Richard M. Merrill[1]

Corresponding authors: jose.borrero@lmu.de (J.B.) & merrill@bio.lmu.de (R.M.M)

[1] *Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany*

[2] *School of Biological Science, University of Bristol, Bristol, United Kingdom*

[3] *Universidad Regional Amazónica de Ikiam, Tena, Ecuador*

[4] *Cellular and Organismic Networks, Faculty of Biology, LMU Munich, Germany*

Abstract

Sexual dimorphism in foraging behaviour is widespread in insects and may arise from differences in nutritional demands, sensory systems, or cognition. *Heliconius* pollen-feeding is an evolutionary innovation among butterflies that supports extended lifespans and sustained reproduction. However, how foraging behaviour varies between the sexes and how it relates to sexually dimorphic traits remains poorly understood. We investigated sex-specific foraging strategies in wild *Heliconius himera*, a highland specialist from southern Ecuador, using field surveys and DNA metabarcoding. Females carried more pollen than males, consistent with higher nutritional demands, yet we observed no differences in plant richness and composition. This indicates that sex differences reflect effort in foraging behaviour rather than shifts in plant choice. Gut samples revealed greater pollen diversity and a more consistent community profile than proboscis samples, suggesting they better capture cumulative foraging history. We also quantified sexually dimorphic sensory traits and found that males had larger eyes and more ommatidia, whereas females had larger mushroom bodies. While the functional significance of these differences remains unclear, these patterns are consistent with sexual dimorphism reported across *Heliconius* and suggest males and females may be under divergent selective

27   pressures. Our findings highlight how sex-specific foraging differences can arise from differential

28   effort on shared floral resources and co-occur with divergent sensory and neural investment,

29   offering insights into the ecological basis of intraspecific variation in pollen use.

Introduction

Sexual dimorphism is widespread across animals and often involves divergence in morphology, physiology, and behaviour. Among insects, sex differences in foraging behaviour may arise from distinct dietary needs or sensory biases. Many species exhibit sex-specific specializations in visual and olfactory systems that support different ecological roles, such as mate location in males and resource acquisition in females (Smith et al. 2019). For instance, in the oil bee *Macropis fulvipes*, both sexes forage on the same flowers, but males rely more on visual cues and only collect pollen, while females show a stronger preference for olfactory cues and collect both pollen and floral oils (Dötterl et al. 2011). Similarly, in the fly *Bibio rufiventris*, males and females differ in their responses to floral volatiles over the course of the day, reflecting sex-specific attraction to scent cues (Zhu et al. 2025). Despite such examples, the ecological and sensory drivers of sex-specific foraging behaviour remain poorly understood in most insect taxa.

Pollen-feeding is a widespread and ecologically important behaviour in insects, particularly among bees, flies, and beetles, where it plays a central role in adult nutrition, reproduction, and pollination (Romeis et al. 2005). These lineages have evolved diverse morphological and behavioural adaptations to collect and digest nutrient-rich pollen (Roulston and Cane 2000; Ruedenauer et al. 2019). Unlike most butterflies that feed only on nectar, *Heliconius* uniquely show a derived pollen-feeding behaviour, a strategy central to their nutrition, reproduction, and mutualism with the plants they pollinate (Gilbert 1972; Young and Montgomery 2020). Sex differences in pollen feeding have been documented in some *Heliconius* species (Boggs et al. 1981; Cardoso 2001; Mendoza-Cuenca and Macías-Ordóñez 2005), but how these differences align with sexually dimorphic sensory traits, and whether they involve shifts in pollen load versus composition, remains understudied.

Insects that feed on pollen gain access to amino acids and lipids otherwise scarce in nectar

(Romeis et al. 2005). In *Heliconius*, pollen feeding has been linked to enlarged mushroom

bodies that support sensory integration and learning, possibly facilitating this unique behaviour

(Sivinski 1989; Montgomery et al. 2016). Both sexes benefit from pollen-derived nutrients that

extend lifespan and sustain reproduction (Dunlap-Pianka et al. 1977; Pinheiro de Castro et al.

2020), but the direct benefits differ: in females, amino acids and lipids directly support

oogenesis and sustained egg production, and in older females, pollen-feeding helps maintain

adult cyanogenic defences that contribute to Müllerian mimicry (O'Brien et al. 2003; O'Brien et

al. 2004; Pinheiro de Castro et al. 2025). In males direct benefits appear more limited, where it

mainly supports spermatophore production and mating effort (Boggs and Gilbert 1979; Boggs et

al. 1981). These differences in direct benefits likely drive a female bias in pollen foraging.

Multiple studies have found evidence for sex differences in pollen feeding in *Heliconius*. Field

observations of *H. charithonia*, a species within the *erato/sara/sapho* clade, have revealed sex-

specific pollen use: females are more often recorded on *Hamelia patens* plants, while males

favor *Lantana camara* (Mendoza-Cuenca and Macías-Ordóñez 2005). Similarly, broader studies

across *Heliconius* have documented larger pollen loads and longer foraging duration in females

than males (Boggs et al. 1981; Gilbert 1991; Cardoso 2001). These patterns mirror broader

trends in other Lepidoptera, where females actively seek out nectar with higher amino acid

content than males (Alm et al. 1990; Erhardt and Rusterholz 1998; Mevi☐Schütz and Erhardt

2005).

In *Heliconius*, differences in foraging may be influenced by sexually dimorphic visual systems.

Unlike most butterflies, *Heliconius* possess two UV-sensitive opsins, and in the erato/sara/sapho

clade one of these (UVRh1) is linked to the sex chromosome (W) and expressed only in

females, enabling UV wavelength discrimination in females, which may enhance floral detection

capabilities (Finkbeiner and Briscoe 2021; Chakraborty et al. 2023). However, the ecological

relevance remains unclear; for instance, UV sensitivity does not appear to affect oviposition in

H. erato and *H. himera* (Borrero et al. 2023). Beyond opsin expression, other visual traits such

as facet number, facet diameter, and corneal area shape acuity and sensitivity and can differ

between sexes (Land 1997; Wright et al. 2023). As such, integrated studies of foraging

behaviours and sensory traits may help elucidate the basis of sex-specific pollen foraging.

Heliconius himera is a highland specialist inhabiting seasonally dry forests of southern Ecuador

(~1500 m asl). Compared with its lowland relatives, it occupies more open habitats with higher

light intensity, stronger temperature fluctuations, and distinct floral communities, conditions that

likely shape both resource availability and foraging strategies (Descimon and Mast 1984;

Jiggins et al. 1996; Dell'Aglio, Mena, et al. 2022). Reflecting adaptation to this environment, *H.*

himera differs from lowland *H. erato* in brain morphology and sensory processing, investing

more in antennal lobes and less in visual neuropils (Montgomery and Merrill 2017). When visual

and olfactory cues conflict, *H. himera* relies more heavily on olfactory input than *H. erato*

(Dell'Aglio, McMillan, et al. 2022), a pattern also seen in other Lepidoptera with variable neural

investment (Stöckl et al. 2016; Borrero et al. 2024). As an incipient species within the

erato/sara/sapho clade, *H. himera* likely shares the sexually dimorphic opsin expression and

vision reported in related taxa (McCulloch et al. 2017; Chakraborty et al. 2023). This

combination of distinct sensory weighting and potential sexually dimorphic vision makes *H.*

himera a compelling model for exploring how environmental adaptation and sex-specific

sensory traits shape foraging behaviour.

Despite its ecological importance, pollen use in *Heliconius* remains incompletely understood.

Traditional methods such as field observations and palynological keys have revealed diverse

diets dominated by *Psychotria*, *Psiguria*, and *Lantana* (Boggs et al. 1981; Cardoso 2001;

Estrada and Jiggins 2002; Mendoza-Cuenca and Macías-Ordóñez 2005). However, these

methods are slow, expertise-dependent, and limited in taxonomic resolution. DNA

metabarcoding provides a powerful alternative: targeting regions such as ITS2, enables high-

throughput, species-level identification of mixed pollen samples (Keller et al. 2015; Bell et al.

2016). Already widely applied in bee pollination studies (Piko et al. 2021; Wilson et al. 2021;

Martins et al. 2023), this method can open new perspectives on *Heliconius* foraging ecology and

sex-specific pollen use.

In this study, we integrate field surveys, DNA metabarcoding, and morphological analyses to

investigate sex differences in foraging behaviour in *H. himera*. Specifically, we tested the

hypothesis that sexes differ in the amount and diversity of pollen collected, reflecting distinct

nutritional demands or foraging effort. We also quantified eye and brain morphology to

document sexually dimorphic sensory and neural traits. This study provides one of the first

molecular characterizations of pollen resource use in *Heliconius* and highlights how sex-specific

differences in pollen collection can co-occur with divergent sensory and neural investment.

Materials & Methods:

Study sites:

Field surveys were conducted at five sites in Loja Province, Ecuador, around the city of

Vilcabamba (4°15′S, 79°13′W), at elevations of 1500–1700 m (Fig. 1A–B). All sites are

characterized by thorn scrub dry forest dominated by *Acacia macracantha* and xerophytic

vegetation (Jiggins et al., 1996). Rumi Wilco is a 40-ha private reserve containing ~700 plant

species (Jorgensen & Leon-Yanez, 1999), while Cerro Mandango is a nearby mountain site

dominated by Acacia. The other sites (Quebrada Puliche, Agua de Hierro, and San Pedro) occur

Figure 1*: **Sampling region and pollen foraging behavior in H. himera.** (**A**) Study area in southern Ecuador. (**B**) Overview map showing the location of butterfly survey sites in Loja Province. Points represent individual collection records, with colour indicating sex (blue = female, orange = male). (**C**) Satellite map of survey sites, including Rumi Wilco, Cerro Mandango, Quebrada Puliche, and San Pedro. (**D**) Close-up photograph of a H. himera with a visibly pollen-loaded proboscis.*

along small streams with similar vegetation, adjacent to coffee plantations and pastures.

Butterfly sampling:

From April to September 2023 spanning the wet-to-dry season transition, we sampled *H. himera*

along trails at all five sites, with most effort concentrated at Rumi Wilco and Cerro Mandango,

which were visited at least five days per week. Daily sampling took place between 08:00 and

17:00. This sampling window corresponds to the main period of *Heliconius* activity: *H. himera* is

active early in the morning and remains active into the late afternoon (Davison et al. 1999) and

Heliconius species in Panama also show sustained activity across 08:00–16:00 (Dell'Aglio et al.

2024). Recording the exact capture time, allowed us to examine potential sex differences in

activity pattern.

Each butterfly was captured, marked on the forewing, and released after recording GPS

location, time of capture, sex, body length, pollen load and observer identity. Pollen load was

visually classified into categories based on the proportion of the proboscis covered with pollen:

none (0%), small (<50%), and large (>50%) (Fig. 1D). Two observers collected these data, and

observer identity was recorded for all individuals to account for potential observer effects. Our

categorical scoring system is comparable to pollen-load ranking approaches used in previous

Heliconius field studies (Boggs et al. 1981; Cardoso 2001; Mendoza-Cuenca and Macías-

Ordóñez 2005). Recaptured individuals were noted for movement analyses. In total, 877

butterflies were recorded (371 females, 506 males).

Microclimate measurements were collected for a subset of butterflies during routine fieldwork.

Temperature (°C) and relative humidity (%) were recorded at capture sites using HOBO

Temp/RH U23-002 data loggers placed opportunistically on days when loggers were available.

Loggers were deployed directly at each capture location and recorded continuously for at least

24 h post-capture. The 49 individuals with microclimate measurements (24 females, 25 males)

represent an opportunistic subset determined by logger availability, collected between 26 April

and 12 September 2023, spanning 20 sampling days across April, May, June, July, August, and

September. Sex was sampled randomly across this period, with males and females collected

throughout the full sampling period. These individuals did not overlap with the butterflies used

for pollen metabarcoding, and microclimate data were used only for exploratory analyses of

microhabitat differences between sexes.

Pollen collection and ITS2 sequencing:

To identify pollen sources, we collected 71 *H. himera* (25 females, 46 males) with hand nets

between July and August 2023. The sample contained more males than females because, as is

common in *Heliconius* field surveys, males are more frequently encountered and captured

(Boggs et al. 1981; Cardoso 2001; Jiggins 2017). Additionally, a subset of females was retained

alive to maintain stock cages for parallel experiments. Specimens were placed in entomological

envelopes; proboscises were dissected and stored in PCR tubes, while bodies (containing guts)

were preserved in DMSO/EDTA/NaCl solution. Samples were kept at –15 °C until DNA

extraction and sequencing in February 2024. All proboscises (25 females and 46 males), and a

subset of guts (13 females, 26 males) were analysed using ITS2 DNA metabarcoding, each as

an individual sample (not pooled). DNA metabarcoding enables the identification of pollen to the

genus or species level from mixed samples, without the need for specialized expertise in

traditional palynology (Keller et al. 2015).

DNA was extracted using the Macherey–Nagel NucleoSpin Food Kit (Düren, Germany), exactly

following the ITS2 metabarcoding protocol of Sickel *et al.*, (2015), with additional practical

details on reagents and laboratory steps drawn from (Campos et al. 2021) (Section 8: Standard

method for identification of bee pollen mixtures through meta-barcoding). We included nine

negative controls (DNA-free molecular-grade water) and four positive controls (pre-established

1:3 mixtures of *Musa acuminata* and *Mangifera indica* plant DNA) throughout laboratory

processing to detect potential contamination. PCR amplification, dual-indexing, library

preparation, and sequencing were performed in-house at LMU. Sequencing was carried out on

an Illumina MiSeq using the MiSeq Reagent Kit v2 (2 × 250 bp) (Illumina Inc., San Diego, CA,

USA), with custom read and index primers added to the reagent cartridge following Campos et

al. 2021.

Raw sequence reads were processed combining open-source tools with a custom

bioinformatics pipeline (https://github.com/chiras/metabarcoding_pipeline). Reads were

demultiplexed, primer-trimmed, and merged using VSEARCH (Rognes et al. 2016), with short

reads (<150 bp) discarded. Amplicon sequence variants (ASVs) were inferred via dereplication

and denoising, and chimeric sequences removed. Taxonomic assignments were performed via

global alignment (at least 97% identity threshold, best hit selection) to a curated regional ITS2

plant reference database built using BCdatabaser (Keller et al. 2020). Unclassified reads were

further annotated first with global alignments against a curated worldwide database (Quaresma

et al., 2024), then if still unclassified with the same database hierarchically using the SINTAX

algorithm as taxonomically deep as possible (Edgar 2016). Taxa were retained if they reached

≥1% relative abundance. We restricted analyses to vascular plants using a curated Streptophyta

reference database (Quaresma et al. 2024) that excludes algae and fungi.

Sequencing generated a mean raw throughput of 30,401 ± 14,445 reads per sample (range:

170–101,135, including negative controls). After quality filtering, merging, dereplication, chimera

removal, and taxonomic assignment, samples retained an average of 12,175 reads per sample

(range: 141–31,662). Negative controls showed only very low read counts, while positive ITS2

mock controls amplified as expected, confirming run quality and supporting the removal of

potential contaminants. All remaining samples retained sufficient read depth for downstream

ecological analyses.

Wing reflectance spectrometry and age estimation:

To estimate the relative age of wild individuals, we adapted a spectrometric method developed

by Dell'Aglio et al. (2017), who used wing colour metrics to infer butterfly age. We collected full-

spectrum reflectance data collected from the red and yellow wing bands of insectary-reared

individuals of known age. Reflectance spectra were measured using a Flame-T-XR1-ES

Miniature Spectrometer (Ocean Optics Inc.) connected to a PX-2 Pulsed Xenon light source

(spectral range 220-750 nm) and an Ocean Optics OCF-109053 UV/Vis bifurcated reflection

probe with 200μm fibre diameter. All reflectance measurements were standardized with a WS-1

diffuse white reflectance standard. For each recording (integration time: 2500 ms; three-scan

average), the probe was positioned at a 45° angle and held 1 mm from the wing surface using

an in-house probe mount. This configuration produced an illuminated sampling spot

approximately 2 mm in diameter. For each individual, we collected three replicate spectra per

colour patch (red or yellow), each taken from a different spot within the patch to capture within-

patch variation.

Reflectance was analysed in R with the pavo package (Maia et al. 2019), and the area under

the reflectance curve (AUC) was calculated for each colour patch. In total, we measured

reflectance and age data for 320 insectary-reared individuals across seven *Heliconius* species

with red and/or yellow coloration, including 108 insectary reared *H. himera.* Age was calculated

from known eclosion and sampling dates and linked to reflectance data via standardized

individual identifiers.

Age was predicted using linear models with log-transformed AUC values as predictors. A

combined model including both red and yellow AUCs provided the best fit (adjusted R^2 = 0.13, P

< 0.001) and was used to estimate the age of wild *H. himera*. Predicted values were back-

transformed to age in days, and each individual was assigned a relative age score. This

continuous relative age metric provided a standardized measure across individuals and was

used to test how pollen foraging behaviour varies with age in the wild population.

Eye cuticle morphology:

To test for sexual dimorphism in eye cuticle morphology, we analysed wild-caught and insectary-

reared *H. himera* following established protocols (Wright et al. 2024; Wright and Merrill 2025a;

Wright and Merrill 2025b). Eyes were dissected and incubated in 20% sodium hydroxide

(NaOH) for 18–24 hours to loosen internal tissues beneath the corneal cuticle. After incubation,

each eye was carefully cleaned, mounted in Euparal (Carl Roth GmbH) on a microscope slide,

and left to cure overnight.

Slides were imaged at 25× magnification using a Leica M80 stereomicroscope equipped with a

Flexacam C1 camera and Leica Application Suite X software. All images included a 1 mm scale

bar for size calibration. Corneal surface area was measured using the Freehand Selection and

Measure tools in ImageJ/Fiji (Schindelin et al. 2012), while total ommatidia (facet) number was

estimated via image thresholding and the Analyze Particles function. To account for variation in

body size, we additionally measured inter-ocular distance and hind tibia length using the

Straight-Line tool. Based on the strong bilateral symmetry previously reported for these traits

(Wright et al. 2024), we measured the left eye for all analyses unless it was damaged or

missing, in which case the right eye was used.

Brain immunohistochemistry and Imaging:

We examined sexual dimorphism in brain morphology in 43 *H. himera* (21 females, 22 males),

combining previously analysed wild individuals from Vilcabamba (8 females, 8 males) and two

cohorts   of   common-garden   reared   butterflies,   from   Cambridge   (5   females,   5

males)(Montgomery and Merrill 2017) and Tena, Ecuador (8 females, 9 males). All insectary-

reared butterflies (both Cambridge and Tena cohorts) originated from stocks established from

wild individuals from the same source population in Vilcabamba. The Tena cohort was included,

to increase sample size and to further test whether any sex-specific patterns are robust to

differences in rearing environment.

Brains were labelled using anti-synapsin immunofluorescence (Ott 2008; Montgomery et al.

2016) imaged with confocal microscopy, and neuropils segmented in Amira-Avizo 2023 (Thermo

Fisher Scientific). We measured volumes of key optic and central brain regions previously

shown to be ecologically and evolutionarily relevant in *Heliconius* (Montgomery and Merrill

2017; Montgomery et al. 2021), including the medulla, lamina, lobula, lobula plate, accessory

medulla, ventral lobula, antennal lobe, anterior optic tubercle, and mushroom body calyx and

lobes plus pedunculus. The rest-of-central-brain (rCBR) volume was used as an allometric

control. Paired structures were quantified in one hemisphere and doubled, and volumetric data

were z-axis corrected using a scaling factor of 1.52 (Montgomery and Ott 2015) and $\log_{10}$-

transformed before analysis. We measured brain volumes for the 17 newly collected butterflies

and combined these data with the previously published measurements for the 26 individuals

from Montgomery and Merrill (2017).

Statistical analysis:

All analyses were conducted in R v.4.5.1 (R Core Team 2025). Generalized linear mixed-effects

models (GLMMs) and linear mixed-effects models (LMMs) were fitted using the *lme4* package

(Bates et al. 2015), and all figures were produced with *ggplot2* (Wickham 2011). Estimated

marginal means were computed using the *emmeans* package (Lenth et al. 2019).

Pollen presence was modelled with binomial GLMMs including sex and time of day and their

interaction as fixed effects and random effects for individual and date. We used likelihood-ratio

tests (LRTs) to assess the significance of sex, time of day and their interaction by comparing

each full model to a reduced model lacking the predictor of interest. Pollen load categories were

analysed with cumulative link models using the *ordinal* package (Christensen 2015), with sex

and time of day as fixed effects. Activity patterns, inferred using time of capture, were analysed

by modelling the probability of capturing males versus females as a function of hour of day via

multinomial regression (*nnet* package; (Venables and Ripley 2002). Body size was tested with

linear mixed models including sex as a fixed effect and observer as a random effects.

Movement distances of recaptured individuals were calculated with Vincenty ellipsoid methods

implemented in the *geosphere* package (Hijmans 2024), and analysed with log-transformed

linear models including sex and body length as fixed effects. Microhabitat associations

(Temperature (°C) and relative humidity (%)) with sex were examined with binomial GLMMs

including temperature and relative humidity as continuous predictors and date as a random

effect.

To test effects of sex, body part (proboscis vs. gut), and predicted age on pollen composition,

we used PERMANOVA on Bray–Curtis dissimilarities with the *adonis2* function in vegan

(Oksanen et al. 2025). Pollen richness (observed taxa) and diversity (Shannon index) were

modelled with GLMMs, including date as a random effect. Age was modelled as a continuous

predictor; results were compared with likelihood ratio tests and visualized with estimated

marginal means.

Sexual dimorphism in eye and brain morphology was tested using linear models and

standardized major axis (SMA) regression. For eyes, log_{10}-transformed corneal area and facet

number were modelled with sex, log_{10}-transformed hind tibia length, and log_{10}-transformed

interocular width as fixed effects. For neuropils, log_{10}- transformed volumes were modelled with

sex, rearing location (wild, Cambridge, or Tena), and log_{10} rest-of-central-brain volume (rCBR)

as fixed effects. Because our dataset included individuals from three environments but all

originating from the same source population, we explicitly tested whether sexual dimorphism

was robust to differences in rearing environment by including the Sex × Location interaction in

all models. Model residuals were checked for normality, and fixed effects were evaluated with

Type II ANOVA. Likelihood ratio tests assessed the contribution of predictors, and post hoc

contrasts were performed with *emmeans* using Bonferroni correction. SMA regression (smatr

package (Warton et al. 2012)) was used to test sex-specific differences in slopes, intercepts

(elevation shifts), and major axis shifts along a common slope. We report test statistics, p-

values, and effect sizes as correlation coefficients (r) derived from the reported statistics; for

elevation and shift tests, direction indices (DI) are provided where applicable. P-values from

SMA elevation and shift tests were adjusted using the Benjamini–Hochberg false discovery rate

(FDR) procedure. Results from both linear models and SMA analyses are summarized in Table

S1-S2. Mushroom body calyx and lobes plus pedunculus were analysed separately, consistent

with previous *Heliconius* studies (Montgomery et al. 2016; Montgomery and Merrill 2017).

Results:

Sex Differences in Pollen Presence and Load

Our field collections resulted in records from 371 females and 506 males, which we used to test whether the probability of carrying pollen differed between the sexes. Overall, females were more likely to both carry pollen and tend to carry greater quantities compared to males. Likelihood-ratio tests comparing binomial GLMM models with and without each predictor showed that sex had a significant effect on pollen presence ($\chi^2 = 16.59$, df =1, $P < 0.001$), whereas time of day did not ($\chi^2 = 0.46$, df = 1, P = 0.50) and the Sex × Capture Time interaction did not improve model fit ($\chi^2 = 0.03$, df = 1, P = 0.86). Fitting a cumulative link model analysing

Figure 2: Pollen loads and temporal activity patterns in H. himera. (A) Estimated marginal means of predicted probabilities for each pollen load category by sex, derived from a cumulative link mixed model. Letters denote significant pairwise differences between sexes within each category (α = 0.05). (B Proportional distribution of female (blue) and male (orange) captures by hour of day, with absolute counts shown inside bar segments. (C) H. himera actively foraging on Lantana flowers.

pollen load categories, "no", "small" and "large" (Figure 2A), also revealed that females were

significantly more likely to carry both small (estimate = 0.09, z = 4.31, P < 0.001) and large

pollen loads (estimate = 0.06, z = 4.00, P = 0.0001) than males.

Pollen richness and diversity differ by body part but not by sex

To investigate the diversity of pollen resources used by *H. himera* and assess potential

differences between sexes, sample types, and individual age, we performed DNA

metabarcoding on pollen collected from 25 females and 46 males proboscis samples, as well as

gut samples from a subset of these individuals (13 females and 26 males). Metabarcoding

yielded 1,309,788 high-quality, taxonomically classified reads (mean = 11,907 per sample),

detecting 54 families, 122 genera, and 163 species. All samples passed quality filtering and

were retained for downstream analysis. The most common taxa included *Lantana camara*

(Verbenaceae)(Fig. 2C), *Stellaria media* (Caryophyllaceae), and several Asteraceae species:

Chromolaena odorata, *Kaunia rufescens*, and *Fleischmannia pycnocephala* (Fig. 3A; Fig S1-

S2).

Non-metric multidimensional scaling (NMDS) ordination revealed substantial overlap in pollen

composition, with no significant differences between sexes (PERMANOVA: R^2 = 0.0096, F =

1.05, P = 0.378). In contrast, pollen composition differed significantly between body parts (R^2 =

0.0325, F = 3.63, P = 0.001; Fig. 3B), suggesting some differentiation in the pollen sources

detected externally versus internally, possibly reflecting variation in sampling windows or

digestion. We found no evidence that predicted age explained variation in pollen composition

(R^2 = 0.014, F = 1.03, P = 0.415; Fig. S3).

Generalized linear and linear mixed models confirmed that observed plant species richness and

Shannon diversity differed significantly by body part, with gut samples containing greater

richness (χ^2 = 29.80, df = 1, P < 0.001) and diversity (χ^2 = 15.56, df = 1, P < 0.001) than

proboscis samples (Fig. 3C). In contrast, sex had no effect on either metric (richness: $\chi^2 = 1.00$,

P = 0.32; Shannon: $\chi^2 = 0.002$, P = 0.97). Predicted age did not influence richness or diversity

when modelled as a continuous variable (richness: $\chi^2 = 0.19$, P = 0.66; Shannon: $\chi^2 = 1.78$, P =

0.18).

Figure 3: Diversity and composition of pollen collected from different sexes and tissues. (**A**) *Relative read abundance of the most common plant taxa identified in proboscis and gut samples. Bars represent the mean relative abundance across individuals, with colors indicating plant families. (**B**) NDMS plot across samples, showing clustering by body part (proboscis = purple, gut = green) and sex (female = circles, male = triangles). 95% confidence ellipses are shown for each group. (**C**) Observed richness and Shannon diversity of pollen samples collected from proboscis and gut tissues of H. himera, separated by sex (females blue and males orange).*

Daily activity, movement, habitat use and age

Given the sex differences in pollen presence and load size, we next tested for variation in body size, daily activity, movement, and microhabitat use (Fig. S4). Females averaged 24.4 mm in length (SD = 1.66, IQR = 23–26), while males averaged 25.3 mm (SD = 1.55, IQR = 24–26). Linear models confirmed males were larger (χ^2 = 129.73, df =1, P < 0.001; Fig. S4)

Daily activity also differed between the sexes (LR = 19.11, P < 0.001), with females initiating foraging behaviour earlier in the day (Fig. 2B). To test whether time of day influenced pollen presence, we included capture time as a fixed effect in the binomial GLMM. After accounting for sex (χ^2 = 16.59, df=1, P < 0.001), capture time did not improve the model (χ^2 = 0.46, df=1, P = 0.50), and the Sex × Capture Time interaction was also non-significant (χ^2 = 0.03, P = 0.86). Thus, earlier female activity did not translate into increased pollen loads.

Movement was analysed from 109 recaptured females and 162 males. Recaptured distances varied widely but were comparable: females moved a mean of 87.2 m (median = 24.8, SD = 176.0) and males 66.9 m (median = 36.2, SD = 95.1). A linear model of log-transformed distances showed no effect of sex on movement distance (χ^2 = 0.63, P = 0.61; Fig. S4).

Microclimate data were available for an opportunistic subset of individuals and were used to explore potential sex differences in local microclimatic conditions. Both sexes were recaptured under similar microclimatic conditions. Females were recaptured at a mean temperature of 25.8 °C (median = 25.9, SD = 3.5) and 50.9% relative humidity (median = 48.0, SD = 16.8), while males were recaptured at 26.4 °C (median = 26.2, SD = 3.4) and 49.5% relative humidity (median = 48.3, SD = 15.0). These distributions overlapped strongly, with interquartile ranges of 23.3–27.9 °C for females and 24.1–29.3 °C for males. Binomial GLMMs confirmed that neither temperature (χ^2 = 1.05, df = 1, P = 0.31) nor humidity (χ^2 = 0.11, P = 0.75) significantly affected the probability of capturing either sex (Fig. S4).

Sexual dimorphism in eye and brain morphology

We tested for sexual dimorphism in eye and brain morphology (Fig. 4). Males had larger eyes

than females ($F_{1,39}$ = 26.25, P < 0.001). SMATR analysis showed no sex difference in slope

between corneal area and tibia length but a significant elevation shift (Wald χ^2 = 68.82, FDR-p <

0.001), with males consistently having larger eyes for a given tibia length (Fig. 4A). Facet

number was also higher in males ($F_{1,40}$ = 6.07, P = 0.018). Using corneal area as the allometric

control, SMATR again detected no sex difference in slope or elevation but a significant major

axis shift (Wald χ^2 = 21.23, FDR-p < 0.001), showing that males more facets (Fig. 4D).

We also found sex differences in brain morphology, particularly in higher-order integrative

regions (Fig. 4B–F). For the mushroom body calyces (MBCA), linear models revealed a strong

effect of sex (χ^2_1 = 28.51, P < 0.001). Rearing location did not significantly affect MBCA volume

(χ^2_2 = 2.81, P = 0.28), and the Sex × Location interaction was also non-significant (χ^2_2 = 2.99, P

= 0.27), indicating that the observed sexual dimorphism was consistent across rearing

environments. SMATR analysis confirmed a significant elevation shift in the scaling relationship

between MBCA volume and rCBR (Wald χ^2_1 = 34.25, FDR-p < 0.001), with females having

larger MBCA for a given brain size (Fig. 4C). A similar pattern, with larger female investment

was observed for the mushroom body lobes and pedunculus (MBPED). Sex showed a

significant effect on MBPED volume (χ^2_1 = 29.38, P < 0.001), with females having larger

MBPED. Rearing location showed a main effect (χ^2_2 = 37.51, P < 0.001), but the Sex × Location

interaction remained non-significant (χ^2_2 = 3.96, P = 0.18). Linear models showed significant

effects of sex (χ^2_1 = 29.5, P < 0.001), after accounting for rCBR and rearing condition location

(χ^2_2 = 49.8, P < 0.0001). SMATR analysis detected a significant elevation shift in the scaling

relationship (Wald χ^2_1 = 8.17, FDR-p =0.023), again indicating larger investment in these

mushroom body regions in females (Fig. 4F).

By contrast, we found no evidence of sexual dimorphism in the optic lobes.

Sex was not a significant predictor of optic lobe volume (χ^2_1 = 1.02, P = 0.34), and the Sex ×

Location interaction was likewise non-significant (χ^2_2 = 6.19, P = 0.06). Rearing location showed

a modest main effect (χ^2_2 = 8.33, P = 0.02), driven by overall size differences between cohorts,

but this did not translate into sex-specific variation (Fig. 4B). Similarly, antennal lobe volumes

did not differ between sexes (χ^2_1 = 3.92, P = 0.06), and the Sex × Location interaction was not

significant (χ^2_2 = 3.70, P = 0.20). As with the optic lobes, rearing location had a main effect (χ^2_2 =

52.61, P < 0.001), but this reflected overall size variation rather than sexual dimorphism (Fig.

4E). Full model results are provided in Supplementary Tables S1-S2.

Figure 4: Sexual dimorphism in eye and brain morphology. *Females = blue; Males = orange. All traits are $\log_{10}$-transformed. (**A**) Corneal area shows a significant elevation shift, with males having larger eyes. (**B, E**) Optic lobe (OL) and antennal lobe (AL) volumes scale with rest-of-central-brain volume (rCBR) but show no sex differences. (**C, F**) Mushroom body calyx (MBCA) and mushroom body lobes and pedunculus (MBLOBES+PED) exhibit a significant elevation shift, with females having larger volumes than males. (**D**) Facet count shows a significant major axis shift, with males having more facets. Lines indicate sex-specific SMATR regressions (solid) or common slopes (dashed). Insets show the corresponding anatomical structures: eye (gray), OL (yellow/orange), MB (red), AL (blue).*

Discussion

Systematic field sampling combined with DNA metabarcoding revealed high floral diversity and

clear sex differences in pollen foraging behaviour in wild *H. himera*. Females carried larger

pollen loads than males, consistent with sex-specific nutritional demands; yet we observed no

differences we observed no differences plant richness and composition. This indicates that sex

differences reflect variation in foraging effort rather than shifts in plant choice. Notably, gut

samples contained higher pollen diversity and more consistent community profiles than

proboscis samples, suggesting they better capture cumulative foraging history. We further

confirmed sexual dimorphism in eye morphology and mushroom body volume, consistent more

generally with patterns reported for other aspects of vision across *Heliconius* (McCulloch et al.

2016; McCulloch et al. 2017; Chakraborty et al. 2023; Couto et al. 2023; Wright et al. 2023;

Wright et al. 2024; Buerkle et al. 2025; VanKuren et al. 2025). While we did not directly test

functional links between sensory traits and foraging effort, our study provides one of the first

molecular characterizations of pollen resource use in *Heliconius*, showing that sex-specific

foraging reflects differential effort on shared floral resources rather than partitioning of plant

species.

The floral resources used by *H. himera* differed markedly from those of lowland *Heliconius*.

Metabarcoding revealed a diverse diet of 163 plant species spanning, 122 genera and 54

families. While *Lantana camara* (*Verbenaceae*) was the most frequent resource, *Asteraceae*

were also well represented, particularly *Chromolaena* and *Kaunia*, which produce amino acid–

rich nectar and are widely visited by butterflies and other pollinators (Wist and Davis 2008;

Rathnayake and Wijetunga 2016; Layek et al. 2022). By contrast, *Gurania* and *Psiguria*

(*Cucurbitaceae*), central to the foraging ecology of lowland *Heliconius (Estrada and Jiggins*

2002; Jiggins 2017), are entirely absent in the range and diet of *H. himera*. Instead, *H. himera*

relied heavily on *Lantana*, highlighting its role as a core floral resource in montane dry forests.

Larval resource use also reflects this environment: *H. himera* oviposits on *Passiflora* species of

the *Decaloba* subgenus, which tolerate dry conditions and occur in open habitats (Ulmer and

MacDougal 2004). Thus, both larval and adult foraging are shaped by the montane dry-forest

environment: larvae exploit drought-tolerant host plants, while adults rely on a broad array of

floral resources. Together, these patterns suggest that *erato* clade species, including *H. himera,*

have retained a generalist strategy across life stages, facilitating colonisation of high-altitude

habitats.

In our study, both sexes of *H. himera* experienced similar environmental conditions and

movement patterns, indicating shared microhabitat use and strong spatial overlap in access to

floral resources. In patchy, resource-limited montane habitats, such overlap may favour a

generalist foraging strategy over sex-specific specialization. Accordingly, sex-specific nutritional

demands appear to be met through differences in foraging effort rather than distinct floral

species. Females for example, may accumulate larger pollen loads through greater persistence

at flowers; recent behavioural work shows that *Heliconius* females often revisit flowers

repeatedly, leading to increased pollen accumulation (Dell'Aglio et al. 2024). In other *Heliconius*,

microhabitat differences often underlie floral preferences rather than intrinsic species identity

(Cardoso 2001; Estrada and Jiggins 2002). For example, *H. melpomene* and *H. cydno* rely on

Lantana and *Psychotria*, respectively, due to habitat use rather than innate preference (Jiggins

2017). Likewise, long-term studies of *Argynnis idalia* show that sex differences in floral visitation

diminish when males and females forage during overlapping phenological windows

(Chmielewski et al. 2023). By contrast, *H. charithonia* exhibits clear sex-specific floral

preferences (Mendoza-Cuenca and Macías-Ordóñez 2005). The absence of strong floral

partitioning in *H. himera* therefore suggests that sex-specific strategies are expressed mainly

through pollen collection effort, as an adaptive response to ecological constraints of montane

environments.

Our comparison of gut and proboscis samples confirmed that tissue type strongly affects pollen

detection. Gut samples showed higher richness and diversity, likely reflecting a longer

integration window and cumulative foraging history or more passive uptake from nectar.

Because whole abdomens were processed, we cannot exclude the possibility that pollen on the

abdomen exterior contributed to gut-sample detections; however, we note that this might still be

informative for assessing habitat use. Proboscis samples were often dominated by *Lantana*,

suggesting its importance as a core pollen source, actively processed through prolonged

proboscis manipulation and digestion. In *Heliconius*, pollen digestion occurs externally on the

proboscis surface, and only dissolved nutrients are ingested (Gilbert 1972; Penz and Krenn

2000; Eberhard et al. 2007; Hikl and Krenn 2011; Harpel et al. 2015). As a result, pollen from

nectar-only visits or brief contacts may be detected externally, which could explain why some

taxa (e.g., *Impatiens walleriana*) appeared exclusively in proboscis samples. Together, these

patterns suggest that gut samples may reflect longer-term foraging, whereas proboscis samples

more closely reflect recent plant interactions and active pollen processing.

Beyond these tissue-specific insights, our results highlight the power of DNA metabarcoding for

ecological research in butterflies. This approach enabled species-level identification of complex

pollen mixtures from small tissue samples, detecting a broad taxonomic spectrum not easily

resolved by classical palynology (Bell et al. 2016). The successful use of DMSO-preserved

samples further demonstrates its potential for archived collections, opening opportunities to

compare historical and spatial variation in pollen use, shifts in foraging behaviour, and

interspecific differences across the *Heliconius* clade. For example, seasonal or local changes in

pollen availability may influence investment in cyanogenic glycosides or egg-laying effort

(Jiggins 2017; Young and Montgomery 2020). Leveraging DNA metabarcoding across temporal

and environmental gradients can thus deepen our understanding of the ecological and

evolutionary dynamics structuring pollinator–plant interactions in tropical ecosystems.

Higher pollen loads in females likely reflect reproductive and physiological demands. This

pattern aligns with earlier observations in other *Heliconius* species, where females forage more

actively and carry larger pollen loads than males (Boggs et al. 1981; Cardoso 2001; Mendoza-

Cuenca and Macías-Ordóñez 2005). Female *Heliconius* require a steady intake of amino acids

and lipids for oogenesis and to maintain cyanogenic glycosides that serve as chemical defenses

for adults and offspring (Gilbert 1972; Dunlap-Pianka et al. 1977; O'Brien et al. 2003; Pinheiro

de Castro et al. 2025). These demands cannot be met through nectar alone, making pollen

essential for replenishing protein reserves. Males also benefit from amino acids for

spermatophore production and maintenance (Boggs and Gilbert 1979; Boggs et al. 1981), but

carried lower pollen loads, possibly reflecting weaker nutritional demands, greater time

investment in mate search, or greater efficiency in resource extraction. Female-biased foraging

extends beyond *Heliconius*: across Lepidoptera, females preferentially use nutrient-rich

resources supplying amino acids, lipids, and sterols required for egg production (Alm et al.

1990; Erhardt and Rusterholz 1998; O'Brien et al. 2004; Mevi□Schütz and Erhardt 2005; Keller

et al. 2015; Chmielewski et al. 2023). This sex-specific investment pattern is consistent with

optimal foraging theory, where individuals adjust investment in foraging behaviour according to

the relative fitness payoffs of nutrient acquisition (Houston 1995). For females, the high returns

of pollen amino acids for egg production outweigh the time and energetic costs of prolonged

foraging, while males balance pollen collection against investment in mate-searching behaviour.

Although females began foraging earlier in the day, time of day did not predict pollen presence,

nor did its effect differ between sexes, indicating that earlier activity alone does not explain the

observed female bias in pollen load. More broadly, sex-based differences in resource use are an

important source of intraspecific variation that may influence pollination networks and

community dynamics (Smith et al. 2019; Kishi and Kakutani 2020; Smith et al. 2021).

We used full-spectrum wing reflectance to estimate the relative age of wild butterflies, adapting

a model trained on known-age individuals. Although wing wear and colouration are widely used

as age proxies in Lepidoptera (Kemp 2006; Dell'Aglio et al. 2017), they are influenced by

environmental exposure and behaviour, so reflectance-based estimates should be interpreted

cautiously. In our dataset, predicted age did not explain variation in pollen richness or diversity.

This contrasts with earlier work on different *Heliconius* species from Trinidad and Costa Rica

(Boggs et al. 1981) which reported age-related peaks in pollen load, suggesting that age effects

may be weak or context dependent.

Our analyses revealed pronounced sexual dimorphism in the visual system, with males

possessing significantly larger eyes and more facets than females. Similar male-biased visual

investment has been widely reported in insects and is often associated with mate detection or

mate-search behaviour (Ziemba and Rutowski 2000; Somanathan et al. 2017). In contrast,

males and females visited the same floral species, indicating that variation in eye morphology

does not correspond to differences in floral resource breadth. *H. himera* females had

significantly larger mushroom bodies, consistent with the *Heliconius*-wide patterns (Couto et al.

2023), but functional basis of this dimorphism remains unclear: laboratory assays of learning

and memory in *Heliconius* have not detected sex effects (Toure et al. 2020; Young et al. 2024;

Hodge et al. 2025). We therefore treat the neuroanatomical patterns as descriptive rather than

explanatory. Future work should directly test whether males and females differ in colour

discrimination, odour sensitivity, or flower-approach behaviour, and whether the observed visual

dimorphism is primarily shaped by sexual selection rather than foraging ecology.

Together, our findings show that female *H. himera* carry larger pollen loads than males, yet both
sexes exploit a similar diversity and composition of floral resources, indicating sexes adopt
divergent behavioural strategies on effort rather than specialization on different floral species. In
patchy montane environments, such a generalist strategy may facilitate colonisation of high-
altitude habitats. Our use of DNA metabarcoding highlights the value of molecular tools for
reconstructing butterfly diets and linking sex-specific foraging to broader questions of
intraspecific variation and pollination ecology. Gut samples proved particularly informative,
capturing a broader and more integrated record of pollen use than proboscis samples and
underscoring the importance of tissue choice in dietary inference. We also documented sexual
dimorphism in eye and brain morphology, consistent with patterns across *Heliconius*, although
the   functional   significance   of   these   traits   remains   unresolved.
538

Acknowledgements

We are grateful to the Ministerio del Ambiente, Agua y Transición Ecológica for permission to

collect butterflies in Ecuador (MAATE-DBI-CM-2021-0176; MAATE-DNB-CM-2021-0176). We

thank Alicia Cerchiai for allowing us access to Bosque y Vegetación Protectores Rumi Wilco.

Alicia Cerchiai, Alejandro Mancino, Verónica Romero, and Veronika Schottenheim contributed to

plant and butterfly stock maintenance. Yi Peng Toh assisted with brain imaging.

Portions of the text were revised with assistance from OpenAI's ChatGPT (GPT-4, accessed

July 2025) to improve clarity and style. All content was critically reviewed and finalized by the

authors.

Funding information

Research Funded by ERC Starter Grant 851040 to R.M.M. and a scholarship to D.F.R.S. from

the Ministerio de Ciencia, Tecnología e Innovación (Convocatoria No. 860).

Conflict of interest

The authors have no conflict of interest to declare.

Author Contributions

J.B.: conceptualization, data curation, formal analysis, investigation, visualization, writing—

original draft, writing—review and editing; D.F.R.S.: conceptualization, investigation, writing—

review and editing; D.S.W.: methodology, writing—review and editing; S.H.M.:

conceptualization, supervision, writing—review and editing; A. K.: resources, formal analysis,

writing—review and editing.; R.M.M.: conceptualization, resources, supervision, writing—original

draft, writing—review and editing.

Data Availability Statement

All datasets, analysis scripts, and metadata associated with this study are publicly available on

Zenodo. The full repository can be accessed at:

https://doi.org/10.5281/zenodo.17777739

References

Alm J, Ohnmeiss TE, Lanza J, Vriesenga L. 1990. Preference of cabbage white butterflies and honey bees for nectar that contains amino acids. Oecologia. 84(1):53–57. https://doi.org/10.1007/BF00665594

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. J Stat Soft. 67(1):1–48. https://doi.org/10.18637/jss.v067.i01

Bell KL et al. 2016. Pollen DNA barcoding: current applications and future prospects. Genome. 59(9):629–640. https://doi.org/10.1139/gen-2015-0200

Boggs C, Gilbert LE. 1979. Male Contribution to Egg Production in Butterflies: Evidence for Transfer of Nutrients at Mating. Science. 206(4414):83–84. https://doi.org/10.1126/science.206.4414.83

Boggs C, Smiley J, Gilbert L. 1981. Patterns of pollen exploitation by Heliconius butterflies. Oecologia. 48:284–289. https://doi.org/10.1007/BF00347978

Borrero J et al. 2024. Weighting of sensory cues reflect changing patterns of visual investment during ecological divergence in Heliconius butterflies. Biology Letters. 20(10):20240377. https://doi.org/10.1098/rsbl.2024.0377

Borrero J, Wright DS, Bacquet CN, Merrill RM. 2023. Oviposition behavior is not affected by ultraviolet light in a butterfly with sexually-dimorphic expression of a UV-sensitive opsin. Ecology and Evolution. 13(7):e10243. https://doi.org/10.1002/ece3.10243

Buerkle NP et al. 2025. Sex-limited diversification of the eye in Heliconius cydno butterflies. J Comp Physiol A. [published online ahead of print] [accessed 2025 Nov 3]. https://link.springer.com/10.1007/s00359-025-01768-z. https://doi.org/10.1007/s00359-025-01768-z

Campos MG et al. 2021. Standard methods for pollen research. Journal of Apicultural Research. 60(4):1–109. https://doi.org/10.1080/00218839.2021.1948240

Cardoso M. 2001. Patterns of Pollen Collection and Flower Visitation by Heliconius Butterflies in Southeastern Mexico. Journal of Tropical Ecology. 17(5):763–768. https://doi.org/10.1017/S0266467401001572

Chakraborty M et al. 2023. Sex-linked gene traffic underlies the acquisition of sexually dimorphic UV color vision in Heliconius butterflies. Proceedings of the National Academy of Sciences. 120(33):e2301411120. https://doi.org/10.1073/pnas.2301411120

Chmielewski MW et al. 2023. Phenology and foraging bias contribute to sex-specific foraging patterns in the rare declining butterfly Argynnis idalia idalia. Ecology and Evolution. 13(7):e10287. https://doi.org/10.1002/ece3.10287

Christensen RHB. 2015. Package 'ordinal'. Stand. 19(2016)

Couto A et al. 2023. Rapid expansion and visual specialisation of learning and memory centres in the brains of Heliconiini butterflies. Nat Commun. 14(1):4024. https://doi.org/10.1038/s41467-023-39618-8

Davison A et al. 1999. Behavioral and physiological differences between two parapatric Heliconius species. Biotropica. 31(4):661–668. https://doi.org/10.1111/j.1744-7429.1999.tb00415.x

Dell'Aglio D, Mena S, et al. 2022. Divergence in *Heliconius* flight behaviour is associated with local adaptation to different forest structures. Journal of Animal Ecology. 91(4):727–737. https://doi.org/10.1111/1365-2656.13675

Dell'Aglio D, Akkaynak D, McMillan WO, Jiggins CD. 2017. Estimating the age of *Heliconius* butterflies from calibrated photographs. PeerJ. 5:e3821. https://doi.org/10.7717/peerj.3821

Dell'Aglio D, McMillan OW, Montgomery S. 2024. Using motion-detection cameras to monitor foraging behaviour of individual butterflies. Ecology and Evolution. 14(7):e70032. https://doi.org/10.1002/ece3.70032

Dell'Aglio D, McMillan WO, Montgomery SH. 2022. Shifting balances in the weighting of sensory modalities are predicted by divergence in brain morphology in incipient species of Heliconius butterflies. Animal Behaviour. 185:83–90. https://doi.org/10.1016/j.anbehav.2022.01.003

Descimon H, Mast DM. 1984. Semispecies relationships between Heliconius erato cyrbia Godt. and H. himera Hew. in southwestern Ecuador. The Journal of Research on the Lepidoptera. 22(4):229–237. https://doi.org/10.5962/p.266821

Dötterl S, Milchreit K, Schäffler I. 2011. Behavioural plasticity and sex differences in host finding of a specialized bee species. J Comp Physiol A. 197(12):1119–1126. https://doi.org/10.1007/s00359-011-0673-2

Dunlap-Pianka H, Boggs CL, Gilbert LE. 1977. Ovarian Dynamics in Heliconiine Butterflies: Programmed Senescence versus Eternal Youth. Science. 197(4302):487–490. https://doi.org/10.1126/science.197.4302.487

Eberhard SH, Hrassnigg N, Crailsheim K, Krenn HW. 2007. Evidence of protease in the saliva of the butterfly *Heliconius melpomene* (L.) (Nymphalidae, Lepidoptera). Journal of Insect Physiology. 53(2):126–131. https://doi.org/10.1016/j.jinsphys.2006.11.001

Edgar RC. 2016. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. 074161 [accessed 2025 June 3]. https://www.biorxiv.org/content/10.1101/074161v1. https://doi.org/10.1101/074161

Erhardt A, Rusterholz H-P. 1998. Do peacock butterflies (Inachis io L.) detect and prefer nectar amino acids and other nitrogenous compounds? Oecologia. 117:536–542

Estrada C, Jiggins CD. 2002. Patterns of pollen feeding and habitat preference among Heliconius species. Ecological Entomology. 27(4):448–456. https://doi.org/10.1046/j.1365-2311.2002.00434.x

Finkbeiner SD, Briscoe AD. 2021. True UV color vision in a female butterfly with two UV opsins.
Journal of Experimental Biology. 224(18):jeb242802. https://doi.org/10.1242/jeb.242802

Gilbert LE. 1972. Pollen Feeding and Reproductive Biology of *Heliconius* Butterflies. Proc Natl
Acad Sci USA. 69(6):1403–1407. https://doi.org/10.1073/pnas.69.6.1403

Gilbert LE. 1991. Biodiversity of a Central American Heliconius community: pattern, process,
and problems. Plant-animal interactions: evolutionary ecology in tropical and temperate regions.
403–427

Harpel D et al. 2015. Pollen feeding proteomics: Salivary proteins of the passion flower butterfly,
Heliconius melpomene. Insect Biochemistry and Molecular Biology. 63:7–13.
https://doi.org/10.1016/j.ibmb.2015.04.004

Hijmans RJ. 2024. geosphere: Spherical Trigonometry. https://github.com/rspatial/geosphere

Hikl A-L, Krenn HW. 2011. Pollen processing behavior of Heliconius butterflies: A derived
grooming behavior. J Insect Sci. 11(1):99. https://doi.org/10.1673/031.011.9901

Hodge EA et al. 2025. Modality-specific long-term memory enhancement in *Heliconius*
butterflies. Phil Trans R Soc B. 380(1929):20240119. https://doi.org/10.1098/rstb.2024.0119

Houston AI. 1995. Energetic constraints and foraging efficiency. Behavioral Ecology. 6(4):393–
396. https://doi.org/10.1093/beheco/6.4.393

Jiggins CD. 2017. The Ecology and Evolution of Heliconius Butterflies. Oxford University Press.
[accessed 2023 Apr 26]. https://academic.oup.com/book/26557.
https://doi.org/10.1093/acprof:oso/9780199566570.001.0001

Jiggins CD, McMillan WO, Neukirchen W, Mallet J. 1996. What can hybrid zones tell us about
speciation? The case of Heliconius erato and H. himera (Lepidoptera: Nymphalidae). Biological
Journal of the Linnean Society. 59(3):221–242. https://doi.org/10.1111/j.1095-
8312.1996.tb01464.x

Keller A et al. 2015. Evaluating multiplexed next-generation sequencing as a method in
palynology for mixed pollen samples Mock H -P., editor. Plant Biol J. 17(2):558–566.
https://doi.org/10.1111/plb.12251

Keller A et al. 2020. BCdatabaser: on-the-fly reference database creation for (meta-)barcoding
Valencia A, editor. Bioinformatics. 36(8):2630–2631.
https://doi.org/10.1093/bioinformatics/btz960

Kemp DJ. 2006. Heightened phenotypic variation and age-based fading of ultraviolet butterfly
wing coloration. Evolutionary Ecology Research. 8(3):515–527

Kishi S, Kakutani T. 2020. Male Visitors May Decrease Modularity in Flower–Visitor Networks.
Front Ecol Evol. 8 [accessed 2025 July 10]. https://www.frontiersin.org/journals/ecology-and-
evolution/articles/10.3389/fevo.2020.00124/full. https://doi.org/10.3389/fevo.2020.00124

Land M. 1997. Visual Acuity in Insects. Annual review of entomology. 42:147–77.
https://doi.org/10.1146/annurev.ento.42.1.147

Layek U, Das A, Das U. 2022. Floral biology, floral volatile organic compounds and floral visitors
of Chromolaena odorata, an invasive alien species in West Bengal, India. Biodiversitas Journal
of Biological Diversity. 23(4) [accessed 2025 July 9]. https://smujo.id/biodiv/article/view/10495.
https://doi.org/10.13057/biodiv/d230447

Lenth R et al. 2019. Package 'emmeans'.

Maia R, Gruson H, Endler JA, White TE. 2019. pavo 2: New tools for the spectral and spatial
analysis of colour in r. Methods in Ecology and Evolution. 10(7):1097–1107.
https://doi.org/10.1111/2041-210X.13174

Martins AC et al. 2023. Contrasting patterns of foraging behavior in neotropical stingless bees
using pollen and honey metabarcoding. Sci Rep. 13(1):1–13. https://doi.org/10.1038/s41598-
023-41304-0

McCulloch KJ et al. 2017. Sexual Dimorphism and Retinal Mosaic Diversification following the
Evolution of a Violet Receptor in Butterflies. Molecular biology and evolution. 34(9):2271–2284.
https://doi.org/10.1093/molbev/msx163

McCulloch KJ, Osorio D, Briscoe AD. 2016. Sexual dimorphism in the compound eye of
Heliconius erato: A nymphalid butterfly with at least five spectral classes of photoreceptor.
Journal of Experimental Biology. 219(15):2377–2387. https://doi.org/10.1242/jeb.136523

Mendoza-Cuenca L, Macías-Ordóñez R. 2005. Foraging polymorphism in *Heliconius charitonia*
(Lepidoptera: Nymphalidae): morphological constraints and behavioural compensation. J Trop
Ecol. 21(4):407–415. https://doi.org/10.1017/S0266467405002385

Mevi-Schütz J, Erhardt A. 2005. Amino Acids in Nectar Enhance Butterfly Fecundity: A
Long-Awaited Link. The American Naturalist. 165(4):411–419. https://doi.org/10.1086/429150

Montgomery SH, Merrill RM. 2017. Divergence in brain composition during the early stages of
ecological specialization in Heliconius butterflies. Journal of Evolutionary Biology. 30(3):571–
582. https://doi.org/10.1111/jeb.13027

Montgomery SH, Merrill RM, Ott SR. 2016. Brain composition in Heliconius butterflies,
posteclosion growth and experience-dependent neuropil plasticity. Journal of Comparative
Neurology. 524(9):1747–1769. https://doi.org/10.1002/cne.23993

Montgomery SH, Ott SR. 2015. Brain composition in Godyris zavaleta, a diurnal butterfly,
Reflects an increased reliance on olfactory information. Journal of Comparative Neurology.
523(6):869–891. https://doi.org/10.1002/cne.23711

Montgomery SH, Rossi M, McMillan WO, Merrill RM. 2021. Neural divergence and hybrid
disruption between ecologically isolated *Heliconius* butterflies. Proc Natl Acad Sci USA.
118(6):e2015102118. https://doi.org/10.1073/pnas.2015102118

O'Brien DM, Boggs CL, Fogel ML. 2003. Pollen feeding in the butterfly Heliconius charitonia:
isotopic evidence for essential amino acid transfer from pollen to eggs. Proceedings of the
Royal Society of London Series B: Biological Sciences. 270(1533):2631–2636.
https://doi.org/10.1098/rspb.2003.2552

O'Brien DM, Boggs CL, Fogel ML. 2004. Making eggs from nectar: the role of life history and dietary carbon turnover in butterfly reproductive resource allocation. Oikos. 105(2):279–291. https://doi.org/10.1111/j.0030-1299.2004.13012.x

Oksanen J et al. 2025. vegan: Community Ecology Package. https://vegandevs.github.io/vegan/

Ott SR. 2008. Confocal microscopy in large insect brains: Zinc–formaldehyde fixation improves synapsin immunostaining and preservation of morphology in whole-mounts. Journal of Neuroscience Methods. 172(2):220–230. https://doi.org/10.1016/j.jneumeth.2008.04.031

Penz CM, Krenn HW. 2000. Behavioral Adaptations to Pollen-Feeding in Heliconius Butterflies (Nymphalidae, Heliconiinae): An Experiment Using Lantana Flowers. Journal of Insect Behavior. 13:865–880. https://doi.org/10.1023/A:1007814618149

Piko J et al. 2021. Effects of three flower field types on bumblebees and their pollen diets. Basic and Applied Ecology. 52:95–108. https://doi.org/10.1016/j.baae.2021.02.005

Pinheiro de Castro EC et al. 2020. The dynamics of cyanide defences in the life cycle of an aposematic butterfly: Biosynthesis versus sequestration. Insect Biochemistry and Molecular Biology. 116:103259. https://doi.org/10.1016/j.ibmb.2019.103259

Pinheiro de Castro EC et al. 2025. Pollen-feeding delays reproductive senescence and maintains toxicity of Heliconius erato. Peer Community Journal. 5. https://doi.org/10.24072/pcjournal.546

Quaresma A et al. 2024. Semi-automated sequence curation for reliable reference datasets in ITS2 vascular plant DNA (meta-)barcoding. Sci Data. 11(1):129. https://doi.org/10.1038/s41597-024-02962-5

R Core Team. 2025. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. https://www.R-project.org/

Rathnayake DGRMMK, Wijetunga WMGASTB. 2016. Species Composition and Visiting Frequencies of Flower Visitors of Chromolaena odorata in a Dry Zone Forest Patch of Sri Lanka. Psyche: A Journal of Entomology. 2016(1):8746251. https://doi.org/10.1155/2016/8746251

Rognes T et al. 2016. VSEARCH: a versatile open source tool for metagenomics. PeerJ. 4:e2584. https://doi.org/10.7717/peerj.2584

Romeis J, Städler E, Wäckers F. 2005. Nectar- and pollen-feeding by adult herbivorous insects. In: Environmental Microbiology - ENVIRON MICROBIOL. p 178–219. https://doi.org/10.1017/CBO9780511542220.008

Roulston TH, Cane JH. 2000. Pollen nutritional content and digestibility for animals. Pl Syst Evol. 222(1):187–209. https://doi.org/10.1007/BF00984102

Ruedenauer FA, Spaethe J, van der Kooi CJ, Leonhardt SD. 2019. Pollinator or pedigree: which factors determine the evolution of pollen nutrients? Oecologia. 191(2):349–358. https://doi.org/10.1007/s00442-019-04494-x

Schindelin J et al. 2012. Fiji: an open-source platform for biological-image analysis. Nat Methods. 9(7):676–682. https://doi.org/10.1038/nmeth.2019

Sickel W et al. 2015. Increased efficiency in identifying mixed pollen samples by meta-barcoding with a dual-indexing approach. BMC Ecology. 15(1):20. https://doi.org/10.1186/s12898-015-0051-y

Sivinski J. 1989. Mushroom body development in nymphalid butterflies: A correlate of learning? J Insect Behav. 2(2):277–283. https://doi.org/10.1007/BF01053299

Smith GP et al. 2021. Sex-associated differences in the network roles of pollinators. Ecosphere. 12(12):e03863. https://doi.org/10.1002/ecs2.3863

Smith GP, Bronstein JL, Papaj DR. 2019. Sex differences in pollinator behavior: Patterns across species and consequences for the mutualism. Journal of Animal Ecology. 88(7):971–985. https://doi.org/10.1111/1365-2656.12988

Somanathan H, Borges RM, Warrant EJ, Kelber A. 2017. Visual Adaptations for Mate Detection in the Male Carpenter Bee Xylocopa tenuiscapa Nieh JC, editor. PLoS ONE. 12(1):e0168452. https://doi.org/10.1371/journal.pone.0168452

Stöckl A et al. 2016. Differential investment in visual and olfactory brain areas reflects behavioural choices in hawk moths. Sci Rep. 6(April):1–10. https://doi.org/10.1038/srep26041

Toure MW, Young FJ, McMillan WO, Montgomery SH. 2020. Heliconiini butterflies can learn time-dependent reward associations. Biology Letters. 16(9):20200424. https://doi.org/10.1098/rsbl.2020.0424

Ulmer T, MacDougal JM. 2004. Passiflora: passionflowers of the world. Timber Press (OR).

VanKuren NW et al. 2025. Genetic, developmental, and neural changes underlying the evolution of butterfly mate preference Desplan C, editor. PLoS Biol. 23(3):e3002989. https://doi.org/10.1371/journal.pbio.3002989

Venables WN, Ripley BD. 2002. Modern Applied Statistics with S. Fourth. Springer. https://www.stats.ox.ac.uk/pub/MASS4/

Warton DI, Duursma RA, Falster DS, Taskinen S. 2012. smatr 3– an R package for estimation and inference about allometric lines. Methods in Ecology and Evolution. 3(2):257–259. https://doi.org/10.1111/j.2041-210X.2011.00153.x

Wickham H. 2011. ggplot2. Wiley interdisciplinary reviews: computational statistics. 3(2):180–185. https://doi.org/10.1002/wics.147

Wilson RS et al. 2021. Many small rather than few large sources identified in long-term bee pollen diets in agroecosystems. Agriculture, Ecosystems & Environment. 310:107296. https://doi.org/10.1016/j.agee.2020.107296

Wist TJ, Davis AR. 2008. Floral Structure and Dynamics of Nectar Production in Echinacea pallida var. angustifolia (Asteraceae). International Journal of Plant Sciences. 169(6):708–722. https://doi.org/10.1086/533602

787 Wright DS et al. 2023. Quantifying visual acuity in Heliconius butterflies. Biology Letters.
788 19(12):20230476. https://doi.org/10.1098/rsbl.2023.0476

789 Wright DS et al. 2024. Selection drives divergence of eye morphology in sympatric Heliconius
790 butterflies. Evolution. qpae073. https://doi.org/10.1093/evolut/qpae073

791 Wright DS, Merrill RM. 2025a. Protocol: Lepidoptera Eye Dissection & Mounting. osf.io/ewbs9.
792 https://doi.org/10.17605/OSF.IO/EWBS9

793 Wright DS, Merrill RM. 2025b. Protocol for ImageJ/FIJI analysis of eye, tibia and abdomen
794 measurements of Lepidoptera. osf.io/fm8vp. https://doi.org/10.17605/OSF.IO/FM8VP

795 Young FJ et al. 2024. Enhanced long-term memory and increased mushroom body plasticity in
796 Heliconius butterflies. iScience. 27(2):108949. https://doi.org/10.1016/j.isci.2024.108949

797 Young FJ, Montgomery SH. 2020. Pollen feeding in *Heliconius* butterflies: the singular evolution
798 of an adaptive suite. Proc R Soc B. 287(1938):20201304.
799 https://doi.org/10.1098/rspb.2020.1304

800 Zhu H et al. 2025. Temporal variation in floral scent emission of a woody plant and flower
801 visiting behaviour of male and female flies. Oikos. 2025(5):e10727.
802 https://doi.org/10.1111/oik.10727

803 Ziemba K, Rutowski R. 2000. Sexual Dimorphism in Eye Morphology in a Butterfly
804 (Asterocampa leilia; Lepidoptera, Nymphalidae). Psyche: A Journal of Entomology. 103.
805 https://doi.org/10.1155/2000/54503

806