Spectral network analysis illuminates coordinated trait adaptation across plant populations

Rishav Ray[1,2], Brandie Quarles-Chidyagwai[2,3], Jessica Lyons[1], Sarah Ashlock[3], Jennifer R. Gremer[3], Julin N. Maloof[2], Troy S. Magney[1,4]

1. Department of Plant Sciences, University of California Davis
2. Department of Plant Biology, University of California Davis
3. Department of Ecology and Evolution, University of California Davis
4. Department of Forest Management, University of Montana

Summary

- Understanding how plant populations respond to environmental variation through functional leaf traits remains challenging due to limitations of traditional phenotyping approaches. Hyperspectral reflectance offers a rapid, non-destructive and high-throughput method to capture functional trait variation and detect signatures of local adaptation across populations.

- We combined hyperspectral data, inverse modeling, and network analysis to investigate population-level adaptations in *Streptanthus tortuosus*. Using a common garden experiment with four populations, we applied partial least square discriminant analysis (PLS-DA) and ridge regression for population discrimination, inverse PROSPECT modeling to estimate leaf biochemical traits, and canonical correlation analysis to examine trait-climate relationships across historical (1900-1994) and recent (1995-2024) periods. We developed a novel spectral network approach treating wavelength correlations as biologically meaningful trait networks.

- Populations showed distinct, heritable spectral signatures with high classification accuracy. Significant population differences emerged in anthocyanins, carotenoids, chlorophyll, and water content. Trait-climate correlations shifted between time periods, consistent with historical climate adaptation. Network analysis revealed population-specific integration patterns, with more variable environments displaying greater spectral modularity.

30   ● Hyperspectral signatures provide a high-throughput tool for detecting population-level
31   adaptation and trait coordination. Our findings suggest plant populations respond to
32   climate change through evolved shifts in trait networks rather than isolated traits alone.

Introduction

Climate change is driving unprecedented environmental shifts that are reshaping selective
pressures on plant populations worldwide (Gienapp *et al.*, 2014; Edelsparre *et al.*, 2024), yet our
ability to detect and understand evolutionary responses to these changes remain limited by
traditional approaches of studying adaptation (Franks *et al.*, 2014; Anderson & Gezon, 2015).
While common garden experiments and reciprocal transplants have provided foundational
insights into local adaptation (Leimu & Fischer, 2008; Hereford, 2009; Schwinning *et al.*, 2022),
these methods are often constrained by logistical challenges, long generational times, and the
difficulty of measuring multiple traits simultaneously across large populations (Merilä &
Hendry, 2014; Song *et al.*, 2021). The growing availability of genomic data has revolutionized
our understanding of genetic basis of adaptation (Savolainen *et al.*, 2013; Tigano & Friesen,
2016), but a critical gap remains between high resolution genomic data and the complex, multi
dimensional phenotypic data that are targets of natural selection (Houle *et al.*, 2010; Furbank &
Tester, 2011). This phenotyping bottleneck is particularly acute in the study of climate
adaptation, where responses may involve coordinated changes across multiple physiological and
biochemical traits rather than simple modifications of individual traits (Nicotra *et al.*, 2010;
Anderson *et al.*, 2012; Hein *et al.*, 2021). Consequently, there is an urgent need for high
throughput, integrative approaches that can capture the complexity of adaptive phenotypes while
scaling to the spatio-temporal dimensions over which climate driven evolution operates (Des
Roches *et al.*, 2018; Cavender-Bares *et al.*, 2022).
53
Hyperspectral reflectance analysis offers a promising solution to these challenges by providing a
rapid, non-destructive window into plant functional traits that integrates information across
multiple physiological and biochemical processes (Ustin & Gamon, 2010; Cavender-Bares *et al.*,
2016). Unlike traditional vegetation indices that rely on a few broad spectral bands,
hyperspectral data captures reflectance across hundred to thousands of narrow wavelength
intervals, encoding detailed information about leaf pigment composition, water content, and

structural properties (Kokaly *et al.*, 2009; Serbin *et al.*, 2014). This rich spectral information can

be decoded using models run in inverse mode such as PROSPECT, which enable the quantitative

estimation of specific biochemical traits, such as chlorophyll, carotenoids, and anthocyanins etc.

directly from reflectance spectra (Jacquemoud & Baret, 1990; Féret *et al.*, 2021). Recent

advances have demonstrated the power of hyperspectral approaches for detecting functional

diversity within and among plant communities (Asner *et al.*, 2015; Schweiger *et al.*, 2018; Wong

et al., 2023; Li *et al.*, 2023; Pau *et al.*, 2025), mapping plant stress responses (Gamon *et al.*,

2019), linking spectra to gene expression (Chen *et al.*, 2025), and even predicting ecosystem

function from spectral diversity patterns (Wang *et al.*, 2018; Cavender-Bares *et al.*, 2022).

However, most applications have focused on species-level differences or broad ecological

patterns, with limited exploration of how hyperspectral signatures might reveal population-level

adaptation and the evolutionary processes that shape functional trait variation within species

(Singh *et al.*, 2015; Meireles *et al.*, 2020). Furthermore, traditional analyses treat spectral bands

as independent variables, potentially missing emergent properties that can be gleaned from the

coordinated relationship between them (Gamon *et al.*, 2020).

Understanding how plants respond to environmental variation requires moving beyond

individual traits to consider the coordinated relationships among multiple functional

characteristics—a perspective rooted in the concept of phenotypic integration from quantitative

genetics and evolutionary developmental biology (Cheverud, 1996; Armbruster *et al.*, 2014).

Traits rarely evolve in isolation; instead, they are connected through genetic correlations (linkage

disequilibrium), developmental constraints, and functional dependencies that result in observable

phenotypes (Wagner & Altenberg, 1996; Klingenberg, 2008). The structure of these trait

correlations itself is an evolvable property that can be shaped by natural selection, with different

patterns of integration. This can range from highly modular to tightly integrated – representing

alternative adaptive strategies for responding to environmental challenges (Hansen & Houle,

2008; Murren, 2012). Network theory has emerged as a powerful framework for understanding

such complex systems across biology, from gene regulatory networks that control development

(Barabási & Oltvai, 2004; Davidson & Erwin, 2006) to ecological networks that structure

community interactions (Bascompte & Jordano, 2007; Olesen *et al.*, 2007). In each case, network

topology – the pattern of connections among nodes – provides insights into system function,

stability, and evolutionary potential that are undetectable when components are studied in

isolation (Albert & Barabási, 2002; Proulx *et al.*, 2005). However, applying network approaches

to study phenotypic integration has been limited by the challenge of measuring sufficient

numbers of traits to characterize network structure, particularly in studies of natural populations

where high-throughput phenotyping methods are essential (Murren, 2012; Messier *et al.*, 2017).

Hyperspectral data, with its hundreds of wavelength measurements that reflect coordinated

physiological processes, offers an unprecedented opportunity to examine trait networks at high

resolution and test fundamental hypotheses about how environmental variation shapes the

evolution of phenotypic integration (Cavender-Bares *et al.*, 2016; Schweiger *et al.*, 2018).

The California mountain jewelflower (*Streptanthus tortuosus*) provides an ideal system for

testing these novel approaches, as populations across California's diverse climatic gradients have

been shown to exhibit local adaptation in multiple functional traits including germination time,

drought tolerance, and life history strategies (Gremer *et al.*, 2020b,a). This species' wide

distribution across elevation and precipitation gradients, combined with common garden

experiments, makes it particularly suitable for investigating how environmental variation drives

adaptive landscape of complex trait architecture (Gremer *et al.*, 2020a; Bontrager *et al.*, 2025).

In this study, we combine multivariate hyperspectral reflectance analysis, inverse modeling, and

network theory to investigate population-level climate adaptation in *S. tortuosus.* Using a

common garden experiment spanning four *S. tortuosus* populations, we test whether spectral

reflectance data can detect heritable variation in functional traits and reveal population-specific

patterns of trait coordination. We apply a novel two-step analytical framework – partial least

square discriminant analysis (PLS-DA) for population discrimination followed by ridge

regression analysis to identify key wavelength regions – to characterize population specific

spectral signatures. Through inverse PROSPECT modeling, we estimate six key leaf functional

traits and examine whether their relationships with historical and recent climate variables are

correlated using canonical correlation analysis across two temporal scales. We then developed a

spectral network approach that treats wavelength correlations as biologically meaningful trait

networks, revealing distinct patterns of spectral integration among populations. Finally, we test

whether these network architectures correlate with long-term climate variability patterns,

providing insights into how environmental variation correlates with both individual traits and
their coordination.

Materials and Methods

Plant material and common garden setup

We conducted a common garden experiment at the UC Davis Vegetable Crops fields in Davis,
CA USA, with seeds collected from four populations of *Streptanthus tortuosus*. Seedlings from
these populations were planted at the garden in raised beds with 21-98 replicates per population.
Seeds were stratified for four weeks at 4°C in the dark in a cold room at UC Davis. On October
30, 2023, after stratification, plants were moved to a growth chamber with 12-hour day length
and warm light conditions (light intensity ~100uMol/microeinstein) to stimulate germination.
Plants were initially exposed to 22°C day/8°C night temperatures. To harden plants to the cold
temperatures of the field, temperatures were adjusted accordingly: 20°C day/7°C night 7-days
prior to field transplant, 19°C day/6°C night 6-days prior to field transplant, and 18°C day/5°C
night 5-days prior to field transplant. Seedlings were transplanted into a randomized block design
with 12 blocks spread across 6 raised beds and 30-cm between experimental plants. Beds were
12" tall, 110cm wide, and 14m in length; there was 1m between each bed. Beds were filled with
Ron's mix, sand, and perlite. The beds were watered with drip irrigation the day before
transplant for 5 hours, during the afternoon after planting for 5 hours, and five days after
transplant for 5 hours.

Hyperspectral data collection and processing

Leaf-level hyperspectral reflectance measurements were collected from plants grown in the
common garden using a full-range spectrometer (HR-1024i; Spectra Vista Corporation—SVC).
Spectral reflectance was measured from 400 to 2500 nm with a spectral resolution of 1.5 nm
(250–1000 nm), 3.8 nm (1000–1890 nm), and 2.5 nm (1890–2500 nm). Leaf spectra were
collected with a leaf clip (LC-RP PRO; SVC) attached to a fiber optic cable. The leaf clip houses
an internal tungsten halogen lamp that was standardized for reflectance using a built-in
Spectralon white reference panel (Labsphere Inc.). Leaf measurements were carried out on
flowering plants on young fully expanded leaves that were big enough to fit the leaf clip.

Reference measurements were made before every plant and dark current corrections were

automatically taken following each scan with an internal shutter. Spectra were then averaged

across three measurements per leaf and smoothed using a Savitzky-Golay filter.

Statistical analysis

Partial Least Squares Discriminant Analysis (PLS-DA) was performed (*mixOmics* (Rohart *et al.*,

2017) package in R) to assess population-level differentiation in hyperspectral space. PLS-DA

was chosen because it is specifically designed to handle high-dimensional datasets where the

number of variables (wavelengths) exceeds the number of samples, a common challenge in

hyperspectral analysis. PLS-DA reduces dimensionality while maximizing the separation

between groups, making it ideal for identifying population-specific spectral discrimination. It

was followed by a PERMANOVA test (*vegan* package (Oksanen *et al.*, 2025)) to statistically test

for differences in spectral data among populations. PLS-DA identified canonical variates that

maximally separated the four populations while explaining spectral variance. Broad-sense

heritability (H²) was calculated for each wavelength using variance components from a mixed-

effects model using the *lme4* package with population as a random effect: $H^2 = \sigma^2G/(\sigma^2G + \sigma^2E)$,

where σ^2G is genetic variance among populations and σ^2E is environmental/residual variance

within populations.

Ridge regression models were fitted for each population(*glmnet* package (Tay *et al.*, 2023)) to

identify wavelength-specific contributions to population classification while avoiding overfitting

that commonly occurs with high-dimensional spectral data. The L2 regularization in ridge

regression shrinks coefficients toward zero but retains all wavelengths, allowing us to quantify

the relative importance of different spectral regions for population classification. Spectral

coefficients were extracted across all wavelengths and integrated within six functionally relevant

spectral regions (carotenoids: 450-520 nm, chlorophyll: 650-700 nm, NIR plateau: 700-1300 nm,

red edge: 680-750 nm, water: 1400-1500 nm, water/protein: 1900-2000 nm). Receiver Operating

Characteristic (ROC) curves were generated using cross-validation, and Area Under the Curve

(AUC) values were calculated to assess classification performance.

Inverse PROSPECT modeling was applied using the *prospect* R package (Féret & Boissieu,

2024) to estimate six leaf functional traits: anthocyanin content, carotenoid content, chlorophyll

content, leaf mass area, structural parameter (N), and equivalent water thickness. Climate data

for each population's origin site were obtained from Flint Basin Characterization Model (Flint *et*

al., 2021) (Fig. S2), spanning 1900-2024 and divided into historical (1900-1994) and recent

(1995-2024) periods. Canonical Correlation Analysis (CCA) was performed (*mixOmics* package)

to examine relationships between estimated leaf traits and climate variables across different

temporal periods. CCA examines multivariate relationships between multiple leaf traits and

multiple climate variables simultaneously, capturing complex trait-environment associations that

would be missed by univariate approaches. It identifies linear combinations of traits and climate

variables that are maximally correlated, revealing coordinated trait response to environmental

gradients.

For each population, correlation matrices were calculated between all wavelength pairs using

Pearson correlation coefficients. To account for sample size limitations, we employed a bootstrap

approach, randomly sampling 10 individuals with replacement 100 times and constructing

networks for each iteration. Edges were retained only for correlations exceeding $|r| > 0.7$. Four

key network parameters were calculated using the *igraph* package: (1) Density - the proportion

of possible edges that are actually present in the network, calculated as the number of observed

edges divided by the maximum possible number of edges (n(n-1)/2 for undirected networks).

Values range from 0 (no connections) to 1 (fully connected), with higher values indicating more

integrated spectral responses across wavelengths. (2) Centralization - a measure of how

unequally distributed the connections are across nodes, quantifying the extent to which the

network is dominated by highly connected "hub" wavelengths. Centralization is calculated based

on the variance in node degrees relative to a star network (maximum centralization) and ranges

from 0 (all nodes equally connected) to 1 (one central hub connected to all other nodes). High

centralization indicates that spectral integration is controlled by a few key wavelengths. (3)

Modularity - a measure of community structure that quantifies how well the network can be

divided into distinct groups of densely connected nodes (modules) with sparse connections

between groups. Modularity was calculated using the Louvain algorithm, which optimizes the

modularity function $Q = \Sigma[A_{ij} - (k_i k_j/2m)]\delta(c_i,c_j)$, where A_{ij} is the adjacency matrix, k_i is the

degree of node i, m is the total number of edges, and $\delta(c_i,c_j) = 1$ if nodes i and j belong to the

same community. Values range from -0.5 to 1, with higher values indicating stronger modular

organization of spectral responses. (4) Transitivity - (clustering coefficient) - the probability that

two nodes connected to a common neighbor are also connected to each other, measuring the
tendency for wavelengths to form tightly connected clusters. Calculated as the ratio of observed
triangles to possible triangles in the network: $C = 3 \times$ (number of triangles) / (number of
connected triples). Values range from 0 to 1, with higher values indicating greater local
clustering of spectral correlations.
CCA was performed between population-level network metrics and long-term climate variability
measures (coefficients of variation for temperature, precipitation, and climatic water deficit from
1900-2024) to test relationships between network topology and environmental variation.

All analyses were performed in R (version 4.2.1 (R Core Team, 2022)). Significance testing for
network metrics was conducted using ANOVA with Tukey's HSD post-hoc tests. Bootstrap
confidence intervals were calculated for all network parameters. Statistical significance was set
at $\alpha = 0.05$, with Bonferroni correction applied for multiple comparisons where appropriate.

222 Results

223 Population-level hyperspectral differentiation and genetic variation

Fig. 1: Hyperspectral variation and heritability in *Streptanthus tortuosus* populations across
California. A) Partial Least Squares Discriminant Analysis (PLS-DA) of hyperspectral data showing four
S. tortuosus populations (TM2, CC, DPR, and BH) along two principal variates. The x-variate explains
65.62% of the spectral variation while the y-variate explains 32.31%. **B)** Map of California showing the
geographic distribution of the sampled populations: Table Mountain (TM2), Canyon Creek (CC), Drum
Powerhouse Road (DPR), Ben Hur (BH) and the UC Davis Common Garden (UCD) where all these
populations were planted and measured. **C)** Reflectance spectra by population across wavelengths from
400-2500 nm. Different coloured lines represent individual plant measurements. The red line (labeled
"cv") shows the coefficient of variation across wavelengths (right y-axis). **D)** Broad-sense heritability
(H^2) estimates across wavelengths from 400-2500 nm, showing variation in the genetic contribution to

spectral reflectance traits. The coloured bands indicate spectral regions that correlate with leaf chlorophyll content (green), and water/protein content (blue).

To determine whether *S. tortuosus* populations maintain distinct spectral signatures under standardized common garden conditions, we performed PLS-DA on the hyperspectral data over a multivariate space. The PLS-DA ordination demonstrated significant differences in spectral values among populations (PERMANOVA F:17.43, p-value:0.001), with the first two canonical variates explaining 97.93% of the total spectral variation (Fig. 1A, S1). The populations formed distinct clusters in the ordination space, with TM2 and DPR populations showing greatest

separation along the first canonical variate, while the BH population separated in a distinct

cluster along the second canonical variate consistent with its geographical separation (Fig. 1B).

Despite being measured under the same common garden conditions, the samples showed spectral

variation across key regions of the wavelength range, denoted by the coefficient of variation

(CV) plotted in the second axis (Fig. 1C). The CV across samples peaked in several key spectral

regions, with the highest variability observed around 680 nm (chlorophyll absorption), and water

absorption bands around 1400 nm and 1900 nm, regions that correspond to functionally

important leaf properties. Broad-sense heritability (H^2) – which estimates the amount of

phenotypic variation explained by genotype – revealed substantial genetic control over spectral

reflectance traits (Fig 1D). The H^2 values peak considerably across the chlorophyll and water

absorption regions, suggesting that the genetic variation is concentrated in spectral regions that

reflect key physiological and biochemical leaf properties and that natural selection may have

acted on these functionally relevant traits detected by the spectral measurements. Taken together,

maintenance of population-specific spectral signatures under common garden conditions,

combined with high heritability estimates, provides strong evidence for genetic differentiation in

leaf optical properties among *S. tortuosus* populations.

Ridge regression analysis reveals population-specific spectral signatures

Fig. 2: Spectral characteristics and classification performance across *Streptanthus tortuosus*

populations. A) Spectral coefficients derived from ridge regression across wavelengths (400-2500 nm)

for four *S. tortuosus* populations (TM2, DPR, CC, and BH). Background shading highlights key spectral

regions associated with different leaf properties: carotenoids, chlorophyll, NIR plateau, red edge, water,

and water/protein. **B)** Receiver Operating Characteristic (ROC) curves showing classification

performance for each population based on ridge regression models. All populations demonstrate excellent

classification accuracy with Area Under the Curve (AUC) values ranging from 0.971 (CC) to 1.0 (BH),

indicating high discriminative power of spectral data for population identification. **C)** Integrated

coefficients rank plot showing the relative importance of six key spectral regions in distinguishing

between populations. Each colored line represents a population (TM2, DPR, CC, and BH), with vertical positions indicating their rank (higher = stronger signal) for each spectral region.

Having established that there are heritable differences among populations, we next asked, what are the specific wavelengths that distinguish between the populations? To answer this, we applied ridge regression analysis and quantified the predictive power of hyperspectral data for population classification. Regression coefficients across the spectra (referred to as spectral coefficients hereafter) varied markedly across wavelengths and among populations, with each population exhibiting unique patterns of spectral importance (Fig. 2A). All populations showed pronounced spectral coefficients in the visible region (400-700nm), particularly in areas corresponding to carotenoid (450-520 nm), and chlorophyll (650-700nm) absorption bands. Notable population specific differences emerged in the water absorption regions (1400-1500 and 1900-2000 nm), where DPR displayed stronger spectral coefficients compared to other populations, which varied in those regions (Fig 2C). The near-infrared region (700-1300 nm) showed distinct coefficient patterns, with BH exhibiting particularly strong signals in this

structurally-sensitive spectral region (Fig. 2A). We numerically integrated the spectral
coefficients across six functionally relevant wavelength regions and ranked them to quantify the
population specific spectral "fingerprints" (Fig. 2C). TM2 consistently ranked highest in three
key regions: chlorophyll, carotenoids, and water/protein absorption bands. In contrast, BH was
most sensitive to the near infra-red (NIR) region, indicating unique leaf structural properties that
affect light scattering in this wavelength range. CC and DPR populations showed more moderate
rankings across most spectral regions, with CC ranking highest in the one of the water absorption
regions and DPR showing intermediate values across all spectral domains. Furthermore, the
discriminative power of hyperspectral data for population identification was high (AUC 0.97-1)
across all populations, indicating that hyperspectral reflectance data contain sufficient
population-specific information to achieve high classification accuracy (Fig 2B). In summary,
these results suggest clear population-specific differentiation in spectral signatures, indicating
that populations vary in multiple optical properties..

Leaf trait differences and climate associations across temporal scales

Fig. 3: Leaf trait variation across populations and relationships with climate variables. A) Mean
values (±SE) and F-statistic and p-value of ANOVA test of six key leaf traits across four *S. tortuosus*
populations (BH, CC, DPR, and TM2). Traits include anthocyanins (μg/cm²), carotenoids (μg/cm²),
chlorophyll (μg/cm²), leaf mass area (g/m²), leaf structure constant (N), and water content (g/cm²)**. B-C)**
Canonical Correlation Analysis (CCA) biplot showing relationships between leaf traits (blue vectors) and
climate variables (gray vectors). Timeline at bottom indicates the historical (1900-1994) and recent
(1995-2024) climate periods used in the analysis. **B)** The first canonical variate explains the primary
correlation structure between leaf traits and recent climate conditions. **C)** CCA biplot showing
relationships between leaf traits and historical climate variables (1900-1994). **D)** CCA biplot showing

relationships between leaf traits and climate change variables (difference between recent and historical periods).

We next asked whether the variation in reflectance in different populations can be explained by various home climatic variables that they have experienced. To that end, we applied the inverse PROSPECT (Jacquemoud & Baret, 1990; Féret & Boissieu, 2024) model to estimate six key leaf

functional traits from the hyperspectral data, namely: anthocyanins, carotenoids, chlorophyll, leaf mass area, N (leaf structural complexity), and equivalent water thickness (Fig 3A). The traits show significant population-level variation, e.g. anthocyanin contents were highest in BH and CC populations, exceeding levels in DPR and TM2 populations by about two-fold. Carotenoid content showed the opposite pattern, with TM2 displaying markedly higher concentrations compared to other populations, and not detectable from BH spectra. Chlorophyll content was highest in CC, while BH showed substantially lower levels. Water content also varied significantly among populations, with CC again showing the highest values compared to BH. Leaf mass area and the structural parameter N showed more modest but consistent population differences, indicating coordinated variation in both biochemical and structural leaf properties.

To test for environmental associations with these estimated traits, we applied canonical correlation analysis with populations' home climatic variables going back to 1900, split between recent (1995-2024), and historic (1900-1994) time periods. The analysis revealed distinct patterns of trait-climate associations across different temporal periods. For recent climate conditions (1995-2024), the first canonical correlation explained strong relationships between leaf traits and contemporary environmental variables (Fig. 3B). Chlorophyll content and equivalent water thickness correlated positively with recent actual evapotranspiration and precipitation and negatively with climate water deficit. Historical climate relationships (1900-1994) showed similar trait-environment associations compared to recent patterns (Fig. 3C). However, climate change analysis (difference between recent and historical periods) provided insights into how shifting environmental conditions correlate with current trait variation (Fig. 3D). Anthocyanins showed a strong positive correlation with change in maximum temperature, and interestingly, chlorophyll, and equivalent water thickness all correlated positively with changes in precipitation and minimum temperature patterns. These temporal patterns reveal shifting trait-climate correlations across historical and recent time periods, with different climate variables showing the strength and direction of these statistical associations vary across different time intervals..

Spectral network architecture reveals population-specific network patterns linked

to climate variability

Fig. 4: Spectral network analysis and climate-network relationships in *S. tortuosus* populations. A)

Network visualization of spectral relationships in four *S. tortuosus* populations (TM2, BH, DPR, and CC).

Each node represents a wavelength, and edges represent correlations between wavelengths. **B)**

Comparison of four network parameters (centralization, density, modularity, and transitivity) across

populations. Box plots show the distribution of bootstrapped values (n=100, 10 individuals with

replacement per iteration) with significance indicators for population differences (*p<0.05, **p<0.01,

***p<0.001, ****p<0.0001). **C)** CCA biplot showing relationships between network metrics (blue

vectors) and climate variability parameters (gray vectors) from 1900-2024.

We then asked whether the leaf reflectance patterns are co-ordinated amongst themselves and the

relationship between them are population specific or not. To that end, we applied a network

approach to build a spectral graph, where each node represents a particular wavelength, and each

connecting edge is the correlation coefficient of their reflectance value between the two nodes.

This network analysis of wavelength correlations revealed distinct patterns of spectral integration

that varied systematically among *S. tortuosus* populations (Fig. 4A). Visual inspection of

network topologies showed clear population-specific architectures: TM2 and DPR populations

exhibited dense, highly interconnected networks with numerous edges connecting wavelengths

across the full spectral range, indicating strong coordination among diverse optical traits. In

contrast, BH and CC populations displayed more modular network structures characterized by

distinct clusters of highly connected wavelengths separated by sparser connections, suggesting

compartmentalized spectral responses. Quantitative analysis of network parameters confirmed

significant population differences in spectral integration patterns (Fig. 4B). Both TM2 and DPR

showed significantly higher network density and degree compared to BH and CC (p < 0.0001),

indicating greater overall connectivity among wavelengths. CC exhibited the highest

centralization values, suggesting that spectral integration in this population is dominated by a

few highly connected wavelength "hubs." Modularity was significantly higher in CC and BH

populations compared to TM2 and DPR, confirming the visual impression of more

compartmentalized spectral organization. Transitivity showed more modest but significant

differences, with DPR displaying the highest values, indicating greater local clustering of

wavelength correlations. These results demonstrate that populations have fundamentally different

approaches to organizing spectral trait correlations, ranging from highly integrated (TM2, DPR)

to highly modular (CC, BH) network architectures suggesting population-specific variation in

the coordination of leaf optical properties.

Canonical correlation analysis revealed that these population differences in spectral network

organization are systematically related to long-term climate variability patterns (Fig. 4C). The

first canonical variate separated density from the other network parameters, while the second

canonical variate distinguished centralization and modularity and transitivity. The AET/PET

ratio correlated highly with density, suggesting that populations from environments with higher

water availability (higher AET/PET) tend to have denser, more integrated spectral networks.

Conversely, centralization and modularity both loaded positively on the second canonical variate

and were associated with minimum and maximum temperature variability, indicating that

populations from thermally variable environments have more centralized and modular spectral

integration patterns. Furthermore, The coefficient of variation in climatic water deficit (CWD

CV) loaded negatively on the first canonical variate, suggesting that populations experiencing

greater water stress variability are associated with lower transitivity and higher density in their

spectral networks.

Discussion

This study demonstrates the power of integrating advanced hyperspectral analysis with network

theory to reveal previously hidden patterns of plant trait coordination. Our novel analytical

framework discriminated four *S. tortuosus* populations with good accuracy, while revealing that

spectral traits itself represent a heritable characteristic. The combination of PLS-DA for

population discrimination and ridge regression for identifying key wavelength regions proved

highly effective, with population specific signatures maintained even under common garden

conditions. Beyond individual trait variation, our network analysis uncovered distinct patterns of

spectral coordination among populations, ranging from highly connected (TM2) to modular (CC,

BH) architectures that correlate with long-term climate variability. Critically, these

methodological advances revealed that populations have evolved not only different leaf

biochemical properties—with notable differences in anthocyanins, carotenoids, and chlorophyll

content—but also fundamentally different strategies for coordinating these traits reflected

through their spectral networks. Strong correlations between network topology and historical

climate patterns suggest that environmental variability has shaped both individual traits and their

patterns of integration, representing a new perspective on the evolution of complex trait

architecture in natural populations.

Building on these climate network relationships, specific spectral regions provide additional

insights into the functional basis of trait variation. The near-infrared region (700-1300 nm)

showed distinct coefficient patterns, with BH exhibiting particularly strong signals in this

structurally-sensitive spectral region. The NIR plateau is primarily influenced by leaf internal

structure rather than biochemical composition, with reflectance determined by the scattering of

light at air-cell wall interfaces within the leaf mesophyll (Jacquemoud & Baret, 1990; Féret *et*

al., 2021). Higher NIR reflectance typically indicates greater leaf thickness, increased

intercellular air spaces, or altered mesophyll organization—structural characteristics that can

influence light capture efficiency, gas exchange, and water use efficiency (Onoda *et al.*, 2017).

The strong NIR signal in BH may reflect population-level variation in structural traits that could

be related to environmental conditions at this population's origin site (Niinemets, 2001; Wright

et al., 2004; Karabourniotis *et al.*, 2021). However, it is important to emphasize that our study

cannot distinguish between spectral variation arising from adaptive response to local

environment versus genetic drift, founder effects, or population structure. While the functional

significance of the estimated traits correlating with the environmental variables suggests that

they may be ecologically relevant, we make no causal claims about the evolutionary mechanisms

that drive these patterns.

Our hyperspectral network analysis represents a shift from traditional approaches that treat

spectral bands as independent variables to a systems-level framework that captures the

coordinated relationship between wavelengths. While previous studies have relied primarily on

vegetation indices or individual wavelength analysis (Bannari *et al.*, 1995; Thenkabail *et al.*,

2000; Gamon *et al.*, 2019), our network approach reveals emergent properties of spectral trait

integration that are not captured by conventional methods. Our two-step analytical pipeline –

combining PLS-DA for population discrimination with ridge regression for mechanistic insight –

offers significant advantages over single method approaches by first establishing the existence of

population differences and then identifying the specific wavelength regions driving these

patterns. This framework addresses a key gap in evolutionary ecology, where high throughput

phenotyping methods are needed to match the resolution of genomic tools (Houle *et al.*, 2010;

Furbank & Tester, 2011; Pieruschka & Poorter, 2012; Cavender-Bares *et al.*, 2022).The

integration of canonical correlation analysis across multiple temporal scales (historic vs. recent

climate) further demonstrates how advanced statistical methods can detect trait correlation with
environmental variables. This study provides a template for studying evolutionary responses to
environmental change, which can be further bolstered by addition of genomic and transcriptomic
data , to study adaptation in real-time (Hoffmann & Sgrò, 2011; Anderson *et al.*, 2012; Franks *et*
al., 2014; Merilä & Hendry, 2014).
447
Additionally, our spectral network analysis reveals a previously unrecognized level of biological
organization – the coordinated structure of the reflectance patterns – that parallels network
organization observed across diverse biological systems. While reflectance patterns themselves
are not direct targets of selection, they represent emergent properties of functionally important
traits such as pigment composition, water retention capacity, and leaf structural characteristics
that can be under direct selective pressure. Similar to how gene regulatory networks exhibit
modular architectures that facilitate evolvability while maintaining stability (Wagner *et al.*, 2007;
Espinosa-Soto & Wagner, 2010), our spectral networks show population-specific patterns of
modularity and integration that are consistent with different strategies for coordinating
underlying physiological and biochemical processes.. The dense, highly connected networks in
TM2 and DPR populations mirror the "small-world" properties observed in metabolic networks,
where high clustering facilitates local optimization while maintaining global connectivity (Jeong
et al., 2000; Barabási & Oltvai, 2004). Conversely, the modular architectures in CC and BH
populations resemble the compartmentalized organization of ecological food webs, where
modularity enhances stability against perturbations while allowing independent evolution of
subsystems (Olesen *et al.*, 2007; Stouffer & Bascompte, 2011). This biological network
perspective suggests that natural selection acts not only on individual functional traits but also on
the patterns of coordination among these traits, with spectral networks providing a high-
dimensional window into this complex trait architecture, as proposed in quantitative genetics
theory (Cheverud, 1996; Armbruster *et al.*, 2014). The correlation between network topology and
climate variability patterns supports the hypothesis that environmental unpredictability favors
modular organization—allowing rapid reconfiguration of trait combinations—while stable
environments favor integrated networks that optimize coordinated responses (Schlosser &
Wagner, 2004). Our findings thus extend network thinking beyond its traditional domains in
molecular biology and ecology to encompass the evolution of complex phenotypes in the

spectral sensing domain. This network perspective offers practical applications for conservation

biology, as species with highly integrated spectral networks (low modularity, high density) may

be more vulnerable to environmental disruption due to their tightly coupled trait systems.

While our methodological framework demonstrates significant promise, several limitations

warrant consideration. First, our analysis was restricted to a single growing season under

common garden conditions, which does not fully capture the dynamic of spectral trait expression

across environments or temporal variation. This is highlighted by the absence of detectable

carotenoid levels in the BH population using the inverse modeling. This highlights potential

limitations in our inverse PROSPECT modeling approach for certain biochemical compounds

under specific conditions. This could reflect either genuine biological differences in carotenoid

accumulation strategies or methodological constraints in detecting low-concentration pigments in

certain leaf types. Additionally, while our network analysis reveals compelling patterns of

spectral integration, the biological mechanisms underlying these coordination patterns remain

unclear. Future studies combining transcriptomic and metabolomic approaches with

hyperspectral analysis could provide mechanistic insights into how plants achieve different

integration strategies.

The successful implementation of this analysis framework requires careful consideration of

several methodological parameters and quality control practices that can significantly influence

results. Key analytical choices include spectral smoothing window size (we used Savitzky-Golay

filtering with a 5-point window), correlation threshold for edge retention ($|r| > 0.7$), and number

of bootstrap replicates ($n = 100$). Each of these parameters represent a trade-off between noise

reduction, information retention and computational complexity that should be optimized for

specific dataset and research questions. Sample size remains a critical consideration, as our

analysis with four populations represents a minimum threshold for detecting network patterns.

The small number of population and relatively small sample size used in this study also limits

our ability to generalize across the species' full geographic range, and expand to additional

populations across broader gradients. Future studies with a large sample size would significantly

strengthen our conclusions about climate trait relationships. Field measurement quality control is

another essential component and should include consistent leaf positioning, regular white

reference calibration, and systematic checking for spectral artifacts or outliers that could
propagate through network analyses. Importantly, network metrics (density, modularity,
centralization, transitivity) and biochemical trait estimates should be interpreted comparatively
between treatments or populations rather than as absolute biological values, as their biological
meaning emerges from relative patterns rather than specific numerical thresholds. This analytical
framework shows considerable promise for scaling to larger spatial and temporal dimensions:
canopy-level hyperspectral imaging could extend population comparisons to landscape scales,
while integration with emerging airborne and satellite hyperspectral sensors could enable real-
time monitoring of trait network evolution across entire species ranges.

Conclusion

This study demonstrates how hyperspectral analysis can be integrated with network theory to
understand population-level trait coordination in plants. Our analytical framework reveals that
spectral data encode information about both individual functional traits and their patterns of
coordination, providing a high-throughput approach to studying complex phenotypes in natural
populations. Future integration of this spectral framework with genomic and transcriptomic
datasets could illuminate the molecular mechanisms underlying different coordination strategies,
potentially identifying the genetic basis of trait integration and informing predictive models of
population responses to environmental change. Such multi-scale applications will be essential for
predicting and monitoring how natural populations respond to accelerating environmental
change.

Acknowledgements

525 This research was funded by the National Science Foundation (DEB-2129589 to J.N.M, J.R.G,
526 T.S.M, and Denneal Jamison-McClung). We acknowledge the contribution from Maya Arakaki,
527 technician Kate Ruiz-Cox, and Carlos Perez and all the undergraduate students involved in
528 setting up the field and data collection. J. Schmitt provided valuable guidance on study design,
529 analyses, and manuscript drafts.

Competing interests

The authors declare no competing interests.

Author contribution

Conceptualization (research direction and analytical framework): RR, JNM, TSM, (common garden and data collection) RR, BQ-C, JRG, JNM, TSM; Data curation: RR, JL; Formal Analysis: RR, JNM, TSM; Funding acquisition: JRG, JNM, TSM; Investigation: RR, JNM, TSM; Visualization: RR; Writing – original draft: RR; Writing – review & editing: RR, BQ-C, JL, SA, JRG, JNM, TSM

Data availability

Raw data and source code to reproduce all the analysis and figures would be available in Github upon publication.

Reference

Albert R, Barabási A-L. **2002**. Statistical mechanics of complex networks. *Reviews of Modern Physics* **74**: 47–97.

Anderson JT, Gezon ZJ. **2015**. Plasticity in functional traits in the context of climate change: a case study of the subalpine forb Boechera stricta (Brassicaceae). *Global Change Biology* **21**: 1689–1703.

Anderson JT, Inouye DW, McKinney AM, Colautti RI, Mitchell-Olds T. **2012**. Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. *Proceedings of the Royal Society B: Biological Sciences* **279**: 3843–3852.

Armbruster WS, Pélabon C, Bolstad GH, Hansen TF. **2014**. Integrated phenotypes: understanding trait covariation in plants and animals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**: 20130245.

Asner GP, Martin RE, Anderson CB, Knapp DE. **2015**. Quantifying forest canopy traits: Imaging spectroscopy versus field survey. *Remote Sensing of Environment* **158**: 15–27.

Bannari A, Morin D, Bonn F, Huete AR. **1995**. A review of vegetation indices. *Remote Sensing Reviews* **13**: 95–120.

Barabási A-L, Oltvai ZN. **2004**. Network biology: understanding the cell's functional
organization. *Nature Reviews Genetics* **5**: 101–113.

Bascompte J, Jordano P. **2007**. Plant-Animal Mutualistic Networks: The Architecture of
Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**: 567–593.

Bontrager M, Worthy SJ, Cacho NI, Leventhal L, Maloof JN, Gremer JR, Schmitt J,
Strauss SY. **2025**. Herbarium specimens reveal a constrained seasonal climate niche despite
diverged annual climates across a wildflower clade. *Proceedings of the National Academy of*
Sciences **122**: e2503670122.

Cavender-Bares J, Meireles JE, Couture JJ, Kaproth MA, Kingdon CC, Singh A, Serbin
SP, Center A, Zuniga E, Pilz G, *et al.* 2016. Associations of Leaf Spectra with Genetic and
Phylogenetic Variation in Oaks: Prospects for Remote Detection of Biodiversity. *Remote Sensing*
8: 221.

Cavender-Bares J, Schneider FD, Santos MJ, Armstrong A, Carnaval A, Dahlin KM,
Fatoyinbo L, Hurtt GC, Schimel D, Townsend PA, *et al.* 2022. Integrating remote sensing
with ecology and evolution to advance biodiversity conservation. *Nature Ecology & Evolution* **6**:
506–519.

Chen Y, Monks L, Rubio VE, Cox AJ, Swenson NG. **2025**. Linking leaf hyperspectral
reflectance to gene expression. *Communications Earth & Environment* **6**: 694.

Cheverud JM. **1996**. Developmental Integration and the Evolution of Pleiotropy. *American*
Zoologist **36**: 44–50.

Davidson EH, Erwin DH. **2006**. Gene Regulatory Networks and the Evolution of Animal Body
Plans. *Science* **311**: 796–800.

Des Roches S, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, Schweitzer JA,
Palkovacs EP. **2018**. The ecological importance of intraspecific variation. *Nature Ecology &*
Evolution **2**: 57–64.

Edelsparre AH, Fitzpatrick MJ, Saastamoinen M, Teplitsky C. **2024**. Evolutionary
adaptation to climate change. *Evolution Letters* **8**: 1–7.

Espinosa-Soto C, Wagner A. **2010**. Specialization Can Drive the Evolution of Modularity.
PLOS Computational Biology **6**: e1000719.

Féret J-B, Berger K, de Boissieu F, Malenovský Z. **2021**. PROSPECT-PRO for estimating
content of nitrogen-containing leaf proteins and other carbon-based constituents. *Remote Sensing*
of Environment **252**: 112173.

Féret J-B, Boissieu F de. **2024**. prospect: an R package to link leaf optical properties with their
chemical and structural properties with the leaf model PROSPECT. *Journal of Open Source*
Software **9**: 6027.

Flint LE, Flint AL, Stern MA. **2021**. *The basin characterization model—A regional water*
balance software package. U.S. Geological Survey.

Franks SJ, Weber JJ, Aitken SN. **2014**. Evolutionary and plastic responses to climate change
in terrestrial plant populations. *Evolutionary Applications* **7**: 123–139.

Furbank RT, Tester M. **2011**. Phenomics – technologies to relieve the phenotyping bottleneck.
Trends in Plant Science **16**: 635–644.

Gamon JA, Somers B, Malenovský Z, Middleton EM, Rascher U, Schaepman ME. **2019**.
Assessing Vegetation Function with Imaging Spectroscopy. *Surveys in Geophysics* **40**: 489–513.

Gamon JA, Wang R, Gholizadeh H, Zutta B, Townsend PA, Cavender-Bares J. **2020**.
Consideration of Scale in Remote Sensing of Biodiversity. In: Cavender-Bares J, Gamon JA,
Townsend PA, eds. Remote Sensing of Plant Biodiversity. Cham: Springer International
Publishing, 425–447.

Gienapp P, Reed TE, Visser ME. **2014**. Why climate change will invariably alter selection
pressures on phenology. *Proceedings of the Royal Society B: Biological Sciences* **281**: 20141611.

Gremer JR, Chiono A, Suglia E, Bontrager M, Okafor L, Schmitt J. **2020a**. Variation in the
seasonal germination niche across an elevational gradient: the role of germination cueing in
current and future climates. *American Journal of Botany* **107**: 350–363.

Gremer JR, Wilcox CJ, Chiono A, Suglia E, Schmitt J. **2020b**. Germination timing and
chilling exposure create contingency in life history and influence fitness in the native wildflower
Streptanthus tortuosus. *Journal of Ecology* **108**: 239–255.

Hansen TF, Houle D. **2008**. Measuring and comparing evolvability and constraint in
multivariate characters. *Journal of Evolutionary Biology* **21**: 1201–1219.

Hein NT, Ciampitti IA, Jagadish SVK. **2021**. Bottlenecks and opportunities in field-based
high-throughput phenotyping for heat and drought stress. *Journal of Experimental Botany* **72**:
5102–5116.

Hereford J. **2009**. A Quantitative Survey of Local Adaptation and Fitness Trade-Offs. *The*
American Naturalist **173**: 579–588.

Hoffmann AA, Sgrò CM. **2011**. Climate change and evolutionary adaptation. *Nature* **470**: 479–
485.

Houle D, Govindaraju DR, Omholt S. **2010**. Phenomics: the next challenge. *Nature Reviews*
Genetics **11**: 855–866.

Jacquemoud S, Baret F. **1990**. PROSPECT: A model of leaf optical properties spectra. *Remote*
Sensing of Environment **34**: 75–91.

Jeong H, Tombor B, Albert R, Oltvai ZN, Barabási A-L. **2000**. The large-scale organization

of metabolic networks. *Nature* **407**: 651–654.

Karabourniotis G, Liakopoulos G, Bresta P, Nikolopoulos D. **2021**. The Optical Properties of
Leaf Structural Elements and Their Contribution to Photosynthetic Performance and
Photoprotection. *Plants* **10**: 1455.

Klingenberg CP. **2008**. Morphological Integration and Developmental Modularity. *Annual*
Review of Ecology, Evolution, and Systematics **39**: 115–132.

Kokaly RF, Asner GP, Ollinger SV, Martin ME, Wessman CA. **2009**. Characterizing canopy
biochemistry from imaging spectroscopy and its application to ecosystem studies. *Remote*
Sensing of Environment **113**: S78–S91.

Leimu R, Fischer M. **2008**. A Meta-Analysis of Local Adaptation in Plants. *PLOS ONE* **3**:
e4010.

Li C, Czyż EA, Halitschke R, Baldwin IT, Schaepman ME, Schuman MC. **2023**. Evaluating
potential of leaf reflectance spectra to monitor plant genetic variation. *Plant Methods* **19**: 108.

Meireles JE, Cavender-Bares J, Townsend PA, Ustin S, Gamon JA, Schweiger AK,
Schaepman ME, Asner GP, Martin RE, Singh A, *et al.* 2020. Leaf reflectance spectra capture
the evolutionary history of seed plants. *New Phytologist* **228**: 485–493.

Merilä J, Hendry AP. **2014**. Climate change, adaptation, and phenotypic plasticity: the problem
and the evidence. *Evolutionary Applications* **7**: 1–14.

Messier J, McGill BJ, Enquist BJ, Lechowicz MJ. **2017**. Trait variation and integration across
scales: is the leaf economic spectrum present at local scales? *Ecography* **40**: 685–697.

Murren CJ. **2012**. The Integrated Phenotype. *Integrative and Comparative Biology* **52**: 64–76.

Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U, Poot P,
Purugganan MD, Richards CL, Valladares F, *et al.* 2010. Plant phenotypic plasticity in a
changing climate. *Trends in Plant Science* **15**: 684–692.

Niinemets Ü. **2001**. Global-Scale Climatic Controls of Leaf Dry Mass Per Area, Density, and
Thickness in Trees and Shrubs. *Ecology* **82**: 453–469.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,
Solymos P, Stevens MHH, Szoecs E, *et al.* 2025. *vegan: Community Ecology Package*.

Olesen JM, Bascompte J, Dupont YL, Jordano P. **2007**. The modularity of pollination
networks. *Proceedings of the National Academy of Sciences* **104**: 19891–19896.

Pau S, Slapikas R, Ho C-L, Bayliss SLJ, Donnelly RC, Abdullahi A, Helliker BR, Nippert
JB, Riley WJ, Still CJ, *et al.* 2025. Hyperspectral leaf reflectance of grasses varies with
evolutionary lineage more than with site. *Ecosphere* **16**: e70257.

Pieruschka R, Poorter H. **2012**. Phenotyping plants: genes, phenes and machines. *Functional*
plant biology: FPB **39**: 813–820.

Proulx SR, Promislow DEL, Phillips PC. **2005**. Network thinking in ecology and evolution.
Trends in Ecology & Evolution **20**: 345–353.

R Core Team. **2022**. *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing.

Rohart F, Gautier B, Singh A, Lê Cao K-A. **2017**. mixOmics: An R package for 'omics
feature selection and multiple data integration. *PLoS Computational Biology* **13**: e1005752.

Savolainen O, Lascoux M, Merilä J. **2013**. Ecological genomics of local adaptation. *Nature*
Reviews Genetics **14**: 807–820.

Schlosser G, Wagner GP (Eds). **2004**. *Modularity in Development and Evolution*. Chicago, IL:
University of Chicago Press.

Schweiger AK, Cavender-Bares J, Townsend PA, Hobbie SE, Madritch MD, Wang R,
Tilman D, Gamon JA. **2018**. Plant spectral diversity integrates functional and phylogenetic
components of biodiversity and predicts ecosystem function. *Nature Ecology & Evolution* **2**:
976–982.

Schwinning S, Lortie CJ, Esque TC, DeFalco LA. **2022**. What common-garden experiments
tell us about climate responses in plants. *Journal of Ecology* **110**: 986–996.

Serbin SP, Singh A, McNeil BE, Kingdon CC, Townsend PA. **2014**. Spectroscopic
determination of leaf morphological and biochemical traits for northern temperate and boreal tree
species. *Ecological Applications* **24**: 1651–1669.

Singh A, Serbin SP, McNeil BE, Kingdon CC, Townsend PA. **2015**. Imaging spectroscopy
algorithms for mapping canopy foliar chemical and morphological traits and their uncertainties.
Ecological Applications **25**: 2180–2197.

Song P, Wang J, Guo X, Yang W, Zhao C. **2021**. High-throughput phenotyping: Breaking
through the bottleneck in future crop breeding. *The Crop Journal* **9**: 633–645.

Stouffer DB, Bascompte J. **2011**. Compartmentalization increases food-web persistence.
Proceedings of the National Academy of Sciences **108**: 3648–3652.

Tay JK, Narasimhan B, Hastie T. **2023**. Elastic Net Regularization Paths for All Generalized
Linear Models. *Journal of Statistical Software* **106**: 1–31.

Thenkabail PS, Smith RB, De Pauw E. **2000**. Hyperspectral Vegetation Indices and Their
Relationships with Agricultural Crop Characteristics. *Remote Sensing of Environment* **71**: 158–
182.

Tigano A, Friesen VL. **2016**. Genomics of local adaptation with gene flow. *Molecular Ecology*

25: 2144–2164.

Ustin SL, Gamon JA. **2010**. Remote sensing of plant functional types. *New Phytologist* **186**:
795–816.

Wagner GP, Altenberg L. **1996**. Perspective: Complex Adaptations and the Evolution of
Evolvability. *Evolution* **50**: 967–976.

Wagner GP, Pavlicev M, Cheverud JM. **2007**. The road to modularity. *Nature Reviews*
Genetics **8**: 921–931.

Wang R, Gamon JA, Cavender-Bares J, Townsend PA, Zygielbaum AI. **2018**. The spatial
sensitivity of the spectral diversity–biodiversity relationship: an experimental test in a prairie
grassland. *Ecological Applications* **28**: 541–556.

Wong CY, Gilbert ME, Pierce MA, Parker TA, Palkovic A, Gepts P, Magney TS, Buckley
TN. **2023**. Hyperspectral Remote Sensing for Phenotyping the Physiological Drought Response
of Common and Tepary Bean. *Plant Phenomics* **5**: 0021.

Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J,
Chapin T, Cornelissen JHC, Diemer M, *et al.* **2004**. The worldwide leaf economics spectrum.
Nature **428**: 821–827.

Yearly variation of climate variables (1900-2020)

Fig. S2 Yearly variation of five climate variables used in the study. Each dot represents a year for each individual population.

A PLS-DA of hyperspectral data

B Map of California

Table mountain
Canyon Creek
Drum Powerhouse Rd.
UCD Garden
Ben Hur

C Population — BH — CC — DPR — TM2

D Spectral Heritability Across Wavelengths