Spectral network analysis illuminates coordinated plant traits across a climate gradient

Rishav Ray[1,2], Brandie Quarles-Chidyagwai[2,3], Jessica Lyons[1], Sarah Ashlock[3], Jennifer R. Gremer[3], Julin N. Maloof[2], Troy S. Magney[1,4]

1. Department of Plant Sciences, University of California Davis
2. Department of Plant Biology, University of California Davis
3. Department of Ecology and Evolution, University of California Davis
4. Department of Forest Management, University of Montana

Summary

- Understanding how plant populations respond to environmental variation through functional leaf traits remains challenging due to limitations of traditional phenotyping approaches. Hyperspectral reflectance offers a rapid, non-destructive and high-throughput method to capture functional trait variation and detect signatures of local adaptation across populations.

- We combined hyperspectral data, inverse modeling, and network analysis to investigate population-level variation in *Streptanthus tortuosus*. Using a common garden experiment with four geographically distinct populations, we applied partial least square discriminant analysis (PLS-DA) and ridge regression for population discrimination, inverse PROSPECT modeling to estimate leaf biochemical traits, and canonical correlation analysis to examine trait-climate relationships across historical (1900-1994) and recent (1995-2024) periods. We developed a spectral network approach treating wavelength correlations as biologically meaningful trait networks.

- Populations showed distinct, heritable spectral signatures with high classification accuracy. Significant population differences emerged in anthocyanins, carotenoids, chlorophyll, and water content. Trait-climate correlations shifted between time periods, consistent with historical climate adaptation. Network analysis revealed population-specific integration patterns, with more variable environments displaying greater spectral modularity.

30   ● Hyperspectral signatures provide a high-throughput tool for detecting population-level
31   adaptation and trait coordination. Our findings provide a framework to investigate how
32   plant populations respond to climate change through evolved shifts in trait networks
33   rather than isolated traits alone.

Introduction

Climate change is driving unprecedented environmental shifts that are reshaping selective

pressures on plant populations worldwide (Gienapp *et al.*, 2014; Edelsparre *et al.*, 2024), yet our

ability to detect and understand evolutionary responses to these changes remain limited by

traditional approaches of studying adaptation (Franks *et al.*, 2014; Anderson & Gezon, 2015;

Anderson *et al.*, 2025). While common garden experiments and reciprocal transplants have

provided foundational insights into local adaptation (Leimu & Fischer, 2008; Hereford, 2009;

Schwinning *et al.*, 2022), these methods are often constrained by logistical challenges, long

generational times, and the difficulty of measuring multiple traits simultaneously across large

populations (Merilä & Hendry, 2014; Song *et al.*, 2021). The growing availability of genomic

data has revolutionized our understanding of genetic basis of adaptation (Savolainen *et al.*, 2013;

Tigano & Friesen, 2016), but a critical gap remains between high resolution genomic data and

the complex, multi dimensional phenotypic data that are targets of natural selection (Houle *et al.*,

2010; Furbank & Tester, 2011). This phenotyping bottleneck is particularly acute in the study of

climate adaptation, where responses may involve coordinated changes across multiple

physiological and biochemical traits rather than simple modifications of individual traits (Nicotra

et al., 2010; Anderson *et al.*, 2012; Hein *et al.*, 2021). Consequently, there is an urgent need for

high throughput, integrative approaches that can capture the complexity of adaptive phenotypes

while scaling to the spatio-temporal dimensions over which climate driven evolution operates

(Des Roches *et al.*, 2018; Cavender-Bares *et al.*, 2022).

Hyperspectral reflectance analysis offers a promising solution to these challenges by providing a

rapid, non-destructive window into plant functional traits that integrates information across

multiple physiological and biochemical processes (Ustin & Gamon, 2010; Cavender-Bares *et al.*,

2016; Wong *et al.*, 2023; Wong, 2023; Magney, 2025). Unlike traditional vegetation indices that

rely on a few broad spectral bands, hyperspectral data captures reflectance across hundreds to

thousands of narrow wavelengths, encoding detailed information about leaf biochemistry, water

content, and structural properties (Kokaly *et al.*, 2009; Serbin *et al.*, 2014). This rich spectral

information can be used with models run in inverse mode such as PROSPECT, which enable the

quantitative estimation of specific biochemical traits, such as chlorophyll, carotenoids,

anthocyanins, leaf-mass per area (LMA), water content, and leaf structure from reflectance

spectra (Jacquemoud & Baret, 1990; Féret *et al.*, 2021). Recent advances have demonstrated the

power of hyperspectral approaches for detecting functional diversity within and among plant

communities (Asner *et al.*, 2015; Schweiger *et al.*, 2018; Wong *et al.*, 2023; Li *et al.*, 2023; Pau

et al., 2025), mapping plant stress responses (Gamon *et al.*, 2019), linking spectra to gene

expression (Chen *et al.*, 2025), and even predicting ecosystem function from spectral diversity

patterns (Wang *et al.*, 2018; Cavender-Bares *et al.*, 2022). However, most applications have

focused on species-level differences or broad ecological patterns, with limited exploration of

how hyperspectral signatures might reveal population-level adaptation and the evolutionary

processes that shape functional trait variation within species (Singh *et al.*, 2015; Meireles *et al.*,

2020). Furthermore, traditional analyses treat spectral bands as independent variables, potentially

missing emergent properties that can be gleaned from the coordinated relationship between them

(Gamon *et al.*, 2020).

Understanding how plants respond to environmental variation requires moving beyond

individual traits to consider the coordinated relationships among multiple functional

characteristics, a perspective rooted in the concept of phenotypic integration from quantitative

genetics and evolutionary developmental biology (Cheverud, 1996; Armbruster *et al.*, 2014).

Traits rarely evolve in isolation; instead, they are connected through genetic correlations (linkage

disequilibrium), developmental constraints, and functional dependencies that result in observable

phenotypes (Wagner & Altenberg, 1996; Klingenberg, 2008; He *et al.*, 2020; Li & He, 2024).

The structure of these trait correlations itself is an evolvable property that can be shaped by

natural selection, with different patterns of integration. This can range from highly modular to

tightly integrated, representing alternative adaptive strategies for responding to environmental

challenges (Hansen & Houle, 2008; Murren, 2012; Wang *et al.*, 2023). Network theory has

emerged as a powerful framework for understanding such complex systems across biology, from

gene regulatory networks that control development (Barabási & Oltvai, 2004; Davidson &

Erwin, 2006), to ecological networks that structure community interactions (Bascompte &

Jordano, 2007; Olesen *et al.*, 2007; Gladstone-Gallagher *et al.*, 2023). In each case, network

topology, that is the pattern of connections among nodes, provides insights into system function,

stability, and evolutionary potential that are undetectable when components are studied in

isolation (Albert & Barabási, 2002; Proulx *et al.*, 2005). However, applying network approaches

to study phenotypic integration has been limited by the challenge of measuring sufficient

numbers of traits to characterize network structure, particularly in studies of natural populations

where high-throughput phenotyping methods are essential (Murren, 2012; Messier *et al.*, 2017;

Li & He, 2024). Hyperspectral data offers an unprecedented opportunity to examine trait

networks at high resolution and test fundamental hypotheses about how environmental variation

shapes the evolution of phenotypic integration (Cavender-Bares *et al.*, 2016; Schweiger *et al.*,

2018).

The California mountain jewelflower, *Streptanthus tortuosus* (Brassicaceae), provides an ideal

system for testing these novel approaches. While most research on local adaptation in

Streptanthus has focused on other members of the clade, few studies have characterized

population-level variation in *S. tortuosus* across California's diverse climatic gradients (Preston,

1991, 1994; Gremer *et al.*, 2020b,a). This species' wide distribution across elevation and

precipitation gradients, combined with common garden experiments, makes it particularly

suitable for investigating how environmental variation drives complex trait architecture (Gremer

et al., 2020a; Bontrager *et al.*, 2025). In this study, we combine multivariate hyperspectral

reflectance analysis, inverse modeling, and network theory to investigate how population-level

variation in spectral traits associates with climate of origin in *S. tortuosus*. Using a common

garden experiment spanning four *S. tortuosus* populations, we test whether spectral reflectance

data can detect heritable variation in functional traits and reveal population-specific patterns of

trait coordination. We apply a novel two-step analytical framework – partial least square

discriminant analysis (PLS-DA) for population discrimination followed by ridge regression

analysis to identify key wavelength regions – to characterize population specific spectral

signatures. Through inverse PROSPECT modeling, we estimate six key leaf functional traits and

examine whether their relationships with historical and recent climate variables using canonical

correlation analysis across two temporal scales. We then developed a spectral network approach

that treats wavelength correlations as biologically meaningful trait networks, revealing distinct
patterns of spectral integration among populations. Finally, we test whether these network
architectures correlate with long-term climate variability patterns, providing insights into how
environmental variation shapes both individual traits and their coordination.

Materials and Methods

Plant material and common garden

We conducted a common garden experiment at the UC Davis Vegetable Crops fields in Davis,
CA USA, with seeds collected from four populations of *S. tortuosus* (Table 1). Seedlings from
these populations were planted at the garden in raised beds with 21 replicates for BH, CC, DPR
populations, and 98 replicates for TM2 population. Seeds were stratified for four weeks at 4°C in
the dark in a cold room at UC Davis. On October 30, 2023, after stratification, plants were
moved to a growth chamber with 12-hour day length and warm light conditions (light intensity
~100uMol/microeinstein) to stimulate germination. Plants were initially exposed to 22°C
day/8°C night temperatures. To harden plants to the cold temperatures of the field, temperatures
were adjusted accordingly: 20°C day/7°C night 7-days prior to field transplant, 19°C day/6°C
night 6-days prior to field transplant, and 18°C day/5°C night 5-days prior to field transplant.
Seedlings were transplanted into a randomized block design with 12 blocks spread across 6
raised beds and 30-cm between experimental plants. The raised beds were 12" tall, 110cm wide,
and 14m in length; there was 1m between each bed. Ron's mix, sand, and perlite were used to fill
each bed. The beds were watered with drip irrigation the day before transplant for 5 hours,
during the afternoon after planting for 5 hours, and five days after transplant for 5 hours.

Population Name	Population Code	Latitude	Longitude
Table mountain 2	TM2	39.59255	-121.55072

Ben Hur	BH	37.40985	-119.96458
Canyon Creek	CC	39.58597	-121.43311
Drum Powerhouse Road	DPR	39.22846	-120.81518

148 **Table 1: *S. tortuosus* population names and codes used in the study along with their**
149 **geographic coordinates.**

Hyperspectral data collection and processing

Leaf-level hyperspectral reflectance measurements were collected from plants grown in the
common garden using a full-range spectrometer (HR-1024i; Spectra Vista Corporation—SVC).
Spectral reflectance was measured from 400 to 2500 nm with a spectral resolution of 1.5 nm
(250–1000 nm), 3.8 nm (1000–1890 nm), and 2.5 nm (1890–2500 nm). Leaf spectra were
collected with a leaf clip (LC-RP PRO; SVC) attached to a fiber optic cable. The leaf clip houses
an internal tungsten halogen lamp that was standardized for reflectance using a built-in
Spectralon white reference panel (Labsphere Inc.). Leaf measurements were carried out on
flowering plants with young fully expanded leaves that were big enough to fit the leaf clip; three
measurements were taken per plant. This resulted in a total of 153 measurements (BH: 5x3, CC:
7x3, DPR: 7x3, TM2: 32x3). Reference measurements were made before every plant and dark
current corrections were automatically taken following each scan with an internal shutter.
Spectra were then averaged across the three measurements per plant and smoothed using a
Savitzky-Golay filter with a 5-point moving window.
164

Statistical analysis

Partial Least Squares Discriminant Analysis (PLS-DA) was performed (*mixOmics* package in R;
Rohart *et al.*, 2017) to assess population-level differentiation in hyperspectral space. PLS-DA
was chosen because it is specifically designed to handle high-dimensional datasets where the
number of variables (wavelengths) exceeds the number of samples, a common challenge in
hyperspectral analysis. PLS-DA reduces dimensionality while maximizing the separation
between groups, making it ideal for identifying population-specific spectral discrimination. It
was followed by a PERMANOVA test (*vegan* package; Oksanen *et al.*, 2025) to statistically test

for differences in spectral data among populations. We used PLS-DA to identify canonical

variates that maximally separated the four populations while explaining spectral variance. Broad-

sense heritability (H^2) was calculated for each wavelength using variance components from a

mixed-effects model using the *lme4* package with population as a random effect: $H^2 = \sigma^2_G/(\sigma^2_G$

$+ \sigma^2_E)$, where σ^2_G is genetic variance among populations and σ^2_E is environmental/residual

variance within populations.

Ridge regression models were fitted for each population(*glmnet* package; Tay *et al.*, 2023) to

identify wavelength-specific contributions to population classification while avoiding overfitting

that commonly occurs with high-dimensional spectral data. The L2 loss function in ridge

regression shrinks coefficients toward zero but retains all wavelengths, allowing us to quantify

the relative importance of different spectral regions for population classification. Spectral

coefficients were extracted across all wavelengths and integrated within six functionally relevant

spectral regions (carotenoids: 450-520 nm, chlorophyll: 650-700 nm, NIR plateau: 700-1300 nm,

red edge: 680-750 nm, SWIR (short wave infra-red) 1 : 1400-1500 nm, SWIR 2: 1900-2000 nm).

To avoid overfitting and provide unbiased performance estimates, we used leave-one-out cross-

validation (LOOCV). For each sample, the ridge regression model was trained on all remaining

samples, and predictions were made on the held-out sample. Area under the ROC curve (AUC)

was calculated from these out-of-fold predictions. Confidence intervals were estimated using

1000 bootstrap replicates of the LOOCV procedure (Fig. S1).

Inverse PROSPECT modeling was applied using the *prospect* R package (Féret & Boissieu,

2024) to estimate six leaf functional traits: anthocyanin content, carotenoid content, chlorophyll

content, leaf mass area, structural parameter (N), and equivalent water thickness (estimated water

mass per leaf area). Climate data for each population's origin site were obtained from Flint Basin

Characterization Model (Flint *et al.*, 2021; Fig. S2), spanning 1900-2024 and divided into

historical (1900-1994) and recent (1995-2024) periods. Canonical Correlation Analysis (CCA)

was performed (*mixOmics* package) to examine relationships between estimated leaf traits and

climate variables across different temporal periods. CCA examines multivariate relationships

between multiple leaf traits and multiple climate variables simultaneously, capturing complex

trait-environment associations that would be missed by univariate approaches. It identifies linear

combinations of traits and climate variables that are maximally correlated, revealing coordinated

trait response to environmental gradients.

To address spectral autocorrelation and reduce redundancy for the network analysis, we

implemented a filtering strategy based on spectral information content. We calculated the

Shannon diversity index ($H = -\Sigma(p_i \times \ln(p_i))$) for each spectrum to estimate the effective number

of independent wavelengths, where p_i represents the proportion of total reflectance at wavelength

i (Fig. S3). The mean Shannon index across all samples was 7.3, yielding an effective number of

wavelengths (eH) of approximately 1495. Combined with our heritability filtering ($H^2 > 0.5$),

this approach retained 987 wavelengths with both genetic differentiation and non-redundant

spectral information, acknowledging that adjacent wavelengths are inherently correlated due to

the physical properties of light-matter interactions. This filtering strategy reduced computational

burden while preserving biologically meaningful spectral variation for network construction. For

each population, correlation matrices were calculated between all wavelength pairs using

Pearson correlation coefficients. To account for sample size limitations, we employed a bootstrap

approach, randomly sampling 10 individuals with replacement 100 times and constructing

networks for each iteration. Edges were retained only for correlations exceeding $|r| > 0.7$. Four

key network parameters were calculated using the *igraph* package: (1) Density - the proportion

of possible edges that are actually present in the network, calculated as the number of observed

edges divided by the maximum possible number of edges (n(n-1)/2 for undirected networks).

Values range from 0 (no connections) to 1 (fully connected), with higher values indicating more

integrated spectral responses across wavelengths. (2) Centralization - a measure of how

unequally distributed the connections are across nodes, quantifying the extent to which the

network is dominated by highly connected "hub" wavelengths. Centralization is calculated based

on the variance in node degrees relative to a star network (maximum centralization) and ranges

from 0 (all nodes equally connected) to 1 (one central hub connected to all other nodes). High

centralization indicates that spectral integration is controlled by a few key wavelengths. (3)

Modularity - a measure of community structure that quantifies how well the network can be

divided into distinct groups of densely connected nodes (modules) with sparse connections

between groups. Modularity was calculated using the Louvain algorithm, which optimizes the

modularity function $Q = \Sigma[A_{ij} - (k_i k_j/2m)]\delta(c_i, c_j)$, where Aij is the adjacency matrix, k_i is the

degree of node i, m is the total number of edges, and $\delta(c_i,c_j) = 1$ if nodes i and j belong to the

same community. Values range from -0.5 to 1, with higher values indicating stronger modular

organization of spectral responses. (4) Transitivity - clustering coefficient - the probability that

two nodes connected to a common neighbor are also connected to each other, measuring the

tendency for wavelengths to form tightly connected clusters. Calculated as the ratio of observed

triangles to possible triangles in the network: $C = 3 \times$ (number of triangles) / (number of

connected triples). Values range from 0 to 1, with higher values indicating greater local

clustering of spectral correlations. CCA was performed between population-level network

metrics and long-term climate variability measures (coefficients of variation for temperature,

precipitation, and climatic water deficit from 1900-2024) to test relationships between network

topology and environmental variation.

All analyses were performed in R (version 4.2.1; R Core Team, 2022). Significance testing for

network metrics was conducted using ANOVA with Tukey's HSD post-hoc tests. Bootstrap

confidence intervals were calculated for all network parameters. Statistical significance was set

at $\alpha = 0.05$, with Bonferroni correction applied for multiple comparisons where appropriate.

Results

Population-level hyperspectral differentiation and genetic variation

Fig. 1: Hyperspectral variation and heritability in *Streptanthus tortuosus* populations across

California. A) Partial Least Squares Discriminant Analysis (PLS-DA) of hyperspectral data showing four

S. tortuosus populations (TM2, CC, DPR, and BH) along two principal variates. The x-variate explains

65.62% of the spectral variation while the y-variate explains 32.31%. **B)** Map of California showing the

geographic distribution of the sampled populations: Table Mountain (TM2), Canyon Creek (CC), Drum

Powerhouse Road (DPR), Ben Hur (BH) and the UC Davis Common Garden (UCD) where all these

populations were planted and measured. **C)** Reflectance spectra by population across wavelengths from

400-2500 nm. Different coloured lines represent individual plant measurements. The red line (labeled

"cv") shows the coefficient of variation across wavelengths (right y-axis). **D)** Broad-sense heritability

(H²) estimates across wavelengths from 400-2500 nm, showing variation in the genetic contribution to

spectral reflectance traits. The coloured bands indicate spectral regions that correlate with leaf chlorophyll content (green), and SWIR regions (blue).

To determine whether *S. tortuosus* populations maintain distinct spectral signatures under standardized common garden conditions, we performed PLS-DA on the hyperspectral data over a multivariate space. The PLS-DA ordination demonstrated significant differences in spectral values among populations (PERMANOVA F:17.43, p-value:0.001), with the first two canonical variates explaining 97.93% of the total spectral variation (Fig. 1A, S4). The populations formed distinct clusters in the ordination space, with TM2 and DPR populations showing greatest

separation along the first canonical variate, while the BH population separated in a distinct

cluster along the second canonical variate consistent with its geographical separation (Fig. 1B).

Despite being measured under the same common garden conditions, the samples showed spectral

variation across key regions of the wavelength range, denoted by the coefficient of variation

(CV) plotted in the second axis (Fig. 1C). The CV across samples peaked in several key spectral

regions, with the highest variability observed around 680 nm (chlorophyll absorption), and water

absorption bands around 1400 nm and 1900 nm; regions that correspond to functionally

important leaf properties. Broad-sense heritability (H^2), which estimates the amount of

phenotypic variation explained by genotype, revealed substantial genetic control over spectral

reflectance traits (Fig 1D). The H^2 values peak considerably across the chlorophyll and SWIR

regions, suggesting that the genetic variation is concentrated in spectral regions that reflect key

physiological and biochemical leaf properties. Taken together, maintenance of population-

specific spectral signatures under common garden conditions, combined with high heritability

estimates, provides strong evidence for genetic differentiation in leaf optical properties among *S.*

tortuosus populations.

Ridge regression analysis reveals population-specific spectral signatures

Fig. 2: Spectral characteristics and classification performance across *Streptanthus tortuosus*

populations. A) Spectral coefficients derived from ridge regression across wavelengths (400-2500 nm)

for four *S. tortuosus* populations (TM2, DPR, CC, and BH). Background shading highlights key spectral

regions associated with different leaf properties: carotenoids, chlorophyll, NIR plateau, red edge, SWIR 1,

and SWIR 2. **B)** Receiver Operating Characteristic (ROC) curves showing classification performance for

each population based on ridge regression models. All populations demonstrate excellent classification

accuracy with Area Under the Curve (AUC) (Fig. S1) values ranging from 0.838 (CC) to 0.99 (BH),

indicating high discriminative power of spectral data for population identification. **C)** Integrated

coefficients rank plot showing the relative importance of six key spectral regions in distinguishing

between populations. Each colored line represents a population (TM2, DPR, CC, and BH), with vertical positions indicating their rank (higher = stronger signal) for each spectral region.

Having established that there are heritable differences among populations, we next asked, what are the specific wavelengths that distinguish between the populations? To answer this, we applied ridge regression analysis and quantified the predictive power of hyperspectral data for population classification. Regression coefficients across the spectra (referred to as spectral coefficients hereafter) varied markedly across wavelengths and among populations, with each population exhibiting unique patterns of spectral importance (Fig. 2A). All populations showed pronounced spectral coefficients in the visible region (400-700nm), particularly in areas corresponding to carotenoid (450-520 nm), and chlorophyll (400-250 & 650-700nm) absorption bands. Notable population specific differences emerged in the water absorption regions (1400-1500 and 1900-2000 nm), where DPR displayed stronger spectral coefficients compared to other populations, which varied in those regions (Fig 2C). The near-infrared region (700-1300 nm) is influenced by leaf internal structure and showed distinct coefficient patterns, with BH exhibiting

particularly strong signals in this spectral region (Fig. 2A). We numerically integrated the

spectral coefficients across six functionally relevant wavelength regions and ranked them to

quantify the population specific spectral "fingerprints" (Fig. 2C). TM2 consistently ranked

highest in three key regions: chlorophyll, carotenoids, and SWIR 2 absorption bands. In contrast,

BH was most sensitive to the near infra-red (NIR) region. CC and DPR populations showed

more moderate rankings across most spectral regions, with CC ranking highest in the SWIR 1

region and DPR showing intermediate values across all spectral domains. Furthermore, the

discriminative power of hyperspectral data for population identification was high (AUC 0.838-

0.99) across all populations, indicating that hyperspectral reflectance data contain sufficient

population-specific information to achieve high classification accuracy (Fig 2B, S1). In

summary, these results suggest clear population-specific differentiation in spectral signatures,

indicating that populations vary in multiple optical properties.

Leaf trait differences and climate associations across temporal scales

Fig. 3: Leaf trait variation across populations and relationships with climate variables. A) Mean

values (±SE) and F-statistic and p-value of ANOVA test for six key leaf traits across four *S. tortuosus*

populations (BH, CC, DPR, and TM2). Traits include anthocyanins ($\mu g/cm^2$), carotenoids ($\mu g/cm^2$),

chlorophyll ($\mu g/cm^2$), leaf mass area (g/m^2), leaf structure constant (N), and water content (g/cm^2). The

Timeline at the bottom indicates the historical (1900-1994) and recent (1995-2024) climate periods used

in the analysis. **B-D)** Canonical Correlation Analysis (CCA) biplots showing relationships between leaf

traits (blue vectors) and climate variables (gray vectors) during recent (**B**) and historical (**C**) time periods,

and the difference between the two time periods (**D**). The first canonical variate explains the primary

correlation structure between leaf traits and climate conditions.

We next asked whether the variation in reflectance in different populations can be explained by

various home climatic variables that they have experienced. To that end, we applied the inverse

PROSPECT (Jacquemoud & Baret, 1990; Féret & Boissieu, 2024) model to estimate six key leaf

functional traits from the hyperspectral data, namely: anthocyanins, carotenoids, chlorophyll, leaf mass area, N (leaf structural complexity), and equivalent water thickness (EWT) (Fig 3A). The traits show significant population-level variation, e.g. anthocyanin contents were highest in BH and CC populations, exceeding levels in DPR and TM2 populations by about two-fold. Carotenoid content showed the opposite pattern, with TM2 displaying markedly higher concentrations compared to other populations, and not detectable from BH spectra. Chlorophyll content was highest in CC, while BH showed substantially lower levels. EWT also varied significantly among populations, with CC again showing the highest values compared to BH. Leaf mass area and the structural parameter N showed more modest but consistent population differences, indicating coordinated variation in both biochemical and structural leaf properties.

To test for environmental associations with these estimated traits, we applied canonical correlation analysis with populations' home climatic variables going back to 1900, split between recent (1995-2024), and historic (1900-1994) time periods. The analysis revealed distinct patterns of trait-climate associations across different temporal periods. For recent climate conditions, the first canonical correlation explained strong relationships between leaf traits and contemporary environmental variables (Fig. 3B). Chlorophyll content and EWT correlated positively with recent actual evapotranspiration and precipitation and negatively with climate water deficit. Historical climate relationships showed similar trait-environment associations compared to recent patterns (Fig. 3C). However, climate change analysis (difference between recent and historical periods) provided insights into how shifting environmental conditions correlate with current trait variation (Fig. 3D). Anthocyanins showed a strong positive correlation with change in maximum temperature, and interestingly, chlorophyll, and EWT both correlated positively with changes in precipitation and minimum temperature patterns. These temporal patterns reveal that trait-climate correlations differ between historical and recent periods, with the strength and direction of associations varying across time intervals.

Fig. 4: Spectral network analysis and climate-network relationships in *S. tortuosus* populations. A)

Network visualization of spectral relationships in four *S. tortuosus* populations (TM2, BH, DPR, and CC).

Each node represents a wavelength, and edges represent correlations between wavelengths. **B)**

Comparison of four network parameters (centralization, density, modularity, and transitivity) across

populations. Box plots show the distribution of bootstrapped values (n=100, 10 individuals with

replacement per iteration) with significance indicators for population differences (*p<0.05, **p<0.01,

***p<0.001, ****p<0.0001). **C)** CCA biplot showing relationships between network metrics (blue

vectors) and climate variability parameters (gray vectors) from 1900-2024.

We then asked whether the leaf reflectance patterns are co-ordinated amongst themselves and if

the relationship between them was population specific or not. To that end, we applied a network

approach to build a spectral graph, where each node represents a particular wavelength, and each

connecting edge is the correlation coefficient of their reflectance value between the two nodes.

This network analysis revealed distinct population-specific network topology with quantitatively

significant differences in spectral integration patterns (Fig. 4 A-B). TM2 and DPR populations

exhibited dense, highly interconnected networks with numerous edges connecting across

wavelengths, reflected in significantly higher network density and degree compared to BH and

CC (p < 0.0001). In contrast, BH and CC populations displayed more modular network

structures characterized by distinct clusters of highly connected wavelengths separated by

sparser connections, with modularity values significantly higher than TM2 and DPR, confirming

compartmentalized spectral organization. CC exhibited the highest centralization values,

suggesting that spectral integration in this population is dominated by a few highly connected

wavelength "hubs." Transitivity showed more modest but significant differences, with DPR

displaying the highest values, indicating greater local clustering of wavelength correlations.

These results demonstrate that populations have fundamentally different approaches to

organizing spectral trait correlations, ranging from highly integrated (TM2, DPR) to highly

modular (CC, BH) network architectures suggesting population-specific variation in the

coordination of leaf optical properties.

Canonical correlation analysis revealed that these population differences in spectral network

organization are systematically related to long-term climate variability patterns, rather than

conditions in specific historical or recent periods (Fig. 4C). The first canonical variate separated

density from the other network parameters, while the second canonical variate distinguished

centralization and modularity and transitivity. The AET/PET ratio correlated highly with density,

suggesting that populations from environments with higher water availability (higher AET/PET)

tend to have denser, more integrated spectral networks. Conversely, centralization and

modularity both loaded positively on the second canonical variate and were associated with

minimum and maximum temperature variability, indicating that populations from thermally

variable environments have more centralized and modular spectral integration patterns.

Furthermore, The coefficient of variation in climatic water deficit (CWD CV) loaded negatively

on the first canonical variate, suggesting that populations experiencing greater water stress

variability are associated with lower transitivity and higher density in their spectral networks.

Discussion

This study demonstrates the power of integrating advanced hyperspectral analysis with network

theory to reveal patterns of plant trait coordination. Our novel analytical framework

discriminated four *S. tortuosus* populations with good accuracy (AUC ranging from 0.838 -

0.99), while revealing that spectral traits represent heritable characteristics. The combination of

PLS-DA for population discrimination and ridge regression for identifying key wavelength

regions proved highly effective, with population specific signatures observed even under

common garden conditions. Beyond individual trait variation, our network analysis uncovered

distinct patterns of spectral coordination among populations, ranging from highly connected

(TM2) to modular (CC, BH) architectures that correlate with long-term climate variability.

Critically, these methodological advances revealed that populations have evolved not only

different leaf biochemical properties, with notable differences in anthocyanins, carotenoids, and

chlorophyll content, but also fundamentally different strategies for coordinating these traits

reflected through their spectral networks. Strong correlations between network topology and

historical climate patterns suggest that environmental variability has shaped both individual traits

and their patterns of integration, representing a new perspective on the evolution of complex trait

architecture in natural populations.

Building on these climate network relationships, specific spectral regions provide additional

insights into the functional basis of trait variation. The near-infrared region (700-1300 nm)

showed distinct coefficient patterns, with BH exhibiting particularly strong signals in this

structurally-sensitive spectral region. The NIR plateau is primarily influenced by leaf internal

structure rather than biochemical composition, with reflectance determined by the scattering of

light at air-cell wall interfaces within the leaf mesophyll (Jacquemoud & Baret, 1990; Féret et al.,

2021). Higher NIR reflectance typically indicates greater leaf thickness, increased intercellular

air spaces, or altered mesophyll organization—structural characteristics that can influence light

capture efficiency, gas exchange, and water use efficiency (Onoda et al., 2017). The strong NIR

signal in BH may reflect population-level variation in structural traits that could be related to

environmental conditions at this population's origin site (Niinemets, 2001; Wright *et al.*, 2004;

Karabourniotis *et al.*, 2021). However, it is important to emphasize that our study cannot

distinguish between spectral variation arising from adaptive response to local environment

versus genetic drift, founder effects, or population structure. While the functional significance of

the estimated traits correlating with the environmental variables suggests that they may be

ecologically relevant, we make no causal claims about the evolutionary mechanisms that drive

these patterns.

Our hyperspectral network analysis represents a shift from traditional approaches that treat

spectral bands as independent variables to a systems-level framework that captures the

coordinated relationship between wavelengths. While previous studies have relied primarily on

vegetation indices or individual wavelength analysis (Bannari *et al.*, 1995; Thenkabail *et al.*,

2000; Gamon *et al.*, 2019), our network approach reveals emergent properties of spectral trait

integration that are not captured by conventional methods. Our two-step analytical pipeline,

combining PLS-DA for population discrimination with ridge regression for mechanistic insight,

offers significant advantages over single method approaches by first establishing the existence of

population differences and then identifying the specific wavelength regions driving these

patterns. This framework addresses a key gap in evolutionary ecology, where high throughput

phenotyping methods are needed to match the resolution of genomic tools (Houle *et al.*, 2010;

Furbank & Tester, 2011; Pieruschka & Poorter, 2012; Cavender-Bares *et al.*, 2022).The

integration of canonical correlation analysis across multiple temporal scales (historic vs. recent

climate) further demonstrates how advanced statistical methods can detect trait correlation with

environmental variables. This study provides a template for studying evolutionary responses to

environmental change, which can be further bolstered by addition of genomic and transcriptomic

data ,to study adaptation in real-time (Hoffmann & Sgrò, 2011; Anderson *et al.*, 2012; Franks *et*

al., 2014; Merilä & Hendry, 2014).

Additionally, our spectral network analysis reveals a previously unrecognized level of biological

organization, i.e. the coordinated structure of the reflectance patterns, that parallels network

organization observed across diverse biological systems. While reflectance patterns themselves

are not direct targets of selection, they represent emergent properties of functionally important

traits such as pigment composition, water retention capacity, and leaf structural characteristics

that can be under direct selective pressure. Similar to how gene regulatory networks exhibit

modular architectures that facilitate evolvability while maintaining stability (Wagner *et al.*, 2007;

Espinosa-Soto & Wagner, 2010), our spectral networks show population-specific patterns of

modularity and integration that are consistent with different strategies for coordinating

underlying physiological and biochemical processes. The dense, highly connected networks in

TM2 and DPR populations mirror the "small-world" properties observed in metabolic networks,

where high clustering facilitates local optimization while maintaining global connectivity (Jeong

et al., 2000; Barabási & Oltvai, 2004). Conversely, the modular architectures in CC and BH

populations resemble the compartmentalized organization of ecological food webs, where

modularity enhances stability against perturbations while allowing independent evolution of

subsystems (Olesen *et al.*, 2007; Stouffer & Bascompte, 2011). This biological network

perspective suggests that natural selection acts not only on individual functional traits but also on

the patterns of coordination among these traits, with spectral networks providing a high-

dimensional window into this complex trait architecture, as proposed in quantitative genetics

theory (Cheverud, 1996; Armbruster *et al.*, 2014). The correlation between network topology and

climate variability patterns supports the hypothesis that environmental unpredictability favors

modular organization, thereby allowing rapid reconfiguration of trait combinations, while stable

environments favor integrated networks that optimize coordinated responses (Schlosser &

Wagner, 2004). Our findings thus extend network thinking beyond its traditional domains in

molecular biology and ecology to encompass the evolution of complex phenotypes in the

spectral sensing domain. This network perspective offers practical applications for conservation

biology, as species with highly integrated spectral networks (low modularity, high density) may

be more vulnerable to environmental disruption due to their tightly coupled trait systems.

While our methodological framework demonstrates significant promise, several limitations

warrant consideration. First, our analysis was restricted to a single growing season under

common garden conditions, which does not fully capture the dynamic of spectral trait expression

across environments or temporal variation. This is highlighted by the absence of detectable

carotenoid levels in the BH population using the inverse modeling. This highlights potential

limitations in our inverse PROSPECT modeling approach for certain biochemical compounds

under specific conditions (Féret & Boissieu, 2024). This could reflect either genuine biological

differences in carotenoid accumulation strategies or methodological constraints in detecting low-

concentration pigments in certain leaf types (Wong, 2023). Additionally, while our network

analysis reveals compelling patterns of spectral integration, the biological mechanisms

underlying these coordination patterns remain unclear. Future studies combining transcriptomic

and metabolomic approaches with hyperspectral analysis could provide mechanistic insights into

how plants achieve different integration strategies.

The successful implementation of this analysis framework requires careful consideration of

several methodological parameters and quality control practices that can significantly influence

results. Key analytical choices include spectral smoothing window size (we used Savitzky-Golay

filtering with a 5-point window), correlation threshold for edge retention ($|r| > 0.7$), and number

of bootstrap replicates ($n = 100$). Each of these parameters represent a trade-off between noise

reduction, information retention, and computational complexity that should be optimized for

specific dataset and research questions. The small number of populations and relatively small

sample size used in this study also limits our ability to generalize across the species' full

geographic range. Future studies with a larger sample size and more populations would

significantly strengthen our conclusions about climate trait relationships. Field measurement

quality control is another essential component and should include consistent leaf positioning,

regular white reference calibration, and systematic checking for spectral artifacts or outliers that

could propagate through network analyses (Burnett *et al.*, 2021). Importantly, network metrics

(density, modularity, centralization, transitivity) and biochemical trait estimates should be

interpreted comparatively between treatments or populations rather than as absolute biological

values, as their biological meaning emerges from relative patterns rather than specific numerical

thresholds. This analytical framework shows considerable promise for scaling to larger spatial

and temporal dimensions: canopy-level hyperspectral imaging could extend population
comparisons to landscape scales (Kaur *et al.*, 2024; Pierrat *et al.*, 2025)(, while integration with
emerging airborne (Cawse-Nicholson *et al.*, 2025) and satellite hyperspectral sensors (Cawse-
Nicholson *et al.*, 2025; Caplan & Huemmrich, 2025) could enable real-time monitoring of trait
network evolution across entire species ranges.

Conclusion

This study demonstrates how hyperspectral analysis can be integrated with network theory to
understand population-level trait coordination in plants. Our analytical framework reveals that
spectral data encode information about both individual functional traits and their patterns of
coordination, providing a high-throughput approach to studying complex phenotypes in natural
populations. Future integration of this spectral framework with genomic and transcriptomic
datasets could illuminate the molecular mechanisms underlying different coordination strategies,
potentially identifying the genetic basis of trait integration and informing predictive models of
population responses to environmental change. Such multi-scale applications will be essential for
predicting and monitoring how natural populations respond to accelerating environmental
change.

Acknowledgements

This research was funded by the National Science Foundation (DEB-2129589 to J.N.M, J.R.G,
T.S.M, and Denneal Jamison-McClung) and The National Institute of Food and Agriculture,
United States Department of Agriculture (USDA-NIFA) award CA-D-PLB-2795-H to JNM. We
acknowledge the contribution from Maya Arakaki, technician Kate Ruiz-Cox, and Carlos Perez
and all the undergraduate students involved in setting up the field and data collection. J. Schmitt
provided valuable guidance on study design, analyses, and manuscript drafts.

Competing interests

550 The authors declare no competing interests.

Author contribution

Conceptualization (research direction and analytical framework): RR, JNM, TSM, (common garden and data collection): BQ-C, JRG, JNM, TSM; Data curation: RR, JL; Formal Analysis: RR, JNM, TSM; Funding acquisition: JRG, JNM, TSM; Investigation: RR, JNM, TSM; Visualization: RR; Writing – original draft: RR; Writing – review & editing: RR, BQ-C, JL, SA, JRG, JNM, TSM

Data availability

Raw data and source code are available in the following Github repository.

https://github.com/rishavray/spectral-network

References

Albert R, Barabási A-L. **2002**. Statistical mechanics of complex networks. *Reviews of Modern Physics* **74**: 47–97.

Anderson JT, DeMarche ML, Denney DA, Breckheimer I, Santangelo J, Wadgymar SM. **2025**. Adaptation and gene flow are insufficient to rescue a montane plant under climate change. *Science*.

Anderson JT, Gezon ZJ. **2015**. Plasticity in functional traits in the context of climate change: a case study of the subalpine forb Boechera stricta (Brassicaceae). *Global Change Biology* **21**: 1689–1703.

Anderson JT, Inouye DW, McKinney AM, Colautti RI, Mitchell-Olds T. **2012**. Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. *Proceedings of the Royal Society B: Biological Sciences* **279**: 3843–3852.

Armbruster WS, Pélabon C, Bolstad GH, Hansen TF. **2014**. Integrated phenotypes: understanding trait covariation in plants and animals. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**: 20130245.

Asner GP, Martin RE, Anderson CB, Knapp DE. **2015**. Quantifying forest canopy traits:

Imaging spectroscopy versus field survey. *Remote Sensing of Environment* **158**: 15–27.

Bannari A, Morin D, Bonn F, Huete AR. **1995**. A review of vegetation indices. *Remote*
Sensing Reviews **13**: 95–120.

Barabási A-L, Oltvai ZN. **2004**. Network biology: understanding the cell's functional
organization. *Nature Reviews Genetics* **5**: 101–113.

Bascompte J, Jordano P. **2007**. Plant-Animal Mutualistic Networks: The Architecture of
Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**: 567–593.

Bontrager M, Worthy SJ, Cacho NI, Leventhal L, Maloof JN, Gremer JR, Schmitt J,
Strauss SY. **2025**. Herbarium specimens reveal a constrained seasonal climate niche despite
diverged annual climates across a wildflower clade. *Proceedings of the National Academy of*
Sciences **122**: e2503670122.

Burnett AC, Anderson J, Davidson KJ, Ely KS, Lamour J, Li Q, Morrison BD, Yang D,
Rogers A, Serbin SP. **2021**. A best-practice guide to predicting plant traits from leaf-level
hyperspectral data using partial least squares regression. *Journal of Experimental Botany* **72**:
6175–6189.

Caplan S, Huemmrich KF. **2025**. Unveiling PACE OCI's hyperspectral terrestrial data
products. *Remote Sensing Letters* **16**: 422–433.

Cavender-Bares J, Meireles JE, Couture JJ, Kaproth MA, Kingdon CC, Singh A, Serbin
SP, Center A, Zuniga E, Pilz G, *et al.* **2016**. Associations of Leaf Spectra with Genetic and
Phylogenetic Variation in Oaks: Prospects for Remote Detection of Biodiversity. *Remote Sensing*
8: 221.

Cavender-Bares J, Schneider FD, Santos MJ, Armstrong A, Carnaval A, Dahlin KM,
Fatoyinbo L, Hurtt GC, Schimel D, Townsend PA, *et al.* **2022**. Integrating remote sensing
with ecology and evolution to advance biodiversity conservation. *Nature Ecology & Evolution* **6**:
506–519.

Cawse-Nicholson K, Chadwick KD, Brodrick PG, Kiper M, Thompson DR, Schimel D,

Miller CE, Townsend PA, Alves LF, Shiklomanov AN, *et al.* 2025. Intrinsic dimensionality as
a metric for temporal plant diversity evaluation: Case study from the SHIFT campaign.
Ecosphere **16**: e70213.

Chen Y, Monks L, Rubio VE, Cox AJ, Swenson NG. 2025. Linking leaf hyperspectral
reflectance to gene expression. *Communications Earth & Environment* **6**: 694.

Cheverud JM. 1996. Developmental Integration and the Evolution of Pleiotropy. *American
Zoologist* **36**: 44–50.

Davidson EH, Erwin DH. 2006. Gene Regulatory Networks and the Evolution of Animal Body
Plans. *Science* **311**: 796–800.

**Des Roches S, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, Schweitzer JA,
Palkovacs EP. 2018**. The ecological importance of intraspecific variation. *Nature Ecology &
Evolution* **2**: 57–64.

Edelsparre AH, Fitzpatrick MJ, Saastamoinen M, Teplitsky C. 2024. Evolutionary
adaptation to climate change. *Evolution Letters* **8**: 1–7.

Espinosa-Soto C, Wagner A. 2010. Specialization Can Drive the Evolution of Modularity.
PLOS Computational Biology **6**: e1000719.

Féret J-B, Berger K, de Boissieu F, Malenovský Z. 2021. PROSPECT-PRO for estimating
content of nitrogen-containing leaf proteins and other carbon-based constituents. *Remote Sensing
of Environment* **252**: 112173.

Féret J-B, Boissieu F de. 2024. prospect: an R package to link leaf optical properties with their
chemical and structural properties with the leaf model PROSPECT. *Journal of Open Source
Software* **9**: 6027.

Flint LE, Flint AL, Stern MA. 2021. *The basin characterization model—A regional water
balance software package*. U.S. Geological Survey.

Franks SJ, Weber JJ, Aitken SN. 2014. Evolutionary and plastic responses to climate change

in terrestrial plant populations. *Evolutionary Applications* **7**: 123–139.

Furbank RT, Tester M. **2011**. Phenomics – technologies to relieve the phenotyping bottleneck.
Trends in Plant Science **16**: 635–644.

Gamon JA, Somers B, Malenovský Z, Middleton EM, Rascher U, Schaepman ME. **2019**.
Assessing Vegetation Function with Imaging Spectroscopy. *Surveys in Geophysics* **40**: 489–513.

Gamon JA, Wang R, Gholizadeh H, Zutta B, Townsend PA, Cavender-Bares J. **2020**.
Consideration of Scale in Remote Sensing of Biodiversity. In: Cavender-Bares J, Gamon JA,
Townsend PA, eds. Remote Sensing of Plant Biodiversity. Cham: Springer International
Publishing, 425–447.

Gienapp P, Reed TE, Visser ME. **2014**. Why climate change will invariably alter selection
pressures on phenology. *Proceedings of the Royal Society B: Biological Sciences* **281**: 20141611.

Gladstone-Gallagher RV, Hewitt JE, Siwicka E, Gammal JM, Brustolin MC, Norkko A,
Pilditch CA, Thrush SF. **2023**. Ecological network analysis of traits reveals variable response
capacity to stress. *Proceedings of the Royal Society B: Biological Sciences* **290**: 20230403.

Gremer JR, Chiono A, Suglia E, Bontrager M, Okafor L, Schmitt J. **2020a**. Variation in the
seasonal germination niche across an elevational gradient: the role of germination cueing in
current and future climates. *American Journal of Botany* **107**: 350–363.

Gremer JR, Wilcox CJ, Chiono A, Suglia E, Schmitt J. **2020b**. Germination timing and
chilling exposure create contingency in life history and influence fitness in the native wildflower
Streptanthus tortuosus. *Journal of Ecology* **108**: 239–255.

Hansen TF, Houle D. **2008**. Measuring and comparing evolvability and constraint in
multivariate characters. *Journal of Evolutionary Biology* **21**: 1201–1219.

He N, Li Y, Liu C, Xu L, Li M, Zhang J, He J, Tang Z, Han X, Ye Q, *et al.* 2020. Plant Trait
Networks: Improved Resolution of the Dimensionality of Adaptation. *Trends in Ecology &*
Evolution **35**: 908–918.

Hein NT, Ciampitti IA, Jagadish SVK. **2021**. Bottlenecks and opportunities in field-based

high-throughput phenotyping for heat and drought stress. *Journal of Experimental Botany* **72**:

5102–5116.

Hereford J. **2009**. A Quantitative Survey of Local Adaptation and Fitness Trade-Offs. *The*

American Naturalist **173**: 579–588.

Hoffmann AA, Sgrò CM. **2011**. Climate change and evolutionary adaptation. *Nature* **470**: 479–

485.

Houle D, Govindaraju DR, Omholt S. **2010**. Phenomics: the next challenge. *Nature Reviews*

Genetics **11**: 855–866.

Jacquemoud S, Baret F. **1990**. PROSPECT: A model of leaf optical properties spectra. *Remote*

Sensing of Environment **34**: 75–91.

Jeong H, Tombor B, Albert R, Oltvai ZN, Barabási A-L. **2000**. The large-scale organization

of metabolic networks. *Nature* **407**: 651–654.

Karabourniotis G, Liakopoulos G, Bresta P, Nikolopoulos D. **2021**. The Optical Properties of

Leaf Structural Elements and Their Contribution to Photosynthetic Performance and

Photoprotection. *Plants* **10**: 1455.

Kaur S, Kakani VG, Carver B, Jarquin D, Singh A. **2024**. Hyperspectral imaging combined

with machine learning for high-throughput phenotyping in winter wheat. *The Plant Phenome*

Journal **7**: e20111.

Klingenberg CP. **2008**. Morphological Integration and Developmental Modularity. *Annual*

Review of Ecology, Evolution, and Systematics **39**: 115–132.

Kokaly RF, Asner GP, Ollinger SV, Martin ME, Wessman CA. **2009**. Characterizing canopy

biochemistry from imaging spectroscopy and its application to ecosystem studies. *Remote*

Sensing of Environment **113**: S78–S91.

Leimu R, Fischer M. **2008**. A Meta-Analysis of Local Adaptation in Plants. *PLOS ONE* **3**:

e4010.

Li C, Czyż EA, Halitschke R, Baldwin IT, Schaepman ME, Schuman MC. **2023**. Evaluating
potential of leaf reflectance spectra to monitor plant genetic variation. *Plant Methods* **19**: 108.

Li Y, He N. **2024**. Innovations and prospectives of multidimensional trait integration. *New*
Phytologist **244**: 337–340.

Magney TS. **2025**. Hyperspectral reflectance integrates key traits for predicting leaf metabolism.
New Phytologist **246**: 383–385.

Meireles JE, Cavender-Bares J, Townsend PA, Ustin S, Gamon JA, Schweiger AK,
Schaepman ME, Asner GP, Martin RE, Singh A, *et al.* 2020. Leaf reflectance spectra capture
the evolutionary history of seed plants. *New Phytologist* **228**: 485–493.

Merilä J, Hendry AP. **2014**. Climate change, adaptation, and phenotypic plasticity: the problem
and the evidence. *Evolutionary Applications* **7**: 1–14.

Messier J, McGill BJ, Enquist BJ, Lechowicz MJ. **2017**. Trait variation and integration across
scales: is the leaf economic spectrum present at local scales? *Ecography* **40**: 685–697.

Murren CJ. **2012**. The Integrated Phenotype. *Integrative and Comparative Biology* **52**: 64–76.

Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U, Poot P,
Purugganan MD, Richards CL, Valladares F, *et al.* 2010. Plant phenotypic plasticity in a
changing climate. *Trends in Plant Science* **15**: 684–692.

Niinemets Ü. **2001**. Global-Scale Climatic Controls of Leaf Dry Mass Per Area, Density, and
Thickness in Trees and Shrubs. *Ecology* **82**: 453–469.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,
Solymos P, Stevens MHH, Szoecs E, *et al.* 2025. *vegan: Community Ecology Package*.

Olesen JM, Bascompte J, Dupont YL, Jordano P. **2007**. The modularity of pollination
networks. *Proceedings of the National Academy of Sciences* **104**: 19891–19896.

Pau S, Slapikas R, Ho C-L, Bayliss SLJ, Donnelly RC, Abdullahi A, Helliker BR, Nippert
JB, Riley WJ, Still CJ, *et al.* 2025. Hyperspectral leaf reflectance of grasses varies with
evolutionary lineage more than with site. *Ecosphere* **16**: e70257.

Pierrat ZA, Magney TS, Richardson WP, Runkle BRK, Diehl JL, Yang X, Woodgate W,
Smith WK, Johnston MR, Ginting YRS, *et al.* 2025. Proximal remote sensing: an essential
tool for bridging the gap between high-resolution ecosystem monitoring and global ecology. *New*
Phytologist **246**: 419–436.

Pieruschka R, Poorter H. 2012. Phenotyping plants: genes, phenes and machines. *Functional*
plant biology: FPB **39**: 813–820.

Preston RE. 1991. The Intrafloral Phenology of Streptanthus Tortuosus (brassicaceae).
American Journal of Botany **78**: 1044–1053.

Preston RE. 1994. Pollination Biology of Streptanthus Tortuosus (brassicaceae). *Madroño* **41**:
138–147.

Proulx SR, Promislow DEL, Phillips PC. 2005. Network thinking in ecology and evolution.
Trends in Ecology & Evolution **20**: 345–353.

R Core Team. 2022. *R: A Language and Environment for Statistical Computing*. Vienna,
Austria: R Foundation for Statistical Computing.

Rohart F, Gautier B, Singh A, Lê Cao K-A. 2017. mixOmics: An R package for 'omics
feature selection and multiple data integration. *PLoS Computational Biology* **13**: e1005752.

Savolainen O, Lascoux M, Merilä J. 2013. Ecological genomics of local adaptation. *Nature*
Reviews Genetics **14**: 807–820.

Schlosser G, Wagner GP (Eds). 2004. *Modularity in Development and Evolution*. Chicago, IL:
University of Chicago Press.

Schweiger AK, Cavender-Bares J, Townsend PA, Hobbie SE, Madritch MD, Wang R,
Tilman D, Gamon JA. 2018. Plant spectral diversity integrates functional and phylogenetic

components of biodiversity and predicts ecosystem function. *Nature Ecology & Evolution* **2**:
976–982.

Schwinning S, Lortie CJ, Esque TC, DeFalco LA. **2022**. What common-garden experiments
tell us about climate responses in plants. *Journal of Ecology* **110**: 986–996.

Serbin SP, Singh A, McNeil BE, Kingdon CC, Townsend PA. **2014**. Spectroscopic
determination of leaf morphological and biochemical traits for northern temperate and boreal tree
species. *Ecological Applications* **24**: 1651–1669.

Singh A, Serbin SP, McNeil BE, Kingdon CC, Townsend PA. **2015**. Imaging spectroscopy
algorithms for mapping canopy foliar chemical and morphological traits and their uncertainties.
Ecological Applications **25**: 2180–2197.

Song P, Wang J, Guo X, Yang W, Zhao C. **2021**. High-throughput phenotyping: Breaking
through the bottleneck in future crop breeding. *The Crop Journal* **9**: 633–645.

Stouffer DB, Bascompte J. **2011**. Compartmentalization increases food-web persistence.
Proceedings of the National Academy of Sciences **108**: 3648–3652.

Tay JK, Narasimhan B, Hastie T. **2023**. Elastic Net Regularization Paths for All Generalized
Linear Models. *Journal of Statistical Software* **106**: 1–31.

Thenkabail PS, Smith RB, De Pauw E. **2000**. Hyperspectral Vegetation Indices and Their
Relationships with Agricultural Crop Characteristics. *Remote Sensing of Environment* **71**: 158–
182.

Tigano A, Friesen VL. **2016**. Genomics of local adaptation with gene flow. *Molecular Ecology*
25: 2144–2164.

Ustin SL, Gamon JA. **2010**. Remote sensing of plant functional types. *New Phytologist* **186**:
795–816.

Wagner GP, Altenberg L. **1996**. Perspective: Complex Adaptations and the Evolution of
Evolvability. *Evolution* **50**: 967–976.

Wagner GP, Pavlicev M, Cheverud JM. **2007**. The road to modularity. *Nature Reviews*

Genetics **8**: 921–931.

Wang R, Gamon JA, Cavender-Bares J, Townsend PA, Zygielbaum AI. **2018**. The spatial

sensitivity of the spectral diversity–biodiversity relationship: an experimental test in a prairie

grassland. *Ecological Applications* **28**: 541–556.

Wang X, Ji M, Zhang Y, Zhang L, Akram MA, Dong L, Hu W, Xiong J, Sun Y, Li H, *et al.*

2023. Plant trait networks reveal adaptation strategies in the drylands of China. *BMC Plant*

Biology **23**: 266.

Wong CYS. **2023**. Plant optics: underlying mechanisms in remotely sensed signals for

phenotyping applications. *AoB PLANTS* **15**: plad039.

Wong CY, Gilbert ME, Pierce MA, Parker TA, Palkovic A, Gepts P, Magney TS, Buckley

TN. **2023**. Hyperspectral Remote Sensing for Phenotyping the Physiological Drought Response

of Common and Tepary Bean. *Plant Phenomics* **5**: 0021.

Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J,

Chapin T, Cornelissen JHC, Diemer M, *et al.* 2004. The worldwide leaf economics spectrum.

Nature **428**: 821–827.

Supplementary Figures

AUC Values with 95% Confidence Intervals

768   **Fig. S1: Area under the curve values for each population with 95% confidence intervals**
769   **calculated with 1000 bootstrap iterations.** Bootstrap confidence intervals for AUC values were
770   narrow across all populations, indicating stable and reliable classification performance despite
771   modest sample sizes.
772
773

Yearly variation of climate variables (1900-2020)

 Fig. S2: Yearly variation of the five climate variables used in the study. Each dot represents

 a year for each individual population.

776   **Fig. S3: Shannon diversity index calculated on the reflectance data for all the samples used**

777   **in the study. Each dot represents an individual sample for each population.**

Fig. S4: **Importance value of each wavelength across the 4 PLS-DA variates from Fig. 1A.**

A PLS-DA of hyperspectral data

B Map of California

Table mountain
Canyon Creek
Drum Powerhouse Rd.
UCD Garden
Ben Hur

C

D Spectral Heritability Across Wavelengths

Population — BH — CC — DPR — TM2

A Spectral Coefficients Across Populations

B ROC Curves for Population Classification

AUC values:
TM2 = 0.985
BH = 0.991
DPR = 0.938
CC = 0.838

C Integrated Coefficients Rank