Duck slurry as a matrix for avian influenza virus detection and genetic characterization in
domestic flocks

Clément Castille[a*], Laura Lebouteiller[a*], Mathilda Walch[a], Guillaume Croville[a], Jean-Luc Guérin[a]
and Sébastien Mathieu Soubies[a#]

[a] IHAP, Université de Toulouse, INRAE, ENVT, Toulouse, France

Running Head: Detection and sequencing of avian influenza in duck slurry

#Address correspondence to sebastien.soubies@envt.fr.

*Clément Castille and Laura Lebouteiller contributed equally to this work. Author order was
determined by tossing a coin.

Keywords: avian influenza, environmental detection, duck, slurry, wastewater

Abstract

Avian influenza viruses (AIVs) circulate widely in domestic and wild birds and continue to pose
major risks to animal and public health, in particular those of high pathogenicity (HP). Ducks
play a key role in AIV ecology due to frequent subclinical infections and high viral shedding.
Efficient environmental surveillance approaches are needed to complement costly individual
testing, especially in the context of large-scale anti-H5 vaccination as implemented in France.
We evaluated duck slurry, the wastewater produced in duck fattening units, as a matrix for AIV
detection and optimized molecular protocols for sensitive molecular detection and genetic
characterization. Slurry samples (n = 172) were collected from 37 duck farms in southwestern
France between May 2024 and March 2025. Among several extraction strategies, RNA
extraction from the solid fraction using TRIzol LS followed by magnetic bead purification yielded
the highest sensitivity. Using this optimized protocol, 82% of samples tested positive for AIV
RNA, with 30% of positives being H5-positive; no H7 viruses were detected. Whole-segment
RT-PCR and sequencing were successful for shorter genomic segments, enabling subtype
determination, although virus isolation consistently failed. These findings demonstrate that slurry
is a promising matrix for AIV detection, providing valuable molecular data. Slurry-based
surveillance may therefore serve as an effective complement to individual testing and improve
early warning capacities for influenza surveillance in a One Health perspective.

Importance

In France, current surveillance of high pathogenicity avian influenza in ducks relies mainly on
individual swabs, which may miss subclinical or transient infections, particularly with low-
pathogenicity strains. By targeting slurry, a wastewater generated in duck fattening units, we
developed an affordable and practical approach for environmental surveillance. Slurry samples
captured high levels of avian influenza RNA and allowed partial genomic characterization
despite failed virus isolation. This strategy is directly relevant to monitoring avian influenza
viruses dynamics under current anti-H5 vaccination and can be extended to other livestock
systems. Duck slurry-based surveillance thus represents an efficient One Health tool to
strengthen preparedness against avian influenza outbreaks.

Introduction

Avian influenza viruses (AIVs) are enveloped, negative-sense RNA viruses belonging to the *Orthomyxoviridae* family. Their genome consists of eight segments and evolves through both mutation and reassortment. AIVs are subtyped depending on the combination of hemagglutinin (HA) and neuraminidase (NA), the two major surface glycoproteins of the viral envelope. Nineteen HA subtypes and 11 NA subtypes have been described so far, including the recently described H19 subtype (1, 2). All of these subtypes except H17, H18, N10 and N11 circulate among aquatic birds.

AIVs are additionally classified into pathotypes, namely low (LP) and high pathogenicity (HP) depending on the sequence of HA (3). LPAIVs typically replicate in the digestive tract of *Anatidae* without clinical expression (4). In contrast, HPAIVs display systemic replication in ducks with high respiratory, digestive (5) and feather (6) shedding along with variable clinical symptoms depending on viral genotype (7), species (8), age (9, 10) and vaccinal status of birds. Pre-symptomatic shedding in ducks may explain their role as amplifiers of viral outbreaks in high-density poultry areas (11).

Since their emergence, H5Nx HPAIVs of the Goose/Guangdong/96 lineage, particularly those belonging to clade 2.3.4.4b, have caused hundreds of millions of deaths in both wild and domestic birds, infected various mammalian species (including cats and cattle) and led to human cases (12, 13). Extensive reassortment with LPAIVs present in wild birds has resulted in a wide range of genotypes with diverse segment constellations (14).

France is the leading European producer of farmed ducks, that are in particular used for fatty liver production. Ducklings intended for this purpose typically spend their first 80 days of life in farms with outdoor access starting around 40 days of life. Afterwards, they are transferred into indoor fattening units, with a capacity of few hundred individuals each, for a final period of about 15 days. These units, typically equipped with slatted floors, generate a liquid waste - referred to as "slurry" - from faeces and feed residues washed away during cleaning**.** Slurry is stored on-site and can then be used either as fertilizer or for biogas production.

The high number of duck farms likely explains the numerous HPAIVs outbreaks in poultry farms that France experienced during the recent years. As a result, a nationwide anti-H5 vaccination program for farmed ducks was initiated in autumn 2023. This vaccination scheme is accompanied by thorough but costly virological surveillance to avoid silent circulation of HPAIVs in vaccinated flocks.

Alongside individual testing approaches, environmental virus detection approaches have recently gained momentum, particularly inspired by the success of wastewater-based surveillance of SARS-CoV-2. (15).

82   The aim of this study was to investigate the interest of duck slurry, a wastewater
83 produced by ducks in fattening units, as an environmental matrix for detecting AIVs infecting
84 ducks to get a better understanding of AIVs circulation in those animals.

Material and methods

<u>Samples and collection strategy:</u>

Slurry samples originated from thirty-seven duck fattening units situated within 50 km of
a methanation plant (Méthalandes, France) in southwestern France. Slurry was serially sampled
between May 2024 and March 2025. Slurry from each fattening unit was temporarily stored on-
site in slurry pits before being pumped into tanker trucks and transported to the methanation
plant. Each truck collected and transported slurry from a single farm at a time. Samples used for
analyses were collected before unloading each tanker truck into the plant system, using a
manual valve at the back of the tank. Slurry samples (500 mL), collected with a minimal
frequency of one collection per month, were kept at 4°C, for a maximal duration of 15 days until
transport and processing at the laboratory.

<u>Optimization of sample treatment and RNA extraction</u>

Aliquots of 1 mL of raw slurry samples were centrifuged at 11,000 g for 1 hour, 4 °C.
Supernatant were transferred into new tubes and 5 sample groups were processed (Figure 1A):
1) 125 µL of supernatant underwent RNA extraction using MAGFAST-768 RNA extraction kits
(IDVet, Grabels, France) following the manufacturer's instructions (Figure 1A, "Sup Beads"), 2)
250 µL of supernatant was diluted into 750 µL of TRIzol LS (ThermoFisher) and was treated
according to the manufacturer's instructions until phase separation, when the aqueous phase
was either subjected to magnetic bead-based RNA extraction as described above (120 µL
aqueous phase) (Figure 1A, "Sup Trizol Beads") or 3) subjected to RNA precipitation as
described below(Figure 1A, "Sup Trizol PP"). Pelleted solids were resuspended into 750 µL of
TRIzol LS and incubated for 5 minutes at room temperature. Two hundred microlitres of
chloroform were added, tubes were briefly vortexed and incubated for 3 minutes at room
temperature, followed by centrifugation for 15 minutes at 12,000 g, 4 °C. The aqueous phase
was transferred into new tubes and then treated by either 4) magnetic bead-based RNA
extraction (Figure 1A, "Pellet Trizol Beads") as described above (120 µL aqueous phase) or 5)
subjected to RNA precipitation Figure 1A, "Pellet Trizol PP") as described below.

<u>RNA precipitation</u>

Five hundred microlitres of isopropanol were added to the aqueous phase. Tubes were
incubated for 10 minutes at room temperature and centrifuged at 12,000 g for 10 minutes, 4 °C.
Supernatants were transferred and pellets were resuspended into 1 mL of 75% ethanol,

followed by brief vortexing and centrifugation for 5 minutes at 7500 g, 4 °C. Supernatant were

removed and pellets were dried at room temperature until complete evaporation of any residual

ethanol. Pellets were then resuspended into 50 µL of ultrapure water.

Real-time RT-PCR (rRT-PCR)

Real-time RT-PCR (rRT-PCR) was carried out on all extracted RNA using a commercial

kit simultaneously detecting the matrix gene as well as an avian reference transcript (reference

IDFLUATRI-100 Innovative Diagnostics SAS, Grabels, France). Positive samples were further

analysed by rRT-PCR using the IDFLUH5H7- kit (Innovative Diagnostics SAS, Grabels, France)

to detect H5 or H7 AIV. All rRT-PCR were performed on a LightCycler96 thermocycler (Roche

SAS, Boulogne-Billancourt, France) following the respective kit cycling instructions.

Full segments RT-PCR and sequencing

Each of the viral segments was amplified by 2-step reverse transcription-PCR (RT-PCR).

For reverse transcription, 10 µL of RNA was treated with the RevertAid™ First Strand cDNA

Synthesis Kit (ThermoScientific, reference 10387979) using the Uni12 primer (16) according to

the manufacturer's protocol. The obtained cDNAs were used for full-segment amplification using

Phusion hot start flex 2x master mix kit (New England Biolabs, reference M0536S) and a set of

gene specific primers (16). Eight distinct PCR mixes, corresponding to the 8 genes, were

prepared and contained 12.5 µL 2X buffer,7.5µL RNase free water, 2.5 µL of 10 µM primers and

2.5 µL cDNA. The PCR program consisted of an initial 30 s step at 98°C followed by 35 cycles

with the following conditions: 98°C for 10 s, 61°C for 30 s, and 72°C for 2 min 20 s. The

program ended with 1 step at 72°C for 10 min. PCR products were then excised after agarose

gel electrophoresis and sent for "ONT Lite Clonal Amplicon Sequencing" to Eurofins genomics

(Ebersberg, Germany).

Alignment to reference sequences

FASTQ files obtained after Nanopore sequencing were aligned to a set of reference AIV

genomes using Geneious Prime 2024.0.7.

Virus isolation

For virus isolation, 1 mL aliquots of slurry were centrifuged for 1 h at 11 000 g, 4 °C.

Three hundred microlitres were collected and mixed with 300 µL of Penicillin/streptomycin

(10.000 µg/mL, ThermoFisher), 35 µL of 10% Bovine Serum Albumin (m/V in PBS) and 70 µL of

amphotericine B (250 µg/mL, ThermoFisher). Two hundred microlitres of this solution were then

inoculated into the chorioallantoic sac (CAS) of 9 to 11 day-old embryonated chicken eggs (2 to

3 eggs per sample). Inoculated eggs were candled daily. After 72 h or until embryo mortality

was observed, allantoic fluids were collected and tested by rRT-PCR (M detection) as described
above.

Results

Based on previous data from the laboratory as well as the literature, detection of avian
influenza viruses (AIV) RNA by real-time RT-PCR (rRT-PCR) from the supernatant of
centrifuged slurry seemed achievable. Initial attempts were therefore performed on slurry
samples without viral spiking. This first method, hereafter referred to as the "reference protocol",
involved magnetic bead-based purification of the slurry supernatant after treatment with lysis
buffer.

Initial positive results for M gene detection (Figure 1B, column 1 "sup beads", median
Cq: 30,12) demonstrated the feasibility of using naturally contaminated slurry as a matrix for
optimizing detection protocols in the absence of viral spiking. Those same samples were then
used to compare several extraction strategies (Figure 1A). TRIzol LS - based RNA extraction of
slurry supernatant followed by isopropanol precipitation resulted in Cq values comparable to
those obtained with the reference protocol (Figure 1B, column 3 "sup trizol pp", median Cq:
30.59). In contrast, RNA extracted from slurry solid fraction (pellet) resuspended in TRIzol LS
and subjected to isopropanol precipitation yielded significantly lower Cq values (Figure 1B,
column 5 "pellet trizol pp", median Cq: 25.95). To circumvent the lengthy steps associated with
isopropanol precipitation, a variation of the TRIzol LS protocol described above was tested, both
on slurry supernatant and pellet. In this modified protocol, the aqueous phase obtained after
chloroform extraction was subjected to magnetic bead-based RNA purification: the resulting Cq
values were comparable to those obtained after isopropanol RNA precipitation (Figure 1B,
column 2 and 4 "sup trizol beads " and "pellet trizol beads", median Cq of 31.1 and 25.99,
respectively). The "pellet trizol beads" protocol, combining high analytical sensitivity with ease of
implementation, was considered as the "optimized protocol" and used in subsequent
experiments.

Real-time RT-PCR analysis of the endogenous RNA detected using the commercial kit
showed a trend consistent with that observed for AIV RNA, albeit with a milder decrease in Cq
between the reference protocol and the optimized protocol, (Figure 1C, column 1 versus 4).

A total of 172 slurry samples collected from 37 duck farms between May 2024 and
March 2025 were subsequently analysed using the optimized protocol. Among the 37 farms
from which slurry was analysed, 23 farms provided at least 3 samples during the study period.
Overall, 142 samples (83%) tested positive for AIV RNA (Figure 2A). Among those positive
samples, 43 tested positive for H5, corresponding to 30% of AIV positive samples and 25% of
all analysed samples. No sample tested positive for H7 AIV. A significant lower median Ct was
observed for M+H5+ samples (median value : 28.22) when compared with M+H5- samples
(median value: 31.55, p=2.1e-5 using Wilcoxon test, Figure 2A).

Temporal variations in Ct values were observed (Figure 2B), with mean values around

25-30 observed in June-July 2024 followed by an increase above 30, between August and

December 2024, consistent with lower genomic viral loads. Mean Ct values decreased again in

January - February 2025. Farm-to-farm variations were observed in the number of collected

samples (Figure 2C). Thirty-four out of the 37 studied farms provided at least one positive slurry

sample, and 16 farms provided at least one H5-positive sample. Among farms with at least five

analyzed samples (n = 15), distinct patterns of AIV detection were observed. Six farms (F24,

F26, F27, F30, F31and F34) were positive for AIV in all collected samples. Among those, four

farms (F26, F27, F30 and F34) yielded both M+,H5- and M+,H5+ samples (Figure 2C).

Longitudinal analysis of individual farms with at least 5 samples further revealed that for

8 out 15 farms, at least 2 series of positive samples separated by at least one negative sample

were observed (Figure 2D).

Whole-genome RT-PCR amplification was attempted on a subset of samples with M

gene Cq values below an empirical cut-off of 25, using RNA extracted with the optimized

method. In several cases, RT-PCR yielded visible amplicons on gel electrophoresis for

segments 4 to 8 while amplification failed for segments 1 to 3 (Figure 3A). Shorter segments

generally yielded higher amplification efficiency, although amplicons for segments 4 and 6 were

occasionally undetectable. Visible PCR products were excised, gel-purified and subjected to

Nanopore sequencing. Consensus sequences were obtained after alignment with a set of

reference influenza genomes, in particular for segments 4 and 6, which encode HA and NA,

respectively, for subtype determination. In the representative result shown (Figure 3B),

sequencing depths of several hundred reads were obtained for segments 4 to 8, enabling

robust consensus sequence generation.

Virus isolation was attempted on samples whose Cq values for M gene were lower than

an empirical cut-off of 32. Out of 39 sampled tested during the study period, none yielded AIV

isolates.

Discussion

In this work, we optimized the detection of avian influenza viruses (AIVs) in duck slurry to

enable viral surveillance and genetic characterization.

Molecular detection using the slurry supernatant had previously been performed by our

laboratory (unpublished results) and others (17). Based on other reports on influenza detection

in soil (18) and in wastewater (19), we found that RNA extraction from the solid fraction of slurry

using TRIzol LS resulted in enhanced detection sensitivity by rRT-PCR. Combining TRIzol LS

extraction with magnetic bead-based RNA purification further improved sample workflow. The

reasons for preferential detection of influenza RNA in the solid fraction remains unclear,

although influenza viral particles have been reported to adsorb on particulate matters (20).

Interestingly, the endogenous duck RNA detected using the rRT-PCR kit was also more
abundant in the solid fraction.

Longitudinal analysis of slurry collected in the methanation plant revealed a surprisingly
high frequency of AIV-positive samples, including approximately one-third that were H5-positive.
Although each truck collected slurry from a single farm at a time before unloading into the plant
system, the absence of tank cleaning between successive collections may have led to spillover
contamination, thereby artificially inflating the observed positivity rate. For instance, assuming a
hypothetical residual volume of positive slurry of 2 % of the tank capacity, the next sample
collected might yield a false positive result with a Cq value approximately 5 cycles higher.
Careful analysis of truck collection sequences may provide more accurate insights into the true
positivity rate of the samples, which is beyond the scope of this study.

The high detection rate of H5 sequences in the collected samples raises the question of
the pathotype of the detected viruses. Although formal classification would require sequencing
of the HA cleavage site, which is beyond the scope of this work, several lines of evidence
suggest that the detected viruses were most likely of low pathogenicity: i) only four H5 HPAIV
outbreaks were reported in isolated duck farms in southwestern France during the study period,
ii) the thorough official surveillance scheme enforced following the implementation of anti-H5
vaccination in ducks did not report H5 HPAIV positive detection in the area surrounding the
methanation plant during the period of our study, iii) partial HA sequencing from one H5-positive
sample revealed a monobasic cleavage site consistent with a LPAIV (data not shown). The
absence of H5-positive reports in the region—despite numerous positive detections in our
samples—may seem paradoxical. A likely explanation is that individual surveillance in
vaccinated duck flocks relies primarily on oropharyngeal or tracheal swabs. These samples are
excellent indicators of respiratory shedding of H5 HPAIVs, particularly those of clade 2.3.4.4b,
which are the main target of current surveillance programs in France. However, in the case of
LPAIV infection in ducks, respiratory shedding is only transient (21), and such infections may be
missed using oropharyngeal swabs alone. Considering that H5 LPAIV circulation in domestic
ducks may be a prelude to the emergence of HPAIVs, a scenario already experienced in France
(22), slurry-based detection may serve as a low-cost, effective complement to individual
surveillance.

Using slurry as a matrix to detect AIVs suffers from several limitations. First, it is
applicable only to farm systems that produce liquid waste, such as ducks in fattening units,
thereby excluding earlier production stages when birds are not housed under such systems.
Second, as mentioned above, cross-contamination between successive samples or over time
on a given farm may occur. Third, in our hands, virus isolation and amplification, which would
render complete genome sequencing possible, has consistently failed. The failure to isolate
virus - in contrast with the findings of Schmitz et al. - may be explained by several factors: i) the

likely LPAIV nature of the detected viruses, which may results in lower levels of viral shedding in
the faeces compared with HPAIVs, ii) the acidic pH of some slurry samples (pH <6, data not
shown), which may rapidly inactivate virions, iii) warm ambient temperatures during parts of the
study period (spring to autumn 2024), which may have accelerated viral degradation, and iv)
delays of up to 20 days between sample collection and virus isolation attempts. Despite these
limitations, slurry-based AIV detection appears to be a promising surveillance tool. It can
provide valuable genetic information and potentially track the emergence of HPAIVs from
LPAIVs progenitors. Future work will aim to generate more detailed temporal data on AIV
dynamics at the farm level, along with deeper genetic characterization of the detected viruses.
More broadly, such approaches could be extended to other livestock systems and pathogens,
contributing to integrated One Health surveillance strategies.

Acknowledgement

The authors wish to thank Mr. Xavier Labat, owner of the methanation plant, and Méthalandes
technical staff who helped for slurry sample collection.

Funding

This work was supported by funding from from the French government, managed by the
National Research Agency (ANR), under the France 2030 program OBEPINE+ (reference ANR-
24-MIEM-0004) , "Chaire de professor junior " held by Sebastien Mathieu Soubies, as well as by
the French "Région Nouvelle-Aquitaine" (project EMERG) and the SISP&EaU project (reference
ANRS-23-PEPR-MIE-0003), funded under the PEPR-MIE program and managed by ANRS-MIE
(French Agency for Research on AIDS and Viral Hepatitis – Emerging Infectious Diseases).

Data availablity

The raw data supporting the findings of this study are available from the corresponding author
upon reasonable request.

Author contribution

Conceptualization: SMS, JLG. Methodology: CC, LL, MW, GC, SMS. Investigation (sample
collection, laboratory work, sequencing): CC, LL, MW. Data curation & Formal analysis: CC, LL,
SMS, MW, GC. Writing – original draft: CC, LL, SMS. Writing – review & editing: CC, LL, SMS,
GC, JLG, MW. Supervision: SMS. Funding acquisition: SMS, JLG.

Conflict of interest

The authors declare no conflict of interest.

Ethics statement

Slurry samples were collected at the methanation plant with the consent of the owner. No live animals were sampled for this study. Virus isolation was performed by inoculation of embryonated chicken eggs, a standard method that does not require institutional ethics approval under European and French regulations for laboratory diagnostics.

References

1. Fereidouni S, Starick E, Karamendin K, Genova CD, Scott SD, Khan Y, Harder T, Kydyrmanov A. 2023. Genetic characterization of a new candidate hemagglutinin subtype of influenza A viruses. Emerg Microbes Infect 2225645.

2. Karakus U, Mena I, Kottur J, El Zahed SS, Seoane R, Yildiz S, Chen L, Plancarte M, Lindsay L, Halpin R, Stockwell TB, Wentworth DE, Boons G-J, Krammer F, Stertz S, Boyce W, de Vries RP, Aggarwal AK, García-Sastre A. 2024. H19 influenza A virus exhibits species-specific MHC class II receptor usage. Cell Host & Microbe 32:1089-1102.e10.

3. Lee D-H, Torchetti MK, Killian ML, Brown I, Swayne DE. 2024. Genome sequences of haemagglutinin cleavage site predict the pathogenicity phenotype of avian influenza virus: statistically validated data for facilitating rapid declarations and reducing reliance on in vivo testing. Avian Pathol 53:242–246.

4. Kida H, Yanagawa R, Matsuoka Y. 1980. Duck influenza lacking evidence of disease signs and immune response. Infect Immun 30:547–553.

5. Beerens N, Germeraad EA, Venema S, Verheij E, Pritz-Verschuren SBE, Gonzales JL. 2021. Comparative pathogenicity and environmental transmission of recent highly pathogenic avian influenza H5 viruses. Emerg Microbes Infect 10:97–108.

6. Gaide N, Filaire F, Bertran K, Crispo M, Dirat M, Secula A, Foret-Lucas C, Payré B, Perlas
A, Cantero G, Majó N, Soubies S, Guérin J-L. 2023. The Feather Epithelium Contributes to
the Dissemination and Ecology of clade 2.3.4.4b H5 High Pathogenicity Avian Influenza
Virus in Ducks. bioRxiv https://doi.org/10.1101/2023.07.26.550633.

7. Sturm-Ramirez KM, Hulse-Post DJ, Govorkova EA, Humberd J, Seiler P, Puthavathana P,
Buranathai C, Nguyen TD, Chaisingh A, Long HT, Naipospos TSP, Chen H, Ellis TM,
Guan Y, Peiris JSM, Webster RG. 2005. Are ducks contributing to the endemicity of highly
pathogenic H5N1 influenza virus in Asia? J Virol 79:11269–11279.

8. Pantin-Jackwood M, Swayne DE, Smith D, Shepherd E. 2013. Effect of species, breed and
route of virus inoculation on the pathogenicity of H5N1 highly pathogenic influenza (HPAI)
viruses in domestic ducks. Veterinary Research 44:62.

9. Pantin-Jackwood MJ, Suarez DL, Spackman E, Swayne DE. 2007. Age at infection affects
the pathogenicity of Asian highly pathogenic avian influenza H5N1 viruses in ducks. Virus
Research 130:151–161.

10. Cagle C, Wasilenko J, Adams SC, Cardona CJ, To TL, Nguyen T, Spackman E, Suarez
DL, Smith D, Shepherd E, Roth J, Pantin-Jackwood MJ. 2012. Differences in
pathogenicity, response to vaccination, and innate immune responses in different types of
ducks infected with a virulent H5N1 highly pathogenic avian influenza virus from Vietnam.
Avian Dis 56:479–487.

11. Vergne T, Gubbins S, Guinat C, Bauzile B, Delpont M, Chakraborty D, Gruson H, Roche
B, Andraud M, Paul M, Guérin J-L. 2021. Inferring within-flock transmission dynamics of
highly pathogenic avian influenza H5N8 virus in France, 2020. Transbound Emerg Dis
68:3151–3155.

12. Abdelwhab EM, Beer M. 2024. Panzootic HPAIV H5 and risks to novel mammalian hosts.
npj Viruses 2:1–3.

13. Bellido-Martín B, Rijnink WF, Iervolino M, Kuiken T, Richard M, Fouchier RAM. 2025.

Evolution, spread and impact of highly pathogenic H5 avian influenza A viruses. Nat Rev

Microbiol 1–16.

14. Fusaro A, Zecchin B, Giussani E, Palumbo E, Agüero-García M, Bachofen C, Bálint Á,

Banihashem F, Banyard AC, Beerens N, Bourg M, Briand F-X, Bröjer C, Brown IH,

Brugger B, Byrne AMP, Cana A, Christodoulou V, Dirbakova Z, Fagulha T, Fouchier RAM,

Garza-Cuartero L, Georgiades G, Gjerset B, Grasland B, Groza O, Harder T, Henriques

AM, Hjulsager CK, Ivanova E, Janeliunas Z, Krivko L, Lemon K, Liang Y, Lika A, Malik P,

McMenamy MJ, Nagy A, Nurmoja I, Onita I, Pohlmann A, Revilla-Fernández S, Sánchez-

Sánchez A, Savic V, Slavec B, Smietanka K, Snoeck CJ, Steensels M, Svansson V,

Swieton E, Tammiranta N, Tinak M, Van Borm S, Zohari S, Adlhoch C, Baldinelli F,

Terregino C, Monne I. 2024. High pathogenic avian influenza A(H5) viruses of clade

2.3.4.4b in Europe—Why trends of virus evolution are more difficult to predict. Virus

Evolution 10:veae027.

15. Wurtzer S, Marechal V, Mouchel JM, Maday Y, Teyssou R, Richard E, Almayrac JL,

Moulin L. 2020. Evaluation of lockdown effect on SARS-CoV-2 dynamics through viral

genome quantification in waste water, Greater Paris, France, 5 March to 23 April 2020.

Eurosurveillance 25:2000776.

16. Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. 2001. Universal primer set for the

full-length amplification of all influenza A viruses. Arch Virol 146:2275–2289.

17. Schmitz A, Pertusa M, Le Bouquin S, Rousset N, Ogor K, LeBras M-O, Martenot C, Daniel

P, Belen Cepeda Hontecillas A, Scoizec A, Morin H, Massin P, Grasland B, Niqueux E,

Eterradossi N. 2020. Natural and Experimental Persistence of Highly Pathogenic H5

Influenza Viruses in Slurry of Domestic Ducks, with or without Lime Treatment. Appl

Environ Microbiol 86:e02288-20.

18. Deboosere N, Horm SV, Delobel A, Gachet J, Buchy P, Vialette M. 2012. Viral elution and
concentration method for detection of influenza A viruses in mud by real-time RT-PCR. J
Virol Methods 179:148–153.

19. Mercier E, D'Aoust PM, Thakali O, Hegazy N, Jia J-J, Zhang Z, Eid W, Plaza-Diaz J, Kabir
MP, Fang W, Cowan A, Stephenson SE, Pisharody L, MacKenzie AE, Graber TE, Wan S,
Delatolla R. 2022. Municipal and neighbourhood level wastewater surveillance and
subtyping of an influenza virus outbreak. 1. Sci Rep 12:15777.

20. Dong Z, Ma J, Qiu J, Ren Q, Shan Q, Duan X, Li G, Zuo YY, Qi Y, Liu Y, Liu G, Lynch I,
Fang M, Liu S. 2023. Airborne fine particles drive H1N1 viruses deep into the lower
respiratory tract and distant organs. Sci Adv 9:eadf2165.

21. França M, Stallknecht DE, Poulson R, Brown J, Howerth EW. 2012. The pathogenesis of
low pathogenic avian influenza in mallards. Avian Dis 56:976–980.

22. Briand F-X, Schmitz A, Ogor K, Le Prioux A, Guillou-Cloarec C, Guillemoto C, Allée C, Le
Bras M-O, Hirchaud E, Quenault H, Touzain F, Cherbonnel-Pansart M, Lemaitre E,
Courtillon C, Gares H, Daniel P, Fediaevsky A, Massin P, Blanchard Y, Eterradossi N, van
der Werf S, Jestin V, Niqueux E. 2017. Emerging highly pathogenic H5 avian influenza
viruses in France during winter 2015/16: phylogenetic analyses and markers for zoonotic
potential. Euro Surveill 22:30473.

393
394

 Figure Legends

Figure 1. Comparison of influenza RNA (M gene, A) and bird endogenous RNA (B)
detection by rRT-PCR using the indicated protocols. Sup: supernatant; PP: precipitation.
Boxplots present median, interquartile range and extreme values. Points represent individual
slurry samples and lines link values from one slurry sample.

Figure 2. rRT-PCR detection of avian influenza virus (AIV) in slurry samples from 37
farms between May 2024 and March 2025. A: Distribution of M gene Ct values in analyzed
samples (n=172); boxplots show distributions for M positive, H5-negative samples (left) and M
positive, H5-positive samples (right) ; the p-value corresponds to a two-tailed Wilcoxon test. B:
Temporal variations of mean M gene Ct values (line). The shaded area represents the standard
deviation and points represent individual samples. C: Number of samples provided by individual
farms, showing total samples (grey), M-positive samples (blue), and H5-positive samples (red).
D: Temporal patterns of AIV detection in individual farms; the colour scale indicates M gene Ct
values and H5 positive samples are indicated. For all panels, negative samples were empirically
assigned a Ct value of 41.

Figure 3. Genome RT-PCR amplification (A) and sequencing (B) of avian influenza
viruses from slurry samples. A: representative agarose gel electrophoresis of sets of PCR
amplicons from four slurry samples (a) to d)). 1-8: viral segment number; M: molecular ladder.
B: example of sequencing coverage plots presenting read number per segment position from
segments   4   to   8.   Reads   were   mapped   to   reference
A/Mallard/Sweden/SVA250129SZ0398/JM026682/OT/2024.