HCFC1 and YY1 mediate recruitment of COMPASS and Integrator to initiate X chromosome inactivation.

Jeffrey Boeren[1*], Beatrice F. Tan[1*], Sarra Merzouk[1,2], Eveline Rentmeester[1], Rien van Haperen[1], Samuel J. Luchsinger-Morcelle[1], Cheryl Maduro[1], Richelle Rietdijk[1], Büşra Göynük[1], Wilfred F.J. van IJcken[3], Jeroen A.A. Demmers[4], Cristina Gontan[1,†], Hegias Mira Bontenbal[1,5,†,✉], Joost Gribnau[1,†,✉]

Affiliations

[1]Department of Developmental Biology, Erasmus MC, 3015CN, Rotterdam, The Netherlands,

[2]Current address: Broad Institute of MIT and Harvard, Cambridge, MA, United States,

[3]Center for Biomics, Erasmus MC, 3015CN, Rotterdam, The Netherlands,

[4]Center for Proteomics, Erasmus MC, PO Box 2040, 3000 CA, Rotterdam, The Netherlands,

[5]Current address: IGMM, University of Montpellier, CNRS, Montpellier, France.

*These authors contributed equally.

†These authors contributed equally.

✉Correspondence: hegias.mira-bontenbal@igmm.cnrs.fr, j.gribnau@erasmusmc.nl

Abstract

The evolution of mammalian sex chromosomes has driven the emergence of mechanisms that balance X-linked gene dosage between male (XY) and female (XX) cells. In females, dosage compensation is achieved through X chromosome inactivation (XCI), initiated by upregulation of the long non-coding RNA *Xist*, which spreads in *cis*, recruiting chromatin modifiers to silence gene expression on one X chromosome. Here, we conducted a forward genetic screen and identified X-encoded Host Cell Factor 1 (HCFC1), a member of the COMPASS H3K4 methyltransferase complex, as a dose-dependent XCI-activator. HCFC1 loss results in genome-wide reduction of H3K4me3 at specific regulatory elements and downregulation of nearby genes, including *Xist*. We show that HCFC1 and YY1 are co-recruited genome-wide to gene regulatory elements. Mass spectrometry analysis confirmed an interaction of HCFC1 and YY1 and uncovered the Integrator complex as another prominent YY1 partner. YY1 depletion results in genome-wide loss of Integrator recruitment at gene regulatory elements and reduced expression of nearby genes, including *Xist cis*-regulatory genes *Jpx* and *Ftx*. These results highlight a co-regulatory role for HCFC1 in COMPASS recruitment and *Xist* activation, alongside YY1-mediated recruitment of Integrator to *Xist* regulatory elements and genes to activate female-exclusive XCI.

Introduction

X chromosome inactivation (XCI) provides a paradigm for studying gene regulation in a chromosomal context. In mice and other mammals, random XCI is initiated in all female somatic cells shortly after implantation and can be modeled in female (XX) mouse embryonic stem cells (ESCs), which trigger XCI upon *in vitro* differentiation. This process is orchestrated by *Xist*, a long non-coding RNA (lncRNA) encoded within the X inactivation center (XIC), a genomic locus enriched for lncRNA genes that regulate *Xist* expression. Upon initiation of XCI, *Xist* coats the X chromosome in *cis* and recruits chromatin-modifying complexes to silence X-linked genes. Prior to and during early XCI, *Xist* transcription is repressed on the future active X by antisense transcription of the lncRNA *Tsix* (reviewed in [1,2]).

Female-exclusive XCI is regulated by X-encoded activators of the XCI process directing dose-dependent activation of *Xist*[3,4]. The X-encoded E3 ubiquitin-ligase RNF12 was identified as an XCI-activator through dose-dependent degradation of pluripotency factor REX1[5,6]. YY1 and its

homologue REX1 compete for binding at the same gene regulatory element of *Xist*, where REX1 recruitment prevents YY1-mediated activation of *Xist*[7]. Homozygous *Rnf12* knockout (KO) ESCs fail to initiate XCI during differentiation, whereas heterozygous loss of *Rnf12* leads to delayed XCI initiation with skewing towards inactivation of the X chromosome carrying the mutated *Rnf12* allele in *vitro*[8]. While the maternally inherited *Rnf12* allele is essential for imprinted XCI in extra-embryonic tissues in mice[9,10], a conditional *Rnf12* KO mouse model showed that *Rnf12* is dispensable for the initiation of random XCI in the post-implantation epiblast[11]. Additionally, compound homozygous KO ESCs and mice of *Rnf12* and its primary target *Zfp42* (encoding REX1) display a partial rescue of the XCI phenotype, restoring it to near wild-type levels[10].

These findings highlight the important role RNF12 plays in XCI regulation. Nonetheless, the fact that XCI is still initiated in *Rnf12*[+/−] and *Rnf12*[−/−]:*Zfp42*[−/−] differentiating female ESCs, but not in male ESCs, points to the presence of additional XCI activators. The lncRNAs *Jpx* and *Ftx*, respectively located 10 kb and 90 kb upstream of *Xist*, have been implicated as XCI-activators, playing important roles in *Xist* regulation (reviewed in[1,2]) However, deletion studies indicate that most of this activity is mediated in *cis*[8,12,13]. Also, the X-linked demethylase KDM5C has been described as an XCI-activator by catalyzing the conversion of H3K4me2/3 to H3K4me1 at the YY1-REX1 regulatory element of *Xist*[14]. Although *Kdm5c* KO female embryos are non-viable, *Xist* expression is unaffected in *Kdm5c* heterozygous KO females supporting the hypothesis that multiple XCI activators are at play[14]. Finally, the GATA transcription factor family has been implicated in XCI regulation; however, the only X-linked family member, *Gata1,* is not expressed in ESCs or the epiblast, indicating that other XCI-activators remain to be identified[15].

Here, we used a forward genetic screen deleting large segments of the X chromosome to identify novel regulators of XCI and uncover the X-linked Host Cell Factor 1 (HCFC1) as a potent XCI activator. HCFC1, a component of the COMPASS H3K4 methyltransferase complex, promotes XCI by recruiting COMPASS to deposit H3K4me3 at the *Xist* regulatory element. Acute HCFC1 depletion during mouse ESC differentiation abolishes *Xist* and *Jpx* expression, preventing XCI. We found that genome-wide recruitment of HCFC1 is partly mediated by the transcription factor YY1, which we confirmed to physically interact with HCFC1. YY1 interactome analysis further identified the Integrator complex, a key mediator of transcriptional pause release, as a major YY1-associated partner. Loss of YY1 reduces Integrator binding at thousands of regulatory sites,

including *Xist cis*-regulatory elements, without affecting H3K4me3 deposition. Together, these

findings reveal a dual role for HCFC1 and YY1 in XCI initiation: HCFC1-mediated recruitment

of COMPASS to establish H3K4me3 and YY1-mediated recruitment of Integrator to promote

transcriptional activation of *Xist*.

Results

A forward genetic screen to identify novel XCI-activators

To identify new XCI-activators that act in parallel with *Rnf12* in the activation of XCI, we

performed a CRISPR/Cas9-based screen deleting large regions of the X chromosome (Fig. 1a).

We made use of female heterozygous $Rnf12^{+/-}$ F1 hybrid *Mus musculus castaneus* (Cast): *Mus*

musculus musculus 129/Sv (129) ESCs providing a powerful SNP-based readout to differentiate

the two X chromosomes. These cells also display complete skewed XCI of the *Rnf12* mutant X_{129}

chromosome to retain one active copy of *Rnf12* X_{cast} throughout SC differentiation[8]. Due to this

phenotype, large heterozygous deletions of the X_{129} chromosome are tolerated during

differentiation except for deletions that affect XCI (Fig. 1a). The deletions generated covered all

regions of the early evolutionary strata, which are expected to retain key regulatory elements and

genes involved in XCI (Fig. 1b)[16–18]. Proper deletion of regions from the X_{129} chromosome was

confirmed by PCR of allele-specific repeat amplifications, whole-genome sequencing (WGS)

detecting SNPs between the X_{129} and X_{Cast} chromosomes, and DNA-FISH (Extended Data Fig.

1a-e and Extended Data Fig. 2a). Differentiation followed by *Xist* quantitative PCR (qPCR)

analyses of at least two biological replicates per deletion indicated significant loss of *Xist*

expression at day 8 of differentiation in deletion E (ΔE), located 29.5Mb centromeric to *Xist* (Fig.

1b,c). ΔE's inability to upregulate *Xist* was subsequently confirmed by RNA fluorescent in situ

hybridization (RNA-FISH) (Extended Data Fig. 2b-d). Deletion of regions harboring putative

activators *Gata1* (located in ΔA and ΔAB) and *Kdm5c* (located in ΔK) did not result in a significant

loss of *Xist* expression (Fig. 1c)[14,15]. To identify the gene(s) involved in XCI activation, we

generated incremental and convergent deletions within ΔE (Fig. 1d) and determined the effect on

Xist expression at day 4 and day 8 of ESC differentiation. ΔE2 and ΔE3 showed a reduction of *Xist*

expression, unlike ΔE1, implying that our target gene(s) must be contained within ΔE2 and ΔE3

(Extended Data Fig. 2e). Next, we narrowed down the search by creating additional deletions

within regions E2 and E3, confirmed by WGS and RNA-seq (Fig. 1e and Extended Data Fig. 1e)

and discovered that ΔE2B and ΔE3B displayed a *Xist* phenotype at day 4 and day 8 of

differentiation, but not ΔE2A and ΔE3A (Fig. 1f). RNA-seq analysis revealed that loss of an 80kb

region encompassing *Hcfc1* and *Irak1* severely affects XCI at both analysed timepoints, whereas

downregulation of pluripotency factors and upregulation of factors associated with ESC

differentiation were broadly uncompromised compared to the parental cell line (Fig. 1e and

Extended Data Fig. 2f). This analysis confirmed downregulation of *Xist* in ESC lines harboring

ΔE2B and ΔE3B encompassing *Hcfc1* and *Irak1* (Fig. 1g,h) and as expected, impaired inactivation

of X-linked genes (Extended Data Fig. 2g). Analysis of previously published datasets show that,

similar to *Rnf12*, *Hcfc1* is upregulated between days 1 and 3 of ESC differentiation when XCI is

initiated (Fig. 1i and Extended Data Fig. 2g) [5,19], whereas *Irak1* is not upregulated. HCFC1 has

also been shown to interact with the crucial *Xist*-regulator YY1, pointing at *Hcfc1* as a candidate

XCI activator [7,20].

Hcfc1 is a dose-dependent activator of XCI

To test whether *Hcfc1* possesses XCI-activator activity, we restored *Hcfc1* expression in ΔE2B

and ΔE3B ESC lines that showed an XCI phenotype by introducing a wild-type *Hcfc1* (129 origin)

or a control transgene (Fig. 2a, Extended Data Fig. 3a). RT-qPCR analysis showed that *Xist* levels

were rescued upon restoration of *Hcfc1* expression levels at day 4 of ESC differentiation in a

transfected pool of cells, and in individual clones with a stably integrated transgene, confirming a

dose-dependent regulatory role for *Hcfc1* in XCI (Fig. 2b,c, Extended Data Fig. 3b,c). To test

whether HCFC1 is required for activation of XCI, we studied the effect of dTAG-13-mediated

depletion of HCFC1 (Fig. 2d). Correct 3' integration of a 2xFLAG-V5-tag combined with

FKBP12^{F36V} in both *Hcfc1* (HCFC1-FKBP) alleles was confirmed by PCR (Extended Data Fig.

3d). HCFC1 is glycosylated and cleaved by OGT to form dimers that stay associated[21,22]. We

confirmed loss of expression of both the N- and C-terminal domains after 48 hours of dTAG13

treatment (Fig. 2e, Extended Data Fig. 3e and 4e). Loss of HCFC1 occurs very rapidly, as

examination of the C-terminal peptide showed a complete loss within 2 hours after addition of

dTAG-13 (Fig. 2e). Depletion of HCFC1 during different time intervals of ESC differentiation

resulted in a significant reduction of *Xist* expression (Fig. 2d,f). *Xist* RNA-FISH analysis

confirmed loss of *Xist* expression and a severely reduced percentage of *Xist*-coated X

chromosomes at day 3 of differentiation (Fig. 2g,h). Allele-specific RNA-seq analysis also showed

that positive *Xist cis*-regulator *Jpx* was significantly downregulated, and *Ftx* showing a

downregulation trend, whereas expression of pluripotency factors *Nanog*, *Pou5f1*, *Sox2* and *Rex1*

remained unaffected, indicating that ESC differentiation proceeded normally in the absence of

HCFC1 within the analysed time windows (Fig. 2i, Extended Data Fig. 3f). On the contrary,

expression of several *trans*-regulators of *Xist*, including *Rnf12, Yy1, Kdm5c* and *Rif1* was

significantly increased upon loss of HCFC1, suggesting a compensatory mechanism in *Xist*

regulation (Extended Data Fig. 3g). dTAG-mediated loss of HCFC1 did not affect negative cis-

regulators of *Xist*, such as *Tsx* and *Tsix* (Fig. 2i). Consistent with the observed substantial *Xist*

downregulation, allele-specific expression analysis of X-linked genes indicated loss of XCI in

HCFC1-depleted ESCs at day 3 of differentiation (Fig. 2j). To discern if this loss of silencing is

due to *Xist* loss or HCFC1 loss, we removed HCFC1 and forced *Xist* expression in undifferentiated

ESCs by means of a doxycyclin inducible promoter at the endogenous *Xist* locus. We observed

that XCI still happens properly since X-linked *Rnf12* was silenced while a gene escaping XCI,

Kdm6a, was not (Extended Data Fig. 3h-k). These findings identify HCFC1 as a dose-dependent

activator of XCI and indicate that *Hcfc1* is required for upregulation of *Xist* and positive *Xist*

regulatory genes.

Overlapping genome-wide enrichment of HCFC1, YY1 and Integrator

HCFC1 is a member of the Set1A/B and MLL1/2 COMPASS complexes, mediating H3K4me3

deposition at gene regulatory elements which promotes transcription activation through Integrator

complex recruitment and promoter-proximal eviction of paused RNA Pol II[23–26]. To

mechanistically understand the role of HCFC1 in XCI and gene regulation in general, we re-

analysed published ChIP-seq datasets examining the genome-wide distribution of HCFC1 and the

COMPASS core member DPY30, as well as YY1, H3K4me3 and Ser-5 phosphorylated Pol II in

undifferentiated mouse ESCs[25,27,28]. This analysis confirmed genome-wide co-enrichment of

DPY30 with HCFC1 and their catalytic product H3K4me3 (Fig. 3a). Of 31,247 HCFC1 peaks in

ESCs, 70.5% were found in genes, of which 48.7% were at promoters (Extended Data Fig. 4a).

Within promoters, 81.5% of YY1 peaks overlapped with HCFC1 (Fig. 3b,c), confirming the previously described overlap in HCFC1 and YY1 binding to chromatin in human cells[20,29]. This analysis also shows that many HCFC1 peaks are not co-bound by YY1, indicating that other transcription factors might be recruiting HCFC1 to those sites.

To understand HCFC1-mediated regulation of *Xist*, we examined HCFC1's distribution by ChIP-seq on day 3 of ESC differentiation when *Xist* expression is upregulated. For this, we generated 2xFLAG-V5-tagged HCFC1 (HCFC1-FLAGV5) female ESC lines through CRISPR/Cas9-mediated integration (Extended Data Fig. 4b). ChIP-seq analysis revealed 20,660 HCFC1 binding peaks, the majority of which (77.1%) were located within genes, of which 44.6% were at promoter regions (Extended Data Fig. 4a). Comparison of HCFC1 binding between day 0 and day 3 revealed only minor changes (Fig. 3c,d), and similarly, the overlap with YY1 binding sites was evident, with 84.5% of YY1 promoter peaks overlapping with HCFC1 peaks at day 3 of differentiation (Fig. 3c,e). As described for human cells, HCFC1 peaks were highly enriched at CpG islands and promoters (Extended Data Fig. 4c)(Michaud 2013). Moreover, HCFC1 peaks showed strong enrichment for the YY1 binding motif and motifs of other transcription factors implicated in HCFC1 recruitment, such as RONIN and ZNF143/STAF (Extended Data Fig. 4d)[29]. Similar to undifferentiated ESCs, H3K4me3 ChIP-seq analysis at day 3 of differentiation revealed a high level of overlap with HCFC1 peaks, consistent with HCFC1's role as a member of the COMPASS complexes responsible for H3K4me3 catalysis (Fig. 3a).

Recent studies have indicated an important role for COMPASS-mediated H3K4me3 in recruitment of Integrator to target genes[25]. To investigate the role of HCFC1-COMPASS-mediated catalysis of H3K4me3 and subsequent recruitment of Integrator, we studied the presence and distribution of H3K4me3 and INTS11, a member of the endonuclease module of the Integrator complex, before and after dTAG-13-mediated depletion of HCFC1. Western blot analysis revealed a global decrease of H3K4me3 presence at day 3 of differentiation upon HCFC1 removal by dTAG-13 treatment (Extended Data Fig. 4e,f). ChIP-seq analysis identified a genome-wide reduction of H3K4me3 enrichment at 4,991 sites (9.8% of all 51,021 H3K4me3 peaks) upon HCFC1 removal, confirming an important role for HCFC1 in H3K4me3 deposition, although the loss was less prominent than reported for the depletion of DPY30 or RBPP5 (Fig. 3f, Extended Data Fig. 4g)[25]. As expected, correlation analyses between RNA-seq and ChIP-seq data showed that gene

expression changes following HCFC1 depletion are positively associated with changes in

H3K4me3 levels, (Fig. 3g, Extended Data Fig. 4h,i). Sites that displayed decreased H3K4me3

enrichment upon HCFC1 depletion did not exhibit or only exhibited a mild loss of INTS11

binding, indicating that INTS11 recruitment to those sites is predominantly H3K4me3 independent

(Fig. 3h, Extended Data Fig. 4g,i). Interestingly, HCFC1 sites that showed reduced H3K4me3

deposition upon HCFC1 removal also displayed very strong YY1 binding (Fig. 3h). Our results

indicate that HCFC1 plays a crucial role in H3K4me3 catalysis at HCFC1-YY1 co-bound sites,

but that the presence of this histone modification is unrelated to the recruitment of Integrator.

YY1-mediated recruitment of the COMPASS and Integrator complexes

Our results suggest a putative interaction between YY1 and HCFC1 in regulating H3K4me3

deposition. To test whether YY1 is indeed involved in recruitment of COMPASS to target

sequences, we generated two female 2xFLAG-V5-tagged YY1 ES cell lines and performed FLAG

pull-downs followed by mass spectrometry analysis (Extended Data Fig. 5a,b). Our data confirmed

strong interactions with the INO80 chromatin remodeling complex, as previously described[30] and

with HCFC1 and COMPASS member RBBP5, and revealed weak interactions with other

COMPASS complex subunits, likely reflecting indirect protein–protein interactions (Fig. 4a).

Interestingly, we also identified most Integrator complex subunits (INTS1, INTS2, INTS4, INTS7,

INTS8, INTS9, INTS11 and INTS12) as strong YY1 interactors, suggesting a role for YY1 in

recruitment of Integrator to gene regulatory elements. This includes all subunits of the RNA

endonuclease module of Integrator, responsible for cleavage of nascent RNA associated with

paused RNA pol II[31].

Interrogation of published ChIP-seq datasets of INTS11 distribution in undifferentiated ESCs

indicates a high level of overlap between genome-wide distribution of YY1 and the Integrator

complex (Fig. 4b, Extended Data Fig. 5c), with the immense majority (95.8%) of YY1 peaks at

promoters overlapping with INTS11 (Fig. 4b,c). Indeed, overlapping YY1-INTS11 peaks are

enriched for the YY1 motif while INTS11-only peaks are not (Extended Data Fig. 5d). These

results suggest a dual role for YY1 in recruitment of both HCFC1 and Integrator complexes.

To functionally test whether HCFC1 and Integrator recruitment is YY1-dependent, we generated

compound knock-in 2xFLAG-V5-Hcfc1:FKBP12^{F36V}– 2xHA-Yy1 female (XX) ESC lines

(HCFC1-FLAGV5/YY1-FKBP ESCs) and analysed the effect of dTAG-13-mediated depletion of YY1 on HCFC1 and Integrator recruitment to regulatory elements (Fig. 4d). Western blot analysis confirmed complete loss of YY1 within 2h of dTAG-13 treatment (Extended Data Fig. 5e). We then proceeded to differentiate these cells for 3 days and treated them with dTAG-13 for the last 2 days of differentiation (Extended Data Fig. 5e). HCFC1 ChIP-seq analysis at day 3 of differentiation revealed that HCFC1 enrichment at the great majority of its peaks does not change upon YY1 removal, indicating that HCFC1 recruitment is predominantly mediated by other transcription factor (TFs) (Fig. 4e). Nevertheless, 7.9% of HCFC1 peaks are YY1-dependent, and analysis of DNA binding motifs at HCFC1 peaks with decreased enrichment in the YY1 dTAG-13 condition compared to peaks with increased enrichment confirmed strong presence of the YY1 consensus binding motif (Fig. 4e,f). At 70 promoter-associated loci within this subset, loss of both YY1 and HCFC1 is associated with reduced gene expression (Fig 4h), indicating that YY1 recruits HCFC1 to a subset of promoters to activate their expression.

Since eight subunits of the Integrator complex were significantly pulled down by YY1[32], we analysed INTS11 recruitment dependence on YY1. Loss of YY1 leads to reduction of INTS11 recruitment at 23.6% of all INTS11 binding sites genome-wide (Fig. 4e). INTS11 peaks with decreased enrichment in the YY1 dTAG-13 condition are strongly enriched for the YY1 binding motif (Extended Data Fig. 5f). Moreover, concomitant loss of YY1 and INTS11 at certain loci is associated with reduction in expression of 93 proximal genes such as *Cnnm2* and *Zfp37* at day 0[33] and day 3 of differentiation (Fig. 4i, Extended Data Fig. 5g-k). Although western blot analysis showed a general decrease in H3K4me3 (Extended Data Fig. 5i,m), H3K4me3 ChIP-seq analysis indicated that 2,057 sites (3.5% of all 58,455 H3K4me3 peaks) showed a reduction of H3K4me3, a lower percentage compared to HCFC1 (Fig. 3f and Fig. 4e,i). Interestingly, most sites displaying a reduction of Integrator recruitment after YY1 depletion show persistent H3K4me3 enrichment (Fig. 4j). Importantly, at these sites INTS11 recruitment is HCFC1 independent (Fig. 4j), in line with our finding that dTAG-13-mediated depletion of HCFC1 revealed limited effects on INTS11 recruitment (Fig. 3h). These results highlight an important dual regulatory role for YY1 in genome-wide recruitment of COMPASS or Integrator to target genes to activate their expression.

COMPASS- and Integrator-mediated regulation of XCI

To understand the XCI phenotype in *Hcfc1* mutant cells, we studied the role of HCFC1/COMPASS, YY1 and Integrator in XCI regulation. ChIP-seq analysis of HCFC1 and YY1 binding in day 0 ESCs showed clear HCFC1 and YY1 overlapping peaks around the *Xist, Jpx* and *Ftx* promoter and regulatory elements (Fig. 5a)[27,28]. Analysis of day 3 differentiated female ESCs revealed HCFC1 binding at the *Xist* regulatory element, as well as at regulatory elements of negative (*Tsix,*) and positive *(Jpx, Ftx)* regulators of *Xist* (Fig. 5a). Importantly, dTAG-13-mediated loss of HCFC1 or YY1 both leads to reduction of H3K4me3 at the *Xist* regulatory element accompanied by a reduction in *Xist* expression at day 3 of ESC differentiation (Fig. 2f and Fig. 5a,b). Integrator recruitment at the *Xist* regulatory element appears largely unaffected by HCFC1 and YY1 depletion and their concomitant loss of H3K4me3. *Jpx* expression is also significantly reduced upon HCFC1 or YY1 removal (Fig. 2i and 5b). In contrast to *Xist*, H3K4me3 deposition at the *Jpx* promoter remains unchanged in both conditions despite loss of HCFC1 recruitment at the *Jpx* promoter following YY1 depletion (Fig. 5a). Notably, INTS11 recruitment to the *Jpx* promoter is completely abrogated upon YY1 loss indicating a crucial role for YY1 in Integrator recruitment to the *Jpx* promoter. Although loss of HCFC1 results in a mild, non-significant reduction in *Ftx* expression, YY1 depletion causes a significant decrease in *Ftx* expression. In both cases, only INTS11 recruitment, and not H3K4me3 deposition, is affected (Fig. 2i and 5a,b), suggesting an indirect effect of HCFC1 on *Ftx* expression, possibly *via* loss of promoter co-regulation and/or mass action. While *Tsix* expression is affected by YY1 removal it is not affected by HCFC1 removal. Also, INTS11 recruitment to Tsix' promoter is YY1-dependent and accompanied by a slight reduction in H3K4me3 deposition (Fig. 2i and 5a,b). Interestingly, the more severe effect of YY1 depletion on *Xist*, *Jpx* and *Ftx* compared to HCFC1 could be explained by the positive effect YY1 has on *Hcfc1* expression (Fig. 5b).

These findings indicate distinct mechanisms in the regulation of XCI. HCFC1- and YY1-mediated H3K4 trimethylation is essential for *Xist* activation, with HCFC1 activity confined to *Xist*, while YY1 plays a dual role by recruiting the Integrator complex to both *Xist*-activating (*Jpx*, *Ftx*) and -repressing (*Tsix*) cis-regulatory elements. Interestingly, both HCFC1 and YY1 are required for H3K4me3 enrichment at the *Xist* locus, despite continued HCFC1 recruitment in the absence of YY1, suggesting an additional role for YY1 in catalyzing H3K4me3 at the *Xist* regulatory element.

Discussion

Here, we provide evidence that HCFC1 plays a critical role in regulating XCI. Knockout and rescue experiments indicate that HCFC1 is a dosage-sensitive X-encoded activator of XCI involved in the regulation of *Xist* and *Xist* regulatory genes. Previous studies involving HCFC1 heterozygous knockout mice described an XCI phenotype with complete skewing of XCI towards inactivation of the mutant allele[34]. As described for *Rnf12*, this skewing phenotype could be explained by a continued requirement for *Hcfc1* during the XCI process but could also be explained by the essential function of HCFC1 in gene regulation in general and cell cycle control[34]. Unfortunately, early lethality of *Hcfc1*[-/-] embryos due to imprinted XCI problems prohibited studying the role of HCFC1 in random XCI in the developing embryo.

Although the effect of the combined heterozygous deletions of large regions containing *Hcfc1* and *Rnf12* on *Xist* expression and XCI is severe, a low percentage of cells is still capable of initiating XCI upon ESC differentiation pointing to the presence of additional XCI-activators located in other regions and not detected in this study. Interestingly, our study did involve deletions encompassing genes previously implicated in regulating XCI including *Kdm5c* and *Gata1*, yet these deletions had no detectable effect on XCI in our system[14,15]. This result may be due to limited expression (*Gata1*) or sensitivity of our model system and readout (*Kdm5c*), which might not have been able to capture subtle effects on XCI.

HCFC1 is a member of the SET1A/B-COMPASS and MLL1/2-COMPASS complexes in mammals (reviewed in[35]). Our study confirms the importance of HCFC1-mediated recruitment of COMPASS to target sequences, H3K4me3 deposition, and proximal gene expression. However, in line with its non-core function within the complex, the effect of loss of HCFC1 on H3K4me3 deposition is less severe than depletion of core COMPASS subunits such as DPY30 or RBBP5[25,35]. Moreover, we show that Integrator recruitment to HCFC1-dependent H3K4me3 promoters is largely invariant, indicating other mechanisms at play to recruit Integrator to those sites. Interestingly, we noticed strong enrichment of most Integrator subunits in our YY1 pulldown studies, suggesting a direct role for YY1 in recruiting Integrator to target sites. Indeed, many YY1 sites showed reduced INTS11 recruitment upon YY1 removal and as with HCFC1, this seemed to be independent of H3K4me3, while the great majority of proximal promoters were downregulated.

Also, although most HCFC1 binding seemed to be YY1 independent and thus mediated by other TFs, we observed a subset of YY1-dependent promoter-proximal HCFC1 sites that correlate with gene expression. Our study highlights the versatile role of YY1 in mediating recruitment of Integrator and HCFC1-COMPASS genome-wide. Although previous studies indicated a role for histone modification H3K4me3 in recruitment of the Integrator complex, our studies reveal a role for YY1 in recruitment of Integrator that seems independent of H3K4me3[25].

By dissecting the contribution of YY1 and HCFC1 to XCI regulation, we show that YY1 and HCFC1 direct dose-dependent catalysis of H3K4 tri-methylation at the *Xist* regulatory element and activation of *Xist* transcription (Fig. 5c). Interestingly, although HCFC1 recruitment to this element appears unaffected in the absence of YY1, HCFC1 acute depletion still leads to loss of H3K4me3 suggesting a role for YY1 in recruiting COMPASS complexes independent of HCFC1, or indirectly by recruitment of other factors that facilitate H3K4me3 deposition. HCFC1 depletion also leads to expression reduction of the positive regulators of *Xist, Jpx*, *Ftx*, all located in the *Xist* topological associated domain (TAD). Since HCFC1 depletion does not lead to a loss of H3K4me3 at these co-activators, downregulation of *Jpx* and *Ftx* might be an indirect effect driven through loss of *Xist* co-activation. As HCFC1 has also been described as a member of the NSL complex, its loss may affect histone acetylation and thereby decrease gene expression of genes involved in XCI, a hypothesis that was not tested in the present study but warrants further investigation[36,37]. In contrast to the effects of HCFC1 loss, YY1 reduction decreases expression of genes in both the Xist-TAD and the neighbouring Tsix-TAD, which harbours negative regulators of *Xist*. Beyond its role in catalysing H3K4me3 at the *Xist* regulatory element, YY1 recruits both HCFC1 and INTS11 to the *Jpx* and *Tsix* promoters, and INTS11 alone to the *Ftx* regulatory element. Interestingly, YY1 depletion affects Integrator recruitment to these elements but loss of HCFC1 recruitment seems to have limited effect on H3K4me3 deposition pointing to an additional mechanism to recruit the core COMPASS complex independently of YY1 and HCFC1. Our results therefore indicate a role for YY1 in regulating *Xist* expression through H3K4me3 catalysis at its regulatory element, and regulating *Jpx, Ftx* and *Tsix* expression through recruitment of Integrator.

Our findings highlight the cooperative activities of HCFC1 and RNF12 in female exclusive initiation of XCI (Fig. 5d); through dose-dependent breakdown of REX1, RNF12 facilitates increased recruitment of YY1 to different regulatory elements at the XIC in female cells[7]. The effect on *Xist* expression is more pronounced in female cells likely through the dual action of YY1

on *Xist* and its positive regulators *Jpx* and *Ftx*, and *Hcfc1* itself. HCFC1 on the contrary, only

appears to act through dose-dependent recruitment of COMPASS to the *Xist* regulatory region,

where the combined action of YY1 and HCFC1 is required for catalysis of H3K4me3. In female

cells, the increased activity of HCFC1, RNF12 and other XCI-activators amplifies the activation

of *Xist* to sufficient levels to spread locally in *cis* and silence its negative regulator *Tsix*. Negative

feedback will prevent inactivation of all X chromosomes except one. These findings highlight the

complex and concerted action of RNF12 and HCFC1 in female exclusive initiation of XCI by

dose-dependent action converging on the same XCI regulatory elements.

Methods

Cells and Cell Culture

Cell culture

F1 2-1 mESCs were grown on irradiated male mouse embryonic fibroblasts (MEFs) in medium containing DMEM (Gibco, 11995065) supplemented with 15% fetal calf serum, 100 U ml^{-1} penicillin, 100 µg mL^{-1} streptomycin (Sigma-Aldrich, P0781), 0.1 mM non-essential amino acids (Lonza, BE13-114E), 0.1 mM 2-mercaptoethanol (Gibco, 31350010) and 5.000 U mL^{-1} home-made LIF. Cells were passaged at least two times prior to differentiation. mESC were depleted of MEFs by pre-plating dissociated cells for 45 minutes at 37°C in non-gelatinized plates. The supernatant containing mESCs was counted and subsequently spun down for 5 min at 300 × g. Specific cell numbers were plated for different time points for monolayer differentiation in medium consisting of IMDM GlutaMAX™ (Gibco, 31980030) supplemented with 15% FCS, 0.1 mM NEAA, 100 U mL−1 penicillin, 100 µg mL−1 streptomycin, 23,75 µL L−1 monothioglycerol (Sigma-Aldrich, M6145) and 42 µg mL−1 ascorbic acid (Sigma-Aldrich, A92902). Degradation of HCFC1 or YY1 in mESCs or differentiation was achieved by refreshing the media with addition of 500 nM dTAG-13 (Tocris, 6605) or equivalent amounts of DMSO at the indicated time points.

Generating Rnf12 lines

Rnf12$^{+/-}$ mESC lines were generated in F1 2–1 hybrid (129/Sv-Cast/Ei) mESC by CRISPR/Cas9 mediated targeting using different sets of gRNAs. The Rnf12$^{+/-}$ ESC line Δ1 was generated by targeting mESCs with PX459 vectors with gRNAs targeting intron 2 and the 3'UTR using Gene Pulser 0.2 cm-gap cuvettes (Biorad, 1652086) at 118 kV, 1200 µF and ∞Ω in a Gene Pulser xCell electroporation system (Biorad). Twenty-four hours post-transfection, medium was refreshed and supplemented with 1 µg mL−1 puromycin (Sigma-Aldrich, P8833) for 36h. Drug-resistant clones were picked, expanded and characterized by PCR. Due to chromosomal instability upon inducing ΔJ, a second Rnf12$^{+/-}$ line was generated (Rnf12$^{+/-}$ Δ2). Rnf12$^{+/-}$ Δ2 was generated by transfecting 2×1 µg PX459 alternative gRNA expressing vectors as well as 2 µg homology arm vector with lipofectamine 2000 (Invitrogen, 11668019). Twenty-four hours post-transfection, the medium was refreshed and supplemented with 1 µg mL−1 puromycin for 48h. Drug-resistant clones were picked, expanded and characterized by PCR.

Generating deletion lines

Deletion regions were designed so that genes within the same evolutionary strata and syntenic blocks were removed together[17,18]. As described previously, deletion lines were generated by transfecting 2×1 µg PX459 gRNA expressing vectors as well as 2 µg homology arm vector (containing a neomycin resistance cassette) with lipofectamine. 24h post-transfection, the medium was refreshed and supplemented with 1 µg mL−1 puromycin for 48h followed by 285 µg mL−1 Geneticin (G418) (Gibco, 11811-031) for 4-6 days for ΔA through ΔK. Combinations of 5′ and 3′ gRNAs were used to generate the incremental deletions ΔE1 to Δ E3 and ΔE2A to ΔE3B. No Geneticin selection was performed for the incremental deletions. Drug-resistant clones were picked, expanded, and characterized by microsatellite PCR. ΔA through ΔI and ΔK as well as all incremental deletions were generated in Rnf12$^{+/-}$ Δ1. ΔJ was generated in Rnf12$^{+/-}$ Δ2. No deletions were generated between ΔK and the telomere due to genomic instability.

Generating rescue lines

The coding sequence of *Hcfc1* was amplified from 129/Sv mESC cDNA and cloned into the pCAG-2×FLAG-V5-IRES-Puro expression vector. Rnf12$^{+/}$, ΔE2B and ΔE3B mESCs were transfected with 2 µg of either the *Hcfc1* expression vector or an empty vector control, as described previously. 24h post-transfection, the culture medium was refreshed and supplemented with 1 µg mL^{-1} puromycin for 5 days to select for stable integration. After two passages without selection, cells were split: one fraction was used to isolate individual clones, and the remaining pool was subjected to differentiation. Cells were harvested at day 4 of differentiation and analyzed by RT-qPCR.

Generation of knock-in ESC lines

Female mESCs harboring a heterozygous doxycycline responsive endogenous *Xist* promoter[38] were used as a parental cell line for generation of endogenously tagged *Hcfc1* lines. In short, mESCs were transfected with gRNAs targeting the 3' end and HCFC1-2×FLAG-V5-P2A-mTagBFP2 or HCFC1-2×FLAG-V5-FKBP^{F36V}-P2A-mTagBFP2 constructs for homology-directed repair. 24 hours post-transfection cells were selected with medium supplemented with puromycin for 48 hours. Drug-resistant clones were picked on EVOS M5000 Imaging System (Thermo Scientific, AMF5000SV) for highest fluorescent intensity. To generate endogenous C-terminally tagged YY1 ESCs for dTAG-inducible degradation, HCFC1-2×FLAG-V5-P2A-mTagBFP2 cells were transfected with gRNAs targeting the 3'UTR and using a

pAW62.YY1.FKBP.knock-in.mCherry vector[33] (Addgene #10437) for homology-directed repair.

Due to a SNP in the 3'UTR of *Yy1*, two separate gRNAs targeting either 129/Sv or Cast/Ei were

selected. Drug-resistant clones were picked on EVOS M5000 Imaging System. All positive clones

were expanded and screened for homozygous insertions flanking the integration site by PCR and

western blotting.

Female ESCs expressing 2xFLAG-V5-YY1 were generated in a *Tsix-STOP* background by

random integration of a pCAG 2xFLAG V5 YY1 construct. Cells were electroporated and

subjected to puromycin selection 24 h later. After one week, individual clones were isolated and

expanded. Transgene expression was confirmed by western blotting, and karyotypes were

assessed. Two independent clones with normal karyotypes and YY1 expression comparable to

endogenous levels were selected for FLAG pull down experiments.

Western Blot

Nuclear protein extracts were performed as described previously[39] with modifications. All buffers

were supplemented with 1× protease inhibitors (Roche, 11836170001). Cells were harvested by

scraping in cold PBS on ice and subsequently centrifuged for 5 min, 300 x g at 4°C. The pellet

was resuspended in 5× pellet volume Buffer A (10 mM Hepes pH 7.6, 1.5 mM MgCl2 and 10 mM

KCl, 0.5 mM DTT), incubated for 10 min at 4°C, vortexed for 30 sec and spun down at 900 x g,

for 5 min, 4°C. The supernatant was transferred and the nuclear pellet was resuspended in 2× pellet

volume Buffer C (20 mM Hepes pH 7.6, 25% glycerol, 420 mM NaCl, 1.5 mM MgCl2, and

0.2 mM EDTA, 0.5 mM DTT, 0.05% NP-40). After rotating the solution for 30 min, 4°C, it was

subsequently spun down for 10 min at 18.000 x g, 4°C. The supernatant containing the nuclear

fraction was collected and the concentration was determined using Nanodrop or The Pierce BCA

Protein Assay Kit (Thermo Scientific, 23227). For western blot analysis, lysates were diluted 1:1

with 2× Laemmli Sample Buffer (0.125M Tris-HCl pH 6.8, 4% SDS, 20% (w/v) glycerol, 0.01%

bromophenol blue, 150 mM DTT) and boiled for 5 min at 95°C. Equal concentration of protein

lysates were loaded in the 4–15% Criterion™ TGX™ Precast Midi Protein Gel, 18-well (Biorad,

#5671084) with Precision Plus Protein Dual Color Standards (Biorad, #1610374) and run in 1x

Tris/Glycine/SDS buffer (Biorad, #1610772). Samples were transferred to a Trans-Blot Turbo

PVDF membrane (Biorad, #1704157) using semi-dry transfer in the Trans-Blot Turbo Transfer

System for 7 min at 2.5 A constant up to 25V. After transfer, membranes were blocked with 5%

milk powder in 1×TBS (0.02M Tris-HCl, 0.15M NaCl, pH 7.6). The membrane was washed 3×

with 1×TBST (0.02M Tris-HCl, 0.15M NaCl, 0.1% Tween-20, pH 7.6), incubated with the
appropriate primary antibodies, HCFC1 Carboxy-terminal Antigen antibody (Cell signaling,
50708S, 1:1000), V5 antibody (Invitrogen, R96025, 1:5000), YY1 antibody (Santa Cruz, sc-7341,
1:200) and/or H3K4me3 antibody (Sigma-Aldrich, 05-1339, 1:1000). Membranes were washed
3× with 1× TBST and incubated with the appropriate secondary antibodies, IRDye 800CW Goat
anti-Mouse IgG (LI-COR Biosciences, 926-32210), IRDye 800CW Donkey anti-Mouse IgG (LI-
COR Biosciences, 926-32212) and/or IRDye® 680RD Donkey anti-Rabbit IgG (LI-COR
Biosciences, 926-68073) and visualised on the Odyssey CLx Imager. Last, the membranes were
incubated with β-Actin−Peroxidase (Sigma-Aldrich, A3854), stained with ECL Western blotting
detection reagents (Cytyva, RPN2209) and visualised on the AI600 Imager.

Expression analysis and RNA-seq

mESCs were depleted of MEFs as described above, supernatants were spun down and cell pellets
were lysed. Differentiated cells were washed 1× in PBS and lysed directly in the culture dish prior
to extraction. Total RNA was extracted using the ReliaPrep™ RNA Miniprep kit (Promega,
Z6012) with increased DNase treatment (25 min at room temperature). Reverse transcription was
performed on 500 ng RNA using Superscript III (Invitrogen, 18080093), random hexamers
(Invitrogen, N8080127) and RNaseOUT (Invitrogen, 10777019). The cDNA was diluted 1:1 in
water prior to amplification. RT-qPCRs, allele-specific RT-qPCRs and qPCRs were performed in
triplicate in the GoTaq qPCR Master Mix (Promega, A6002) in a CFX384 Touch Real-Time PCR
Detection System (BioRad, LJB22YE8Z). Allele-specific primer concentrations were determined
and optimized using pure gDNA of 129/Sv and Cast/Ei mice. Rplp0 was used as reference gene
for relative quantification of HCFC1-FKBP cells. Hist2h2aa1 was used as reference gene for all
other qPCRs. All primers are included in Extended Data Table 1.

Sequencing libraries were prepared using the Illumina TruSeq stranded mRNA library preparation
kit. Libraries from the deletion cell lines were sequenced on a HiSeq 2500 sequencer, generating
single-end 50 bp reads. Libraries from the HCFC1 dTAG-13 and DMSO-treated samples were
sequenced on an Illumina NextSeq 2000 sequencer, resulting in paired-end read clusters of 50
bases in length.

  RNA-seq analysis

RNA-seq reads were processed allele-specifically. First, we created an N-masked reference

genome from mm10 using SNPsplit v0.3.4[40], masking all SNPs between the 129/Sv and Cast/Ei

strains based on the dbSNP v142 database[41]. Reads were trimmed using TrimGalore v0.6.7[42] and

aligned to the N-masked reference genome using HISAT2 v2.2.1[43]. Mapped reads were then

assigned to either the 129/Sv, Cast/Ei, or unassigned BAM file based on the best alignment using

SNPsplit (default settings with --paired for the paired-end data). All allele-specific and unassigned

reads were merged into a composite BAM file using Samtools v1.10[44].

Transcript-level counts were obtained from each BAM file using featureCounts v2.0.6[45] with the

Ensembl v98 gene annotations (-t exon -s 2). We normalized the raw counts to transcripts per

million (TPM) using a custom Python script. Gene expression levels were visualized in bar plots,

showing TPM values normalized to the average TPM of DMSO-treated or untreated control

samples. To validate the deleted regions, $\log_2(\text{TPM}+1)$ values were calculated per allele and

visualized in a heatmap across all samples.

Differential expression analyses were performed using DESeq2 v1.30.1[46] with parametric

dispersion fitting. Fold changes were shrunk using the ashr v2.5-54 method[47]. Differentially

expressed genes (DEGs) were defined as those with an adjusted p-value < 0.05 and an absolute

$\log_2(\text{fold change}) > 1$. Results were visualized in volcano plots, with autosomal and X-linked

DEGs coloured separately.

To assess X-chromosome silencing, we calculated allelic ratios for $Rnf12^{+/-}$, ΔE, ΔE2B and ΔE3B

as well as HCFC1 dTAG-13 and DMSO-treated samples. For each condition, genes with more

than 20 allele-specific reads across all replicates were included. The allelic ratio was defined as:

505
$$\frac{Xi}{Xi + Xa}$$

where Xi represents 129/Sv and Xa represents Cast/Ei. Allelic ratios of X-linked genes were

visualized using violin plots combined with boxplots. Statistical significance was assessed using a

two-sided Mann-Whitney U test (α < 0.05).

We reanalysed YY1-FKBP and wild-type RNA-seq data from GSE99521[33] using the same allele-

specific pipeline as described above, except that the N-masked reference genome was generated

using SNPs between C57BL/6 and 129/Sv. Additionally, to assess expression dynamics of

candidate genes during early differentiation, we analysed a bulk RNA-seq dataset from

GSE151009[19]. Allele-specific count tables were downloaded and normalized to TPM values. For

both ΔXic^{B6} and ΔXic^{Cast} cell lines, allelic expression of *Hcfc1*, *Irak1, Rnf12* and *Xist* was plotted

over time in line plots, displaying the mean and 95% confidence interval. Statistical significance

between day 0 and day 1 was assessed using one-sided independent t-tests, followed by multiple

testing correction with the Benjamini-Hochberg method.

RNA-FISH

Xist probes were made from a 19 Kb Xist/Tsix plasmid (p510) using the Nick Translation Kit

(Abbott, 07J00-001) with red dUTPs (Abbott 02N34-050) according to the manufacturer's

instructions. Per coverslip, 3 µl of Xist probe was mixed with 1 µl salmon sperm DNA, 0.5 µl

NaOAc 3M and 100 µl 100% EtOH and centrifuged 30 min, 12.800 rpm at 4°C. Pellet was washed

with 500 µl cold 70% EtOH and let dry. 7 µl of formamide was added to the probe, incubated 10

min at 75°C and subsequently quenched on ice. 7 µl of hybridization buffer (20% Dextran sulfate,

2 mg/ml BSA, 20 mM VRC, 4×SSC), mixed and added to the coverslip.

ESCs were grown on gelatin coated coverslips at different time points for monolayer

differentiation. Cells were fixed in 4% PFA in PBS for 10 min while rocking and rinsed 3× with

PBS. Cells were subsequently permeabilized in 0.5% Triton, 20 mM VRC (NEB, S6639) in PBS

for 4 min on ice. Cells were sequentially dehydrated by addition of 70, 90 and 100% ethanol,

followed by airdrying. The probe, dissolved in hybridization buffer was added to the coverslip and

incubated overnight at 37°C in humid chamber with 50% formamide in 2× SSC. After

hybridization, coverslips were washed 3 × with 50% formamide in 2×SSC at 42°C, 2 × with 2×SSC

at 42°C and once with 2 × SSC supplemented with 4 µM Hoechst (Thermo Scientific, 62249)

before mounting with ProLong™ Gold Antifade (Invitrogen, P36930).

WGS

ESCs were depleted of MEFs by pre-plating dissociated cells for 45 minutes at 37°C in non-

gelatinized plates and subsequently grown for several days in a 6-well dish without MEFs. Cells

were collected and lysed overnight in low-SDS lysis buffer (50 mM Tris-HCl pH 8.0, 100 mM

EDTA, 0.2% SDS, 100 mM NaCl) with Proteinase K for Phenol-Chloroform DNA extraction.

Following DNA extraction, 10 µg DNA was treated with 4 µg/ml RNase A treated for 1 hour at

37°C followed by clean up with the QIAamp DNA Micro Kit (QIAGEN, 56304). Sequencing

libraries were prepared using the Nextera DNA Flex protocol from Illumina using 50 ng of

genomic DNA and 5 cycles of PCR. Libraries were sequenced on an Illumina HiSeq 2500

sequencer, generating 50 bp single reads. For ΔE, over 50 million reads were obtained, while all

other samples yielded at least 20 million reads.

WGS analysis

Reads were trimmed using TrimGalore v0.6.7[42] and aligned to the N-masked reference genome

(see RNA-seq analysis) using Bowtie2 v2.5.1[48]. Mapped reads were assigned to either the 129/Sv

or Cast/Ei BAM file using SNPsplit v0.3.4[40], based on the best alignment. Allele-specific read

counts were calculated for genomic windows of 100 kb using deepTools v3.5.5[49] bamCoverage (-

bs 100000 -of bedgraph --normalizeUsing None). For each window with more than 10 allele-

specific reads, the $\log_2$(fold change) between the 129/Sv and Cast/Ei allele was computed.

Statistical significance between both alleles was assessed using two-sided binomial tests with

p=0.5, assuming equal read counts from both alleles in non-deleted regions. Resulting p-values

were corrected for multiple testing using the Benjamini-Hochberg method. Data were visualized

in scatter plots showing the $\log_2$(fold change) across the entire X-chromosome and within the 67-

76.5 Mb region. For visualization, extreme $\log_2$(fold change) values were capped at ±2.

Statistically significant windows ($\alpha < 0.05$) were highlighted, as well as the window containing

the *Hcfc1* locus.

DNA FISH

BACs/Fosmids were selected inside (MeCP2; WI1-1364A8) and outside the deleted region

(HPRT; RP23-412J16 and Xist; WI1-2363H9). Probes were labeled using the Nick Translation

Kit (Abbott, 07J00-001) with red (Abbott 02N34-050) or green (Abbott 02N32-050) dUTPs

according to the manufacturer's instructions.

DNA FISH was performed as described previously[50]. Cells were blocked in metaphase by addition

of 12 µl/ml KaryoMAX Colcemid (Gibco, 15210040) to the culture medium for 2 hours. Cells

were trypsinized and resuspended in hypotonic buffer (0.075 M KCl) followed by fixation by

addition of 3:1 methanol-acetic acid solution. Metaphase spreads were made by dropping fixed

cells on coverslips and aged overnight at 37°C. Coverslips were equilibrated in 2×SSC (1× SSC:

0.15 M NaCl, 0.015 M sodium citrate), treated with RNAse A (1 µg/ml in 2×SSC) for 5 min at

37°C in humid chamber and washed 3× with 2×SSC at room temperature. Cells were
permeabilized in pre-warmed 1% pepsin in 0.01M HCl solution for 7.5 min at 37°C. After
permeabilization, cells were washed 2× with PBS and prefixed in 1% formaldehyde for 5 min at
room temperature. Coverslips were washed 2× with PBS and sequentially dehydrated by addition
of 70, 90 and 100% ethanol and airdrying.

5 µl of Nick-translated DNA probes was combined with 3 µg Mouse Cot-1 DNA (Invitrogen,
18440016) and 10 µg UltraPure™ Salmon Sperm DNA (Roche, 11467140001) and precipitated
using NaAc and ethanol. The pellets were washed with 70% ethanol, spun down and air dried. The
dried pellet was dissolved in formamide and denatured by incubating for 10 min at 75°C while
shaking (1000 Rpm) and subsequently mixed 1:1 with cold 2×hybridization buffer (50%
formamide, 2× SSC, 50 mM phosphate buffer (pH 7.0), 10% dextran sulfate and 1 ng/µl mouse
Cot DNA). Following preparation of the probe mix, it was added to the coverslips and incubated
on heatplate for 2 min at 75°C. Slides were transferred to a humid chamber with 50% formamide
in 2× SSC and incubated overnight at 37°C. The next day, slides were washed 2× in 50%
formamide in 2× SSC at 42°C, washed 2× in 2× SSC at 42°C, air dried and mounted with ProLong
Gold Antifade with DAPI (Invitrogen P36931).

ChIP-seq
ChIP-seq was performed as previously described[39] with several modifications. Up to 1×10^8
attached cells were fixed with 20mM disuccinimidyl glutarate (ThermoFisher, 20593-50MG) in
PBS for 45min RT, washed twice with PBS followed by fixing in 1% PFA (Sigma-Aldrich, F1635-
25ML) in PBS for 10min RT for the V5-HCFC1 and Integrator ChIPs, or only 1% PFA in medium
for 10min RT for the H3K4me3 ChIPs (V5-HCFC1 ChIP was performed in FLAGV5-HCFC1
YY1-FKBP ESCs). After crosslinking, cells were quenched by addition of 125 mM glycine for 5
min. Cells were washed 1× with PBS before scraping in PBS with 1x protease inhibitors (Roche,
11836170001). From this point onwards, all buffers contained protease inhibitors (Roche,
4693132001) unless stated. Fixed cells were pelleted and subsequently resuspended in LB1 buffer
(50 mM HEPES pH 8.0, 140 mM NaCL, 1 mM EDTA, 10% glycerol, 0.5% NP-40, 0.25% Triton
X-100) and incubated for 10 min at 4 °C while rotating, then in LB2 buffer (10 mM Tris-HCl pH
8.0, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA) and last in LB3 buffer (10 mM Tris-HCl pH
8.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1% Sodium Deoxycholate, 0.5% N-

laurolylsarcosine). DNA was sheared using a Bioruptor Pico (Diagenode, B01060001) for 10

cycles of 30 sec on and 30 sec off (single crosslink samples) or 14 cycles of 30 sec on and

30 sec off (double crosslink samples) to an average fragment length of 300-1000. 1% triton

X-100 was added to sheared chromatin and incubated for 10 minutes at 4°C and spun down

for 10 min at 4500 g. Chromatin was aliquoted, flash frozen in liquid nitrogen and stored at -

80°C until further use. Different amounts of chromatin were used for the H3K4me3 (125 ug),

HCFC1-V5 and INTS11 ChIP (150 ug). Each chromatin sample was diluted to 500 ul using ChIP

dilution buffer (1.1% Triton X-100, 0.01% SDS, 167 mM NaCl, 16.7 mM Tris-HCl and 1.2 mM

EDTA) and 1% of it was substracted (INPUT) and frozen at -20°C. A specific amount of

antibody was added (Ab; 2 ug H3K4me3 Ab, Cell Signaling Technologies, 9751S [C42D8]; 10

ug INTS11 Ab, Bethyl Laboratories, A301-274A; 7ul V5 Ab, Invitrogen R960-25) to each

chromatin sample and incubated with rotation overnight at 4°C. The next day, 30 ul Dynabeads

protein G beads were washed (Invitrogen, 10004D) 2x with 500 ul ChIP dilution buffer. Chromatin

with antibody was added to the beads and incubated 2h at 4°C. Afterwards, by magnetic separation,

beads were washed 5 min 4°C 3x with Low salt wash buffer (20 mM Tris-HCl, 2 mM EDTA, 1%

Triton X-100, 150 mM NaCl, 0.1% SDS), 1x with High salt wash buffer (20 mM Tris-HCl, 2 mM

EDTA, 1% Triton X-100, 500 mM NaCl, 0.1% SDS), 1x with LiCl wash buffer (10 mM Tris-

HCl, 1 mM EDTA, 0.25 M LiCl, 1% NP-40, 1% deoxycholate). Beads were finally washed with

TE without protease inhibitors. 200 ul ChIP elution buffer (0.1 M NaHCO3, 10 mM EDTA and

1% SDS) were added to the beads with 22 ml proteinase K (10 mg/mL) and 5 ml RNase (10

mg/mL), and shaken 1,000 rpm on an orbital thermomixer for 2 h at 37°C and finally at 65°C

overnight. DNA was then cleaned using Zymo ChIP DNA clean concentrator kit (Zymo

Research, D5202) and eluted in 12 ul. 0.5 ul was used to confirm ChIP efficiency. The

concentration of the ChIP samples was measured, and sequencing libraries were prepared using a

ThruPLEX DNA-seq Kit (Takara Bio, R400674). Libraries were sequenced on an Illumina Nextseq 2000 sequencer, generating 50 bp paired reads.

were sequenced on an Illumina Nextseq 2000 sequencer, generating 50 bp paired reads.

ChIP-seq analysis

Reads were trimmed using TrimGalore v0.6.7[42] and aligned to the masked reference genome

(see RNAseq analysis) using Bowtie2 v2.5.1[43]. For datasets containing Drosophila melanogaster

spike-ins, reads were also aligned to the dm6 genome, and those with higher alignment scores to

dm6 than to mm10 were separated into a dedicated FASTQ file. Remaining reads were assigned

to either the 129/Sv, Cast/Ei, or unassigned BAM file based on the best alignment using SNPsplit

(--paired). Allele-specific and unassigned reads were merged into a composite BAM file using

Samtools v1.10[44].

Peak calling was performed on the composite BAM files using MACS3 v3.0.1[51] with the callpeak

function (-f BAMPE), comparing ChIP samples to their corresponding input controls. Bigwigs

with signal tracks were generated using MACS3 bdgcmp with the ppois method. Peaks were

annotated using the annotatePeaks.pl script from Homer v4.11[52], and motif enrichment analysis

was conducted with findMotifsGenome.pl (-size 200). Small annotation categories were merged

for clarity: 3' UTR and TTS were combined into "3' UTR - TTS", and exon and non-coding into

"exon".

Several ChIP-seq datasets were reanalysed using the same approach as described above. Reads

from a YY1 ChIP-seq dataset in day 0 ESCs were obtained from GSE240684[27] and analysed allele-

specifically. In addition, we downloaded ChIP-seq data for H3K4me3, DPY30, PolIISer5, and

INTS11 in day 0 ESCs from GSE181714[25], as well as a HCFC1 ChIP-seq dataset in day 0 ESCs

from GSE49847[28]. As these datasets were generated from non-hybrid cells, reads were mapped to

the mm10 genome, and the resulting BAM files were used as composite BAM files for all

subsequent analyses.

Overlap between peak locations

To assess overlap between HCFC1 binding and other chromatin features, HCFC1 day 0 peaks

were used as a reference to evaluate enrichment of HCFC1 (day 3), H3K4me3 (day 0 and 3), YY1,

DPY30, PolIISer5, and INTS11 (all day 0). Heatmaps were generated using deepTools v3.5.5 with

computeMatrix (reference-point mode, averageTypeBins mean, 10bp bins, ±3kb) followed by

plotHeatmap.

Overlap between HCFC1 (day 0 and 3) and YY1 (day 0) peaks was quantified using intervene

v0.6.5[53] venn (--type genomic --save-overlaps). Resulting BED files for each overlap category

were annotated with Homer annotatePeaks.pl. The proportion of peaks per annotation category

was calculated for each overlap group and visualized in stacked bar plots. To specifically evaluate

promoter-associated binding, peaks were filtered for those located in promoter regions, and their

overlap was visualized in a Venn diagram using matplotlib-venn v1.1.1. Overlap between YY1

and INTS11 binding on day 0 and 3 was assessed following the same approach.

Correlation between HCFC1 binding at day 0 and day 3 was further assessed using deepTools

multiBigwigSummary in BED-file mode, with a BED file containing all TSSs ±500 bp in mm10.

Counts (obtained with outRawCounts setting) were compared in a scatter plot using log1p-

transformed values. Pearson correlation was calculated using the plotCorrelation function with the

removeOutliers option.

Differential peak calling in H3K4me3 enrichment

To detect global changes in H3K4me3 levels, *Drosophila melanogaster* spike-ins were

incorporated into all H3K4me3 samples. The number of reads mapping better to dm6 than mm10

(see ChIP-seq analysis) were counted and used to scale CPM-normalized tracks. Initial

visualization of these tracks revealed large variability between replicates (data not shown).

Consequently, the spike-in normalization was excluded from the final analysis. Instead, all

H3K4me3 tracks were normalized using the ChIPseqSpikeInFree v1.2.4 method[54]. Composite

BAM files were processed to compute read coverage in sliding windows, which were normalized

to CPM and used to calculate scaling factors for each sample.

Differential peak calling between dTAG-13 and DMSO conditions was performed using DESeq2

v1.30.1[46]. Peaks identified by MACS3 in each condition were merged using BEDTools v2.31.1

merge[55], and the resulting BED file was converted to SAF format. Read counts per merged peak

were quantified with featureCounts v2.0.6[45] and provided to DESeq2. The estimated size factors

were adjusted using the scaling factors computed by ChIPseqSpikeInFree. A Wald test was

performed to compare dTAG-13 and DMSO conditions, and fold changes were shrunk using the

ashr v2.5-54 method[47]. Differential peaks were defined as those with an adjusted p-value < 0.05

and an absolute $\log_2$(fold change) > 1.

Tracks for each sample were scaled using the size factors computed by ChIPseqSpikeInFree.

Average tracks per condition were generated with deepTools bigwigAverage. Moreover, tracks

with $\log_2$(fold change) between dTAG-13 and DMSO were computed using deepTools

bigwigCompare. To show HCFC1- and YY1-associated changes in H3K4me3 enrichment, we

generated heatmaps with deepTools computeMatrix (reference-point mode, averageTypeBins

mean, 100bp bins, ±3kb) and plotHeatmap. Differential peaks with significantly reduced

H3K4me3 signal (as identified by DESeq2) were compared to unaffected peaks, which were

randomly downsampled to match the number of reduced peaks. Both log$_2$(fold change) tracks and
DMSO tracks were plotted for comparison.

Differential peak calling in HCFC1 and INTS11 binding

Changes in HCFC1 and INTS11 binding in YY1-FKBP cells after dTAG-13 treatment were
evaluated using MACS3 bdgdiff. The genome-wide number of unique and overlapping peaks were
visualized in Venn diagrams using matplotlib-venn. Peaks from each category were annotated
using the annotatePeaks.pl script from Homer v4.11 (Heinz et al. 2010), after which small
annotation categories were merged for clarity: 3' UTR and TTS were combined into "3' UTR -
TTS", and exon and non-coding into "exon". Motif enrichment analyses were conducted on each
peak category separately with Homer findMotifsGenome.pl (-size 200). Enrichment results for
known motifs were compared using scatter plots, where p-values for each motif were plotted
between two categories, and motifs were coloured by TF family.

Correlation between RNA-seq and ChIP-seq in HCFC1-FKBP cells

To correlate RNA-seq changes between HCFC1-FKBP dTAG-13 and DMSO treatments with
changes to H3K4me3 and INTS11 ChIP-seq signal under the same conditions, we first selected
DEGs from the RNA-seq analysis (adjusted p-value < 0.05, absolute log$_2$(fold change) > 1). To
allow comparison with unaffected expressed genes, we matched each gene in the downregulated
group to a gene in the non-significant group with a similar expression level. This was done using
the NearestNeighbors algorithm (n_neighbors=1) from scikit-learn v1.5.1 (Pedregosa et al., 2011),
applying the fit and kneighbors functions. To avoid duplicate matches, each selected nearest
neighbor was removed from the pool after selection. This resulted in a non-significant gene subset
of equal size and similar expression distribution as the downregulated group. Finally, we visualized
fold changes in H3K4me3 and INTS11 enrichment following dTAG-13 treatment, as well as
baseline enrichment of HCFC1, YY1, H3K4me3, and INTS11 at TSSs of upregulated,
downregulated, and matched unaffected genes.

We quantified H3K4me3 changes in the promoter region using deepTools multiBigwigSummary
in BED-file mode, with a bed file containing regions spanning TSS + 1kb, based on the enrichment
pattern observed in the ChIP-seq heatmap. The resulting mean signal change per promoter region
was visualized in violin plots for each gene category separately. Statistical significance between

groups was assessed using a two-sided Mann-Whitney U test, with p-values corrected for multiple

testing using the Benjamini-Hochberg method ($\alpha < 0.05$).

Correlation between YY1-FKBP dTAG-13 vs DMSO RNA-seq and INTS11 ChIP-seq

To assess the relationship between transcriptional changes upon YY1 degradation and alterations

in INTS11 binding, we compared RNA-seq data from YY1-FKBP dTAG-13 and DMSO-treated

samples with changes in the INTS11 ChIP-seq signal under the same conditions. INTS11 peaks

that were significantly reduced upon YY1 dTAG-13 treatment (see 'Changes in HCFC1 and

INTS11 binding') were intersected with YY1 ChIP-seq peaks using intervene venn (type genomic,

save-overlaps). Overlapping peaks were annotated with Homer annotatePeaks.pl, and genes with

shared peaks located within 1 kb of a transcription start site (TSS) were selected. The resulting

gene list was compared to the set of DEGs from the RNA-seq analysis. The number of up- and

downregulated genes among these shared targets was visualized in a bar plot. Statistical

significance was assessed using a Pearson's chi-squared (χ^2) goodness-of-fit test, comparing the

observed number of up- and downregulated genes to the expected distribution based on the overall

proportions of up- and downregulated genes.

Genome browser views

Genome browser views were created using IGV v2.18.2[56]. ChIP-seq signal tracks were generated

using the bdgcmp function in MACS3 (see ChIP-seq analysis). Log_2(fold change) tracks

comparing dTAG-13 and DMSO conditions were computed based on CPM-normalized coverage

for HCFC1 and INTS11 binding, or ChIPseqSpikeInFree-scaled CPM-normalized tracks for

H3K4me3 (see "Differential peak calling in H3K4me3 enrichment") using deepTools

bigwigCompare.

Mass Spectometry

Nuclear extracts preparation

All subsequent steps for nuclear extract preparation and FLAG-purification were performed at 4°C

using buffers supplemented with 1× protease inhibitors (Roche, 4693132001) and 15 µM MG-132

proteasome inhibitor (Sigma, C2211). Pre-chilled buffers and equipment were used throughout,

and low protein-binding Falcon and Eppendorf tubes were employed to minimize sample loss. One

day prior to harvest, two independently generated FLAG-V5-YY1 and 4 control mESC lines were

trypsinized, MEFs were removed by pre-plating for 45 min in non-gelatinized culture dishes. The

resulting mESCs were cultured without MEFs for 16h before processing. Cells were washed with cold PBS, collected by scraping on ice in PBS+ (1×PBS, 0.5 mM DTT), transferred to a Falcon tube and pelleted by centrifugation at 1500 rpm for 5 min. Pellets were resuspended in 2 pellet volumes of PBS+ and centrifuged at 1500 rpm for 5 min. The pellets were resuspended in 5 pellet volumes of buffer A (10 mM Hepes pH 7.6, 1.5 mM $MgCl_2$, 10 mM KCl, 0.5 mM DTT), incubated on ice for 10 min and centrifuged at 3000 rpm for 5 min. The supernatant was removed, and the pellets were resuspended in 2 pellet volumes of buffer A. Cells were lysed with 10 strokes of a Dounce homogenizer (type A pestle) on ice. The homogenate was transferred to Falcon tubes and centrifuged at 3000 rpm for 5 min. The supernatant, containing the cytoplasmic fraction, was discarded. The nuclear pellet was resuspended in 1.5 pellet volumes of Buffer C (20 mM Hepes pH 7.6, 20% glycerol, 420 mM NaCl, 1.5 mM $MgCl_2$, 0.2 mM EDTA, 0.5 mM DTT) and homogenized with 10 strokes of a Dounce homogenizer (type B pestle) on ice. The nuclear suspension was transferred to a Falcon tube, incubated while rotating for 30 min and centrifuged at 14000 rpm for 15min. The supernatant, containing the nuclear fraction, was collected and protein concentration was determined using a Nanodrop.

FLAG pull down

Protein concentrations of nuclear extracts were adjusted to the lowest concentration using buffer C. For each sample, 1 mL nuclear extract was diluted with 1 mL buffer C-0 (20 mM Hepes pH 7.6, 20% glycerol, 1.5 mM $MgCl_2$, 0.2 mM EDTA), to reduce the salt concentration to approximately 150 mM NaCl. Samples were incubated on ice for 15 min, centrifuged at 4000 rpm for 15 min, and the supernatant was transferred to a new tube. Benzonase nuclease (200 U; Novagen, 70664) was added to prevent RNA- or DNA-mediated interactions. Anti-Flag M2 Affinity Gel (Sigma-Aldrich, A2220) beads slurry (120 µl per sample) was washed four times with 1 ml Buffer C-150 (20 mM Hepes pH 7.6, 20% glycerol, 150 mM KCl, 1.5 mM $MgCl_2$, 0.2 mM EDTA, 0.02% NP40, 0.5 mM DTT) by centrifugation at 1000 rpm for 1 min. Nuclear extracts were added to the washed beads, resuspended, and incubated with gentle rotation for 3 h. Anti-FLAG-beads were pelleted by centrifugation at 2000 rpm for 2 min and washed five times with 1 mL buffer C-100 (20 mM Hepes pH 7.6, 10% glycerol, 100 mM KCl, 1.5 mM MgCl2, 0.2 mM EDTA, 0.02% NP40, 0.5 mM DTT). Each wash was performed with 5 min rotation (10 min for the final wash), followed by centrifugation at 2000 rpm for 2 min. The bound fraction was eluted by resuspending the Anti-FLAG-beads with 70 µL 0.2 mg/mL FLAG tripeptide peptide (Sigma,

F4799) in buffer 150 (without NP40), incubating for 15 min on ice, and centrifuging at 200

rpm for 2 min. The supernatant was collected as the eluted fraction, and the elution ste

repeated four times. An aliquot of the eluted fraction was analyzed by Western blot to confi

presence of FLAG 5-YY1 protein. Eluted fractions were combined and centrifuged at 2000 rp

for 2 min to remove residual FLAG beads. Protein eluates were concentrated by precipitation

with trichloroacetic acid (w/v) / 0.1% sodium deoxycholate (w/v). Pellets were resuspended

Laemmli sample buffer, boiled for 5 min at 95°C, separated SDS PAGE by 10% NuPAGE

Bis-Tris Gel (Invitrogen, NP0315BOX) and stained using a colloidal blue staining kit (Invitroge

LC6025).

Sample preparation

Complete lanes of SDS PAGE gels were excised and divided into smaller slices. Proteins were

reduced with 10 mM dithiothreitol (DTT) for 1 h at room temperature, alkylated with 55

chloroacetamide (CAM) for 1 h in the dark, and digested overnight at room temperature

sequencing grade trypsin (1:100, w/w; Roche). Peptides were desalted using C18 Sep Pak

cartridges (Waters) and eluted with 80% acetonitrile (AcN).

MS analysis

Peptides were analyzed by nanoflow MS/MS using an EASY LC system (Thermo) coupled

to a Fusion Tribrid Orbitrap mass spectrometer (Thermo) operating in positive ion mode

equipped with a nanospray source. Peptide mixtures were first trapped on a ReproSil C18 rev

S K D V H F R O X P Q   ' U O D L V F K * P E - k R X V H F P W D I O R P Z U D FW

then separated on a ReproSil C18 re S K D V H F R O X P Q   ' U O D L V F K * P E +

packed in house) using a linear gradient 80% to buffer B (0.1% FA, 80% v/v AcN) over 70 or

120 min at a constant flow rate of ~200 nL/min. The column effluent was directly sprayed i

ESI source of the mass spectrometer. Spectra were acquired in profile mode with an

resolution of 70,000 (AGC target: 3E6) across an m/z range of 350 to 1700. Data dependent

acquisition (Top15) was performed with HCD fragmentation, using an isolation window of

m/z and a normalized collision energy of 26–28%. MS2 spectra were recorded in the ion trap.

Singly charged precursors were excluded, dynamic exclusion was set to 20 s, and the inte

threshold was set to 8.0E3.

Data analysis

YY1

2.   Luchsinger-Morcelle, S. J., Gribnau, J. & Barentenbal, H. Orchestrating Asymmetric
Expression: Mechanisms behind Xist Regulation. *Epigenomes* **8**, 6 (2024).

3.   Schwämmle, T. & Schulz, E. G. Regulatory principles and mechanisms governing the
onset of random X-chromosome inactivation. *Curr. Opin. Genet. Dev.* **81**, 102063 (2023).

4.   Monkhorst, K., Jonkers, I., Rentmeester, E., Grosveld, F. & Gribnau, J. X inactivation
counting and choice is a stochastic process: evidence for involvement of an X-linked activator.
Cell **132**, 410–421 (2008).

5.   Jonkers, I. *et al.* RNF12 is an X-Encoded dose-dependent activator of X chromosome
inactivation. *Cell* **139**, 999–1011 (2009).

6.   Gontan, C. *et al.* RNF12 initiates X-chromosome inactivation by targeting REX1 for
degradation. *Nature* **485**, 386–390 (2012).

7.   Makhlouf, M. *et al.* A prominent and conserved role for YY1 in Xist transcriptional
activation. *Nat. Commun.* **5**, 4878 (2014).

8.   Barakat, T. S. *et al.* The trans-activator RNF12 and cis-acting elements effectuate X
chromosome inactivation independent of X-pairing. *Mol. Cell* **53**, 965–978 (2014).

9.   Shin, J. *et al.* Maternal Rnf12/RLIM is required for imprinted X-chromosome
inactivation in mice. *Nature* **467**, 977–981 (2010).

10.   Gontan, C. *et al.* REX1 is the critical target of RNF12 in imprinted X chromosome
inactivation in mice. *Nat. Commun.* **9**, 4752 (2018).

11.   Shin, J. *et al.* Rnf12/RLIM is dispensable for X-chromosome inactivation in the mouse
embryonic epiblast. *Nature* **511**, 86–89 (2014).

12.   Tian, D., Sun, S. & Lee, J. T. The long noncoding RNA, Jpx, is a molecular switch for X
chromosome inactivation. *Cell* **143**, 390–403 (2010).

13.   Chureau, C. *et al.* Ftx is a non-coding RNA which affects Xist expression and chromatin
structure within the X-inactivation center region. *Hum. Mol. Genet.* **20**, 705–718 (2011).

14.   Samanta, M. K. *et al.* Activation of Xist by an evolutionarily conserved function of
KDM5C demethylase. *Nat. Commun.* **13**, 2602 (2022).

15.   Ravid Lustig, L. *et al.* GATA transcription factors drive initial Xist upregulation after
fertilization through direct activation of long-range enhancers. *Nat. Cell Biol.* **25**, 1704–1715
(2023).

16.   Gribnau, J. & Grootegoed, J. A. Origin and evolution of X chromosome inactivation.
Curr. Opin. Cell Biol. **24**, 397–404 (2012).

17.   Disteche, C. M. Escapees on the X chromosome. *Proc. Natl. Acad. Sci. U. S. A.* **96**,
14180–14182 (1999).

18. Prada, C. F. & Laissue, P. A high resolution map of mammalian X chromosome fragile
regions assessed by large-scale comparative genomics. *Mamm. Genome Off. J. Int. Mamm.*
Genome Soc. **25**, 618–635 (2014).

19. Pacini, G. *et al.* Integrated analysis of Xist upregulation and X-chromosome inactivation
with single-cell and single-allele resolution. *Nat. Commun.* **12**, 3638 (2021).

20. Yu, H. *et al.* The ubiquitin carboxyl hydrolase BAP1 forms a ternary complex with YY1
and HCF-1 and is a critical regulator of gene expression. *Mol. Cell. Biol.* **30**, 5071–5085 (2010).

21. Capotosti, F. *et al.* O-GlcNAc transferase catalyzes site-specific proteolysis of HCF-1.
Cell **144**, 376–388 (2011).

22. Wilson, A. C., LaMarco, K., Peterson, M. G. & Herr, W. The VP16 accessory protein
HCF is a family of polypeptides processed from a large precursor protein. *Cell* **74**, 115–125
(1993).

23. Wysocka, J., Myers, M. P., Laherty, C. D., Eisenman, R. N. & Herr, W. Human Sin3
deacetylase and trithorax-related Set1/Ash2 histone H3-K4 methyltransferase are tethered
together selectively by the cell-proliferation factor HCF-1. *Genes Dev.* **17**, 896–911 (2003).

24. Shilatifard, A. The COMPASS family of histone H3K4 methylases: mechanisms of
regulation in development and disease pathogenesis. *Annu. Rev. Biochem.* **81**, 65–95 (2012).

25. Wang, H. *et al.* H3K4me3 regulates RNA polymerase II promoter-proximal pause-
release. *Nature* **615**, 339–348 (2023).

26. van Nuland, R. *et al.* Quantitative Dissection and Stoichiometry Determination of the
Human SET1/MLL Histone Methyltransferase Complexes. *Mol. Cell. Biol.* **33**, 2067–2077
(2013).

27. Bowness, J. S., Almeida, M., Nesterova, T. B. & Brockdorff, N. YY1 binding is a gene-
intrinsic barrier to Xist-mediated gene silencing. *EMBO Rep.* **25**, 2258–2277 (2024).

28. Yue, F. *et al.* A comparative encyclopedia of DNA elements in the mouse genome.
Nature **515**, 355–364 (2014).

29. Michaud, J. *et al.* HCFC1 is a common component of active human CpG-island
promoters and coincides with ZNF143, THAP11, YY1, and GABP transcription factor
occupancy. *Genome Res.* **23**, 907–916 (2013).

30. Cai, Y. *et al.* YY1 functions with INO80 to activate transcription. *Nat. Struct. Mol. Biol.*
14, 872–874 (2007).

31. Beckedorff, F. *et al.* The Human Integrator Complex Facilitates Transcriptional
Elongation by Endonucleolytic Cleavage of Nascent Transcripts. *Cell Rep.* **32**, 107917 (2020).

32. Mendoza-Figueroa, M. S., Tatomer, D. C. & Wilusz, J. E. The Integrator Complex in
Transcription and Development. *Trends Biochem. Sci.* **45**, 923–934 (2020).

33.   Weintraub, A. S. *et al.* YY1 Is a Structural Regulator of Enhancer-Promoter Loops. *Cell*
171, 1573-1588.e28 (2017).

34.   Minocha, S. *et al.* Epiblast-specific loss of HCF-1 leads to failure in anterior-posterior
axis specification. *Dev. Biol.* **418**, 75–88 (2016).

35.   Cenik, B. K. & Shilatifard, A. COMPASS and SWI/SNF complexes in development and
disease. *Nat. Rev. Genet.* **22**, 38–58 (2021).

36.   Mendjan, S. *et al.* Nuclear pore components are involved in the transcriptional regulation
of dosage compensation in Drosophila. *Mol. Cell* **21**, 811–823 (2006).

37.   Cai, Y. *et al.* Subunit composition and substrate specificity of a MOF-containing histone
acetyltransferase distinct from the male-specific lethal (MSL) complex. *J. Biol. Chem.* **285**,
4268–4272 (2010).

38.   Loda, A. *et al.* Genetic and epigenetic features direct differential efficiency of Xist-
mediated silencing at X-chromosomal and autosomal locations. *Nat. Commun.* **8**, 690 (2017).

39.   Robert-Finestra, T. *et al.* SPEN is required for Xist upregulation during initiation of X
chromosome inactivation. *Nat. Commun.* **12**, 7000 (2021).

40.   Krueger, F. & Andrews, S. R. SNPsplit: Allele-specific splitting of alignments between
genomes with known SNP genotypes. *F1000Research* **5**, 1479 (2016).

41.   Sherry, S. T. *et al.* dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* **29**,
308–311 (2001).

42.   Krueger, F., James, F., Ewels, P., Afyounian, E. & Schuster-Boeckler, B.
FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. Zenodo
https://doi.org/10.5281/zenodo.5127899 (2021).

43.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915
(2019).

44.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma. Oxf. Engl.*
25, 2078–2079 (2009).

45.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinforma. Oxf. Engl.* **30**, 923–930 (2014).

46.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

47.   Stephens, M. False discovery rates: a new deal. *Biostat. Oxf. Engl.* **18**, 275–294 (2017).

48.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–359 (2012).

49.   Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic Acids Res.* **42**, W187-191 (2014).

50.   Jonkers, I. *et al.* Xist RNA is confined to the nuclear territory of the silenced X
chromosome throughout the cell cycle. *Mol. Cell. Biol.* **28**, 5583–5594 (2008).

51.   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137
(2008).

52.   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589
(2010).

53.   Khan, A. & Mathelier, A. Intervene: a tool for intersection and visualization of multiple
gene or genomic region sets. *BMC Bioinformatics* **18**, 287 (2017).

54.   Jin, H. *et al.* ChIPseqSpikeInFree: a ChIP-seq normalization approach to reveal global
changes in histone modifications without spike-in. *Bioinforma. Oxf. Engl.* **36**, 1270–1272 (2020).

55.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinforma. Oxf. Engl.* **26**, 841–842 (2010).

56.   Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. *Brief. Bioinform.* **14**,
178–192 (2013).

57.   Tyanova, S., Temu, T. & Cox, J. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. *Nat. Protoc.* **11**, 2301–2319 (2016).

Acknowledgements

We thank Mariana Pelicano de Almeida and Debbie van den Berg for the INTS11 ChIP protocol

and feedback. We also thank Stanley van Herk for access to the Bioruptor Pico. Finally, we thank

Danilo Remmers and Koert van den Bergh for technical assistance.

Funding

This study was supported by a ZonMW-TOP subsidy (nr 91215046) to JG.

Author Information

Author contribution

J.B., B.T., S.M., C.G., H.M.-B. and J.G. conceptualised and designed experiments. J.B, S.M., E.R., R.v.H., S.J.L.-M., C.M., R.R., B.G., C.G. and H.M.-B. performed experiments and analysed data. B.T. performed all bioinformatic analyses. J.D. and W.v.I performed the mass spectrometry and sequencing, respectively. J.B., B.T., C.G., H.M.-B. and J.G wrote the manuscript and all authors edited it. J.B., S.M., C.M. and B.G. generated all cell lines. C.G. performed the pulldowns. J.B., S.M. and R.v.H. genotyped cell lines. J.B., S.J.L.-M. and H.M.-B. performed ChIP-seq experiments. J.B., E.R. and R.v.H. performed WB and RNA-FISH experiments. C.G., H.M.-B. and J.G. jointly supervised the project.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information

The online version contains supplementary material available at XXX.

Materials and correspondence

All requests for materials and related questions should be addressed to Hegias Mira-Bontenbal and Joost Gribnau.

Figure 1

    Legends

Figure 1: Large mega-domain deletions on the X chromosome identify HCFC1 as a potential XCI
activator.

a, Targeting and differentiation strategies. *Rnf12*[+/-] cells were transiently transfected with

CRISRPR-Cas9 and gRNAs targeting different locations on the X chromosomes. Correctly

targeted clones were differentiated for several days. **b**, X chromosome scheme depicting

evolutionary strata 1, 2, 3, 4 and the pseudo-autosomal region (PAR)[17,18]. The locations of *Xist* and

Rnf12 are depicted with a black and grey line, respectively, while the deleted *Rnf12* allele on the

129 chromosome is also shown. The different deletions (ΔA to ΔK) are shown with different colors

underneath. **c**, Relative *Xist* expression at day 8 of differentiation of WT, *Rnf12*[+/-] and the different

Rnf12[+/-] cell lines with X chromosomal deletions, normalized to *Rnf12*[+/-]. Average ± SD, n=2 to 5

biological replicates. A one-sided Welch's t-test was used to test significant differences between

Rnf12[+/-] and WT or deletion cell lines. P-values were corrected for multiple testing with the

Benjamini-Hochberg method. **d**, Zoom-in on a specific region of ΔE. The locations of different

overlapping deletions (ΔE1, ΔE2, ΔE3) are shown with black lines; the smaller incremental

deletions (ΔE2A, ΔE2B, ΔE3A, ΔE3B) are shown in green. The blue rectangle highlights the

location of *Hcfc1* and *Irak1*. **e**, Expression heatmap of X-linked genes, sorted by genomic location

and split by allele (129 or Cast) for 2 WT, 2 *Rnf12*[+/-], 1 ΔE2A, 2 ΔE2B, 2 ΔE3A, 2 ΔE3B and 2

ΔE clones. Notice the deleted genes per clone on the 129 alleles. **f**, Relative *Xist* expression at day

0 and day 8 of WT, *Rnf12*[+/-], ΔE and smaller ΔE deletion cell lines. Average expression ± SD; n=2

(WT, *Rnf12*[+/-] and ΔE2A lines), n=4 (ΔE2B, ΔE3A and ΔE3B lines) and n=5 (ΔE lines) biological

replicates. A one-sided Welch's t-test was used to test significant differences per day of

differentiation between *Rnf12*[+/-] and WT or deletions cell lines. P-values were corrected for

multiple testing with the Benjamini-Hochberg method. **g**, Volcano plot portraying gene expression

changes between ΔE2B and *Rnf12*[+/-] cells. Downregulated and upregulated genes in ΔE2B are

shown in blue and red, respectively. Notice *Xist* is depicted on the left, as the most significant

downregulated gene. **h**, Same as **g** but for ΔE3B vs *Rnf12*[+/-] cells. **i**, Expression (TPM) of *Hcfc1*

(green), *Irak1* (grey), *Rnf12* (blue) and *Xist* (red) in ΔXic_Cast ESCs from Pacini et al., 2021[19]. P-

values were calculated between day 0 and 1 for *Hcfc1* (green) and *Irak1* (grey) using one-sided

independent t-tests and corrected for multiple testing with the Benjamini-Hochberg method.

Figure 2

 Figure 2: HCFC1 is an XCI activator.

a, HCFC1 rescue and differentiation strategy of control (Ctrl; parental Rnf12$^{+/-}$), ΔE2B and ΔE3B

ESCs. **b**, RT-qPCR analysis of *Hcfc1* expression upon *Hcfc1* rescue in pools of transfected cells

after 4 days of differentiation of in Ctrl (parental *Rnf12$^{+/-}$*), ΔE2B and ΔE3B. The average of n=2

biological replicates is shown. **c**, RT-qPCR analysis of *Xist* expression in pools of transfected cells

after 4 days of differentiation. The average of n=2 biological replicates is shown. **d**, 2xFLAG-V5-

FKBP12^{F36V} tagging strategy of HCFC1 and differentiation schemes with different dTAG-13

treatment windows. **e**, Western blot of the C-terminal fragments of HCFC1 upon dTAG-13

treatment in undifferentiated ESCs. **f**, RT-qPCR analysis of *Xist* expression at different timepoints

after start of dTAG-13 treatment and differentiation normalised to DMSO control; see Fig. 2a. A

two-sided Welch's t-test was used to assess significant differences between DMSO- and dTAG-13

treated cells. P-values were corrected for multiple testing using the Benjamini-Hochberg method.

g, *Xist* RNA-FISH analysis in HCFC1-FKBP ESCs differentiated for 3 days and treated with

dTAG-13 for the last 48h vs DMSO and parental ESCs. **h**, Quantification of **g**. Average ± SD,

n=3-6 biological replicates, n=157-236. A two-sided Welch's t-test was used to assess significant

differences between DMSO- and dTAG-13 treated cells. P-values were corrected for multiple

testing using the Benjamini-Hochberg method. **i**, Gene expression levels of several genes located

in the XIC determined by RNA-seq analysis of HCFC1-FKBP ESCs differentiated for 3 days and

treated with dTAG-13 for the last 48h. Shown in green are genes residing in the *Xist* TAD and

involved in its activation. In red, genes located in the *Tsix* TAD and involved in *Xist* repression.

Average TPMs normalized to 48h DMSO control ± SD, n= 3 biological replicates. P-values were

calculated using DESeq2, which corrects for multiple testing using the Benjamini-Hochberg

method. **j**, Violin plot of allelic ratios of X-linked gene expression in the DMSO and dTAG-13

conditions. The box plots display the median (black line), the interquartile range (box limits) and

1.5x of the interquartile range (whiskers). Dashed line indicates ratio of 0.5 as expected ratio

without silencing. P-value was calculated using a two-sided Mann-Whitney U test.

1049

Figure 3

 Figure 3: HCFC1 binding correlates with promoters, H3K4me3 deposition and YY1 binding.

a, Heatmap of HCFC1, YY1, H3K4me3, DPY30, PolIISer5 and INTS11 enrichment at HCFC1

peaks in undifferentiated ESCs (left). The right panels show HCFC1 and H3K4me3 enrichment at

day 3 of differentiation at HCFC1 peaks identified at day 0. **b**, HCFC1-only and YY1-only and

shared HCFC1-YY1 peak distribution at different genomic locations at day 0. **c**, Venn diagram

with the overlap of HCFC1 and YY1 peaks in promoter regions at day 0 and 3 of differentiation.

d, Correlation between HCFC1 binding at day 0 and day 3 in promoter regions, with r representing

the Pearson correlation coefficient. **e**, Genome browser views showing examples of HCFC1-YY1

overlapping peaks in promoter regions. **f**, MA plot depicting H3K4me3 enrichment changes upon

dTAG-13-mediated removal of HCFC1. In red, peaks with increased H3K4me3 enrichment; in

grey, unaffected peaks and in blue, peaks with reduced H3K4me3 enrichment. **g**, Violin plots

showing H3K4me3 changes in promoter regions of differentially expressed genes from the RNA-

seq comparison between HCFC1-FKBP dTAG-13 and DMSO. P-values were calculated using

two-sided Mann-Whitney U tests, corrected for multiple testing using the Benjamini-Hochberg

approach. **h**, Heatmaps showing differences in H3K4me3 and INTS11 enrichment at H3K4me3

peaks that are lost upon dTAG-13-mediated removal of HCFC1 and unaffected H3K4me3 peaks.

For both H3K4me3 and INTS11, the fold change between dTAG-13 and DMSO treatment is

shown. HCFC1 binding (day 3) and YY1 binding (day 0) are displayed on the right. For the

unaffected peaks, a random subset of 4,991 peaks is shown.

1069

Figure 4

Figure 4: YY1 interacts with the Integrator complex.

a, YY1 interactome identified by FLAG pull-down and mass spectrometry. Volcano plot showing proteins enriched in YY1 pull-downs compared to control. Known members of the Integrator (light blue), INO80 (dark blue), and COMPASS (green) complexes are highlighted. Selected proteins, including YY1 itself and HCFC1, are labeled in black. Dashed lines denote thresholds for significance ($p < 0.05$, $\log_2$ fold change > 1). Data are based on two biological replicates for FLAG-YY1 and four biological replicates for the control. **b**, Percentages of shared YY1-INTS11, YY1-only or INTS11-only peaks at different genomic locations at day 0. **c**, Venn diagram showing the overlap of YY1 and INTS11 binding in promoter regions. **d**, Generation strategy of the FKBP12^{F36V}-HA-Yy1:2xFLAG-V5-Hcfc1 compound knockin ESC lines by homologous recombination. **e**, Stacked barplot showing percentages of H3K4me3, HCFC1 and INTS11 peaks that significantly lost or gained enrichment or remained unchanged, upon HCFC1 or YY1 loss. **f,** MA plot depicting H3K4me3 enrichment changes upon dTAG-13-mediated removal of YY1. In red, peaks with increased H3K4me3 enrichment; in grey, unaffected peaks and in blue, peaks with reduced H3K4me3 enrichment. **g**, Scatter plot comparing motif enrichment between gained and lost HCFC1 peaks shown in **e**. Each point represents a motif, plotted by its -$\log_{10}$(p-value) in both analyses, and motifs with the lowest p-values in either condition are labeled. **h**, Barplot showing the number of up- and downregulated genes with reduced HCFC1 binding in the promoter region upon YY1 depletion. P-value was calculated using a Pearson's chi-squared (χ^2) goodness-of-fit test, based on the overall proportion of up- and downregulated genes. **i**, Same as **h**, but for promoter regions with reduced INTS11 binding. **j**, Heatmap showing changes in INTS11 and H3K4me3 enrichment upon dTAG-13-mediated depletion of YY1 or HCFC1, alongside YY1 and HCFC1 binding at INTS11 peaks. Peaks are grouped based on whether INTS11 enrichment is reduced (top) or unaffected (bottom) by YY1 depletion.

Figure 5

a

b

48h DMSO   48h YY1 dTAG-13

fold change over DMSO

p = 1.12·10⁻³ p = 8.70·10⁻⁵ p = 4.56·10⁻⁴ p = 0.0173 p = 1.12·10⁻³ ns

Tsix Xist Jpx Ftx Hcfc1 Rex1

day 3 differentiation

c

X-linked activity

autosomal activity

Tsix

COMPASS
HCFC1

Xist

H3K4me3

RNF12 Jpx

Ftx

YY1
Integrator

YY1 REX1

YY1
Integrator

YY1
Integrator

d

probability

WT

Rnf12⁺/⁻ or Hcfc1⁺/⁻

Rnf12⁺/⁻ + Hcfc1⁺/⁻

activation level

threshold

♀

♂

ES differentiation →

Figure 5: HCFC1 binds to and regulates the expression of several non-coding genes in the XIC.

a, Genome browser views showing ChIP-seq tracks of H3K4me3 in HCFC1-depleted or YY1-

depleted versus DMSO controls at day 3 of differentiation, HCFC1 ChIP-seq in YY1-depleted

versus its DMSO control at day 3 of differentiation, INTS11 ChIP-seq in YY1-depleted versus its

DMSO control at day 3, and YY1 and HCFC1 ChIP-seq in female ESCs at day 0 around the *Tsix*,

Xist, *Jpx* and *Ftx* promoter regions. **b,** Relative expression of *Tsix, Xist*, *Jpx, Ftx, Hcfc1* and

pluripotency factor *Rex1* at day 3 of differentiation of YY1-FKBP ESC cells normalised to DMSO

control. A two-sided Welch's t-test was used to test significant differences between DMSO- and

dTAG-13 treated cells. P-values were corrected for multiple testing using the Benjamini-Hochberg

method. **c,** Gene-regulatory network controlling *Xist* expression. RNF12 promotes YY1 binding

by targeting the *Xist* repressor REX1 for proteasomal degradation. HCFC1/COMPASS and YY1

are both required for H3K4me3 catalysis at the *Xist* regulatory element. YY1 also recruits

Integrator to activate transcription of *Xist* cis-regulatory genes within the XIC. **d**, Dose-dependent

activation of *Xist* expression. Elevated levels of X-linked activators in XX cells raise the

probability for *Xist* transcription initiation above threshold; this probability is reduced in $Rnf12^{+/-}$

or $Hcfc1^{+/-}$ cells and further diminished in double-heterozygous $Rnf12^{+/-};Hcfc1^{+/-}$ cells.

1110

Extended Data Figure 1

Extended Data Figure 1: Megadomain deletion confirmation by PCR and WGS.

a, Overview of the locations of the different deletions generated in this study. Different X-linked

length polymorphisms between 129 and Cast are shown (DX amplicons). **b-c,** PCR analysis of

different DX amplicons for the different deletions. **d**, Location of the different DX amplicons with

respect to the deleted regions, and length polymorphism analysis inside and around several

incremental deletions within region E. **e**, Scatter plots of allelic ratios along the X-chromosome

from shallow whole-genome sequencing for different ΔE, ΔE2A, ΔE2B, ΔE3A and ΔE3B clones

(left). Y-axis represents the $\log_2$(fold change) between the 129 and Cast alleles for each window.

Windows with significant signal differences between alleles are highlighted in green. The zoomed-

in plots on the right show the region around 67-77 Mb, with the *Hcfc1*-containing window

indicated by a star.

1122

1123

Extended Data Figure 2

Extended Data Figure 2: Deletions containing *Hcfc1* show reduced Xist upregulation but can differentiate.

a, DNA FISH analysis on metaphase spreads of the parental and three ΔE clones detecting *Hprt* and *Xist* in green, and *Mecp2* in red. Notice red is inside the deleted region. White arrowheads indicate chromosome X. Arrowheads with an asterisk in ΔE clone 3 indicate the signal within an interphase nucleus. **b**, Representative images of *Xist* RNA FISH (red) in WT, *Rnf12*[+/-] and three ΔE clones at day 8 of differentiation. **c**, Quantification of **b**, average number of cells with a *Xist* cloud ± SD, n=3 biological replicas, n=201-231, statistical significance was calculated using a two-sided Welch T-test and corrected for multiple testing using the Benjamini-Hochberg procedure. P-values are indicated, ns = not significant. **d**, Allelic gDNA PCR of *Tsix* in WT, *Rnf12*[+/-] and 3 independent ΔE clones on day 8 of differentiation to confirm the presence of two X chromosomes. Average ± SD, n=1-3 biological replicates. **e**, Relative *Xist* expression in WT, *Rnf12*[+/-], ΔE, ΔE1, ΔE2 and ΔE3 ESCs at day 0 and 8 of differentiation. A one-sided Welch's t-test was used to test significant differences between *Rnf12*[+/-] and WT or deletion cell lines. P-values were corrected for multiple testing with the Benjamini-Hochberg method. **f**, Heatmap of gene expression (in TPMs) for pluripotency markers (*Pou5f1*, *Nanog*, *Zfp42/Rex1*, *Klf4*) and differentiation markers (*Kdr/Flk-1*, *Gata4*, *Gata6*, *T* and *Nodal*) across WT, *Rnf12*[+/-] and ΔE, ΔE2A, ΔE2B, ΔE3A and ΔE3B clones. **g**, Violin plot of allelic ratios of X-linked gene expression in *Rnf12*[+/-], ΔE, ΔE2B and ΔE3B. The box plots display the median (black line), the interquartile range (box limits) and 1.5x of the interquartile range (whiskers). Dashed line indicates ratio of 0.5 as expected ratio without silencing. P-values were calculated using a two-sided Mann-Whitney U test and corrected for multiple testing using the Benjamini-Hochberg approach. **h**, Expression (TPM) of *Hcfc1* (green), *Irak1* (grey), *Rnf12* (blue) and *Xist* (red) in ΔXic_B6 ESCs from Pacini et al., 2021[19]. P-values were calculated between day 0 and 1 for Hcfc1 (green) and Irak1 (grey) using one-sided independent t-tests and corrected for multiple testing with the Benjamini-Hochberg method.

Extended Data Figure 3

Extended Data Figure 3: HCFC1 rescue and conditional knockout confirmation.

a, Relative *Hcfc1* expression in control (*Rnf12*[+/-]), ΔE2B and ΔE3B ESCs upon stable integration,

Hcfc1 expression vs control plasmid in different individual undifferentiated clones; Average ± SD,

n=4-5 independent clones. **b**, Relative *Xist* expression in Control (*Rnf12*[+/-]), ΔE2B and ΔE3B upon

stable integration, *Hcfc1* expression vs control plasmid at day 4 of differentiation in different

individual clones; Average ± SD, n=4-5 independent clones. **c**, Same as in **b**, but for differentiation

marker *Nestin*. **d**, PCR analysis of HCFC1-FKBP ESCs, top. DX95 PCR to control for the presence

of both X chromosomes, bottom. **e**, HCFC1 C-term and V5 WB in HCFC1-FKBP ESCs after

treatment with dTAG-13 or DMSO control in the ESC state. Actin is used as loading control. **f**,

Expression of several pluripotency markers at day 3 of differentiation after 48h of dTAG-13 or

DMSO treatment. TPM values are normalized to the average expression in the DMSO condition

for each gene. P-values were calculated using DESeq2, which corrects for multiple testing using

the Benjamini-Hochberg method **g**, Same as in **f**, but for genes involved in *Xist* regulation. **h**,

Schematic of HCFC1-FKBP ESCs with a doxycycline-inducible *Xist* promoter and strategy to

induce *Xist* expression in the ESC stage in the presence of dTAG-13. **i**, RT-qPCR analysis of *Xist*

expression in HCFC1-FKBP ESCs upon 24h of dox treatment and/or 30h dTAG-13 treatment.

Average ± SD, n=3 biological replicates. **j**, Allelic expression of X-linked gene *Rnf12* in HCFC1-

FKBP ESCs upon 24h Dox and/or 30h dTAG-13 treatment. Average ± SD, n=3 biological

replicates. **k**, Allelic expression of XCI escapee *Kdm6a* upon forced *Xist* expression (+Dox) for

24h in the presence of dTAG-13 or DMSO. For (i–k), p-values from two-sided Welch's t-test

between DMSO and dTAG-13 treatment, corrected using the Benjamini–Hochberg procedure.

1173

1174

Extended Data Figure 4

 Extended Data Figure 4: HCFC1 binding sites analysis and genome-wide H3K4me3 and INTS11
recruitment upon dTAG-13-mediated depletion of HCFC1.

a, Genomic annotation of HCFC1 peaks at day 0 and day 3 of ESC differentiation. **b**, PCR analysis

confirming proper integration of HCFC1-V5- and the YY-FKBP-tags and presence of both X

chromosomes (DX95). **c**, Feature enrichment analysis at HCFC1 peaks at day 3 of differentiation.

Features are grouped by over- and underrepresentation, and the top 10 with the lowest p-values

are labeled. **d**, Motif enrichment analysis at HCFC1 peaks at day 3 of differentiation. The 10 most

significantly enriched motifs are labeled. **e**, Western blot analysis of H3K4me3 in HCFC1-FKBP

cells treated with DMSO or dTAG-13 (48h) at day 3 of differentiation. **f**, Quantification of global

H3K4me3 WB signal in Ctrl and HCFC1-FKBP cells treated with DMSO or dTAG-13 (48h) at

day 3 of differentiation. ns, not significant, p-values were calculated using a two-sided Welch's t-

test and corrected *via* Benjamini-Hochberg. **g**, Heatmap of H3K4me3 and INTS11 enrichment at

H3K4me3 peaks lost upon HCFC1 depletion, and at unaffected peaks (random subset of 4,991

peaks). **h**, MA plot showing gene expression changes upon dTAG-13-mediated depletion of

HCFC1. Genes are colored based on significant upregulation (red), downregulation (blue) or no

change (grey). **j**, Heatmap of H3K4me3 and INTS11 changes upon HCFC1 depletion, as well as

HCFC1 and YY1 binding and H3K4me3 and INTS11 enrichment in the DMSO condition, at

promoters of upregulated, unaffected and downregulated genes.

Extended Data Figure 5

Extended Data Figure 5: YY1 and INTS11 binding sites analysis.

a, Generation strategy of the 2xFLAG-V5-Yy1 ESC lines by random integration in XX *Tsix*-STOP

ESCs. **b**, Western blotting analysis of YY1 expression in 2xFLAG-V5-YY1 ESCs using an

endogenous YY1 or V5 antibody, and Actin as loading control. **c**, Venn diagram showing the

genome-wide overlap of YY1 and INTS11 peaks at day 0 of differentiation. **d**, Scatter plot

comparing motif enrichment between YY1-INTS11 overlapping peaks and INTS11-unique peaks

identified in **c**. Each point represents a motif, plotted by its -$\log_{10}$(p-value) in both analyses, and

motifs with the lowest p-values in either condition are labeled. **e**, WB analysis of YY1 degradation

upon dTAG-13 treatment vs DMSO for different hours of YY1-FKBP ESCs at day 0 of

differentiation. **f**, Scatter plot comparing motif enrichment between reduced and unaffected

INTS11 peaks in YY1-FKBP ESCs in the dTAG-13 condition at day 3 of differentiation. Each

point represents a motif, plotted by its -$\log_{10}$(p-value) in both analyses, and motifs with the lowest

p-values in either condition are labeled. **g**, Volcano plot showing gene expression changes upon

YY1 depletion. Genes are colored based on significant upregulation (red), downregulation (blue)

or no change (grey). **h**, Expression of *Cnnm2* and *Zfp37* upon 24h YY1 removal in undifferentiated

ESCs[33], average of n=2 biological replicates. TPM values are normalized to the average expression

in the DMSO condition for each gene. P-values were calculated using Wald tests and corrected for

multiple testing using the Benjamini-Hochberg method, as implemented by DESeq2. **i**, *Cnnm2* and

Zfp37 expression analysis by qRT-PCR in KFBP-YY1 ESCs upon 48h DMSO or dTAG-13

treatment at day 3 of differentiation. A two-sided Welch's t-test was used to assess significant

differences between DMSO- and dTAG-13 treated cells. P-values were corrected for multiple

testing using the Benjamini-Hochberg method. **j**, Views of ChIP-seq tracks for H3K4me3 (green)

and INTS11 (blue) in YY1-depleted ESCs versus its DMSO control at day 3 of differentiation,

and YY1 ChIP-seq track in female ESCs at day 0 (black), around the *Cnnm2* promoter region. **k**,

Same as **j**, for *Zfp37*. **l**, WB analysis of YY1 and H3K4me3 of YY1-FKBP ESCs differentiated

for 3 days and treated with dTAG-13 or DMSO for the last 48h. **m**, Quantification of global

H3K4me3 WB signal in Ctrl and YY1-FKBP cells treated with DMSO or dTAG-13 (48h) at day

3 of differentiation. ns, not significant. P-values were calculated using a two-sided Welch's t-test

and corrected *via* Benjamini-Hochberg.

1222

1223