METABOLIC AND TRANSCRIPTOMIC ANALYSES IDENTIFY COORDINATED RESOURCE REALLOCATION IN RESPONSE TO PHOSPHATE SUPPLY IN HEMP

Benjamin Wee Y.[1,2], Oliver Berkowitz[1,2], Sophia Ng[1,2], Amelia Pegg[1,2], James Whelan[1,3], Ricarda Jost[1,2*]

[1]ARC Research Hub for Medicinal Agriculture, Department of Animal, Plant and Soil Sciences, School of Agriculture, Biomedicine and Environment, La Trobe University, Bundoora VIC 3086, Australia
[2]La Trobe Institute for Sustainable Agriculture & Food, La Trobe University, Bundoora VIC 3086, Australia
[3]College of Life Science, Zhejiang University, Hangzhou, Zhejiang, 310058, P.R. China

Author for correspondence:
Ricarda Jost: r.jost@latrobe.edu.au

Email addresses of authors:
Benjamin Wee Y.: 20826392@students.latrobe.edu.au
Oliver Berkowitz: o.berkowitz@latrobe.edu.au
Sophia Ng: jinqtyan@me.com
Amelia Pegg: a.pegg@latrobe.edu.au
James Whelan: jimwhelan@zju.edu.cn
Ricarda Jost: r.jost@latrobe.edu.au

Number of figures: 6
Number of tables: 0
Word count: 7424
Supplementary data: 2 figures and 7 tables

1 **HIGHLIGHT (30 WORDS)**

2   Contrasting human selection in *Cannabis sativa* enables the analysis of
3 divergent nutrient-use strategies. Our integrated transcriptomic and physiological
4 data revealed adaptive resource allocation, informing the development of nutrient-
5 efficient cultivars.

6 ABSTRACT

Cannabis sativa L., one of the earliest domesticated crops, was selectively bred into drug-type (high phytocannabinoid yield) and hemp-type (fibre and seed production) varieties, resulting in divergent traits including nutrient requirements. To address the limited molecular understanding of nutrient use regulation in Cannabis, this study investigated the response of dual-purpose hemp to varying phosphate availability. Physiological and transcriptomic profiling across source and sink organs revealed that biomass partitioning to flowers was sustained unless phosphate was completely withdrawn. Analyses of key phosphate-responsive genes highlighted the importance of P reallocation to flowers via glycolytic bypass reactions and release of phosphate from organic pools in source organs. Phospholipid biosynthesis and other P-dependent metabolic pathways remained active in sink organs. While the transcriptional response of key regulators, such as SPX DOMAIN GENE (SPX), PHOSPHATE STARVATION RESPONSE1 (PHR1) and NITRATE-INDUCIBLE, GARP-TYPE TRANSCRIPTIONAL REPRESSOR1 (NIGT1) followed canonical patterns, our data suggest a lack of suppression of phosphate uptake and root-to-shoot translocation at higher phosphate supply levels. Subsequent depletion of nitrate pools restricts further growth. These findings uncover conserved and unique aspects of nutrient signalling in hemp, shedding light on its adaptation to nutrient-poor soils through distinct source–sink regulation.

26 KEYWORDS

Cannabis sativa, hemp, female flower, nitrogen, phosphorus, sink, source, nutrient use, transcriptomics.

INTRODUCTION

Phosphorus (P) is a critical nutrient for plant growth, influencing a range of physiological processes from gene regulation to energy transfer and flowering (Cho *et al.*, 2025; Ojeda-Rivera *et al.*, 2022). It is exclusively taken up by plant roots in the form of inorganic phosphate (Pi, $H_2PO_4^-$). Despite its importance for metabolism and plant development, research on the molecular regulation of Pi acquisition and use in Cannabis is limited. Understanding the impact of Pi on Cannabis reproductive growth or specialised metabolism is essential to improve biomass and product yield (Lambers, 2021; Xie *et al.*, 2023). P deficiency results in reduced growth, accelerated senescence, and lower cannabinoid content, while excess Pi supply does not enhance biomass production or specialised metabolite accumulation (Cockson *et al.*, 2020; Hershkowitz *et al.*, 2025; Shiponi and Bernstein, 2021b). Notably, cannabinoid concentration in tissues is negatively correlated with Pi supply, while total cannabinoid yield per plant increases with Pi supply due to higher flower biomass (Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b; Wee Y. *et al.*, 2024).

Despite its impact on overall product yield, integration of knowledge gained from physiological responses with available molecular data to characterize the phosphate starvation response (PSR) in Cannabis is limited (Jost *et al.*, 2025; Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b; Wee Y. *et al.*, 2024). The PSR describes a highly complex and conserved regulatory system in plants that controls the coordinated acclimation to low Pi availability through changes in nutrient uptake, root system architecture, and shoot metabolism which has been studied extensively in model plants such as *Arabidopsis*, rice and tomato (Paz-Ares *et al.*, 2022; Péret *et al.*, 2014; Plaxton and Tran, 2011; Yang *et al.*, 2024). MYB transcription factors PHOSPHATE STARVATION RESPONSE1 (PHR1) and PHR1-like orthologs play a key role in PSR, controlling the expression of hundreds of genes containing the P1BS (GnATATnC) cis-regulatory motif in their promoter and enhancer regions (Bustos *et al.*, 2010; Guo *et al.*, 2015; Jiang *et al.*, 2019; Linn *et al.*, 2017; Rubio *et al.*, 2001; Safi *et al.*, 2017). Target genes are associated with processes such as Pi acquisition, e.g., members of the high affinity *PHOSPHATE TRANSPORTER1* (*PHT1*) family, Pi scavenging, e.g., purple acid phosphatases (*PAP*s), and ribonucleases (RNS), assimilation of other nutrients such as iron,

sulfate and nitrate as well as membrane lipid remodelling. Release of Pi by phospholipid hydrolysis requires a shift to alternative lipids such as sulfo- and galactolipids. Their synthesis is catalysed by UDP-sulfoquinovose synthases (SQD1 and SQD2), or type B monogalactosyldiacylglycerol (MGD2 and MGD3) and digalactosyldiacylglycerol (DGD2) synthases, respectively (Kelly and Dormann, 2002; Kobayashi *et al.*, 2009; Okazaki *et al.*, 2013; Tjellstrom *et al.*, 2008; Yu *et al.*, 2002). PHR1 itself is regulated by SYG1/Pho81/XPR1 (SPX) domain proteins, which bind higher-order inositol pyrophosphates (IPs) to signal cellular phosphorus status (Collins *et al.*, 2024). Other notable transcription factors include WRKY6 and WRKY42, which negatively regulate expression of transcripts encoding Pi exporter and mediator of Pi root-to-shoot translocation, PHOSPHATE1 (PHO1), via binding to the W-box motif in the *PHO1* promoter (Chen *et al.*, 2009; Su *et al.*, 2015). At the protein level, PHO1 is negatively regulated by ubiquitin-conjugating E2 enzyme, PHOSPHATE 2 (PHO2) and E3 ligase PRU1 (Liu *et al.*, 2012; Ye *et al.*, 2018).

Deciphering the interplay between these molecular determinants of PSR and associated physiological responses is essential for advancing development of crop cultivars with high P use efficiency (Gu *et al.*, 2016; Heuer *et al.*, 2017). Given drug-type Cannabis is a high-value medicinal plant largely grown in controlled environments, there has been little effort to improve its P use. This is reflected by differences in nutrient response between usage-types, with drug-type Cannabis exhibiting a tolerance for higher nutrient supplies without yield penalties as opposed to hemp-types, which are generally more nutrient efficient due to being traditionally grown on marginal lands (Scalabrin *et al.*, 2024; Struik *et al.*, 2000; Tang *et al.*, 2018; Wee Y. *et al.*, 2024). These differences underscore the divergent selection of Cannabis for either seed, hurd and fibre or pharmaceutical production. Introgression of a functional *CANNABIDIOLIC ACID SYNTHASE* (*CBDAS*) gene from hemp- into drug-type Cannabis for production of cannabidiol (CBD), has resulted in reintroduction of previously segregated gene combinations (Burgarella *et al.*, 2019; Grassa *et al.*, 2021), with noticeable consequences for nutrient use between chemotypes (Jost *et al.*, 2024). Comparison of two distinct genotypes representing CBD- and tetrahydrocannabinol (THC)-dominant drug-type cultivars revealed shifts in the expression of some key genes linked to nutrient transport and assimilation, indicating potential trade-offs between cannabinoid biosynthesis and nutrient efficiency. Cannabinoids can represent up to 28% of the dry weight of the female

flower as a result of selective breeding for highly productive strains of drug-type
Cannabis. In addition, flowers account for up to 70 % of total aboveground biomass,
suggesting resource allocation in Cannabis is weighted towards sink.

To establish a foundation for our molecular understanding of the PSR in
Cannabis, we characterised the morphological, physiological and transcriptomic
response of source and sink organs in a dual-purpose hemp cultivar to varying Pi
supply. We find that PSR-driven acclimation redirects nutrients and resources to
support flower development. In contrast to established crop species, hemp does not
downregulate Pi uptake at luxurious supply levels and depletes vacuolar nitrate
pools to support P assimilation and growth. Lack of suppression of key molecular
PSR components is accompanied by upregulation of high affinity nitrate transporters
and genes associated with nitrate assimilation. Our data provide a detailed
understanding of the PSR and crosstalk with N assimilation in Cannabis to support
the development of strategies for optimizing nutrient use in this multifaceted crop.

MATERIALS AND METHODS

Plant material and growth conditions

Photoperiod sensitive Chinese industrial hemp cultivar Han NW bred for seed
and fibre production was used to generate mother plant stock via single-seed decent
(Clarke, 2020; Welling *et al.*, 2021; Zhang *et al.*, 2021).

Female Han NW plants were clonally propagated using 15-cm long cuttings
taken from lateral shoots of a mother plant. The base of each cutting was dipped in
auxin-enriched gel (Purple Clonex Rooting Hormone Gel, Yates) before transfer into
propagation cubes (EasyPlug, Netherlands) pre-saturated with reverse osmosis
(RO)-purified water. Cuttings were placed in a seedling tray covered by a humidity
dome. Rooting occurred at 24 °C in a controlled environment growth room under
long-day conditions (18 h day / 6 h night cycle), with a light intensity of 120 µmol m^{-2}
s^{-1}.

After two weeks, the most uniform rooted cuttings were transplanted into 0.25
L pots containing medium-grade Perlite (P400, Chillagoe Pty Ltd, Australia) and
grown under long-day conditions with a light intensity of 200 µmol m^{-2} s^{-1}. Daily
fertigation was applied as needed using a modified Hoagland's solution (7.25 mM
NO_3^-, 0.25 mM NH_4^+, 4.5 mM K, 1 mM Mg, 2 mM Ca, 1 mM S, 20 µM Fe, 2 µM Mn,

2 µM Zn, 25 µM Cu, 0.5 µM Co, 0.5 µM Mo, and 25 µM B, with a pH of 6.2)

containing 1 mM Pi, with clonal plants receiving a total of 1.8 L in feed and 1.8 mmol

Pi.

Thirteen days after the first transplant (13 DAT), 15 plants were transferred

into 4.5 L pots with Perlite P400 substrate and grown for another four weeks under

short-day conditions (12 h day / 12 h day-night cycle) with a week-long ramp-up to a

final light intensity of 600 µmol m^{-2} s^{-1} to induce flowering. Five Pi treatments were

initiated immediately after the transplant, with modified Hoagland's solution

containing either 0, 0.25, 0.5, 1, or 2 mM KH_2PO_4. Across treatments, potassium

levels were maintained at a concentration of 4.5 mM using potassium chloride. Three

plants were used in each treatment group (n=3). Fertigation events and volume per

event were adjusted to plant development and transpiration, and all plants received

the same volume of fertigation mix, which totalled 22.5 L over the entire course of

the experiment. All plants were harvest at 42 DAT when female flowers were still

immature with white pistillate flower clusters (Punja and Holmes, 2020).

Sampling

Six different plant organ types (Top Flower (terminal inflorescence), TF;

Young Stem, YS; Mature Stem, MS; Young Leaf, YL; Mature Leaf, ML; Old Leaf, OL;

Lateral Root, LR) were sampled from each plant at 42 DAT. To ensure consistent

phenological definition of leaf types, leaves were pre-selected the day before

sampling to ensure similar size, level of lobe expansion, and nodal position along the

main stem. YS were sampled from nodes below the TF basal node to the next four

lower nodes, while MS were sampled from the node where ML was collected

including the next two lower nodes. LR samples were excised from the exterior of the

root ball, washed twice with RO water, and dried on paper towels. All samples were

shock-frozen in liquid nitrogen, then stored in 15 mL polycarbonate tubes (OPS

Diagnostics, USA) at -80 °C, prior to adding a metal grinding ball (⌀ 0.5 cm) and

processing the frozen material using a Geno/Grinder automated tissue homogenizer

(SPEX SamplePrep, USA) for 2 x 1 min at 1450 rpm. Frozen, ground material was

stored at -80 °C before being sub-sampled under liquid nitrogen for downstream

assays.

Morphological measurements

Over the course of the experiment, and starting at 1 DAT, measurements of plant height, node count, plant width (widest points of fully extended branches) were taken at indicated time points, while ramification points were recorded from 13 DAT onwards. Ramification points are a measure of shoot branching, as described by Pérez-Harguindeguy *et al.* (2016). For this experiment the 3[rd] branch from the base of each plant was chosen as representative node.

Biomass quantification and nutrient efficiency calculations

Plants harvested at 42 DAT were at early flowering stage (terminal racemose inflorescences with white hair-like pistils) (Punja and Holmes, 2020). Plants were separated into flower clusters, leaf and stem for fresh biomass quantification. Root fresh weight was determined following removal of Perlite by two rinses in RO water, followed by a 12-h period of air-drying at room temperature (24 °C). For determination of dry matter, separated organ types were collected in paper bags and dried at 65 °C for approximately seven days to constant weight.

From fresh weight, agronomic phosphate-use efficiency (APE) was calculated as plant biomass produced per unit Pi supplied, expressed as g FW mmol^{-1} Pi, using a modified formula adapted from (Neto *et al.*, 2016):

$$APE = \frac{(FW_{x\,Pi} - FW_{no\,Pi})}{Difference\ in\ the\ total\ available\ Pi\ between\ x\ Pi\ and\ no\ Pi}$$

$FW_{x\,Pi}$ refers to the average organ fresh weight at a given Pi supply which is compared to that of the respective organ type of a plant that received no Pi. The difference in total Pi supplied to each plant over the experimental period was 5.6, 11.2, 22.5 and 44.9 mmol for 0.25, 0.50, 1.0, and 2.0 mM Pi treatments, respectively. Total available Pi (mmol) for the respective Pi treatment was calculated by multiplying the treatment concentration (mM) with the total applied volume (22.5 L), also accounting for 1.8 mmol Pi provided during the 2 weeks of growth in 0.25 L pots prior to transplant at 13 DAT when treatments were initiated.

Determination of nutrient composition and key metabolites

Aliquots of the frozen ground plant material were used for downstream assays as described in (Jost *et al.*, 2024), unless stated otherwise.

One aliquot was used to determine phosphate, sulfate and anthocyanin
concentration following a 1% (v/v) acetic extraction at 4 °C in the dark, and
centrifugation for 10 min at 20,400 x g and 4 °C. Anthocyanin concentration was
determined via the pH differential method (Wrolstad *et al.*, 2005). Free phosphate
was quantified using the molybdenum blue method, while sulfate ions were
quantified measuring turbidity following precipitation with barium chloride (Ames,
1966; Coutinho, 1996).

Nitrate concentration was determined from a second aliquot following water
extraction at 80 °C and 500 rpm (ThermoMixer C, Eppendorf). The mixture was then
centrifuged for 10 min at 20,400 x g and 22 °C, and supernatants used to quantify
nitrate by measuring the yellow complex formed during the nitration of salicylic acid
under acidic conditions at 410 nm (Hachiya and Okamoto, 2017).

A third aliquot was used to determine chlorophyll, protein and starch following
80 % (v/v) ethanol extraction in 10 mM MES buffer pH 5.0 at 70 °C and a 10 min
centrifugation at 11,000 x g and 4 °C, carried out in the dark (Hendriks *et al.*, 2003).
The supernatant then was used to determine chlorophyll (Lichtenthaler, 1987). The
remaining pellet was resuspended in 0.1 mM NaOH at 80 °C and centrifuged as
above, with the supernatant used for protein quantification at 595 nm using
Coomassie Plus Protein Assay Reagent (Thermo Fisher Scientific, Australia). The
remaining pellet was resuspended in 0.5 M HCl in 0.1 M sodium acetate buffer pH
4.9, followed by ON starch hydrolysis at 37 °C. Starch-derived glucose was then
quantified in a coupled enzymatic assay using the NADH released by ATP-
dependent hexokinase (Hendriks *et al.*, 2003).

Elemental composition was analysed via inductively coupled plasma mass
spectrometry (ICP-MS) following acid digestion of 10 mg of dried sample in *aqua*
regia for 3 hours at 70 °C (Foroughi *et al.*, 2014).

Total carbon and nitrogen were quantified using 2 mg of dry plant material,
combusted at 1000 °C, with subsequent chemiluminescence detection of nitrous
oxide in a 2400 Series II CHNS/O Analyzer (Perkin Elmer, Australia). The reference
material for elemental composition and C/N ratio analysis was NIST1573A tomato
leaf (Sigma-Aldrich).

RNA-seq analyses

Total RNA was extracted from ca. 100 mg aliquots of ground frozen material, using the Spectrum Plant Total RNA kit (Sigma-Aldrich) according to the manufacturer's protocol. Removal of gDNA was performed with an On-Column DNaseI digestion kit (Sigma-Aldrich) prior to RNA elution with RNase-free water. RNA-seq library generation and sequencing was performed at the Australian Genome Research Facility (Melbourne) as paired-end 150 bp reads. FastQC software (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) identified that samples had 95% of reads with an average quality score above 35 and an average of 38 million reads. Transcript read counts were quantified as transcripts per million (TPM) by pseudo-aligning reads against a k-mer index built from the transcript models for the *Cannabis sativa* cs10 reference genome downloaded from the Ensembl Plants database (https://plants.ensembl.org/Cannabis_sativa_female) using the kallisto program with 100 bootstraps (Bray *et al.*, 2016). Differentially expressed genes (DEGs) were then quantified using the sleuth package (Pimentel *et al.*, 2017) and genes with TPM > 10 for at least one organ type across treatments were kept for further downstream analyses. Genes were considered differentially expressed if they had a false discovery rate FDR < 0.05 and the expression difference was greater than two-fold ($|\log_2 FC| > 1$). The ComplexHeatmap package was used to generate heatmaps (Gu, 2022).

Statistical analyses

All statistical analyses were performed in R (https://www.r-project.org/) by one-way analysis of variance, followed by Tukey's honestly significant difference (HSD) post-hoc test to compare means. To summarize pairwise comparisons from Tukey's HSD test, multcompView package was used to generate compact letter displays (Graves *et al.*, 2015). For comparison of P fractions with plant metabolites, a Pearson correlation matrix was calculated and visualized with corrplot package (Wei and Simko, 2024).

RESULTS

Biomass prioritization under varying Pi supply leads to differences in use efficiencies

To characterise the acclimation response to different Pi supply in Cannabis, we firstly determined the biomass allocation to different organs in hemp plants under different Pi supplies. At harvest, visible Pi deficiency symptoms such as stunted growth, chlorosis, and fully senescent mature fan leaves were evident only following complete Pi withdrawal. Plants grown with the lowest Pi supply of 0.25 mM showed limited growth suppression – with growth of younger organs likely supported by higher resource mobilisation from the oldest fan leaves which displayed early senescence as evident by their yellow purplish colour (Figure 1A). Plants grown with higher inputs of Pi showed no premature leaf senescence – but some of the oldest fan leaves along the main stem of plants grown at 1 mM and 2 mM Pi supply were yellowing suggesting nitrogen limitation (Cockson *et al.*, 2019). Plants at the highest Pi supply of 2 mM had smaller leaves which was supported by an average 12% reduction in organ biomass (Supplementary Tables S1 and S2). Root development was similarly constrained, with the largest root systems found in plants treated with 1 mM Pi (Figure 1B, Supplementary Table S1 and 2). The highest shoot-to-root biomass ratio of 2.3 was observed at 0.5 mM Pi supply, while Pi withdrawal reduced this ratio to 1.0 indicating a higher relative resource investment into roots (Supplementary Table S1 and S2).

Time-course analysis revealed significant reductions in height and width due to complete Pi withdrawal which became apparent at between 21 and 29 DAT, highlighting delayed response to Pi deficiency (Figure 2A). This contrasts with the faster response time reported in medicinal cannabis (Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b), but aligns with observations in another hemp cultivar Cockson *et al.* (2020). Branching, quantified as the number of ramification points, was particularly sensitive to Pi supply with more pronounced differences between treatments from 35 DAT onwards. Excess Pi supply (2 mM) reduced branching significantly whilst plants supplied with 1 mM Pi continued with lateral shoot outgrowth. Together with the reductions in overall biomass this suggests potential growth constraints due to overaccumulation of Pi.

Pi deprivation-triggered resource reallocation towards the root is a trait
associated with the PSR in many plant species to enhance Pi acquisition (Lambers
et al., 2006; Péret *et al.*, 2014). In hemp, a relative increase in root over shoot
biomass was only observed under complete Pi withdrawal (Figure 2B,
Supplementary Table S1), indicating a low Pi threshold for suppressing foraging
responses and confirming the high inherent PUE of hemp (Wee Y. *et al.*, 2024). Pi
supply of 0.25 mM was sufficient to shift biomass allocation towards the shoot
(Figure 2B), significantly increasing above ground (leaf, stem, flower) over root
biomass. At this Pi input, vegetative (leaf, stem) biomass was almost four-fold higher
than in P-starved plants, and only 21% lower than in plants treated with 1 mM Pi
(Supplementary Table S1). In comparison to Pi-starved plants, flower biomass
increased significantly by between six- to eight-fold with Pi supply higher than 0.25
mM. There were no significant differences in flower biomass between treatments
consistent with high inherent PUE in this hemp cultivar – but also indicating that Pi
use plateaued at relatively low Pi supplies (Figure 2B, Supplementary Table S1).

Analysis of agronomic phosphate efficiency (APE) found that 0.5 mM Pi
yielded the greatest biomass increase per unit of Pi supplied across all organs
except for roots (Figure 2C), corroborating findings from other Cannabis studies
(Cockson *et al.*, 2020; Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b).
These plants also had the most compact growth habit (Figures 1A and 1B). With
higher Pi supply, APE declined for all organ types except flowers, suggesting
diminishing returns in Pi utilization, likely due to metabolic constraints – and possibly
N.

Organ-specific N-limitation with increasing Pi supply

Next, we assessed how varying Pi supply impacted different fractions of P
(organic P, Po; inorganic P, Pi) and allocation of other nutrients (nitrate, total
nitrogen, sulfate) between different organs (TF, terminal flower; YL, young leaf; ML,
mature leaf; OL, old leaf; LR, lateral root; YS, young stem). Significant differences in
Pi and Po fractions were observed across organs and treatments ($p < 0.05$) (Figure
3A, Supplementary Table S3). Following Pi withdrawal, plants maintained a
minimum intracellular Pi concentration of around 5 µmol g^{-1} FW in aboveground
organs regardless of their source or sink identity, indicating that efficient internal Pi
recycling and conservation sustain critical metabolic processes under low Pi

availability in hemp. In Pi-limited LR however, Pi concentration was as low as 1.1 ±
0.3 µmol g^{-1} FW (Figure 3A), likely due to translocation to the shoot. Compared with
Pi withdrawal, Pi concentrations increased significantly in most organs under optimal
Pi supply (1 mM), reaching 18.4 ± 1.8 and 8.5 ± 1.5 µmol g^{-1} FW in shoots and roots,
respectively. OL contained the lowest amount of total P, indicating resource
remobilization (Figure 3A). TF were the strongest sinks, with prioritized Pi allocation
to flowers maintaining Po levels at 11.8 ± 0.4 µmol g^{-1} FW even under Pi withdrawal
– already reaching half (23.8 ± 3.7 µmol g^{-1} FW) of the maximum concentration
under limiting Pi supply (0.25 mM), and rising to 54.5 ± 6.2 µmol g^{-1} FW under
optimal Pi supply (1 mM). Po levels in different leaf types under Pi limitation ($\leq$ 0.25
mM Pi) remained below 2.1 µmol g^{-1} FW, reflecting restricted assimilation of Pi.
Preferential Pi translocation into floral organs was further evident from
elevated concentrations in YS which retained 14.0 ± 1.2 µmol Pi g^{-1} FW even under
Pi withdrawal (Figure 3B), serving as a proxy of sustained active Pi transport to floral
organs. By comparison, Pi levels were at least 50% lower in all other organs. Beyond
1 mM Pi supply, shoot organ concentrations of Pi and Po plateaued, indicating luxury
Pi supply and therefore additional metabolic constraints on further Pi utilization.
However, in LR, both P fractions steadily increased with Pi supply, suggesting a lack
of downregulation in Pi uptake despite sufficient local and systemic P reserves. Lack
of downregulation of root Pi uptake and accumulation under sufficient to luxury Pi
supply in drug-types was found to be genotype-specific (Jost *et al.*, 2024; Shiponi
and Bernstein, 2021b; Wee Y. *et al.*, 2024), while plants such as Arabidopsis,
ryegrass and maize effectively restrict further uptake (Penn *et al.*, 2023; Sharma and
Sahi, 2005; Shukla *et al.*, 2017).
Phosphate and sulfate uptake are often tightly and antagonistically
coordinated (Clarkson *et al.*, 1992; de Jager *et al.*, 2023; Rouached, 2011). In crop
plants, both are usually found at a leaf concentration range of 10 to 20 µmol g^{-1} FW
with sulfate linked to PSR-induced lipid remodelling (Blake-Kalff *et al.*, 2000; Cuyas
et al., 2023; Prodhan *et al.*, 2017; Suriyagoda *et al.*, 2023; Torres-Rodriguez *et al.*,
2021). Across the Pi supply range, hemp roots contained an average 26 ± 2 µmol g^{-1}
FW sulfate suggesting that sulfate uptake and / or translocation are coordinated to
maintain a high basal level of sulfate, irrespective of P status (Figure 3B).
Conversely, in aboveground organs, sulfate concentrations were trending opposite to

phosphate concentrations. P-limited sink organs, particularly TF, accumulated
sulfate with concentrations sharply rising from close to detection limit at 0.5 mM Pi
supply to 19 ± 1 µmol g^{-1} FW upon Pi withdrawal. In YL, sulfate concentrations were
likewise significantly higher at limiting (≤ 0.25 mM) compared with sufficient Pi
supplies (≥ 0.5 mM), but did not exceed 11 ± 2 µmol g^{-1} FW (Figure 3B). With
increasing Pi supply, sulfate concentrations in most shoot organs decreased to less
than 1.0 µmol g^{-1} FW, with the notable exception of YS which retained sulfate
concentrations of greater than 10 µmol g^{-1} FW up until 0.5 mM Pi supply. Increased
translocation can also be deduced from the bell-shaped curve in OL with sulfate
concentration peaking at 0.5 mM Pi supply (Figure 3B) – most likely due to the slow
remobilisation of S during leaf senescence under non-limiting S conditions
(Dubousset *et al.*, 2009; Maillard *et al.*, 2015). Higher rates of translocation may
indicate that sulfate is still used for partial substitution of phospholipids by sulfolipids
under the 0.5 mM Pi treatment. Sulfate concentrations in ML followed a similar Pi
supply-dependent trend as those in sink organs but only reached a maximum of 5.2
± 0.3 µmol g^{-1} FW, which is about four-fold lower than in P-limited TF. In contrast to
hemp, sulfate concentrations in *Arabidopsis* increase in P-limited roots, but decrease
in the respective shoots (Rouached, 2011; Rouached *et al.*, 2011).

N availability in plants is known to affect regulation of PSR genes through
crosstalk between nitrate and Pi signalling with low N status suppressing PSR (Hu
and Chu, 2020; Poza-Carrion and Paz-Ares, 2019; Torres-Rodriguez *et al.*, 2021;
Zhang *et al.*, 2023). Following Pi withdrawal, nitrate concentrations in all but sink
organs were high – ranging from 18 ± 8 µmol g^{-1} FW in LR to 80 ± 36 µmol g^{-1} FW in
ML - likely due to a higher root-to-shoot ratio and decreased metabolic activity in
shoots (Figure 3B). Lower concentrations of 6.0 ± 1.5 µmol g^{-1} FW in TF and $3.5 \pm$
0.9 µmol g^{-1} FW in YL indicate continued N assimilation in P-limited sink organs. In
the presence of as little as 0.25 mM Pi in the nutrient solution, nitrate concentrations
dropped sharply across all aboveground organs, ranging from $0.9 – 4.2$ µmol g^{-1} FW
(Figure 3B), well below adequate concentrations reported in drug-type Cannabis
(Jost *et al.*, 2024). Only transport organs (LR, YS) showed a less severe decline in
nitrate levels reflecting preferential translocation towards sink organs.

Overall, levels of free anions (nitrate, phosphate, and sulfate) suggest N-
limited growth across the entire Pi supply range. To further investigate this, we

analysed total elemental composition, as well as distribution profiles of key pigments
(anthocyanins, chlorophyll) and metabolites (RNA, protein, starch) across organs
and treatments (Supplementary Table S3). There were no treatment-specific
differences in total C or N concentrations in sink (TF, YL) organs and stems again
highlighting active buffering against resource fluctuations (Supplementary
(Supplementary Figure S1). Lower total N concentrations in ML at the lowest Pi
supply level (0.25 mM) indicate P-limited metabolic activity of source leaves.
Accumulation of total N (5-fold) and total C (3-fold) in OL upon Pi withdrawal likely
reflects reduced demand from growing sink organs – and storage of N-rich
metabolites (e.g., N-rich amino acids, polyamines, proteins) or starch, respectively
(Supplementary Figure S1). In LR, a significant two-fold increase in total N under Pi
withdrawal is also observed ($p < 0.05$), likely due to increased biomass partitioning
towards roots for foraging as previously shown in Figures 1B and 1D. Total C,
protein and starch levels in roots remain relatively stable across treatments
(Supplementary Table S3).

Pearson's correlation analysis was used to explore the relationship between
quantified metabolites and P fractions, aiming to identify links between P status and
the accumulation of these metabolites across organs (Figure 4). In YS, there was a
strong positive correlation between Pi and RNA, protein, or starch, showing
dependence of metabolic activity on Pi status. This was also true for the correlation
between starch and Pi or Po in sink organs (TF, YL), highlighting strong association
between Pi availability and carbon fixation. Starch is also known to accumulate
because of metabolic constraints under Pi deficiency (MacNeill *et al.*, 2017), but this
expected negative correlation with P status was only found in roots (MacNeill *et al.*,
2017). Anthocyanin accumulation, a common stress marker in plants (Vance *et al.*,
2003), was negatively correlated with Pi in all shoot organs, although this was only
statistically significant in ML (Vance *et al.*, 2003). Notably, sink organs (TF, YL)
produced more significant correlations compared to source leaves and roots,
suggesting a stronger impact of Pi on their metabolic status. This again highlights
preferential Pi allocation to these organs (Figure 3B). In sink organs, a strong
negative correlation between Pi and Fe was observed, in line with the reported
accumulation of chelated iron complexes in P-limited Arabidopsis shoots (Hirsch *et*
al., 2006). In terminal flowers, Ca had an overall negative correlation to Pi, while K
and Mg had positive correlations to Pi and / or Po in YL and TF, respectively. These

relationships reflect the complex and often antagonistic interactions between Pi and
these cations (Devau *et al.*, 2011; Liu *et al.*, 2011; Rietra *et al.*, 2017; Xie *et al.*,
2021; Xie *et al.*, 2019). Positive correlation between P and K status in sink organs,
especially in YL, agrees with findings in hemp (Cockson *et al.*, 2020) and drug-type
Cannabis (Shiponi and Bernstein, 2021b).

Transcriptional regulation of Phosphate Starvation Response genes implies
coordinated metabolic adjustments across source and sink organs

To gain insight into the underlying transcriptional regulation of observed
physiological responses, RNA-seq analysis was performed and identified a total of
10,327 differentially expressed genes (DEGs) across all organs and treatments
compared with Pi supply of 1 mM. Over half of these genes were unique to OL under
Pi withdrawal (Supplementary Figure S2 A,B; Supplementary Table S5). GO term
enrichment analysis for these genes showed that they are involved in catabolic
processes in agreement with the senescence phenotype of OL (Supplementary
Figure S2C, Figure 1). To focus on specific responses to Pi availability, these
senescence-associated genes were excluded from further analyses.

Pi uptake and translocation

To characterise the transcriptional response in Cannabis, a curated list of
genes with known function in the homeostasis and metabolism of Pi in other species
was generated (Supplementary Table S6). Six out of nine *PHT1* family homologs
identified in hemp were differentially expressed in response to Pi supply in at least
one organ type (Figure 5, Supplementary Tables 5 and 6). Their expression profiles
suggested a function in increasing root uptake from the soil solution as well as
enabling translocation from root to shoot (Wang *et al.*, 2017). The expression of
PHT1 homologs was dependent on Pi supply, with two homologs (*LOC115715182*,
LOC115721935) being responsive across all organs. Their induction was strongest
(up to 962-fold) in Pi-limited source organs (OL and ML), indicating involvement in Pi
mobilisation, with weaker induction observed in transport (LR, YS) and sink organs
(TF). Pi responses of the other *PHT1* genes were organ-specific, *i.e.*, limited to the
root (*LOC115721895*, *LOC115722577*) or to roots and leaves (*LOC115721968*,
LOC115721288) (Wang *et al.,* 2017). Except for *LOC115721935* and
LOC115722577, the other Pi responsive *PHT1* homologs maintained a basal level of

expression under sufficient Pi supply (≥ 1 mM) in LR, likely contributing to the
continuous accumulation of Pi (Figure 3A). Supporting PHT1-mediated Pi
translocation is the expression of the only *PHOSPHATE TRANSPORTER*
FACILITATOR1 (*PHF1*) hemp homolog (*LOC115707553*), which is essential for
intracellular trafficking of PHT1 from the endoplasmic reticulum to the plasma
membrane (Bayle *et al.*, 2011). Under Pi limitation, *CsPHF1* was upregulated by 3-
to 9-fold in LR, OL, ML and YS. In line with the sink status of TF, a low, basal
expression level was maintained (Figure 5, Supplementary Table S5). In hemp,
seven of ten *PHO1(-like)* family members were differentially expressed
(Supplementary Table S5), with *LOC115722063* being the closest homolog of the
functionally best characterised PHO1 in Arabidopsis (Arpat *et al.*, 2012; Hamburger
et al., 2002). Even under optimal Pi supply, this gene showed sustained expression
in LR and YS, underscoring the sustained transport of Pi towards sink organs driven
by PHO1 (Figure 5).

Vacuolar Phosphate Efflux (VPE) and SPX- and MFS-domain containing
Vacuolar Phosphate Transporter (VPT) / PHT5 family members are responsible for
vacuolar Pi efflux and influx, respectively, and play a vital role in intracellular Pi
homeostasis. Their expression is induced upon Pi withdrawal in a PHR1-dependent
manner (Sun *et al.*, 2023). In hemp, we identified two *VPE* and three *VPT / PHT5*
homologs. One hemp *VPE* homolog (*LOC115706709*) was substantially expressed
across organs and responsive to Pi limitation (Figure 5, Supplementary Table S5).
Its observed upregulation in Pi-starved hemp roots hints at fine tuning of cytosolic Pi
pools to ensure that sufficient amounts of Pi are available for xylem loading, which is
also observed in rice (Luan *et al.*, 2019; Sun *et al.*, 2023). In rice, *VPE1* and *VPE2*
are upregulated under Pi-deficient conditions (Xu *et al.*, 2019). In contrast to
Arabidopsis, two of the three *CsVPT* homologs (*LOC115717942*, *LOC115717117*)
were mildly Pi responsive in all organs apart from TF. In Arabidopsis, *VPT*s are
responsible for vacuolar Pi sequestration and are highly upregulated under Pi-stress
to participate in long distance systemic Pi homeostasis (Luan *et al.*, 2019). The
response of the third isoform, *LOC115708335*, was more similar to Arabidopsis
AtVPT1, and showed a more than 1,300-fold induction in Pi-starved roots (Liu *et al.*,
2015; Rose *et al.*, 2016; Yang *et al.*, 2024).

Sensing and Regulation of P status

PHO2, a ubiquitin-conjugating E2 enzyme, is a key regulator of Pi
homeostasis suppressing Pi uptake and translocation under P-replete conditions by
targeting Pi transporters such as PHO1 and PHT1 for ubiquitin-mediated
degradation (Aung *et al.*, 2006; Bari *et al.*, 2006; Liu *et al.*, 2012). The E3 ubiquitin
ligase NITROGEN LIMITATION ADAPTATION (NLA) works alongside PHO2 to
degrade Pi transporters (Lin et al. 2013; Park et al. 2014), as well as PHR1 in
response to changes in inositol polyphosphate levels (Park *et al.*, 2023). Out of two
PHO2 homologs in hemp, we identified one (*LOC115698240*) with lower transcript
abundance in P-limited LR and YS (Figure 5, Supplementary Table S5). Lower
transcript levels are consistent with higher Pi transporter and PHR1 activity in these
organs, as well as systemic, post-transcriptional regulation of *PHO2* mRNA by
miRNA399 observed in Arabidopsis roots (Chiou *et al.*, 2005; Fujii *et al.*, 2005).
However, in P-limited source organs (ML, OL), *PHO2* transcript abundance was 3- to
4-fold higher than under P-replete conditions (Figure 5, Supplementary Table S5).
This may indicate an additional role of PHO2 in suppressing Pi import into these
organs. Like *PHO2*, transcripts encoding the two hemp *NLA* isoforms showed a 3- to
7-fold suppression in P-limited roots, while they were largely unresponsive to Pi-
supply in shoots, with one isoform (*LOC115695922*) showing higher constitutive
expression across organs.

In Arabidopsis and rice, transcript abundance of *PHR1*, the master regulator
of the PSR, is largely unresponsive to Pi supply and subject to post-translational
regulation by SPX proteins and / or ubiquitin-dependent turnover (Bustos *et al.*,
2010; Nilsson *et al.*, 2007; Park *et al.*, 2023). The closest homolog to *AtPHR1*,
LOC115714052, was also not transcriptionally responsive to changes in Pi supply
(Supplementary Table S7). Two close *PHR1-LIKE* (*PHL*) homologs (*CsPHL13* /
LOC115719175, *CsPHL1* / *LOC115701526*) were identified in hemp. CsPHL13 had
a higher constitutive transcript abundance, especially in roots and leaves, and was
responsive to Pi supply with a two-fold induction in P-limited YS (Supplementary
Table S5 and S7). By contrast, CsPHL1 expression was very low across organs but
showed a 60-fold increased expression in roots supplied with 1 and 2 mM Pi (Figure
5, Supplementary Table S7). Two out of the three *SPX* homologs (*LOC115725567* /
CsSPX1, *LOC115697978* / *CsSPX3*) were transcriptionally induced across P-limited
organs by about 30-fold (Figure 5), similar to corresponding homologs in

Arabidopsis, rice and tomato (Puga *et al.*, 2014; Satheesh *et al.*, 2022; Wang *et al.*,
2014). In TF, expression of both *SPX* homologs was five- to seven-fold induced
under complete Pi withdrawal only, indicating Pi sufficiency in TF at a Pi supply
greater than 0.25 mM (Figure 5, Supplementary Table S5). Both genes are therefore
sensitive indicators of plant P status. Inositol polyphosphates (InsP) influence Pi
signalling and homeostasis by regulating SPX-PHR1 interactions (Collins *et al.*,
2024; Ried *et al.*, 2021). In hemp, two putative INOSITOL 1,3,4-TRISPHOSPHATE
5/6-KINASE (ITPK) encoding homologs (*LOC115695287, LOC115716459*) were
identified, showing organ-specific responses to Pi availability. Notably, transcripts
encoding two out of four putative VIP HOMOLOG1 (VIH1) and one VIH2 homolog in
hemp showed constitutive expression across organs and were only differentially
expressed in P-starved OL (Supplementary Table S5 and S7) (Dong *et al.*, 2019;
Laha *et al.*, 2015). In Arabidopsis, ITPK1 and ITPK2 function as key InsP6 kinases to
produce inositol pyrophosphates $InsP_7$ and $InsP_8$ in P-sufficient shoots (Riemer *et*
al., 2021). High $InsP_8$ levels stabilize the SPX-PHR1 interaction to keep PHR1 in an
inactive state, while Pi stress reduces shoot $InsP_8$ concentration to disrupt SPX-
PHR1 binding, thereby allowing PHR1 to bind P1BS motifs and activate PSR genes.
In hemp, transcriptional upregulation of the two ITPK homologs in P-limited LR, ML,
OL and YS (3- to 11- fold change) is counter-intuitive as this should be associated
with reduced PHR1 activity (Figure 5, Supplementary Table S5) (Park *et al.*, 2023).

Mobilisation of Pi from organic compounds by Pi scavenging and lipid remodelling
P-limited plants mobilise Pi by the replacement of membrane phospholipids
and breakdown of organic compounds (e.g. phosphate esters), the latter by
increasing the activity of phosphatases (PAPs) and ribonucleases (RNSs) and their
secretion into the soil (Bariola *et al.*, 1994; Siebers *et al.*, 2015; Tran *et al.*, 2010).
Homologs of several genes involved in lipid remodelling were identified and their
expression followed a Pi supply dependent pattern across organ types, with some
showing the strongest P-limitation responses among all DEGs (Figure 5). Transcripts
encoding a phospholipase D (*PLD, LOC115718607*) homolog were induced 55- to
3,000-fold in P-limited transport (R, YS) and source (ML, OL) organs, respectively
(Supplementary Table 5). The latter corroborates the importance of lipid remodelling
in leaves (Li *et al.*, 2006). *GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE*
(*GDPD)* isoforms are involved in the lipid acyl hydrolase (LAH) pathway of

phospholipid hydrolysis and are upregulated under Pi limitation (Mehra *et al.*, 2019).
GDPD homologs encoded by *LOC115701510* and *LOC115708092* are responsive to
Pi supply across all organ types with the former showing more than 3,000-fold
induction in P-limited source organs (ML, OL) and YS (Figure 5) and attenuated
suppression in P-replete roots compared with other organs. Together with the two
CsSPX isoforms, *LOC115701510* is thus another sensitive expression marker of P
status in Cannabis. *PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE*
PHOSPHATASE1 (*PECP1*) homologs (*LOC115706703*, *LOC115704970*,
LOC115704930, *LOC115704928*) encode haloacid dehydrogenase (HAD)-type
phosphatases involved in cleaving of Pi from phospholipid polar head groups
(Hanchi *et al.*, 2018; Tannert *et al.*, 2018). All three show upregulation under limiting
Pi supply across organ types, with strongest induction observed for LOC115704970
and *LOC115704930* in leaves. *LOC115706703* showed a limited transcriptional
response to Pi supply but had a higher basal level of expression across organ types,
indicating a role in ongoing membrane lipid turnover (Mehra *et al.*, 2019).

The protection of sink organs against phospholipid depletion under low Pi
supply in hemp is illustrated by the expression patterns of
PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE (PEAMT) encoding
homologs *LOC115709566* and *LOC115709567*. In *Arabidopsis*, two PEAMT
isoforms are involved in *de novo* phospholipid synthesis (Chen *et al.*, 2018). In
hemp, transcripts of both *PEAMT* homologs are downregulated by up to 1,200-fold in
P-limited sink organs (ML, OL) and YS (Figure 5; Supplementary Table S5). Yet in
TF, where Pi is preferentially translocated to (Figure 3A, 3B), expression is largely
unaffected, especially for *LOC115709566* which shows a very high transcript
abundance in both TF and YS irrespective of external Pi supply. Interestingly, this is
opposite to what is observed in Arabidopsis*,* where *PEAMT* (*NMT1 / PMT1*)
expression is upregulated in both roots and shoots to maintain phospholipid
synthesis under Pi starvation (Ngo *et al.*, 2022). In aboveground organs of Cannabis,
phospholipid synthesis via *PEAMT* is supported only in the presence of sufficient Pi.
In P-limited hemp organs where *de novo* phospholipid synthesis via *PEAMT* is
limited and phospholipid hydrolysis (*PLD*, *PECP*, *GDPD*) is induced, genes involved
in alternative lipid synthesis are also activated. Galactolipid and sulfolipid
synthesizing enzymes encoded by *MONOGALACTOSYL DIACYLGLYCEROL*
SYNTHASE (*MGD*), DIGALCTOSYLDIACYLGLYCEROL SYNTHASE (*DGD*) and

SULFOQUINOVOSYLDIACYLGLYCEROL SYNTHASE (*SQD*) genes, respectively,
exhibited lower transcriptional activity in sink organs, where phospholipid hydrolysis
was similarly constrained (Figure 5). By contrast, roots and leaves showed increased
transcriptional activity with a 3- to 720-fold induction of select homologs under
limiting Pi supply. Like Arabidopsis, transcripts encoding for one *DGD* homolog
(*LOC115706653*) were largely unresponsive to Pi supply, except for P-starved OL –
and like AtDGD1*,* it probably contributes to constitutive galactolipid synthesis in
chloroplasts (Kobayashi *et al.*, 2014; Kobayashi *et al.*, 2013), while transcripts of two
other homologs (CsMGD2, *LOC115695927*; CsDGD2, *LOC115698658*) are highly Pi
responsive in roots and leaves, just like *AtMGD2* and *AtDGD2* (Kelly and Dörmann,
2002; Kobayashi *et al.*, 2009). Surprisingly, transcripts encoding the type A MGD
synthase CsMGD1 (*LOC115707728*) are induced in P-limited source organs, with a
more constitutive expression in YS and TF. This reminiscent of the simultaneous
induction of both type A and type B MGD synthase genes observed in sesame plants
which are highly tolerant to P impoverished soils (Shimojima *et al.*, 2013). In contrast
to the Arabidopsis genes *AtSQD1* and *AtSQD2*, only one *SQD* homolog (*CsSQD2*,
LOC115697608) was identified in hemp with strong 14- to 30-fold induction in P-
limited roots and leaves, respectively. From these expression profiles, *CsSQD2* and
CsMGD2 are another pair of sensitive markers of organ P status (Figure 5,
Supplementary Table S5).

Pi can also be (re-)mobilised from organic P compounds such as
phosphoesters and RNA catalysed by phosphatases (PAP) and ribonucleases
(RNS). In Arabidopsis, class I ribonucleases encoding *RNS1* and *RNS2* are
upregulated under Pi starvation (Gho *et al.*, 2020; Taylor *et al.*, 1993; Tran *et al.*,
2010). In hemp, two out of six *RNS* homologs were differentially expressed (Figure
5, Supplementary Table S5). Based on their expression profile in hemp, one of the
homologs identified, *CsRNS3* (*LOC115698752*), is more similar to Arabidopsis
RNS1, with strong induction in R, OL and ML, while *CsRNS1* (*LOC115698919*) is
more like *AtRNS2* as transcripts are most abundant in P-limited leaves (Bariola *et
al.*, 1994; Gho *et al.*, 2020; Taylor *et al.*, 1993). Both isoforms exhibit low
transcriptional activity in TF, while in YS, a three- to four-fold induction was evident
upon Pi withdrawal (Figure 5, Supplementary Table S5). Purple acid phosphatases
(PAP*)* are upregulated under Pi stress and play a role in Pi release from intracellular
and / or external organic P pools (Bhadouria and Giri, 2022). The differentially

expressed 15 *PAP* homologs identified in hemp have a similar expression profile to
the *RNS* isoforms (Figure 5), with some showing a strong PSR across roots and
leaves, while others are more responsive in P-limited leaves. Three *PAP* genes,
CsPAP3 / LOC115723670, *CsPAP8 / LOC115723241*, and *CsPAP17 /*
LOC115709102 are good indicators of Pi stress with up to 200-fold induction.

Altered Pi supply impacts N requirements which is reflected in transcriptional
adjustment of nitrate signalling

Pi limitation impacts the homeostasis of other nutrients, especially N, with
coordinated regulation at the transcriptional level (Hu and Chu, 2020; Krouk and
Kiba, 2020; Raven, 2015). In our study, nitrate accumulated significantly in leaves
and young stems following Pi withdrawal (Figure 3B). This is unusual given that – in
most plants studied - nitrate uptake by roots and its translocation to the shoot are
suppressed under Pi limiting conditions (de Magalhães *et al.*, 1998; Gniazdowska
and Rychter, 2001; Jeschke *et al.*, 1997; Krouk and Kiba, 2020; Maeda *et al.*, 2018;
Ueda and Yanagisawa, 2023; Wang *et al.*, 2020). Total N was significantly increased
in P-starved LR and OL only (Supplementary Figure S1), suggesting redirection of N
resources to support the observed increase in root-to-shoot ratio (Figure 2B). Under
sufficient Pi supply, nitrate concentrations were unexpectedly low, indicative of
preferential assimilation into the organic fraction, and likely becoming limiting at
higher Pi supplies (Figure 3B, Supplementary Figure S1).

We next examined the expression of N regulatory genes as well as of genes
encoding steps central to nitrogen uptake and assimilation. An increased demand for
N with Pi supply was particularly evident in LR, where expression of *LOC115708988*,
one of two *NITRATE TRANSPORTER 3* (*NRT3*) genes encoding a key component
of high-affinity nitrate uptake (Okamoto *et al.*, 2006), was steadily induced from 3- to
6-fold as Pi supply was raised from 0.25 mM to 2 mM (Figure 6, Supplementary
Table S5 and S6). Transcript profiles of three out of four hemp homologs encoding
MADS box transcription factor *ARABIDOPSIS NITRATE REGULATED 1* (*ANR1*) are
matching those of *NRT3.1 / LOC115708988* showing root-specific induction with Pi
supply (Zhang and Forde, 1998). This induction of nitrate uptake is most likely
brought about to match the increased capacity for Pi assimilation and consequently
growth (Figure 1). Two-fold induction of a *NITRATE TRANSPORTER 1*
(NRT1)/PEPTIDE TRANSPORTER (*NPF*) homolog (*NRT1.2, LOC115707825*) in YS

with Pi supply points to active transport of nitrate to floral sink organs (Figure 6,
Supplementary Table S5 and S7). Increased expression of *NITRATE REDUCTASE*
(*NIA*, *LOC115713111*) and *GLUTAMINE SYNTHETASE2* (*GLN2*, LOC115722196)
in leaves indicates upregulation of the primary nitrate assimilation pathway to
support new growth when P is adequate, which in turn depletes nitrate levels.
Transcripts encoding NPF transporter homologs *LOC115699885* (*NPF4.5*, root),
LOC115700848 (*NPF6.4*, leaf), LOC115709842 (NPF2.9, leaf), LOC115710913
(NPF7.3, leaf) were more abundant in P-limited organs. Two nitrite reductase
encoding *CsNIR* isoforms (*LOC115702926*, *LOC115702931*), the alpha-subunit of
GLUTAMATE DEHYDROGENASE1 encoding *CsGDH1* (*LOC115705444*), and one
out of two mitochondrial aspartate aminotransferases encoding loci (*AAT1*,
LOC115699642) had predominant expression in source organs under Pi withdrawal
when nitrate concentrations were high. Reduced suppression of *CsGDH1* in source
organs at the highest Pi supply level could be a response to nitrate depletion and
increased N metabolic turnover under excess Pi supply. GDH isoforms catalyse both
the synthesis and catabolism of glutamate, allowing the (re-)assimilation of ammonia
under P or C starvation conditions as well as the generation of N-storage
compounds such as N-rich amino acids and polyamines, particularly under high
nitrate availability and / or a lack of assimilatory capacity (Forde and Lea, 2007).
Contrasts in *CsGLN2* and *CsGDH1* transcript abundance are particularly useful in
determining what constitutes a balanced supply of Pi and N for hemp cultivation.

Stimulation of nitrate uptake by high P status in hemp is also evident from the
transcriptional profile of *NITRATE-INDUCIBLE, GARP-TYPE TRANSCRIPTIONAL*
REPRESSOR1 (*NIGT1*, *LOC115700564*) which suppresses high affinity nitrate
acquisition and is also linked to stimulating PSR in Arabidopsis (Medici *et al.*, 2015;
Ueda *et al.*, 2020a; Ueda *et al.*, 2020b; Wang *et al.*, 2020). AtNIGT1 shows
antagonistic regulation of nitrate transporter gene expression compared with NIN-
LIKE PROTEIN (NLP) transcription factors (Maeda *et al.*, 2018). In hemp, *NIGT1*
transcript abundance is very low in P-replete organs, particularly in source leaves
(ML, OL), while upon Pi withdrawal its expression is upregulated by 8- to 170-fold in
ML, OL, and LR, consistent with its role in suppressing nitrate uptake (Figures 3B
and 6). In TF and YS, weaker (4.3- to 8-fold) *NIGT1* induction is observed only with
Pi withdrawal. As reported in Arabidopsis, *LOC115702149* encoding a NLP6
homolog in hemp shows the opposite expression profile to *NIGT1* (Maeda *et al.*,

2018; Okamoto *et al.*, 2018). Overall, the upregulation of *CsANR1*, *CsNLP6*, and
CsNRT3.1 in P-replete roots, together with *CsNIA*, and *CsGLN2* expression in P-
replete shoots is matched by suppression of *CsNIGT1* and *CsGDH1* across those
organs. This suggests that in hemp, nitrate uptake and N assimilation are highly
responsive to Pi availability. In model organisms such as Arabidopsis, nitrate uptake
is likewise counteracted by PHR1-mediated induction of *NIGT1* upon Pi withdrawal,
whilst excessive Pi uptake is counteracted by down-regulation of *NIGT1* and
promotion of SPX sequestration of PHR1 and NLP (Maeda *et al.*, 2018; Ueda *et al.*,
2020a). In hemp, this seems not to be the case since Pi acquisition is not
suppressed by low organ nitrate levels and one putative *PHR1* homolog, *CsPHL13*,
even showing transcriptional upregulation with increasing Pi supply in nitrate-
depleted source organs (Figure 5).

DISCUSSION

P utilization efficiency attributed to high Pi recycling and conservation in hemp

Developing optimized fertigation strategies and cultivars balancing nutrient uptake and internal utilization capacity is necessary towards building high-performance Cannabis germplasm (Cockson *et al.*, 2020; Hershkowitz *et al.*, 2025; Heuer *et al.*, 2017; Shiponi and Bernstein, 2021b; Tripathi *et al.*, 2022). In our study, the dual-purpose hemp cultivar showed a remarkable resilience to growth inhibition, requiring complete Pi withdrawal for biomass reallocation away from shoot and reproductive organs towards roots (Figure 2B).

The ability of hemp to grow on marginal lands allowed us to study traits for its nutrient-use efficiency (Scalabrin *et al.*, 2024; Wee Y. *et al.*, 2024). This identified key metabolic and transcriptomic acclimation responses in hemp that likely underpin the high PUE. The PSR supports Pi being actively remobilized from roots and source organs towards sink organs, such as flowers (Figure 5). Notably, the young stem serves as a conduit for the efficient translocation of Pi under limiting conditions, with expression closely mirroring that of flowers. These transcriptional changes align very closely with shifts in Pi and Po metabolism across organ types (Figure 3A). Even under Pi limitation (0.25 mM Pi), Po concentration in flowers was 70-fold higher than in all other organ types, indicating their protection from nutrient stress as a hallmark of high PUE in hemp. Similar trends have been observed in other sink-heavy crops such as wheat and rice, where reproductive organs accumulate higher P levels to support grain development in genotypes with high PUE (Adem *et al.*, 2020; Aziz *et al.*, 2014; Hayes *et al.*, 2022; Irfan *et al.*, 2020; Julia *et al.*, 2016; Suriyagoda *et al.*, 2023; Veneklaas *et al.*, 2012).

To maintain central processes such as ATP production, nitrogen and carbon assimilation under P limitation, it is essential that alternative pathways are activated in tandem with intracellular Pi-recycling mechanisms (Albinsky *et al.*, 2010; Carstensen *et al.*, 2018; O'Leary *et al.*, 2011). Under Pi stress, we observed a preferential increase in sulfate accumulation in shoot organs. This is in contrast to other plants such as Arabidopsis where sulfate concentrations decrease in shoot organs while increasing in roots (Rouached, 2011). This increased sulfate concentration in shoots under Pi-stress is indicative of how the hemp cultivar complements ongoing Pi reallocation from source to sink organs by transitioning to

Pi-free glycerolipids such as sulfo- and galactolipids (Hanchi *et al.*, 2018; Nakamura,
2013; Shimojima *et al.*, 2013; Tjellstrom *et al.*, 2008). As opposed to Arabidopsis
where the rate-limiting phospholipid biosynthesis gene *PEAMT* is upregulated under
Pi-stress, its expression was positively correlated with Pi supply in hemp (Figure 5)
(Chen *et al.*, 2018). Considering that Po levels are maintained even under Pi-
limitation in floral organs but not in source organs (Figure 3A), high *PEAMT*
transcript abundance irrespective of Pi supply implies *de novo* phospholipid
synthesis is supported by preferential Pi translocation to reproductive organs (Aziz *et*
al., 2014). In source organs (ML, OL) where Pi is limiting under Pi stress, PEAMT-
mediated phospholipid synthesis is compromised, and expression of genes encoding
alternative lipid synthesis dominates. The latter is supported by the decrease in
sulfate concentration (Figure 3B) relative to reproductive organs at the same level of
Pi supply (Figure 5).

Collectively, these two complementary acclimation responses, i.e. enhanced
Pi allocation to reproductive sinks and heightened release and turnover of organic P
pools through lipid remodelling and phosphatase / nuclease activities, likely underpin
the key adaptations that contribute to the high PUE and sustained floral production
under Pi-stress in hemp.

Nutrient cross-talk and regulatory implications
This study also revealed that hemp adopts a distinctive strategy for balancing
P and N nutrition, differing significantly from canonical models established in, e.g.,
Arabidopsis and rice. Our results suggest that hemp does not down-regulate Pi
uptake and hyperaccumulates Pi in roots and aboveground organs, a trait that is
atypical and suggests an alternative mechanism for Pi sensing and translocation that
is reminiscent of plants adapted to P impoverished soils like the Australian
Proteacea and some legumes (Pang *et al.*, 2009; Prodhan *et al.*, 2016; Shane *et al.*,
2004a; Shane *et al.*, 2004b). This result is supported by other studies in hemp and
drug-type Cannabis (Cockson *et al.*, 2020; Jost *et al.*, 2024; Shiponi and Bernstein,
2021b; Wee Y. *et al.*, 2024). The abundance of transcripts encoding the Pi exporter
responsible for Pi loading into the xylem, PHO1, supports this observed difference in
Pi uptake regulation under excess supply (Figure 5). The dominant *PHO1* gene
remains highly expressed in roots and young stems even under high Pi supply,
underscoring the strong sink strength of reproductive organs (TF) that drives

continuous Pi uptake in roots (Dai *et al.*, 2024; Wang *et al.*, 2004). Continued Pi
accumulation in roots and stems is in spite of *CsPHO2* upregulation under high Pi
supply, which in other plants leads to PHO1 degradation to decrease Pi translocation
to the shoots - pointing towards a decoupling of the PHO2-PHO1 regulatory axis in
hemp (Ouyang *et al.*, 2016). Given that *PHO1* regulates floral transition in both
Arabidopsis and rice, further investigation into its potential role as a Pi-dependent
modulator of flowering time in hemp and drug-type Cannabis is warranted –
particularly in cultivars that exhibit elevated *PHO1* expression (Dai *et al.*, 2024).

Sustained organ Pi accumulation also raises the important question whether
excess Pi uptake in hemp is the result of N limitation as indicated by depleted nitrate
storage reserves under Pi-sufficient conditions (Figure 3B). In many plant species,
efficient biomass production relies on coordinated nitrate and Pi acquisition, as both
are essential for nucleotide and protein biosynthesis (Medici *et al.*, 2019; Pueyo *et
al.*, 2021; Raven, 2015). Ample supply of nitrate is essential towards activation of
PSR genes, whereas nitrate limitation strongly inhibits induction and leads to Pi
accumulation along with downregulation of typical PSR genes in crops such as
maize, and Arabidopsis (Kant *et al.*, 2011; Maeda *et al.*, 2018; Medici *et al.*, 2015;
Schluter *et al.*, 2012; Torres-Rodriguez *et al.*, 2021; Wang *et al.*, 2020). In the hemp
cultivar, Pi and Po status were negatively correlated with nitrate (Figure 4), despite
upregulation of key high affinity nitrate transport system (HATS) component
CsNRT3.1 and corresponding downregulation of its negative GARP-type repressor
CsNIGT1 with increasing Pi supply (Figure 6). Increased N demand with Pi supply is
reflected in the upregulation of key nitrate assimilation genes (*NIA, GLN1*) (Figure 6)
(Liu *et al.*, 2024; Marschner, 2012; Wang *et al.*, 2002). Yet, remarkably, total N and
total C remained unresponsive to Pi supply across aboveground sink organs (Figure
3, Supplementary Figure S1), while the total P and Po fractions – increased with Pi
supply (Figure 3).

These results suggest that in hemp an increase in Pi supply needs to be
carefully balanced with a corresponding higher input of nitrate – as well as
photosynthetic activity, to promote synergistic interactions between N and P to
support higher stem and / or flower biomass and ultimately target product yield. The
observed decoupling of key transcriptional circuits such as PHR1-driven PSR and
the PHO2-NLA regulon in hemp, alongside its inverted Pi–N coordination, extends
current models of nutrient regulation obtained in other species.

Deeper understanding of genotype and usage-type specific nutrient dynamics, helped by the expression profiles of indicative genes identified in our study, has the potential to predict their nutrient use characteristics, enhance cannabinoid production, and support the development of more sustainable cultivation strategies tailored to individual Cannabis germplasm.

SUPPLEMENTARY DATA

Supplementary Figure 1. Total carbon and nitrogen concentrations in response to Pi supply. Shown are total carbon and nitrogen concentrations across organs and Pi treatments (mean ± SE, n = 3 plants per treatment). Letters denote statistical significance between Pi treatments (ANOVA, Tukey HSD, $p < 0.05$).

Supplementary Figure 2. Analysis of differentially expressed genes in OL across Pi treatments. Upset plot for differentially expressed genes (DEGs) shared between P-starved organs (**A**) and Venn diagram of DEGs in OL across treatments (**B**) visualise the disproportionate number of DEGs in the OL samples after withdrawal of Pi (P0 treatment) due to starvation induced premature senescence. (**C**) Gene ontology (GO) term enrichment analysis for the DEGs specific for the OL of P-starved plants highlights senescence- and proteolysis-related terms in upregulated DEGs, while downregulated DEGs are related to photosynthesis as expected of leaves in the later stages of senescence.

Supplementary Table 1: Summary of recorded morphological and physiological data. Sample means and standard errors (SE) were calculated for each variable across the three biological replicates (n) across five treatment groups (0, 0.25, 0.5, 1.0 and 2.0 mM Pi). The residual degrees of freedom (df) from one-way ANOVA was used to calculate the upper and lower 95 % confidence limits (CLs) for each variable's pairwise difference in Tukey's HSD test ($p < 0.05$). To summarize pairwise comparisons from Tukey's HSD, compact letter displays (group) were assigned using the *cld* function from package 'multcompView' to generate letter codes, where alpha = 0.5 was used to separate significantly different treatments. df = N-p, where N is the total number of observations (15) and p is the number of parameters estimated (five treatment levels). ADI = Apical dominance index, *i.e.*, number of ramifications per centimetre of representative branch, APE = agronomic phosphate efficiency, DW = dry weight, FW = fresh weight, Internodal Spacing = average distance between nodes along the main stem, SHI = shoot harvest index, S:R = shoot-to-root biomass ratio, Veg = vegetative biomass. Percent scores represent the biomass (FW or DW) of a given organ type as a % of total biomass.

Supplementary Table 2: **Detailed summary of statistical outputs from Tukey's HSD following one-way ANOVA for recorded morphological and physiological data.** Per variable, treatment pairwise contrasts are given, supplying estimates (difference in group means, *i.e.*, first treatment minus second), pooled standard errors (SE(pooled)) for recorded morphological and physiological data from emmeans Tukey output (pooled, df (residual degrees of freedom), t-statistic (estimate divided by SEpooled)), and adjusted p-values (family-wise error-controlled probabilities from Tukey's HSD for each pairwise contrast). A pooled SE is used because the design is balanced and an assumption of equal variances is assumed. ADI = Apical dominance index, *i.e.*, number of ramifications per centimeter of representative branch, APE = agronomic phosphate efficiency, DW = dry weight, FW = fresh weight, Internodal Spacing = average distance between nodes along the main stem, SHI = shoot harvest index, S:R = shoot-to-root biomass ratio, Veg = vegetative biomass. Percent scores represent the biomass (FW or DW) of a given organ type as a % of total biomass.

Supplementary Table 3: Summary of assayed nutrients and metabolites alongside statistical analysis by ANOVA followed by Tukey's pairwise comparisons, across hemp organs at final destructive harvest. The residual degrees of freedom (df) from one-way ANOVA was used to calculate the upper and lower 95 % confidence limits (CLs) for each variable's pairwise difference in Tukey's HSD test (p < 0.05). To summarize pairwise comparisons from Tukey's HSD, compact letter displays (group) were assigned using the *cld* function from package 'multcompView' to generate letter codes, where at alpha=0.05 to separate significantly different treatments. df= N-p, where N is the total number of observations (15) and p is the number of treatment levels (five). C-N ratio = total carbon to nitrogen ratio, organic P= organic phosphorus.

Supplementary Table 4: **Detailed summary of statistical outputs from Tukey's HSD following one-way ANOVA for assayed metabolite data, across hemp organs at final destructive harvest.** Per variable, pairwise treatment (mM Pi) contrasts are given, supplying estimates (difference in group means, *i.e.*, first treatment minus second), pooled standard errors (SE(pooled)) for recorded morphological and physiological data from emmeans Tukey output, df (residual

degrees of freedom), t-statistic (estimate divided by SEpooled)), and adjusted p-
values (family-wise error-controlled probabilities from Tukey's HSD for each pairwise
contrast). A pooled SE is used because the design is balanced and an assumption of
equal variances is assumed. C-N ratio = total carbon to nitrogen ratio, organic P =
organic phosphorus.

Supplementary Table 5: Summary of DEGs across organs and Pi treatments in
hemp. Gene IDs and descriptions are derived from the cs10 genome annotation.
Selection criteria applied were as follows: FDR adj. p-value < 0.05, $|log_2FC| > 1$,
TPM > 10 in at least 1 organ type across treatments. Values are provided as log_2
fold-change (logFC) relative to organs of plant treated with 1 mM phosphate
(control). Feature = RNA species (m = messenger, nc = non-coding, r = ribosomal,
misc = miscellaneous), FDR = false discovery rate with 'NA' indicating lack of
statistical significance. "gene family" highlights gene families of special interest,
"Pathway annotation" refers to custom annotation.. NA = not available, TF = top
flower / terminal inflorescence, ML = mature fan leaf, OL = old fan leaf, YL = young
fan leaf, YS = young stem, LR = lateral root. P0 = no phosphate supply, P025 = 0.25
mM phosphate supply, P2 = 2 mM phosphate supply.

Supplementary Table 6: Curated list of PSR and other nutrient-response genes
based on known function in other species. Gene IDs and descriptions are derived
from the cs10 genome annotation. Gene annotations have been verified against
Arabidopsis homologs, selecting for the lowest E score (cut-off E<10^-50), and
corresponding gene families have also been provided after curation.

Supplementary Table 7: Transcripts per million (TPM) values across treatment
and organ types. Gene IDs and gene annotations are derived from the cs10
genome annotation. Accession numbers are NCBI RefSeq mRNA transcript
identifiers for each predicted transcript model in the cs10 genome annotation.
Feature = RNA species (m = messenger, nc = non-coding, r = ribosomal, misc =
miscellaneous), TPM = transcripts per million, TF = top flower / terminal
inflorescence; ML = mature fan leaf, OL = old fan leaf, YL = young fan leaf, YS =
young stem, LR = lateral root. P0 = no phosphate supply, P025 = 0.25 mM
phosphate supply, P2 = 2 mM phosphate supply.

ACKNOWLEDGEMENTS

The authors would like to thank Stefanie Kabitz, Filippa Brugliera (both Cann
Group Limited) and colleagues within the ARC Industrial Transformation Research
Hub for Medicinal Agriculture for useful discussions and feedback.

AUTHOR CONTRIBUTIONS

RJ, OB, JW and BWY conceived the project. BWY conducted the experiment,
performed the various assays outlined in the methods as well as data analyses and
visualisation. BWY, OB and RJ analysed the data. AP and SN assisted with
experimental design and data acquisition. BWY, RJ and OB wrote the manuscript
with revision by all authors.
908

CONFLICT OF INTEREST

The authors declare no conflicting interests.

FUNDING

This work was supported by the ARC Industrial Transformation Research Hub
for Medicinal Agriculture (JW, grant ID IH180100006).

DATA AVAILABILITY

RNA-seq read data were deposited at the NCBI SRA database under project
ID PRJNA1278883.

REFERENCES

Adem GD, Ueda Y, Hayes PE, Wissuwa M. 2020. Genetic and physiological traits for internal phosphorus utilization efficiency in rice. PLoS One **15**, e0241842.

Albinsky D, Kusano M, Higuchi M, Hayashi N, Kobayashi M, Fukushima A, Mori M, Ichikawa T, Matsui K, Kuroda H, Horii Y, Tsumoto Y, Sakakibara H, Hirochika H, Matsui M, Saito K. 2010. Metabolomic screening applied to rice FOX Arabidopsis lines leads to the identification of a gene-changing nitrogen metabolism. Mol Plant **3**, 125-142.

Ames B. 1966. Methods in enzymology. Bd. VIII Complex Carbohydrates: Assay of Inorganic Phosphate, Total Phosphate and Phosphatases.

Arpat AB, Magliano P, Wege S, Rouached H, Stefanovic A, Poirier Y. 2012. Functional expression of PHO1 to the Golgi and trans-Golgi network and its role in export of inorganic phosphate. The Plant Journal **71**, 479-491.

Aung K, Lin SI, Wu CC, Huang YT, Su CL, Chiou TJ. 2006. pho2, a phosphate overaccumulator, is caused by a nonsense mutation in a microRNA399 target gene. Plant Physiol **141**, 1000-1011.

Aziz T, Finnegan PM, Lambers H, Jost R. 2014. Organ-specific phosphorus-allocation patterns and transcript profiles linked to phosphorus efficiency in two contrasting wheat genotypes. Plant, Cell & Environment **37**, 943-960.

Bari R, Datt Pant B, Stitt M, Scheible WR. 2006. PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants. Plant Physiol **141**, 988-999.

Bariola PA, Howard CJ, Taylor CB, Verburg MT, Jaglan VD, Green PJ. 1994. The Arabidopsis ribonuclease gene RNS1 is tightly controlled in response to phosphate limitation. Plant J **6**, 673-685.

Bayle V, Arrighi J-F, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E, Paz-Ares J, Nussaume L. 2011. Arabidopsis thaliana High-Affinity Phosphate Transporters Exhibit Multiple Levels of Posttranslational Regulation The Plant Cell **23**, 1523-1535.

Bhadouria J, Giri J. 2022. Purple acid phosphatases: roles in phosphate utilization and new emerging functions. Plant Cell Rep **41**, 33-51.

Blake-Kalff MMA, Hawkesford MJ, Zhao FJ, McGrath SP. 2000. Diagnosing sulfur deficiency in field-grown oilseed rape (Brassica napus L.) and wheat (Triticum aestivum L.). Plant and Soil **225**, 95-107.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol **34**, 525-527.

Burgarella C, Barnaud A, Kane NA, Jankowski F, Scarcelli N, Billot C, Vigouroux Y, Berthouly-Salazar C. 2019. Adaptive Introgression: An Untapped Evolutionary Mechanism for Crop Adaptation. Front Plant Sci **10**, 4.

Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Perez-Perez J, Solano R, Leyva A, Paz-Ares J. 2010. A central regulatory system largely controls transcriptional activation and repression responses to phosphate starvation in Arabidopsis. PLoS Genet **6**, e1001102.

Carstensen A, Herdean A, Schmidt SB, Sharma A, Spetea C, Pribil M, Husted S. 2018. The Impacts of Phosphorus Deficiency on the Photosynthetic Electron Transport Chain. Plant Physiol **177**, 271-284.

Chen W, Salari H, Taylor MC, Jost R, Berkowitz O, Barrow R, Qiu D, Branco R, Masle J. 2018. NMT1 and NMT3 N-Methyltransferase Activity Is Critical to Lipid Homeostasis, Morphogenesis, and Reproduction. Plant Physiol **177**, 1605-1628.

Chen YF, Li LQ, Xu Q, Kong YH, Wang H, Wu WH. 2009. The WRKY6
transcription factor modulates PHOSPHATE1 expression in response to low Pi
stress in Arabidopsis. Plant Cell **21**, 3554-3566.
Chiou T-J, Aung K, Lin S-I, Wu C-C, Chiang S-F, Su C-I. 2005. Regulation of
Phosphate Homeostasis by MicroRNA in Arabidopsis. The Plant Cell **18**, 412-421.
Cho H, Choi I, Bouain N, Nawaz A, Zheng L, Shahzad Z, Brandizzi F, Rhee SY,
Rouached H. 2025. Phosphorus Availability Modulates Flowering Time Through
Subcellular Reprogramming of bGLU25 and GRP7 in Flowering Plants. bioRxiv,
2025.2001.2002.631137.
Clarke RC. 2020. Tasmania Hemp (Cannabis) Fiber and Seed Cultivar Field Trials—
2018-2019. International Journal of Food Science and Agriculture **4**, 470-481.
Clarkson DT, Hawkesford MJ, Davidian J-C, Grignon C. 1992. Contrasting
responses of sulphate and phosphate transport in barley (Hordeum vulgare L.) roots
to protein-modifying reagents and inhibition of protein synthesis. Planta **187**, 306-
314.
Cockson P, Landis H, Smith T, Hicks K, Whipker BE. 2019. Characterization of
Nutrient Disorders of Cannabis sativa. Applied Sciences **9**, 4432.
Cockson P, Schroeder-Moreno M, Veazie P, Barajas G, Logan D, Davis M,
Whipker BE. 2020. Impact of Phosphorus on Cannabis sativa Reproduction,
Cannabinoids, and Terpenes. Applied Sciences **10**, 1-18.
Collins E, Shou H, Mao C, Whelan J, Jost R. 2024. Dynamic interactions between
SPX proteins, the ubiquitination machinery, and signalling molecules for stress
adaptation at a whole-plant level. Biochem J **481**, 363-385.
Coutinho J. 1996. Automated method for sulphate determination in soil-plant
extracts and waters. Communications in Soil Science and Plant Analysis **27**, 727-
740.
Cuyas L, David P, de Craieye D, Ng S, Arkoun M, Plassard C, Faharidine M,
Hourcade D, Degan F, Pluchon S, Nussaume L. 2023. Identification and interest
of molecular markers to monitor plant Pi status. BMC Plant Biol **23**, 401.
Dai S, Chen H, Shi Y, Xiao X, Xu L, Qin C, Zhu Y, Yi K, Lei M, Zeng H. 2024.
PHOSPHATE1-mediated phosphate translocation from roots to shoots regulates
floral transition in plants. J Exp Bot **75**, 5054-5075.
de Jager N, Shukla V, Koprivova A, Lyčka M, Bilalli L, You Y, Zeier J, Kopriva
S, Ristova D. 2023. Traits linked to natural variation of sulfur content in Arabidopsis
thaliana. Journal of Experimental Botany **75**, 1036-1050.
de Magalhães JV, Alves VMC, de Novais RF, Mosquim PR, Magalhães JR, Filho
AFCB, Hubert DM. 1998. Nitrate uptake by corn under increasing periods of
phosphorus starvation. Journal of Plant Nutrition **21**, 1753-1763.
Devau N, Hinsinger P, Le Cadre E, Gérard F. 2011. Root-induced processes
controlling phosphate availability in soils with contrasted P-fertilized treatments. Plant
and Soil **348**, 203-218.
Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang TE,
Wittwer C, Jessen HJ, Zhang H, An GY, Chao DY, Liu D, Lei M. 2019. Inositol
Pyrophosphate InsP(8) Acts as an Intracellular Phosphate Signal in Arabidopsis. Mol
Plant **12**, 1463-1473.
Dubousset L, Abdallah M, Desfeux AS, Etienne P, Meuriot F, Hawkesford MJ,
Gombert J, Segura R, Bataille MP, Reze S, Bonnefoy J, Ameline AF, Ourry A,
Le Dily F, Avice JC. 2009. Remobilization of leaf S compounds and senescence in
response to restricted sulphate supply during the vegetative stage of oilseed rape
are affected by mineral N availability. J Exp Bot **60**, 3239-3253.

Forde BG, Lea PJ. 2007. Glutamate in plants: metabolism, regulation, and signalling. J Exp Bot **58**, 2339-2358.

Foroughi S, Baker AJM, Roessner U, Johnson AAT, Bacic A, Callahan DL. 2014. Hyperaccumulation of zinc by Noccaea caerulescens results in a cascade of stress responses and changes in the elemental profile. Metallomics **6**, 1671-1682.

Fujii H, Chiou T-J, Lin S-I, Aung K, Zhu J-K. 2005. A miRNA Involved in Phosphate-Starvation Response in Arabidopsis. Current Biology **15**, 2038-2043.

Gho YS, Choi H, Moon S, Song MY, Park HE, Kim DH, Ha SH, Jung KH. 2020. Phosphate-Starvation-Inducible S-Like RNase Genes in Rice Are Involved in Phosphate Source Recycling by RNA Decay. Front Plant Sci **11**, 585561.

Gniazdowska A, Rychter AM. 2001. Nitrate uptake by bean (Phaseolus vulgaris L.) roots under phosphate deficiency. In: Gašparíková O, Čiamporová M, Mistrík I, Baluška F, eds. *Recent Advances of Plant Root Structure and Function: Proceedings of the 5th International Symposium on Structure and Function of Roots. Stará Lensná, Slovakia, 30 August – 4 September, 1998*. Dordrecht: Springer Netherlands, 111-117.

Grassa CJ, Weiblen GD, Wenger JP, Dabney C, Poplawski SG, Timothy Motley S, Michael TP, Schwartz CJ. 2021. A new Cannabis genome assembly associates elevated cannabidiol (CBD) with hemp introgressed into marijuana. New Phytol **230**, 1665-1679.

Graves S, Piepho H-P, Selzer ML. 2015. Package 'multcompView'. Visualizations of paired comparisons **451**, 452.

Gu M, Chen A, Sun S, Xu G. 2016. Complex Regulation of Plant Phosphate Transporters and the Gap between Molecular Mechanisms and Practical Application: What Is Missing? Mol Plant **9**, 396-416.

Gu Z. 2022. Complex heatmap visualization. Imeta **1**, e43.

Guo M, Ruan W, Li C, Huang F, Zeng M, Liu Y, Yu Y, Ding X, Wu Y, Wu Z, Mao C, Yi K, Wu P, Mo X. 2015. Integrative Comparison of the Role of the PHOSPHATE RESPONSE1 Subfamily in Phosphate Signaling and Homeostasis in Rice. Plant Physiol **168**, 1762-1776.

Hachiya T, Okamoto Y. 2017. Simple Spectroscopic Determination of Nitrate, Nitrite, and Ammonium in Arabidopsis thaliana. Bio Protoc **7**, e2280.

Hamburger D, Rezzonico E, MacDonald-Comber Petetot J, Somerville C, Poirier Y. 2002. Identification and characterization of the Arabidopsis PHO1 gene involved in phosphate loading to the xylem. Plant Cell **14**, 889-902.

Hanchi M, Thibaud MC, Legeret B, Kuwata K, Pochon N, Beisson F, Cao A, Cuyas L, David P, Doerner P, Ferjani A, Lai F, Li-Beisson Y, Mutterer J, Philibert M, Raghothama KG, Rivasseau C, Secco D, Whelan J, Nussaume L, Javot H. 2018. The Phosphate Fast-Responsive Genes PECP1 and PPsPase1 Affect Phosphocholine and Phosphoethanolamine Content. Plant Physiol **176**, 2943-2962.

Hayes PE, Adem GD, Pariasca-Tanaka J, Wissuwa M. 2022. Leaf phosphorus fractionation in rice to understand internal phosphorus-use efficiency. Ann Bot **129**, 287-302.

Hendriks JH, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose pyrophosphorylase is activated by posttranslational redox-modification in response to light and to sugars in leaves of Arabidopsis and other plant species. Plant Physiol **133**, 838-849.

Hershkowitz JA, Westmoreland FM, Bugbee B. 2025. Elevated root-zone P and nutrient concentration do not increase yield or cannabinoids in medical cannabis. Front Plant Sci **16**, 1433985.

Heuer S, Gaxiola R, Schilling R, Herrera-Estrella L, López-Arredondo D, Wissuwa M, Delhaize E, Rouached H. 2017. Improving phosphorus use efficiency: a complex trait with emerging opportunities. The Plant Journal **90**, 868-885.

Hirsch J, Marin E, Floriani M, Chiarenza S, Richaud P, Nussaume L, Thibaud MC. 2006. Phosphate deficiency promotes modification of iron distribution in Arabidopsis plants. Biochimie **88**, 1767-1771.

Hu B, Chu C. 2020. Nitrogen-phosphorus interplay: old story with molecular tale. New Phytol **225**, 1455-1460.

Irfan M, Aziz T, Maqsood MA, Bilal HM, Siddique KHM, Xu M. 2020. Phosphorus (P) use efficiency in rice is linked to tissue-specific biomass and P allocation patterns. Sci Rep **10**, 4278.

Jeschke WD, Kirkby EA, Peuke AD, Pate JS, Hartung W. 1997. Effects of P deficiency on assimilation and transport of nitrate and phosphate in intact plants of castor bean (Ricinus communis L.). Journal of Experimental Botany **48**, 75-91.

Jiang M, Sun L, Isupov MN, Littlechild JA, Wu X, Wang Q, Wang Q, Yang W, Wu Y. 2019. Structural basis for the Target DNA recognition and binding by the MYB domain of phosphate starvation response 1. FEBS J **286**, 2809-2821.

Jost R, Berkowitz O, Pegg A, Hurgobin B, Tamiru-Oli M, Welling MT, Deseo MA, Noorda H, Brugliera F, Lewsey MG, Doblin MS, Bacic AT, Whelan J. 2024. Sink strength, nutrient allocation, cannabinoid yield and associated transcript profiles vary in two drug-type Cannabis chemovars. Journal of Experimental Botany.

Jost R, Schilling S, Melzer R. 2025. Improving medicinal plant cultivation through in-depth understanding of environmental, physiological, metabolic, and genetic constraints. J Exp Bot **76**, 1-4.

Julia C, Wissuwa M, Kretzschmar T, Jeong K, Rose T. 2016. Phosphorus uptake, partitioning and redistribution during grain filling in rice. Ann Bot **118**, 1151-1162.

Kant S, Peng M, Rothstein SJ. 2011. Genetic regulation by NLA and microRNA827 for maintaining nitrate-dependent phosphate homeostasis in arabidopsis. PLoS Genet **7**, e1002021.

Kelly AA, Dormann P. 2002. DGD2, an arabidopsis gene encoding a UDP-galactose-dependent digalactosyldiacylglycerol synthase is expressed during growth under phosphate-limiting conditions. J Biol Chem **277**, 1166-1173.

Kelly AA, Dörmann P. 2002. DGD2, an Arabidopsis Gene Encoding a UDP-Galactose-dependent Digalactosyldiacylglycerol Synthase Is Expressed during Growth under Phosphate-limiting Conditions*. Journal of Biological Chemistry **277**, 1166-1173.

Kobayashi K, Awai K, Nakamura M, Nagatani A, Masuda T, Ohta H. 2009. Type-B monogalactosyldiacylglycerol synthases are involved in phosphate starvation-induced lipid remodeling, and are crucial for low-phosphate adaptation. Plant J **57**, 322-331.

Kobayashi K, Fujii S, Sasaki D, Baba S, Ohta H, Masuda T, Wada H. 2014. Transcriptional regulation of thylakoid galactolipid biosynthesis coordinated with chlorophyll biosynthesis during the development of chloroplasts in Arabidopsis. Frontiers in Plant Science **Volume 5 - 2014**.

Kobayashi K, Narise T, Sonoike K, Hashimoto H, Sato N, Kondo M, Nishimura M, Sato M, Toyooka K, Sugimoto K, Wada H, Masuda T, Ohta H. 2013. Role of galactolipid biosynthesis in coordinated development of photosynthetic complexes

and thylakoid membranes during chloroplast biogenesis in Arabidopsis. Plant J **73**,
250-261.
Krouk G, Kiba T. 2020. Nitrogen and Phosphorus interactions in plants: from
agronomic to physiological and molecular insights. Curr Opin Plant Biol **57**, 104-109.
Laha D, Johnen P, Azevedo C, Dynowski M, Weiß M, Capolicchio S, Mao H,
Iven T, Steenbergen M, Freyer M, Gaugler P, de Campos MKF, Zheng N,
Feussner I, Jessen HJ, Van Wees SCM, Saiardi A, Schaaf G. 2015. VIH2
Regulates the Synthesis of Inositol Pyrophosphate InsP8 and Jasmonate-Dependent
Defenses in Arabidopsis. The Plant Cell **27**, 1082-1097.
Lambers H. 2021. Phosphorus Acquisition and Utilization in Plants. Annu Rev Plant
Biol.
Lambers H, Shane MW, Cramer MD, Pearse SJ, Veneklaas EJ. 2006. Root
structure and functioning for efficient acquisition of phosphorus: Matching
morphological and physiological traits. Ann Bot **98**, 693-713.
Li M, Qin C, Welti R, Wang X. 2006. Double knockouts of phospholipases Dzeta1
and Dzeta2 in Arabidopsis affect root elongation during phosphate-limited growth but
do not affect root hair patterning. Plant Physiol **140**, 761-770.
Lichtenthaler HK. 1987. [34] Chlorophylls and carotenoids: Pigments of
photosynthetic biomembranes. *Methods in Enzymology*, Vol. 148: Academic Press,
350-382.
Linn J, Ren M, Berkowitz O, Ding W, van der Merwe MJ, Whelan J, Jost R.
2017. Root Cell-Specific Regulators of Phosphate-Dependent Growth. Plant Physiol
174, 1969-1989.
Liu J, Yang L, Luan M, Wang Y, Zhang C, Zhang B, Shi J, Zhao FG, Lan W,
Luan S. 2015. A vacuolar phosphate transporter essential for phosphate
homeostasis in Arabidopsis. Proc Natl Acad Sci U S A **112**, E6571-6578.
Liu T-Y, Aung K, Tseng C-Y, Chang T-Y, Chen Y-S, Chiou T-J. 2011. Vacuolar
Ca2+/H+ Transport Activity Is Required for Systemic Phosphate Homeostasis
Involving Shoot-to-Root Signaling in Arabidopsis   Plant Physiology **156**, 1176-1189.
Liu TY, Huang TK, Tseng CY, Lai YS, Lin SI, Lin WY, Chen JW, Chiou TJ. 2012.
PHO2-dependent degradation of PHO1 modulates phosphate homeostasis in
Arabidopsis. Plant Cell **24**, 2168-2183.
Liu Z, Huang S, Zhu L, Li C, Zhang D, Chen M, Liu Y, Zhang Y. 2024. PHR1
negatively regulates nitrate reductase activity by directly inhibiting the transcription of
NIA1 in Arabidopsis. Journal of Plant Physiology **303**, 154382.
Luan M, Zhao F, Han X, Sun G, Yang Y, Liu J, Shi J, Fu A, Lan W, Luan S. 2019.
Vacuolar Phosphate Transporters Contribute to Systemic Phosphate Homeostasis
Vital for Reproductive Development in Arabidopsis. Plant Physiol **179**, 640-655.
MacNeill GJ, Mehrpouyan S, Minow MAA, Patterson JA, Tetlow IJ, Emes MJ.
2017. Starch as a source, starch as a sink: the bifunctional role of starch in carbon
allocation. J Exp Bot **68**, 4433-4453.
Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y,
Sakakibara H, Yanagisawa S. 2018. A NIGT1-centred transcriptional cascade
regulates nitrate signalling and incorporates phosphorus starvation signals in
Arabidopsis. Nat Commun **9**, 1376.
Maillard A, Diquelou S, Billard V, Laine P, Garnica M, Prudent M, Garcia-Mina
JM, Yvin JC, Ourry A. 2015. Leaf mineral nutrient remobilization during leaf
senescence and modulation by nutrient deficiency. Front Plant Sci **6**, 317.
Marschner H. 2012. *Marschner's Mineral Nutrition of Higher Plants*.

Medici A, Marshall-Colon A, Ronzier E, Szponarski W, Wang R, Gojon A, Crawford NM, Ruffel S, Coruzzi GM, Krouk G. 2015. AtNIGT1/HRS1 integrates nitrate and phosphate signals at the Arabidopsis root tip. Nat Commun **6**, 6274.

Medici A, Szponarski W, Dangeville P, Safi A, Dissanayake IM, Saenchai C, Emanuel A, Rubio V, Lacombe B, Ruffel S, Tanurdzic M, Rouached H, Krouk G. 2019. Identification of Molecular Integrators Shows that Nitrogen Actively Controls the Phosphate Starvation Response in Plants. Plant Cell **31**, 1171-1184.

Mehra P, Pandey BK, Verma L, Giri J. 2019. A novel glycerophosphodiester phosphodiesterase improves phosphate deficiency tolerance in rice. Plant Cell Environ **42**, 1167-1179.

Nakamura Y. 2013. Phosphate starvation and membrane lipid remodeling in seed plants. Prog Lipid Res **52**, 43-50.

Neto AP, Favarin JL, Hammond JP, Tezotto T, Couto HT. 2016. Analysis of Phosphorus Use Efficiency Traits in Coffea Genotypes Reveals Coffea arabica and Coffea canephora Have Contrasting Phosphorus Uptake and Utilization Efficiencies. Front Plant Sci **7**, 408.

Ngo AH, Angkawijaya AE, Lin YC, Liu YC, Nakamura Y. 2022. The phospho-base N-methyltransferases PMT1 and PMT2 produce phosphocholine for leaf growth in phosphorus-starved Arabidopsis. J Exp Bot **73**, 2985-2994.

Nilsson L, Muller R, Nielsen TH. 2007. Increased expression of the MYB-related transcription factor, PHR1, leads to enhanced phosphate uptake in Arabidopsis thaliana. Plant Cell Environ **30**, 1499-1512.

O'Leary B, Park J, Plaxton WC. 2011. The remarkable diversity of plant PEPC (phosphoenolpyruvate carboxylase): recent insights into the physiological functions and post-translational controls of non-photosynthetic PEPCs. Biochem J **436**, 15-34.

Ojeda-Rivera JO, Alejo-Jacuinde G, Najera-Gonzalez HR, Lopez-Arredondo D. 2022. Prospects of genetics and breeding for low-phosphate tolerance: an integrated approach from soil to cell. Theor Appl Genet.

Okamoto M, Kumar A, Li W, Wang Y, Siddiqi MY, Crawford NM, Glass ADM. 2006. High-Affinity Nitrate Transport in Roots of Arabidopsis Depends on Expression of the NAR2-Like Gene AtNRT3.1. Plant Physiology **140**, 1036-1046.

Okamoto Y, Suzuki T, Sugiura D, Kiba T, Sakakibara H, Hachiya T. 2018. Shoot nitrate underlies a perception of nitrogen satiety to trigger local and systemic signaling cascades in
 Arabidopsis thaliana. Soil Science and Plant Nutrition **65**, 56-64.

Okazaki Y, Otsuki H, Narisawa T, Kobayashi M, Sawai S, Kamide Y, Kusano M, Aoki T, Hirai MY, Saito K. 2013. A new class of plant lipid is essential for protection against phosphorus depletion. Nat Commun **4**, 1510.

Ouyang X, , Hong X, , Zhao X, , Zhang W, , He X, , Ma W, , Teng W, , Tong Y, . 2016. Knock out of the PHOSPHATE 2 Gene TaPHO2-A1 Improves Phosphorus Uptake and Grain Yield under Low Phosphorus Conditions in Common Wheat. Scientific Reports **6**, 1-11.

Pang J, Tibbett M, Denton MD, Lambers H, Siddique KHM, Bolland MDA, Revell CK, Ryan MH. 2009. Variation in seedling growth of 11 perennial legumes in response to phosphorus supply. Plant and Soil **328**, 133-143.

Park SH, Jeong JS, Huang CH, Park BS, Chua NH. 2023. Inositol polyphosphates-regulated polyubiquitination of PHR1 by NLA E3 ligase during phosphate starvation response in Arabidopsis. New Phytol **237**, 1215-1228.

Paz-Ares J, Puga MI, Rojas-Triana M, Martinez-Hevia I, Diaz S, Poza-Carrión C, Miñambres M, Leyva A. 2022. Plant adaptation to low phosphorus availability: Core signaling, crosstalks, and applied implications. Mol Plant **15**, 104-124.

Penn CJ, Camberato JJ, Wiethorn MA. 2023. How Much Phosphorus Uptake Is Required for Achieving Maximum Maize Grain Yield? Part 1: Luxury Consumption and Implications for Yield. Agronomy **13**, 95.

Péret B, Desnos T, Jost R, Kanno S, Berkowitz O, Nussaume L. 2014. Root Architecture Responses: In Search of Phosphate. Plant Physiol **166**, 1713-1723.

Pérez-Harguindeguy N, Díaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE, Urcelay C, Veneklaas EJ, Reich PB, Poorter L, Wright IJ, Ray P, Enrico L, Pausas JG, de Vos AC, Buchmann N, Funes G, Quétier F, Hodgson JG, Thompson K, Morgan HD, ter Steege H, Sack L, Blonder B, Poschlod P, Vaieretti MV, Conti G, Staver AC, Aquino S, Cornelissen JHC. 2016. Corrigendum to: New handbook for standardised measurement of plant functional traits worldwide. Australian Journal of Botany **64**.

Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. 2017. Differential analysis of RNA-seq incorporating quantification uncertainty. Nat Methods **14**, 687-690.

Plaxton WC, Tran HT. 2011. Metabolic adaptations of phosphate-starved plants. Plant Physiol **156**, 1006-1015.

Poza-Carrion C, Paz-Ares J. 2019. When nitrate and phosphate sensors meet. Nat Plants **5**, 339-340.

Prodhan MA, Jost R, Watanabe M, Hoefgen R, Lambers H, Finnegan PM. 2016. Tight control of nitrate acquisition in a plant species that evolved in an extremely phosphorus-impoverished environment. Plant Cell Environ **39**, 2754-2761.

Prodhan MA, Jost R, Watanabe M, Hoefgen R, Lambers H, Finnegan PM. 2017. Tight control of sulfur assimilation: an adaptive mechanism for a plant from a severely phosphorus-impoverished habitat. New Phytol **215**, 1068-1079.

Pueyo JJ, Quinones MA, Coba de la Pena T, Fedorova EE, Lucas MM. 2021. Nitrogen and Phosphorus Interplay in Lupin Root Nodules and Cluster Roots. Front Plant Sci **12**, 644218.

Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, de Lorenzo L, Irigoyen ML, Masiero S, Bustos R, Rodriguez J, Leyva A, Rubio V, Sommer H, Paz-Ares J. 2014. SPX1 is a phosphate-dependent inhibitor of Phosphate Starvation Response 1 in Arabidopsis. Proc Natl Acad Sci U S A **111**, 14947-14952.

Punja ZK, Holmes JE. 2020. Hermaphroditism in Marijuana (Cannabis sativa L.) Inflorescences – Impact on Floral Morphology, Seed Formation, Progeny Sex Ratios, and Genetic Variation. Front Plant Sci **11**, 718.

Raven JA. 2015. Interactions between Nitrogen and Phosphorus metabolism. *Annual Plant Reviews Volume 48*, 187-214.

Ried MK, Wild R, Zhu J, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA, Hothorn LA, Fiedler D, Hiller S, Hothorn M. 2021. Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. Nat Commun **12**, 384.

Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei MR, Laha NP, Krusenbaum L, Schneider R, Saiardi A, Fiedler D, Jessen HJ, Schaaf G, Giehl RFH. 2021. ITPK1 is an InsP(6)/ADP phosphotransferase that controls phosphate signaling in Arabidopsis. Mol Plant **14**, 1864-1880.

Rietra RPJJ, Heinen M, Dimkpa CO, Bindraban PS. 2017. Effects of Nutrient Antagonism and Synergism on Yield and Fertilizer Use Efficiency. Communications in Soil Science and Plant Analysis **48**, 1895-1920.

Rose T, Kretzschmar T, Liu L, Lancaster G, Wissuwa M. 2016. Phosphorus Deficiency Alters Nutrient Accumulation Patterns and Grain Nutritional Quality in Rice. Agronomy **6**.

Rouached H. 2011. Multilevel coordination of phosphate and sulfate homeostasis in plants. Plant Signal Behav **6**, 952-955.

Rouached H, Secco D, Arpat B, Poirier Y. 2011. The transcription factor PHR1 plays a key role in the regulation of sulfate shoot-to-root flux upon phosphate starvation in Arabidopsis. BMC Plant Biol **11**, 19.

Rubio V, Linhares F, Solano R, Martin AC, Iglesias J, Leyva A, Paz-Ares J. 2001. A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. Genes Dev **15**, 2122-2133.

Safi A, Medici A, Szponarski W, Ruffel S, Lacombe B, Krouk G. 2017. The world according to GARP transcription factors. Curr Opin Plant Biol **39**, 159-167.

Satheesh V, Zhang J, Li J, You Q, Zhao P, Wang P, Lei M. 2022. High transcriptome plasticity drives phosphate starvation responses in tomato. Stress Biol **2**, 18.

Scalabrin E, Radaelli M, Capodaglio G, Pierobon M, Del Vecchio S, Buffa G. 2024. Hemp cultivation opportunities for marginal lands development. PLoS One **19**, e0299981.

Schluter U, Mascher M, Colmsee C, Scholz U, Brautigam A, Fahnenstich H, Sonnewald U. 2012. Maize source leaf adaptation to nitrogen deficiency affects not only nitrogen and carbon metabolism but also control of phosphate homeostasis. Plant Physiol **160**, 1384-1406.

Shane MW, McCully ME, Lambers H. 2004a. Tissue and cellular phosphorus storage during development of phosphorus toxicity in Hakea prostrata (Proteaceae). Journal of Experimental Botany **55**, 1033-1044.

Shane MW, Szota C, Lambers H. 2004b. A root trait accounting for the extreme phosphorus sensitivity of Hakea prostrata (Proteaceae). Plant, Cell & Environment **27**, 991-1004.

Sharma NC, Sahi SV. 2005. Characterization of Phosphate Accumulation in Lolium multiflorum for Remediation of Phosphorus-Enriched Soils. Environmental Science & Technology **39**, 5475-5480.

Shimojima M, Watanabe T, Madoka Y, Koizumi R, Yamamoto MP, Masuda K, Yamada K, Masuda S, Ohta H. 2013. Differential regulation of two types of monogalactosyldiacylglycerol synthase in membrane lipid remodeling under phosphate-limited conditions in sesame plants. Front Plant Sci **4**, 469.

Shiponi S, Bernstein N. 2021a. Response of medical cannabis (Cannabis sativa L.) genotypes to P supply under long photoperiod: Functional phenotyping and the ionome. Industrial Crops and Products **161**.

Shiponi S, Bernstein N. 2021b. The Highs and Lows of P Supply in Medical Cannabis: Effects on Cannabinoids, the Ionome, and Morpho-Physiology. Front Plant Sci **12**.

Shukla D, Rinehart CA, Sahi SV. 2017. Comprehensive study of excess phosphate response reveals ethylene mediated signaling that negatively regulates plant growth and development. Scientific Reports **7**, 1-16.

Siebers M, Dörmann P, Hölzl G. 2015. Membrane remodelling in phosphorus-deficient plants. *Annual Plant Reviews Volume 48*, 237-263.

Struik PC, Amaducci S, Bullard MJ, Stutterheim NC, Venturi G, Cromack HTH. 2000. Agronomy of fibre hemp (Cannabis sativa L.) in Europe. Industrial Crops and Products **11**, 107-118.

Su T, Xu Q, Zhang FC, Chen Y, Li LQ, Wu WH, Chen YF. 2015. WRKY42 modulates phosphate homeostasis through regulating phosphate translocation and acquisition in Arabidopsis. Plant Physiol **167**, 1579-1591.

Sun G, Luan M, Wen J, Wang B, Lan W. 2023. Genetically controlling VACUOLAR PHOSPHATE TRANSPORTER 1 contributes to low-phosphorus seeds in Arabidopsis. Plant Signal Behav **18**, 2186641.

Suriyagoda LDB, Ryan MH, Gille CE, Dayrell RLC, Finnegan PM, Ranathunge K, Nicol D, Lambers H. 2023. Phosphorus fractions in leaves. New Phytol **237**, 1122-1135.

Tang K, Fracasso A, Struik PC, Yin X, Amaducci S. 2018. Water- and Nitrogen-Use Efficiencies of Hemp (Cannabis sativa L.) Based on Whole-Canopy Measurements and Modeling. Front Plant Sci **9**, 951.

Tannert M, May A, Ditfe D, Berger S, Balcke GU, Tissier A, Kock M. 2018. Pi starvation-dependent regulation of ethanolamine metabolism by phosphoethanolamine phosphatase PECP1 in Arabidopsis roots. J Exp Bot **69**, 467-481.

Taylor CB, Bariola PA, delCardayré SB, Raines RT, Green PJ. 1993. RNS2: a senescence-associated RNase of Arabidopsis that diverged from the S-RNases before speciation. Proceedings of the National Academy of Sciences **90**, 5118-5122.

Tjellstrom H, Andersson MX, Larsson KE, Sandelius AS. 2008. Membrane phospholipids as a phosphate reserve: the dynamic nature of phospholipid-to-digalactosyl diacylglycerol exchange in higher plants. Plant Cell Environ **31**, 1388-1398.

Torres-Rodriguez JV, Salazar-Vidal MN, Chavez Montes RA, Massange-Sanchez JA, Gillmor CS, Sawers RJH. 2021. Low nitrogen availability inhibits the phosphorus starvation response in maize (Zea mays ssp. mays L.). BMC Plant Biol **21**, 259.

Tran HT, Hurley BA, Plaxton WC. 2010. Feeding hungry plants: The role of purple acid phosphatases in phosphate nutrition. Plant Science **179**, 14-27.

Tripathi R, Tewari R, Singh KP, Keswani C, Minkina T, Srivastava AK, De Corato U, Sansinenea E. 2022. Plant mineral nutrition and disease resistance: A significant linkage for sustainable crop protection. Front Plant Sci **13**, 883970.

Ueda Y, Kiba T, Yanagisawa S. 2020a. Nitrate-inducible NIGT1 proteins modulate phosphate uptake and starvation signalling via transcriptional regulation of SPX genes. Plant J **102**, 448-466.

Ueda Y, Nosaki S, Sakuraba Y, Miyakawa T, Kiba T, Tanokura M, Yanagisawa S. 2020b. NIGT1 family proteins exhibit dual mode DNA recognition to regulate nutrient response-associated genes in Arabidopsis. PLoS Genet **16**, e1009197.

Ueda Y, Yanagisawa S. 2023. Transcription factor module NLP-NIGT1 fine-tunes NITRATE TRANSPORTER2.1 expression. Plant Physiol **193**, 2865-2879.

Vance CP, Stone CU, Allan DL. 2003. Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource. New Phytol **157**, 423-447.

Veneklaas EJ, Lambers H, Bragg J, Finnegan PM, Lovelock CE, Plaxton WC, Price CA, Scheible WR, Shane MW, White PJ, Raven JA. 2012. Opportunities for improving phosphorus-use efficiency in crop plants. New Phytol **195**, 306-320.

Wang D, Lv S, Jiang P, Li Y. 2017. Roles, Regulation, and Agricultural Application of Plant Phosphate Transporters. Front Plant Sci **8**, 817.

Wang X, Wang HF, Chen Y, Sun MM, Wang Y, Chen YF. 2020. The Transcription Factor NIGT1.2 Modulates Both Phosphate Uptake and Nitrate Influx during Phosphate Starvation in Arabidopsis and Maize. Plant Cell **32**, 3519-3534.

Wang Y, Ribot C, Rezzonico E, Poirier Y. 2004. Structure and expression profile of the Arabidopsis PHO1 gene family indicates a broad role in inorganic phosphate homeostasis. Plant Physiol **135**, 400-411.

Wang YH, Garvin DF, Kochian LV. 2002. Rapid induction of regulatory and transporter genes in response to phosphorus, potassium, and iron deficiencies in tomato roots. Evidence for cross talk and root/rhizosphere-mediated signals. Plant Physiol **130**, 1361-1370.

Wang Z, Ruan W, Shi J, Zhang L, Xiang D, Yang C, Li C, Wu Z, Liu Y, Yu Y, Shou H, Mo X, Mao C, Wu P. 2014. Rice SPX1 and SPX2 inhibit phosphate starvation responses through interacting with PHR2 in a phosphate-dependent manner. Proc Natl Acad Sci U S A **111**, 14953-14958.

Wee Y. B, Berkowitz O, Whelan J, Jost R. 2024. Same, Yet Different: Towards Understanding Nutrient Use In Hemp- And Drug-Type Cannabis. Journal of Experimental Botany.

Wei T, Simko V. 2024. R package 'corrplot': Visualization of a Correlation Matrix.

Welling MT, Deseo MA, Bacic A, Doblin MS. 2021. Untargeted Metabolomic Analyses Reveal Chemical Complexity of Dioecious *Cannabis* Flowers. Australian Journal of Chemistry **74**, 463-479.

Wrolstad RE, Durst RW, Lee J. 2005. Tracking color and pigment changes in anthocyanin products. Trends in food science & technology **16**, 423-428.

Xie K, Cakmak I, Wang S, Zhang F, Guo S. 2021. Synergistic and antagonistic interactions between potassium and magnesium in higher plants. The Crop Journal **9**, 249-256.

Xie X, Hu W, Fan X, Chen H, Tang M. 2019. Frontiers | Interactions Between Phosphorus, Zinc, and Iron Homeostasis in Nonmycorrhizal and Mycorrhizal Plants. Frontiers in Plant Science **10**.

Xie Z, Mi Y, Kong L, Gao M, Chen S, Chen W, Meng X, Sun W, Chen S, Xu Z. 2023. Cannabis sativa: origin and history, glandular trichome development, and cannabinoid biosynthesis. Hortic Res **10**, uhad150.

Xu L, Zhao H, Wan R, Liu Y, Xu Z, Tian W, Ruan W, Wang F, Deng M, Wang J, Dolan L, Luan S, Xue S, Yi K. 2019. Identification of vacuolar phosphate efflux transporters in land plants. Nat Plants **5**, 84-94.

Yang SY, Lin WY, Hsiao YM, Chiou TJ. 2024. Milestones in understanding transport, sensing, and signaling of the plant nutrient phosphorus. Plant Cell **36**, 1504-1523.

Ye Q, Wang H, Su T, Wu W-H, Chen Y-F. 2018. The Ubiquitin E3 Ligase PRU1 Regulates WRKY6 Degradation to Modulate Phosphate Homeostasis in Response to Low-Pi Stress in Arabidopsis. The Plant Cell **30**, 1062-1076.

Yu B, Xu C, Benning C. 2002. Arabidopsisdisrupted in SQD2encoding sulfolipid synthase is impaired in phosphate-limited growth. Proceedings of the National Academy of Sciences **99**, 5732-5737.

Zager JJ, Lange I, Srividya N, Smith A, Lange BM. 2019. Gene Networks Underlying Cannabinoid and Terpenoid Accumulation in Cannabis. Plant Physiol **180**, 1877-1897.

Zhang H, Forde BG. 1998. An *Arabidopsis* MADS Box Gene That Controls Nutrient-
Induced Changes in Root Architecture. Science **279**, 407-409.
Zhang M, Anderson SL, Brym ZT, Pearson BJ. 2021. Photoperiodic Flowering
Response of Essential Oil, Grain, and Fiber Hemp (Cannabis sativa L.) Cultivars.
Front Plant Sci **12**, 694153.
Zhang Z, Zhong Z, Xiong Y. 2023. Sailing in complex nutrient signaling networks:
Where I am, where to go, and how to go? Mol Plant.

FIGURE LEGENDS:

Figure 1. Phenotypic response of female hemp plants to changes in Pi supply. Shown are images of representative clonal plants for each treatment prior to final destructive harvest at 42 DAT. (**A**) Plant morphology across treatments. Chlorosis gradient increased in senescing leaves along the main stem with decreasing Pi supply. Scale bar represents 20 cm. (**B**) Root ball morphology across treatments. Scale bar represents 5 cm. Organ biomass in g FW plant^{-1} is given below each panel, as mean ± SE (n = 3 plants per treatment).

Figure 2. Morphological response to Pi supply across timepoints. (**A**) Changes in plant height, plant width and node count were determined at indicated time points across treatments. DAT refers to <u>d</u>ays <u>a</u>fter <u>t</u>ransfer of rooted female cuttings into pots. Arrows indicate timepoint when Pi treatments were initiated. (**B**) Biomass partitioning across organ types, as a percentage of total fresh biomass for each treatment. Vegetative organs are stem and leaves combined. (**C**) Agronomic phosphate efficiency (APE) across treatments in comparison to 0 mM Pi supply. APE peaks at 1 mM Pi supply, before decreasing at excess Pi supply across organs. Note that flowers are shielded from these effects. Given are means ± SE, n = 3 plants per treatment. Letters denote significant difference (ANOVA, Tukey's HSD, $p < 0.05$) between treatments and / or organ types. Also refer to Supplementary Table S1 and S2 for supporting datasets.

Figure 3. Pi supply affects Pi assimilation as well as inorganic storage pools across organ types. (**A**) Concentrations of inorganic and organic P across organ types and Pi treatments. (**B**) Concentrations of nitrate, phosphate and sulfate across organ types and Pi treatments. Letters denote statistical significance across treatments (ANOVA/TUKEY HSD, $p < 0.05$). Given are means ± SE (n = 3 plants per treatment). See Supplementary Table S3 for data and statistical analyses. TF, terminal flower; YL, young leaf; ML, mature leaf; OL, old leaf; LR, lateral root; YS, young stem.

Figure 4. Correlation matrix of measured metabolites and P fractions across organ types. Pearson's correlation to assess relationship between concentrations

of indicated metabolites and different P fractions for given organs. Asterisks denote significant correlations ($p < 0.05$) and colour scale indicates correlation coefficient. ND: not determined.

Figure 5. Transcriptional response of curated PSR genes. Cannabis homologs of known PSR genes in Arabidopsis were categorised into functional groups. PSR transcript abundances across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower) and Pi treatments (0, 0.25, 1, 2 mM Pi) are expressed as $\log_2$ transformed TPM values. Transcriptomic data and curated gene list are provided in Supplementary Table S5 to S7. The cs10 reference referenc gene IDs are shown on the right. Abbreviations for gene family names shown to the left of the panel: *PHT, PHOSPHATE TRANSPORTER; PHF1, PHOSHATE TRANSPORT FACILITATOR1; PHO1, PHOSPHATE1; VPE, VACUOLAR PHOSPHATE EFFLUX TRANSPORTER; VPT, VACUOLAR PHOSPHATE TRANSPORTER; PHO2, PHOSPHATE2; NLA, NITROGEN LIMITATION ADAPTATION; PHR1, PHOSPHATE STARVATION RESPONSE1; SPX, SPX DOMAIN GENE; ITPK, INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE; WRKY, WRKY DNA-BINDING PROTEIN; MYB, MYB-DOMAIN PROTEIN; RNS, RIBONUCLEASE; PAP, PURPLE ACID PHOSPHATASE; PLD, PHOSPHOLIPASE D; GDPD, GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE; PECP1, PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE PHOSPHATASE1; PEAMT, PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE; MGD, MONO-GALACTOSYL DIACYLGLYCEROL SYNTHASE; SQD, SULFOQUINOVOSYL-DIACYLGLYCEROL SYNTHASE.*

Figure 6. Transcriptionalresponse of curated genes involved in N assimilation and metabolism in hemp. Shown are transcript abundances for Cannabis homologs of genes with a known role in N assimilation, metabolism and their transcriptional regulation in Arabidopsis. The heatmap represents log-transformed TPM ($\log_2(\text{TPM}+1)$) across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower). Transcriptomic dataset and fully curated gene list is provided in Supplementary Table S5-S7. cs10 genome gene IDs are shown on the right. Abbreviations for gene family names shown on the left of the panel: *AMT, AMMONIUM TRANSPORTER; NRT, NITRATE TRANSPORTER;*

ANR1, ARABIDOPSIS NITRATE REGULATED1; NITRITE REDUCTASE, NIR; NITRATE REDUCTASE, NIA; GLUTAMINE SYNTHETASE, GLN; NITRATE-INDUCIBLE GARP-TYPE TRANSCRIPTIONAL REPRESSOR1, NIGT1; NIN-LIKE PROTEIN, NLP; NRT1/PEPTIDE TRANSPORTER, NPF; GLUTAMATE DEHYDROGENASE, GDH; ASPARTATE AMINOTRANSFERASE, AAT; C-TERMINALLY ENCODED PEPTIDE, CEP; CALCINEURIN B-LIKE PROTEIN-INTERACTING PROTEIN KINASE, CIPK.

Figure 1. Phenotypic response of female hemp plants to changes in Pi supply. Shown are images of representative clonal plants for each treatment prior to final destructive harvest at 42 DAT. (**A**) Plant morphology across treatments. Chlorosis gradient increased in senescing leaves along the main stem with decreasing Pi supply. Scale bar represents 20 cm. (**B**) Root ball morphology across treatments. Scale bar represents 5 cm. Organ biomass in g FW plant^{-1} is given below each panel, as mean ± SE (n = 3 plants per treatment).

Figure 2. Morphological response to Pi supply across timepoints. (A) Changes in plant height, plant width and node count were determined at indicated time points across treatments. DAT refers to days after transfer of rooted female cuttings into pots. Arrows indicate timepoint when Pi treatments were initiated. **(B)** Biomass partitioning across organ types, as a percentage of total fresh biomass for each treatment. Vegetative organs are stem and leaves combined. **(C)** Agronomic phosphate efficiency (APE) across treatments in comparison to 0 mM Pi supply. APE peaks at 1 mM Pi supply, before decreasing at excess Pi supply across organs. Note that flowers are shielded from these effects. Given are means ± SE, n = 3 plants per treatment. Letters denote significant difference (ANOVA, Tukey's HSD, p < 0.05) between treatments and / or organ types. Also refer to Supplementary Tables 1 and 2 for supporting datasets.

Figure 3. Pi supply affects Pi assimilation as well as inorganic storage pools across organ types.
(A) Concentrations of inorganic and organic P across organ types and Pi treatments. **(B)** Concentrations of nitrate, phosphate and sulfate across organ types and Pi treatments. Letters denote statistical significance across treatments (ANOVA/TUKEY HSD, $p < 0.05$). Given are means ± SE (n = 3 plants per treatment). See Supplementary Table S3 for data and statistical analyses. TF, terminal flower; YL, young leaf; ML, mature leaf; OL, old leaf; LR, lateral root; YS, young stem.

Figure 4. Correlation matrix of measured metabolites and P fractions across organ types. Pearson's correlation to assess relationship between concentrations of indicated metabolites and different P fractions for given organs. Asterisks denote significant correlations ($p < 0.05$) and colour scale indicates correlation coefficient. ND: not determined.

Figure 5. Transcriptional response of curated PSR genes. *Cannabis* homologs of known PSR genes in Arabidopsis were categorised into functional groups. PSR transcript abundance across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower) is expressed as log2 transformed TPM values. Transcriptomic data and curated gene list are provided in Supplemental Tables 5 to 7. The cs10 reference gene IDs are shown on the right. Abbreviations for gene family names shown to the left of the panel: *PHT, PHOSPHATE TRANSPORTER; PHF1, PHOSHATE TRANSPORT FACILITATOR1; PHO1, PHOSPHATE1; VPE, VACUOLAR PHOSPHATE EFFLUX TRANSPORTER; VPT, VACUOLAR PHOSPHATE TRANSPORTER; PHO2, PHOSPHATE2; NLA, NITROGEN LIMITATION ADAPTATION; PHR1, PHOSPHATE STARVATION RESPONSE1; SPX, SPX DOMAIN GENE; ITPK, INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE; WRKY, WRKY DNA-BINDING PROTEIN; MYB, MYB-DOMAIN PROTEIN; RNS, RIBONUCLEASE; PAP, PURPLE ACID PHOSPHATASE; PLD, PHOSPHOLIPASE D; GDPD, GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE; PECP1, PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE PHOSPHATASE1; PEAMT, PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE; MGD, MONO-GALACTOSYL DIACYLGLYCEROL SYNTHASE; SQD, SULFOQUINOVOSYL-DIACYLGLYCEROL SYNTHASE.*

Figure 6. Transcriptional response of curated genes involved in N assimilation and metabolism in hemp. Shown are transcript abundances for *Cannabis* homologs of genes with a known role in N assimilation, metabolism and their transcriptional regulation in Arabidopsis. The heatmap represents log-transformed TPM ($\log_2(TPM+1)$) across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower). Transcriptomic dataset and fully curated gene list is provided in Supplemental Tables 5-7. cs10 genome gene IDs are shown on the right. Abbreviations for gene family names shown on the left of the panel: *AMT, AMMONIUM TRANSPORTER; NRT, NITRATE TRANSPORTER; ANR1, ARABIDOPSIS NITRATE REGULATED1; NITRITE REDUCTASE, NIR; NITRATE REDUCTASE, NIA; GLUTAMINE SYNTHETASE, GLN; NITRATE-INDUCIBLE GARP-TYPE TRANSCRIPTIONAL REPRESSOR1, NIGT1; NIN-LIKE PROTEIN, NLP; NRT1/PEPTIDE TRANSPORTER, NPF; GLUTAMATE DEHYDROGENASE, GDH; ASPARTATE AMINOTRANSFERASE, AAT; C-TERMINALLY ENCODED PEPTIDE, CEP; CALCINEURIN B-LIKE PROTEIN-INTERACTING PROTEIN KINASE, CIPK.*