METABOLIC AND TRANSCRIPTOMIC ANALYSES IDENTIFY COORDINATED RESOURCE REALLOCATION IN RESPONSE TO PHOSPHATE SUPPLY IN HEMP

Benjamin Wee Y.[1,2], Oliver Berkowitz[1,2], Sophia Ng[1,2], Amelia Pegg[1,2], James Whelan[1,3], Ricarda Jost[1,2*]

[1]ARC Research Hub for Medicinal Agriculture, Department of Animal, Plant and Soil Sciences, School of Agriculture, Biomedicine and Environment, La Trobe University, Bundoora VIC 3086, Australia
[2]La Trobe Institute for Sustainable Agriculture & Food, La Trobe University, Bundoora VIC 3086, Australia
[3]College of Life Science, Zhejiang University, Hangzhou, Zhejiang, 310058, P.R. China

Author for correspondence:
Ricarda Jost: r.jost@latrobe.edu.au

Email addresses of authors:
Benjamin Wee Y.: 20826392@students.latrobe.edu.au
Oliver Berkowitz: o.berkowitz@latrobe.edu.au
Sophia Ng: jinqtyan@me.com
Amelia Pegg: a.pegg@latrobe.edu.au
James Whelan: jimwhelan@zju.edu.cn
Ricarda Jost: r.jost@latrobe.edu.au

Number of figures: 6
Number of tables: 0
Word count: 7424
Supplementary data: 2 figures and 7 tables

1   **HIGHLIGHT (30 WORDS)**

2   Contrasting human selection in *Cannabis sativa* enables the analysis of
3   divergent nutrient-use strategies. Our integrated transcriptomic and physiological data
4   revealed adaptive resource allocation, informing the development of nutrient-efficient
5   cultivars.

ABSTRACT

Cannabis sativa L., one of the earliest domesticated crops, was selectively bred into drug-type (high phytocannabinoid yield) and hemp-type (fibre and seed production) varieties, resulting in divergent traits including nutrient requirements. To address the limited molecular understanding of nutrient use regulation in Cannabis, this study investigated the response of dual-purpose hemp to varying phosphate availability. Physiological and transcriptomic profiling across source and sink organs revealed that biomass partitioning to flowers was sustained unless phosphate was completely withdrawn. Analyses of key phosphate-responsive genes highlighted the importance of P reallocation to flowers via glycolytic bypass reactions and release of phosphate from organic pools in source organs. Phospholipid biosynthesis and other P-dependent metabolic pathways remained active in sink organs. While the transcriptional response of key regulators, such as SPX DOMAIN GENE (SPX), PHOSPHATE STARVATION RESPONSE1 (PHR1) and NITRATE-INDUCIBLE, GARP-TYPE TRANSCRIPTIONAL REPRESSOR1 (NIGT1) followed canonical patterns, our data suggest a lack of suppression of phosphate uptake and root-to-shoot translocation at higher phosphate supply levels. Subsequent depletion of nitrate pools restricts further growth. These findings uncover conserved and unique aspects of nutrient signalling in hemp, shedding light on its adaptation to nutrient-poor soils through distinct source–sink regulation.

KEYWORDS

Cannabis sativa, hemp, female flower, nitrogen, phosphorus, sink, source, nutrient use, transcriptomics.

INTRODUCTION

Phosphorus (P) is a critical nutrient for plant growth, influencing a range of physiological processes from gene regulation to energy transfer and flowering (Cho *et al.*, 2025; Ojeda-Rivera *et al.*, 2022). It is exclusively taken up by plant roots in the form of inorganic phosphate (Pi, $H_2PO_4^-$). Despite its importance for metabolism and plant development, research on the molecular regulation of Pi acquisition and use in Cannabis is limited. Understanding the impact of Pi on Cannabis reproductive growth or specialised metabolism is essential to improve biomass and product yield (Lambers, 2021; Xie *et al.*, 2023). P deficiency results in reduced growth, accelerated senescence, and lower cannabinoid content, while excess Pi supply does not enhance biomass production or specialised metabolite accumulation (Cockson *et al.*, 2020; Hershkowitz *et al.*, 2025; Shiponi and Bernstein, 2021b). Notably, cannabinoid concentration in tissues is negatively correlated with Pi supply, while total cannabinoid yield per plant increases with Pi supply due to higher flower biomass (Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b; Wee Y. *et al.*, 2024).

Despite its impact on overall product yield, integration of knowledge gained from physiological responses with available molecular data to characterize the phosphate starvation response (PSR) in Cannabis is limited (Jost *et al.*, 2025; Shiponi and Bernstein, 2021a; Shiponi and Bernstein, 2021b; Wee Y. *et al.*, 2024). The PSR describes a highly complex and conserved regulatory system in plants that controls the coordinated acclimation to low Pi availability through changes in nutrient uptake, root system architecture, and shoot metabolism which has been studied extensively in model plants such as *Arabidopsis*, rice and tomato (Paz-Ares *et al.*, 2022; Péret *et al.*, 2014; Plaxton and Tran, 2011; Yang *et al.*, 2024). MYB transcription factors PHOSPHATE STARVATION RESPONSE1 (PHR1) and PHR1-like orthologs play a key role in PSR, controlling the expression of hundreds of genes containing the P1BS (GnATATnC) cis-regulatory motif in their promoter and enhancer regions (Bustos *et al.*, 2010; Guo *et al.*, 2015; Jiang *et al.*, 2019; Linn *et al.*, 2017; Rubio *et al.*, 2001; Safi *et al.*, 2017). Target genes are associated with processes such as Pi acquisition, e.g., members of the high affinity *PHOSPHATE TRANSPORTER1* (*PHT1*) family, Pi scavenging, e.g., purple acid phosphatases (*PAP*s), and ribonucleases (*RNS*), assimilation of other nutrients such as iron, sulfate and nitrate as well as membrane lipid remodelling. Release of Pi by phospholipid hydrolysis requires a shift to

alternative lipids such as sulfo- and galactolipids. Their synthesis is catalysed by UDP-
sulfoquinovose synthases (SQD1 and SQD2), or type B monogalactosyldiacylglycerol
(MGD2 and MGD3) and digalactosyldiacylglycerol (DGD2) synthases, respectively
(Kelly and Dormann, 2002; Kobayashi *et al.*, 2009; Okazaki *et al.*, 2013; Tjellstrom *et*
al., 2008; Yu *et al.*, 2002). PHR1 itself is regulated by SYG1/Pho81/XPR1 (SPX)
domain proteins, which bind higher-order inositol pyrophosphates (IPs) to signal
cellular phosphorus status (Collins *et al.*, 2024). Other notable transcription factors
include WRKY6 and WRKY42, which negatively regulate expression of transcripts
encoding Pi exporter and mediator of Pi root-to-shoot translocation, PHOSPHATE1
(PHO1), via binding to the W-box motif in the *PHO1* promoter (Chen *et al.*, 2009; Su
et al., 2015). At the protein level, PHO1 is negatively regulated by ubiquitin-conjugating
E2 enzyme, PHOSPHATE 2 (PHO2) and E3 ligase PRU1 (Liu *et al.*, 2012; Ye *et al.*,
2018).

Deciphering the interplay between these molecular determinants of PSR and
associated physiological responses is essential for advancing development of crop
cultivars with high P use efficiency (Gu *et al.*, 2016; Heuer *et al.*, 2017). Given drug-
type Cannabis is a high-value medicinal plant largely grown in controlled
environments, there has been little effort to improve its P use. This is reflected by
differences in nutrient response between usage-types, with drug-type Cannabis
exhibiting a tolerance for higher nutrient supplies without yield penalties as opposed
to hemp-types, which are generally more nutrient efficient due to being traditionally
grown on marginal lands (Scalabrin *et al.*, 2024; Struik *et al.*, 2000; Tang *et al.*, 2018;
Wee Y. *et al.*, 2024). These differences underscore the divergent selection of
Cannabis for either seed, hurd and fibre or pharmaceutical production. Introgression
of a functional *CANNABIDIOLIC ACID SYNTHASE* (*CBDAS*) gene from hemp- into
drug-type Cannabis for production of cannabidiol (CBD), has resulted in reintroduction
of previously segregated gene combinations (Burgarella *et al.*, 2019; Grassa *et al.*,
2021), with noticeable consequences for nutrient use between chemotypes (Jost *et*
al., 2024). Comparison of two distinct genotypes representing CBD- and
tetrahydrocannabinol (THC)-dominant drug-type cultivars revealed shifts in the
expression of some key genes linked to nutrient transport and assimilation, indicating
potential trade-offs between cannabinoid biosynthesis and nutrient efficiency.
Cannabinoids can represent up to 28% of the dry weight of the female flower as a
result of selective breeding for highly productive strains of drug-type Cannabis. In

addition, flowers account for up to 70 % of total aboveground biomass, suggesting resource allocation in Cannabis is weighted towards sink.

To establish a foundation for our molecular understanding of the PSR in Cannabis, we characterised the morphological, physiological and transcriptomic response of source and sink organs in a dual-purpose hemp cultivar to varying Pi supply. We find that PSR-driven acclimation redirects nutrients and resources to support flower development. In contrast to established crop species, hemp does not downregulate Pi uptake at luxurious supply levels and depletes vacuolar nitrate pools to support P assimilation and growth. Lack of suppression of key molecular PSR components is accompanied by upregulation of high affinity nitrate transporters and genes associated with nitrate assimilation. Our data provide a detailed understanding of the PSR and crosstalk with N assimilation in Cannabis to support the development of strategies for optimizing nutrient use in this multifaceted crop.

MATERIALS AND METHODS

Plant material and growth conditions

Photoperiod sensitive Chinese industrial hemp cultivar Han NW bred for seed and fibre production was used to generate mother plant stock via single-seed decent (Clarke, 2020; Welling *et al.*, 2021; Zhang *et al.*, 2021).

Female Han NW plants were clonally propagated using 15-cm long cuttings taken from lateral shoots of a mother plant. The base of each cutting was dipped in auxin-enriched gel (Purple Clonex Rooting Hormone Gel, Yates) before transfer into propagation cubes (EasyPlug, Netherlands) pre-saturated with reverse osmosis (RO)-purified water. Cuttings were placed in a seedling tray covered by a humidity dome. Rooting occurred at 24 °C in a controlled environment growth room under long-day conditions (18 h day / 6 h night cycle), with a light intensity of 120 µmol m^{-2} s^{-1}.

After two weeks, the most uniform rooted cuttings were transplanted into 0.25 L pots containing medium-grade Perlite (P400, Chillagoe Pty Ltd, Australia) and grown under long-day conditions with a light intensity of 200 µmol m^{-2} s^{-1}. Daily fertigation was applied as needed using a modified Hoagland's solution (7.25 mM NO_3^-, 0.25 mM NH_4^+, 4.5 mM K, 1 mM Mg, 2 mM Ca, 1 mM S, 20 µM Fe, 2 µM Mn, 2 µM Zn, 25 µM Cu, 0.5 µM Co, 0.5 µM Mo, and 25 µM B, with a pH of 6.2) containing 1 mM Pi, with clonal plants receiving a total of 1.8 L in feed and 1.8 mmol Pi.

Thirteen days after the first transplant (13 DAT), 15 plants were transferred into

4.5 L pots with Perlite P400 substrate and grown for another four weeks under short-

day conditions (12 h day / 12 h day-night cycle) with a week-long ramp-up to a final

light intensity of 600 µmol m^{-2} s^{-1} to induce flowering. Five Pi treatments were initiated

immediately after the transplant, with modified Hoagland's solution containing either

0, 0.25, 0.5, 1, or 2 mM KH$_2$PO$_4$. Across treatments, potassium levels were maintained

at a concentration of 4.5 mM using potassium chloride. Three plants were used in

each treatment group (n=3). Fertigation events and volume per event were adjusted

to plant development and transpiration, and all plants received the same volume of

fertigation mix, which totalled 22.5 L over the entire course of the experiment. All plants

were harvest at 42 DAT when female flowers were still immature with white pistillate

flower clusters (Punja and Holmes, 2020).

Sampling

Six different plant organ types (Top Flower (terminal inflorescence), TF; Young

Stem, YS; Mature Stem, MS; Young Leaf, YL; Mature Leaf, ML; Old Leaf, OL; Lateral

Root, LR) were sampled from each plant at 42 DAT. To ensure consistent phenological

definition of leaf types, leaves were pre-selected the day before sampling to ensure

similar size, level of lobe expansion, and nodal position along the main stem. YS were

sampled from nodes below the TF basal node to the next four lower nodes, while MS

were sampled from the node where ML was collected including the next two lower

nodes. LR samples were excised from the exterior of the root ball, washed twice with

RO water, and dried on paper towels. All samples were shock-frozen in liquid nitrogen,

then stored in 15 mL polycarbonate tubes (OPS Diagnostics, USA) at -80 °C, prior to

adding a metal grinding ball ($\varnothing$ 0.5 cm) and processing the frozen material using a

Geno/Grinder automated tissue homogenizer (SPEX SamplePrep, USA) for 2 x 1 min

at 1450 rpm. Frozen, ground material was stored at -80 °C before being sub-sampled

under liquid nitrogen for downstream assays.

Morphological measurements

Over the course of the experiment, and starting at 1 DAT, measurements of

plant height, node count, plant width (widest points of fully extended branches) were

taken at indicated time points, while ramification points were recorded from 13 DAT

onwards. Ramification points are a measure of shoot branching, as described by

Pérez-Harguindeguy et al. (2016). For this experiment the 3rd branch from the base of

each plant was chosen as representative node.

Biomass quantification and nutrient efficiency calculations

Plants harvested at 42 DAT were at early flowering stage (terminal racemose

inflorescences with white hair-like pistils) (Punja and Holmes, 2020). Plants were

separated into flower clusters, leaf and stem for fresh biomass quantification. Root

fresh weight was determined following removal of Perlite by two rinses in RO water,

followed by a 12-h period of air-drying at room temperature (24 °C). For determination

of dry matter, separated organ types were collected in paper bags and dried at 65 °C

for approximately seven days to constant weight.

From fresh weight, agronomic phosphate-use efficiency (APE) was calculated

as plant biomass produced per unit Pi supplied, expressed as g FW mmol^{-1} Pi, using

a modified formula adapted from (Neto et al., 2016):

174
$$APE = \frac{(FW_{x\,Pi} - FW_{no\,Pi})}{Difference\ in\ the\ total\ available\ Pi\ between\ x\ Pi\ and\ no\ Pi}$$

$FW_{x\,Pi}$ refers to the average organ fresh weight at a given Pi supply which is

compared to that of the respective organ type of a plant that received no Pi. The

difference in total Pi supplied to each plant over the experimental period was 5.6, 11.2,

22.5 and 44.9 mmol for 0.25, 0.50, 1.0, and 2.0 mM Pi treatments, respectively. Total

available Pi (mmol) for the respective Pi treatment was calculated by multiplying the

treatment concentration (mM) with the total applied volume (22.5 L), also accounting

for 1.8 mmol Pi provided during the 2 weeks of growth in 0.25 L pots prior to transplant

at 13 DAT when treatments were initiated.

Determination of nutrient composition and key metabolites

Aliquots of the frozen ground plant material were used for downstream assays

as described in (Jost et al., 2024), unless stated otherwise.

One aliquot was used to determine phosphate, sulfate and anthocyanin

concentration following a 1% (v/v) acetic extraction at 4 °C in the dark, and

centrifugation for 10 min at 20,400 x g and 4 °C. Anthocyanin concentration was

determined via the pH differential method (Wrolstad et al., 2005). Free phosphate was

quantified using the molybdenum blue method, while sulfate ions were quantified

measuring turbidity following precipitation with barium chloride (Ames, 1966;
Coutinho, 1996).

Nitrate concentration was determined from a second aliquot following water
extraction at 80 °C and 500 rpm (ThermoMixer C, Eppendorf). The mixture was then
centrifuged for 10 min at 20,400 x g and 22 °C, and supernatants used to quantify
nitrate by measuring the yellow complex formed during the nitration of salicylic acid
under acidic conditions at 410 nm (Hachiya and Okamoto, 2017).

A third aliquot was used to determine chlorophyll, protein and starch following
80 % (v/v) ethanol extraction in 10 mM MES buffer pH 5.0 at 70 °C and a 10 min
centrifugation at 11,000 x g and 4 °C, carried out in the dark (Hendriks *et al.*, 2003).
The supernatant then was used to determine chlorophyll (Lichtenthaler, 1987). The
remaining pellet was resuspended in 0.1 mM NaOH at 80 °C and centrifuged as
above, with the supernatant used for protein quantification at 595 nm using Coomassie
Plus Protein Assay Reagent (Thermo Fisher Scientific, Australia). The remaining pellet
was resuspended in 0.5 M HCl in 0.1 M sodium acetate buffer pH 4.9, followed by ON
starch hydrolysis at 37 °C. Starch-derived glucose was then quantified in a coupled
enzymatic assay using the NADH released by ATP-dependent hexokinase (Hendriks
et al., 2003).

Elemental composition was analysed via inductively coupled plasma mass
spectrometry (ICP-MS) following acid digestion of 10 mg of dried sample in *aqua regia*
for 3 hours at 70 °C (Foroughi *et al.*, 2014).

Total carbon and nitrogen were quantified using 2 mg of dry plant material,
combusted at 1000 °C, with subsequent chemiluminescence detection of nitrous oxide
in a 2400 Series II CHNS/O Analyzer (Perkin Elmer, Australia). The reference material
for elemental composition and C/N ratio analysis was NIST1573A tomato leaf (Sigma-
Aldrich).

RNA-seq analyses

Total RNA was extracted from ca. 100 mg aliquots of ground frozen material,
using the Spectrum Plant Total RNA kit (Sigma-Aldrich) according to the
manufacturer's protocol. Removal of gDNA was performed with an On-Column
DNaseI digestion kit (Sigma-Aldrich) prior to RNA elution with RNase-free water. RNA-
seq library generation and sequencing was performed at the Australian Genome
Research Facility (Melbourne) as paired-end 150 bp reads. FastQC software

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) identified that samples had 95% of reads with an average quality score above 35 and an average of 38 million reads. Transcript read counts were quantified as transcripts per million (TPM) by pseudo-aligning reads against a k-mer index built from the transcript models for the *Cannabis sativa* cs10 reference genome downloaded from the Ensembl Plants database (https://plants.ensembl.org/Cannabis_sativa_female) using the kallisto program with 100 bootstraps (Bray *et al.*, 2016). Differentially expressed genes (DEGs) were then quantified using the sleuth package (Pimentel *et al.*, 2017) and genes with TPM > 10 for at least one organ type across treatments were kept for further downstream analyses. Genes were considered differentially expressed if they had a false discovery rate FDR < 0.05 and the expression difference was greater than two-fold ($|log_2FC| > 1$). The ComplexHeatmap package was used to generate heatmaps (Gu, 2022).

Statistical analyses

All statistical analyses were performed in R (https://www.r-project.org/) by one-way analysis of variance, followed by Tukey's honestly significant difference (HSD) post-hoc test to compare means. To summarize pairwise comparisons from Tukey's HSD test, multcompView package was used to generate compact letter displays (Graves *et al.*, 2015). For comparison of P fractions with plant metabolites, a Pearson correlation matrix was calculated and visualized with corrplot package (Wei and Simko, 2024).

RESULTS

Biomass prioritization under varying Pi supply leads to differences in use efficiencies

To characterise the acclimation response to different Pi supply in Cannabis, we firstly determined the biomass allocation to different organs in hemp plants under different Pi supplies. At harvest, visible Pi deficiency symptoms such as stunted growth, chlorosis, and fully senescent mature fan leaves were evident only following complete Pi withdrawal. Plants grown with the lowest Pi supply of 0.25 mM showed limited growth suppression – with growth of younger organs likely supported by higher resource mobilisation from the oldest fan leaves which displayed early senescence as evident by their yellow purplish colour (Figure 1A). Plants grown with higher inputs of

Pi showed no premature leaf senescence – but some of the oldest fan leaves along

the main stem of plants grown at 1 mM and 2 mM Pi supply were yellowing suggesting

nitrogen limitation (Cockson *et al.*, 2019). Plants at the highest Pi supply of 2 mM had

smaller leaves which was supported by an average 12% reduction in organ biomass

(Supplementary Tables S1 and S2). Root development was similarly constrained, with

the largest root systems found in plants treated with 1 mM Pi (Figure 1B,

Supplementary Table S1 and 2). The highest shoot-to-root biomass ratio of 2.3 was

observed at 0.5 mM Pi supply, while Pi withdrawal reduced this ratio to 1.0 indicating

a higher relative resource investment into roots (Supplementary Table S1 and S2).

Time-course analysis revealed significant reductions in height and width due to

complete Pi withdrawal which became apparent at between 21 and 29 DAT,

highlighting delayed response to Pi deficiency (Figure 2A). This contrasts with the

faster response time reported in medicinal cannabis (Shiponi and Bernstein, 2021a;

Shiponi and Bernstein, 2021b), but aligns with observations in another hemp cultivar

Cockson *et al.* (2020). Branching, quantified as the number of ramification points, was

particularly sensitive to Pi supply with more pronounced differences between

treatments from 35 DAT onwards. Excess Pi supply (2 mM) reduced branching

significantly whilst plants supplied with 1 mM Pi continued with lateral shoot outgrowth.

Together with the reductions in overall biomass this suggests potential growth

constraints due to overaccumulation of Pi.

Pi deprivation-triggered resource reallocation towards the root is a trait

associated with the PSR in many plant species to enhance Pi acquisition (Lambers *et*

al., 2006; Péret *et al.*, 2014). In hemp, a relative increase in root over shoot biomass

was only observed under complete Pi withdrawal (Figure 2B, Supplementary Table

S1), indicating a low Pi threshold for suppressing foraging responses and confirming

the high inherent PUE of hemp (Wee Y. *et al.*, 2024). Pi supply of 0.25 mM was

sufficient to shift biomass allocation towards the shoot (Figure 2B), significantly

increasing above ground (leaf, stem, flower) over root biomass. At this Pi input,

vegetative (leaf, stem) biomass was almost four-fold higher than in P-starved plants,

and only 21% lower than in plants treated with 1 mM Pi (Supplementary Table S1). In

comparison to Pi-starved plants, flower biomass increased significantly by between

six- to eight-fold with Pi supply above 0.25 mM. There were no significant differences

in flower biomass between treatments consistent with high inherent PUE in this hemp

cultivar – but also indicating that Pi use plateaued at relatively low Pi supplies (Figure
2B, Supplementary Table S1).

Analysis of agronomic phosphate efficiency (APE) found that 0.25 mM Pi
yielded the greatest flower biomass increase per unit of Pi supplied. Plants receiving
0.5 mM P did not invest more in root biomass, and their leaves had no visible
symptoms of nutrient stress (Figure 2C), corroborating findings from other Cannabis
studies (Cockson *et al.*, 2020; Shiponi and Bernstein, 2021a; Shiponi and Bernstein,
2021b). These plants also had the most compact growth habit (Figures 1A and 1B).
At the two highest Pi supply levels, APE declined for all organ types, suggesting
diminishing returns in Pi utilization, likely due to metabolic constraints – and possibly
N limitation. Overall, these results indicate that a Pi supply between 0.25 mM and 0.5
mM optimises the balance for flower production over vegetative biomass.

Organ-specific N-limitation with increasing Pi supply
Next, we assessed how varying Pi supply impacted different fractions of P
(organic P, Po; inorganic P, Pi) and allocation of other nutrients (nitrate, total nitrogen,
sulfate) between different organs (TF, terminal flower; YL, young leaf; ML, mature leaf;
OL, old leaf; LR, lateral root; YS, young stem). Significant differences in Pi and Po
fractions were observed across organs and treatments ($p < 0.05$) (Figure 3A,
Supplementary Table S3). Following Pi withdrawal, plants maintained a minimum
intracellular Pi concentration of around 5 µmol g^{-1} FW in aboveground organs
regardless of their source or sink identity, indicating that efficient internal Pi recycling
and conservation sustain critical metabolic processes under low Pi availability in hemp.
In Pi-limited LR however, Pi concentration was as low as 1.1 ± 0.3 µmol g^{-1} FW (Figure
3A), likely due to translocation to the shoot. Compared with Pi withdrawal, Pi
concentrations increased significantly in most organs under optimal Pi supply (0.5
mM), reaching 18.4 ± 1.8 and 8.5 ± 1.5 µmol g^{-1} FW in shoots and roots, respectively.
OL contained the lowest amount of total P, indicating resource remobilization (Figure
3A). TF were the strongest sinks, with prioritized Pi allocation to flowers maintaining
Po levels at 11.8 ± 0.4 µmol g^{-1} FW even under Pi withdrawal – already reaching half
(23.8 ± 3.7 µmol g^{-1} FW) of the maximum concentration under limiting Pi supply ($\leq$
0.25 mM), and rising to 54.5 ± 6.2 µmol g^{-1} FW under optimal Pi supply (0.5mM). Po
levels in different leaf types under Pi limitation (≤ 0.25 mM Pi) remained below 2.1
µmol g^{-1} FW, reflecting restricted assimilation of Pi.

Preferential Pi translocation into floral organs was further evident from elevated

concentrations in YS which retained 14.0 ± 1.2 µmol Pi g^{-1} FW even under Pi

withdrawal (Figure 3B), serving as a proxy of sustained active Pi transport to floral

organs. By comparison, Pi levels were at least 50% lower in all other organs. Beyond

0.5 mM Pi supply, shoot organ concentrations of Pi and Po plateaued, indicating luxury

Pi supply and therefore additional metabolic constraints on further Pi utilization.

However, in LR, both P fractions steadily increased with Pi supply, suggesting a lack

of downregulation in Pi uptake despite sufficient local and systemic P reserves. Lack

of downregulation of root Pi uptake and accumulation under sufficient to luxury Pi

supply in drug-types was found to be genotype-specific (Jost *et al.*, 2024; Shiponi and

Bernstein, 2021b; Wee Y. *et al.*, 2024), while plants such as Arabidopsis, ryegrass

and maize effectively restrict further uptake (Penn *et al.*, 2023; Sharma and Sahi,

2005; Shukla *et al.*, 2017).

Phosphate and sulfate uptake are often tightly and antagonistically coordinated

(Clarkson *et al.*, 1992; de Jager *et al.*, 2023; Rouached, 2011). In crop plants, both are

usually found at a leaf concentration range of 10 to 20 µmol g^{-1} FW with sulfate linked

to PSR-induced lipid remodelling (Blake-Kalff *et al.*, 2000; Cuyas *et al.*, 2023; Prodhan

et al., 2017; Suriyagoda *et al.*, 2023; Torres-Rodriguez *et al.*, 2021). Across the Pi

supply range, hemp roots contained an average 26 ± 2 µmol g^{-1} FW sulfate suggesting

that sulfate uptake and / or translocation are coordinated to maintain a high basal level

of sulfate, irrespective of P status (Figure 3B). Conversely, in aboveground organs,

sulfate concentrations were trending opposite to phosphate concentrations. P-limited

sink organs, particularly TF, accumulated sulfate with concentrations sharply rising

from close to detection limit at 0.5 mM Pi supply to 19 ± 1 µmol g^{-1} FW upon Pi

withdrawal. In YL, sulfate concentrations were likewise significantly higher at limiting

(≤ 0.25 mM) compared with sufficient Pi supplies (≥ 0.5 mM), but did not exceed $11 \pm$

2 µmol g^{-1} FW (Figure 3B). With increasing Pi supply, sulfate concentrations in most

shoot organs decreased to less than 1.0 µmol g^{-1} FW, with the notable exception of

YS which retained sulfate concentrations of greater than 10 µmol g^{-1} FW up until 0.5

mM Pi supply. Increased translocation can also be deduced from the bell-shaped

curve in OL with sulfate concentration peaking at 0.5 mM Pi supply (Figure 3B) – most

likely due to the slow remobilisation of S during leaf senescence under non-limiting S

conditions (Dubousset *et al.*, 2009; Maillard *et al.*, 2015). Higher rates of translocation

may indicate that sulfate is still used for partial substitution of phospholipids by

sulfolipids under the 0.5 mM Pi treatment. Sulfate concentrations in ML followed a

similar Pi supply-dependent trend as those in sink organs but only reached a maximum

of 5.2 ± 0.3 µmol g^{-1} FW, which is about four-fold lower than in P-limited TF. In contrast

to hemp, sulfate concentrations in *Arabidopsis* increase in P-limited roots, but

decrease in the respective shoots (Rouached, 2011; Rouached *et al.*, 2011).

N availability in plants is known to affect regulation of PSR genes through

crosstalk between nitrate and Pi signalling with low N status suppressing PSR (Hu and

Chu, 2020; Poza-Carrion and Paz-Ares, 2019; Torres-Rodriguez *et al.*, 2021; Zhang

et al., 2023). Following Pi withdrawal, nitrate concentrations in all but sink organs were

high – ranging from 18 ± 8 µmol g^{-1} FW in LR to 80 ± 36 µmol g^{-1} FW in ML - likely due

to a higher root-to-shoot ratio and decreased metabolic activity in shoots (Figure 3B).

Lower concentrations of 6.0 ± 1.5 µmol g^{-1} FW in TF and 3.5 ± 0.9 µmol g^{-1} FW in YL

indicate continued N assimilation in P-limited sink organs. In the presence of as little

as 0.25 mM Pi in the nutrient solution, nitrate concentrations dropped sharply across

all aboveground organs, ranging from $0.9 – 4.2$ µmol g^{-1} FW (Figure 3B), well below

adequate concentrations reported in drug-type Cannabis (Jost *et al.*, 2024). Only

transport organs (LR, YS) showed a less severe decline in nitrate levels reflecting

preferential translocation towards sink organs.

Overall, levels of free anions (nitrate, phosphate, and sulfate) suggest N-limited

growth across the entire Pi supply range. To further investigate this, we analysed total

elemental composition, as well as distribution profiles of key pigments (anthocyanins,

chlorophyll) and metabolites (RNA, protein, starch) across organs and treatments

(Supplementary Table S3). There were no treatment-specific differences in total C or

N concentrations in sink (TF, YL) organs and stems again highlighting active buffering

against resource fluctuations (Supplementary (Supplementary Figure S1). Lower total

N concentrations in ML at the lowest Pi supply level (0.25 mM) indicate P-limited

metabolic activity of source leaves. Accumulation of total N (5-fold) and total C (3-fold)

in OL upon Pi withdrawal likely reflects reduced demand from growing sink organs –

and storage of N-rich metabolites (e.g., N-rich amino acids, polyamines, proteins) or

starch, respectively (Supplementary Figure S1). In LR, a significant two-fold increase

in total N under Pi withdrawal is also observed ($p < 0.05$), likely due to increased

biomass partitioning towards roots for foraging as previously shown in Figures 1B and

1D. Total C, protein and starch levels in roots remain relatively stable across treatments (Supplementary Table S3).

Pearson's correlation analysis was used to explore the relationship between quantified metabolites and P fractions, aiming to identify links between P status and the accumulation of these metabolites across organs (Figure 4). In YS, there was a strong positive correlation between Pi and RNA, protein, or starch, showing dependence of metabolic activity on Pi status. This was also true for the correlation between starch and Pi or Po in sink organs (TF, YL), highlighting strong association between Pi availability and carbon fixation. Starch is also known to accumulate because of metabolic constraints under Pi deficiency (MacNeill *et al.*, 2017), but this expected negative correlation with P status was only found in roots (MacNeill *et al.*, 2017). Anthocyanin accumulation, a common stress marker in plants (Vance *et al.*, 2003), was negatively correlated with Pi in all shoot organs, although this was only statistically significant in ML (Vance *et al.*, 2003). Notably, sink organs (TF, YL) produced more significant correlations compared to source leaves and roots, suggesting a stronger impact of Pi on their metabolic status. This again highlights preferential Pi allocation to these organs (Figure 3B). In sink organs, a strong negative correlation between Pi and Fe was observed, in line with the reported accumulation of chelated iron complexes in P-limited Arabidopsis shoots (Hirsch *et al.*, 2006). In terminal flowers, Ca had an overall negative correlation to Pi, while K and Mg had positive correlations to Pi and / or Po in YL and TF, respectively. These relationships reflect the complex and often antagonistic interactions between Pi and these cations (Devau *et al.*, 2011; Liu *et al.*, 2011; Rietra *et al.*, 2017; Xie *et al.*, 2021; Xie *et al.*, 2019). Positive correlation between P and K status in sink organs, especially in YL, agrees with findings in hemp (Cockson *et al.*, 2020) and drug-type Cannabis (Shiponi and Bernstein, 2021b).

Transcriptional regulation of Phosphate Starvation Response genes implies coordinated metabolic adjustments across source and sink organs

To gain insight into the underlying transcriptional regulation of observed physiological responses, RNA-seq analysis was performed and identified a total of 10,327 differentially expressed genes (DEGs) across all organs and treatments compared with Pi supply of 1 mM. Over half of these genes were unique to OL under Pi withdrawal (Supplementary Figure S2 A,B; Supplementary Table S5). GO term

enrichment analysis for these genes showed that they are involved in catabolic
processes in agreement with the senescence phenotype of OL (Supplementary Figure
S2C, Figure 1). To focus on specific responses to Pi availability, these senescence-
associated genes were excluded from further analyses.

Pi uptake and translocation

To characterise the transcriptional response in Cannabis, a curated list of genes
with known function in the homeostasis and metabolism of Pi in other species was
generated (Supplementary Table S6). Six out of nine *PHT1* family homologs identified
in hemp were differentially expressed in response to Pi supply in at least one organ
type (Figure 5, Supplementary Tables 5 and 6). Their expression profiles suggested a
function in increasing root uptake from the soil solution as well as enabling
translocation from root to shoot (Wang *et al.*, 2017). The expression of *PHT1*
homologs was dependent on Pi supply, with two homologs (*LOC115715182*,
LOC115721935) being responsive across all organs. Their induction was strongest
(up to 962-fold) in Pi-limited source organs (OL and ML), indicating involvement in Pi
mobilisation, with weaker induction observed in transport (LR, YS) and sink organs
(TF). Pi responses of the other *PHT1* genes were organ-specific, *i.e.,* limited to the
root (*LOC115721895*, *LOC115722577*) or to roots and leaves (*LOC115721968*,
LOC115721288) (Wang *et al.*, 2017). Except for *LOC115721935* and *LOC115722577*,
the other Pi responsive *PHT1* homologs maintained a basal level of expression under
surplus Pi supply (≥ 1 mM) in LR, likely contributing to the continuous accumulation of
Pi (Figure 3A). Supporting PHT1-mediated Pi translocation is the expression of the
only *PHOSPHATE TRANSPORTER FACILITATOR1* (*PHF1*) hemp homolog
(*LOC115707553*), which is essential for intracellular trafficking of PHT1 from the
endoplasmic reticulum to the plasma membrane (Bayle *et al.*, 2011). Under Pi
limitation, *CsPHF1* was upregulated by 3- to 9-fold in LR, OL, ML and YS. In line with
the sink status of TF, a low, basal expression level was maintained (Figure 5,
Supplementary Table S5). In hemp, seven of ten *PHO1(-like)* family members were
differentially expressed (Supplementary Table S5), with *LOC115722063* being the
closest homolog of the functionally best characterised PHO1 in Arabidopsis (Arpat *et*
al., 2012; Hamburger *et al.*, 2002). Even under surplus Pi supply, this gene showed
sustained expression in LR and YS, underscoring the sustained transport of Pi towards
sink organs driven by PHO1 (Figure 5).

Vacuolar Phosphate Efflux (VPE) and SPX- and MFS-domain containing
Vacuolar Phosphate Transporter (VPT) / PHT5 family members are responsible for
vacuolar Pi efflux and influx, respectively, and play a vital role in intracellular Pi
homeostasis. Their expression is induced upon Pi withdrawal in a PHR1-dependent
manner (Sun *et al.*, 2023). In hemp, we identified two *VPE* and three *VPT / PHT5*
homologs. One hemp *VPE* homolog (*LOC115706709*) was substantially expressed
across organs and responsive to Pi limitation (Figure 5, Supplementary Table S5). Its
observed upregulation in Pi-starved hemp roots hints at fine tuning of cytosolic Pi pools
to ensure that sufficient amounts of Pi are available for xylem loading, which is also
observed in rice (Luan *et al.*, 2019; Sun *et al.*, 2023). In rice, *VPE1* and *VPE2* are
upregulated under Pi-deficient conditions (Xu *et al.*, 2019). In contrast to Arabidopsis,
two of the three *CsVPT* homologs (*LOC115717942*, *LOC115717117*) were mildly Pi
responsive in all organs apart from TF. In Arabidopsis, *VPT*s are responsible for
vacuolar Pi sequestration and are highly upregulated under Pi-stress to participate in
long distance systemic Pi homeostasis (Luan *et al.*, 2019). The response of the third
isoform, *LOC115708335*, was more similar to Arabidopsis *AtVPT1*, and showed a
more than 1,300-fold induction in Pi-starved roots (Liu *et al.*, 2015; Rose *et al.*, 2016;
Yang *et al.*, 2024).

Sensing and Regulation of P status

PHO2, a ubiquitin-conjugating E2 enzyme, is a key regulator of Pi homeostasis
suppressing Pi uptake and translocation under P-replete conditions by targeting Pi
transporters such as PHO1 and PHT1 for ubiquitin-mediated degradation (Aung *et al.*,
2006; Bari *et al.*, 2006; Liu *et al.*, 2012). The E3 ubiquitin ligase NITROGEN
LIMITATION ADAPTATION (NLA) works alongside PHO2 to degrade Pi transporters
(Lin et al. 2013; Park et al. 2014), as well as PHR1 in response to changes in inositol
polyphosphate levels (Park *et al.*, 2023). Out of two PHO2 homologs in hemp, we
identified one (*LOC115698240*) with lower transcript abundance in P-limited LR and
YS (Figure 5, Supplementary Table S5). Lower transcript levels are consistent with
higher Pi transporter and PHR1 activity in these organs, as well as systemic, post-
transcriptional regulation of *PHO2* mRNA by miRNA399 observed in Arabidopsis roots
(Chiou *et al.*, 2005; Fujii *et al.*, 2005). However, in P-limited source organs (ML, OL),
PHO2 transcript abundance was 3- to 4-fold higher than under P-replete conditions
(Figure 5, Supplementary Table S5). This may indicate an additional role of PHO2 in

suppressing Pi import into these organs. Like *PHO2*, transcripts encoding the two

hemp *NLA* isoforms showed a 3- to 7-fold suppression in P-limited roots, while they

were largely unresponsive to Pi-supply in shoots, with one isoform (*LOC115695922*)

showing higher constitutive expression across organs.

In Arabidopsis and rice, transcript abundance of *PHR1*, the master regulator of

the PSR, is largely unresponsive to Pi supply and subject to post-translational

regulation by SPX proteins and / or ubiquitin-dependent turnover (Bustos *et al.*, 2010;

Nilsson *et al.*, 2007; Park *et al.*, 2023). The closest homolog to *AtPHR1*,

LOC115714052, was also not transcriptionally responsive to changes in Pi supply

(Supplementary Table S7). Two close *PHR1-LIKE* (*PHL*) homologs (*CsPHL13 /*

LOC115719175, *CsPHL1 / LOC115701526*) were identified in hemp. CsPHL13 had a

higher constitutive transcript abundance, especially in roots and leaves, and was

responsive to Pi supply with a two-fold induction in P-limited YS (Supplementary Table

S5 and S7). By contrast, CsPHL1 expression was very low across organs but showed

a 60-fold increased expression in roots supplied with 1 and 2 mM Pi (Figure 5,

Supplementary Table S7). Two out of the three *SPX* homologs (*LOC115725567 /*

CsSPX1, *LOC115697978 / CsSPX3*) were transcriptionally induced across P-limited

organs by about 30-fold (Figure 5), similar to corresponding homologs in Arabidopsis,

rice and tomato (Puga *et al.*, 2014; Satheesh *et al.*, 2022; Wang *et al.*, 2014). In TF,

expression of both *SPX* homologs was five- to seven-fold induced under complete Pi

withdrawal only, indicating Pi sufficiency in TF at a Pi supply of 0.25 mM (Figure 5,

Supplementary Table S5). Both genes are therefore sensitive indicators of plant P

status. Inositol polyphosphates (InsP) influence Pi signalling and homeostasis by

regulating SPX-PHR1 interactions (Collins *et al.*, 2024; Ried *et al.*, 2021). In hemp,

two putative INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE (ITPK) encoding

homologs (*LOC115695287*, *LOC115716459*) were identified, showing organ-specific

responses to Pi availability. Notably, transcripts encoding two out of four putative VIP

HOMOLOG1 (VIH1) and one VIH2 homolog in hemp showed constitutive expression

across organs and were only differentially expressed in P-starved OL (Supplementary

Table S5 and S7) (Dong *et al.,* 2019; Laha *et al.*, 2015). In Arabidopsis, ITPK1 and

ITPK2 function as key InsP6 kinases to produce inositol pyrophosphates $InsP_7$ and

$InsP_8$ in P-sufficient shoots (Riemer *et al.*, 2021). High $InsP_8$ levels stabilize the SPX-

PHR1 interaction to keep PHR1 in an inactive state, while Pi stress reduces shoot

$InsP_8$ concentration to disrupt SPX-PHR1 binding, thereby allowing PHR1 to bind

P1BS motifs and activate PSR genes. In hemp, transcriptional upregulation of the two

ITPK homologs in P-limited LR, ML, OL and YS (3- to 11- fold change) is counter-

intuitive as this should be associated with reduced PHR1 activity (Figure 5,

Supplementary Table S5) (Park *et al.*, 2023).

Mobilisation of Pi from organic compounds by Pi scavenging and lipid remodelling

P-limited plants mobilise Pi by the replacement of membrane phospholipids and

breakdown of organic compounds (e.g. phosphate esters), the latter by increasing the

activity of phosphatases (PAPs) and ribonucleases (RNSs) and their secretion into the

soil (Bariola *et al.*, 1994; Siebers *et al.*, 2015; Tran *et al.*, 2010). Homologs of several

genes involved in lipid remodelling were identified and their expression followed a Pi

supply dependent pattern across organ types, with some showing the strongest P-

limitation responses among all DEGs (Figure 5). Transcripts encoding a

phospholipase D (*PLD, LOC115718607*) homolog were induced 55- to 3,000-fold in

P-limited transport (R, YS) and source (ML, OL) organs, respectively (Supplementary

Table 5). The latter corroborates the importance of lipid remodelling in leaves (Li *et*

al., 2006). *GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE* (*GDPD)*

isoforms are involved in the lipid acyl hydrolase (LAH) pathway of phospholipid

hydrolysis and are upregulated under Pi limitation (Mehra *et al.*, 2019). *GDPD*

homologs encoded by *LOC115701510* and *LOC115708092* are responsive to Pi

supply across all organ types with the former showing more than 3,000-fold induction

in P-limited source organs (ML, OL) and YS (Figure 5) and attenuated suppression in

P-replete roots compared with other organs. Together with the two *CsSPX* isoforms,

LOC115701510 is thus another sensitive expression marker of P status in Cannabis.

PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE PHOSPHATASE1 (*PECP1*)

homologs (*LOC115706703, LOC115704970, LOC115704930, LOC115704928*)

encode haloacid dehydrogenase (HAD)-type phosphatases involved in cleaving of Pi

from phospholipid polar head groups (Hanchi *et al.*, 2018; Tannert *et al.*, 2018). All

three show upregulation under limiting Pi supply across organ types, with strongest

induction observed for LOC115704970 and *LOC115704930* in leaves.

LOC115706703 showed a limited transcriptional response to Pi supply but had a

higher basal level of expression across organ types, indicating a role in ongoing

membrane lipid turnover (Mehra *et al.*, 2019).

The protection of sink organs against phospholipid depletion under low Pi
supply in hemp is illustrated by the expression patterns of
PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE (PEAMT) encoding
homologs *LOC115709566* and *LOC115709567*. In *Arabidopsis*, two PEAMT isoforms
are involved in *de novo* phospholipid synthesis (Chen *et al.*, 2018). In hemp,
transcripts of both *PEAMT* homologs are downregulated by up to 1,200-fold in P-
limited sink organs (ML, OL) and YS (Figure 5; Supplementary Table S5). Yet in TF,
where Pi is preferentially translocated to (Figure 3A, 3B), expression is largely
unaffected, especially for *LOC115709566* which shows a very high transcript
abundance in both TF and YS irrespective of external Pi supply. Interestingly, this is
opposite to what is observed in Arabidopsis*,* where *PEAMT* (*NMT1 / PMT1*)
expression is upregulated in both roots and shoots to maintain phospholipid synthesis
under Pi starvation (Ngo *et al.*, 2022). In aboveground organs of Cannabis,
phospholipid synthesis via *PEAMT* is supported only in the presence of sufficient Pi.
In P-limited hemp organs where *de novo* phospholipid synthesis via *PEAMT* is limited
and phospholipid hydrolysis (*PLD*, *PECP*, *GDPD*) is induced, genes involved in
alternative lipid synthesis are also activated. Galactolipid and sulfolipid synthesizing
enzymes encoded by *MONOGALACTOSYL DIACYLGLYCEROL SYNTHASE*
(*MGD*), DIGALCTOSYLDIACYLGLYCEROL SYNTHASE (*DGD*) and
SULFOQUINOVOSYLDIACYLGLYCEROL SYNTHASE (*SQD*) genes, respectively,
exhibited lower transcriptional activity in sink organs, where phospholipid hydrolysis
was similarly constrained (Figure 5). By contrast, roots and leaves showed increased
transcriptional activity with a 3- to 720-fold induction of select homologs under limiting
Pi supply. Like Arabidopsis, transcripts encoding for one *DGD* homolog
(*LOC115706653*) were largely unresponsive to Pi supply, except for P-starved OL –
and like AtDGD1*,* it probably contributes to constitutive galactolipid synthesis in
chloroplasts (Kobayashi *et al.*, 2014; Kobayashi *et al.*, 2013), while transcripts of two
other homologs (CsMGD2, *LOC115695927*; CsDGD2, *LOC115698658*) are highly Pi
responsive in roots and leaves, just like *AtMGD2* and *AtDGD2* (Kelly and Dörmann,
2002; Kobayashi *et al.*, 2009). Surprisingly, transcripts encoding the type A MGD
synthase CsMGD1 (*LOC115707728*) are induced in P-limited source organs, with a
more constitutive expression in YS and TF. This reminiscent of the simultaneous
induction of both type A and type B MGD synthase genes observed in sesame plants
which are highly tolerant to P impoverished soils (Shimojima *et al.*, 2013). In contrast

to the Arabidopsis genes *AtSQD1* and *AtSQD2*, only one *SQD* homolog (*CsSQD2*,

LOC115697608) was identified in hemp with strong 14- to 30-fold induction in P-limited

roots and leaves, respectively. From these expression profiles, *CsSQD2* and *CsMGD2*

are another pair of sensitive markers of organ P status (Figure 5, Supplementary Table

S5).

Pi can also be (re-)mobilised from organic P compounds such as

phosphoesters and RNA catalysed by phosphatases (PAP) and ribonucleases (RNS).

In Arabidopsis, class I ribonucleases encoding *RNS1* and *RNS2* are upregulated

under Pi starvation (Gho *et al.*, 2020; Taylor *et al.*, 1993; Tran *et al.*, 2010). In hemp,

two out of six *RNS* homologs were differentially expressed (Figure 5, Supplementary

Table S5). Based on their expression profile in hemp, one of the homologs identified,

CsRNS3 (*LOC115698752*), is more similar to Arabidopsis *RNS1*, with strong induction

in R, OL and ML, while *CsRNS1* (*LOC115698919*) is more like *AtRNS2* as transcripts

are most abundant in P-limited leaves (Bariola *et al.*, 1994; Gho *et al.*, 2020; Taylor *et*

al., 1993). Both isoforms exhibit low transcriptional activity in TF, while in YS, a three-

to four-fold induction was evident upon Pi withdrawal (Figure 5, Supplementary Table

S5). Purple acid phosphatases (PAP*)* are upregulated under Pi stress and play a role

in Pi release from intracellular and / or external organic P pools (Bhadouria and Giri,

2022). The differentially expressed 15 *PAP* homologs identified in hemp have a similar

expression profile to the *RNS* isoforms (Figure 5), with some showing a strong PSR

across roots and leaves, while others are more responsive in P-limited leaves. Three

PAP genes, *CsPAP3 / LOC115723670*, *CsPAP8 / LOC115723241*, and *CsPAP17 /*

LOC115709102 are good indicators of Pi stress with up to 200-fold induction.

Altered Pi supply impacts N requirements which is reflected in transcriptional

adjustment of nitrate signalling

Pi limitation impacts the homeostasis of other nutrients, especially N, with

coordinated regulation at the transcriptional level (Hu and Chu, 2020; Krouk and Kiba,

2020; Raven, 2015). In our study, nitrate accumulated significantly in leaves and

young stems following Pi withdrawal (Figure 3B). This is unusual given that – in most

plants studied - nitrate uptake by roots and its translocation to the shoot are

suppressed under Pi limiting conditions (de Magalhães *et al.*, 1998; Gniazdowska and

Rychter, 2001; Jeschke *et al.*, 1997; Krouk and Kiba, 2020; Maeda *et al.*, 2018; Ueda

and Yanagisawa, 2023; Wang *et al.*, 2020). Total N was significantly increased in P-

starved LR and OL only (Supplementary Figure S1), suggesting redirection of N
resources to support the observed increase in root-to-shoot ratio (Figure 2B). Under
sufficient Pi supply, nitrate concentrations were unexpectedly low, indicative of
preferential assimilation into the organic fraction, and likely becoming limiting at higher
Pi supplies (Figure 3B, Supplementary Figure S1).

We next examined the expression of N regulatory genes as well as of genes
encoding steps central to nitrogen uptake and assimilation. An increased demand for
N with Pi supply was particularly evident in LR, where expression of *LOC115708988*,
one of two *NITRATE TRANSPORTER 3* (*NRT3*) genes encoding a key component of
high-affinity nitrate uptake (Okamoto *et al.*, 2006), was steadily induced from 3- to 6-
fold as Pi supply was raised from 0.25 mM to 2 mM (Figure 6, Supplementary Table
S5 and S6). Transcript profiles of three out of four hemp homologs encoding MADS
box transcription factor *ARABIDOPSIS NITRATE REGULATED 1* (*ANR1*) are
matching those of *NRT3.1 / LOC115708988* showing root-specific induction with Pi
supply (Zhang and Forde, 1998). This induction of nitrate uptake is most likely brought
about to match the increased capacity for Pi assimilation and consequently growth
(Figure 1). Two-fold induction of a *NITRATE TRANSPORTER 1 (NRT1)/PEPTIDE
TRANSPORTER* (*NPF*) homolog (*NRT1.2, LOC115707825*) in YS with Pi supply
points to active transport of nitrate to floral sink organs (Figure 6, Supplementary Table
S5 and S7). Increased expression of *NITRATE REDUCTASE* (*NIA, LOC115713111*)
and *GLUTAMINE SYNTHETASE2* (*GLN2*, LOC115722196) in leaves indicates
upregulation of the primary nitrate assimilation pathway to support new growth when
P is adequate, which in turn depletes nitrate levels. Transcripts encoding NPF
transporter homologs *LOC115699885* (*NPF4.5*, root), *LOC115700848* (*NPF6.4*, leaf),
LOC115709842 (NPF2.9, leaf), LOC115710913 (NPF7.3, leaf) were more abundant
in P-limited organs. Two nitrite reductase encoding *CsNIR* isoforms (*LOC115702926*,
LOC115702931), the alpha-subunit of GLUTAMATE DEHYDROGENASE1 encoding
CsGDH1 (*LOC115705444*), and one out of two mitochondrial aspartate
aminotransferases encoding loci (*AAT1*, *LOC115699642*) had predominant
expression in source organs under Pi withdrawal when nitrate concentrations were
high. Reduced suppression of *CsGDH1* in source organs at the highest Pi supply level
could be a response to nitrate depletion and increased N metabolic turnover under
excess Pi supply. GDH isoforms catalyse both the synthesis and catabolism of
glutamate, allowing the (re-)assimilation of ammonia under P or C starvation

conditions as well as the generation of N-storage compounds such as N-rich amino acids and polyamines, particularly under high nitrate availability and / or a lack of assimilatory capacity (Forde and Lea, 2007). Contrasts in *CsGLN2* and *CsGDH1* transcript abundance are particularly useful in determining what constitutes a balanced supply of Pi and N for hemp cultivation.

Stimulation of nitrate uptake by high P status in hemp is also evident from the transcriptional profile of *NITRATE-INDUCIBLE, GARP-TYPE TRANSCRIPTIONAL REPRESSOR1* (*NIGT1*, *LOC115700564*) which suppresses high affinity nitrate acquisition and is also linked to stimulating PSR in Arabidopsis (Medici *et al.*, 2015; Ueda *et al.*, 2020a; Ueda *et al.*, 2020b; Wang *et al.*, 2020). AtNIGT1 shows antagonistic regulation of nitrate transporter gene expression compared with NIN-LIKE PROTEIN (NLP) transcription factors (Maeda *et al.*, 2018). In hemp, *NIGT1* transcript abundance is very low in P-replete organs, particularly in source leaves (ML, OL), while upon Pi withdrawal its expression is upregulated by 8- to 170-fold in ML, OL, and LR, consistent with its role in suppressing nitrate uptake (Figures 3B and 6). In TF and YS, weaker (4.3- to 8-fold) *NIGT1* induction is observed only with Pi withdrawal. As reported in Arabidopsis, *LOC115702149* encoding a NLP6 homolog in hemp shows the opposite expression profile to *NIGT1* (Maeda *et al.*, 2018; Okamoto *et al.*, 2018). Overall, the upregulation of *CsANR1*, *CsNLP6*, and *CsNRT3.1* in P-replete roots, together with *CsNIA*, and *CsGLN2* expression in P-replete shoots is matched by suppression of *CsNIGT1* and *CsGDH1* across those organs. This suggests that in hemp, nitrate uptake and N assimilation are highly responsive to Pi availability. In model organisms such as Arabidopsis, nitrate uptake is likewise counteracted by PHR1-mediated induction of *NIGT1* upon Pi withdrawal, whilst excessive Pi uptake is counteracted by down-regulation of *NIGT1* and promotion of SPX sequestration of PHR1 and NLP (Maeda *et al.*, 2018; Ueda *et al.*, 2020a). In hemp, this seems not to be the case since Pi acquisition is not suppressed by low organ nitrate levels and one putative *PHR1* homolog, *CsPHL13*, even showing transcriptional upregulation with increasing Pi supply in nitrate-depleted source organs (Figure 5).

DISCUSSION

P utilization efficiency attributed to high Pi recycling and conservation in hemp

Developing optimized fertigation strategies and cultivars balancing nutrient
uptake and internal utilization capacity is necessary towards building high-
performance Cannabis germplasm (Cockson *et al.*, 2020; Hershkowitz *et al.*, 2025;
Heuer *et al.*, 2017; Shiponi and Bernstein, 2021b; Tripathi *et al.*, 2022). In our study,
the dual-purpose hemp cultivar showed a remarkable resilience to growth inhibition,
requiring complete Pi withdrawal for biomass reallocation away from shoot and
reproductive organs towards roots (Figure 2B).

The ability of hemp to grow on marginal lands allowed us to study traits for its
nutrient-use efficiency (Scalabrin *et al.*, 2024; Wee Y. *et al.*, 2024). This identified key
metabolic and transcriptomic acclimation responses in hemp that likely underpin the
high PUE. The PSR supports Pi being actively remobilized from roots and source
organs towards sink organs, such as flowers (Figure 5). Notably, the young stem
serves as a conduit for the efficient translocation of Pi under limiting conditions, with
expression closely mirroring that of flowers. These transcriptional changes align very
closely with shifts in Pi and Po metabolism across organ types (Figure 3A). Even under
limiting Pi supplyng (≤ 0.25 mM Pi), Po concentration in flowers was 70-fold higher
than in all other organ types, indicating their protection from nutrient stress as a
hallmark of high PUE in hemp. Similar trends have been observed in other sink-heavy
crops such as wheat and rice, where reproductive organs accumulate higher P levels
to support grain development in genotypes with high PUE (Adem *et al.*, 2020; Aziz *et*
al., 2014; Hayes *et al.*, 2022; Irfan *et al.*, 2020; Julia *et al.*, 2016; Suriyagoda *et al.*,
2023; Veneklaas *et al.*, 2012).

To maintain central processes such as ATP production, nitrogen and carbon
assimilation under P limitation, it is essential that alternative pathways are activated in
tandem with intracellular Pi-recycling mechanisms (Albinsky *et al.*, 2010; Carstensen
et al., 2018; O'Leary *et al.*, 2011). Under Pi stress, we observed a preferential increase
in sulfate accumulation in shoot organs. This is in contrast to other plants such as
Arabidopsis where sulfate concentrations decrease in shoot organs while increasing
in roots (Rouached, 2011). This increased sulfate concentration in shoots under Pi-
stress is indicative of how the hemp cultivar complements ongoing Pi reallocation from
source to sink organs by transitioning to Pi-free glycerolipids such as sulfo- and

galactolipids (Hanchi *et al.*, 2018; Nakamura, 2013; Shimojima *et al.*, 2013; Tjellstrom

et al., 2008). As opposed to Arabidopsis where the rate-limiting phospholipid

biosynthesis gene *PEAMT* is upregulated under Pi-stress, its expression was

positively correlated with Pi supply in hemp (Figure 5) (Chen *et al.*, 2018). Considering

that Po levels are maintained even under Pi-limitation in floral organs but not in source

organs (Figure 3A), high *PEAMT* transcript abundance irrespective of Pi supply implies

de novo phospholipid synthesis is supported by preferential Pi translocation to

reproductive organs (Aziz *et al.*, 2014). In source organs (ML, OL) where Pi is limiting

under Pi stress, PEAMT-mediated phospholipid synthesis is compromised, and

expression of genes encoding alternative lipid synthesis dominates. The latter is

supported by the decrease in sulfate concentration (Figure 3B) relative to reproductive

organs at the same level of Pi supply (Figure 5).

Collectively, these two complementary acclimation responses, i.e. enhanced Pi

allocation to reproductive sinks and heightened release and turnover of organic P

pools through lipid remodelling and phosphatase / nuclease activities, likely underpin

the key adaptations that contribute to the high PUE and sustained floral production

under Pi-stress in hemp.

Nutrient cross-talk and regulatory implications

This study also revealed that hemp adopts a distinctive strategy for balancing

P and N nutrition, differing significantly from canonical models established in, e.g.,

Arabidopsis and rice. Our results suggest that hemp does not down-regulate Pi uptake

and hyperaccumulates Pi in roots and aboveground organs, a trait that is atypical and

suggests an alternative mechanism for Pi sensing and translocation that is reminiscent

of plants adapted to P impoverished soils like the Australian *Proteacea* and some

legumes (Pang *et al.*, 2009; Prodhan *et al.*, 2016; Shane *et al.*, 2004a; Shane *et al.*,

2004b). This result is supported by other studies in hemp and drug-type Cannabis

(Cockson *et al.*, 2020; Jost *et al.*, 2024; Shiponi and Bernstein, 2021b; Wee Y. *et al.*,

2024). The abundance of transcripts encoding the Pi exporter responsible for Pi

loading into the xylem, PHO1, supports this observed difference in Pi uptake regulation

under excess supply (Figure 5). The dominant *PHO1* gene remains highly expressed

in roots and young stems even under high Pi supply, underscoring the strong sink

strength of reproductive organs (TF) that drives continuous Pi uptake in roots (Dai *et*

al., 2024; Wang *et al.*, 2004). Continued Pi accumulation in roots and stems is in spite

of *CsPHO2* upregulation under high Pi supply, which in other plants leads to PHO1

degradation to decrease Pi translocation to the shoots - pointing towards a decoupling

of the PHO2-PHO1 regulatory axis in hemp (Ouyang *et al.*, 2016). Given that *PHO1*

regulates floral transition in both *Arabidopsis* and rice, further investigation into its

potential role as a Pi-dependent modulator of flowering time in hemp and drug-type

Cannabis is warranted – particularly in cultivars that exhibit elevated *PHO1* expression

(Dai *et al.*, 2024).

Sustained organ Pi accumulation also raises the important question whether

excess Pi uptake in hemp is the result of N limitation as indicated by depleted nitrate

storage reserves under Pi-sufficient conditions (Figure 3B). In many plant species,

efficient biomass production relies on coordinated nitrate and Pi acquisition, as both

are essential for nucleotide and protein biosynthesis (Medici *et al.*, 2019; Pueyo *et al.*,

2021; Raven, 2015). Ample supply of nitrate is essential towards activation of PSR

genes, whereas nitrate limitation strongly inhibits induction and leads to Pi

accumulation along with downregulation of typical PSR genes in crops such as maize,

and Arabidopsis (Kant *et al.*, 2011; Maeda *et al.*, 2018; Medici *et al.*, 2015; Schluter *et*

al., 2012; Torres-Rodriguez *et al.*, 2021; Wang *et al.*, 2020). In the hemp cultivar, Pi

and Po status were negatively correlated with nitrate (Figure 4), despite upregulation

of key high affinity nitrate transport system (HATS) component *CsNRT3.1* and

corresponding downregulation of its negative GARP-type repressor *CsNIGT1* with

increasing Pi supply (Figure 6). Increased N demand with Pi supply is reflected in the

upregulation of key nitrate assimilation genes (*NIA, GLN1*) (Figure 6) (Liu *et al.*, 2024;

Marschner, 2012; Wang *et al.*, 2002). Yet, remarkably, total N and total C remained

unresponsive to Pi supply across aboveground sink organs (Figure 3, Supplementary

Figure S1), while the total P and Po fractions – increased with Pi supply (Figure 3).

These results suggest that in hemp an increase in Pi supply needs to be

carefully balanced with a corresponding higher input of nitrate – as well as

photosynthetic activity, to promote synergistic interactions between N and P to support

higher stem and / or flower biomass and ultimately target product yield. The observed

decoupling of key transcriptional circuits such as PHR1-driven PSR and the PHO2-

NLA regulon in hemp, alongside its inverted Pi–N coordination, extends current

models of nutrient regulation obtained in other species.

Deeper understanding of genotype and usage-type specific nutrient dynamics,

helped by the expression profiles of indicative genes identified in our study, has the

potential to predict their nutrient use characteristics, enhance cannabinoid production,

and support the development of more sustainable cultivation strategies tailored to

individual Cannabis germplasm.

SUPPLEMENTARY DATA

Supplementary Figure 1. Total carbon and nitrogen concentrations in response to Pi supply. Shown are total carbon and nitrogen concentrations across organs and Pi treatments (mean ± SE, n = 3 plants per treatment). Letters denote statistical significance between Pi treatments (ANOVA, Tukey HSD, $p < 0.05$).

Supplementary Figure 2. Analysis of differentially expressed genes in OL across Pi treatments. Upset plot for differentially expressed genes (DEGs) shared between P-starved organs (**A**) and Venn diagram of DEGs in OL across treatments (**B**) visualise the disproportionate number of DEGs in the OL samples after withdrawal of Pi (P0 treatment) due to starvation induced premature senescence. (**C**) Gene ontology (GO) term enrichment analysis for the DEGs specific for the OL of P-starved plants highlights senescence- and proteolysis-related terms in upregulated DEGs, while downregulated DEGs are related to photosynthesis as expected of leaves in the later stages of senescence.

Supplementary Table 1: Summary of recorded morphological and physiological data. Sample means and standard errors (SE) were calculated for each variable across the three biological replicates (n) across five treatment groups (0, 0.25, 0.5, 1.0 and 2.0 mM Pi). The residual degrees of freedom (df) from one-way ANOVA was used to calculate the upper and lower 95 % confidence limits (CLs) for each variable's pairwise difference in Tukey's HSD test ($p < 0.05$). To summarize pairwise comparisons from Tukey's HSD, compact letter displays (group) were assigned using the *cld* function from package 'multcompView' to generate letter codes, where alpha = 0.5 was used to separate significantly different treatments. df = N-p, where N is the total number of observations (15) and p is the number of parameters estimated (five treatment levels). ADI = Apical dominance index, *i.e.*, number of ramifications per centimetre of representative branch, APE = agronomic phosphate efficiency, DW = dry weight, FW = fresh weight, Internodal Spacing = average distance between nodes along the main stem, SHI = shoot harvest index, S:R = shoot-to-root biomass ratio, Veg = vegetative biomass. Percent scores represent the biomass (FW or DW) of a given organ type as a % of total biomass.

Supplementary Table 2: **Detailed summary of statistical outputs from Tukey's HSD following one-way ANOVA for recorded morphological and physiological data.** Per variable, treatment pairwise contrasts are given, supplying estimates (difference in group means, *i.e.*, first treatment minus second), pooled standard errors (SE(pooled)) for recorded morphological and physiological data from emmeans Tukey output (pooled, df (residual degrees of freedom), t-statistic (estimate divided by SEpooled)), and adjusted p-values (family-wise error-controlled probabilities from Tukey's HSD for each pairwise contrast). A pooled SE is used because the design is balanced and an assumption of equal variances is assumed. ADI = Apical dominance index, *i.e.*, number of ramifications per centimeter of representative branch, APE = agronomic phosphate efficiency, DW = dry weight, FW = fresh weight, Internodal Spacing = average distance between nodes along the main stem, SHI = shoot harvest index, S:R = shoot-to-root biomass ratio, Veg = vegetative biomass. Percent scores represent the biomass (FW or DW) of a given organ type as a % of total biomass.

Supplementary Table 3: Summary of assayed nutrients and metabolites alongside statistical analysis by ANOVA followed by Tukey's pairwise comparisons, across hemp organs at final destructive harvest. The residual degrees of freedom (df) from one-way ANOVA was used to calculate the upper and lower 95 % confidence limits (CLs) for each variable's pairwise difference in Tukey's HSD test (p < 0.05). To summarize pairwise comparisons from Tukey's HSD, compact letter displays (group) were assigned using the *cld* function from package 'multcompView' to generate letter codes, where at alpha=0.05 to separate significantly different treatments. df= N-p, where N is the total number of observations (15) and p is the number of treatment levels (five). C-N ratio = total carbon to nitrogen ratio, organic P= organic phosphorus.

Supplementary Table 4: **Detailed summary of statistical outputs from Tukey's HSD following one-way ANOVA for assayed metabolite data, across hemp organs at final destructive harvest.** Per variable, pairwise treatment (mM Pi) contrasts are given, supplying estimates (difference in group means, *i.e.*, first treatment minus second), pooled standard errors (SE(pooled)) for recorded morphological and physiological data from emmeans Tukey output, df (residual degrees of freedom), t-statistic (estimate divided by SEpooled)), and adjusted p-values

(family-wise error-controlled probabilities from Tukey's HSD for each pairwise
contrast). A pooled SE is used because the design is balanced and an assumption of
equal variances is assumed. C-N ratio = total carbon to nitrogen ratio, organic P =
organic phosphorus.

Supplementary Table 5: Summary of DEGs across organs and Pi treatments in
hemp. Gene IDs and descriptions are derived from the cs10 genome annotation.
Selection criteria applied were as follows: FDR adj. p-value < 0.05, $|\log_2 FC| > 1$, TPM >
10 in at least 1 organ type across treatments. Values are provided as $\log_2$ fold-change
(logFC) relative to organs of plant treated with 1 mM phosphate (control). Feature =
RNA species (m = messenger, nc = non-coding, r = ribosomal, misc = miscellaneous),
FDR = false discovery rate with 'NA' indicating lack of statistical significance. "gene
family" highlights gene families of special interest, "Pathway annotation" refers to
custom annotation.. NA = not available, TF = top flower / terminal inflorescence, ML =
mature fan leaf, OL = old fan leaf, YL = young fan leaf, YS = young stem, LR = lateral
root. P0 = no phosphate supply, P025 = 0.25 mM phosphate supply, P2 = 2 mM
phosphate supply.

Supplementary Table 6: Curated list of PSR and other nutrient-response genes
based on known function in other species. Gene IDs and descriptions are derived
from the cs10 genome annotation. Gene annotations have been verified against
Arabidopsis homologs, selecting for the lowest E score (cut-off E<10^-50), and
corresponding gene families have also been provided after curation.

Supplementary Table 7: Transcripts per million (TPM) values across treatment
and organ types. Gene IDs and gene annotations are derived from the cs10 genome
annotation. Accession numbers are NCBI RefSeq mRNA transcript identifiers for each
predicted transcript model in the cs10 genome annotation. Feature = RNA species (m
= messenger, nc = non-coding, r = ribosomal, misc = miscellaneous), TPM =
transcripts per million, TF = top flower / terminal inflorescence; ML = mature fan leaf,
OL = old fan leaf, YL = young fan leaf, YS = young stem, LR = lateral root. P0 = no
phosphate supply, P025 = 0.25 mM phosphate supply, P2 = 2 mM phosphate supply.

ACKNOWLEDGEMENTS

The authors would like to thank Stefanie Kabitz, Filippa Brugliera (both Cann
Group Limited) and colleagues within the ARC Industrial Transformation Research
Hub for Medicinal Agriculture for useful discussions and feedback.

AUTHOR CONTRIBUTIONS

RJ, OB, JW and BWY conceived the project. BWY conducted the experiment,
performed the various assays outlined in the methods as well as data analyses and
visualisation. BWY, OB and RJ analysed the data. AP and SN assisted with
experimental design and data acquisition. BWY, RJ and OB wrote the manuscript with
revision by all authors.

CONFLICT OF INTEREST

The authors declare no conflicting interests.

FUNDING

This work was supported by the ARC Industrial Transformation Research Hub
for Medicinal Agriculture (JW, grant ID IH180100006).

DATA AVAILABILITY

RNA-seq read data were deposited at the NCBI SRA database under project
ID PRJNA1278883.

REFERENCES

Adem GD, Ueda Y, Hayes PE, Wissuwa M. 2020. Genetic and physiological traits for internal phosphorus utilization efficiency in rice. PLoS One **15**, e0241842.

Albinsky D, Kusano M, Higuchi M, Hayashi N, Kobayashi M, Fukushima A, Mori M, Ichikawa T, Matsui K, Kuroda H, Horii Y, Tsumoto Y, Sakakibara H, Hirochika H, Matsui M, Saito K. 2010. Metabolomic screening applied to rice FOX Arabidopsis lines leads to the identification of a gene-changing nitrogen metabolism. Mol Plant **3**, 125-142.

Ames B. 1966. Methods in enzymology. Bd. VIII Complex Carbohydrates: Assay of Inorganic Phosphate, Total Phosphate and Phosphatases.

Arpat AB, Magliano P, Wege S, Rouached H, Stefanovic A, Poirier Y. 2012. Functional expression of PHO1 to the Golgi and trans-Golgi network and its role in export of inorganic phosphate. The Plant Journal **71**, 479-491.

Aung K, Lin SI, Wu CC, Huang YT, Su CL, Chiou TJ. 2006. pho2, a phosphate overaccumulator, is caused by a nonsense mutation in a microRNA399 target gene. Plant Physiol **141**, 1000-1011.

Aziz T, Finnegan PM, Lambers H, Jost R. 2014. Organ-specific phosphorus-allocation patterns and transcript profiles linked to phosphorus efficiency in two contrasting wheat genotypes. Plant, Cell & Environment **37**, 943-960.

Bari R, Datt Pant B, Stitt M, Scheible WR. 2006. PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants. Plant Physiol **141**, 988-999.

Bariola PA, Howard CJ, Taylor CB, Verburg MT, Jaglan VD, Green PJ. 1994. The Arabidopsis ribonuclease gene RNS1 is tightly controlled in response to phosphate limitation. Plant J **6**, 673-685.

Bayle V, Arrighi J-F, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E, Paz-Ares J, Nussaume L. 2011. Arabidopsis thaliana High-Affinity Phosphate Transporters Exhibit Multiple Levels of Posttranslational Regulation The Plant Cell **23**, 1523-1535.

Bhadouria J, Giri J. 2022. Purple acid phosphatases: roles in phosphate utilization and new emerging functions. Plant Cell Rep **41**, 33-51.

Blake-Kalff MMA, Hawkesford MJ, Zhao FJ, McGrath SP. 2000. Diagnosing sulfur deficiency in field-grown oilseed rape (Brassica napus L.) and wheat (Triticum aestivum L.). Plant and Soil **225**, 95-107.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol **34**, 525-527.

Burgarella C, Barnaud A, Kane NA, Jankowski F, Scarcelli N, Billot C, Vigouroux Y, Berthouly-Salazar C. 2019. Adaptive Introgression: An Untapped Evolutionary Mechanism for Crop Adaptation. Front Plant Sci **10**, 4.

Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Perez-Perez J, Solano R, Leyva A, Paz-Ares J. 2010. A central regulatory system largely controls transcriptional activation and repression responses to phosphate starvation in Arabidopsis. PLoS Genet **6**, e1001102.

Carstensen A, Herdean A, Schmidt SB, Sharma A, Spetea C, Pribil M, Husted S. 2018. The Impacts of Phosphorus Deficiency on the Photosynthetic Electron Transport Chain. Plant Physiol **177**, 271-284.

Chen W, Salari H, Taylor MC, Jost R, Berkowitz O, Barrow R, Qiu D, Branco R, Masle J. 2018. NMT1 and NMT3 N-Methyltransferase Activity Is Critical to Lipid Homeostasis, Morphogenesis, and Reproduction. Plant Physiol **177**, 1605-1628.

Chen YF, Li LQ, Xu Q, Kong YH, Wang H, Wu WH. 2009. The WRKY6 transcription factor modulates PHOSPHATE1 expression in response to low Pi stress in Arabidopsis. Plant Cell **21**, 3554-3566.

Chiou T-J, Aung K, Lin S-I, Wu C-C, Chiang S-F, Su C-I. 2005. Regulation of Phosphate Homeostasis by MicroRNA in Arabidopsis. The Plant Cell **18**, 412-421.

Cho H, Choi I, Bouain N, Nawaz A, Zheng L, Shahzad Z, Brandizzi F, Rhee SY, Rouached H. 2025. Phosphorus Availability Modulates Flowering Time Through Subcellular Reprogramming of bGLU25 and GRP7 in Flowering Plants. bioRxiv, 2025.2001.2002.631137.

Clarke RC. 2020. Tasmania Hemp (Cannabis) Fiber and Seed Cultivar Field Trials—2018-2019. International Journal of Food Science and Agriculture **4**, 470-481.

Clarkson DT, Hawkesford MJ, Davidian J-C, Grignon C. 1992. Contrasting responses of sulphate and phosphate transport in barley (Hordeum vulgare L.) roots to protein-modifying reagents and inhibition of protein synthesis. Planta **187**, 306-314.

Cockson P, Landis H, Smith T, Hicks K, Whipker BE. 2019. Characterization of Nutrient Disorders of Cannabis sativa. Applied Sciences **9**, 4432.

Cockson P, Schroeder-Moreno M, Veazie P, Barajas G, Logan D, Davis M, Whipker BE. 2020. Impact of Phosphorus on Cannabis sativa Reproduction, Cannabinoids, and Terpenes. Applied Sciences **10**, 1-18.

Collins E, Shou H, Mao C, Whelan J, Jost R. 2024. Dynamic interactions between SPX proteins, the ubiquitination machinery, and signalling molecules for stress adaptation at a whole-plant level. Biochem J **481**, 363-385.

Coutinho J. 1996. Automated method for sulphate determination in soil-plant extracts and waters. Communications in Soil Science and Plant Analysis **27**, 727-740.

Cuyas L, David P, de Craieye D, Ng S, Arkoun M, Plassard C, Faharidine M, Hourcade D, Degan F, Pluchon S, Nussaume L. 2023. Identification and interest of molecular markers to monitor plant Pi status. BMC Plant Biol **23**, 401.

Dai S, Chen H, Shi Y, Xiao X, Xu L, Qin C, Zhu Y, Yi K, Lei M, Zeng H. 2024. PHOSPHATE1-mediated phosphate translocation from roots to shoots regulates floral transition in plants. J Exp Bot **75**, 5054-5075.

de Jager N, Shukla V, Koprivova A, Lyčka M, Bilalli L, You Y, Zeier J, Kopriva S, Ristova D. 2023. Traits linked to natural variation of sulfur content in Arabidopsis thaliana. Journal of Experimental Botany **75**, 1036-1050.

de Magalhães JV, Alves VMC, de Novais RF, Mosquim PR, Magalhães JR, Filho AFCB, Hubert DM. 1998. Nitrate uptake by corn under increasing periods of phosphorus starvation. Journal of Plant Nutrition **21**, 1753-1763.

Devau N, Hinsinger P, Le Cadre E, Gérard F. 2011. Root-induced processes controlling phosphate availability in soils with contrasted P-fertilized treatments. Plant and Soil **348**, 203-218.

Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang TE, Wittwer C, Jessen HJ, Zhang H, An GY, Chao DY, Liu D, Lei M. 2019. Inositol Pyrophosphate InsP(8) Acts as an Intracellular Phosphate Signal in Arabidopsis. Mol Plant **12**, 1463-1473.

Dubousset L, Abdallah M, Desfeux AS, Etienne P, Meuriot F, Hawkesford MJ, Gombert J, Segura R, Bataille MP, Reze S, Bonnefoy J, Ameline AF, Ourry A, Le Dily F, Avice JC. 2009. Remobilization of leaf S compounds and senescence in response to restricted sulphate supply during the vegetative stage of oilseed rape are affected by mineral N availability. J Exp Bot **60**, 3239-3253.

Forde BG, Lea PJ. 2007. Glutamate in plants: metabolism, regulation, and signalling. J Exp Bot **58**, 2339-2358.

Foroughi S, Baker AJM, Roessner U, Johnson AAT, Bacic A, Callahan DL. 2014. Hyperaccumulation of zinc by Noccaea caerulescens results in a cascade of stress responses and changes in the elemental profile. Metallomics **6**, 1671-1682.

Fujii H, Chiou T-J, Lin S-I, Aung K, Zhu J-K. 2005. A miRNA Involved in Phosphate-Starvation Response in Arabidopsis. Current Biology **15**, 2038-2043.

Gho YS, Choi H, Moon S, Song MY, Park HE, Kim DH, Ha SH, Jung KH. 2020. Phosphate-Starvation-Inducible S-Like RNase Genes in Rice Are Involved in Phosphate Source Recycling by RNA Decay. Front Plant Sci **11**, 585561.

Gniazdowska A, Rychter AM. 2001. Nitrate uptake by bean (Phaseolus vulgaris L.) roots under phosphate deficiency. In: Gašparíková O, Čiamporová M, Mistrík I, Baluška F, eds. *Recent Advances of Plant Root Structure and Function: Proceedings of the 5th International Symposium on Structure and Function of Roots. Stará Lensná, Slovakia, 30 August – 4 September, 1998*. Dordrecht: Springer Netherlands, 111-117.

Grassa CJ, Weiblen GD, Wenger JP, Dabney C, Poplawski SG, Timothy Motley S, Michael TP, Schwartz CJ. 2021. A new Cannabis genome assembly associates elevated cannabidiol (CBD) with hemp introgressed into marijuana. New Phytol **230**, 1665-1679.

Graves S, Piepho H-P, Selzer ML. 2015. Package 'multcompView'. Visualizations of paired comparisons **451**, 452.

Gu M, Chen A, Sun S, Xu G. 2016. Complex Regulation of Plant Phosphate Transporters and the Gap between Molecular Mechanisms and Practical Application: What Is Missing? Mol Plant **9**, 396-416.

Gu Z. 2022. Complex heatmap visualization. Imeta **1**, e43.

Guo M, Ruan W, Li C, Huang F, Zeng M, Liu Y, Yu Y, Ding X, Wu Y, Wu Z, Mao C, Yi K, Wu P, Mo X. 2015. Integrative Comparison of the Role of the PHOSPHATE RESPONSE1 Subfamily in Phosphate Signaling and Homeostasis in Rice. Plant Physiol **168**, 1762-1776.

Hachiya T, Okamoto Y. 2017. Simple Spectroscopic Determination of Nitrate, Nitrite, and Ammonium in Arabidopsis thaliana. Bio Protoc **7**, e2280.

Hamburger D, Rezzonico E, MacDonald-Comber Petetot J, Somerville C, Poirier Y. 2002. Identification and characterization of the Arabidopsis PHO1 gene involved in phosphate loading to the xylem. Plant Cell **14**, 889-902.

Hanchi M, Thibaud MC, Legeret B, Kuwata K, Pochon N, Beisson F, Cao A, Cuyas L, David P, Doerner P, Ferjani A, Lai F, Li-Beisson Y, Mutterer J, Philibert M, Raghothama KG, Rivasseau C, Secco D, Whelan J, Nussaume L, Javot H. 2018. The Phosphate Fast-Responsive Genes PECP1 and PPsPase1 Affect Phosphocholine and Phosphoethanolamine Content. Plant Physiol **176**, 2943-2962.

Hayes PE, Adem GD, Pariasca-Tanaka J, Wissuwa M. 2022. Leaf phosphorus fractionation in rice to understand internal phosphorus-use efficiency. Ann Bot **129**, 287-302.

Hendriks JH, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose pyrophosphorylase is activated by posttranslational redox-modification in response to light and to sugars in leaves of Arabidopsis and other plant species. Plant Physiol **133**, 838-849.

Hershkowitz JA, Westmoreland FM, Bugbee B. 2025. Elevated root-zone P and nutrient concentration do not increase yield or cannabinoids in medical cannabis. Front Plant Sci **16**, 1433985.

Heuer S, Gaxiola R, Schilling R, Herrera-Estrella L, López-Arredondo D, Wissuwa M, Delhaize E, Rouached H. 2017. Improving phosphorus use efficiency: a complex trait with emerging opportunities. The Plant Journal **90**, 868-885.

Hirsch J, Marin E, Floriani M, Chiarenza S, Richaud P, Nussaume L, Thibaud MC. 2006. Phosphate deficiency promotes modification of iron distribution in Arabidopsis plants. Biochimie **88**, 1767-1771.

Hu B, Chu C. 2020. Nitrogen-phosphorus interplay: old story with molecular tale. New Phytol **225**, 1455-1460.

Irfan M, Aziz T, Maqsood MA, Bilal HM, Siddique KHM, Xu M. 2020. Phosphorus (P) use efficiency in rice is linked to tissue-specific biomass and P allocation patterns. Sci Rep **10**, 4278.

Jeschke WD, Kirkby EA, Peuke AD, Pate JS, Hartung W. 1997. Effects of P deficiency on assimilation and transport of nitrate and phosphate in intact plants of castor bean (Ricinus communis L.). Journal of Experimental Botany **48**, 75-91.

Jiang M, Sun L, Isupov MN, Littlechild JA, Wu X, Wang Q, Wang Q, Yang W, Wu Y. 2019. Structural basis for the Target DNA recognition and binding by the MYB domain of phosphate starvation response 1. FEBS J **286**, 2809-2821.

Jost R, Berkowitz O, Pegg A, Hurgobin B, Tamiru-Oli M, Welling MT, Deseo MA, Noorda H, Brugliera F, Lewsey MG, Doblin MS, Bacic AT, Whelan J. 2024. Sink strength, nutrient allocation, cannabinoid yield and associated transcript profiles vary in two drug-type Cannabis chemovars. Journal of Experimental Botany.

Jost R, Schilling S, Melzer R. 2025. Improving medicinal plant cultivation through in-depth understanding of environmental, physiological, metabolic, and genetic constraints. J Exp Bot **76**, 1-4.

Julia C, Wissuwa M, Kretzschmar T, Jeong K, Rose T. 2016. Phosphorus uptake, partitioning and redistribution during grain filling in rice. Ann Bot **118**, 1151-1162.

Kant S, Peng M, Rothstein SJ. 2011. Genetic regulation by NLA and microRNA827 for maintaining nitrate-dependent phosphate homeostasis in arabidopsis. PLoS Genet **7**, e1002021.

Kelly AA, Dormann P. 2002. DGD2, an arabidopsis gene encoding a UDP-galactose-dependent digalactosyldiacylglycerol synthase is expressed during growth under phosphate-limiting conditions. J Biol Chem **277**, 1166-1173.

Kelly AA, Dörmann P. 2002. DGD2, an Arabidopsis Gene Encoding a UDP-Galactose-dependent Digalactosyldiacylglycerol Synthase Is Expressed during Growth under Phosphate-limiting Conditions*. Journal of Biological Chemistry **277**, 1166-1173.

Kobayashi K, Awai K, Nakamura M, Nagatani A, Masuda T, Ohta H. 2009. Type-B monogalactosyldiacylglycerol synthases are involved in phosphate starvation-induced lipid remodeling, and are crucial for low-phosphate adaptation. Plant J **57**, 322-331.

Kobayashi K, Fujii S, Sasaki D, Baba S, Ohta H, Masuda T, Wada H. 2014. Transcriptional regulation of thylakoid galactolipid biosynthesis coordinated with chlorophyll biosynthesis during the development of chloroplasts in Arabidopsis. Frontiers in Plant Science **Volume 5 - 2014**.

Kobayashi K, Narise T, Sonoike K, Hashimoto H, Sato N, Kondo M, Nishimura M, Sato M, Toyooka K, Sugimoto K, Wada H, Masuda T, Ohta H. 2013. Role of galactolipid biosynthesis in coordinated development of photosynthetic complexes and thylakoid membranes during chloroplast biogenesis in Arabidopsis. Plant J **73**, 250-261.

Krouk G, Kiba T. 2020. Nitrogen and Phosphorus interactions in plants: from agronomic to physiological and molecular insights. Curr Opin Plant Biol **57**, 104-109.

Laha D, Johnen P, Azevedo C, Dynowski M, Weiß M, Capolicchio S, Mao H, Iven T, Steenbergen M, Freyer M, Gaugler P, de Campos MKF, Zheng N, Feussner I, Jessen HJ, Van Wees SCM, Saiardi A, Schaaf G. 2015. VIH2 Regulates the

Synthesis of Inositol Pyrophosphate InsP8 and Jasmonate-Dependent Defenses in Arabidopsis. The Plant Cell **27**, 1082-1097.

Lambers H. 2021. Phosphorus Acquisition and Utilization in Plants. Annu Rev Plant Biol.

Lambers H, Shane MW, Cramer MD, Pearse SJ, Veneklaas EJ. 2006. Root structure and functioning for efficient acquisition of phosphorus: Matching morphological and physiological traits. Ann Bot **98**, 693-713.

Li M, Qin C, Welti R, Wang X. 2006. Double knockouts of phospholipases Dzeta1 and Dzeta2 in Arabidopsis affect root elongation during phosphate-limited growth but do not affect root hair patterning. Plant Physiol **140**, 761-770.

Lichtenthaler HK. 1987. [34] Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. *Methods in Enzymology*, Vol. 148: Academic Press, 350-382.

Linn J, Ren M, Berkowitz O, Ding W, van der Merwe MJ, Whelan J, Jost R. 2017. Root Cell-Specific Regulators of Phosphate-Dependent Growth. Plant Physiol **174**, 1969-1989.

Liu J, Yang L, Luan M, Wang Y, Zhang C, Zhang B, Shi J, Zhao FG, Lan W, Luan S. 2015. A vacuolar phosphate transporter essential for phosphate homeostasis in Arabidopsis. Proc Natl Acad Sci U S A **112**, E6571-6578.

Liu T-Y, Aung K, Tseng C-Y, Chang T-Y, Chen Y-S, Chiou T-J. 2011. Vacuolar Ca2+/H+ Transport Activity Is Required for Systemic Phosphate Homeostasis Involving Shoot-to-Root Signaling in Arabidopsis   Plant Physiology **156**, 1176-1189.

Liu TY, Huang TK, Tseng CY, Lai YS, Lin SI, Lin WY, Chen JW, Chiou TJ. 2012. PHO2-dependent degradation of PHO1 modulates phosphate homeostasis in Arabidopsis. Plant Cell **24**, 2168-2183.

Liu Z, Huang S, Zhu L, Li C, Zhang D, Chen M, Liu Y, Zhang Y. 2024. PHR1 negatively regulates nitrate reductase activity by directly inhibiting the transcription of NIA1 in Arabidopsis. Journal of Plant Physiology **303**, 154382.

Luan M, Zhao F, Han X, Sun G, Yang Y, Liu J, Shi J, Fu A, Lan W, Luan S. 2019. Vacuolar Phosphate Transporters Contribute to Systemic Phosphate Homeostasis Vital for Reproductive Development in Arabidopsis. Plant Physiol **179**, 640-655.

MacNeill GJ, Mehrpouyan S, Minow MAA, Patterson JA, Tetlow IJ, Emes MJ. 2017. Starch as a source, starch as a sink: the bifunctional role of starch in carbon allocation. J Exp Bot **68**, 4433-4453.

Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H, Yanagisawa S. 2018. A NIGT1-centred transcriptional cascade regulates nitrate signalling and incorporates phosphorus starvation signals in Arabidopsis. Nat Commun **9**, 1376.

Maillard A, Diquelou S, Billard V, Laine P, Garnica M, Prudent M, Garcia-Mina JM, Yvin JC, Ourry A. 2015. Leaf mineral nutrient remobilization during leaf senescence and modulation by nutrient deficiency. Front Plant Sci **6**, 317.

Marschner H. 2012. *Marschner's Mineral Nutrition of Higher Plants*.

Medici A, Marshall-Colon A, Ronzier E, Szponarski W, Wang R, Gojon A, Crawford NM, Ruffel S, Coruzzi GM, Krouk G. 2015. AtNIGT1/HRS1 integrates nitrate and phosphate signals at the Arabidopsis root tip. Nat Commun **6**, 6274.

Medici A, Szponarski W, Dangeville P, Safi A, Dissanayake IM, Saenchai C, Emanuel A, Rubio V, Lacombe B, Ruffel S, Tanurdzic M, Rouached H, Krouk G. 2019. Identification of Molecular Integrators Shows that Nitrogen Actively Controls the Phosphate Starvation Response in Plants. Plant Cell **31**, 1171-1184.

Mehra P, Pandey BK, Verma L, Giri J. 2019. A novel glycerophosphodiester
phosphodiesterase improves phosphate deficiency tolerance in rice. Plant Cell
Environ **42**, 1167-1179.
Nakamura Y. 2013. Phosphate starvation and membrane lipid remodeling in seed
plants. Prog Lipid Res **52**, 43-50.
Neto AP, Favarin JL, Hammond JP, Tezotto T, Couto HT. 2016. Analysis of
Phosphorus Use Efficiency Traits in Coffea Genotypes Reveals Coffea arabica and
Coffea canephora Have Contrasting Phosphorus Uptake and Utilization Efficiencies.
Front Plant Sci **7**, 408.
Ngo AH, Angkawijaya AE, Lin YC, Liu YC, Nakamura Y. 2022. The phospho-base
N-methyltransferases PMT1 and PMT2 produce phosphocholine for leaf growth in
phosphorus-starved Arabidopsis. J Exp Bot **73**, 2985-2994.
Nilsson L, Muller R, Nielsen TH. 2007. Increased expression of the MYB-related
transcription factor, PHR1, leads to enhanced phosphate uptake in Arabidopsis
thaliana. Plant Cell Environ **30**, 1499-1512.
O'Leary B, Park J, Plaxton WC. 2011. The remarkable diversity of plant PEPC
(phosphoenolpyruvate carboxylase): recent insights into the physiological functions
and post-translational controls of non-photosynthetic PEPCs. Biochem J **436**, 15-34.
Ojeda-Rivera JO, Alejo-Jacuinde G, Najera-Gonzalez HR, Lopez-Arredondo D.
2022. Prospects of genetics and breeding for low-phosphate tolerance: an integrated
approach from soil to cell. Theor Appl Genet.
Okamoto M, Kumar A, Li W, Wang Y, Siddiqi MY, Crawford NM, Glass ADM. 2006.
High-Affinity Nitrate Transport in Roots of Arabidopsis Depends on Expression of the
NAR2-Like Gene AtNRT3.1. Plant Physiology **140**, 1036-1046.
Okamoto Y, Suzuki T, Sugiura D, Kiba T, Sakakibara H, Hachiya T. 2018. Shoot
nitrate underlies a perception of nitrogen satiety to trigger local and systemic signaling
cascades in
Arabidopsis thaliana. Soil Science and Plant Nutrition **65**, 56-64.
Okazaki Y, Otsuki H, Narisawa T, Kobayashi M, Sawai S, Kamide Y, Kusano M,
Aoki T, Hirai MY, Saito K. 2013. A new class of plant lipid is essential for protection
against phosphorus depletion. Nat Commun **4**, 1510.
Ouyang X, , Hong X, , Zhao X, , Zhang W, , He X, , Ma W, , Teng W, , Tong Y, .
2016. Knock out of the PHOSPHATE 2 Gene TaPHO2-A1 Improves Phosphorus
Uptake and Grain Yield under Low Phosphorus Conditions in Common Wheat.
Scientific Reports **6**, 1-11.
Pang J, Tibbett M, Denton MD, Lambers H, Siddique KHM, Bolland MDA, Revell
CK, Ryan MH. 2009. Variation in seedling growth of 11 perennial legumes in response
to phosphorus supply. Plant and Soil **328**, 133-143.
Park SH, Jeong JS, Huang CH, Park BS, Chua NH. 2023. Inositol polyphosphates-
regulated polyubiquitination of PHR1 by NLA E3 ligase during phosphate starvation
response in Arabidopsis. New Phytol **237**, 1215-1228.
Paz-Ares J, Puga MI, Rojas-Triana M, Martinez-Hevia I, Diaz S, Poza-Carrión C,
Miñambres M, Leyva A. 2022. Plant adaptation to low phosphorus availability: Core
signaling, crosstalks, and applied implications. Mol Plant **15**, 104-124.
Penn CJ, Camberato JJ, Wiethorn MA. 2023. How Much Phosphorus Uptake Is
Required for Achieving Maximum Maize Grain Yield? Part 1: Luxury Consumption and
Implications for Yield. Agronomy **13**, 95.
Péret B, Desnos T, Jost R, Kanno S, Berkowitz O, Nussaume L. 2014. Root
Architecture Responses: In Search of Phosphate. Plant Physiol **166**, 1713-1723.

Pérez-Harguindeguy N, Díaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P,
Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE, Urcelay C, Veneklaas EJ,
Reich PB, Poorter L, Wright IJ, Ray P, Enrico L, Pausas JG, de Vos AC,
Buchmann N, Funes G, Quétier F, Hodgson JG, Thompson K, Morgan HD, ter
Steege H, Sack L, Blonder B, Poschlod P, Vaieretti MV, Conti G, Staver AC,
Aquino S, Cornelissen JHC. 2016. Corrigendum to: New handbook for standardised
measurement of plant functional traits worldwide. Australian Journal of Botany **64**.
Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. 2017. Differential analysis of
RNA-seq incorporating quantification uncertainty. Nat Methods **14**, 687-690.
Plaxton WC, Tran HT. 2011. Metabolic adaptations of phosphate-starved plants.
Plant Physiol **156**, 1006-1015.
Poza-Carrion C, Paz-Ares J. 2019. When nitrate and phosphate sensors meet. Nat
Plants **5**, 339-340.
Prodhan MA, Jost R, Watanabe M, Hoefgen R, Lambers H, Finnegan PM. 2016.
Tight control of nitrate acquisition in a plant species that evolved in an extremely
phosphorus-impoverished environment. Plant Cell Environ **39**, 2754-2761.
Prodhan MA, Jost R, Watanabe M, Hoefgen R, Lambers H, Finnegan PM. 2017.
Tight control of sulfur assimilation: an adaptive mechanism for a plant from a severely
phosphorus-impoverished habitat. New Phytol **215**, 1068-1079.
Pueyo JJ, Quinones MA, Coba de la Pena T, Fedorova EE, Lucas MM. 2021.
Nitrogen and Phosphorus Interplay in Lupin Root Nodules and Cluster Roots. Front
Plant Sci **12**, 644218.
Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, de Lorenzo L,
Irigoyen ML, Masiero S, Bustos R, Rodriguez J, Leyva A, Rubio V, Sommer H,
Paz-Ares J. 2014. SPX1 is a phosphate-dependent inhibitor of Phosphate Starvation
Response 1 in Arabidopsis. Proc Natl Acad Sci U S A **111**, 14947-14952.
Punja ZK, Holmes JE. 2020. Hermaphroditism in Marijuana (Cannabis sativa L.)
Inflorescences – Impact on Floral Morphology, Seed Formation, Progeny Sex Ratios,
and Genetic Variation. Front Plant Sci **11**, 718.
Raven JA. 2015. Interactions between Nitrogen and Phosphorus metabolism. *Annual*
Plant Reviews Volume 48, 187-214.
Ried MK, Wild R, Zhu J, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA,
Hothorn LA, Fiedler D, Hiller S, Hothorn M. 2021. Inositol pyrophosphates promote
the interaction of SPX domains with the coiled-coil motif of PHR transcription factors
to regulate plant phosphate homeostasis. Nat Commun **12**, 384.
Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei
MR, Laha NP, Krusenbaum L, Schneider R, Saiardi A, Fiedler D, Jessen HJ,
Schaaf G, Giehl RFH. 2021. ITPK1 is an InsP(6)/ADP phosphotransferase that
controls phosphate signaling in Arabidopsis. Mol Plant **14**, 1864-1880.
Rietra RPJJ, Heinen M, Dimkpa CO, Bindraban PS. 2017. Effects of Nutrient
Antagonism and Synergism on Yield and Fertilizer Use Efficiency. Communications in
Soil Science and Plant Analysis **48**, 1895-1920.
Rose T, Kretzschmar T, Liu L, Lancaster G, Wissuwa M. 2016. Phosphorus
Deficiency Alters Nutrient Accumulation Patterns and Grain Nutritional Quality in Rice.
Agronomy **6**.
Rouached H. 2011. Multilevel coordination of phosphate and sulfate homeostasis in
plants. Plant Signal Behav **6**, 952-955.
Rouached H, Secco D, Arpat B, Poirier Y. 2011. The transcription factor PHR1 plays
a key role in the regulation of sulfate shoot-to-root flux upon phosphate starvation in
Arabidopsis. BMC Plant Biol **11**, 19.

Rubio V, Linhares F, Solano R, Martin AC, Iglesias J, Leyva A, Paz-Ares J. 2001.
A conserved MYB transcription factor involved in phosphate starvation signaling both
in vascular plants and in unicellular algae. Genes Dev **15**, 2122-2133.
Safi A, Medici A, Szponarski W, Ruffel S, Lacombe B, Krouk G. 2017. The world
according to GARP transcription factors. Curr Opin Plant Biol **39**, 159-167.
Satheesh V, Zhang J, Li J, You Q, Zhao P, Wang P, Lei M. 2022. High transcriptome
plasticity drives phosphate starvation responses in tomato. Stress Biol **2**, 18.
Scalabrin E, Radaelli M, Capodaglio G, Pierobon M, Del Vecchio S, Buffa G. 2024.
Hemp cultivation opportunities for marginal lands development. PLoS One **19**,
e0299981.
Schluter U, Mascher M, Colmsee C, Scholz U, Brautigam A, Fahnenstich H,
Sonnewald U. 2012. Maize source leaf adaptation to nitrogen deficiency affects not
only nitrogen and carbon metabolism but also control of phosphate homeostasis. Plant
Physiol **160**, 1384-1406.
Shane MW, McCully ME, Lambers H. 2004a. Tissue and cellular phosphorus storage
during development of phosphorus toxicity in Hakea prostrata (Proteaceae). Journal
of Experimental Botany **55**, 1033-1044.
Shane MW, Szota C, Lambers H. 2004b. A root trait accounting for the extreme
phosphorus sensitivity of Hakea prostrata (Proteaceae). Plant, Cell & Environment **27**,
991-1004.
Sharma NC, Sahi SV. 2005. Characterization of Phosphate Accumulation in Lolium
multiflorum for Remediation of Phosphorus-Enriched Soils. Environmental Science &
Technology **39**, 5475-5480.
Shimojima M, Watanabe T, Madoka Y, Koizumi R, Yamamoto MP, Masuda K,
Yamada K, Masuda S, Ohta H. 2013. Differential regulation of two types of
monogalactosyldiacylglycerol synthase in membrane lipid remodeling under
phosphate-limited conditions in sesame plants. Front Plant Sci **4**, 469.
Shiponi S, Bernstein N. 2021a. Response of medical cannabis (Cannabis sativa L.)
genotypes to P supply under long photoperiod: Functional phenotyping and the
ionome. Industrial Crops and Products **161**.
Shiponi S, Bernstein N. 2021b. The Highs and Lows of P Supply in Medical
Cannabis: Effects on Cannabinoids, the Ionome, and Morpho-Physiology. Front Plant
Sci **12**.
Shukla D, Rinehart CA, Sahi SV. 2017. Comprehensive study of excess phosphate
response reveals ethylene mediated signaling that negatively regulates plant growth
and development. Scientific Reports **7**, 1-16.
Siebers M, Dörmann P, Hölzl G. 2015. Membrane remodelling in phosphorus-
deficient plants. *Annual Plant Reviews Volume 48*, 237-263.
Struik PC, Amaducci S, Bullard MJ, Stutterheim NC, Venturi G, Cromack HTH.
2000. Agronomy of fibre hemp (Cannabis sativa L.) in Europe. Industrial Crops and
Products **11**, 107-118.
Su T, Xu Q, Zhang FC, Chen Y, Li LQ, Wu WH, Chen YF. 2015. WRKY42 modulates
phosphate homeostasis through regulating phosphate translocation and acquisition in
Arabidopsis. Plant Physiol **167**, 1579-1591.
Sun G, Luan M, Wen J, Wang B, Lan W. 2023. Genetically controlling VACUOLAR
PHOSPHATE TRANSPORTER 1 contributes to low-phosphorus seeds in
Arabidopsis. Plant Signal Behav **18**, 2186641.
Suriyagoda LDB, Ryan MH, Gille CE, Dayrell RLC, Finnegan PM, Ranathunge K,
Nicol D, Lambers H. 2023. Phosphorus fractions in leaves. New Phytol **237**, 1122-
1135.

Tang K, Fracasso A, Struik PC, Yin X, Amaducci S. 2018. Water- and Nitrogen-Use Efficiencies of Hemp (Cannabis sativa L.) Based on Whole-Canopy Measurements and Modeling. Front Plant Sci **9**, 951.

Tannert M, May A, Ditfe D, Berger S, Balcke GU, Tissier A, Kock M. 2018. Pi starvation-dependent regulation of ethanolamine metabolism by phosphoethanolamine phosphatase PECP1 in Arabidopsis roots. J Exp Bot **69**, 467-481.

Taylor CB, Bariola PA, delCardayré SB, Raines RT, Green PJ. 1993. RNS2: a senescence-associated RNase of Arabidopsis that diverged from the S-RNases before speciation. Proceedings of the National Academy of Sciences **90**, 5118-5122.

Tjellstrom H, Andersson MX, Larsson KE, Sandelius AS. 2008. Membrane phospholipids as a phosphate reserve: the dynamic nature of phospholipid-to-digalactosyl diacylglycerol exchange in higher plants. Plant Cell Environ **31**, 1388-1398.

Torres-Rodriguez JV, Salazar-Vidal MN, Chavez Montes RA, Massange-Sanchez JA, Gillmor CS, Sawers RJH. 2021. Low nitrogen availability inhibits the phosphorus starvation response in maize (Zea mays ssp. mays L.). BMC Plant Biol **21**, 259.

Tran HT, Hurley BA, Plaxton WC. 2010. Feeding hungry plants: The role of purple acid phosphatases in phosphate nutrition. Plant Science **179**, 14-27.

Tripathi R, Tewari R, Singh KP, Keswani C, Minkina T, Srivastava AK, De Corato U, Sansinenea E. 2022. Plant mineral nutrition and disease resistance: A significant linkage for sustainable crop protection. Front Plant Sci **13**, 883970.

Ueda Y, Kiba T, Yanagisawa S. 2020a. Nitrate-inducible NIGT1 proteins modulate phosphate uptake and starvation signalling via transcriptional regulation of SPX genes. Plant J **102**, 448-466.

Ueda Y, Nosaki S, Sakuraba Y, Miyakawa T, Kiba T, Tanokura M, Yanagisawa S. 2020b. NIGT1 family proteins exhibit dual mode DNA recognition to regulate nutrient response-associated genes in Arabidopsis. PLoS Genet **16**, e1009197.

Ueda Y, Yanagisawa S. 2023. Transcription factor module NLP-NIGT1 fine-tunes NITRATE TRANSPORTER2.1 expression. Plant Physiol **193**, 2865-2879.

Vance CP, Stone CU, Allan DL. 2003. Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource. New Phytol **157**, 423-447.

Veneklaas EJ, Lambers H, Bragg J, Finnegan PM, Lovelock CE, Plaxton WC, Price CA, Scheible WR, Shane MW, White PJ, Raven JA. 2012. Opportunities for improving phosphorus-use efficiency in crop plants. New Phytol **195**, 306-320.

Wang D, Lv S, Jiang P, Li Y. 2017. Roles, Regulation, and Agricultural Application of Plant Phosphate Transporters. Front Plant Sci **8**, 817.

Wang X, Wang HF, Chen Y, Sun MM, Wang Y, Chen YF. 2020. The Transcription Factor NIGT1.2 Modulates Both Phosphate Uptake and Nitrate Influx during Phosphate Starvation in Arabidopsis and Maize. Plant Cell **32**, 3519-3534.

Wang Y, Ribot C, Rezzonico E, Poirier Y. 2004. Structure and expression profile of the Arabidopsis PHO1 gene family indicates a broad role in inorganic phosphate homeostasis. Plant Physiol **135**, 400-411.

Wang YH, Garvin DF, Kochian LV. 2002. Rapid induction of regulatory and transporter genes in response to phosphorus, potassium, and iron deficiencies in tomato roots. Evidence for cross talk and root/rhizosphere-mediated signals. Plant Physiol **130**, 1361-1370.

Wang Z, Ruan W, Shi J, Zhang L, Xiang D, Yang C, Li C, Wu Z, Liu Y, Yu Y, Shou H, Mo X, Mao C, Wu P. 2014. Rice SPX1 and SPX2 inhibit phosphate starvation

responses through interacting with PHR2 in a phosphate-dependent manner. Proc Natl Acad Sci U S A **111**, 14953-14958.

Wee Y. B, Berkowitz O, Whelan J, Jost R. 2024. Same, Yet Different: Towards Understanding Nutrient Use In Hemp- And Drug-Type Cannabis. Journal of Experimental Botany.

Wei T, Simko V. 2024. R package 'corrplot': Visualization of a Correlation Matrix.

Welling MT, Deseo MA, Bacic A, Doblin MS. 2021. Untargeted Metabolomic Analyses Reveal Chemical Complexity of Dioecious <i>Cannabis</i> Flowers. Australian Journal of Chemistry **74**, 463-479.

Wrolstad RE, Durst RW, Lee J. 2005. Tracking color and pigment changes in anthocyanin products. Trends in food science & technology **16**, 423-428.

Xie K, Cakmak I, Wang S, Zhang F, Guo S. 2021. Synergistic and antagonistic interactions between potassium and magnesium in higher plants. The Crop Journal **9**, 249-256.

Xie X, Hu W, Fan X, Chen H, Tang M. 2019. Frontiers | Interactions Between Phosphorus, Zinc, and Iron Homeostasis in Nonmycorrhizal and Mycorrhizal Plants. Frontiers in Plant Science **10**.

Xie Z, Mi Y, Kong L, Gao M, Chen S, Chen W, Meng X, Sun W, Chen S, Xu Z. 2023. Cannabis sativa: origin and history, glandular trichome development, and cannabinoid biosynthesis. Hortic Res **10**, uhad150.

Xu L, Zhao H, Wan R, Liu Y, Xu Z, Tian W, Ruan W, Wang F, Deng M, Wang J, Dolan L, Luan S, Xue S, Yi K. 2019. Identification of vacuolar phosphate efflux transporters in land plants. Nat Plants **5**, 84-94.

Yang SY, Lin WY, Hsiao YM, Chiou TJ. 2024. Milestones in understanding transport, sensing, and signaling of the plant nutrient phosphorus. Plant Cell **36**, 1504-1523.

Ye Q, Wang H, Su T, Wu W-H, Chen Y-F. 2018. The Ubiquitin E3 Ligase PRU1 Regulates WRKY6 Degradation to Modulate Phosphate Homeostasis in Response to Low-Pi Stress in Arabidopsis. The Plant Cell **30**, 1062-1076.

Yu B, Xu C, Benning C. 2002. Arabidopsisdisrupted in SQD2encoding sulfolipid synthase is impaired in phosphate-limited growth. Proceedings of the National Academy of Sciences **99**, 5732-5737.

Zager JJ, Lange I, Srividya N, Smith A, Lange BM. 2019. Gene Networks Underlying Cannabinoid and Terpenoid Accumulation in Cannabis. Plant Physiol **180**, 1877-1897.

Zhang H, Forde BG. 1998. An <i>Arabidopsis</i> MADS Box Gene That Controls Nutrient-Induced Changes in Root Architecture. Science **279**, 407-409.

Zhang M, Anderson SL, Brym ZT, Pearson BJ. 2021. Photoperiodic Flowering Response of Essential Oil, Grain, and Fiber Hemp (Cannabis sativa L.) Cultivars. Front Plant Sci **12**, 694153.

Zhang Z, Zhong Z, Xiong Y. 2023. Sailing in complex nutrient signaling networks: Where I am, where to go, and how to go? Mol Plant.

FIGURE LEGENDS:

Figure 1. Phenotypic response of female hemp plants to changes in Pi supply.
Shown are images of representative clonal plants for each treatment prior to final
destructive harvest at 42 DAT. (**A**) Plant morphology across treatments. Chlorosis
gradient increased in senescing leaves along the main stem with decreasing Pi supply.
Scale bar represents 20 cm. (**B**) Root ball morphology across treatments. Scale bar
represents 5 cm. Organ biomass in g FW plant^{-1} is given below each panel, as mean
± SE (n = 3 plants per treatment).

Figure 2. Morphological response to Pi supply across timepoints. (**A**) Changes
in plant height, plant width and node count were determined at indicated time points
across treatments. DAT refers to days after transfer of rooted female cuttings into pots.
Arrows indicate timepoint when Pi treatments were initiated. (**B**) Biomass partitioning
across organ types, as a percentage of total fresh biomass for each treatment.
Vegetative organs are stem and leaves combined. (**C**) Agronomic phosphate
efficiency (APE) across treatments in comparison to 0 mM Pi supply. APE is highest
at 0.25 mM Pi supply and decreases with higher Pi supply across organs. Given are
means ± SE, n = 3 plants per treatment. Letters denote significant difference (ANOVA,
Tukey's HSD, p < 0.05) between treatments and / or organ types. Also refer to
Supplementary Table S1 and S2 for supporting datasets.

**Figure 3. Pi supply affects Pi assimilation as well as inorganic storage pools
across organ types.** (**A**) Concentrations of inorganic and organic P across organ
types and Pi treatments. (**B**) Concentrations of nitrate, phosphate and sulfate across
organ types and Pi treatments. Letters denote statistical significance across
treatments (ANOVA/TUKEY HSD, p < 0.05). Given are means ± SE (n = 3 plants per
treatment). See Supplementary Table S3 for data and statistical analyses. TF, terminal
flower; YL, young leaf; ML, mature leaf; OL, old leaf; LR, lateral root; YS, young stem.

**Figure 4. Correlation matrix of measured metabolites and P fractions across
organ types.** Pearson's correlation to assess relationship between concentrations of
indicated metabolites and different P fractions for given organs. Asterisks denote

significant correlations ($p < 0.05$) and colour scale indicates correlation coefficient. ND: not determined.

Figure 5. Transcriptional response of curated PSR genes. Cannabis homologs of known PSR genes in Arabidopsis were categorised into functional groups. PSR transcript abundances across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower) and Pi treatments (0, 0.25, 1, 2 mM Pi) are expressed as $\log_2$ transformed TPM values. Transcriptomic data and curated gene list are provided in Supplementary Table S5 to S7. The cs10 reference referenc gene IDs are shown on the right. Abbreviations for gene family names shown to the left of the panel: *PHT, PHOSPHATE TRANSPORTER; PHF1, PHOSHATE TRANSPORT FACILITATOR1; PHO1, PHOSPHATE1; VPE, VACUOLAR PHOSPHATE EFFLUX TRANSPORTER; VPT, VACUOLAR PHOSPHATE TRANSPORTER; PHO2, PHOSPHATE2; NLA, NITROGEN LIMITATION ADAPTATION; PHR1, PHOSPHATE STARVATION RESPONSE1; SPX, SPX DOMAIN GENE; ITPK, INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE; WRKY, WRKY DNA-BINDING PROTEIN; MYB, MYB-DOMAIN PROTEIN; RNS, RIBONUCLEASE; PAP, PURPLE ACID PHOSPHATASE; PLD, PHOSPHOLIPASE D; GDPD, GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE; PECP1, PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE PHOSPHATASE1; PEAMT, PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE; MGD, MONO-GALACTOSYL DIACYLGLYCEROL SYNTHASE; SQD, SULFOQUINOVOSYL-DIACYLGLYCEROL SYNTHASE.*

Figure 6. Transcriptionalresponse of curated genes involved in N assimilation and metabolism in hemp. Shown are transcript abundances for Cannabis homologs of genes with a known role in N assimilation, metabolism and their transcriptional regulation in Arabidopsis. The heatmap represents log-transformed TPM ($\log_2(\text{TPM}+1)$) across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower). Transcriptomic dataset and fully curated gene list is provided in Supplementary Table S5-S7. cs10 genome gene IDs are shown on the right. Abbreviations for gene family names shown on the left of the panel: *AMT, AMMONIUM TRANSPORTER; NRT, NITRATE TRANSPORTER; ANR1, ARABIDOPSIS NITRATE REGULATED1; NITRITE REDUCTASE, NIR; NITRATE*

REDUCTASE, NIA; GLUTAMINE SYNTHETASE, GLN; NITRATE-INDUCIBLE GARP-TYPE TRANSCRIPTIONAL REPRESSOR1, NIGT1; NIN-LIKE PROTEIN, NLP; NRT1/PEPTIDE TRANSPORTER, NPF; GLUTAMATE DEHYDROGENASE, GDH; ASPARTATE AMINOTRANSFERASE, AAT; C-TERMINALLY ENCODED PEPTIDE, CEP; CALCINEURIN B-LIKE PROTEIN-INTERACTING PROTEIN KINASE, CIPK.

Figure 1. Phenotypic response of female hemp plants to changes in Pi supply. Shown are images of representative clonal plants for each treatment prior to final destructive harvest at 42 DAT. (**A**) Plant morphology across treatments. Chlorosis gradient increased in senescing leaves along the main stem with decreasing Pi supply. Scale bar represents 20 cm. (**B**) Root ball morphology across treatments. Scale bar represents 5 cm. Organ biomass in g FW plant^{-1} is given below each panel, as mean ± SE (n = 3 plants per treatment).

Figure 2. Morphological response to Pi supply across timepoints. (A) Changes in plant height, plant width and node count were determined at indicated time points across treatments. DAT refers to days after transfer of rooted female cuttings into pots. Arrows indicate timepoint when Pi treatments were initiated. **(B)** Biomass partitioning across organ types, as a percentage of total fresh biomass for each treatment. Vegetative organs are stem and leaves combined. **(C)** Agronomic phosphate efficiency (APE) across treatments in comparison to 0 mM Pi supply. APE across is highest at 0.25 mM Pi supply and decreases with higher Pi supply across organs. Given are means ± SE, n = 3 plants per treatment. Letters denote significant difference (ANOVA, Tukey's HSD, p < 0.05) between treatments and / or organ types. Also refer to Supplemental Tables 1 and 2 for supporting datasets.

Figure 3. Pi supply affects Pi assimilation as well as inorganic storage pools across organ types.
(A) Concentrations of inorganic and organic P across organ types and Pi treatments. **(B)** Concentrations of nitrate, phosphate and sulfate across organ types and Pi treatments. Letters denote statistical significance across treatments (ANOVA/TUKEY HSD, p < 0.05). Given are means ± SE (n = 3 plants per treatment). See Supplementary Table S3 for data and statistical analyses. TF, terminal flower; YL, young leaf; ML, mature leaf; OL, old leaf; LR, lateral root; YS, young stem.

Figure 4. Correlation matrix of measured metabolites and P fractions across organ types. Pearson's correlation to assess relationship between concentrations of indicated metabolites and different P fractions for given organs. Asterisks denote significant correlations (p < 0.05) and colour scale indicates correlation coefficient. ND: not determined.

Figure 5. Transcriptional response of curated PSR genes. *Cannabis* homologs of known PSR genes in Arabidopsis were categorised into functional groups. PSR transcript abundance across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower) is expressed as log2 transformed TPM values. Transcriptomic data and curated gene list are provided in Supplemental Tables 5 to 7. The cs10 reference gene IDs are shown on the right. Abbreviations for gene family names shown to the left of the panel: *PHT, PHOSPHATE TRANSPORTER; PHF1, PHOSHATE TRANSPORT FACILITATOR1; PHO1, PHOSPHATE1; VPE, VACUOLAR PHOSPHATE EFFLUX TRANSPORTER; VPT, VACUOLAR PHOSPHATE TRANSPORTER; PHO2, PHOSPHATE2; NLA, NITROGEN LIMITATION ADAPTATION; PHR1, PHOSPHATE STARVATION RESPONSE1; SPX, SPX DOMAIN GENE; ITPK, INOSITOL 1,3,4-TRISPHOSPHATE 5/6-KINASE; WRKY, WRKY DNA-BINDING PROTEIN; MYB, MYB-DOMAIN PROTEIN; RNS, RIBONUCLEASE; PAP, PURPLE ACID PHOSPHATASE; PLD, PHOSPHOLIPASE D; GDPD, GLYCEROPHOSPHODIESTER PHOSPHODIESTERASE; PECP1, PHOSPHOETHANOLAMINE/PHOSPHOCHOLINE PHOSPHATASE1; PEAMT, PHOSPHOETHANOLAMINE N-METHYLTRANSFERASE; MGD, MONO-GALACTOSYL DIACYLGLYCEROL SYNTHASE; SQD, SULFOQUINOVOSYL-DIACYLGLYCEROL SYNTHASE.*

Figure 6. Transcriptional response of curated genes involved in N assimilation and metabolism in hemp. Shown are transcript abundances for *Cannabis* homologs of genes with a known role in N assimilation, metabolism and their transcriptional regulation in Arabidopsis. The heatmap represents log-transformed TPM ($\log_2(TPM+1)$) across organ types (LR, lateral root; OL, old leaf, ML, mature leaf; YS, young stem; TF, terminal flower). Transcriptomic dataset and fully curated gene list is provided in Supplemental Tables 5-7. cs10 genome gene IDs are shown on the right. Abbreviations for gene family names shown on the left of the panel: *AMT, AMMONIUM TRANSPORTER; NRT, NITRATE TRANSPORTER; ANR1, ARABIDOPSIS NITRATE REGULATED1; NITRITE REDUCTASE, NIR; NITRATE REDUCTASE, NIA; GLUTAMINE SYNTHETASE, GLN; NITRATE-INDUCIBLE GARP-TYPE TRANSCRIPTIONAL REPRESSOR1, NIGT1; NIN-LIKE PROTEIN, NLP; NRT1/PEPTIDE TRANSPORTER, NPF; GLUTAMATE DEHYDROGENASE, GDH; ASPARTATE AMINOTRANSFERASE, AAT; C-TERMINALLY ENCODED PEPTIDE, CEP; CALCINEURIN B-LIKE PROTEIN-INTERACTING PROTEIN KINASE, CIPK.*