Tumors mimic the niche to inhibit neighboring stem cell differentiation

Yang Zhang[1], Yuejia Wang[1], Jinqiao Song[1], Lizhong Yan[1], Ziguang Wang[1], Dongze

Song[1], Yudi Zhao[1], and Shaowei Zhao[1,2]*

[1]Department of Genetics and Cell Biology, College of Life Sciences, Nankai University,

Tianjin, 300071, People's Republic of China

[2]Lead contact

* For correspondence: swzhao@nankai.edu.cn

Abstract

Although it is well-established that stem cells maintain tissue homeostasis while tumors

disrupt it, the mechanisms by which tumors influence the development of nearby stem

cells remain poorly understood. Using *Drosophila* ovaries as a model system, here we

discovered that *bam* or *bgcn* mutant germline tumors inhibit the differentiation of

neighboring wild-type germline stem cells (GSCs). Mechanistically, these tumor cells

mimic the stem cell niche by secreting the BMP ligands Dpp and Gbb, but at reduced

levels, resulting in moderate BMP signaling activation in adjacent GSCs. Such BMP

signaling activation is sufficient to repress *bam* transcription, thereby blocking GSC

differentiation. To our knowledge, this is the first example that tumors can functionally

mimic a stem cell niche to inhibit the differentiation of neighboring wild-type stem cells.

Similar regulatory paradigms may operate in mammalian tissues, including humans,

during tumorigenesis.

Key words: *bam*, BMP signaling, tumors, germline stem cell, stem cell niche,

Drosophila

Introduction

The homeostasis of many tissues in our bodies is maintained by adult stem cells, but this balance can be disrupted by tumor cells. What occurs when tumorigenesis intersects with stem cell development? To address this question, a mosaic-analysis model system is essential, where wild-type stem cells develop alongside tumor cells. *Drosophila* offers an exceptional model for such studies, as it allows for the efficient generation of mosaic clones through various established methods *(Germani et al., 2018; Pastor-Pareja and Xu, 2013)*.

In *Drosophila* ovaries, germline stem cells (GSCs) play a crucial role in sustaining normal oogenesis and maintaining fertility *(Fuller and Spradling, 2007; Lin, 1997)*. These GSCs reside in a specialized microenvironment known as the stem cell niche (hereafter referred to as niche) *(Xie and Spradling, 2000)*. Typically, a GSC undergoes asymmetric division, generating two distinct daughter cells: one remains in the niche to self-renew as a GSC, while the other, called a cystoblast, exits the niche and initiates differentiation. During the differentiation process, each cystoblast performs exactly four rounds of mitotic division with incomplete cytokinesis to produce16 interconnected cystocytes, forming a germline cyst. In each germline cyst, only one germ cell is destined to become the oocyte, while the remaining 15 differentiate into nurse cells that support the development of the oocyte (*Figure 1A*) *(Fuller and Spradling, 2007; Lin, 1997)*. The principal niche signals are bone morphogenetic protein (BMP) ligands, including Decapentaplegic (Dpp) and Glass bottom boat (Gbb), which are secreted predominantly by cap cells, the major somatic niche cells (Chen and McKearin, 2003a; Li et al., 2016; Song et al., 2004; Xie and Spradling, 1998, 2000). These ligands activate BMP signaling in GSCs, leading to the transcriptional repression of *bag of marbles* (*bam*), a key gene that promotes differentiation. In contrast, BMP signaling is inactive in cystoblasts, allowing Bam to be expressed and drive their

49   differentiation (**Chen and McKearin, 2003a; Song et al., 2004**). Bam carries out this

50   function in collaboration with its partner, Benign gonial cell neoplasm (Bgcn) (**Li et al.,**

51   **2009; Ohlstein et al., 2000**).

52   GSCs mutant for *bam* or *bgcn* fail to differentiate and instead hyper-proliferate,

forming a well-established *Drosophila* germline tumor model (**Lavoie et al., 1999;**

McKearin and Ohlstein, 1995; McKearin and Spradling, 1990; Niki and Mahowald,

2003). Notably, these germline tumor cells competitively displace wild-type GSCs from

the niche (**Jin et al., 2008**). The resulting displacement creates a microenvironment

where wild-type GSCs are surrounded by tumor cells, providing an excellent model

system to study stem cell behavior in tumor neighborhoods.

59   Here, we demonstrate that *bam* or *bgcn* mutant germline tumors inhibit the

60   differentiation of neighboring wild-type GSCs by functionally mimicking the stem cell

61   niche. This mechanism may be conserved in mammals, including humans, during

62   tumorigenesis, where malignant cells could similarly disrupt normal stem cell

63   development.

Results

Germline tumors inhibit the differentiation of neighboring wild-type GSCs

Using the *nos>FLP/FRT* or *hs-FLP/FRT* methods we previously established (***Zhang et al., 2023; Zhao et al., 2018***), we generated *bam* or *bgcn* mutant germline clones in *Drosophila* ovaries (***Figure 1B***). Surprisingly, we found that many wild-type germ cells located outside the niche retained a GSC-like single-germ-cell (SGC) morphology, even when encapsulated within egg chambers (***Figure 1C, D, Figure 1—figure supplement 1***). Under normal conditions, GSCs that exit the niche differentiate into interconnected germline cysts, where germ cells are linked rather than remaining as individual, isolated cells (***Fuller and Spradling, 2007; Xie and Spradling, 2000***). To rule out the possibility that the SGC phenotype is an artifact caused by GFP expression, we repeated the experiments using RFP and arm-lacZ as alternative mosaic-analysis markers. Consistent results were observed (***Figure 1E, F***), confirming that the phenotype is not attributable to GFP.

To further confirm that these SGCs exhibit GSC-like characteristics, we conducted anti-α-Spectrin immunofluorescent staining, a method that labels a germline-specific organelle known as the spectrosome in GSCs and cystoblasts, and the fusome in cystocytes. GSCs perform complete cell division, whereas cystocytes undergo incomplete cytokinesis, remaining interconnected through fusomes and ring canals. Consequently, spectrosomes appear as dot-like structures, while fusomes exhibit branched morphologies (***Figure 1G***) (***Lin et al., 1994***). To accurately capture the three-dimensional (3D) architecture of spectrosomes and fusomes, we acquired z-stack images using confocal microscopy. Strikingly, these SGCs displayed dot-like spectrosomes, closely resembling those observed in wild-type GSCs and *bam* or *bgcn* mutant GSC-like tumor cells (***Figure 1H, I***). We also considered the possibility that SGCs might arise through the dedifferentiation of the cystocytes in germline cysts

surrounded by germline tumors. If this were the case, such cystocytes would initially

undergo complete cell division, leaving behind midbodies as markers of the late

cytokinesis stage. When visualized by anti-α-Spectrin immunofluorescence, midbody

appears as a central sphere that is slightly connected to two larger flanking structures,

resembling a variant of nunchucks (**Mathieu et al., 2022**). Notably, in our analyses of

over 50 germline cysts surrounded by *bam* mutant germline tumors, none contained

midbodies, suggesting that dedifferentiation is unlikely to be the primary mechanism

responsible for the SGC phenotype. Together, these findings indicate that *bam* or *bgcn*

mutant germline tumors inhibit the differentiation of neighboring wild-type GSCs.

The inhibition of differentiation in SGCs relies on the lack of Bam expression

Given that Bam is the key factor promoting GSC differentiation (**McKearin and**

Ohlstein, 1995; Ohlstein and McKearin, 1997), we were very curious about the

expression of Bam in SGCs. At first, we assessed Bam protein levels using

immunofluorescent staining with an anti-BamC antibody (**McKearin and Ohlstein,**

1995). Strikingly, none of the SGCs examined (n > 100) were BamC-positive (**Figure**

2A). Then, we analyzed *bam* transcription levels using a *bamP*-GFP reporter (**Chen and**

McKearin, 2003b). Under normal conditions, 100% of GSCs within the niche (n = 153)

were GFP-negative, while 98% of cystoblasts (n = 106) were GFP-positive (**Figure 2B,**

C), confirming that *bam* transcription is associated with the initiation of GSC

differentiation (**McKearin and Ohlstein, 1995**). Notably, 74% of SGCs (n = 132) were

GFP-negative, while the remaining 26% were GFP-positive (**Figure 2B, C**). This

suggests that SGCs can be categorized into two distinct groups: those resembling GSCs

(GSC-like) and those resembling cystoblasts (cystoblast-like). The cystoblast-like SGCs

may have already initiated their differentiation program toward becoming cystocytes.

Since *bam* transcription initiates in cystoblasts (**McKearin and Spradling, 1990**) but

Bam proteins accumulate predominantly in cystocytes (**McKearin and Ohlstein, 1995**),

the Bam protein levels in these cystoblast-like SGCs are likely below the detection

threshold at this early stage.

Next, we asked whether ectopic expression of Bam can drive SGCs to

differentiate. To address this, we established two experimental scenarios: one with the

hs-bam element and one without as the control. In the *hs-bam* scenario (*with hs-bam*),

GFP-positive germ cells are wild-type (carrying *hs-bam*), while GFP-negative cells are

bam$^{-/-}$ (lacking *hs-bam*). In the control scenario (*without hs-bam*), GFP-positive cells are

wild-type, and GFP-negative cells are *bam$^{-/-}$* (**Figure 2D**, see genotypes in **Source data**

1). To quantify the SGC phenotype, we calculated the percentage of SGCs relative to the

total number of SGCs and germline cysts, considering the out-of-niche germ cells that

are either fully enclosed by germline tumors (e.g., the right SGC in **Figure 1I** and the

marked germline cyst in **Figure 2G**) or in contact with wild-type germ cells or somatic

cells on just one side (e.g., the left SGC in **Figure 1I** and the outlined germline cyst in

Figure 4C). Although the SGC phenotype was detectable after fly eclosion, we

quantified it in 14-day-old flies to ensure robust germline tumor growth, improving

quantification efficiency. To induce ectopic Bam expression, 12-day-old female flies were

subjected to heatshock treatment, which involved heating at 37°C for two hours, twice

daily with a 6-hour interval, and over two consecutive days. In the absence of heatshock

treatment, the percentage of SGCs in ovaries of both genotypes showed no significant

difference at either 12 or 14 days (**Figure 2E-H**), indicating that the *hs-bam* element

alone, without heatshock, does not affect the phenotype. However, following heatshock

treatment, the percentage of SGCs in ovaries with *hs-bam* was markedly reduced

compared to those without *hs-bam* (**Figure 2E-H**), suggesting that ectopic Bam

expression can drive SGCs to differentiate. Collectively, these results support that the

differentiation defects of SGCs are due to the lack of Bam expression.

SGCs retain moderate BMP signaling activation

Within the niche, BMP signaling functions to repress *bam* transcription to inhibit GSC

differentiation (**Chen and McKearin, 2003a; Song et al., 2004**). To investigate BMP

signaling activation in SGCs, we employed immunofluorescent staining for pMad, a well-

characterized marker of BMP signaling activity (**Kai and Spradling, 2003**). Surprisingly,

we observed undetectable pMad levels in all SGCs examined (n > 100) (**Figure 3A, B**).

To investigate this further, we examined the activity of *Dad*-lacZ, a highly sensitive BMP

signaling reporter known to be activated not only in GSCs but also in cystoblasts (**Kai**

and Spradling, 2003; Song et al., 2004). Notably, 73% of SGCs were lacZ-positive (n =

107), a proportion lower than that of GSCs within the niche, which showed 100% lacZ

positivity (n = 122) (**Figure 3C, D**). Furthermore, when comparing *Dad*-lacZ expression

levels exclusively in lacZ-positive cells, we found that SGCs exhibited significantly lower

expression levels than GSCs within the niche (**Figure 3C, E**). These findings indicate

that BMP signaling is activated in SGCs but at lower levels than those in GSCs within

the niche.

Beyond maintaining *Drosophila* female GSCs in the niche, BMP signaling also

promotes their division (**Xie and Spradling, 1998**). Since the activation levels of BMP

signaling in SGCs were lower than those in GSCs within the niche, we hypothesized that

SGCs would exhibit slower proliferation rates than GSCs. To test this hypothesis, we

performed BrdU incorporation assays. The results revealed that only 4.5% of SGCs

were BrdU-positive (n = 1034), a significantly lower proportion than the 7.8% observed in

GSCs within the niche (n = 1337) (**Figure 3F-H**). These findings further corroborate the

reduced activation of BMP signaling in SGCs relative to GSCs.

BMP signaling inhibits SGC differentiation

Then we investigated whether BMP signaling functions to inhibit SGC differentiation. The

BMP type II receptor Punt and the co-Smad Medea (Med) are essential for maintaining

GSC stemness within the niche (**Xie and Spradling, 1998**). To investigate whether they

are also required to inhibit SGC differentiation, we established a genetic scenario, in

which GFP$^{+/+}$ RFP$^{-/-}$ germ cells are *punt$^{-/-}$* or *med$^{-/-}$*; GFP$^{+/-}$ RFP$^{+/-}$ germ cells are *punt$^{+/-}$*

bam$^{+/-}$ or *med$^{+/-}$ bam$^{+/-}$* (similar to wild-type); and GFP$^{-/-}$ RFP$^{+/+}$ germ cells are *bam$^{-/-}$*. In

control experiments (with no *punt* or *med* mutation), GFP$^{+/+}$ RFP$^{-/-}$ germ cells are wild-

type; GFP$^{+/-}$ RFP$^{+/-}$ germ cells are *bam$^{+/-}$* (similar to wild-type); and GFP$^{-/-}$ RFP$^{+/+}$ germ

cells are *bam$^{-/-}$* (**Figure 4A**, see genotypes in **Source data 1**). Strikingly, the proportion

of *punt$^{-/-}$* or *med$^{-/-}$* SGCs relative to total SGCs was significantly lower than in controls

(**Figure 4B-E**). Conversely, among *punt$^{-/-}$* or *med$^{-/-}$* germ cells meeting our established

criteria for SGC phenotype quantification, germline cysts constituted a higher percentage

compared to controls (**Figure 4F**). These results indicate that Punt and Med function to

inhibit SGC differentiation.

Mothers against dpp (Mad) is the primary transcription factor of BMP signaling,

and it is also essential for GSC maintenance in *Drosophila* ovaries (**Xie and Spradling,**

1998). Unlike *punt* and *med*, which reside on the same chromosome arm (3R) as *bam*,

mad is located on a separate chromosome arm (2L). To investigate whether Mad is

required to inhibit SGC differentiation, we established a genetic scenario, in which GFP$^{-/-}$

germ cells are *mad$^{-/-}$* and GFP$^{+/+}$ germ cells are *bam$^{-/-}$*. In control experiments (with no

mad mutation), GFP$^{-/-}$ germ cells are wild-type and GFP$^{+/+}$ germ cells are *bam$^{-/-}$* (**Figure**

4G, see genotypes in **Source data 1**). Notably, *mad* mutation significantly decreased the

SGC proportion relative to controls (**Figure 4H-J**). These results suggest that, like Punt

and Med, Mad also plays a crucial role in suppressing SGC differentiation. Together,

these findings demonstrate that BMP signaling contributes to inhibiting SGC

differentiation, despite at reduced activation levels.

Germline tumors secret Dpp and Gbb

The next question we sought to address was which cells surrounding SGCs secrete

BMP ligands, such as Dpp and Gbb. Guided by the Occam's Razor principle, we

focused on *bam* or *bgcn* mutant germline tumor cells, as the SGC phenotype is

dependent on their presence. Initial attempts to detect expression of these two ligand

genes using *in situ* hybridization proved unsuccessful. We therefore employed an

established *dpp*-lacZ (*P4*-lacZ) reporter (***Li et al., 2016***) to monitor Dpp expression.

Consistent with previous findings, we observed robust lacZ signals in cap cells (***Figure***

5A), the primary niche cells that are responsible for BMP ligand secretion (***Chen and***

McKearin, 2003a; Song et al., 2004; Xie and Spradling, 2000). Notably, *bam* mutant

germline tumor cells displayed lower lacZ expression levels than cap cells but

significantly higher levels than differentiating cystocytes (***Figure 5A-C***).

To more sensitively assess Dpp and Gbb expression, we performed real-time

quantitative PCR (RT-qPCR) analyses in *bam* or *bgcn* mutant ovaries, comparing

samples with and without germline-specific knockdown of *dpp* or *gbb*. Detection of

reduced transcript levels in knockdown conditions would confirm active expression of

these genes in the respective genetic backgrounds. Consistent with the essential roles

of these two genes in fly viability, ubiquitous knockdown using *act-GAL4* with either *dpp-*

RNAi or *gbb-RNAi* caused lethality, which results also validated the efficacy of these

RNAi lines. Notably, germline-specific knockdown of *dpp* or *gbb* significantly reduced

their transcript levels compared to *yellow* (*y*) or *white* (*w*) knockdown controls (***Figure***

5D, E). These findings demonstrate that *bam* or *bgcn* mutant germline tumors secrete

the BMP ligands, albeit at lower levels than cap cells.

Dpp and Gbb secreted by germline tumors are required to inhibit SGC

differentiation

Finally, we investigated whether Dpp and Gbb secreted by germline tumors are required

to inhibit SGC differentiation. Using a previously established double-mutant mosaic-

analysis strategy for two genes on different chromosomes (***Zhang et al., 2024; Zhang***

et al., 2023), we generated *dpp bam* or *gbb bam* double-mutant germline clones using

two *dpp* mutant alleles, *dpp^{d6}*, *dpp^{d12}*, and one *gbb* allele, *gbb^1* (**Figure 6A, B**, see

genotypes in **Source data 1**). Heterozygotes in any of these alleles did not affect GSC

maintenance, germ cell differentiation, and female fly fertility (**Figure 6—figure

supplement 1**). However, both *dpp bam* and *gbb bam* double-mutant germline tumor

cells exhibited reduced proliferation rates compared to *bam* single-mutant controls

(**Figure 6—figure supplement 2**), indicating that autocrine BMP signaling promotes

bam mutant tumor growth. As mentioned earlier, our evaluation focused on germ cells

that have exited the niche and are surrounded by germline tumors to quantify the SGC

phenotype. However, this raises the question of whether the extent of tumor

encirclement (i.e., being surrounded by more or fewer tumor cells) influences the

phenotype. To investigate this, we compared the SGC phenotype in bigger and smaller

bam mutant germline tumors. Using confocal microscopy, we analyzed 70 germaria

containing *bam* mutant germline clones that met our criteria for SGC phenotype

quantification. For each *bam* mutant germline clone, we scanned and measured its

maximal 2D cross-sectional area as a proxy for its 3D size. The 35 bigger and 35

smaller clones were categorized as 'bigger' and 'smaller' tumors, respectively. Strikingly,

we observed no significant difference in the SGC phenotype between these two groups

of tumors (**Figure 6—figure supplement 3**), suggesting that direct contact between

tumorous and wild-type germ cells, rather than tumor size, is the primary determinant of

this phenotype.

The results above demonstrate that comparing the severity of the SGC

phenotype is feasible between germ cells surrounded by smaller *dpp bam* or *gbb bam*

double-mutant germline tumors and those surrounded by larger *bam* single-mutant

germline tumors. Remarkably, both *dpp bam* and *gbb bam* double-mutant germline

tumors enclosed fewer SGCs but more germline cysts than their *bam* single-mutant

counterparts (**Figure 6C-I**). Thus, we concluded that the BMP ligands from directly-

contacting germline tumor cells mediate the dominant inhibition of SGC differentiation.

This amazingly parallels the mechanism observed in normal stem cell niche, where only

germ cells in direct contact with cap cells are maintained as GSCs (***Chen and***

McKearin, 2003a; Song et al., 2004; Xie and Spradling, 2000).

Discussion

Our study reveals that *bam* or *bgcn* mutant germline tumors in *Drosophila* ovaries

secrete lower levels of BMP ligands Dpp and Gbb than cap cells, resulting in moderate

BMP signaling activation in adjacent wild-type GSCs (called SGCs in this study). Such

BMP signaling activation is sufficient to repress *bam* transcription, thereby blocking SGC

differentiation (see our working model in **Figure 7**). Strikingly, this mechanism closely

recapitulates the normal niche signaling program mediated by cap cells (**Chen and**

McKearin, 2003a; Song et al., 2004; Xie and Spradling, 1998, 2000). To our

knowledge, this represents the first evidence that tumor cells can functionally mimic a

stem cell niche to arrest neighboring wild-type stem cells in an undifferentiated state.

bam or *bgcn* mutant germline tumors consist of GSC-like cells (**Lavoie et al.,**

1999; McKearin and Ohlstein, 1995), which are expected to resemble SGCs. However,

out-of-niche *bgcn* mutant germline tumor cells did not activate BMP signaling to the

same extent as SGCs, as evidenced by much lower *Dad*-lacZ expression in GFP-

negative *bgcn* mutant tumor cells than in neighboring GFP-positive SGCs (**Figure 3C**).

Consistent with this, while only 26% of SGCs expressed *bamP*-GFP, most of out-of-

niche *bam* mutant germline tumor cells were *bamP*-GFP-positive (**Figure 2B, C**), further

supporting reduced BMP signaling in these tumor cells relative to SGCs. These findings

suggest that SGCs are more responsive to BMP signals secreted by germline tumors

than the tumors themselves. Future studies are needed to elucidate the underlying

mechanisms.

One interesting finding is that *bam* or *bgcn* mutant germline tumors secrete lower

levels of BMP ligands than cap cells (**Figure 5A-C**). This aligns with earlier microarray

data showing that purified *Drosophila* female GSCs express minimal Dpp and Gbb (**Kai**

et al., 2005). However, our work reveals that such BMP levels in germline tumors are

functionally critical to suppress SGC differentiation (**Figure 6**). Unlike normal GSCs,

which receive unidirectional BMP ligands from cap cells (*Chen and McKearin, 2003a;*

Li et al., 2016; Song et al., 2004; Xie and Spradling, 2000), SGCs are often fully

surrounded by *bam* or *bgcn* mutant germline tumors. This spatial advantage likely

enables tumors to inhibit SGC differentiation efficiently without matching the high BMP

output of cap cells. Moreover, since BMP signaling is known to both inhibit normal GSC

differentiation and promote their proliferation (*Xie and Spradling, 1998*), it should

similarly stimulate SGC expansion, which is detrimental for *bam* or *bgcn* mutant germline

tumors. We propose that these tumor cells finely regulate BMP secretion to balance

these opposing demands: maintaining differentiation blockade of SGCs while avoiding

stimulation of their excessive proliferation.

A well-established principle in oncology is that tumor aggressiveness correlates

with poor differentiation, with less-differentiated tumors exhibiting enhanced

transformative capacity and metastatic potential (*Jogi et al., 2012; Lytle et al., 2018*). In

Drosophila ovaries, *bam* or *bgcn* mutant germline tumors consist of GSC-like cells that

may resemble these poorly differentiated human tumors (*Lavoie et al., 1999; McKearin*

and Ohlstein, 1995). This similarity raises the possibility that stem cell-like human

tumors may similarly inhibit the differentiation of adjacent wild-type stem cells. By

blocking differentiation, such tumors could deplete terminally differentiated cell

populations, potentially exacerbating patient mortality. This mechanism may contribute to

the heightened lethality of poorly differentiated tumors. Further investigation is needed to

test this hypothesis.

The differentiation of a single GSC into a 16-cell germline cyst, comprising 15

polyploid nurse cells and one developing oocyte, represents a substantial metabolic

investment (*Fuller and Spradling, 2007; Lin, 1997*). We propose that *bam* or *bgcn*

mutant germline tumors block this process to divert nutrients toward their own

uncontrolled growth. This phenomenon could have broad implications, as many human

tissues and organs (intestine, muscle, skin, blood system, male germline, etc.) similarly

depend on adult stem cells for homeostasis (**Blanpain and Fuchs, 2006; Gehart and**

Clevers, 2019; Sousa-Victor et al., 2022; Spradling et al., 2011; Wilkinson et al.,

2020). Notably, these stem cell-dependent tissues and organs are frequent sites of

tumorigenesis, raising the possibility that human cancers may similarly impair

neighboring stem cell differentiation to optimize nutrient allocation for malignant growth.

A key limitation of our study is that the evidence is derived solely from *Drosophila*

germline. Future work should explore whether similar regulatory paradigms operate in

mammalian tissues during tumorigenesis.

Materials and Methods

Key resources table

Reagent or Resource	Source	Identifier
Antibodies		
mouse anti-α-Spectrin	Developmental Studies Hybridoma Bank, DSHB	3A9
mouse anti-BamC	DSHB	AB_10570327
mouse anti-β-Gal	DSHB	JIE7
mouse anti-BrdU	Sigma	B5002
rabbit anti-pMad	Ed Laufer	N/A
rabbit anti-Vasa	Zhaohui Wang	(*Chen et al., 2014*)
goat anti-mouse 546	Invitrogen	A11003
goat anti-rabbit 488	Apexbio	K1206
goat anti-rabbit 546	Invitrogen	A11035
cDNA clone		
bam cDNA clone	Berkeley *Drosophila* Genome Project, BDGP	FI05606
***Drosophila* strains**		
act-GAL4	Zhaohui Wang	N/A
bam^{BG} (strong loss-of-function allele)	Zhaohui Wang	(*Chen and McKearin, 2005*)
bam^{Δ86} (null allele)	Michael Buszczak	(*Bopp et al., 1993*)

bamP-GFP	Ting Xie	(***Chen and McKearin, 2003b***)
bgcn[1] (null allele)	Bloomington *Drosophila* Stock Center, BDSC	6054
Canton-S	BDSC	64349
Dad-lacZ	Zheng Guo	(***Kai and Spradling, 2003***)
dpp[d6] (hypomorphic allele)	BDSC	2062
dpp[d12] (hypomorphic allele)	BDSC	2070
dpp(P4)-lacZ	Rongwen Xi	(***Li et al., 2016***)
EGFP FRT40A	BDSC	5629
FRT40A	BDSC	1816
FRT42D	BDSC	1802
FRT42D EGFP	BDSC	5626
FRT82B	BDSC	86313
FRT82B arm-lacZ	BDSC	7369
FRT82B EGFP	BDSC	32655
FRT82B RFP	BDSC	30555
gbb[1] (null allele)	BDSC	98344
hs-bam on chromosome 3R	this study	N/A
mad[12] (null allele)	BDSC	51301
med[1] (null allele)	BDSC	9033
nos-GAL4-VP16	Ruth Lehmann	(***Van Doren et al., 1998***)
P{bam+}	Shaowei Zhao	(***Zhang et al., 2023***)
punt[135] (strong loss-of-function allele)	BDSC	3100

UASp-dpp-RNAi-1	TsingHua Fly Center, THFC	TH201500984.S
UASp-dpp-RNAi-2	THFC	THU5880
UASp-FLP	Shaowei Zhao	(*Zhang et al., 2023*)
UASp-gbb-RNAi	THFC	THU1480
UASp-GFP	Michael Buszczak	N/A
UASp-yellow-RNAi	THFC	TH03150.N
UASz-FLP	Shaowei Zhao	(*Zhang et al., 2023*)
Kits		
ChamQ SYBR qPCR Master Mix	Vazyme	Q311
HiFiScript cDNA Synthesis Kit	CWBIO	CW2569M
RNeasy Micro Kit	Qiagen	74004
Plasmids		
bam cDNA clone	Berkeley *Drosophila* Genome Project	FI05606
pCaSpeR-hs	Eric Baehrecke	N/A
attB-pCaSpeR-hs-bam	this study	N/A
Software		
Adobe Photoshop 2022	San Jose, CA, USA	
ImageJ	NIH	
Prism	Graphpad Software, Inc.	
ZEN 3.0 SR imaging software	Carl Zeiss	

Data availability

All genotypes are described in **Source data 1**, and the raw quantification data are

included in **Source data 2**.

Fly husbandry

Flies were raised at 25°C on standard cornmeal/molasses/agar media.

Transgenic flies

hs-bam on chromosome 3R: The coding sequence of the *bam* gene, amplified from

the cDNA clone, was cloned into the *BglII-XbaI* sites of the *pCaSpeR-hs* vector (a gift

from Eric Baehrecke), while the *attB* sequence was inserted into the *XhoI* site. The

resulting *attB-pCaSpeR-hs-bam* plasmid was then microinjected into the *attP154*

(Chromosome 3R*, 97D2*) fly strain to generate site-specific transgenic flies.

Heatshock method to induce germline clones

To ensure developmental synchrony and maintain low-density growth, eggs within 8

hours of laying were collected for heatshock treatment. The animals (late-Larva 3/early-

Pupa stage) were subjected to twice-daily heatshocks at 37°C (2 hours per session, with

a 6-hour interval between the two sessions) for 6 consecutive days.

Fertility test

For each genotype, three independent crosses were performed. Each cross vial

contained two females and four w^{1118} (wild-type) males, all aged three days old. The

crosses were transferred to fresh vials every two days, with five replicate vials quantified

per genotype. After all adult flies eclosed, offspring production was assessed by counting

the number of empty pupae on the vial walls.

BrdU labeling

Ovaries were dissected in Schneider's insect medium (SIM) and incubated in freshly

prepared BrdU solution (100 µg/mL in SIM) for five hours at 25°C. After washing with

PBS (30 min), samples were fixed in 4% paraformaldehyde (PBS-diluted) for three

hours, followed by another PBS wash (30 min). Samples were then treated with RQ1

DNase reaction solution (Promega, Madison, WI, USA) for one hour, washed with PBST

(0.3% Triton X-100 in PBS, 30 min), and incubated overnight at 4°C with mouse anti-

BrdU antibody. Following a PBST wash (one hour), ovaries were incubated with goat

anti-mouse 546 and DAPI (0.1 µg/mL) in PBST for three hours, washed again in PBST

(one hour), and mounted in 70% glycerol (autoclaved).

Immuno-florescent staining, image collection, and data Processing

Ovaries were dissected in PBS, fixed in 4% paraformaldehyde (in PBS) for three hours,

washed with PBST (0.3% Triton X-100 in PBS) for 30 min, and then incubated overnight

at 4°C with primary antibodies diluted in PBST as follows: mouse anti-BamC (1:5),

mouse anti-β-Gal (1:200), mouse anti-α-Spectrin (1:100), rabbit anti-pMad (1:500), and

rabbit anti-Vasa (1:2000). After washing with PBST (one hour), samples were incubated

with Alexa Fluor-conjugated secondary antibodies (1:1000) and 0.1 µg/mL DAPI (in

PBST) for three hours, followed by a final PBST wash (one hour). Ovaries were mounted

in 70% glycerol and imaged using a Zeiss LSM 710 confocal microscope (Carl Zeiss AG,

Baden-Württemberg, Germany). Images were processed with ZEN 3.0 SR imaging

software (Carl Zeiss) and Adobe Photoshop 2022. The quantification data were

processed by Microsoft Excel or GraphPad Prism.

RT-qPCR

Ovaries from 14-day-old flies were dissected, and total RNA was extracted using the

RNeasy Micro Kit. Equal amounts of RNA were reverse-transcribed into cDNA using the

HiFiScript cDNA Synthesis Kit. RT-qPCR was performed on a CFX Connect Real-Time

PCR System (Bio-Rad) with ChamQ SYBR qPCR Master Mix. The PCR protocol

consisted of an initial denaturation at 95°C for 30 min, followed by 40 cycles of 95°C for

10 sec and 60°C for 30 sec. Relative gene expression was calculated using the $2^{-\Delta\Delta C_T}$

method (**Livak and Schmittgen, 2001**). The primers described previously were used

(**Huang et al., 2017**):

dpp primer-1: 5'-TACCACGCCATCCACTCAAC-3'

dpp primer-2: 5'-GCTCGTTACTCGATACGGCT-3'

gbb primer-1: 5'-CTGGATCATCGCACCAGAGG-3'

gbb primer-2: 5'-GTCTGGACGATCGCATGGTT-3'

rp49 (internal control) primer-1: 5'-CACCGGATTCAAGAAGTTCC-3'

rp49 (internal control) primer-2: 5'-GACAATCTCCTTGCGCTTCT-3'

Acknowledgements

We thank Eric Baehrecke, Michael Buszczak, Zheng Guo, Ed Laufer, Ruth Lehmann,

Rongwen Xi, Ting Xie, Zhaohui Wang, BDGP, BDSC, DSHB, and THFC for providing

antibodies, plasmids, and fly strains. This study was supported by grants 32270841 and

32070871 from the National Natural Science Foundation of China (NSFC) to Shaowei

Zhao.

Author contributions

S.Z. conceived and supervised this study. Y.Z., Y.W., J.S., L.Y., Z.W., D.S., Y.D.Z., and

S.Z. performed the experiments. S.Z. wrote the manuscript and all authors commented

on it.

Declaration of interests

The authors declare no competing interests.

References

Blanpain, C., and Fuchs, E. (2006). Epidermal stem cells of the skin. Annu Rev Cell Dev
Biol *22*, 339-373.
Bopp, D., Horabin, J.I., Lersch, R.A., Cline, T.W., and Schedl, P. (1993). Expression of
the Sex-lethal gene is controlled at multiple levels during Drosophila oogenesis.
Development *118*, 797-812.
Chen, D., and McKearin, D. (2003a). Dpp signaling silences bam transcription directly to
establish asymmetric divisions of germline stem cells. Curr Biol *13*, 1786-1791.
Chen, D., and McKearin, D. (2005). Gene circuitry controlling a stem cell niche. Curr Biol
15, 179-184.
Chen, D., and McKearin, D.M. (2003b). A discrete transcriptional silencer in the bam
gene determines asymmetric division of the Drosophila germline stem cell. Development
130, 1159-1170.
Chen, D., Wu, C., Zhao, S., Geng, Q., Gao, Y., Li, X., Zhang, Y., and Wang, Z. (2014).
Three RNA binding proteins form a complex to promote differentiation of germline stem
cell lineage in Drosophila. PLoS Genet *10*, e1004797.
Fuller, M.T., and Spradling, A.C. (2007). Male and female Drosophila germline stem
cells: two versions of immortality. Science *316*, 402-404.
Gehart, H., and Clevers, H. (2019). Tales from the crypt: new insights into intestinal stem
cells. Nat Rev Gastroenterol Hepatol *16*, 19-34.
Germani, F., Bergantinos, C., and Johnston, L.A. (2018). Mosaic Analysis in Drosophila.
Genetics *208*, 473-490.
Huang, J., Reilein, A., and Kalderon, D. (2017). Yorkie and Hedgehog independently
restrict BMP production in escort cells to permit germline differentiation in the Drosophila
ovary. Development *144*, 2584-2594.
Jin, Z., Kirilly, D., Weng, C., Kawase, E., Song, X., Smith, S., Schwartz, J., and Xie, T.
(2008). Differentiation-defective stem cells outcompete normal stem cells for niche
occupancy in the Drosophila ovary. Cell Stem Cell *2*, 39-49.
Jogi, A., Vaapil, M., Johansson, M., and Pahlman, S. (2012). Cancer cell differentiation
heterogeneity and aggressive behavior in solid tumors. Ups J Med Sci *117*, 217-224.
Kai, T., and Spradling, A. (2003). An empty Drosophila stem cell niche reactivates the
proliferation of ectopic cells. Proc Natl Acad Sci U S A *100*, 4633-4638.
Kai, T., Williams, D., and Spradling, A.C. (2005). The expression profile of purified
Drosophila germline stem cells. Dev Biol *283*, 486-502.
Lavoie, C.A., Ohlstein, B., and McKearin, D.M. (1999). Localization and function of Bam
protein require the benign gonial cell neoplasm gene product. Dev Biol *212*, 405-413.
Li, X., Yang, F., Chen, H., Deng, B., Li, X., and Xi, R. (2016). Control of germline stem
cell differentiation by Polycomb and Trithorax group genes in the niche
microenvironment. Development *143*, 3449-3458.
Li, Y., Minor, N.T., Park, J.K., McKearin, D.M., and Maines, J.Z. (2009). Bam and Bgcn
antagonize Nanos-dependent germ-line stem cell maintenance. Proc Natl Acad Sci U S
A *106*, 9304-9309.
Lin, H. (1997). The tao of stem cells in the germline. Annu Rev Genet *31*, 455-491.
Lin, H., Yue, L., and Spradling, A.C. (1994). The Drosophila fusome, a germline-specific
organelle, contains membrane skeletal proteins and functions in cyst formation.
Development *120*, 947-956.
Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.
Lytle, N.K., Barber, A.G., and Reya, T. (2018). Stem cell fate in cancer growth,
progression and therapy resistance. Nat Rev Cancer *18*, 669-680.

Mathieu, J., Michel-Hissier, P., Boucherit, V., and Huynh, J.R. (2022). The deubiquitinase
USP8 targets ESCRT-III to promote incomplete cell division. Science *376*, 818-823.
McKearin, D., and Ohlstein, B. (1995). A role for the Drosophila bag-of-marbles protein in
the differentiation of cystoblasts from germline stem cells. Development *121*, 2937-2947.
McKearin, D.M., and Spradling, A.C. (1990). bag-of-marbles: a Drosophila gene required
to initiate both male and female gametogenesis. Genes Dev *4*, 2242-2251.
Niki, Y., and Mahowald, A.P. (2003). Ovarian cystocytes can repopulate the embryonic
germ line and produce functional gametes. Proc Natl Acad Sci U S A *100*, 14042-14045.
Ohlstein, B., Lavoie, C.A., Vef, O., Gateff, E., and McKearin, D.M. (2000). The
Drosophila cystoblast differentiation factor, benign gonial cell neoplasm, is related to
DExH-box proteins and interacts genetically with bag-of-marbles. Genetics *155*, 1809-
1819.
Ohlstein, B., and McKearin, D. (1997). Ectopic expression of the Drosophila Bam protein
eliminates oogenic germline stem cells. Development *124*, 3651-3662.
Pastor-Pareja, J.C., and Xu, T. (2013). Dissecting social cell biology and tumors using
Drosophila genetics. Annu Rev Genet *47*, 51-74.
Song, X., Wong, M.D., Kawase, E., Xi, R., Ding, B.C., McCarthy, J.J., and Xie, T. (2004).
Bmp signals from niche cells directly repress transcription of a differentiation-promoting
gene, bag of marbles, in germline stem cells in the Drosophila ovary. Development *131*,
1353-1364.
Sousa-Victor, P., García-Prat, L., and Muñoz-Cánoves, P. (2022). Control of satellite cell
function in muscle regeneration and its disruption in ageing. Nat Rev Mol Cell Biol *23*,
204-226.
Spradling, A., Fuller, M.T., Braun, R.E., and Yoshida, S. (2011). Germline stem cells.
Cold Spring Harb Perspect Biol *3*, a002642.
Van Doren, M., Williamson, A.L., and Lehmann, R. (1998). Regulation of zygotic gene
expression in Drosophila primordial germ cells. Curr Biol *8*, 243-246.
Wilkinson, A.C., Igarashi, K.J., and Nakauchi, H. (2020). Haematopoietic stem cell self-
renewal in vivo and ex vivo. Nat Rev Genet *21*, 541-554.
Xie, T., and Spradling, A.C. (1998). decapentaplegic is essential for the maintenance and
division of germline stem cells in the Drosophila ovary. Cell *94*, 251-260.
Xie, T., and Spradling, A.C. (2000). A niche maintaining germ line stem cells in the
Drosophila ovary. Science *290*, 328-330.
Zhang, Q., Li, L., Zhang, Q., Zhang, Y., Yan, L., Wang, Y., Wang, Y., and Zhao, S. (2024).
Genetic circuitry controlling Drosophila female germline overgrowth. Dev Biol *515*, 160-
168.
Zhang, Q., Zhang, Y., Zhang, Q., Li, L., and Zhao, S. (2023). Division promotes adult
stem cells to perform active niche competition. Genetics *224*.
Zhao, S., Fortier, T.M., and Baehrecke, E.H. (2018). Autophagy Promotes Tumor-like
Stem Cell Niche Occupancy. Curr Biol *28*, 3056-3064.e3053.
475

 Figures and figure legends

Figure 1. *bam* or *bgcn* **mutant germline tumors inhibit the differentiation of neighboring wild-type GSCs.** (A) Schematic cartoon for early oogenesis. The red dots and branches indicate spectrosomes and fusomes, respectively. GSC: germline stem cell. (B) Mosaic analysis strategy. The FLP recombinase triggers mitotic recombination by targeting *FRT* sequences. The *nos>FLP* method restricts FLP expression to the germline, while the *hs-FLP* method enables heatshock-inducible FLP expression. (C-F) Representative samples. The asterisks mark the niche cap cells, and the arrows indicate SGCs that have exited the niche and are surrounded by germline tumors mutant for *bam* or *bgcn*. Vasa, a germ cell marker, should label all germ cells. However, due to poor tumor permeability, staining often fails to detect tumorous germ cells in the central region (see Vasa panels in D-F). (G-I) Representative samples (z-stack projections). In (G), the

arrowheads and arrow respectively mark two GSCs and one cystoblast, all containing

dot-like spectrosomes, while the dotted lines delineate cystocytes with branched

fusomes. In (H) and (I), the arrows denote SGCs that also contain dot-like

spectrosomes, akin to GSCs and the adjacent GSC-like tumor cells.

The online version of this article includes the following source data and figure

supplement for figure 1:

Source data 1. All genotypes.

Figure supplement 1. SGCs appear in egg chambers.

Figure 2. The inhibition of SGC differentiation depends on the lack of Bam

expression. (A) Representative sample. The arrowhead marks a BamC-positive 4-

cystocyte germline cyst, while the arrows indicate BamC-negative SGCs. (B)

Representative sample. The asterisk denotes niche cap cells, and the dotted circles

outline *bamP*-GFP-negative GSCs. The solid circle marks a *bamP*-GFP-positive

cystoblast. The arrow and arrowhead point to *bamP*-GFP-negative and -positive SGCs,

respectively. (C) Quantification data. 14-day-old flies were used for the analyses. CBs:

cystoblasts. (D) Schematic of the experimental strategy for (E-H). In "*with hs-bam*" flies

(E and G), wild-type germ cells (both *bam*$^{+/+}$ and *bam*$^{+/-}$) carry the *hs-bam* transgene,

while control "*without hs-bam*" flies (F) lack this element in their wild-type germ cells. (E-

G) Representative samples. The arrows mark SGCs with dot-like spectrosomes, while

the arrowhead indicates a 4-cystocyte germline cyst containing branched fusomes. (H)

Quantification data. For each experiment, three independent replicates were performed,

with over 100 SGCs and germline cysts quantified per replicate. Data represent mean ±

SEM, and statistical significance was determined by t test. *n.s.* ($P > 0.05$).

The online version of this article includes the following source data for figure 2:

Source data 1. All genotypes.

Source data 2. Raw quantification data.

Figure 3. SGCs maintain lower BMP signaling levels than GSCs within the niche.
(A and B) Representative samples. The asterisks mark niche cap cells, arrowheads indicate pMad-positive GSCs, and arrows point to pMad-negative SGCs. (C) Representative samples. The asterisks denote niche cap cells, arrowheads mark *Dad*-lacZ-positive GSCs, and arrows highlight *Dad*-lacZ-positive SGCs. The dotted cycles outline one *Dad*-lacZ-negative SGC. (D and E) Quantification data. 14-day-old flies were used for the analyses. In (E), data represent mean ± SEM, and statistical significance was determined by t test. (F) Representative sample. The asterisk marks the niche cap cell, while the arrows indicate a BrdU+ GSC within the niche. (G) Representative sample. The arrow indicates a BrdU+ SGC surrounded by germline tumors. (H) Quantification data. 14-day-old flies were used for the analyses. Statistical significance was determined by chi-squared test.

The online version of this article includes the following source data for figure 3:

Source data 1. All genotypes.

531 **Source data 2.** Raw quantification data.

Figure 4. BMP signaling inhibits SGC differentiation. (A) Schematic of the

experimental strategy for (B-F). Genotypes were unambiguously distinguished using a

triple-color system (red, yellow, and green). (B-D) Representative samples. The dotted

cycles mark an SGC, while the solid lines outline germline cysts containing

differentiating cystocytes. (E and F) Quantification data. 14-day-old flies were used for

the analyses. (G) Schematic of the experimental strategy for (H-J). (H and I)

Representative samples. The dotted lines mark an SGC, while the solid lines outline a

germline cyst containing differentiating cystocytes. (J) Quantification data. 14-day-old

flies were used for the analyses. For each experiment, three independent replicates

were performed, with over 100 SGCs and germline cysts quantified per replicate. Data

represent mean ± SEM, and statistical significance was determined by t test.

The online version of this article includes the following source data for figure 4:

Source data 1. All genotypes.

Source data 2. Raw quantification data.

Figure 5. Germline tumors secret the BMP ligands. (A and B) Representative

samples. In (A), the asterisks denote *dpp*-lacZ-high niche cap cells, and solid lines

outline GFP-negative *bam* mutant germline tumor cells. In (B), the dotted lines highlight

GFP-positive wild-type cystocytes. (C) Quantification data. 14-day-old flies were used for

the analyses. (D and E) Quantification data for RT-qPCR assays. 14-day-old flies were

used for the analyses. For each experiment, three independent replicates were

performed. Statistical significance in (C) was determined by one-way ANOVA and in (D

and E) by t test.

The online version of this article includes the following source data for figure 5:

Source data 1. All genotypes.

Source data 2. Raw quantification data.

Figure 6. Dpp and Gbb secreted by germline tumors are required to inhibit SGC differentiation. (A) Schematic of the experimental strategy for (C-F). (B) Schematic of the experimental strategy for (G-I). (C-E, G, and H) Representative samples. The arrows mark SGCs containing dot-like spectrosomes, while the arrowheads denote germline cysts with differentiating cystocytes that possess branched fusomes. (F and I) Quantification data for the SGC phenotype. 14-day-old flies were used for the analyses. For each experiment, three independent replicates were performed, with over 100 SGCs and germline cysts quantified per replicate. Data represent mean ± SEM. Statistical significance in (F) was determined by one-way ANOVA and in (I) by t test.

The online version of this article includes the following source data and figure

supplement for figure 6:

Source data 1. All genotypes.

Source data 2. Raw quantification data.

Figure supplement 1. Monoallelic deletion of *dpp* or *gbb* does not affect GSC

maintenance, germ cell differentiation, and female fly fertility.

Figure supplement 2. *dpp bam* or *gbb bam* double-mutant germline tumor cells divide

more slowly than *bam* single-mutant ones.

Figure supplement 3. The SGC phenotype is unchanged irrespective of the number of

surrounding germline tumors.

Figure 7. A working model. *bam* or *bgcn* mutant germline tumors secrete the BMP ligands Dpp and Gbb to activate BMP signaling in out-of-niche GSCs (called SGCs in this study) to inhibit their differentiation (left panel). In contrast, *dpp bam* and *gbb bam* double-mutant germline tumors exhibit a significant loss of this differentiation-inhibiting ability (right panel).

Figure 1-figure supplement 1. SGCs appear in egg chambers. (A, C) Representative

samples. The arrows indicate SGCs enclosed within egg chambers. (B) Schematic of

the experimental strategy for (C). See also **Source data 1**.

Figure 6-figure supplement 1. Monoallelic deletion of *dpp* or *gbb* does not affect

GSC maintenance, germ cell differentiation, and female fly fertility. (A-D)

Representative samples. (E) Quantification data. 14-day-old flies were used for the

analyses. GSCs are germ cells that are located within the niche and contain dot-like

spectrosomes. Cystoblasts are germ cells that have exited the niche, remain in contact

with GSCs, and maintain dot-like spectrosomes. Cystocytes in germline cysts are germ

cells that are characterized by branched fusomes. (F) Fertility test. 3-day-old flies were

used for the analyses. For each genotype, three independent replicates were performed.

Data represent mean ± SEM, and statistical significance was determined by one-way

ANOVA. *n.s.* ($P > 0.05$). See also **Source data 1 and 2**.

Figure 6-figure supplement 2. *dpp bam* or *gbb bam* double-mutant germline tumor cells divide more slowly than *bam* single-mutant ones. (A, B, D, and E) Representative samples. The arrows indicate BrdU⁺ germline tumor cells mutant for *bam*, *dpp bam*, or *gbb bam*. (C and F) Quantification data. 14-day-old flies were used for the analyses. Statistical significance was determined by chi-squared test. See also **Source data 1 and 2**.

607

Figure 6-figure supplement 3. The SGC phenotype is unchanged irrespective of

the number of surrounding germline tumors. (A) Representative samples. The

arrows denote SGCs. Both images are of the same magnification. (B) Quantification

data for tumor size. Data represent mean ± SEM. (C) Quantification data for the SGC

phenotype. In (B and C), 14-day-old flies were used for the analyses. Statistical

significance in (B) was determined by t test and in (C) by chi-squared test. *n.s.* (P >

0.05). See also **Source data 1 and 2**.