Gli3R-mediated inhibition of hedgehog signaling alters the embryonic transcriptome in zebrafish

Anna J. Moyer[1], Summer B. Thyme[1*]

[1] Department of Biochemistry and Molecular Biotechnology, The University of Massachusetts

Chan Medical School, Worcester, MA 01605, USA

* Corresponding author

Email: Summer.Thyme@umassmed.edu

Keywords: zebrafish, embryonic development, RNA-seq, hedgehog, Gli3R, Gli, ionocytes, ion homeostasis, Foxi3, H+-ATPase-rich

ABSTRACT

Hedgehog signaling is a conserved developmental pathway that patterns diverse tissues during

vertebrate embryogenesis. In zebrafish, disruptions to the hedgehog pathway cause well-

characterized defects in specific cell types including neurons and glia derived from the ventral

neural tube. We inhibited hedgehog signaling by overexpressing the Gli3 repressor ubiquitously

and performed bulk RNA-seq of 30 hours post-fertilization zebrafish embryos. Consistent with

known roles of hedgehog signaling, we observed reduced expression of genes marking lateral

floor plate, motor neurons, Kolmer-Agduhr cells, dopaminergic neurons, slow muscle cells, and

anterior pituitary. Gene set enrichment analysis using marker genes derived from the Daniocell

atlas also revealed downregulation of genes marking H+-ATPase-rich and Na+-K+-ATPase-rich

ionocytes, which are located in the embryonic skin and are responsible for osmotic homeostasis.

Reduced expression of ionocyte-specific transporter genes and the transcription factor *foxi3a*

suggests that Gli activity may play a previously unrecognized role in the development of this cell

type.

INTRODUCTION

The hedgehog signaling pathway plays a critical role in vertebrate patterning (Ingham

and McMahon 2001) and is also implicated in regeneration (Imokawa and Yoshizato 1997) and

oncogenesis (Taipale et al. 2000). Depending on the context, hedgehog can act as a morphogen

(Heemskerk and DiNardo 1994) or mitogen (Wechsler-Reya and Scott 1999) and specifies

multiple distinct cell types across organ systems. In the canonical pathway, hedgehog ligands

(Sonic, Indian, and Desert hedgehog) (Echelard et al. 1993) bind to the receptor Patched (Marigo

et al. 1996; Stone et al. 1996), which relieves inhibition of the transducer Smoothened (Chen et

al. 2001; Zhang et al. 2001). Activation of Smoothened shifts the balance between repressor and

activator forms of the Gli transcription factors, which in turn modulates the transcription of

target genes (Ruiz i Altaba 1999; Wang et al. 2000).

Experimental manipulation of hedgehog in the zebrafish model system has revealed

essential roles in development (Hammerschmidt et al. 1996; Huang and Schier 2009; Shen et al.

2013). In the presumptive spinal cord, hedgehog secreted by the notochord and medial floor plate

exposes the neighboring lateral floor plate (p3), pMN, and p0-p2 progenitor domains to a

gradient of hedgehog levels, which contributes to specification of neural cell types including

interneurons, motor neurons, Kolmer-Agduhr cells, oligodendrocytes, and astroglia [reviewed in

(Danesin and Soula 2017)]. In addition to gross morphological phenotypes like cyclopia, ventral

tail curvature, and rounded somites, inhibition of hedgehog in zebrafish can expand or reduce

these ventral spinal cord progenitor domains and their derivative cells (Brand et al. 1996; Chen et

al. 2001; Karlstrom et al. 1996).

However, subtle differences in phenotypes result from the specific method used to disrupt

hedgehog signaling (i.e., mutations, morpholinos, overexpression of Gli repressors, and chemical

inhibitors). These phenotypes may vary due to compensation between paralogs, incomplete or

off-target effects of antagonists, maternal contribution of mRNA in homozygous mutants, and

non-canonical or hedgehog ligand-independent Gli activity. For example, knockdown of *gli1* or

gli2b in maternal-zygotic *smoothened* mutants reduces motor neuron number compared to

mutants, suggesting that basal Gli activity contributes to motoneurogenesis (Mich and Chen

2011). Similarly, overexpression of the Gli3 repressor (Gli3R) in embryos treated with the

Smoothened antagonist cyclopamine reduces the number of motor neurons compared to

smoothened mutants (England et al. 2011). These experiments highlight how changes in the

expression and processing of Gli can modify specification of cells in the ventral spinal cord even

in the absence of upstream hedgehog signaling.

While previous work has manipulated Gli activity in targeted phenotypic assays, the

transcriptome-wide consequences of Gli3R overexpression in zebrafish remain unexplored. To

address this gap, we overexpressed a zebrafish *Gli3R* (*zGli3R*) transgene (Huang and Schier

2009) ubiquitously and performed bulk RNA sequencing (RNA-seq) of 30 hours post-

fertilization (hpf) zebrafish embryos. Our findings suggest that Gli activity may be involved in

the development of ionocytes. We anticipate that our dataset will be most relevant to zebrafish

and developmental biologists who are interested in hedgehog signaling and cell-type

specification during vertebrate development.

MATERIALS & METHODS

Zebrafish husbandry and transgenesis

Animal experiments were approved by the UMass Chan Institutional Animal Care and

Use Committee (IACUC protocol 202300000053). Transgenic animals were generated in an

Ekkwill (EK)-based wildtype strain, and both larvae and adult animals were maintained on a 14

hour/10 hour light/dark cycle at 28°C. The *zGli3R* transgene was subcloned from *pT2-hsp-*

zGli3R-EGFP into Gateway *pDONR221* (Invitrogen) and then into a Gateway-compatible *Tol2*

destination vector containing the *ubb* promoter (Mosimann et al. 2011) and the *myl7:GFP*

transgenesis marker. The annotated plasmid map including full sequence is available in

Supplementary File 1. To produce transgenic animals using *Tol2*, one-cell embryos were injected

with 1 nL of 10 ng/µL *Tol2* mRNA (Kawakami et al. 2004) and 20 ng/µL plasmid. F0 embryos

with expression of the *myl7:GFP* transgenesis marker were grown to adults.

Sample collection

A mosaic F0 male was crossed to a wildtype female, and F1 embryos from one clutch

were collected at 30 hpf for RNA-seq. Transgenic embryos and wildtype control siblings were

sorted based on the *myl7:GFP* transgenesis marker. Four dechorionated embryos were pooled

and frozen on dry ice for each biological replicate, and four biological replicates were collected

per genotype. For live imaging, 2 days post-fertilization (dpf) and 3 dpf animals were

anesthetized with MS-222 and live mounted in 2% low-melt agarose (Fisher Bioreagents,

BP165-25) made in embryo medium with methylene blue. Images were obtained with a Leica

M165 FC stereo microscope and K3C color camera.

RNA-seq

RNA extraction, quality control, library preparation, and sequencing were conducted at

Azenta Life Sciences (South Plainfield, NJ, USA) as follows: Total RNA was extracted from

frozen tissue samples using Qiagen RNeasy Plus Universal mini kit following manufacturer's

instructions (Qiagen, Hilden, Germany). RNA samples were quantified using Qubit 3.0

Fluorometer (ThermoFisher Scientific, Waltham, MA, USA) and RNA integrity was checked

with 4200 TapeStation (Agilent Technologies, Palo Alto, CA, USA).

RNA-seq libraries were prepared using the NEBNext Ultra II RNA Library Prep Kit for

Illumina using manufacturer's instructions (NEB, Ipswich, MA, USA). Briefly, mRNAs were

initially enriched with Oligo d(T) beads. Enriched mRNAs were fragmented for 15 minutes at

94 °C. First strand and second strand cDNA were subsequently synthesized. cDNA fragments

were end repaired and adenylated at 3′ ends, and universal adapters were ligated to cDNA

fragments, followed by index addition and library enrichment by PCR with limited cycles. The

sequencing library was validated on the Agilent TapeStation (Agilent Technologies, Palo Alto,

CA, USA), and quantified by using Qubit 3.0 Fluorometer (Invitrogen, Carlsbad, CA) as well as

by quantitative PCR (KAPA Biosystems, Wilmington, MA, USA).

The sequencing libraries were multiplexed and clustered onto a flowcell on the Illumina

NovaSeq instrument according to manufacturer's instructions. The samples were sequenced

using a 2x150bp Paired End (PE) configuration. Image analysis and base calling were conducted

by the NovaSeq Control Software (NCS). Raw sequence data (.bcl files) generated from Illumina

NovaSeq was converted into fastq files and de-multiplexed using Illumina bcl2fastq 2.20

software. One mismatch was allowed for index sequence identification.

RNA-seq analysis

Alignment and differential gene expression analysis of RNA-seq data were performed as

previously described (Capps et al. 2025). Reads were aligned with the STAR aligner (2.7.3a-

GCC 6.4.0-2.28) (Dobin et al. 2013) to GRCz11 release 104 using the Zebrafish Transcriptome

Annotation version 4.3.2 (Lawson et al. 2020). Uniquely mapped reads ranged from 21.5-29.9M

per sample. Transcripts with zero counts in four or more of eight total samples were removed,

and raw counts were normalized using rlog counts in DESeq2 (Love et al. 2014). The default

DESeq2 method was used for determining significance, and raw and adjusted (Benjamini-

Hochberg) P values are in Supplementary Table 1. DESeq2 script is available on GitHub:

https://github.com/thymelab/BulkRNASeq.

Downstream analysis of bulk RNA-seq data was conducted using R (Version 4.4.2).

Gene set enrichment analysis (GSEA) was performed as previously described (Capps et al. 2025)

using the 'GSEA' function from clusterProfiler (Version 4.14.6) (Wu et al. 2021). For

chromosomal location GSEA, the gene set was generated using RefSeq Genes and Gene

Predictions from the GRCz11 assembly, and genes were sorted by abs(log2 fold change) before

GSEA. For Daniocell GSEA, the gene set was generated as previously described (Dang et al.

2025) and genes were sorted by log2 fold change before GSEA. Volcano plot, GSEA bar plot,

and individual gene bar plots were created using ggplot2 (Version 3.5.2) (Wickham 2016).

Heatmap was generated using the 'heatmap.2' function from gplots (Version 3.2.0) (Warnes et

al. 2025). Code to generate figures is available in Supplementary File 2.

RESULTS

Fig. 1. Morphological and transcriptomic phenotypes of *Tg(ubb:zGli3R)* embryos. a) Representative transgenic and wildtype animals at 2 dpf and 3 dpf. Scale bar = 200 μm. b) Volcano plot comparing gene expression in 30 hpf *Tg(ubb:zGli3R)* embryos and wildtype siblings. Genes with -log10(padj) greater than ten are plotted at ten. Transcripts encoded by chromosome 4 are denoted with X symbols and transcripts encoded by all other chromosomes are denoted with circles. n=4 replicates of 4 pooled embryos for transgenic and 4 replicates of 4 pooled embryos for wildtype siblings. c) Heat map displaying expression of selected marker genes. From left to right, genes mark progenitor domains and interneurons in the ventral spinal cord, Kolmer-Agduhr cells, motor neurons, radial glia/astrocytes, dopaminergic neurons, slow muscle cells, and anterior pituitary. d) Cell types identified with GSEA using gene sets from the Daniocell single-cell atlas (Sur et al. 2023). Red bars represent terms with a negative normalized enrichment score and blue bars represent terms with a positive score. e) Bar plots of normalized counts for genes expressed in ionocytes. P values and padj for all genes are available in Supplementary Table 1.

Differential gene expression in *Tg(ubb:zGli3R)* embryos

We cloned *zGli3R* downstream of the ubiquitin B (*ubb*) promoter and co-injected F0 embryos with *Tol2* to produce transgenic lines. Transgenic F1 offspring possessed severe

morphological defects and died at larval stages (Fig. 1a). We isolated a mosaic F0 parent with

high germline transmission and performed bulk RNA-seq on 30 hpf embryos pooled from a

single clutch. Analysis of differentially expressed genes identified 888 upregulated genes and

1018 downregulated genes at a threshold of P adjusted (padj) <0.05 (Fig. 1b; Supplementary

Table 1). Transgene insertion can cause enrichment of differentially expressed genes based on

chromosomal location (White et al. 2022), and we observed a significant enrichment of genes on

chromosome 4 (Supplementary Table 2).

We next examined the expression of marker genes for cell types known to involve

hedgehog signaling (Fig. 1c; Supplementary Table 1). Progenitor domains in the ventral spinal

cord give rise to specific subtypes of neurons and glia, including interneurons, motor neurons,

Kolmer-Agduhr cells, oligodendrocyte precursor cells (OPCs), and astrocytes [Reviewed in

(Cucun et al. 2024; Danesin and Soula 2017)]. Genes marking most of these cell populations

were downregulated or unchanged, but *gfap*, which is expressed in radial glia and astrocyte-like

cells (Agius et al. 2004; Marcus and Easter 1995), was upregulated. Consistent with the

literature, marker genes for dopaminergic neurons (Chen et al. 2001; Holzschuh et al. 2003),

slow muscle cells (Blagden et al. 1997), and anterior pituitary (adenohypophysis) (Herzog et al.

2003) were also downregulated.

Functional enrichment analysis of cell types using Daniocell single-cell atlas

GSEA using gene sets derived from single-cell expression data can reveal potential

changes in cell composition (Capps et al. 2025; Dang et al. 2025; Moyer et al. 2025). GSEA with

gene sets derived from the Daniocell single-cell atlas (Sur et al. 2023) highlighted 28 clusters

with negative or positive enrichment compared to control (Fig. 1d; Supplementary Table 3).

Several of the identified clusters with negative enrichment scores, including floor plate (glia.15

and glia.2), Kolmer-Agduhr cells (glia.30), spinal OPCs (glia.20 and glia.26), dopaminergic

neurons of the diencephalon (neur.34), and anterior pituitary (tast.10) are consistent with the

known roles of hedgehog during zebrafish development. Other cell types with negative

enrichment scores included ventrolateral mesoderm (hema.20), hemangioblasts (hema.23), type

III taste buds (tast.2), and multiple subtypes of H+-ATPase-rich ionocytes and ionocyte

progenitors (iono.17, iono.15, iono.10, and iono.12). In contrast, olfactory sensory neurons

(tast.9), lymphoid hematopoietic stem cells (hema.15), and primitive erythroblasts (hema.32)

showed positive enrichment scores. Further investigation of known markers of ionocytes

revealed downregulation of the master regulators *foxi3a* (Hsiao et al. 2007) and *gcm2* (Esaki et

al. 2009; Shono et al. 2011) as well as genes expressed in H+-ATPase-rich ionocytes (*slc4a1b*,

atp6v0a1a, *ca15a*, *ceacam1*, and *si:dkey-192d15.2*), and Na+-K+-ATPase-rich ionocytes (*trpv6*

and *s100a11*) (Fig. 1e) (Sur et al. 2023).

DISCUSSION

Although activation of hedgehog signaling removes Gli3R repressor activity in some

contexts (Litingtung and Chiang 2000; Litingtung et al. 2002), *gli3* does not appear to be

expressed before 12 hpf in zebrafish (England et al. 2011; Sur et al. 2023), and activator forms of

Gli may possess basal levels of activity that are hedgehog ligand independent (Huang and Schier

2009; Karlstrom et al. 2003; Mich and Chen 2011). In this study, we explored the transcriptomic

consequences of overexpressing *zGli3R* ubiquitously from the earliest stages of embryonic

development. In contrast to experimental manipulations that inhibit hedgehog signaling, such as

mutations in *smoothened* or cyclopamine treatment, ectopic expression of *zGli3R* has the

potential to shift the balance of Gli activator and repressor forms even in the absence of

hedgehog signaling.

We first examined marker genes for cell types previously shown to depend on hedgehog

signaling (Fig. 1c) and then applied unbiased GSEA with Daniocell terms to discover new cell

types that may be regulated by Gli (Fig. 1d). The validity of our experimental approach is

supported by misexpression of known class II genes, which are activated by hedgehog signaling

and include *nkx6.1*, *nkx6.2*, and *olig2* (Fig. 1c) (Briscoe et al. 2000; England et al. 2011; Guner

and Karlstrom 2007). Genes marking Kolmer-Agduhr cells (*sst1.1*, *urp1*, and *urp2*), motor

neurons (*phox2a* and *isl2a*), dopaminergic neurons (*th* and *slc6a3*), slow muscle cells (*notum2*

and *tnnt2d*), and anterior pituitary (*pomcb*) were also significantly downregulated (Sur et al.

2023). GSEA using Daniocell gene sets identified several of the same populations, including

anterior pituitary (tast.10), dopaminergic neurons (neur.34), and Kolmer-Agduhr cells (glia.30).

However, GSEA also nominated cell types with less literature support for hedgehog involvement

in zebrafish, including taste buds (tast.2), hemangioblasts (hema.23), and H+-ATPase-rich

ionocytes (iono.17, iono.15, iono.10, and iono.12).

Ionocytes are specialized cells found in the embryonic skin that respond to osmotic

changes in the aquatic environment and are functionally analogous to amniote renal tubular cells

[Reviewed in (Hwang and Chou 2013)]. Progenitors derived from non-neural ectoderm give rise

to both ionocytes and keratinocytes (Janicke et al. 2007). Expression of the transcription factors

foxi3a and *foxi3b* in ionocyte progenitors activates Notch1 in neighboring cells, which in turn

suppresses ionocyte cell fate and promotes keratinocyte development (Esaki et al. 2009; Hsiao et

al. 2007; Janicke et al. 2007). The transcription factor *gcm2* also plays a role in the specification

of ionocytes (Chang et al. 2009), and we observed reduced expression of *foxi3a* and *gcm2* but

not of *foxi3b* (Fig. 1e).

Although hedgehog signaling has not been implicated in the specification of zebrafish

ionocytes, it does regulate *gcm2* expression during development of the amniote parathyroid

(Grevellec et al. 2011; Moore-Scott and Manley 2005). Moreover, the Daniocell single-cell atlas

shows detectable expression of hedgehog components *smo*, *ptch2*, *gli1*, *gli2a*, *gli2b*, and *gli3* in

non-neural ectoderm (neur.26). However, a recent study that used single-cell RNA-seq to track

changes in cell abundance reported that cyclopamine treatment increases the abundance of non-

keratinocyte epidermal progenitors and Na+-K+-ATPase-rich ionocytes (Barkan et al. 2025).

Future hypothesis-based approaches will determine whether *zGli3R* overexpression affects

ionocyte development via hedgehog signaling or by suppressing basal Gli activity.

DATA AVAILABILITY STATEMENT

Plasmids are available upon request. Raw RNA-seq data and gene counts have been deposited in

GEO (GSE307979). Processed data are available in Supplementary Table 1.

ACKNOWLEDGEMENTS

The authors thank Dr. Peng Huang for providing the *pT2-hsp-zGli3R-EGFP* plasmid and

research technician Vaishnavi Balaji for experimental support.

FUNDING

This work was supported by NIH grant R01 HD115159 to S.B.T. and Jérôme Lejeune

Foundation postdoctoral fellowship to A.J.M.

LITERATURE CITED
Agius E, Soukkarieh C, Danesin C, Kan P, Takebayashi H, Soula C, Cochard P. 2004. Converse
control of oligodendrocyte and astrocyte lineage development by sonic hedgehog in the
chick spinal cord. Dev Biol. 270(2):308-321.
Barkan E, Duran M, Lammers N, Tresenrider A, Jackson D, Lee H, Haagen B, Saunders L,
Abitua P, Kimelman D et al. 2025. Embryo-scale single-cell chemical transcriptomics
reveals dependencies between cell types and signaling pathways. bioRxiv.
Blagden CS, Currie PD, Ingham PW, Hughes SM. 1997. Notochord induction of zebrafish slow
muscle mediated by sonic hedgehog. Genes Dev. 11(17):2163-2175.
Brand M, Heisenberg CP, Warga RM, Pelegri F, Karlstrom RO, Beuchle D, Picker A, Jiang YJ,
Furutani-Seiki M, van Eeden FJ et al. 1996. Mutations affecting development of the
midline and general body shape during zebrafish embryogenesis. Development. 123:129-
142.
Briscoe J, Pierani A, Jessell TM, Ericson J. 2000. A homeodomain protein code specifies
progenitor cell identity and neuronal fate in the ventral neural tube. Cell. 101(4):435-445.
Capps MES, Moyer AJ, Conklin CL, Martina V, Torija-Olson EG, Klein MC, Gannaway WC,
Calhoun CCS, Vivian MD, Thyme SB. 2025. Disrupted diencephalon development and
neuropeptidergic pathways in zebrafish with autism-risk mutations. Proc Natl Acad Sci U
S A. 122(23):e2402557122.

Chang WJ, Horng JL, Yan JJ, Hsiao CD, Hwang PP. 2009. The transcription factor, glial cell
missing 2, is involved in differentiation and functional regulation of h+-atpase-rich cells
in zebrafish (danio rerio). Am J Physiol Regul Integr Comp Physiol. 296(4):R1192-1201.
Chen W, Burgess S, Hopkins N. 2001. Analysis of the zebrafish smoothened mutant reveals
conserved and divergent functions of hedgehog activity. Development. 128(12):2385-
2396.
Cucun G, Kohler M, Pfitsch S, Rastegar S. 2024. Insights into the mechanisms of neuron
generation and specification in the zebrafish ventral spinal cord. FEBS J. 291(4):646-662.
Danesin C, Soula C. 2017. Moving the shh source over time: What impact on neural cell
diversification in the developing spinal cord? J Dev Biol. 5(2).
Dang NDP, Barcus AK, Conklin CL, Truong TQ, Vivian MD, Wang J, Thomas HR, Parant JM,
Yeo NC, Thyme SB. 2025. Disrupted development of sensory systems and the
cerebellum in a zebrafish ebf3a mutant. G3 (Bethesda). 15(7).
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR. 2013. Star: Ultrafast universal rna-seq aligner. Bioinformatics. 29(1):15-21.
Echelard Y, Epstein DJ, St-Jacques B, Shen L, Mohler J, McMahon JA, McMahon AP. 1993.
Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in
the regulation of cns polarity. Cell. 75(7):1417-1430.
England S, Batista MF, Mich JK, Chen JK, Lewis KE. 2011. Roles of hedgehog pathway
components and retinoic acid signalling in specifying zebrafish ventral spinal cord
neurons. Development. 138(23):5121-5134.
Esaki M, Hoshijima K, Nakamura N, Munakata K, Tanaka M, Ookata K, Asakawa K,
Kawakami K, Wang W, Weinberg ES et al. 2009. Mechanism of development of
ionocytes rich in vacuolar-type h(+)-atpase in the skin of zebrafish larvae. Dev Biol.
329(1):116-129.
Grevellec A, Graham A, Tucker AS. 2011. Shh signalling restricts the expression of gcm2 and
controls the position of the developing parathyroids. Dev Biol. 353(2):194-205.
Guner B, Karlstrom RO. 2007. Cloning of zebrafish nkx6.2 and a comprehensive analysis of the
conserved transcriptional response to hedgehog/gli signaling in the zebrafish neural tube.
Gene Expr Patterns. 7(5):596-605.
Hammerschmidt M, Bitgood MJ, McMahon AP. 1996. Protein kinase a is a common negative
regulator of hedgehog signaling in the vertebrate embryo. Genes Dev. 10(6):647-658.
Heemskerk J, DiNardo S. 1994. Drosophila hedgehog acts as a morphogen in cellular patterning.
Cell. 76(3):449-460.
Herzog W, Zeng X, Lele Z, Sonntag C, Ting JW, Chang CY, Hammerschmidt M. 2003.
Adenohypophysis formation in the zebrafish and its dependence on sonic hedgehog. Dev
Biol. 254(1):36-49.
Holzschuh J, Hauptmann G, Driever W. 2003. Genetic analysis of the roles of hh, fgf8, and
nodal signaling during catecholaminergic system development in the zebrafish brain. J
Neurosci. 23(13):5507-5519.
Hsiao CD, You MS, Guh YJ, Ma M, Jiang YJ, Hwang PP. 2007. A positive regulatory loop
between foxi3a and foxi3b is essential for specification and differentiation of zebrafish
epidermal ionocytes. PLoS One. 2(3):e302.
Huang P, Schier AF. 2009. Dampened hedgehog signaling but normal wnt signaling in zebrafish
without cilia. Development. 136(18):3089-3098.
Hwang PP, Chou MY. 2013. Zebrafish as an animal model to study ion homeostasis. Pflugers
Arch. 465(9):1233-1247.

Imokawa Y, Yoshizato K. 1997. Expression of sonic hedgehog gene in regenerating newt limb
blastemas recapitulates that in developing limb buds. Proc Natl Acad Sci U S A.
94(17):9159-9164.
Ingham PW, McMahon AP. 2001. Hedgehog signaling in animal development: Paradigms and
principles. Genes Dev. 15(23):3059-3087.
Janicke M, Carney TJ, Hammerschmidt M. 2007. Foxi3 transcription factors and notch signaling
control the formation of skin ionocytes from epidermal precursors of the zebrafish
embryo. Dev Biol. 307(2):258-271.
Karlstrom RO, Trowe T, Klostermann S, Baier H, Brand M, Crawford AD, Grunewald B,
Haffter P, Hoffmann H, Meyer SU et al. 1996. Zebrafish mutations affecting retinotectal
axon pathfinding. Development. 123:427-438.
Karlstrom RO, Tyurina OV, Kawakami A, Nishioka N, Talbot WS, Sasaki H, Schier AF. 2003.
Genetic analysis of zebrafish gli1 and gli2 reveals divergent requirements for gli genes in
vertebrate development. Development. 130(8):1549-1564.
Lawson ND, Li R, Shin M, Grosse A, Yukselen O, Stone OA, Kucukural A, Zhu L. 2020. An
improved zebrafish transcriptome annotation for sensitive and comprehensive detection
of cell type-specific genes. Elife. 9.
Litingtung Y, Chiang C. 2000. Specification of ventral neuron types is mediated by an
antagonistic interaction between shh and gli3. Nat Neurosci. 3(10):979-985.
Litingtung Y, Dahn RD, Li Y, Fallon JF, Chiang C. 2002. Shh and gli3 are dispensable for limb
skeleton formation but regulate digit number and identity. Nature. 418(6901):979-983.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
rna-seq data with deseq2. Genome Biol. 15(12):550.
Marcus RC, Easter SS, Jr. 1995. Expression of glial fibrillary acidic protein and its relation to
tract formation in embryonic zebrafish (danio rerio). J Comp Neurol. 359(3):365-381.
Marigo V, Davey RA, Zuo Y, Cunningham JM, Tabin CJ. 1996. Biochemical evidence that
patched is the hedgehog receptor. Nature. 384(6605):176-179.
Mich JK, Chen JK. 2011. Hedgehog and retinoic acid signaling cooperate to promote
motoneurogenesis in zebrafish. Development. 138(23):5113-5119.
Moore-Scott BA, Manley NR. 2005. Differential expression of sonic hedgehog along the
anterior-posterior axis regulates patterning of pharyngeal pouch endoderm and
pharyngeal endoderm-derived organs. Dev Biol. 278(2):323-335.
Mosimann C, Kaufman CK, Li P, Pugach EK, Tamplin OJ, Zon LI. 2011. Ubiquitous transgene
expression and cre-based recombination driven by the ubiquitin promoter in zebrafish.
Development. 138(1):169-177.
Moyer AJ, Barcus A, Capps MES, Chrabasz JA, Lalonde RL, Mosimann C, Thyme SB. 2025.
Genetic context of transgene insertion can influence neurodevelopment in zebrafish.
bioRxiv.
Ruiz i Altaba A. 1999. Gli proteins encode context-dependent positive and negative functions:
Implications for development and disease. Development. 126(14):3205-3216.
Shen MC, Ozacar AT, Osgood M, Boeras C, Pink J, Thomas J, Kohtz JD, Karlstrom R. 2013.
Heat-shock-mediated conditional regulation of hedgehog/gli signaling in zebrafish. Dev
Dyn. 242(5):539-549.
Shono T, Kurokawa D, Miyake T, Okabe M. 2011. Acquisition of glial cells missing 2 enhancers
contributes to a diversity of ionocytes in zebrafish. PLoS One. 6(8):e23746.
Stone DM, Hynes M, Armanini M, Swanson TA, Gu Q, Johnson RL, Scott MP, Pennica D,
Goddard A, Phillips H et al. 1996. The tumour-suppressor gene patched encodes a
candidate receptor for sonic hedgehog. Nature. 384(6605):129-134.

Sur A, Wang Y, Capar P, Margolin G, Prochaska MK, Farrell JA. 2023. Single-cell analysis of
shared signatures and transcriptional diversity during zebrafish development. Dev Cell.
58(24):3028-3047 e3012.
Taipale J, Chen JK, Cooper MK, Wang B, Mann RK, Milenkovic L, Scott MP, Beachy PA.
2000. Effects of oncogenic mutations in smoothened and patched can be reversed by
cyclopamine. Nature. 406(6799):1005-1009.
Wang B, Fallon JF, Beachy PA. 2000. Hedgehog-regulated processing of gli3 produces an
anterior/posterior repressor gradient in the developing vertebrate limb. Cell. 100(4):423-
434.
Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, Lumley T, Maechler M,
Magnusson A, Moeller S et al. 2025. Gplots: Various r programming tools for plotting
data. R package version 320.
Wechsler-Reya RJ, Scott MP. 1999. Control of neuronal precursor proliferation in the
cerebellum by sonic hedgehog. Neuron. 22(1):103-114.
White RJ, Mackay E, Wilson SW, Busch-Nentwich EM. 2022. Allele-specific gene expression
can underlie altered transcript abundance in zebrafish mutants. Elife. 11.
Wickham H. 2016. Ggplot2: Elegant graphics for data analysis. Cham, Switzerland: Springer
International Publishing.
Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L et al. 2021.
Clusterprofiler 4.0: A universal enrichment tool for interpreting omics data. Innovation
(Camb). 2(3):100141.
Zhang XM, Ramalho-Santos M, McMahon AP. 2001. Smoothened mutants reveal redundant
roles for shh and ihh signaling including regulation of l/r symmetry by the mouse node.
Cell. 106(2):781-792.
387
388

389 **Supplementary Materials**
390

391 **Gli3R-mediated inhibition of hedgehog signaling alters the embryonic**
392 **transcriptome in zebrafish**
393

394 Anna J. Moyer, Summer B. Thyme
395

396 **Supplementary File 1:** Annotated plasmid map of *pTol2-ubb:zGli3R*.
397

398 **Supplementary File 2:** R code to reproduce figures.
399

400 **Supplementary Table 1:** DESeq2 output of differentially expressed genes.
401

402 **Supplementary Table 2:** GSEA by chromosomal location.
403

404 **Supplementary Table 3:** GSEA using terms derived from Daniocell atlas single-cell data.
405