Within-Patient Evolution of *Pseudomonas aeruginosa* Populations During Antimicrobial Treatment

Giuseppe Fleres [1,2], Ellen G. Kline[1], Kevin M. Squires[1], Tyler Tate[1,4], Hannah M. Creager[1], Ryan K. Shields[1,3,4], Daria Van Tyne[1,2]

[1]Division of Infectious Diseases, University of Pittsburgh School of Medicine; [2]Center for Evolutionary Biology and Medicine, University of Pittsburgh School of Medicine; [3] Center for Innovative Antimicrobial Therapy, University of Pittsburgh School of Medicine; [4] Antibiotic Management Program, University of Pittsburgh Medical Center

Abstract: Multidrug-resistant (MDR) *Pseudomonas aeruginosa* infections pose a major challenge to effective treatment. Understanding genomic adaptations during antimicrobial therapy in patients infected with this pathogen is crucial for preventing therapeutic failure. Here we investigated the population diversity and evolution of *P. aeruginosa* collected longitudinally from six patients who evolved multidrug-resistant infections. Serial *P. aeruginosa* clinical isolates (n=63) and culture-enriched metagenomic population samples (n=39) were collected and subjected to whole-genome sequencing. The resulting data were used to characterize and compare the species composition, multi-locus sequence types (STs), and resistance-associated mutations present within each sample type. Single-isolate sequencing showed that each patient was infected with a single *P. aeruginosa* strain that accumulated mutations and became increasingly more resistant over time. Mutations in genes associated with beta-lactam resistance, including *ampC, ftsI*, and *mexR,* arose over time and corresponded with changes in antimicrobial susceptibility in single isolates. Species profiling of culture-enriched metagenomic populations revealed that all samples contained *P. aeruginosa*, but also additional Gram-negative pathogens. Metagenomic analysis of culture-enriched populations identified

resistance-associated mutations at low frequency, many of which were not identified in single

isolates from the same sample. In some cases, resistance-associated mutations initially

detected at low frequency rose to fixation after antimicrobial treatment. Overall, this study shows

that population-based metagenomic sequencing effectively captures within-patient genomic

diversity of *P. aeruginosa* during antimicrobial therapy, and could aid the detection and

interpretation of resistance-associated mutations in this pathogen.

Importance: *Pseudomonas aeruginosa* infections are notoriously difficult to treat and are

associated with high rates of morbidity and mortality. While the genetic basis of resistance in *P.*

aeruginosa is well documented *in vitro*, less is known about how resistance evolves within

patients during antibiotic therapy. Standard approaches based on analysis of clonal isolates

may miss within-patient diversity, potentially overlooking low-frequency mutations that contribute

to treatment failure. In this study, we integrated single-colony whole-genome sequencing with

culture-enriched metagenomic sequencing to monitor the evolution of *P. aeruginosa* populations

in patients receiving antibiotic therapy. This approach enabled the detection of emerging

resistance mutations, such as low-frequency variants in *ampC* and *ftsI*, before these variants

rose to fixation. It also revealed genetically resistant subpopulations missed by isolate

sequencing alone. Together, our findings highlight the value of population-based metagenomic

sequencing in capturing bacterial adaptation during infection, and underscore its potential to

improve resistance surveillance and guide personalized antimicrobial therapy.

Keywords: culture-enriched metagenomics; *Pseudomonas aeruginosa*; within-host evolution

Observation: *Pseudomonas aeruginosa* is recognized as a serious public health threat by the

Centers for Disease Control and Prevention and has been classified by the World Health

Organization as a critical priority pathogen, emphasizing the urgent need for new antimicrobial

therapies[1]. A variety of intrinsic resistance mechanisms, combined with a remarkable capacity to

acquire additional resistance determinants, make *P. aeruginosa* infections increasingly difficult

to treat[2]. Infections caused by multidrug-resistant (MDR) strains are particularly concerning and

often necessitate the use of newer β-lactam/β-lactamase inhibitor combinations or siderophore

cephalosporins to manage[3,4]. However, the efficacy of these agents is frequently undermined by

the emergence of diverse and often convergent resistance mechanisms. These include

upregulation of multidrug efflux systems (e.g., MexAB-OprM), inactivation or loss of porins

(OprD), mutations affecting chromosomal β-lactamase expression (AmpC), and point mutations

in essential antibiotic target genes like *ftsI* (encoding penicillin-binding protein 3), *gyrA,* and

gyrB[4,5]. Alterations in regulatory genes such as *lasR* further contribute to phenotypic

heterogeneity by modulating quorum sensing pathways, which can affect antibiotic tolerance

and persistence[2]. These adaptive mutations frequently arise under selective therapeutic

pressures, facilitating rapid within-host evolution and complicating effective treatment[5].

Although resistance mechanisms in *P. aeruginosa* have been well characterized *in vitro* and

among serial clinical isolates collected from infected patients[6,7], much less is known about how

infecting *P. aeruginosa* populations evolve within patients during antibiotic therapy[8]. Treatment

decisions are typically guided by antimicrobial susceptibility testing and whole-genome

sequencing analysis of a single bacterial clone, assuming that it reflects the entire infecting

population. However, this approach offers limited insight into within-host diversity and often

misses low-frequency variants that may influence treatment outcomes[9,10]. Culture-enriched

metagenomic sequencing addresses these limitations by pooling all colonies from a selective

culture plate and performing shotgun metagenomic sequencing, enabling a more

comprehensive analysis of population-level diversity and evolution during infection[9,11].

Here we investigated the within-host evolution of *P. aeruginosa* populations in six patients who

developed MDR infections during treatment and compared results obtained from clonal and

culture-enriched metagenomic sequencing. We hypothesized that reliance on single-colony

testing would underestimate the genotypic and phenotypic diversity of *P. aeruginosa* within the

infected host, particularly under antimicrobial selection. A total of 63 clinical isolates and 39

culture-enriched metagenomic populations were collected from six patients, sequenced, and

analyzed (Figure 1) (Supplemental File 1) (Supplemental Methods). In all patients, susceptibility

profiles of *P. aeruginosa* isolates progressively shifted toward increasing resistance over the

course of treatment. This aligns with prior studies documenting the rapid evolution of resistance

in *P. aeruginosa* under antibiotic stress, especially during prolonged therapy in chronic

infections[12,13]. In two patients (Patients 1 and 6), isolates collected at the final time point

exhibited a more susceptible phenotype compared to earlier isolates, suggesting that resistant

bacteria may have been outcompeted or replaced, potentially due to changes in treatment, host

immunity, or fitness costs associated with resistance[14].

Species profiling of culture-enriched metagenomic samples confirmed the presence of *P.*

aeruginosa in all samples (Figure 1). Metagenomic samples from three patients (Patients 1–3)

contained only *P. aeruginosa*, while the other three patients (Patients 4–6) exhibited

polymicrobial infections including *P. aeruginosa* and other organisms (Supplemental Figure 1).

In each case, the same species were identified as part of standard-of-care testing in the clinical

microbiology laboratory, suggesting that this approach accurately captured polymicrobial

infections.

Single-colony sequencing revealed a *P. aeruginosa* sequence type (ST) that was unique to

each patient, and all colonies sequenced from the same patient belonged to the same ST

(Supplemental Figure 2A). Analysis of within-patient single nucleotide polymorphisms (SNP)

among *P. aeruginosa* isolates collected from the same patient revealed low genetic diversity in

five of the six patients, with pairwise SNP differences ranging from 0 to 23. This narrow range

suggested that each patient was infected with a single strain that underwent clonal expansion

during infection. In contrast, one patient (Patient 1) displayed markedly higher within-host

diversity, with SNP distances reaching up to 411 (Supplemental Table 2). This elevated level of

genetic variation could be attributed to the presence of a hypermutator clone, which accelerates

mutation rates and promotes rapid genetic diversification, particularly under selective pressures

such as antibiotic treatment or the host immune response[15]. Supporting this interpretation,

multiple mutations were identified in DNA mismatch repair genes from this patient's isolates,

including W181R and S324P in *mutS* and L35P in *mutL*, indicating a defective mismatch repair

system.

We next explored whether culture-enriched metagenomic sequencing could enhance the

detection of resistance-associated mutations that might be missed by single-colony sequencing.

In our analysis of 11 resistance-associated genes across 6 patients, culture-enriched

metagenomic sequencing provided additional insight in 19/66 (29%) cases. In 84% of these

cases (n=16/19), this was due to the detection of resistance-associated mutations that were not

identified by single-colony sequencing (Supplemental Table 3). Comparison of single-colony

and culture-enriched metagenomic sequencing revealed distinct interpretations depending on

the method used (Figure 2). In Patient 3, single-colony sequencing identified a low-frequency

mutation in *ampC* (P243S) but failed to detect a co-occurring mutation (E247K) that was present

at over 80% frequency in the population (Figure 2A). This highlights a key limitation of the

single-colony approach, which might randomly capture minority variants while missing the

dominant genotype, ultimately leading to an incomplete or misleading resistance profile. In both

Patients 3 and 4, mutations in *ftsI* were identified at low frequency in population samples

collected at earlier time points, and the same variants were later detected in single colonies as

well as in the populations at high frequency (Figure 2A and 2B). This illustrates how population

sequencing can detect emerging mutations before they become dominant. Finally, in Patient 6

we identified mutations at allele frequencies less than 50% that were exclusively detected in

population samples and were never observed in single-colony genomes (Figure 2C). Together

these findings emphasize the higher sensitivity of culture-enriched metagenomic sequencing for

capturing within-host diversity.

Here we focused on a targeted set of genes known to be associated with antibiotic resistance.

While this approach allowed for in-depth analysis of clinically relevant resistance mechanisms, it

likely underrepresents the broader genomic heterogeneity within infecting *P. aeruginosa*

populations. Other adaptive changes (e.g., mutations related to antibiotic tolerance, biofilm

formation, metabolic remodeling, etc.) may also contribute to treatment failure but were not

systematically assessed here. Future studies incorporating unbiased, genome-wide analyses

could provide a more comprehensive view of within-host evolution during infection and

treatment.

In summary, this study demonstrates that culture-enriched metagenomic sequencing provides

enhanced resolution of within-host evolution in *P. aeruginosa* during antimicrobial treatment.

Compared with prior studies, here we used a combined approach of single-colony and culture-

enriched metagenomic sequencing to achieve higher-resolution tracking of within-host evolution

during antibiotic treatment in infected patients. We found that population-level sequencing

consistently identified low-frequency variants that were missed by single-colony sequencing,

revealing a more comprehensive picture of the adaptive landscape in these patients. As WGS

becomes more commonly used in clinical practice, culture-enriched metagenomic sequencing

could provide greater sensitivity for studies of antibiotic resistance and evolutionary dynamics,

which could contribute to improved management of patients with difficult-to-treat infections.

Supplemental Material

Figure S1

Figure S2

Supplemental Materials and Methods

Supplemental Table 1

Supplemental Table 2

Supplemental Table 3

Data Availability

The dataset generated during the current study are available in the Sequence Read Archive

(SRA), BioProject PRJNA1330241.

Acknowledgements

We gratefully acknowledge members of the UPMC Clinical Microbiology Laboratory and

members of the Van Tyne laboratory for assistance throughout this study. This work was

supported by the Cystic Fibrosis Foundation (grant SCON-00006650 to DVT), and by the

Department of Medicine at the University of Pittsburgh (DVT). The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. WHO Bacterial Priority Pathogens List 2024: Bacterial Pathogens of Public Health

Importance, to Guide Research, Development, and Strategies to Prevent and Control

Antimicrobial Resistance. (World Health Organization, Geneva, 2024).

2. Letizia, M., Diggle, S. P. & Whiteley, M. Pseudomonas aeruginosa: ecology, evolution,

pathogenesis and antimicrobial susceptibility. Nat. Rev. Microbiol. (2025)

doi:10.1038/s41579-025-01193-8.

3. Shields, R. K. et al. Effectiveness of ceftazidime-avibactam versus ceftolozane-tazobactam

for multidrug-resistant Pseudomonas aeruginosa infections in the USA (CACTUS): a

multicentre, retrospective, observational study. Lancet Infect. Dis. **25**, 574–584 (2025).

4. Pang, Z., Raudonis, R., Glick, B. R., Lin, T.-J. & Cheng, Z. Antibiotic resistance in

Pseudomonas aeruginosa: mechanisms and alternative therapeutic strategies. Biotechnol.

Adv. **37**, 177–192 (2019).

5. Weimann, A. et al. Evolution and host-specific adaptation of Pseudomonas aeruginosa.

Science **385**, eadi0908 (2024).

6. Tam, V. H. et al. Prevalence, Resistance Mechanisms, and Susceptibility of Multidrug-

Resistant Bloodstream Isolates of Pseudomonas aeruginosa. Antimicrob. Agents

Chemother. **54**, 1160–1164 (2010).

7. Bitar, I. et al. Genomic Characterization of Mutli-Drug Resistant Pseudomonas aeruginosa

Clinical Isolates: Evaluation and Determination of Ceftolozane/Tazobactam Activity and

Resistance Mechanisms. Front. Cell. Infect. Microbiol. **12**, (2022).

8. Spottiswoode, N. et al. In host evolution of beta lactam resistance during active treatment

for Pseudomonas aeruginosa bacteremia. Front. Cell. Infect. Microbiol. **13**, (2023).

9. Raghuram, V. et al. Comparison of genomic diversity between single and pooled

Staphylococcus aureus colonies isolated from human colonization cultures. Microb.

Genomics **9**, 001111 (2023).

10. Deurenberg, R. H. et al. Application of next generation sequencing in clinical microbiology

and infection prevention. J. Biotechnol. **243**, 16–24 (2017).

11. Chiu, C. Y. & Miller, S. A. Clinical metagenomics. Nat. Rev. Genet. **20**, 341–355 (2019).

12. Cabot, G. et al. Evolution of Pseudomonas aeruginosa Antimicrobial Resistance and Fitness

under Low and High Mutation Rates. Antimicrob. Agents Chemother. **60**, 1767–1778 (2016).

13. Winstanley, C., O'Brien, S. & Brockhurst, M. A. Pseudomonas aeruginosa Evolutionary

Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol.

24, 327–337 (2016).

14. Abdelraouf, K., Kabbara, S., Ledesma, K. R., Poole, K. & Tam, V. H. Effect of multidrug

resistance-conferring mutations on the fitness and virulence of Pseudomonas aeruginosa. J.

Antimicrob. Chemother. **66**, 1311–1317 (2011).

15. Liu, H. et al. Time Series Genomics of Pseudomonas aeruginosa Reveals the Emergence of

a Hypermutator Phenotype and Within-Host Evolution in Clinical Inpatients. Microbiol.

Spectr. **10**, e0005722 (2022).

Figure Legends

Figure 1. Timeline of *P. aeruginosa* infection and treatment for each patient. The panel

shows longitudinal sampling of *P. aeruginosa* single colony isolates and culture-enriched

metagenomic populations during recurrent infections for each patient. The upper section of each

panel depicts single colony isolates, while the lower section shows corresponding population

samples. Colored squares indicate the antimicrobial susceptibility profile of each isolate, and

grey bars show the proportion of each population composed of *P. aeruginosa* or other bacterial

species. β-lactam antibiotics tested included aztreonam (ATM), cefepime (FEP), ceftazidime

(CAZ), meropenem (MEM), piperacillin/tazobactam (TZP), ceftolozane/tazobactam (C/T), and

ceftazidime/avibactam (CZA). Ovals in each timeline are colored based on antimicrobial

treatment administered to each patient, including FEP, MEM, TZP, C/T, ciprofloxacin (CIP),

cefiderocol (FDC), levofloxacin (LEV), imipenem/relebactam (IPR), and CZA. Sample identifiers,

source (respiratory tract or blood), and collection timepoints (in days) are indicated above and

below the timeline.

Figure 2. Comparison of resistance-associated mutations detected in culture-enriched

metagenomic populations versus single colonies in three patients with *P. aeruginosa*

infections. Each panel represents a subset of the isolate and population data collected from

each patient, and displays mutations identified in population sequencing (left) and in single

colony isolate whole-genome sequences (right) at corresponding sampling timepoints.

Nonsynonymous mutations, nonsense mutations, and structural variants are shown in beta-

lactam resistance-associated genes *ampC*, *oprD*, and *ftsI*. Colored bars represent the allele

frequency (%) of each mutation in metagenomic populations. Sample ID and collection day are

indicated next to each genotype.