Within-Patient Evolution of *Pseudomonas aeruginosa* Populations During Antimicrobial Treatment

Giuseppe Fleres[1,2], Ellen G. Kline[1], Kevin M. Squires[1], Tyler Tate[1,3], Hannah M. Creager[1], Ryan K. Shields[1, 3, 4], Daria Van Tyne[1,2,4]

[1]Division of Infectious Diseases, University of Pittsburgh School of Medicine; [2]Center for Evolutionary Biology and Medicine, University of Pittsburgh School of Medicine; [3]Antibiotic Management Program, University of Pittsburgh Medical Center; [4]Center for Innovative Antimicrobial Therapy, University of Pittsburgh School of Medicine

Abstract: Multidrug-resistant (MDR) *Pseudomonas aeruginosa* infections pose a major challenge to effective treatment. Understanding genomic adaptations during antimicrobial therapy in patients infected with this pathogen is crucial for preventing therapeutic failure. Here we investigated the population diversity and evolution of *P. aeruginosa* collected longitudinally from six patients who evolved MDR infections. Serial *P. aeruginosa* clinical isolates (n=63) and culture-enriched metagenomic populations (n=39) were collected and subjected to whole-genome sequencing. The resulting data were used to characterize and compare the species composition, multi-locus sequence types (STs), and resistance-associated mutations present within each sample type. Single-isolate sequencing showed that each patient was infected with a single *P. aeruginosa* strain that accumulated mutations and became increasingly antibiotic-resistant over time. Mutations in genes associated with β-lactam resistance, including *ampC, ftsI*, and *oprD,* arose over time and corresponded with changes in antimicrobial susceptibility in single isolates. Species profiling of culture-enriched metagenomic populations revealed that all samples contained *P. aeruginosa*, but several also contained additional Gram-negative pathogens. Metagenomic analysis of culture-enriched populations identified resistance-

associated mutations at variable frequencies, many of which were not identified in single

isolates from the same sample. In some cases, resistance-associated mutations initially

detected at low frequency rose to fixation after antimicrobial treatment. Overall, this study shows

that population-based metagenomic sequencing effectively captures within-patient genomic

diversity of *P. aeruginosa* during antimicrobial therapy, and could aid in the detection and

interpretation of resistance-associated mutations in this pathogen.

Importance: *Pseudomonas aeruginosa* infections are notoriously difficult to treat and are

associated with high rates of morbidity and mortality. While the genetic basis of resistance in *P.*

aeruginosa is well documented *in vitro*, less is known about how resistance evolves within

patients during antibiotic therapy. Standard approaches based on analysis of clonal isolates

may miss within-patient diversity, potentially overlooking low-frequency mutations that contribute

to treatment failure. In this study, we integrated single-colony whole-genome sequencing with

culture-enriched metagenomic sequencing to monitor the evolution of *P. aeruginosa* populations

in patients receiving antibiotic therapy. This approach enabled the detection of emerging

resistance mutations, such as low-frequency variants in *ampC* and *ftsI*, before these variants

rose to fixation. It also revealed genetically resistant subpopulations missed by isolate

sequencing alone. Together, our findings highlight the value of population-based metagenomic

sequencing in capturing bacterial adaptation during infection, and underscore its potential to

improve resistance surveillance and guide personalized antimicrobial therapy.

Keywords: culture-enriched metagenomics; *Pseudomonas aeruginosa*; within-host evolution;

antibiotic resistance

Observation: *Pseudomonas aeruginosa* is recognized as a serious public health threat by the

Centers for Disease Control and Prevention and has been classified by the World Health

Organization as a critical priority pathogen, emphasizing the urgent need for new antimicrobial

therapies[1]. A variety of intrinsic resistance mechanisms, combined with a remarkable capacity to

acquire additional resistance determinants, make *P. aeruginosa* infections increasingly difficult

to treat[2]. Infections caused by multidrug-resistant (MDR) strains are particularly concerning and

often necessitate the use of newer β-lactam/β-lactamase inhibitor combinations or siderophore

cephalosporins to manage[3,4]. However, the efficacy of these agents is frequently undermined by

the emergence of diverse and often convergent resistance mechanisms. These include

upregulation of multidrug efflux systems (e.g., MexAB-OprM), inactivation or loss of porins

(OprD), mutations affecting chromosomal β-lactamase expression (AmpC), and point mutations

in essential antibiotic target genes like *ftsI* (encoding penicillin-binding protein 3), *gyrA*, and

gyrB[4,5]. Alterations in regulatory genes such as *lasR* further contribute to phenotypic

heterogeneity by modulating quorum sensing pathways, which can affect antibiotic tolerance

and persistence[2]. These adaptive mutations frequently arise under selective therapeutic

pressures, facilitating rapid within-host evolution and complicating effective treatment[5].

Although resistance mechanisms in *P. aeruginosa* have been well characterized *in vitro*

and among serial clinical isolates collected from infected patients[6,7], much less is known about

how infecting *P. aeruginosa* populations evolve within patients during antibiotic therapy[8].

Treatment decisions are typically guided by antimicrobial susceptibility testing and whole-

genome sequencing analysis of individual bacterial clones, assuming that these reflect the

entire infecting population. However, this approach offers limited insight into within-host diversity

and often misses low-frequency variants that may influence treatment outcomes[9,10]. Culture-

enriched metagenomic sequencing addresses these limitations by pooling all colonies from a

selective culture plate and performing shotgun metagenomic sequencing, enabling a more

comprehensive analysis of population-level diversity and evolution during infection[9,11].

Here we investigated the within-host evolution of *P. aeruginosa* populations in six

patients who developed MDR infections during treatment and compared between results

obtained from clonal and culture-enriched metagenomic sequencing from the same clinical

specimens. We hypothesized that reliance on single-colony testing would underestimate the

genotypic and phenotypic diversity of *P. aeruginosa* within the infected host, particularly under

antimicrobial selection. A total of 63 clinical isolates and 39 culture-enriched metagenomic

populations were collected from six patients, sequenced, and analyzed (**Figure 1**)

(**Supplemental File 1**) (**Supplemental Methods**). In all patients, susceptibility profiles of *P.*

aeruginosa isolates progressively shifted toward increasing resistance over the course of

antibiotic treatment. This aligns with prior studies documenting the rapid evolution of resistance

in *P. aeruginosa* under antibiotic stress, especially during prolonged therapy in chronic

infections[12,13]. In two patients (Patients 1 and 6), isolates collected at the final time point

exhibited a more antibiotic-susceptible profile compared to earlier isolates, suggesting that

resistant bacteria may have been outcompeted or replaced, potentially due to changes in

treatment, host immunity, or fitness costs associated with resistance[14].

Species profiling of culture-enriched metagenomic samples confirmed the presence of *P.*

aeruginosa in all samples (**Figure 1**). Metagenomic samples from three patients (Patients 1–3)

contained only *P. aeruginosa*, while the other three patients (Patients 4–6) exhibited

polymicrobial infections that included *P. aeruginosa* and other organisms (**Supplemental**

Figure 1). In each case, the same species were identified as part of standard-of-care testing in

the clinical microbiology laboratory, suggesting that this approach accurately captured

polymicrobial infections.

Single-colony sequencing revealed a single *P. aeruginosa* sequence type (ST) that was

unique to each patient, and all colonies sequenced from the same patient belonged to the same

ST (**Supplemental Figure 2**). Analysis of within-patient single nucleotide polymorphisms

(SNPs) among *P. aeruginosa* isolates collected from the same patient revealed low genetic

diversity in five of the six patients, with pairwise SNP differences ranging from 0 to 23. This

narrow range suggested that each patient was infected with a single strain that underwent

clonal expansion during infection. In contrast, one patient (Patient 1) displayed markedly higher

within-host diversity, with SNP distances reaching up to 411 (**Supplemental Table 2**). This

elevated level of genetic variation could be attributed to the presence of hypermutators, which

have increased mutation rates and rapidly diversify, particularly under selective pressures such

as antibiotic treatment or the host immune response[15]. Supporting this interpretation, multiple

mutations were identified in DNA mismatch repair genes from this patient's isolates, including

W181R and S324P in *mutS* and L35P in *mutL,* indicating a defective mismatch repair system.

We next explored whether culture-enriched metagenomic sequencing could enhance the

detection of resistance-associated mutations that might be missed by single-colony sequencing.

In our analysis of 11 resistance-associated genes across 6 patients, culture-enriched

metagenomic sequencing provided additional insight in 19/66 (29%) cases. In 84% of these

cases (n=16/19), this was due to the detection of resistance-associated mutations that were not

identified by single-colony sequencing (**Supplemental Table 3**). Comparison of single-colony

and culture-enriched metagenomic sequencing revealed distinct interpretations depending on

the method used. In Patient 3, single-colony sequencing identified a mutation in *ampC* (P243S)

but failed to detect a co-occurring mutation (E247K) that was present at over 80% frequency in

the population (**Figure 2A**). This highlights a key limitation of the single-colony approach, which

might randomly capture minority variants while missing the dominant genotype, ultimately

leading to an incomplete or misleading resistance profile. In both Patients 3 and 4, mutations in

ftsI were identified at low frequency in population samples collected at earlier time points, and

the same variants were later detected in single colonies as well as in populations at high

frequency (**Figure 2A and 2B**). This illustrates how population sequencing can detect emerging

mutations before they become dominant. Finally, in Patient 6 we identified mutations at allele

frequencies less than 50% that were exclusively detected in population samples and were never

observed in single-colony genomes (**Figure 2C**). Together these findings emphasize the higher

sensitivity of culture-enriched metagenomic sequencing for capturing within-host diversity.

Here we focused on a targeted set of genes known to be associated with antibiotic

resistance in *P. aeruginosa*. While this approach allowed for in-depth analysis of clinically

relevant resistance mechanisms, it likely underrepresents the broader genomic heterogeneity

within infecting *P. aeruginosa* populations. Other adaptive changes (such as mutations related

to antibiotic tolerance, biofilm formation, metabolic remodeling, etc.) may also contribute to

treatment failure, but were not systematically assessed here. Future studies incorporating

unbiased, genome-wide analyses could provide a more comprehensive view of within-host

evolution during bacterial infection and antimicrobial treatment.

In summary, this study demonstrates that culture-enriched metagenomic sequencing

provides enhanced resolution of within-host evolution of *P. aeruginosa* during antimicrobial

treatment in human patients. We found that population-level sequencing consistently identified

low-frequency variants that were missed by single-colony sequencing, revealing a more

comprehensive picture of the adaptive landscape in these patients. As WGS becomes more

commonly used in clinical practice, culture-enriched metagenomic sequencing could provide

greater sensitivity for studies of antibiotic resistance and evolutionary dynamics, which could

contribute to improved management of patients with difficult-to-treat infections.

Supplemental Materials

Supplemental Figure 1

Supplemental Figure 2

Supplemental Methods

Supplemental File 1

Supplemental Table 1

Supplemental Table 2

Supplemental Table 3

Data Availability

Acknowledgements

We gratefully acknowledge members of the UPMC Clinical Microbiology Laboratory and

members of the Van Tyne laboratory for assistance throughout this study. This work was

supported by the Department of Medicine at the University of Pittsburgh. The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

References

1.   WHO Bacterial Priority Pathogens List 2024: Bacterial Pathogens of Public Health

Importance, to Guide Research, Development, and Strategies to Prevent and Control

Antimicrobial Resistance. (World Health Organization, Geneva, 2024).

2.   Letizia, M., Diggle, S. P. & Whiteley, M. Pseudomonas aeruginosa: ecology, evolution,

pathogenesis and antimicrobial susceptibility. Nat. Rev. Microbiol. (2025)

doi:10.1038/s41579-025-01193-8.

3.   Shields, R. K. et al. Effectiveness of ceftazidime-avibactam versus ceftolozane-tazobactam

for multidrug-resistant Pseudomonas aeruginosa infections in the USA (CACTUS): a

multicentre, retrospective, observational study. Lancet Infect. Dis. **25**, 574–584 (2025).

4.   Pang, Z., Raudonis, R., Glick, B. R., Lin, T.-J. & Cheng, Z. Antibiotic resistance in

Pseudomonas aeruginosa: mechanisms and alternative therapeutic strategies. Biotechnol.

Adv. **37**, 177–192 (2019).

5.   Weimann, A. et al. Evolution and host-specific adaptation of Pseudomonas aeruginosa.

Science **385**, eadi0908 (2024).

6.   Tam, V. H. et al. Prevalence, Resistance Mechanisms, and Susceptibility of Multidrug-

Resistant Bloodstream Isolates of Pseudomonas aeruginosa. Antimicrob. Agents

Chemother. **54**, 1160–1164 (2010).

7.   Bitar, I. et al. Genomic Characterization of Mutli-Drug Resistant Pseudomonas aeruginosa

Clinical Isolates: Evaluation and Determination of Ceftolozane/Tazobactam Activity and

Resistance Mechanisms. Front. Cell. Infect. Microbiol. **12**, (2022).

8.   Spottiswoode, N. et al. In host evolution of beta lactam resistance during active treatment

for Pseudomonas aeruginosa bacteremia. Front. Cell. Infect. Microbiol. **13**, (2023).

9.   Raghuram, V. et al. Comparison of genomic diversity between single and pooled

Staphylococcus aureus colonies isolated from human colonization cultures. Microb.

Genomics **9**, 001111 (2023).

10. Deurenberg, R. H. et al. Application of next generation sequencing in clinical microbiology

and infection prevention. J. Biotechnol. **243**, 16–24 (2017).

11. Chiu, C. Y. & Miller, S. A. Clinical metagenomics. Nat. Rev. Genet. **20**, 341–355 (2019).

12. Cabot, G. et al. Evolution of Pseudomonas aeruginosa Antimicrobial Resistance and Fitness

under Low and High Mutation Rates. Antimicrob. Agents Chemother. **60**, 1767–1778 (2016).

13. Winstanley, C., O'Brien, S. & Brockhurst, M. A. Pseudomonas aeruginosa Evolutionary

Adaptation and Diversification in Cystic Fibrosis Chronic Lung Infections. Trends Microbiol.

24, 327–337 (2016).

14. Abdelraouf, K., Kabbara, S., Ledesma, K. R., Poole, K. & Tam, V. H. Effect of multidrug

resistance-conferring mutations on the fitness and virulence of Pseudomonas aeruginosa. J.

Antimicrob. Chemother. **66**, 1311–1317 (2011).

15. Liu, H. et al. Time Series Genomics of Pseudomonas aeruginosa Reveals the Emergence of

a Hypermutator Phenotype and Within-Host Evolution in Clinical Inpatients. Microbiol.

Spectr. **10**, e0005722 (2022).

Figure 1. Timelines of *P. aeruginosa* infection and treatment in six patients with MDR infections. Each panel shows longitudinal sampling of *P. aeruginosa* single colony isolates (above timeline) and culture-enriched metagenomic populations (below timeline) during recurrent infection for each patient. Colored squares indicate the antimicrobial susceptibility profile of each isolate, and grey bars show the proportion of each population composed of *P. aeruginosa* or other bacterial species. β-lactam antibiotics tested included aztreonam (ATM), cefepime (FEP), ceftazidime (CAZ), meropenem (MEM), piperacillin/tazobactam (TZP), ceftolozane/tazobactam (C/T), and ceftazidime/avibactam (CZA). Ovals in each timeline are colored based on antimicrobial treatment administered to each patient, including FEP, MEM, TZP, C/T, ciprofloxacin (CIP), cefiderocol (FDC), levofloxacin (LEV), imipenem/relebactam (IPR), and CZA. Sample identifiers, specimen source (respiratory tract or blood), and collection timepoints (in days) are indicated above and below each timeline.

Figure 2. Comparison of resistance-associated mutations detected in culture-enriched metagenomic populations versus single colonies. Each panel represents a subset of the isolate and population data collected from each patient, and displays mutations identified in populations (left) and in single colony isolates (right) collected from the same specimens at different sampling timepoints. Nonsynonymous mutations, nonsense mutations, and structural variants are shown in β-lactam resistance-associated genes *ampC*, *oprD*, and *ftsI*. Colored bars represent the allele frequency (%) of each mutation in culture-enriched metagenomic populations. Sample ID and collection day are indicated for each comparison.

Supplemental Figure 1. Relative abundance of different bacterial species in culture-enriched metagenomic population samples collected from each patient.

227   **Supplemental Figure 2.** (A) Core genome phylogenetic tree of 63 single-colony *P. aeruginosa*

228   isolates, colored by patient. Tree branches are marked with the multilocus sequence type (ST)

229   that all isolates from the same patient belong to. Isolate source is noted for each isolate. (B-G)

230   Time-dated phylogenetic trees of single-colony *P. aeruginosa* isolates collected from each

231   patient.