Geomagnetic and visual cues guide seasonal migratory orientation in the
nocturnal fall armyworm, the world's most invasive insect

Yi-Bo Ma[1, †], Gui-Jun Wan[1, †], Yi Ji[1], Hui Chen[1,2], Bo-Ya Gao[1], Dai-Hong Yu[3], Eric J.
Warrant[2], Yan Wu[4], Jason W. Chapman[1,5], Gao Hu[1,4 *]

1 State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant
Protection, Nanjing Agricultural University, Nanjing 210095, China
2 Lund Vision Group, Department of Biology, Lund University, Sölvegatan 35, 22362
Lund, Sweden.
3 Plant Protection Station of Yuanjiang County, Yuanjiang 653399, China
4 Key Laboratory of Surveillance and Management of Invasive Alien Species,
Guizhou Education Department, Department of Biology and Engineering of
Environment, Guiyang University, Guiyang 550005, China.
5 Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall
TR10 9FE, United Kingdom.

* Correspondence to: Gao HU, 1 Weigang Road, Nanjing 210095, China. E-mail:
hugao@njau.edu.cn
† These authors have contributed equally to this work

Abstract

The mechanisms guiding nocturnal insect migration remain poorly understood. Although many species are thought to use the geomagnetic field, the sensory basis of magnetic orientation in insects has yet to be clarified. We developed an indoor experimental system to investigate the integration of geomagnetic and visual cues in the seasonal orientation of a globally distributed pest moth, the fall armyworm (*Spodoptera frugiperda*), a highly invasive species which in the past decade has colonized almost all potentially habitable regions of the globe. Our results demonstrate that fall armyworms require both geomagnetic and visual cues for accurate migratory orientation, with visual cues being indispensable for magnetic orientation. When visual and geomagnetic cues are placed in conflict moths become disoriented, although not immediately, indicating that sensory recognition of the conflict requires time to process. We also show that the absence of visual cues leads to a significant loss of flight stability, which likely explains the disruption in orientation. Our findings highlight that visual cues are critical for stable magnetic orientation in the fall armyworm, offering a basis for future investigations of visual-magnetic integration in noctuid migrants.

Introduction

Numerous species of larger nocturnal moths, particularly in the family Noctuidae (hereafter 'noctuid moths'), undertake long-distance multigenerational migrations in the Northern Hemisphere. Every spring across North America and Eurasia, billions of noctuid moths move hundreds of kilometers northwards to summer breeding grounds in the temperate zone, and in the subsequent fall their progeny return to lower latitude wintering areas (Holland et al., 2006; Satterfield et al., 2020; Hu et al., 2016, 2025; Huang et al., 2024). These migrations take place almost exclusively high above the ground, where fast winds facilitate rapid windborne transport leading to population redistribution over long distances (Chapman et al., 2010, 2015; Hu et al., 2016; Huang et al., 2024). Some of the most abundant species involved in these migrations are the world's most destructive agricultural pests, and thus they are of huge importance for food security and economic prosperity (Bauer & Hoye, 2014; Satterfield et al., 2020; Guo et al., 2020; Hu et al., 2025). It is thus of paramount importance to understand all aspects of the migratory patterns of noctuid moths.

Tracking the seasonal progression of resources requires noctuid moths to move in

the appropriate direction (northwards in spring and southwards in fall), a process
entailing three linked steps. Firstly, they determine the seasonally appropriate
direction (north or south) in which to travel. Secondly, high-flying noctuid moths select
transporting winds that are broadly aligned with this direction. Thirdly, they adopt self-
powered flight headings which are more-or-less aligned with the wind but, when
required, offset to some degree to correct for crosswind drift (Chapman et al., 2015;
Hu et al., 2016). Each step requires use of one or more compass senses based on
globally stable cues to determine the required direction and behavioral responses
(Mouritsen, 2014).

However, migration at night poses a considerable navigational challenge for
noctuid moths and other night-flying insects (Warrant et al., 2011; Mouritsen, 2018;
Foster et al., 2018; Gao et al., 2024; Grob et al., 2025). This is because they cannot
rely on the sun for compass information in the way that diurnally migrating butterflies
and hoverflies do (Mouritsen et al., 2002; Srygley et al., 2008; Gao et al., 2020;
Massy et al., 2021; Pakhomov et al., 2025). Despite this challenge, radar, tagging,
and tethered-flight studies demonstrate that many larger nocturnal moths are highly
sophisticated navigators, capable of selecting and maintaining appropriate movement
directions with a high degree of accuracy, even when flying high above the ground
(Chapman et al., 2010; Dreyer et al., 2018ab; Menz et al., 2022; Chen et al., 2023;
Dreyer et al., 2025). Their ability to achieve seasonally appropriate migration
directions typically surpasses that of diurnal windborne insect migrants (Chapman et
al., 2010; Hu et al., 2016; Werber et al., 2025), and even matches the capability of
nocturnal songbird migrants (Alerstam et al., 2011, Chapman et al., 2016). Clearly,
migratory noctuid moths must possess one or more accurate compass senses, but in
all species but one, the source of the compass information has yet to be elucidated
(Warrant et al., 2011; Foster et al., 2018; Mouritsen, 2018; Freas et al., 2023; Grob et
al., 2025).

The one exception to this lack of knowledge of the sensory basis of navigation in
noctuid moths is the Bogong moth (*Agrotis infusa*) of Australia, which uses both a
magnetic compass integrated with visual cues (Dreyer et al., 2018a) and a stellar
compass (Dreyer et al., 2025) to migrate in seasonally appropriate directions. The
Bogong moth is unique, however, among long-range migratory noctuids because a
single generation makes bidirectional movements to and from a highly restricted
geographic location in southeast Australia (Warrant et al., 2016), rather akin to the
migration of eastern North American monarch butterflies to and from a restricted area
of central Mexico (Reppert et al., 2016). This is distinct from most other noctuid moth

migrants, whose multigenerational migrations involve back-and-forth movements
between broad latitudinal zones (Drake & Reynolds, 2012; Chapman et al., 2015;
Gao et al., 2020; Tong et al., 2022; Hu et al., 2025) rather than to precise locations.
Thus other noctuid moths presumably do not require the same navigational precision,
and may therefore be expected to have simpler sensory capabilities than the Bogong
moth. We examine this question using the fall armyworm (*Spodoptera frugiperda*),
one of the world's most serious crop pests, as a model for understanding general
noctuid moth migration and navigation capabilities.

The fall armyworm is a migratory crop pest native to the Americas, where it breeds
year-round in tropical regions and annually migrates into temperate regions of North
America (Nagoshi & Meagher, 2008; Westbrook et al., 2019). Over the past 10 years,
it has invaded and rapidly spread across Africa, Asia, and Australasia (Goergen et al.,
2016; Kenis et al., 2023; Tay et al., 2023), assisted by its high migratory capacity
(Chen et al., 2022, 2025; Gao et al., 2025), where it causes huge yield losses.
Invasive fall armyworm populations now breed year-round in tropical regions of
Southeast Asia and South China; each spring they migrate northwards as far as
Northeast China, and the progeny resulting from summer breeding return southwards
in the fall (Li et al., 2020; Sun et al., 2021; Wu et al., 2021; Wu et al., 2022). We
recently found that fall armyworms from the year-round range in Southwest China
(Yunnan) exhibit seasonally appropriate migratory headings when flown outdoors in
virtual flight simulators, heading northward in the spring and southward in the fall,
and this seasonal reversal is controlled by photoperiod (Chen et al., 2023). However,
the compass mechanism that fall armyworms use to select seasonally appropriate
headings remains unknown. The Earth's geomagnetic field is a ubiquitous and
reliable source of compass information (Mouritsen, 2018), and thus might be
expected to be the primary compass cue used by night-flying migratory moths.
However, despite intriguing indications of magnetoreception in the oriental armyworm,
Mythimna separata (Xu et al., 2017), so far only the highly specialised Bogong moth
has been demonstrated behaviorally to use magnetic information integrated with
visual cues for orientation (Dreyer et al., 2018a, Dreyer et al., 2025). To test whether
more generalist noctuid moth migrants also rely on a magnetic compass for
orientation, we investigated an invasive fall armyworm population in China using a
newly developed tethered-flight assay designed to quantify migratory orientation (Fig.
1). Although highly reductionist relative to natural flight conditions, particularly with
regard to the artificial visual cue, this indoor assay provides a controlled framework
for isolating the contribution of geomagnetic and visual cues as a first step toward

understanding how these signals operate under more realistic ecological scenarios.
The results of these experiments are not only of fundamental interest to sensory
biology, but are also of applied importance given that the fall armyworm is one of the
most mobile and invasive crop pests in the world (Warrant & Dacke, 2011; Kenis et
al., 2023; Tay et al., 2023).

Fig. 1. Schematic of the experimental setup for studying magnetic orientation in fall
armyworms. Moths are tethered to a vertical shaft at the center of the virtual flight
simulator, with an encoder recording their flight heading. A full-spectrum lamp illuminates
the arena, while the computer controlling the experiment is positioned outside the light
field to avoid interference. Moths are free to rotate in any direction during the assay. Full
experimental details in Methods. The cylinder is illustrated as clear in the figure to reveal
the internal setup, but it is opaque in the actual experiment.

Results

Integration of geomagnetic and visual cues in the seasonal migratory
orientation of field-captured fall armyworms

To examine if fall armyworms integrate geomagnetic and visual cues for seasonal
migratory orientation, we measured flight responses of tethered moths within a virtual
flight simulator (Dreyer et al., 2021) using a modified experimental approach used in
Bogong moth studies (Dreyer et al., 2018a; Dreyer et al., 2025). The simulator
consisted of a polyvinyl chloride (PVC) cylinder incorporating a visual cue on the side
(a black triangle rising above a black horizon; Fig. 1). When tethered within the
simulator, the moth is restrained but is free to rotate and thus take up any orientation
it chooses. The simulator was placed within a 3D Helmholtz coil system. Under

control conditions, the coil was switched off, leaving the Earth's natural magnetic field
(NMF) unaltered; under these conditions, magnetic north and geographic north were
closely aligned. However, when the coils were switched on, they reversed the
horizontal direction of the local magnetic field by 180° (the "Changed Magnetic Field",
CMF) relative to the NMF, while maintaining constant field intensity and inclination
angle (Fig. S1). Thus the field direction within the experimental arena temporarily
switches so that its "local magnetic north" is aligned with geomagnetic south (GMS)
and its "local magnetic south" is aligned with geomagnetic north (GMN), but all other
field parameters remain constant. The experimental setup consisted of recording
moth flight headings across five consecutive 5-minute phases (I, II, III, IV, V) of
tethered flight under different experimental conditions that involved changing the
azimuthal alignment of the visual cue and the local horizontal magnetic field
component with respect to GMN and GMS (Fig. 2, Fig. S1). The assay was divided
into these 5-min segments to provide the temporal resolution needed to detect
changes in flight orientation as the relative alignment of magnetic and visual cues
was systematically altered. In phases I and V moths were exposed to the NMF,
whereas in phases II–IV moths were exposed to the CMF (Fig. 2). During spring
migration trials, the visual cue was aligned with GMN (the expected migratory
orientation in spring) at the start of the trial, whereas in fall migration trials the set-up
was reversed, with the visual cue initially aligned with GMS (the expected migratory
orientation in fall; Fig. 2).

The first two experiments involved field-captured moths tested during spring and fall
migration periods (Figs. 2*A* and 2*B*). In phase I, the visual cue was aligned with the
expected seasonal magnetic direction in the NMF, and moths exhibited significant
group orientation towards the visual cue in both seasons (Spring: mean vector [MV] =
347.4° (95% CI: 319.5°, 16.71°), vector strength [R^*] = 1.59, $P < 0.001$, Fig. 2*A-I*; Fall:
MV = 183.6° (95% CI:211°,162°), $R^* = 1.76$, $P < 0.001$; Fig. 2*B-I*). In phase II, the
horizontal component of the geomagnetic field was rotated 180° (the CMF condition),
creating a conflict between the visual cue direction and the expected magnetic
orientation. Despite the shift in the magnetic field direction, moths continued to show
significant group orientation toward the visual cue during this 5-minute period (Spring:
MV = 319.1° (95% CI:351°,283°), $R^* = 1.39$, $P < 0.005$, Fig. 2*A-II*; Fall: MV = 191.0°
(95% CI:235°,153°), $R^* = 1.16$, $P < 0.05$, Fig. 2*B-II*), indicating that, at least initially,
the visual cue was dominant compared to the magnetic cue. During phase III, which
was a second 5-minute period of the experimental conditions applied in phase II,
moths lost their significant group-level orientation in both seasons (Spring: $R^* = 0.67$,

P > 0.05, Fig. 2*A-III*; Fall: *R** = 0.34, *P* > 0.05; Fig. 2*B-III*), indicating that over time

they had become confused due to the conflicting nature of the cues. Although we did

not analyze flight activity level due to technical limitations, the loss of group

orientation is unlikely to be attributable to a decline in the moths' motivation to

maintain orientation during prolonged flight in the simplified experimental

environment, as the Rayleigh test showed no significant differences in the *r*-values of

individual flight vectors across phases I–III (Figs. 2*A* and 2*B*; Table S2; Table S3). In

phase IV, the visual cue was realigned with the expected seasonal magnetic

orientation, but this time in the CMF (i.e., the cues were arranged in the same way as

in phase I but rotated by 180°). Therefore, the moths should not be able to

distinguish between phase I and phase IV and thus are expected to show the same

orientation response in both phases. Indeed, the moths tended to show group

orientation towards the congruent cues once again, albeit not quite reaching

significance in spring (Spring: MV = 245.2° (95% CI: 193.2°, 309.4°), *R** = 0.95, 0.05

< *P* < 0.10, Fig. 2*A-IV*; Fall: MV = 354.3° (95% CI: 331.4°, 27.2°), *R** = 1.49, *P* <

0.005; Fig 2*B-IV*). Finally, in phase V, the initial configuration was restored, and

moths regained significant group orientation toward the congruent visual cue and the

expected magnetic orientation in the NMF (Spring: MV = 343.8° (95% CI: 326.56°,

2.92°), *R** = 1.84, *P* < 0.001, Fig. 2*A-V*; Fall: MV = 167.7° (95% CI: 191°, 146°), *R** =

1.62, *P* < 0.001, Fig 2*B-V*). These results demonstrate that fall armyworm integrates

geomagnetic field information with visual cues to achieve stable orientation. However,

when geomagnetic and visual cues do not align with expected seasonal directions,

moths gradually lose orientation, reinforcing the critical role of cue integration in

maintaining migratory stability.

Integration of Geomagnetic and Visual Cues Is Consistent in Laboratory-
Reared Fall Armyworms

Our previous research showed that fall armyworms reared under artificially simulated

fall conditions and tested indoors with optic flow and starry sky projections exhibited

southward orientation in a flight simulator, consistent with the behavior observed in

wild individuals tested outdoors under natural night skies (Chen et al., 2023). To

determine whether lab-raised moths exposed to simulated seasonal photoperiods

respond the same way to geomagnetic and visual cues, we tested a population

reared under a simulated fall photoperiod in the same experimental setup as that of

the field-captured fall group (Fig. 2*C*). The results closely matched those from the

field-captured wild population (Fig. 2*B*). In phase I, with visual cues aligned to GMS

in the NMF, lab-raised moths oriented significantly southward (MV = 182.9° (95% CI:

203°, 156°), R^* = 1.77, P < 0.001, Fig. 2*C-I*). When the geomagnetic field was

reversed (CMF), moths initially orientated towards the visual cue, showing no

immediate response to the shifted field (MV = 184.9° (95% CI: 222°, 153°), R^* =1.31,

P < 0.05, Fig. 2*C-II*). After an additional 5-minute period without changes, significant

group-level orientation was lost (R^* = 0.69, P > 0.05, Fig. 2*C-III*). When the visual cue

was realigned with the changed GMS, significant orientation was restored (MV = 4.1°

(95% CI: 350.7°, 19.3°), R^* = 2.17, P < 0.001, Fig. 2*C-IV*), though the direction was

opposite that in phase I. Finally, returning to the original setup restored the initial

orientation pattern (MV = 137.6° (95% CI: 189.9°, 94.9°), R^* = 1.04, P < 0.05, Fig.

2*C-V*). These results demonstrate that lab-raised fall armyworms integrate

geomagnetic and visual cues similarly to wild populations, emphasizing the role of

photoperiod during development in shaping seasonal multimodal migratory

orientation.

To confirm that the loss of orientation in experimental groups was due to the cue

conflict rather than diminished motivation to maintain orientation arising from a

prolonged test duration, we conducted a control experiment with the lab-raised

simulated fall population, where the visual cue and GMS remained consistently

aligned. In this group, moths maintained significant orientation in their expected

migratory direction throughout the experiment when exposed to congruently-aligned

cues, irrespective of whether under NMF or CMF conditions (Fig. 2*D*; directional

statistics in Table S1). This confirms that the loss of orientation observed in

experimental groups was driven by misalignment between visual and geomagnetic

cues.

Fig. 2 The Earth's magnetic field and visual cues guide migratory flight behavior in both field-captured and lab-raised fall armyworms. (*A*) Flight orientation behavior of spring field-captured moths ("Spring Exp. Field") in response to visual and geomagnetic cues. (*B*) Flight orientation behavior of fall field-captured moths ("Fall Exp. Field") in response to visual and geomagnetic cues. (*C*) Flight orientation behavior of lab-raised fall-conditioned moths ("Fall Exp. Lab") in response to visual and geomagnetic cues. (*D*) Flight orientation behavior of lab-raised control fall-conditioned moths ("Fall Control"), tested with consistent visual and geomagnetic alignment. For simplicity and consistency, we tested only conditions in which the visual cue pointed in the migratory direction, coinciding with magnetic north in spring and with magnetic south in fall. Each group underwent five sequential 5-minute phases (I–V), with each subplot representing individual moths' flight directions in the simulator. The length of each vector represents individual directedness (*r*), ranging from 0 to 1, where the outer edge of the plot corresponds to *r* = 1. The thick mean vector (MV) arrow represents the weighted average of individual orientations, calculated using Moore's modified Rayleigh test (see

Methods), and is red when there is significant group orientation but grey when it is
not significant. The R^* value quantifies the directedness of the MV. Dashed circles
indicate thresholds for statistical significance, with radii corresponding to $P < 0.05$
and $P < 0.01$. Shaded sections of the outer diameters of the circles represent the
95% confidence limits of the group orientation. The outermost radius represents R^*
= 2.5. The black triangle denotes the position of the visual cue, while the red
triangle indicates the direction of the expected migratory orientation in each season
(north in spring, south in fall). The experimental setup included both the natural
magnetic field (NMF, panels with light green background) and a changed magnetic
field (CMF, panels with light blue background) where the horizontal magnetic field
direction was reversed; further details of the magnetic field parameters in the NMF
and CMF are shown in Fig. S1.

Fall armyworms show a delayed response to changes in magnetic fields

Our results show that migratory fall armyworms rely on both the Earth's magnetic
field and visual cues to determine their flight direction. However, for moths to
maintain constant orientation, the two sensory inputs must remain congruent, and
when they come into conflict the moths become disoriented at the group level (but
not at the individual level; Fig. 2). Notably, this response to the conflict between
visual and geomagnetic cues was not instanteous but delayed until the second 5-
minute experimental phase with this condition, consistent with similar results from
Bogong moths (Dreyer et al., 2018a). Initial analyses using a 5-minute resolution
provided limited temporal detail. To improve resolution, we subdivided each 5-minute
phase into ten 30-second intervals and independently analyzed directional
consistency using Moore's Rayleigh test to calculate the R^* for each 30-second
period (Fig. 3).

Fig. 3. Fall armyworms exhibit a delayed response to changes in magnetic-visual cue alignment. The behavioral data for (*A*) fall field-captured experimental ("Fall Exp. Field", data from Fig. 2B), (*B*) fall lab-raised experimental ("Fall Exp. Lab", data from Fig. 2C), and (*C*) fall lab-raised control ("Fall Control", data from Fig. 2D) groups were analyzed in 30-second time bins, resulting in ten bins over each 5-minute phase of the experiment. For each group, Moore's modified Rayleigh test was applied, and the obtained R^* values were plotted against time. $R^* > 1$ indicates a significant collective orientation within that 30-second interval, while $R^* < 1$ indicates the absence of significant group-level orientation.

During phase I, R^* values in both experimental groups remained >1 (Figs. 3*A-I* and 3*B-I*), indicating significant orientation behavior. When the magnetic field was rotated to conflict with visual cue directions, R^* values did not drop immediately but instead declined gradually over time (Figs. 3*A-II* and 3*B-II*). By phase III, R^* values were consistently <1, indicating a loss of significant orientation (Figs. 3 *A-III* and 3 *B-III*). Upon realigning the magnetic field with visual cues, R^* values showed a gradual increase, eventually exceeding 1 in both groups (Figs. 3*A-IV* and 3*B-IV*). When the original configuration was restored, the field-captured group consistently exhibited R^* values above 1 (Fig. 3*A-V*), while the lab-raised group had R^* values hovering around 1 (Fig. 3*B-V*). These results indicate that fall armyworms require time to resolve conflicting navigational inputs before losing group orientation entirely, or to react to congruent cues and regain group orientation. In the lab-raised control group, where the geomagnetic field and visual cues remained consistently aligned, R^* values consistently remained above 1 (Fig. 3*C*), confirming that the decline in group-

318 level orientation observed in the experimental groups was driven by conflicts
319 between geomagnetic and visual cues, mirroring the results found in Bogong moths
320 (Dreyer et al., 2018a).

Visual cues are essential for magnetic orientation in fall armyworms

Fig. 4. Visual information is essential for maintaining group flight orientation in fall
armyworms. (*A*) The lab-raised, fall-conditioned population lost significant group
orientation under complete darkness ("Fall Exp. Dark"). (*B*) The lab-raised, fall-
conditioned population exhibited a significant loss of group orientation under illuminated
conditions where visual cues were reduced to the bare minimum ("Fall Exp. BMVC"), i.e.,
obvious visual cues such as the black triangle shown in Fig. 1 were not provided;
however, because the experiments were conducted under illuminated conditions,
complete elimination of all visual information is impossible. (*C*) The distribution of
Rayleigh test *r*-values for individual moth orientations across different treatment groups
(*Left*), and the distribution of average directional change per second (*Right*), the latter
reflecting flight stability. The box plots represent the interquartile range (IQR), with the
horizontal line inside indicating the median. Whiskers extend to the most extreme data
points within 1.5 times the IQR. Pairwise comparisons of the *r*-values were performed
using the Wilcoxon rank-sum test and Wilcoxon signed-rank test (The Wilcoxon signed-
rank test is only used to compare non-independent (i.e., paired or related) samples. For
more information , please refer to the Statistical Analysis section.), with multiple

comparisons corrected by the Benjamini-Hochberg method. Detailed statistical results
are provided in Table S4. Comparisons of the average directional change per second
were performed using same method as that used for r value, with statistical results
provided in Table S5. Groups labeled with different letters differ significantly ($P < 0.05$).
We also analyzed the *r*-values and average directional change per second in the Fall Exp.
Field, Fall Exp. Lab, Exp. Fall Control, and the experiment shown in this figure, with
results consistent with those shown here (see Tables S4 and S5, Fig. S4).

Growing evidence supports the idea that geomagnetic cues provide essential
compass information for orientation across a broad range of taxa, supporting diverse
navigational tasks (Mouritsen, 2018; Dreyer et al.,2018a; Fleischmann et al., 2018;
Wan et al., 2021; Grob et al., 2024; Goforth et al., 2025). To determine whether
geomagnetic input alone is sufficient for flight orientation in fall armyworms, we
conducted a series of experiments to evaluate the role of visual cues in their
orientation behavior. In particular, we examined whether these moths could maintain
orientation in the absence of visual cues.

As a control, we first tested lab-raised, fall-conditioned moths in an arena where
visual landmarks were aligned with GMS in the NMF (Fig. 4*A*). When illuminated,
moths showed significant group orientation towards the visual cue (MV = 218.48° (95%
CI: 242°, 187°), $R^* = 1.50$, $P < 0.005$, Fig. 4*A-I*), consistent with our previous results
(Fig. 2*B*). To determine whether moths could maintain orientation without visual
information, we then removed all light sources, allowing moths to continue flying in
total darkness. Under these conditions, moths lost significant group-level orientation
($R^* = 0.5$, $P > 0.05$, Fig. 4*A-II*), suggesting that their orientation behavior is
dependent on the presence of light. The loss of orientation may be due to the
absence of usable visual landmarks, or the lack of light in a dark environment
causing the moths to lose their sense of direction. However, it is still unclear how the
absence of usable visual landmarks in non-dark environments would affect the
moths' orientation behavior.

To further investigate this, we tested a lab-raised, fall-conditioned population of moths
in a simulator, primarily to determine whether moths could maintain orientation under
illuminated conditions where visual cues on the interior walls of the cylindrical flight
arena were minimized. In this experiment, we provided a uniform visual environment
(i.e., under illuminated conditions, visual cues were not provided on the interior walls
of the cylindrical flight arena). After recording 10 minutes of orientation behavior
under the NMF, we reversed the horizontal component of the geomagnetic field using
Helmholtz coils (CMF) and recorded their flight orientation for an additional 10

minutes. Despite the altered magnetic field direction and sufficient time for adaptation,
moths exhibited no significant group orientation across all four testing phases (Fig.
4*B*; directional statistics in Table S1). The lack of group-level orientation under near-
uniform visual conditions suggests two possibilities: individual moths may have
maintained stable flight, but due to the absence of reliable orientation cues, their
group-level directional choices were random; or their flight may have been unstable,
making it difficult to assess individual orientation behavior. We first compared the
differences in the Rayleigh test *r*-values of moths across different experimental
phases (Fig. 4*C, Left*), using the Wilcoxon signed-rank test for the non-independent
Fall Exp and Fall Dark groups, and the Wilcoxon rank-sum test for all other
independent group comparisons. This is because the Rayleigh test evaluates the
concentration of data, with an *r*-value close to 1 indicating highly consistent flight
directions, and close to 0 suggesting more random flight directions. The results
showed that under conditions with visual cues (Fall Exp. Dark-I), moths had
significantly higher *r*-values compared to conditions of complete darkness (Fall Exp.
Dark-II) or visual cues reduced to the bare minimum (Fall Exp. BMVC).

We also considered the limitations of the Rayleigh test, as its *r*-value only reflects the
overall directional tendency and does not reveal real-time flight stability. For example,
a moth exhibiting two stable flight phases, one oriented north and the other south,
would yield a low *r*-value despite maintaining stability within each phase. Conversely,
a moth losing flight stability could also produce a low *r*-value, making it difficult to
distinguish between these two scenarios based on *r*-value alone. Therefore, we
introduced the analysis of per-second angular change (Fig. 4*C, Right*) to assess
dynamic fluctuations in flight behavior, providing a clearer understanding of whether
the observed differences in *r*-values were due to stable orientation or a loss of flight
control. Our analysis showed that the experimental groups tested under conditions
lacking visual cues exhibited significantly lower flight stability than the other
experimental groups (Fig.4C; Fig. S4). Moths in these conditions exhibited
significantly higher angular change rates, indicating a greater tendency to alter flight
direction and an inability to maintain stable orientation. Notably, even though a small
subset of individuals in these groups managed to sustain stable flight, their
directional choices appeared random rather than goal-oriented (Figs. 4*A-II* and 4*B*).

Discussion

The integration of geomagnetic and visual cues for flight orientation has so far been
experimentally demonstrated only in the Australian Bogong moth, a species

undertaking a precisely oriented, long-distance round-trip migration to and from a
highly restricted aestivation site within a single generation (Dreyer et al., 2018a).
Although this life-history strategy is uncommon among migratory insects, our
development of an indoor behavioral paradigm in the fall armyworm, a globally
distributed nocturnal migratory insect with multi-generational and partially migratory
patterns in China, extends this dual-cue orientation strategy to a species exhibiting a
migratory ecology more typical of insects in general. These findings demonstrate that
this orientation mechanism is not unique to the Bogong moth and may instead
represent a conserved orientation mechanism broadly employed across migratory
moth species, though additional species will need to be examined. Crucially, we
show that visual cues are indispensable for magnetic orientation in the fall armyworm,
as directional responses disappeared in the absence of structured visual input.
Furthermore, our findings reinforce recent results on seasonal shifts in orientation
behavior in this species (Chen et al., 2023), highlighting the ecological relevance of
this guidance system.

Our findings thus emphasize the importance of integrating multiple cues for
successful orientation. It is often assumed that animals lack magnetic orientation
capabilities if they fail to orient under changes in the geomagnetic field (Pakhomov et
al., 2025), either in appropriate lighting conditions or in darkness, without additional
cues. However, our study demonstrates that the absence of group seasonal
orientation under geomagnetic fields, in full-spectrum light or in darkness, does not
preclude magnetic orientation when additional sensory cues are integrated.
Specifically in the Northern Hemisphere, a magnetic compass provides global
directional cues for fall armyworms, enabling them to align magnetic north (or
magnetic pole) and south (or magnetic equator) with visual cues during spring and
fall, respectively, to facilitate seasonal orientation—a hallmark of annual long-
distance migration in migratory species. Although the magnetic compass is critical, as
evidenced by the loss of group orientation when the horizontal geomagnetic field was
reversed while intensity and inclination remained unchanged, our results reveal that
magnetic orientation cannot occur without appropriate visual cues. We also show that
the loss of flight stability observed when visual cues are minimal is likely a major
cause of the lack of group magnetic orientation. This highlights the complexity of
magnetic orientation strategies in nocturnal migratory insects, emphasizing the
crucial interaction between the magnetic compass and visual navigation systems.
Further investigation by altering the geomagnetic polarity and vertical geomagnetic
component is required to determine whether the magnetic compass guiding seasonal

orientation in nocturnal migratory fall armyworms is polarity-sensitive, inclination-
sensitive or both, which is pivotal to understanding the biophysical mechanisms
underlying magnetoreception (Mouritsen, 2018; Hore et al., 2016; Grob et al., 2024).

Desert ants and Bogong moths have also been shown to integrate geomagnetic and
visual cues for navigation. Desert ants rely on a magnetic compass during look-back-
to-the-nest behavior, combining magnetic cues with local visual landmarks, such as
their nest (Grob et al., 2024). In contrast, Bogong moths have been demonstrated to
integrate global stellar cues with the Earth's magnetic field for orientation (Dreyer et
al., 2025). Like Bogong moths, fall armyworms can migrate hundreds, or even
thousands, of kilometers over several successive nights (Warrant et al., 2016; Li et
al., 2020). In nature, they are likely to encounter a range of luminance-gradient visual
cues, including relatively stable celestial cues as well as transient or shifting local
features encountered en route. Although such natural cues differ from our simplified
laboratory stimulus, they may represent intermittently sampled visual inputs that can
be optimally integrated with magnetic information, with the congruency between
visual and magnetic cues likely playing a key role in maintaining a stable compass
response. Whether the cues are static or changing, brief periods without them may
still allow the subsequent recovery of a stable long-distance orientation strategy.
Determining which types of natural visual cues support the magnetic-visual compass,
and how they interact with magnetic information, including how their momentary
alignment or angular relationship is integrated and how such visual cue–magnetic
field interactions may require time to influence orientation, together with elucidating
the genetic and ecological bases of multimodal orientation, will be important
objectives for future research.

In conclusion, our findings underscore the indispensability of visual cues in facilitating
seasonal magnetic orientation and, unlike some vertebrates that can rely solely on
magnetic information (Mouritsen, 2018), highlighting the complex, multimodal nature
of navigation in nocturnal migratory insects. This work advances our understanding
of migratory orientation strategies and raises the possibility that similar magnetic-
visual integration mechanisms may be shared among long-distance migratory moths.
Furthermore, the application of genome editing in fall armyworms, combined with the
indoor behavioral paradigm developed here, provides a promising avenue for
dissecting the molecular basis of magnetoreception and the interplay between
sensory systems.

Methods

Field population

All wild populations used in this study were collected from farmland in Yuanjiang Hani and Yi Autonomous County, Yunnan Province, a region primarily characterized by maize cultivation. Fall-generation populations were collected during September to October of 2023 and 2024, while spring-generation populations were collected from April to May in 2024, as mature larvae at the 5th to 6th instar stages. Each larva was individually collected and reared in a cylindrical container, fed with fresh maize leaves from the collection site until pupation. Upon pupation, larvae were transferred to transparent, thin plastic cups sealed with plastic film. A cotton ball moistened with water was placed in each cup to maintain adequate humidity during the pupal stage. After eclosion, adults were individually provided with a 10% honey solution. All experimental individuals were reared at the Yuanjiang Plant Protection Station, a facility located near agricultural fields and well isolated from urban light pollution. To maintain natural environmental conditions, all electrical devices were strictly excluded from the facility, and windows were kept open around the clock to allow for natural light cycles and ambient temperature conditions.

Lab-raised population

The laboratory-reared population of the fall armyworm used in this study was primarily derived from wild individuals collected from a maize field in Yuanjiang, Yunnan Province (Google Maps coordinates: 23.604°N, 101.977°E) during October to November 2022, with the exception of individuals used in experiments Fall Exp. Dark and Fall Exp. BMVC (see Fig. 4), which were collected from the same field location during October to November 2024. All experiments were conducted using individuals from the 3rd to 5th laboratory-reared generations.

For behavioral assays conducted under ambient temperature conditions, a programmed photoperiodic regimen was implemented to simulate the seasonal changes in day length typical of autumn in East Asia. All treatments were maintained at a constant temperature of 27 ± 1°C and a relative humidity of 60% ± 5%. In the fall photoperiod treatment, the initial light cycle at egg hatching was set to 13 hours of light and 11 hours of darkness (13L:11D), followed by a daily reduction of 2 minutes

514 in the light period. By the time adult moths emerged and were subjected to
515 experimental testing approximately one month later, the photoperiod had been
516 adjusted to approximately 12L:12D.

Attachment of tethering stalks on moths

We used the same method for the attachment of tethering stalks on moths as in the
previous study on fall armyworm moth orientation (Chen et al. 2023). Prior to
tethering, unmated 2-day-old adult moths were housed in plastic cups and sedated at
4°C for at least 1 minute to facilitate handling. The moths were then transferred to an
operational platform, where scales were carefully removed from the junction of the
dorsal thorax and abdomen to ensure secure attachment. The tether consisted of a
slender, non-magnetic copper stalk measuring 0.75 mm in diameter and
approximately 2 cm in length. The terminal end of the stalk was bent into a fork-like
structure, which was carefully affixed to the prepared thoracic-abdominal junction
using Pattex PSK12CT-2 glue. To maintain consistency, this standardized tethering
method was applied uniformly across all experimental protocols.

Behavioral apparatus

Our flight simulator system (Dreyer et al., 2021) is consistent with the one used in
previous research on fall armyworm (Chen et al., 2023), utilizing the Flash flight
simulator system; however, in contrast to previous studies, we did not include optic
flow in order to avoid introducing additional visual complexity. The Flash flight
simulator system was developed based on the early design of the Mouritsen-Frost
flight simulator, and adapted for our experiments conducted in Yuanjiang (longitude
101.98°E, latitude 23.60°N). The moths are able to freely rotate on the horizontal
plane within the simulator, with their azimuth measured by an encoder system with a
resolution of 0.9°. The flight direction (relative to magnetic north) is recorded in real
time by the Flash flight simulator data acquisition system developed by Hui Chen,
and the data is saved as angle values in a text file. In our experiments, the encoder
(made of non-magnetic materials) samples the azimuth five times per second and is
equipped with a graphical interface, which allows for continuous monitoring of the
moth's azimuth and flight status to reconstruct the flight path. The encoder is
connected to the axis that holds the moth, with the axis measuring 15 cm in length
and 1 mm in diameter, made of carbon fiber material and wrapped in a carbon fiber
tube to protect its structure.

The arena consists of an opaque PVC cylinder with a diameter of 400 mm, a height

of 500 mm, and a thickness of 5 mm as the main structure, surrounded by white
wallpaper, with a 20 cm high black wallpaper strip at the bottom simulating the
horizon. A visual cue is provided by a black isosceles triangle (10 cm high, 10 cm
base) made from black wallpaper and fixed to the horizon at the bottom of the arena.
The entire arena is placed on a square wooden board with a side length of 50 cm
and a thickness of 1 cm, covered with black blackout cloth. The bottom of the
wooden board is equipped with a plastic base to reduce friction, ensuring that the
arena can rotate smoothly and quickly during the experiment, changing the direction
of the visual cues while avoiding interference with the moth's activity.

In the BMVC experiment, we used an opaque white acrylic cylinder with the same
dimensions as the PVC cylinder as the arena. No modifications were made to the
interior of the arena, primarily because the interior of the white acrylic cylinder is
smooth and uniform enough, and any additional treatment could potentially introduce
noticeable visual cues. We also removed the carbon fiber crossbar used to fix the
simulator and instead fixed it directly to the top acrylic cover to further minimize
potential visual cues that the moth might use. Our goal was to minimize the visual
cues available to the moth as much as possible, rather than attempting to completely
eliminate all visual cues.

A DH1766-series linear, three-channel, programmable DC power supply (Beijing
Dahua Radio Instrument Factory, China) with adjustable current and voltage was
used to power the full-spectrum LED employed in the experiments. The light source
was enclosed within a 90 cm diameter GODOX softbox to ensure even distribution of
light within the experimental arena. The softbox is fixed to a Helmholtz coil, with the
flight simulator located at its center. The top of the arena is equipped with a UV-
transmitting acrylic panel and covered with Lee Filters 250 semi-white diffusion paper
to control the amount of light transmitted. Spectral measurements, which include both
spectral irradiance and composition, were taken at the position corresponding to the
moth's head using an ATP2000P spectrometer (Optosky Technology Co. Ltd.,
Xiamen, China), ensuring consistent lighting conditions across all visual treatments
(see Fig. S3).

Artificial simulation of the Earth's magnetic field

A three-dimensional pair of Helmholtz coils (Nanjing Science Sky Technology Limited)
was used to manipulate the Earth's magnetic field. The coils measured 1200 × 1100
× 1000 mm, with a cold-state direct current resistance of 2Ω per coil and an

insulation resistance exceeding 10MΩ. The magnetic field strength at the central X, Y,

and Z axes exceeded 10 GS@6A (maximum current), with a spatial uniformity of 0.5%

within a φ144 × 144 × 144 mm volume. At the start of each experiment, we used a

TM4300B handheld three-axis fluxgate magnetometer to measure the ambient

magnetic field strength and spatial vector intensity at the experimental site (averaged

over three measurements). A DH1766 series linear three channel output precision

programmable DC power supply (Beijing Dahua Radio Instrument Factory, China)

with adjustable current and voltage was employed to apply a controlled current to the

3D Helmholtz coils, reversing the horizontal geomagnetic component by

approximately 180°. Crucially, we ensured that total magnetic field intensity (E),

horizontal component intensity (H), and magnetic inclination angle (α) remained

statistically unchanged between the controlled magnetic field (CMF) and the normal

magnetic field (NMF) (see Fig. S1A-D). Electromagnetic noise at the experimental

site was quantified by measuring the magnetic component of the time-dependent

electromagnetic field across a 10 kHz resolution bandwidth. Measurements were

performed using a Spectran NF-5035 spectrum analyzer (1 Hz-30 MHz, Aaronia AG)

connected to an MDF-9400 magnetic-field antenna (9 kHz-400 MHz, Aaronia AG) via

a UBBV-MDF960X preamplifier (9 kHz-60 MHz, 25 dB gain) and a 10-m RF cable.

The frequency range from 1 to 10 MHz was measured using the external antenna-

preamplifier configuration, while low-frequency noise between 2 kHz and 1 MHz was

measured using the internal magnetic probe of the spectrum analyzer. During each

measurement session, the spectrum analyzer continuously scanned the frequency

range for 40 minutes, and values were logged using the instrument's peak-hold mode.

Pronounced noise peaks were detected in the 2 kHz-1 MHz range, whereas noise

levels from 1 to 10 MHz were low and relatively stable, ranging between 0.01 and

0.001 nT with only minor fluctuations (see Fig. S1E).

Experimental procedure

The orientation experiment followed the protocols established in our previous study

(Chen et al. 2023). Experiments began after complete darkness in the night sky: the

field population were tested after adapting to the natural dark environment for at least

60 minutes, while the lab-raised population started after adapting to the dark

environment for at least 60 minutes following the light-to-dark cycle switch in the

incubator, in a laboratory away from large electrical equipment, artificial noise, plants,

and other potential sources of interference. The computer screen was positioned

away from and facing away from the experimental setup. Weak red light was used only when attaching the tethering stalks and fixing the moth to the simulator. After the moth was fixed, a waiting period of 20 to 30 seconds was allowed for the subject to stabilize before data recording began. During this process, it was verified whether the insect could switch between clockwise and counterclockwise rotations. If the insect rotated in only one direction and could not fly stably, no further experiment was conducted. The encoder's horizontal placement was also checked to ensure it was not affected by tilting. The moth's wing vibrations were confirmed to be strong, with equal amplitude on both wings (indicating that the contact glue did not interfere with the wings). During the experiment, the insect's flight state was monitored by observing fluctuations in the pointer values on the experimental software. If the values did not change within 10 seconds, the experimenter needed to approach the flight arena and listen for any sounds of wing vibrations. If the insect stopped flying, the experimenter gently tapped the wall to stimulate the insect to continue flying. We analyzed individuals that were able to maintain fewer than four instances of wingbeats cessation during the experiment.

Statistical analysis

All statistical analyses and graph generation were performed using R (version 4.1.3), accessible at https://www.r-project.org/. A custom R script, incorporating bootstrap confidence intervals, was used for Moore's Modified Rayleigh Test (MMRT)—a non-parametric test increasingly applied in orientation behavior studies (Massy et al., 2021). MMRT accounts for both mean directions and directedness (vector lengths) of individuals, making it an alternative to the Rayleigh test. MMRT assesses whether a group exhibits significant directional tendencies, using the parameter R^* (a scaled alternative to r in circular regression), which quantifies the strength of orientation within a group, and the mean vector (MV), which represents the rank-weighted mean direction of the group. Polar plots display individual vectors, derived from azimuth data collected during the experiment by the encoder at a sampling rate of five times per second, and subsequently analyzed using the Rayleigh test. Each vector representing an individual moth's mean flight direction and directedness (r-value). The vector length reflects the proportion of time a moth maintains a particular direction, ranging from $r = 0$ (completely un-oriented) to $r = 1$ (fully oriented). MMRT requires the analysis of each individual's average direction and r-value to determine the group-level mean MV. It is important to note that the magnitude of R^* is dependent on sample size.

To evaluate the directional orientation of moths in different treatment groups, we

performed Rayleigh tests on individual flight directions. Subsequently, we used the Friedman test to examine whether different experimental phases within each experiment influenced the *r*-values of moth orientation, as this test is appropriate for comparing repeated measurements from the same individuals (Table S2). In the initial test, only the Exp. Spring experiment showed significant differences. Therefore, we further compared the flight stability and *r*-values among its different experimental phases (Table S3). Because r value exhibited non-normal distributions, the Wilcoxon signed-rank test was employed. Since these comparisons involved repeated measurements from the same individuals, pairwise comparisons were conducted. Because multiple comparisons were performed, failing to apply a correction would increase the likelihood of false positives; therefore, the Benjamini–Hochberg procedure was used to control the false discovery rate (FDR).

The Rayleigh *r*-values for each individual were also used to compare orientation strength across experiment. We accounted for the non-independence between Exp Dark I and Exp Dark II by using the paired Wilcoxon signed-rank test for their pairwise comparison. All other comparisons among independent experiments were performed using the Wilcoxon rank-sum test. The multiple comparisons corrected using the Benjamini-Hochberg procedure to control the false discovery rate (FDR). Detailed statistical results are provided in Table S4.

To evaluate flight stability, we measured the angular difference between consecutive seconds of recorded flight behavior and calculated the average directional change per second. This metric reflects the consistency and stability of individual flight orientation. Differences among treatment groups were assessed using the Wilcoxon signed-rank test (Table S5) because some of the data did not follow a normal distribution. We also accounted for the non-independence between the Exp Dark I and Exp Dark II experiments, for which paired comparisons were conducted. Multiple comparisons were corrected using the Benjamini–Hochberg procedure to control the false discovery rate (FDR).

Acknowledgements

This work was supported by the National Key Research and Development Program of China (2021YFD1400700) and the Fundamental Research Funds for the Central Universities (KJJQ2025013) to G.H., and the National Natural Science Foundation of China to (32202289) B.Y.G.

References

Alerstam T, Chapman JW, Bäckman J, Smith AD, Karlsson H, Nilsson C, Reynolds DR, Klaassen RHG, Hill JK, 2011. Convergent patterns of long-distance nocturnal migration in noctuid moths and passerine birds. *Proc. R. Soc. B* **278**:3074-3080. DOI: https://doi.org/10.1098/rspb.2011.0058

Bauer S & Hoye BJ, 2014. Migratory animals couple biodiversity and ecosystem functioning worldwide. *Science* **344**:54-62. DOI: https://doi.org/10.1126/science.1242552

Chapman JW, Nesbit RL, Burgin LE, Reynolds DR., Smith AD, Middleton DR, Hill JK. 2010. Flight orientation behaviors promote optimal migration trajectories in high-flying insects. *Science* **327**:682-685. DOI: https://doi.org/10.1126/science.1182990

Chapman JW, Reynolds DR, Wilson K, 2015. Long-range seasonal migration in insects: mechanisms, evolutionary drivers and ecological consequences. *Ecol. Lett.* **18**:287-302. DOI: https://doi.org/10.1111/ele.12407

Chapman JW, Nilsson C, Lim KS, Bäckman J, Reynolds DR, Alerstam T, 2016. Adaptive strategies in nocturnally migrating insects and songbirds: contrasting responses to wind. *J. Anim. Ecol.* **85**:115-124. DOI: https://doi.org/10.1111/1365-2656.12420

Chen H, Wang Y, Huang L, Xu CF, Li JH, Wang FY, Cheng W, Gao BY, Chapman JW, G. Hu, Flight capability and the low temperature threshold of a Chinese field population of the fall armyworm *Spodoptera frugiperda*. *Insects* **13**:422 (2022). DOI: https://doi.org/10.3390/insects13050422

Chen H, Wan GJ, Li JC, Ma YB, Reynolds DR, Dreyer D, Warrant EJ, Chapman JW, G. Hu, Adaptive migratory orientation of an invasive pest on a new continent. *iScience* **26**:108281 (2023). DOI: https://doi.org/10.1016/j.isci.2023.1082 81

Chen H, Xu CF, Wang YH, Li XR, Yu DH, Chen AD, Lyu BQ, Wu YF, Wang YM, Chapman JW, G. Hu, Characteristics and seasonal variation of fall armyworm migratory behavior in their year-round breeding areas in South China. *Pest Manag. Sci.* **81**:4051-4062 (2025). DOI: https://doi.org/10.1002/ps.8772

Drake VA & Reynolds DR. 2012. Radar entomology: observing insect flight and migration. Wallingford, UK: CABI

Dreyer D, Frost B, Mouritsen H, Günther A, Green K, Whitehouse M, Johnsen S, Heinze S, Warrant E, 2018a. The Earth's magnetic field and visual landmarks steer migratory flight behavior in the nocturnal Australian Bogong moth. *Curr. Biol.* **28**:2160-2166. DOI: https://doi.org/10.1016/j.cub.2018.05.030

Dreyer D, el Jundi B, Kishkinev D, Suchentrunk C, Campostrini L, Frost BJ, Zechmeister T, Warrant EJ, 2018b. Evidence for a southward autumn migration of nocturnal noctuid moths in central Europe. *J. Exp. Biol.* **221**:jeb179218. DOI: https://doi.org/10.1242/jeb.179218

Dreyer D, Frost B, Mouritsen H, Lefèvre A, Menz M, Warrant EJ, 2021. A guide for using flight simulators to study the sensory basis of long-distance migration in insects. *Front. Behav. Neurosci.* **15**:678936. DOI: https://doi.org/10.3389/fnbeh.2021.678936

Dreyer D, Adden A, Chen H, Frost B, H. Mouritsen, J. J. Xu, K. Green, M. Whitehouse, J. Chahl, J. Wallace, G. Hu, J. Foster, S. Heinze, E. J. Warrant,

2025. Bogong moths use a stellar compass for long-distance navigation at night. *Nature* **643**:994-1000. https://doi.org/10.1038/s41586-025-09135-3

Fleischmann PN, Grob R, Müller VL, Wehner R, Rössler W, 2018. The geomagnetic field is a compass cue in *Cataglyphis* ant navigation. *Curr. Biol.* **28**:1440-1444. DOI: https://doi.org/10.1016/j.cub.2018.03.043

Freas CA & Spetch ML, 2023. Varieties of visual navigation in insects. *Anim. Cogn.* **26**:319-342. https://doi.org/10.1016/j.conb.2011.05.020

Foster JJ, Smolka J, Nilsson DE, Dacke M, 2018. How animals follow the stars. *Proc. R. Soc. B* **285**:20172322. DOI: https://doi.org/10.1098/rspb.2017.2322

Fu P, Li X, Sun Z, Chen Y, Lu Z, Tian C, Hu G, Gui F, 2025. Population genetics and trajectory simulation reveals the invasion process of the fall armyworm (*Spodoptera frugiperda*) in the Eastern Hemisphere. *Evol. Appl.* **18**:e70086. DOI: https://doi.org/10.1111/eva.70086

Gao BY, Reynolds DR, Zhai BP, Hu G, Chapman JW, 2020. The 'migratory connectivity' concept, and its applicability to insect migrants. *Move. Ecol.* **8**:48. DOI: https://doi.org/10.1073/pnas.2317646121

Gao BY, Wotton KR, Hawkes WLS, Menz MHM, Reynolds DR, Zhai BP, Hu G, Chapman JW, 2020. Adaptive strategies of high-flying migratory hoverflies in response to wind currents. *Proc. R. Soc. B* **287**:20200406. DOI: https://doi.org/10.1098/rspb.2020.0406

Gao BY, Reynolds DR, Hu G, Chapman JW, 2024. Effects of nocturnal celestial illumination on high-flying insect migration. *Phil. Trans. Roy. Soc. B* **379**:20230115. DOI: https://doi.org/10.1098/rstb.2023.0115

Gao FQ, Chen H, Kyerematen R, Hu G, Early R, Chapman JW, 2025. High migratory potential of fall armyworm in West Africa despite stable temperatures and widely available year-round habitats. *Insect Sci.* https://doi.org/10.1111/1744-7917.13502 .

Goergen G, Kumar PL, Sankung SB, Togola A, Tamo M, 2016. First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in west and central Africa. *PLoS One* **11**:e0165632. DOI: https://doi.org/10.1371/journal.pone.0165632

Goforth KM, Lohmann CMF, Gavin A, Henning R, Harvey A, Hinton TL, Lim DS, Lohmann KJ, 2025. Learned magnetic map cues and two mechanisms of magnetoreception in turtles. *Nature* **638**:1015–1022. DOI: https://doi.org/10.1038/s41586-024-08554-y

Grob R, Wegmann JW, Rössler W, Fleischmann PN, 2024. *Cataglyphis* ants have a polarity-sensitive magnetic compass. *Curr. Biol.* **34**:5833-5838. DOI: https://doi.org/10.1016/j.cub.2024.11.012

R, Farnworth MS, Degen J, Warrant E, Montgomery SH, and el Jundi B, 2025. The diversity of lepidopteran spatial orientation strategies - neuronal mechanisms and emerging challenges in a changing world. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*. DOI: https://doi.org/10.1007/s00359-025-01780-3.

Guo J, Fu X, Zhao S, Shen X, Wyckhuys KAG, Wu K, 2020. Long-term shifts in abundance of (migratory) crop-feeding and beneficial insect species in northeastern Asia. *J. Pest Sci.* **93**:583-594. DOI: https://doi.org/10.1007/s10340-019-01191-9

Holland RA, Wikelski M, Wilcove DS. 2006, How and why do insects migrate?
Science, **313**:794-796. https://doi.org/10.1126/science.1127272

Hu G, Lim KS, Horvitz N, Clark SJ, D. R. Reynolds, N. Sapir, J. W. Chapman, Mass
seasonal bioflows of high-flying insect migrants. *Science* **354**:1584-1587 (2016).
DOI: https://doi.org/10.1126/science.aah4379

Hu G, Feng HQ, Otuka A, Reynolds DR, Drake VA, Chapman JW, 2025. The east
Asian Insect flyway: geographical and climatic factors driving migration among
diverse crop pests. *Annu. Rev. Entomol.* **70**:1-22. DOI: https://doi.org/10.1146/
annurev-ento-012524-124018

Huang JR, Feng HQ, Drake VA, Reynolds DR, Gao BY, Chen FJ, Zhang GY, Zhu J,
Gao YB, Zhai BP, Li GP, Tian CH, Huang B, Hu G, Chapman JW, 2024. Massive
seasonal high-altitude migrations of nocturnal insects above the agricultural
plains of East China. *Proc. Nat. Acad. Sci. USA.* **121**:e2317646121.
DOI: https://doi.org/10.1073/pnas.2317646121

Hore PJ, Mouritsen H, 2016. The radical-pair mechanism of magnetoreception. *Annu.*
Rev. Biophys. **45**:299-344. DOI: https://doi.org/10.1146/annurev-biophys-
032116-094545

Kenis M, Benelli G, Biondi A, Calatayud PA, Day R, Desneux N, Harrison RD,
Kriticos D, Rwomushana I, van den Berg J, Verheggen F, Zhang YJ, Agboyi LK,
Ahissou RB, Ba MN, Wu KM, 2023. Invasiveness, biology, ecology, and
management of the fall armyworm, *Spodoptera frugiperda*. *Entomol. Gen.*
43:187–241. DOI: https://doi.org/10.1127/entomologia/2022/1659

Li XJ, Wu MF, Ma J, Gao BY, Wu QL, Chen AD, Liu J, Jiang YY, Zhai BP, Early R,
Chapman JW, Hu G, 2020. Prediction of migratory routes of the invasive fall
armyworm in eastern China using a trajectory analytical approach. *Pest Manag.*
Sci. **76**:454-463. DOI: https://doi.org/10.1002/ps.5530

Massy R, Hawkes WLS, Doyle T, Troscianko J, Menz MHM, Roberts NM, Chapman
JW, Wotton KR, 2021. Hoverflies use a time-compensated sun compass to
orientate during autumn migration. *Proc. R. Soc. B* **288**:20211805. DOI:
https://doi.org/10.1098/rspb.2021.1805

Menz MHM, Scacco M, Bürki-Spycher HM, Williams HJ, Reynolds DR, Chapman JW,
Wikelski M, 2022. Individual tracking reveals long-distance flight-path control in a
nocturnally migrating moth. *Science* **377**:764-768. DOI:
https://doi.org/10.1126/science.abn166

Mouritsen H & Frost B, 2002. Virtual migration in tethered flying monarch butterflies
reveals their orientation mechanisms. *Proc. Nat. Acad. Sci. USA.* **99**:10162-
10166. DOI: https://doi.org/10.1073/pnas.152137299

Mouritsen H, 2018. Long-distance navigation and magnetoreception in migratory
animals. *Nature* **558**:50-59. https://doi.org/10.1038/s41586-018-0176-1

Nagoshi RN & Meagher RL, 2008. Review of fall armyworm (Lepidoptera: Noctuidae)
genetic complexity and migration. *Fla Entomol* **91**:546-554. DOI:
https://doi.org/10.1653/0015-4040-91.4.546

Pakhomov A, Shapoval A, Shapoval N, and Kishkinev D, 2025. Not all butterflies are
monarchs: compass systems in the red admiral (*Vanessa atalanta*), a European
diurnal migrant. *bioRxiv*. DOI: https://doi.org/10.1101/2025.05.01.651646

Reppert SM, Guerra PA, Merlin C, 2016. Neurobiology of monarch butterfly migration.
Annu. Rev. Entomol. **61**:25-42. DOI: https://doi.org/10.1146/annurev-ento-

010814-020855

Srygley RB & Dudley R, 2008. Optimal strategies for insects migrating in the flight
boundary layer: Mechanisms and consequences. *Integr. Comp. Biol.* **48**:119-133.
DOI: https://doi.org/10.1093/icb/icn011

Tay WT, Meagher RL, Czepak C, Groot AT, 2023. *Spodoptera frugiperda*: Ecology,
evolution, and management options of an invasive species. *Annu. Rev. Entomol.*
68:299-317. DOI: https://doi.org/10.1146/annurev-ento-120220-102548

Tong DD, Zhang L, Wu NN, Xie DJ, Fang GJ, Coates BS, Sappington TW, Liu YQ,
Cheng YX, Xia JX, Jiang XF, Zhan S, 2022. The oriental armyworm genome
yields insights into the long-distance migration of noctuid moths. *Cell Rep.*
41:111843. DOI: https://doi.org/10.1016/j.celrep.2022.111843

Wan GJ, Hayden AN, Iiams SE, Merlin C, 2021. Cryptochrome 1 mediates light-
dependent inclination magnetosensing in monarch butterflies. *Nat. Comm.*
12:771. DOI: https://doi.org/10.1038/s41467-021-21002-z

Warrant EJ & Dacke M, 2011. Vision and visual navigation in nocturnal insects. *Ann.*
Rev. Entomol. **56**:239-254. DOI: https://doi.org/10.1146/annurev-ento-120709-
144852

Warrant EJ, Frost B, Green K, Mouritsen H, Dreyer D, Adden A, Brauburger K,
Heinze S, 2016. The Australian Bogong moth *Agrotis infusa*: A long-distance
nocturnal navigator. *Front. Behav. Neurosci.* **10**:77. DOI: https://doi.org/10.3389/
fnbeh.2016.00077

Warrant E & Somanathan H, 2022. Colour vision in nocturnal insects. *Philos Trans R*
Soc Lond B Biol Sci **377**:20210285. DOI: https://doi.org/10.1098/rstb.2021.0285

Werber Y, Adin E, Chapman JW, Reynolds DR, Sapir N, 2025. Active navigation and
meteorological selectivity drive mass intercontinental insect migration patterns
through the Levant. *Proc. R. Soc. B* **292**:20250587. DOI:
https://doi.org/10.1098/rspb.2025.0587

Westbrook J, Fleischer S, Jairam S, Meagher R, Nagoshi R, 2019. Multigenerational
migration of fall armyworm, a pest insect. *Ecosphere* **10**:e02919. DOI:
https://doi.org/10.1002/ecs2.2919

Wu QL, Jiang YY, Liu J, Hu G, Wu KM, 2021. Trajectory modeling revealed a
southwest-northeast migration corridor for fall armyworm *Spodoptera frugiperda*
(Lepidoptera: Noctuidae) emerging from the North China Plain. *Insect Sci.*
28:649-661. DOI: https://doi.org/10.1111/1744-7917.12852

Wu MF, Qi GJ, Chen H, Ma J, Liu J, Jiang YY, Lee GS, Otuka A, Hu G, 2022.
Overseas immigration of fall armyworm, *Spodoptera frugiperda* (Lepidoptera:
Noctuidae) invading Korea and Japan in 2019. *Insect Sci.* **29**:505-520. DOI:
https://doi.org/10.1111/1744-7917.12940

Satterfield DA, Sillett TS, Chapman JW, Altizer S, Marra PP, 2020. Seasonal insect
migrations: massive, influential, and overlooked. *Front. Ecol. Environ.* **18**:335-
344. DOI: https://doi.org/10.1002/fee.2217

Sun XX, Hu CX, Jia HR, Wu QL, Shen XJ, Zhao SY, Jiang YY, Wu KM, 2021. Case
study on the first immigration of fall armyworm, *Spodoptera frugiperda* invading
into China. *J. Integr. Agri.* **20**:664-672. DOI: https://doi.org/10.1016/s2095-
3119(19)62839-x

Xu JJ, Pan W, Zhang YC, Li Y, Wan GJ, Chen FJ, Sword GA, Pan WD, 2017.

Behavioral evidence for a magnetic sense in the oriental armyworm, *Mythimna*

separata. Biol. Open **6**:340-347. DOI: https://doi.org/10.1242/bio.022954

 Supplementary information

876 **Table S1. Detailed data of flight orientation behavior assay.**

Experiment	period	MV (95% CI lower-upper)	R*	n	p
Fall Exp. Field	I	183.6 (211-162)	1.76	29	p < 0.001
	II	191.0 (235-153)	1.16	29	0.01 < p < 0.025
	III	158.1 (316.3-21.2)	0.34	29	0.5 < p < 0.9
	IV	354.3 (331.4-27.2)	1.49	29	0.001 < p < 0.005
	V	167.7 (191-146)	1.62	29	p < 0.001
Spring Exp. Field	I	347.4 (319.5-16.7)	1.59	28	p < 0.001
	II	319.1 (351-283)	1.39	28	0.001 < p < 0.005
	III	293.7 (33.6-201.3)	0.67	28	0.1 < p < 0.5
	IV	245.2 (309.4-193.2)	0.95	28	0.05 < p < 0.1
	V	343.8 (326.56-2.92)	1.84	28	p < 0.001
Fall Exp. Lab	I	182.9 (203-156)	1.77	27	p < 0.001
	II	184.9 (222-153)	1.31	27	0.005 < p < 0.01
	III	111.3 (203.9-36.6)	0.69	27	0.1 < p < 0.5
	IV	4.1 (350.7-19.3)	2.17	27	p < 0.001
	V	137.6 (189.9-94.9)	1.04	27	0.025 < p < 0.05
Fall Control	I	180.3 (196-164)	2.10	26	p < 0.001
	II	184.1 (199-167)	1.94	26	p < 0.001
	III	195.2 (219-160)	1.43	26	0.001 < p < 0.005
	IV	6.1 (28.8-340.4)	1.63	26	p < 0.001
	V	183.0 (205-162)	1.81	26	p < 0.001
Fall Exp. Dark	Control	218.48 (242-187)	1.50	31	0.001 < p < 0.005
	Dark	41.03 (178.51-5.31)	0.55	31	0.1 < p < 0.5
Exp. BMVC	I	114.51 (277-18.8)	0.32	30	0.5 < p < 0.9
	II	215.76 (284-119)	0.65	30	0.1 < p < 0.5
	III	105.72 (264.8-13.9)	0.31	30	0.5 < p < 0.9
	IV	191.21(284-119)	0.66	30	0.1 < p < 0.5

877
878

879 **Table S2. Friedman test results of Rayleigh's r values across experimental**

Experiment	Friedman chi-squared	df	p-value
Spring Exp. Field	16.207	4	0.002753
Fall Exp. Field	1.1714	4	0.8828
Fall Exp. Lab	8.2154	4	0.084
Fall Control	1.952	4	0.7446

881 **Table S3. Pairwise comparison results of the mean Rayleigh's r values of Exp.**
882 **Spring across different experimental phases, using the Wilcoxon signed rank**
883 **test with multiple comparison correction performed by the Benjamini-**
884 **Hochberg method.**

Experiment	Comparison group	P value	W value
Spring Exp. Field	IV – V	0.1545	140
	III-V	0.1545	141
	II-V	0.0519	115
	I-V	0.0021	44
	III-IV	0.9153	184
	II-IV	0.5166	154
	I-IV	0.1545	84
	II-III	0.9153	210
	I-III	0.3691	122
	I-II	0.2615	116

885
886
887 **Table S4. Multiple comparisons of mean Rayleigh test _r_ values across**
888 **experimental conditions using the Wilcoxon rank-sum test (adjusted by the**
889 **Benjamini-Hochberg method).**

Comparison	P value	W value
A vs B	0.32	9084
A vs C	0.01	7632
A vs D	0.54	2407
A vs E	<0.001	3252
A vs F	<0.001	17135
B vs C	0.22	7934
B vs D	0.26	2386
B vs E	<0.001	2958
B vs F	<0.001	15654
C vs D	0.0356	2543
C vs E	<0.001	3186
C vs F	<0.001	16450
D vs E	0.01	400
D vs F	<0.001	3477
E vs F	0.0700	2843

890 **Note:** A: Fall Exp. Field; B: Fall Exp. Lab; C: Fall Control; D: Fall Exp. Dark I; E: Fall Exp.
891 Dark II; F: Fall Exp. BMVC.
892

893   **Table S5. Wilcoxon rank-sum test for mean angular velocity (directional change**

894   **per second) across experimental conditions,adjusted by the Benjamini-**

895   **Hochberg method.**

Comparison group	*P* value	W value
A vs B	0.75	10006
A vs C	0.41	9967
A vs D	0.94	2227
A vs E	0.00	1479
A vs F	0.00	4237
B vs C	0.77	8957
B vs D	0.72	2005
B vs E	0.00	1376
B vs F	0.00	3890
C vs D	0.48	1850
C vs E	0.00	1119
C vs F	0.00	3199
D vs E	0.01	119
D vs F	0.00	929
E vs F	0.02	1332

896   **Note:** A: Fall Exp. Field; B: Fall Exp. Lab; C: Fall Control; D: Fall Exp. Dark-I; E: Fall

897   Exp. Dark-II; F: Fall Exp. BMVC.

Fig. S1. The magnetic field conditions during experimental procedures in 2023 and 2024. A Comparison of total magnetic field strength between the controlled magnetic field (CMF) and normal magnetic field (NMF) (*t*-test, E_{CMF} = 42.21 µT, E_{NMF}

= 42.02 µT, P = 0.7869), showing no significant difference. **B** There is no significant

difference in the intensity of the horizontal component between CMF and NMF (*t*-test,

H_{CMF} = 31.18, H_{NMF} = 31.35, P >0.1). **C** There is no significant difference in magnetic

inclination between CMF and NMF (*Watson-U²*, α_{CMF} = 38.22°, α_{NMF} = 38.28°, P =

0.92). **D** Comparison of magnetic azimuth between CMF and NMF, demonstrating a

highly significant difference (*Watson-U²*, β_{CMF} =175.62°, β_{NMF} = 4.11°, P < 0.001). **E**

The magnetic component B, measured in nanoteslas (nT), of the time-dependent

electromagnetic field across a 10 kHz resolution bandwidth (RBW), was analyzed as

a function of frequency f, expressed in megahertz (MHz). The data shown in the

figure represents the maximum locked values observed over a 40-minute period at

the experimental location (23.604°N, 101.977°E). Measurements were conducted

using the Spectran NF5035 spectrum analyzer (1 Hz-30 MHz, Aaronia AG),

MDF9400 magnetic field antenna (9 kHz-400 MHz, Aaronia AG), UBBV-MDF960X

preamplifier (9 kHz-60 MHz, with a gain of 25 dB), and a 10-meter RF cable. The

spectrum in the frequency range of 1 MHz to 10 MHz was first measured using the

antenna paired with the preamplifier. Subsequently, low-frequency noise within the

range of 2 kHz to 1 MHz was detected using the internal probe of the spectrum

analyzer.

925
926
927

Fig. S2 Year-specific analysis of orientation behavior in field-captured
armyworms during the fall migration season (2023 and 2024). A Orientation
behavior of the field-captured population in fall 2023. **B** Orientation behavior of the
field-captured population in fall 2024. Experiments were conducted using fall
armyworms captured from the field, revealing consistent orientation behavior across
both years. In both 2023 and 2024, moths oriented significantly toward the visual cue
when visual and geomagnetic cues were aligned. However, after reversing the
magnetic field, the strength of directional orientation gradually diminished. Due to
limited sample sizes in each year, data from both years were pooled for a
comprehensive analysis, as presented in Fig. 2B.

Fig. S3. Spectral distribution of light provided by the full-spectrum lamp. The spectral distribution of the full-spectrum lamp used in the experiment was measured using an ATP2000P spectrometer (Optosky Technology Co. Ltd., Xiamen, China). The sensor was positioned vertically upward at the location originally occupied by the moth's head to measure the spectral irradiance of the light after it passed through the softbox, diffusion paper, and transparent acrylic cover. The results indicate that light radiation was detectable across most wavelengths between 200 nm and 850 nm, confirming that our experimental setup provided a full-spectrum light environment from ultraviolet to near-infrared wavelengths. This spectral range encompasses the known visual sensitivity of moths (Warrant & Somanathan, 2022). Consequently, the acrylic plate and diffusion paper did not significantly block light radiation at any specific wavelength within this range. The ambient light level in the experimental environment was measured to be below 1 lux using a Testo 540 lux meter, manufactured by Testo SE & Co. KGaA (Titisee-Neustadt, Germany).

955
956

Fig. S4 The lack of visual cues and other necessary visual information significantly affects moth orientation ability and flight stability. A Box plots show the distribution of Rayleigh test *r* values for individual orientations of moths in different treatment groups. **B** Box plots show the distribution of average directional degree change per second, reflecting flight stability, across treatment groups. Boxes represent the interquartile range (IQR), with the horizontal line inside indicating the median. The "whiskers" extend to the most extreme data points within 1.5 times the IQR. Error bars indicate data variability, facilitating comparison between groups. Pairwise comparisons in **A** were conducted using the Wilcoxon rank-sum test with multiple comparisons corrected by the Benjamini-Hochberg method. Detailed statistical results are provided in Supplementary Table S4. In **B**, comparisons were performed using Wilcoxon rank-sum test. Groups labeled with different letters differ significantly ($P < 0.05$). Detailed statistical results are provided in Supplementary Table S5.