Characterization of a dominant *SmNac-like* gene as a candidate for

photosensitivity in the fruit peel of eggplant

Joaquin Gomis-Cebolla[1,2,†,*], Silvia Manrique[1,†], Andrea Arrones[1], Maria Dolores Toledo-

Tolgar[1], Jose Luna[1], Virginia Baraja-Fonseca[1], Joan Sánchez-Pascual[1], Esther Gimeno-Páez[1],

Mariola Plazas[1], Pietro Gramazio[1], Santiago Vilanova[1], Jaime Prohens[1].

[1]Institute for Conservation and Improvement of Valencian Agrodiversity (COMAV),

Universitat Politècnica de València, Camí de Vera 14, 46022 Valencia, Spain.

[2]I2SysBio, CSIC-University of Valencia, Parc Científic Universitat de València, Calle

Catedrático Agustín Escardino, 9, 46980, Paterna, Spain.

† These authors contributed equally to this work.

* Corresponding author: Joaquin Gomis-Cebolla

17   I2SysBio, CSIC-University of Valencia, Parc Científic Universitat de

18   València, Calle Catedrático Agustín Escardino, 9, 46980, Paterna,

19   Spain. Email: joaquin.gomis@csic.es

23 Running title: *SmNac-like* gene controls photosensitivity trait in eggplant

ABSTRACT

Anthocyanins in the fruit peel of photosensitive eggplants exhibit a different distribution pattern compared to the photo-insensitive ones. The latter exhibits a uniform anthocyanin content, whereas photosensitive eggplants lack anthocyanin accumulation in areas not exposed to light, such as under the calyx, or have lower concentrations in less-exposed areas. In the current research work, genetic analysis of F1 and F2 populations revealed that the non-photosensitive phenotype in eggplants follows an autosomal dominant inheritance with a 3:1 ratio, indicating that the photosensitive trait is regulated by a single dominant gene. To locate and narrow down the position of the gene underlying photosensitivity, an F2 population segregating for photosensitivity and presence of anthocyanins was used for bulked segregant analysis sequencing (BSA-seq), and RNA-seq analysis of bagged photosensitive and non-photosensitive eggplants was performed. As a result, we identified the *SmNAC1-like* gene as a candidate for fruit photosensitivity trait in the eggplant materials used in this study. In the photo-insensitive eggplants (DH-ECAVI and IVIA-371) two different high-impact mutations were identified leading to light-insensitive and uniform anthocyanin content. The mutation in DH-ECAVI causes a premature stop codon, while in IVIA-371 an indel mutation occurs in the 3'-UTR sequence. No high-impact mutations were found in *SmNAC1-like* gene in photo-sensitive eggplants. These findings provide new insight into the regulation of the molecular mechanisms of anthocyanin biosynthesis in non-photosensitive eggplant and have important implications for marker-assisted selection and breeding.

KEYWORDS

Solanum melongena, *SmNAC1-like*, Anthocyanin Biosynthesis and Photosensitivity

INTRODUCTION

Eggplant (*Solanum melongena* L.) is a solanaceous crop with high economic value and characterized by the presence of high concentrations of anthocyanins in the fruit peel of some cultivars. Anthocyanins are water-soluble pigments derived from anthocyanidins by the addition of sugars, which make flowers and fruits appear red, blue or purple (Takos et al., 2006 and Mattioli et al., 2020). In addition, anthocyanins offer several benefits to plants, such as protection against UV radiation, extreme temperatures, and pathogen attacks (Lorenc-Kukula et al., 2005; Liu et al., 2013 and Ren et al., 2014). Finally, anthocyanins contribute to human health, since they present strong antioxidant properties, with beneficial effects on human well-being (Muñoz-Falcón et al., 2009; Speer et al., 2020).

Light plays a major role in the synthesis of anthocyanins in the fruit peel of eggplants. Previous research in photosensitive eggplants demonstrated that light activates the anthocyanin biosynthetic pathway leading to the accumulation of anthocyanins in the fruit peel in a time-exposure manner (Jiang et al., 2016; Li et al., 2017; Sang et al., 2022, Li et al., 2023). However, non-photosensitive eggplants are able to synthesize large amounts of anthocyanins under shaded conditions (Jiang et al., 2016 and Sang et al., 2022). A special case is the fruit peel under the eggplant calyx, which is always shaded and therefore remains uncolored (either partially or completely) in photosensitive fruit peels compared to non-photosensitive ones, indicating that light is required for the development of purple pigmentation in eggplant (Li et al., 2017; Li et al., 2018 and Zhang et al., 2019). Consequently, breeding for non-photosensitive eggplant varieties is an effective method to maintain a stable anthocyanin accumulation independently of environmental conditions. Given the relevance of the non-photosensitivity trait in eggplant breeding, some studies have focused on mapping the genes responsible for photoinsensitivity. These studies have identified different QTLs associated with non-photosensitivity across different chromosomes, likely due to the use of diverse eggplant materials, research backgrounds and eggplant reference genomes (Qiao et al., 2022; He et al., 2022; Mangino et al., 2022; Luo et al., 2023a-2023b and You et al., 2023). Several genes are involved in the regulation of anthocyanin biosynthesis in eggplant where the structural genes in the anthocyanin biosynthetic pathway include enzyme-encoding genes such as *CHS*, *CHI*, *F3H*, *F3'H*, *F3'5'H*, *DFR* and *ANS* (Jaakola et al., 2013 and Liu et al., 2018).

Those genes are regulated by transcription factors (TFs), several of which have been postulated as

candidates to regulate anthocyanin gene expression (Li et al., 2017; Li et al., 2018; Qiao et al., 2022;

He et al., 2022; Mangino et al., 2022; Zhou et al., 2022; Li et al., 2023; Luo et al., 2023a-2023b You et

al., 2023). A common feature of the candidate TFs is that they activate R2R3-MYB, basic helix-loop-

helix (bHLH) or WD repeat proteins (WDR), which form the MYB-bHLH-WDR (MBW) complex to

regulate anthocyanins structural biosynthesis genes (Ramsay et al., 2005; Yan et al., 2021). Researching

features connected to anthocyanins fruit peel pigmentation can help us comprehend this characteristic

of eggplants and create new kinds of eggplants with varied anthocyanin coloration.

In this study, we investigated the photosensitivity trait in local eggplant varieties from País Valencià.

To map non-photosensitivity trait, we evaluated an F2 population derived from a cross between a

photosensitive cultivar (*S. melongena* ASI-S-1) and a non-photosensitive variety (*S. melongena* IVIA-

371). This population also segregates for anthocyanin and clorophyll pigmentation. Bulked segregant

analysis by sequencing (BSA-seq), combined with RNA-seq, was applied to identify candidate loci,

SmNAC-like protein, for photosensitivity (PS) in the F2. The presence of DNA mutations in the locus

identified was in agreement with the genotype of non-photosensitive phenotypes (e.g., IVIA-371 and

DH-ECAVI) and photosensitive phenotypes (e.g., AN-S-26, H15, and ASI-S-1). Two different

mutations in the down-regulated *SmNac1-like* gene, a transcription factor, were found in IVIA-371 and

DH-ECAVI but not in the photosensitive accessions. The identification of *SmNac1-like* as a candidate

gene for the eggplant photosensitive trait is useful to understand the genetic basis of the regulation of

the anthocyanin pigmentation in the eggplant fruit peel, as well as for eggplant breeding.

MATERIAL AND METHODS

1.- Plant materials and cultivation conditions.

The *S. melongena* accessions ASI-S-1 (anthocyanin photosensitive, non-striped and with presence of chlorophyll pigmentation in the peel) and IVIA-371 (anthocyanin non-photosensitive, striped and no chlorophyll pigmentation in the peel) were used as parents to develop an F2 population of 120 individuals segregating for the photosensitivity trait (PS), presence of fruit anthocyanin (FA) and chlorophyll content (CC) (Figure 1 Panels A and B). PS was categorized as PUC (presence of purple pigmentation under the calyx, i.e., negative for PS) or puc (absence of purple pigmentation under the calyx, i.e., positive for PS. In addition to the eggplant accessions ASI-S-1 and IVIA-371, two sub-introgression lines (IL-69(6)5-12-Ch10 and IL-69(6)5-8-Ch10, derived from the introgression line IL-M5-19) plus accessions DH-ECAVI (photoinsensitive-anthocyanin uniform pigmentation), H15 and AN-S-26 (photosensitive- anthocyanins in the fruit peel exposed to sunlight) were used in the current study (Mangino et al., 2022). With regards to the phenotype of IL-69(6)5-12-Ch10 and IL-69(6)5-8-Ch10 showed a complete loss of anthocyanins content in the fruit peel (Figure S1). The introgression line IL-M5-19 introgression line derives from the MEL5 eggplant (photosensitive) with an introgression from the eggplant wild relative *S. insanum* in chromosome 10.

Seeds of all the accessions were germinated in Petri dishes following the Ranil et al. (2015) protocol and transferred to seedling trays in a climatic chamber under a photoperiod 16 h light/ 8 h dark and temperature regime of 22 ± 2 ºC. The plantlets were grown for four weeks (up to the fifth true leaf development stage) in a climatic chamber before acclimatization for 48 hours in a benched glasshouse. Plantlets were then transplanted to 15 L pots filled with coconut fiber and grown in a pollinator-free benched glasshouse of the Universitat Politècnica de València (UPV), Valencia, Spain. Plants were spaced 1.2 m between rows and 1 m within the row, fertirrigated using a drip irrigation system and trained with vertical strings. Pruning was done manually to regulate vegetative growth and flowering. Phytosanitary treatments were performed when necessary.

2.- Generation of a F2 population and BSA-Seq analysis of PUC vs puc bulks.

The F2 population obtained by the intraspecific cross between ASI-S-1 and IVIA-371 accessions was phenotyped for the PS and FA fruit traits. A Chi-square test was performed to assess the goodness-of-

fit to a monogenic dominant inheritance pattern (3:1 segregation model). The F2 population was used for a BSA-Seq for the PS trait. The PS have been scored in 94 plants out of the 120 of the total population, as 26 plants did not display anthocyanin pigmentation. The PUC and puc bulks consisted of 15 F2 individuals. Young leaves were harvested from the entire F2 population (120 F2 individuals) as well as from six individual plants for each of the two parents. Their DNA was isolated, quantified and tested for quality prior to select six plants from each of the parents and 30 individual F2 plants based on their different PS phenotype (PUC vs. puc). Genomic DNA was extracted using the silica matrix (SILEX) protocol (Vilanova et al., 2020) and checked for integrity by agarose electrophoresis. NanoDrop (Thermo Fisher Scientific, Waltham, MA, USA) ratios 260/280 and 260/230 were used to test the DNA quality, while its concentration was estimated with a Qubit 2.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA, USA).

Pools for the PUC and puc phenotypes bulks were made by mixing in a equimolecular from each of selected F2 individual and shipped to Novogene (Novogene Europe, Cambridge, United Kingdom) where genomic libraries (PE 150, insert 350 bp) were constructed and sequenced. All raw reads were trimmed using the fastq-mcf tool from the Ea-utils package (Aronesty, 2013) with "q 30 -l 50" parameters, and the overall quality was checked using FastQC v.0.12.1 (Andrews et al., 2010). Clean reads were mapped against the 67/3 v3 (for QTL comparison and DNA variant analysis in reported eggplant populations with PS trait) and v.4.1 (for data cross-referencing with RNA-seq) eggplant reference genome (Barchi et al., 2019). To map the clean reads BWA-MEM v.0.7.17–r1188 was used with default parameters (Li and Durbin et al., 2009). The ΔSNP-index was estimated using the QTL-seq software v.2.2.3 (Takagi et al., 2013) from BAM files.

3.- RNA-seq analysis of F2 population.

RNA-seq analysis was conducted on bagged fruits from the F2 population to ascertain the gene expression profile of eggplant fruits with and without PS. Eggplant flowers from each plant were manually pollinated and kept in opaque paper bags for five days in the dark. The time lapse was selected, as it was found to be the shortest time for the appearance of anthocyanin pigmentation. Subsequently, a set of eggplant fruit peels was gathered by cutting them with a surgical blade avoiding as much as

possible to collect eggplant flesh, and subsequently frozen in liquid nitrogen before being kept in storage

at -80ºC. The eggplant fruit peels were grouped in three randomized replicates for each phenotype of

PS trait (PUC and puc). For the PUC samples, each replicate consisted of 4 fruit peels. However, for

the puc samples, due to the limitation in the number of fruits available, each replicate consisted of three

samples. The RNA extractions were done with RNAqueous kit and Ambion Plant RNA isolation aid

reagent (Invitrogen™ AM1912 and AM6990) following the manufacturer instructions. The RNA

integrity and quality were checked by 1.5% agarose gel and with the spectrometer Nanodrop 1000

(Thermo Scientific). RNA concentration was measured using a Qubit 2.0 Fluorometer (Invitrogen) and

an AccuBlue ® Broad Range RNA quantitation kit (Biotium). The samples with the highest values of

260/280 and 260/230 ratios were selected, normalized for their concentration and randomly pooled.

RNA samples were shipped to Novogene Europe (Cambridge, UK) for library preparation and

sequencing. Briefly, mRNA was purified from total RNA using poly-T oligo-attached magnetic beads.

Then, mRNA was fragmented and the first strand cDNA was synthesized using random hexamer

primers, followed by the second strand cDNA synthesis. After adapter ligation, fragments were size-

selected, PCR-amplified, and purified. The library was checked with Qubit and real-time PCR for

quantification and bioanalyzer for size distribution detection. Quantified libraries were pooled and

sequenced on Illumina platforms (NovaSeq 6000), according to effective library concentration and data

amount (9 GB of data). With regards to RNA-seq read processing, clean reads were obtained by

removing reads containing adapter, reads containing ploy-N and low-quality reads from raw data

through Novogene in-house perl scripts and Q20, Q30 and GC content were calculated. All the

downstream analyses were based on clean, high-quality data (Table S1). Reference genome (*S.*

melongena 67/3 v4.1) and gene model annotation files were downloaded from Sol Genomics Network

(SGN) website. Index of the reference genome was built and paired-end clean reads were aligned to the

reference genome using Hisat2 v2.0.5 (Zhang et al., 2021). This tool was selected for mapping as it can

generate a database of splice junctions based on the gene model annotation file, giving a better mapping

result than other non-splice mapping tools. Reads numbers mapped to each gene were counted using

featureCounts v1.5.0-p3 (Liao et al., 2014). Then, FPKM was calculated based on the length of each

gene and reads mapped to it.

Differential expression analysis was performed using the DESeq2 R package (1.20.0), which uses a

statistical model based on the negative binomial distribution for determining differential expression

(Love et al., 2014). The resulting P-values were adjusted using the Benjamini and Hochberg's approach

for controlling false discovery rate (Benjamini and Hochberg., 1995). Genes with an adjusted P-value

≤ 0.05 were assigned as differentially expressed. Finally, Gene Ontology (GO) enrichment analysis of

differentially expressed genes (DEG) was implemented by the clusterProfiler R package (Yu et al.,

2012; Wu et al., 2021; Xu et al., 2024), which corrects gene length bias. GO terms with corrected P-

value < 0.05 were considered significantly enriched by differentially expressed genes. The

clusterProfiler R package was used to test the statistical enrichment of differential expression genes in

KEGG pathways database (http://www.genome.jp/kegg/).

4. Evaluation of DNA sequence variants in photo-sensitive and photo-insensitive cultivars

from different eggplant population.

To determine the presence of DNA sequence variants in the DEGs located inside the QTL identified at

chromosome 10 at photo-in/sensitive eggplant cultivars analyzed against v4.1 eggplant reference

genome (Barchi et al., 2019), firstly the genomic DNA chromosome 10 (63.4 – 67.3 Mb) from the

eggplant reference genome "67/3" v4.1 versus v3 were compared with D-Genies (Cabanettes, Klopp.,

2018). As a result, no major breaks and inversions were detected. So, the gene synteny remains constant

at the genomic DNA region under consideration. Secondly of all the clean reads of ASI-S-1 and IVIA-

371 obtained in a previously research (Mangino et al., 2022) was mapped against the eggplant reference

genome "67/3" v3 with BWA-MEM and only the uniquely aligned reads were selected. Regarding the

variant calling was carried out again, due that the reported sequence changes reported by genes Mangino

et al. (2022) were not validated by Sanger Sequencing (Table S2). With regard the new variant calling

filters for the PS trait in the F2 interspecific cross between ASI-S-1 and IVIA-371 were done with

FreeBayes (Garrison and Marth., 2012) with the following parameters: --min coverage 10, --limit

coverage 60 amd –min alternate fraction 0.2. The predicted effects of the sequence changes detected

were evaluated with SnpEff (Cingolaniet et al., 2012). The DNA variants that affect the regulatory

sequences (5'-UTR and 3'-UTR) and/or the coding sequences (i.e., frameshift, indels, splicing variants,

etc) for the PS trait in homozygous allelic variants were classified as "high impact". The Integrative

Genomics Viewer (IGV) tool (Robinson et al., 2022) was used to validate the variants detected by the

combination of FreeBayes + SnpEff softwares. To expand the number of eggplants cultivars evaluated,

eggplants with PS trait (DH-ECAVI, ANS-S-26 and H15) previously reported (Mangino et al., 2022)

were evaluated. As previously indicated clean reads of the respective eggplants accessions was mapped

against the eggplant reference genome "67/3" v3 with BWA-MEM and only the uniquely aligned reads

were selected. Then the variant calling were done with FreeBayes (Garrison and Marth., 2012) with the

following parameters: --min coverage 10, --limit coverage 60 amd –min alternate fraction 0.2. The

predicted effects of the sequence changes detected were evaluated with SnpEff (Cingolaniet et al.,

2012). "High impact" variants were selected and validated with IGV

RESULTS

1.- Identification and inheritance of a QTL linked to photoinsensitivity.

1.1.- Generation and inheritance of the scored traits in the F2 population.

To determine the PS trait segregation pattern, an F2 population was produced by mating the *S.*

melongena parents ASI-S-1 (puc, non-striped, and with presence of chlorophyll pigmentation in the

peel) with IVIA-371 (PUC, striped, and with absence of chlorophyll pigmentation in the peel).

Regarding the presence of plant anthocyanin pigmentation in the parents, ASI-S-1 displayed

anthocyanin in leaves and stem, meanwhile IVIA-371 only showed anthocyanin content in the stem but

not in the leaves (Figure 1 Panel A). Regarding the F1, it displayed the PUC phenotype and FA

phenotype with uniform (non-striped) anthocyanin pigmentation and also showed background

chlorophyll presence in the peel. In the F2 generation, the PS, FA and CC traits segregated resulting in

several phenotypic combinations (Figure 1 Panel B) for the three traits present in the eggplant cultivars

used in this study.

The segregation pattern of the PS trait showed an autosomal dominant inheritance (Table 1), being

eggplants with PUC and non-striped phenotype the most abundant (Figure 1). Plants producing fruits

without anthocyanin pigmentation (i.e., green or white fruits), were not scored for the PS phenotypes

(as the phenotype could not be determined) (Figure 1 Panel B). These results suggested that the PS trait

in our IVIA-371 x ASI-S-1 population are controlled each by one dominant gene. On the other hand,

although the FA and CC traits segregate in the F2 population were not used in this study as they were

independent traits in relation to the PS trait (data not shown).

1.2.- Identification of the genomic loci linked to PS trait by bulk segregation analysis in

the F2 population

To narrow down the genomic loci linked to the PS trait, two DNA pools from the IVIA-371 x ASI-S-

1 F2 population (15 plants with the PUC phenotype and 15 with the puc phenotype) were sequenced.

This sequencing yielded an average of 154 million 150-bp raw reads per bulk, with a mean coverage

of 33.3x. Using ASI-S-1 as the reference sequence, the ΔSNP-index was calculated for 535,083 SNPs,

getting a QTL on chromosome 10 between 62.9-69.7 Mb with a 99 % confidence interval (CI) for the

PS trait. Similarly, when the ΔSNP-index was calculated for 562,169 SNPs using IVIA-371 as the

reference sequence, a region between 63,4-67,5 Mb was identified on the same chromosome 10 with a

99 % CI (Figure 2, Figure S2 and Table S3). As a result, 102 genes were identified within the

common candidate region (Table S4). Variant calling and prediction of the sequence changes were

evaluated in both candidate regions of ASI-S-1 and IVIA-371, no DNA variants in the coding

sequences (frameshift, indels, splicing variants, etc.) were detected**2.- Genetic evidences of QTl**

for photoinsensitivity in different eggplant populations.

In order to determine the relevance of the QTL on chromosome 10 in the photosensitivity trait, it was

compared with previous QTLs identified in different eggplant populations (Table 2). To be able to

compare with the previous data, the BSA-seq analysis was re-analyzed against the v3 of the eggplant

reference genome. As a result of the re-analysis a QTL in the chromosome 10 for ASI-S-1 between

84,1-87,9 Mb with 99 % CI and a region between 83,5-97,7 Mb with 95 % CI for IVIA-371 were

identified (Table S5, Table S6, Figure S2 and Figure S3). Firstly different sub introgression lines

derived from the IL-M5-19 introgression line (introgressed DNA of Solanum insanum into MEL5

eggplant) were evaluated. As a result, the sub introgression lines (IL-69(6)5-12-Ch10 and IL-69(6)5-8-

Ch10) with introgressed DNA at chromosome 10, IL-69(6)5-12-Ch10 at 84,4-102,8 Mb and IL-69(6)5-

8-Ch10 at 105,8-106,6 Mb respectively (Table 2), showed complete loss of anthocyanins content in the

fruit peel (Figure S1). With regards the S3MEGGIC population for FA (91.1-94.8 Mb) and PUC (91.1-

94.8 and 98.2-100.5 Mb) showed significant SNPs stacked at the end of chromosome 10 (Table 2).

Taking all the QTLs identified in the different populations together, suggest that the end of the

chromosome 10 as a hotspot related with anthocyanin related traits. Six of QTL reported showed

overlap (total or partial) along the different eggplant populations (Table 2). Of these six reported QTLs,

two groups can be observed: the first one consists of four overlapping QTLs (IL-69(6)5-12-Ch10 84,4-

102,8 Mb, FA 91.1-94.8 Mb, PUC 91.1-94.8 Mb, ASI-S-1 84,1-87,9 Mb and IVIA-371 83,5-97,7 Mb),

while the second one consists of two QTLs with partial overlap (IL-69(6)5-12-Ch10 at 84,4-102,8 Mb

and PUC 98.2-100.5 Mb). As a summary two common DNA regions across all the QTLs were identified

(QTLs_ASI-S-1 84.1-87.9 (3.80 Mb) and PUC 98.2-100.5 (2.3 Mb)).

3.- Differential expression analysis of bagged eggplants

The gene expression analysis between PUC and puc bagged eggplant fruits of F2 eggplant population was performed by RNA-seq of the fruit peel. The results indicate that from a total of 177 DEGs, 53 DEGs were up-regulated in the fruit peel of eggplants with PUC phenotype, while 124 DEGs were down-regulated in the fruit peel of eggplants with puc phenotype (Figure 3 and Table S7-9). The 177 DEGs identified from the eggplant fruit peel for the PS trait can be grouped in four different subclusters (Table S9 and Figure S4). The subclusters 1 and 3 are made of the DEGs expressed mainly in IVIA-371 (PUC phenotype) with a total of 53 genes, that constitute the whole up-regulated DEGs identified by RNA-seq. The subclusters 2 and 4 contain 124 genes expressed totally in ASI-S-1 (puc phenotype), and constitute all the down-regulated DEGs reported for the eggplant fruit peel (Table S2 and Table S8). Concerning the functional and gene enrichment analysis of up- and down-regulated genes in eggplant fruit peel with PUC and puc phenotypes, few DEGS were identified due to the fruit peel characteristics. For the up-regulated DEGs expressed in IVIA-371, the gene set showed an enrichment in KEGG terms related to flavonoid and phenylpropanoid biosynthesis (Table S7), mostly located in the subcluster 1 (Table S8) and in agreement with the presence of anthocyanins in the fruit peel of eggplant (Figure 3 and Table S8). The following up-regulated genes in flavonoid biosynthesis were identified in IVIA-371: SMEL4.1_09g021180.1, SMEL4.1_05g002700.1, SMEL4.1_08g022350_1, SMEL4.1_02g020840_1, SMEL4.1_05g002760_1, SMEL4.1_05g018170_1 and SMEL4.1_12g016460_1. Regarding the GO annotation terms, an enrichment in terms like small molecule metabolic process, transferase activity of acyl and hexosyl groups, or coenzyme bonding and iron ion binding was observed. Those functions are consistent with the activity of the up-regulated genes identified by the KEGG terms.

Concerning the down-regulated genes in IVIA-371 and, converserly, over-expressed in ASI-S-1, the gene set showed an enrichment in KEGG terms related to MAPK signaling pathway, phenylpropanoid biosynthesis, cysteine and methionine metabolism, zeatin and alanine biosynthesis, and aspartate and glutamate metabolism (Table S7). The enrichment in those KEGG terms in the MAPK signaling pathway and zeatin biosynthesis are involved in plant growth and fruit enlargement (Schäfer et al., 2015

and Jagodzik et al., 2018) in agreement with the fact that the eggplant fruits were growing. About the

KEGG terms of cysteine and methionine metabolism plus alanine, aspartate and glutamate metabolism

are reported to play a role in the generation of secondary metabolites involved in plant growth, fixing

inorganic sulfur, precursors of several biosynthesis pathways, and plant protection against abiotic stress

(Moormann et al., 2022). Finally, the accumulation of KEGG terms in phenylpropanoid biosynthesis

indicated an accumulation of the biosynthesis precursors of flavonoids (Vogt, 2010; Dong et al., 2021

and Liu et al., 2021). With respect to the GO annotation terms an enrichment in terms like coenzyme

binding, carbohydrate metabolic process, regulation of biological quality, or carbon-oxygen lyase

activity was observed. Those functions are consistent with the activity of the up-regulated genes

identified by the KEGG terms since they are involved in biosynthetic pathways.

4.- Identification of genes expressed in photo-insensitive fruits in the dark conditions,

SmNac1

4.1- Evaluation of the DEGs inside of the candidate region acotated in the F2 population

To narrow down the number of putative candidate genes within the common candiate region of ASI-S-

1 and IVIA-371 (102 genes), their expression levels (Table S9) were evaluated given that no DNA

variant were found in the variant calling plus prediction of the sequence changes (see above – section

1.2). As a result, five genes were differentially expressed (*SMEL4.1_10g016630.1*,

SMEL4.1_10g014730.1, *SMEL4.1_10g013610.1*, *SMEL4.1_10g014530.1*, and

SMEL4.1_10g019230.1) (Table 3). The *SMEL4.1_10g016630.1* codifies for a CHI-like protein,

essential enzyme in the anthocyanin synthesis pathway, is up-regulated. This is in agreement with the

fact that the PUC and puc pools showed anthocyanin content in the fruit peel (Figure 2). With regards

the down-regulated genes (*SMEL4.1_10g014730.1*, *SMEL4.1_10g013610.1*, *SMEL4.1_10g014530.1*,

and *SMEL4.1_10g019230.1)*, *SMEL4.1_10g014730.1* and *SMEL4.1_10g014530.1* codifies for a

NAC1-like protein (a transcription factor) RAD5B-like protein (DNA repair protein) respectively. The

remaining down-regulated genes encode proteins of unknown function (Table 3).

Since the differentially expressed genes did not show mutations in the coding sequences, the regulatory

regions (5'UTR and 3'UTR) were analysed. As a result the NAC-1 like protein (*SMEL4.1_10g014730.1*)

showed an indel in the 3'UTR (Reference: ATATTATTATTATTATTATC, Alternate variant:

ATATTATTATTATT---ATC) (File S1 and FileS2).

328   4.2- Presence of DNA mutation of *SmNac1* in other eggplant accessions photosensible

329   (H15 and ANS-S-26) and photoinsenstible (DH-ECAVI)

With the aim to known of the DNA mutation in the *SMEL4.1_10g014730.1* was present in other

eggplant accessions previously reported in the v3 genome eggplant consortium (Mangino et al., 2022).

Firstly the equivalent of the five genes differentially expressed were located in the v3 of the eggplant

refence genome. All of them were inside of the common DNA region of ASI-S-1 (84,1-87,9 Mb) and

IVIA-371 (83,5-97,7 Mb) (Table S6). Secondly the coding sequences,5'UTR and 3'UTR sequences

were reevaluated in photosensitivy (H15, ANS-S-26 and ASI-S-1) and photoinsensitivity (DH-ECAVI

and IVIA-371) eggplant accessions (Table 4).

As a result, in the photoinsensitivity eggplant accessions only *SMEL4.1_10g014730.1* showed the same

disruptive inframe insertion at 3'UTR in IVIA-371, plus DH-ECAVI showed a disruptive in frame

insertion that causes a frameshift in the coding sequence (Table 4 and File S1 and FileS2). With regards

the photosensitivy (H15, ANS-S-26 and ASI-S-1) eggplant no DNA mutations were detected in

SMEL4.1_10g014730.1 and only for H15 *SMEL4.1_10g014530.1* showed a conservative in frame

insertion that do not change the open reading frame of RAD5B-like protein.

DISCUSSION

Plant domestication in the Solanaceae family resulted in a great diversity of fruit shapes, sizes, and fruit colours. Despite this, modern breeding programs have generally favored eggplants with constant and intense dark and purple pigmentation in the fruit peel (Samuels et al., 2015; Wang et al., 2015 and Taher et al., 2017). In western markets, eggplants with black or purple peel are of higher economic value (Muñoz-Falcón et al., 2009). In our work, different eggplant populations were used to evidence that regarding to antocyanin related traits (complete absence of antocyanin, photsensitivy and fruit anthocyanin) the chromosome 10 is a candidate DNA region to take into account. Anthocyanins have a strong antioxidant effect, so their high content in the fruit peel of eggplant improves nutrition. So, a research to determine the genetics basis of the PS is needed to set the basis of a new generation of eggplants cultivars with high content of secondary metabolites beneficial for human health. In the current research the PS showed an inheritance in agreement with a monogenic dominant gene, following 3:1 ratio in and F2 generation, consistent with previous research data (Honda et al., 2014; He et al., 2022; Qiao et al., 2022; Luo et al., 2023ab). These eggplant accessions were selected because, based on the corrected data reported by the GWAS study of the MAGIC population, neither of the two accessions presented mutations in the CDS of the candidate regions, suggesting that expression must have been altered. Regarding the fruit pigmentation of the F2 generation is probably due to the heterozygosity of the parentals for the genes that controlled photosensitivity, plant anthocyanin, and fruit anthocyanin.

The reported studies on the PS trait used commercial varieties of eggplant (Honda et al., 2012) or non-photosensitive eggplants varieties (He et al., 2022; Qiao et al., 2022; Luo et al., 2023ab) as parental donors for the PUC phenotype. Meanwhile for the puc phenotype present in photosensitive eggplants is recessive compared to the dominant allele of non-photosensitive eggplant varieties (Honda et al., 2012; He et al., 2022; Qiao et al., 2022; Luo et al., 2023a and Luo et al., 2023b). It should be noted that two commercial eggplant varieties with the PUC phenotype (IVIA-371) and (ASI-S-1) were used to produce the F2 population.

As opposite what happens with other fruit colour traits (such as the green fruit neeting (Arrones et al., 2022 and 2024, Toledo Garcia, et al., 2024)), that the phenotype present in the commercial eggplant varieties involves loss of gene function and recessive inheritance, so the PS trait results in a dominant

inheritance and a gain of function. That phenomenon is quite uncommon since domestication usually

involves the loss of gene function or regulation (Lester and Hasan, 1991).

Regarding the localization of the reported QTLs (He et al., 2022; Qiao et al., 2022; Luo et al., 2023a,

Luo et al., 2023b) for the PS at the end chromosome 10 using the eggplant genome 67/3 v3 as a reference

with complete different eggplant accession is in agreement with our results, validating that region as a

promising localization to perform fine-tune mapping or expression analysis to identify gene candidates..

However, other QTLs for the PS trait have been reported in chromosome 1 and 3 (Mangino et al., 2022).

Here, we confirmed the end chromosome 10, using v3 eggplant genome as a reference, as candidate

region related with anthocyanin content traits, identifying 84 potential candidates genes for ASI-S-1

and 550 for IVIA-371. To filter the number of candidates genes identified by the BSA-seq analysis, the

list of genes was compared with the identified DEG from the RNA-seq of F2 population. Out of the

five genes identified as differentially expressed, the *SmNAC1-like* (SMEL4.1_10g014730.1) is mutated

in eggplant varieties with PUC phenotype (DH-ECAVI and IVIA-371) but not in puc phenotype (ANS-

S-26. H15 and ASI-S-1) eggplant varieties. All these results together provide strong evidence of

SmNAC1-like gene as candidate gene controlling the PUC trait in our eggplant varieties.

Regarding gene regulation of the anthocyanin biosynthetic pathway, previous studies suggest that light

and cold induce anthocyanin biosynthesis from the moment of exposure until saturation through a

highly dynamic range of TFs (Jiang et al., 2016; Li et al., 2017; Sang et al., 2022, Li et al., 2023 and

Zhou et al., 2020). Specifically, *SmCry1*, *SmCry2*, *SmCry3* and *SmUVR8* are involved as photoreceptors

in response to light. Meanwhile *SmCBF1*, *SmCBF2* and *SmCBF3* have been reported as sensors in

response to low temperature. Both sets of genes activate the signaling pathway to produce anthocyanins

(Li et al., 2017; Li et al., 2023 and Zhou et al., 2020). Regarding the different downstream stages of the

anthocyanin synthesis pathway, there is consensus on the necessary participation of the pathway's

structural genes: *SmCHI, SmF3H, Sm3GT, Sm5GT, SmDFR* and *SmANS*) (Jaakola et al., 2013 and Liu

et al., 2018. For the intermediate steps, several TFs and their interactions have been proposed in

different eggplant populations (R2R3-MYB transcription factor [*SmMYB113*, *SmMYB86*, *SmMYB75*,

SmMYB35], basic helix–loop–helix [*SmbHLH1*, *SmbHLH13*], *SmHY5*, *SmTT8*, *SmGL3*, *SmTTG1*,

SmCIP7, *SmGL2*, *SmGATA26*, *SmCOP1*, *SmSPA* and *SmFTSH10*) (Jiang et al., 2016; Li et al., 2017;

Li et al., 2018; Qiao et al., 2022; He et al., 2022; Mangino et al., 2022; Zhou et al., 2022; Li et al., 2023;

Luo et al., 2023ab and You et al., 2023). Specifically, any mutation in *SmMYB113* or pathway's

structural genes generate phenotypes without anthocyanin content (Mangino et al., 2022; Luo et al.,

2023ab and You et al., 2023). Therefore, for the PS trait the TFs proposed (*SmHY5*, *SmTT8*, *SmGL3*,

SmTTG1, *SmCIP7*, *SmGL2*, *SmGATA26*) are in the first steps of the signaling pathway and always

upstream of the *SmMYB113* (Mangino et al., 2022, Luo et al., 2023ab and You et al., 2023). In that

sense, *SmNAC1-like* agrees with the previous candidate TFs proposed. The homologous of *SmNAC1-*

like in *Prunus persica* (*PpNAC1*) and *Malus domestica* (*MdNAC1*) are located upstream of the

SmMYB113 homologous in *P. persica* (*PpMYB10.1*) and *M.* domestica (MdMYB10) (Liu et al., 2023

and Meng et al., 2023).

By using different eggplant materials, we have identified *SmNAC1-like* gene as a promising candidate

for the PS trait, with disruptive mutations associated with non-photosensitivity phenotypes. The

eggplant PS trait has significant implications for fruit visual quality, as it results in uniform peel

pigmentation, which is highly desirable for fruit commercialization. The study of this relevant trait

could be of interest for future eggplant breeding projects with the aim of increasing the diversity of the

fruit colour palette. In addition, due to a higher concentration of anthocyanin contents and their effect

as an antioxidant, *SmNAC1-like* could also be a valuable target for controlling the gene expression of

the anthocyanin content in the fruit peel of the eggplants.

ACKNOWLEDGEMENTS

This work has been funded by grants Juan de la Cierva (FJC2020-043194-I) funded by Agencia Estatal de Investigación and APOSTD-GVA (CIAPOS/2022/127) funded by Generalitat Valencia. Funding was also received from grant CIPROM/2021/020 funded by Conselleria d'Educació, Universitats i Ocupació of the Generalitat Valenciana, and from grant PID2021-128148OB-I00 funded by MCIN/AEI/10.13039/501100011033/ and by 'ERDF A way of making Europe'. VB-F and J S-P are grateful to Generalitat Valenciana for grants INVEST/2022/146 and INVEST/2022/150, funded by European Union, Next Generation EU. JL is grateful to Spanish Ministerio de Ciencia, Innovación y Universidades for a predoctoral grant (FPU21/01590). SM would like to thank funding support from UPV and the Spanish Ministerio de Ciencia, Innovación y Universidades under the program María Zambrano funded by the European Union Next Generation-EU. PG is grateful to Spanish Ministerio de Ciencia, Innovación y Universidades for a post-doctoral grant (RYC2021–031.999-I) funded by MCIN/AEI /10.13039/501.100.011.033 and the European Union through NextGenerationEU/PRTR.

CONTRIBUTIONS

JG-C, SM, PG, SV, MP and JP conceived the idea. JG-C and SM supervised and designed the manuscript. JG-C, SM, AA and MDT-T developed the F2 population for BSA-seq and RNA-seq. EG-P developed the introgression lines. JG-C, SM, JL, JS-P, MDT-T performed DNA and RNA extraction. JG-C wrote the manuscript. All the authors reviewed and edited the manuscript.

DATA AVAILABILITY STATEMENT

The data presented in the study are deposited in the NCBI SRA repository, BioProject IDs for BSA-seq are the following: F2 photo-insensitive, SRS26640024, F2 photosensitive SRS26640023, ASI -S-1 SRS26640022 and IVIA-371 SRS26640021. With regards theRNA-

se analysis are under the BioProject IDs: PUCR replicate 1 SRR35685429, PUCR replicate 2 SRR35685426, PUC replicate 3 SRR35685424, puc replicate 1 SRR35685427, puc replicate 2 SRR35685425, puc replicate 3 SRR35685428.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

GENERATIVE AI STATEMENT

The authors declare that no Gen AI was used in the creation if this manuscript or in any of the figure and table attached to the manuscript.

SUPPLEMENTARY MATERIALS

File S1. DNA sequence modification in SMEL4.1_10g014730.1 in 67/3 v3, DH-ECAVI and IVIA-371.

File S2. cDNA and protein sequence of SMEL4.1_10g014730.1 in 67/3 v3 and DH-ECAVI.

Table S1. RNA quality data of the samples used in the RNA-seq data of PUC and puc.

Table S2. Evaluation of the gene related with anthocyanin biosynthesis in the QTLs determined by Mangino et al., 2022.

Table S3. Results of bulk segregation analysis in ASI-S-1 and IVIA-371 with a confidence interval of 95% and 99%, respectively. Eggplant genome reference v4.1

Table S4. List of genes in the candidate region acotated by BSA-seq in eggplant varieties ASI-S-1 and IVIA-371. Eggplant genome reference v4.1

Table S5. Results of bulk segregation analysis in ASI-S-1 and IVIA-371 with a confidence interval of 95% and 99%, respectively. Eggplant genome reference v3

Table S6. List of genes in the candidate region acotated by BSA-seq in eggplant varieties ASI-S-1 and IVIA-371. Eggplant genome reference v3

Table S7. List of differentially expressed genes in the eggplant varieties IVIA-371 (PUC

positive trait) versus ASI-S-1 (puc negative trait).

Table S8. List of the subcluster identified in the heatmap of the differentially expressed genes

of light-independent versus light-dependent eggplant varieties.

Table S9. List of the enrichment analysis of the GO and KEGG terms of the up- and down-

regulated genes in PUC versus puc eggplant fruit peel

Figure S1. Phenotype of the two sub-introgression lines (IL-69(6)5-12-Ch10 and IL-69(6)5-8-Ch10)

derived from the introgression line IL-M5-19.

Figure S2. Delta of SNP-Index of the bulk segregation analysis of light-dependent and light-

independent eggplant varieties

Figure S3. Number of differentially expressed genes (DEG) in the four subclusters identified

in the clustering heatmap PUC vs puc. The DEG in the subcluster 1 and 3 were up-regulated

in the light-independent (IVIA-371) eggplant (Panel A and B) meanwhile the DEG in the

subcluster 2 and 4 were up-regulated in the light-dependent (ASI-S-1) eggplant.

REFERENCES

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Available online at: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Aronesty E. 2013. Comparison of sequencing utility programs. The Open Bioinformatics Journal 7, doi: 10.2174/1875036201307010001.

Arrones A, Manrique S, Gomis-Cebolla J, Baraja-Fonseca V, Plazas M, Prohens J, Portis E, Barchi L, Giuliano G, Gramazio P, Vilanova S. 2024. Irregular green netting of eggplant fruit peel: a domestication trait controlled by *SmGLK2* with potential for fruit colour diversification, Journal of Experimental Botany, Volume 75, Issue 22, 4. Pages 7066–7078

Arrones A, Mangino G, Alonso D, Plazas M, Prohens J, Portis E, Barchi L, Giuliano G, Vilanova S, Gramazio P. 2022. Mutations in the SmAPRR2 transcription factor suppressing chlorophyll pigmentation in the eggplant fruit peel are key drivers of a diversified colour palette. Frontiers in Plant Science 13, 1025951.

Barchi L, Pietrella M, Venturini L, et al. 2019. A chromosome-anchored eggplant genome sequence reveals key events in Solanaceae evolution. Scientific Reports 9, 11769.

Benjamini Y and Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Vol 57 Issue 1 Pages 289-300

Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6(2):80-92. doi: 10.4161/fly.19695.

Cabanettes F, Klopp C. (2018) D-GENIES: dot plot large genomes in an interactive, efficient and simple way. PeerJ 6:e4958 https://doi.org/10.7717/peerj.4958

Dong NQ, Lin HX. 2021. Contribution of phenylpropanoid metabolism to plant development

and plant-environment interactions. J Integr Plant Biol. 63 (1):180-209.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read

sequencing.arXiv preprint arXiv:1207.3907 [q-bio.GN]

He Y, Li S, Dong Y, Zhang X, Li D, Liu Y, Chen H. 2022. Fine mapping and characterization

of the dominant gene SmFTSH10 conferring non-photosensitivity in eggplant (Solanum

melongena L.). Theor Appl Genet. 135(7):2187-2196.

Jaakola L. 2013. New insights into the regulation of anthocyanin biosynthesis in fruits. Trends

Plant Sci. 18(9):477-83.

Jagodzik P, Tajdel-Zielinska M, Ciesla A, Marczak M, Ludwikow A. 2018. Mitogen-Activated

Protein Kinase Cascades in Plant Hormone Signaling. Front Plant Sci. 8; 9:1387.

Jiang M, Ren L, Lian H et al (2016) Novel insight into the mechanism underlying light-

controlled anthocyanin accumulation in eggplant (*Solanum melongena* L.). Plant Sci 249:46–

58

Junren Meng, Shihang Sun, Ang Li, Lei Pan, Wenyi Duan, Guochao Cui, Juan Xu, Liang Niu,

Zhiqiang Wang, Wenfang Zeng. 2023. A NAC transcription factor, PpNAC1, regulates the

expression of *PpMYB10.1* to promote anthocyanin biosynthesis in the leaves of peach trees in

autumn. *Horticulture Advances*, 2023, 1(1): 8.

Lestern-Hasan, S.M.Z. (1991), «Origin and Domestication of the Brinjal Eggplant, Solanum

melongena, from S. incanum, in Africa and Asia», en haWKes, J.G. et alii (eds.), Solanaceae

III, Taxonomy, Chemistry, Evolution, Londres, Royal Botanic Gardens, pp.369-388

Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics. 2014 Apr 1;30(7):923-30. doi:

10.1093/bioinformatics/btt656.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. Bioinformatics 25, 1754–1760.

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv, 1303.3997 [q-bio.GN].

Li J, Ren L, Gao Z et al (2017) Combined transcriptomic and proteomic analysis constructs a new model for light-induced anthocyanin biosynthesis in eggplant (*Solanum melongena* L.). Plant Cell Environ 40:3069–3087

Li J, He YJ, Zhou L, Liu Y, Jiang M, Ren L, Chen H. Transcriptome profiling of genes related to light-induced anthocyanin biosynthesis in eggplant (Solanum melongena L.) before purple color becomes evident. BMC Genomics. 2018 May 4;19(1):324. doi: 10.1186/s12864-018-4693-y.

Li J, Jiang S, Yang G, Xu Y, Li L, Yang F. 2023. RNA-sequencing analysis reveals novel genes involved in the different peel color formation in eggplant. Hortic Res.10 (10):181.

Liu M, Li X, Liu Y, Cao B. 2013. Regulation of flavanone 3-hydroxylase gene involved in the flavonoid biosynthesis pathway in response to UV-B radiation and drought stress in the desert plant, Reaumuria soongorica. Plant Physiol Biochem 73:161–167

Liu Y, Tikunov Y, Schouten RE, Marcelis LFM, Visser RGF, Bovy A. 2018. Anthocyanin Biosynthesis and Degradation Mechanisms in *Solanaceous* Vegetables: A Review. Front Chem. 6:52.

Liu W, Feng Y, Yu S, Fan Z, Li X, Li J, Yin H. 2021. The Flavonoid Biosynthesis Network in Plants. Int J Mol Sci. 22(23):12824.

Liu W, Mei Z, Yu L, Gu T, Li Z, Zou Q, Zhang S, Fang H, Wang Y, Zhang Z, Chen X, Wang N. 2023. The ABA-induced NAC transcription factor MdNAC1 interacts with a bZIP-type transcription factor to promote anthocyanin synthesis in red-fleshed apples. Hortic Res 10(5):uhad049.

Lorenc-Kukuła K, Jafra S, Oszmiański J, Szopa J (2005) Ectopic expression of anthocyanin 5-O-glucosyltransferase in potato tuber causes increased resistance to bacteria. J Agric Food Chem 53:272–281

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8.

Luo L, Molthoff J, Li Q, Liu Y, Luo S, Li N, Xuan S, Wang Y, Shen S, Bovy AG, Zhao J, Chen X. Identification of candidate genes associated with less-photosensitive anthocyanin phenotype using an EMS mutant (*pind*) in eggplant (*Solanum melongena* L.). Front Plant Sci. 2023 Dec 11;14:1282661. doi: 10.3389/fpls.2023.1282661. PMID: 38169942; PMCID: PMC10758619.

Mangino G, Arrones A, Plazas M, Pook T, Prohens J, Gramazio P, Vilanova S. 2022. Newly Developed MAGIC Population Allows Identification of Strong Associations and Candidate Genes for Anthocyanin Pigmentation in Eggplant. Front Plant Sci. 7; 13:847789.

Mattioli R, Francioso A, Mosca L, Silva P.2020. Anthocyanins: A Comprehensive Review of Their Chemical Properties and Health Effects on Cardiovascular and Neurodegenerative Diseases. Molecules. 21; 25(17):3809.

Muñoz-Falcón J.E; Prohens J; Vilanova S; Nuez F. (2009). Diversity in commercial varieties and landraces of black eggplants and implications for broadening the breeders' gene pool, 154(3), 453–465. doi:10.1111/j.1744-7348.2009.00314.x

Moormann J, Heinemann B, Hildebrandt TM. 2022. News about amino acid metabolism in plant-microbe interactions. Trends Biochem Sci. 47(10):839-850.

Qiao Jun, Liu Jing, LI Suwen, Wang Liying. 2022. Prediction of Fruit Color Genes Under the Calyx of Eggplant Based on Genome-wide Resequencing in an Extreme Mixing Pool. Acta Horticulturae Sinica, 2022, 49(3): 613-621.

Ramsay NA, Glover BJ. 2005. MYB-bHLH-WD40 protein complex and the evolution of cellular diversity. Trends Plant Sci. 10 (2):63-70.

Ranil RHG, Niran HML, Plazas M, Fonseka RM, Fonseka HH, Vilanova S, Andújar I, Gramazio P, Fita A, Prohens J. 2015. Improving seed germination of the eggplant rootstock Solanum torvum by testing multiple factors using an orthogonal array design. Scientia Horticulturae 193, 174–181.

Ren L, Hu Z, Li Y et al 2014. Heterologous expression of bopap1 in tomato induces stamen specific anthocyanin accumulation and enhances tolerance to a long-term low temperature stress. J Plant Growth Regul 33:757–768

Robinson J.T, Thorvaldsdóttir H, Turner D, Mesirov J.P. 2023. igv.js: an embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV). Bioinformatics, 39(1) (2023), btac830.

Samuels, J. (2015). Biodiversity of Food Species of the Solanaceae Family: A Preliminary Taxonomic Inventory of Subfamily Solanoideae. *Resources*, *4*(2), 277-322.

Sang Z, Zuo J, Wang Q, Fu A, Zheng Y, Ge Y, Qian Z, Cui Y. Determining the Effects of Light on the Fruit Peel Quality of Photosensitive and Nonphotosensitive Eggplant. Plants (Basel). 2022 Aug 12;11(16):2095. doi: 10.3390/plants11162095. PMID: 36015399; PMCID: PMC9416183.

Schäfer M, Brütting C, Meza-Canales ID, Großkinsky DK, Vankova R, Baldwin IT, Meldau S. 2015. The role of cis-zeatin-type cytokinins in plant growth regulation and mediating responses to environmental interactions. J Exp Bot. 66(16):4873-84.

Speer H, D'Cunha NM, Alexopoulos NI, McKune AJ, Naumovski N. Anthocyanins and Human Health-A Focus on Oxidative Stress, Inflammation and Disease. Antioxidants (Basel). 2020 Apr 28;9(5):366. doi: 10.3390/antiox9050366. PMID: 32353990; PMCID: PMC7278778.

Taher D, Solberg SØ, Prohens J, Chou YY, Rakha M, Wu TH. World Vegetable Center

Eggplant Collection: Origin, Composition, Seed Dissemination and Utilization in Breeding.

Front Plant Sci. 2017 Aug 25;8:1484. doi: 10.3389/fpls.2017.01484. PMID: 28970840;

PMCID: PMC5609569.

Takagi H, Abe A, Yoshida K, et al. 2013. QTL-seq: Rapid mapping of quantitative trait loci in

rice by whole genome resequencing of DNA from two bulked populations. The Plant Journal

74, 174–183.

Takos AM, Jaffé FW, Jacob SR et al (2006) Light-induced expression of a myb gene regulates

anthocyanin biosynthesis in red apples. Plant Physiol 142:1216–1232

Toledo Garcia, MD. (2024). Regulación genética de la coloración del fruto de berenjena.

Universitat Politècnica de València. https://riunet.upv.es/handle/10251/207776

Xu S, Hu E, Cai Y, Xie Z, Luo X, Zhan L, Tang W, Wang Q, Liu B, Wang R, Xie W, Wu T,

Xie L, Yu G. Using clusterProfiler to characterize multiomics data. Nat Protoc. 2024

Nov;19(11):3292-3320. doi: 10.1038/s41596-024-01020-z.

Vogt T. Phenylpropanoid biosynthesis. Mol Plant. 2010 Jan;3(1):2-20. doi:

10.1093/mp/ssp106. Epub 2009 Dec 24. PMID: 20035037.

Wang L, Li J, Zhao J, He C. Evolutionary developmental genetics of fruit morphological

variation within the Solanaceae. Front Plant Sci. 2015 Apr 13;6:248. doi:

10.3389/fpls.2015.00248. PMID: 25918515; PMCID: PMC4394660.

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Liu S, Bo

X, Yu G. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innovation (Camb). 2021 Jul 1;2(3):100141. doi: 10.1016/j.xinn.2021.100141.

Yan H, Pei X, Zhang H, Li X, Zhang X, Zhao M, Chiang VL, Sederoff RR, Zhao X. 2021

MYB-Mediated Regulation of Anthocyanin Biosynthesis. Int J Mol Sci. 18;22(6):3103.

You Q, Li H, Wu J, Li T, Wang Y, Sun G, Li Z, Sun B. 2022. Mapping and validation of the

epistatic *D* and *P* genes controlling anthocyanin biosynthesis in the peel of eggplant (*Solanum*

melongena L.) fruit. Hortic Res. 2;10(2)

Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological

themes among gene clusters. OMICS. 2012 May;16(5):284-7. doi: 10.1089/omi.2011.0118.

Zhang Ying, Zhao Yueqi, Chen Yuhui, Lian Yong, and Liu Fuzhong *. Progress in

Molecular Research on Purple Formation of Eggplant[J]. ACTA HORTICULTURAE

SINICA, 2019, 46(9): 1779-1796.

Zhang Y, Park C, Bennett C, Thornton M, Kim D. Rapid and accurate alignment of nucleotide

conversion sequencing reads with HISAT-3N. Genome Res. 2021 Jul;31(7):1290-1295. doi:

10.1101/gr.275193.120.

Zhou X, Liu S, Yang Y, Liu J, Zhuang Y. 2022. Integrated Metabolome and Transcriptome

Analysis Reveals a Regulatory Network of Fruit Peel Pigmentation in Eggplant (*Solanum*

melongena L.). Int J Mol Sci. 23(21):13475.

Zhou L, He Y, Li J, Liu Y, Chen H. 2020. CBFs Function in Anthocyanin Biosynthesis by

Interacting with MYB113 in Eggplant (Solanum melongena L.). Plant Cell Physiol. 61(2):416-

426. doi: 10.1093/pcp/pcz209. PMID: 31725150

TABLES

Table 1. Genetic analysis of the segregation of photosensitivity in the F2 population from the

cross of ASI-S-1 x IVIA-371. For photosensitivity, only the plants yielding fruits with

anthocyanic pigmentation were used.

Cross	Population	Photosensitivity		Total	Expected segregation	Chi-square	p value
		Presence	Absence				
[ASI-S-1xIVIA371] self-crossed	F2	76	18	94	3:1	1.58	0.20

651 **Table 2**. QTLs related with anthocyanin biosynthesis (fruit anthocyanin and purple under calyx traits) in different eggplant materials used in the

652 current research and in Mangino et al. (2022), which used as parents of a MAGIC population some of the accessions used in this study.

Type of material	Trait	Accession	Genome version	Chromosome	QTL anthocyanin pigmentation (Mb)		Size (Mb)	Reported
					Start	End		
Introgression Line	Purple under calyx and fruit anthocyanin	IL-69(6)5-12-Ch10	v3	10	84.38*	102.79*	18.41	Current study
Introgression Line	Purple under calyx and fruit anthocyanin	IL-69(6)5-8-Ch10	v3	10	105.85	106.64	0.79	Current study
MAGIC population	Fruit anthocyanin	S3MEGGIC population	v3	1	5.11	5.88	0.77	Mangino et al., 2022
				9	16.2	17.4	1.20	
				10	91.08*	94.81*	3.73	
	Purple under calyx	S3MEGGIC population	v3	1	5.11	5.88	0.77	
				3	7.22	8.65	1.43	
					31.54	40.09	8.55	
				10	2.08	2.67	0.59	
					3.94	4.94	1.00	
					91.08*	94.81*	3.73	
					98.25*	100.55*	2.30	
F2 ASI-S-1 x IVIA-371	Purple under calyx	ASI-S-1 IVIA-371	v3	10	84.1*	87.9*	3.80	Current study
					83.5*	97.7*	14.20	

653 * The QTLs of the IL-69(6)5-12-Ch10, S3MEGGIC population for fruit anthocyanin and Purple under calyx overlap with the candidate regions obtained by

654 BSA-seq in the eggplant accessions IVIA-371 and ASI-S-1 as a reference.

656 **Table 3**. List of differentially expressed genes in the QTL of chromosome 10 delimited by bulk segregation analysis in the eggplant varieties

657 IVIA-371 (photoinsensitive) and ASI-S-1 (photosensitive).

Regulation	RNA-seq data			Genome version	Gene description
	Gene Name	log2FoldChange	p value		
UP	*SMEL4.1_10g016630.1*	577,158,472,269,854	6.03E-09	v4.1	CHI-like. Chalcone flavonone isomerase of Nicotiana tabacum
DOWN	*SMEL4.1_10g014730.1*	-8,997,230,728,622	1.40E+04	v4.1	NAC1-like. NAC domain-containing protein 1 of Solanum lycopersicum
	SMEL4.1_10g013610.1	-231,064,107,425,366	7.57E+09	v4.1	Protein of unknown function
	SMEL4.1_10g014530.1	-128,031,116,712,945	0.00011	v4.1	RAD5B-like. DNA repair protein RAD5B of Arabidopsis thaliana
	SMEL4.1_10g019230.1	-114,443,348,682,244	0.00031	v4.1	Protein of unknown function

659   **Table 4.** DNA high-impact sequence variants present in the differentially expressed genes located in the QTL of chromosome 10 delimited by

660   bulk segregation analysis in the photoinsensitive and photosensitive eggplant varieties used in this study

Gene Name*	Gene position†		PUC			*puc*	
	Start	End	DH ECAVI	IVIA-371	AN-S-26	H15	ASI-S-1
SMEL4.1_10g016630.1 *(SMEL_010g353630.1)*	96300106	96301916	-	-	-	-	-
SMEL4.1_10g014730.1 **(SMEL_010g350010.1)**	88316013	88317850	Position: 88,316,597-602 Reference: AAAGGT Alternate: AGATCGGAAGAGCGTCGT Type: INDEL; Disruptive inframe insertion	Position: 88,317,680-699 Reference: ATATTATTATTATTATTATC Alternate: ATATTATTATTATTATC Type: INDEL; 3_prime_UTR_variant	-	-	-
SMEL4.1_10g013610.1 *(SMEL_010g349620.1)*	87113226	87117639	-	-	-	-	-
SMEL4.1_10g014530.1 *(SMEL_010g349810.1)*	88067120	88069892	-	-	-	Position: 88,069,753-756 Reference: TAGG Alternate: AGATCGGAAGAGC Type: INDEL; Conservative_inframe_insertion	-
SMEL4.1_10g019230.1 *(SMEL_010g351900.1)*	92381275	92381400	-	-	-	-	-

661   *The gene names between brackets belong to the genome version v3 of *Solanum melongena* from Eggplant Sequencing Consortium

662   † The gene position of the DEG located in the QTL of chromosome 10 belong to the genome version v3 of *Solanum melongena* from Eggplant

663   Sequencing Consortium

FIGURES

Figure 1. Phenotype of the *Solanum melongena* varieties photosensitive (ASI-S-1) and photoinsensitive (IVIA-371) and F2 population. Panel A: Phenotype of the eggplant fruit of ASI-S-1 and IVIA-371. Panel B: Crosses performed to develop the F2 population and the different F2 phenotypes obtained for fruit anthocyanin (FA), purple under calyx (PUC) and chlorophyll content (CC).

Figure 2. Results of the bulk segregation analysis in the F2 population of IVIA-371 (photoinsensitive) x ASI-S-1 (photosensitive). Panel A: Phenotype of the eggplants used to generate the bulks. Panel B: ΔSNP-index and the position of the respective QTL for IVIA-371 and ASI-S-1, respectively.

Figure 3. Differentially expressed genes (DEG) in early bagged photoinsensitive (IVIA-371)
versus photosensitive (ASI-S-1) eggplants. Panel A: Vulcano plot of the up- and down-
regulated of the fruit peel in early developed bagged eggplants. Panel B: Enrichment analysis
of the up- down-regulated terms in GO and KEGG terms. Panel C: Clustering heatmap among
the PUC vs puc and the DEG expressed in the eggplant fruit peel.