System-Level Computations Underlie Visual Field Heterogeneity

Shutian Xue*†[1], Antoine Barbot†[1], Jared Abrams[2], Qingyuan Chen[1] & Marisa Carrasco[1,3]

Affiliations
[1] Department of Psychology, New York University, New York, United States
[2] Applied Research Laboratories, University of Texas at Austin, United States
[3] Center for Neural Science, New York University, New York, United States
* Equal contribution

*Correspondence: Shutian Xue (shutian.xue@nyu.edu)
†These authors contributed equally to this manuscript.

Number of figures: 6/8
Number of tables: 0
Number of supplemental figures: 0
Number of supplemental tables: 0
Number of references: 80/50 (recommended)
Number of words: Total: 4,600/4,500; Abstract = 149/150; Introduction = 990; Results = 2,130; Discussion = 1,560; Online Methods = 3,100.

Keywords
visual field; eccentricity effect; polar angle asymmetries; system-level computations; internal noise; gain; equivalent noise; perceptual template model

Acknowledgment
 We thank David Tu and Zhong-Lin Lu for their helpful comments.

Funding
This research is funded by NIH R01-EY027401 to M.C., Vision Training Grant 5T32EY007136-30 to NYU, and NIH F31-EY036732 to S.X.

Conflict of Interests: The authors declare no competing financial interests.

Author Contributions: S.X., A.B., J.A. and M.C. conceived and designed the study. A.B., J.A. and S.X. wrote the experimental code. J.A. collected preliminary data. S.X., A.B. and Q.C. collected and analyzed
the experimental data. S.X. drafted the manuscript. M.C. administered and supervised the project; provided extensive editorial input and critical revisions throughout the writing

47   process. All authors provided feedback, contributed to editing and approved the final version
48   of the manuscript.
49

Abstract

Visual perception varies with eccentricity and polar angle. We investigated whether and how
system-level computations, which transform visual input into perception, underlie these
heterogeneities. Using the equivalent noise method and perceptual template model, we
estimated gain, internal noise, and nonlinearity for orientation discrimination across
eccentricity (fovea, parafovea and perifovea) and around polar angle. Performance declined
with eccentricity due to decreased gain and nonlinearity and increased internal noise.
Observers with stronger eccentricity effects showed greater gain decrease. Only gain varied
with polar angle—higher along the horizontal than vertical meridian, and lower than upper
vertical meridian—paralleling performance asymmetries. This dissociation aligns with known
variations in neuronal count and tuning, suggesting that neural correlations and neural noise
contribute to these system-level computations. By revealing distinct system-level
computations underlying the eccentricity effect and polar angle asymmetries, our findings
provide a link between perceptual heterogeneity across the visual field and neural
architecture.

Significance Statement

Human visual performance varies throughout the visual field, across eccentricity (distance
from gaze) and around polar angle (circular dimension). Optical and retinal factors as well as
cortical surface area increasingly explain these variations, but only partially. Here, we show
that visual field heterogeneity arises from distinct system-level computations that vary across
eccentricity and polar angle. Eccentricity effects stem from both reduced gain and
nonlinearity and elevated internal noise, whereas polar angle asymmetries arise solely from
gain variations. These findings provide a computational link between behavior and its neural
bases and highlight the importance of considering both eccentricity and polar angle in
modeling human perception. Moreover, they have translational implications for rehabilitation
protocols for visual deficits and ergonomic interfaces.

Introduction

Visual performance declines with eccentricity[1–3] and varies around polar angle at isoeccentric locations. Performance is better along the horizontal than vertical meridian—horizontal-vertical anisotropy (HVA)—and along the lower than upper vertical meridian—vertical meridian asymmetry (VMA) (**Fig. 1A**; review[3]). These polar angle asymmetries are ubiquitous across many visual functions, including early vision, such as contrast sensitivity (e.g[4–8]) and acuity (e.g[8,9]), in mid-level vision, such as texture segmentation[10] and crowding [11], and in high-level vision, such as visual short-term memory[12] and face perception[13].

Figure 1. Schematics of key concepts. **(A)** A schematic of visual field heterogeneity in performance. The size of the dot indicates performance (bigger dots = better performance). From left to right we illustrate the eccentricity effect (fovea is better than farther eccentricities), horizontal-vertical anisotropy (HVA, the horizontal is better than the vertical meridian), and vertical-meridian asymmetry (VMA, the lower is better than the upper vertical meridian). **(B)** The equivalent noise method infers neural computations by mapping performance (e.g., contrast threshold) as a function of external noise and observing how performance is altered by both external (dark green) and internal (orange) noise. The performance decrease with higher external noise levels is captured by the threshold-vs-noise (TvN) curve. **(C)** The Perceptual Template Model (PTM) includes three system-level computations—gain, additive internal noise and nonlinearity. A noisy stimulus is filtered by a perceptual template, and its contrast is scaled by a gain parameter, transformed nonlinearly, and corrupted by internal noise before decision-making. **(D)** Proposed computations influence performance in distinct ways. Differences in gain scale the entire curve without altering its shape, reflecting proportional changes in how the system filters both signal and noise; Changes in internal noise primarily affect thresholds at low external noise levels, where internal variability dominates. As external noise increases, performance converges across locations. Nonlinearity determines the curvature of the function: higher nonlinearity results in a more curved (nonlinear) function.

Both the eccentricity effect and polar angle asymmetries are related to better optical quality[14], higher density of cones[15,16] and midget retinal ganglion cells[17,18] and larger surface area in primary visual cortex (V1) (eccentricity[19–24]; HVA/VMA[21–23,25]). Importantly, V1 surface

area correlates with contrast sensitivity around polar angle[25], indicating a link between
cortical anatomy and visual perception at the level of individual observer.

However, retinal and cortical factors are insufficient to account for polar angle asymmetries.
An encoding model simulating orientation discrimination based on spatial variations in both
optical and retinal factors explained only 10% of VMA and 40% of HVA performance[26,27].
Furthermore, M-scaling the stimulus size as per a well-established cortical magnification
factor eliminates contrast sensitivity differences across eccentricity (e.g[6,28]), but not around
polar angle[6].

Here, we investigated whether system-level computations reflecting the pooled activity of
neuronal populations activated by the stimulated visual field play a critical role. We examined
three computations that limit visual performance in a noisy environment: gain —the system's
ability to amplify the signal[29–31] (or modeled as efficiency[32]), nonlinearity—how the system
transforms inputs in a nonlinear fashion[33,34], and internal noise—the variability intrinsic to the
visual system, independent of external stimulation[33–35]. Both gain and nonlinearity are
template properties reflecting the system's ability to extract task-relevant information from
stimuli. We used the equivalent noise method to probe these system-level computations by
introducing external noise (e.g[33–38]) and infer the three computations by observing how
performance changes (**Fig. 1B**). We constructed threshold-versus-noise curves at different
locations to capture how performance is jointly influenced by external noise and system-level
computations.

The perceptual template model quantifies the three system-level computations (**Fig. 1C**). We
used a simplified version of the original model developed by Lu and Dosher (e.g[33,34,38]),
which retains its explanatory power. First, the noisy stimulus is filtered through a perceptual
template tuned to the signal's orientation and spatial frequency (SF). This step is scaled by
the gain parameter, such that better template-signal alignment yields higher gain. Then, the
template-matched output is passed through a nonlinear transformation, typically a power
function. Next, this transformed signal is corrupted by additive internal noise. Finally, a
decision is made based on the noisy signal.

These neural computations have been linked to neurophysiological properties. Higher gain
reflects stronger template alignment, which may depend on smaller receptive fields and
sharper SF tuning[39], thereby supporting higher sensitivity. Neurons have intrinsic variability
due to factors such as fluctuations in firing rate and membrane potential[40]. At the system
level, internal noise is shaped by neural correlations: when neural activity is more correlated
among neurons, their noise is less likely to cancel out through averaging, leading to greater
internal noise at the system level[41,42]. Nonlinearity is shaped by both retinal and cortical
processes, including synaptic rectification[43] and divisive normalization[44].

Each computation influences performance in distinct ways (**Fig. 1D**). When gain differs
between two locations, the system's ability to filter both signal and noise changes
proportionally, resulting in a vertical shift of the threshold-versus-noise curve without altering
its shape. Differences in internal noise impact performance the most at lower external noise
levels, where internal noise dominates. As external noise becomes more dominant,
performance at higher external noise becomes indistinguishable across locations. Finally,
nonlinearity determines the curvature of the function.

To investigate how these system-level computations shape visual field heterogeneity, we
assessed whether they vary throughout the visual field. Previous studies have shown that
internal noise increases across eccentricity[45–47]. Neuroimaging studies have shown that the
size of surface area decreases with eccentricity (e.g[19,20,24]). These studies averaged data
around polar angle; thus, it is unknown whether polar angle modulates these computations.
Based on the anatomical evidence that the decline of cone and retinal ganglion cell
density[15–18] and cortical surface area[21] across eccentricity is steeper along the vertical than
horizontal meridian, as well as behavioral findings that the eccentricity effect is more
prominent along the vertical than horizontal meridian (e.g[5,48–50]) and a robust HVA (e.g[4,8]),
we hypothesize that the eccentricity effect as well as the gain and internal noise underlying
this effect will vary more along the vertical than horizontal meridian.

It is unknown whether gain and internal noise also vary around polar angle. Anatomical and
neuroimaging measurements, including cone and retinal ganglion density[15,16], as well as V1
cortical surface area[21–25], population receptive field size[21,24], and BOLD signal amplitude[11,51],
contribute to the behavioral polar angle asymmetries[26,27]. Accordingly, both gain and internal
noise may vary around polar angle. However, magnifying stimulus size only reduced
variation around polar angle, suggesting that differences in V1 cortical surface area do not
fully explain polar angle asymmetries[6]. Here, we explored whether in addition to retinal and
cortical differences, polar angle asymmetries also reflect differences in gain and internal
noise, or whether the underlying computations differ from those underlying the eccentricity
effect. We also assessed nonlinearity across eccentricity and around polar angle, as it
reflects how the visual system amplifies or compresses neural responses and plays a critical
role in shaping the threshold-versus-noise curve.

Second, we examined whether variations in gain, internal noise, and nonlinearity across
locations contribute to performance heterogeneity. Specifically, we asked whether the
magnitude of variation in system-level computations—quantified by a location difference
index—predicts corresponding differences in performance. By quantifying the impact of each
computation on performance[47], this analysis provides further evidence that system-level
computations play a mechanistic role in shaping visual perception throughout the visual field.

180   **Results**

Observers performed an orientation discrimination task at nine visual field locations: fovea,
four cardinal meridians (left and right horizontal meridian and upper and lower vertical
meridian) at 4⁰ and 8⁰ eccentricity (**Fig. 2A**). The target signal was a tilted Gabor embedded
in dynamic white noise with a SF of either 4 or 6 cpd. We varied the contrast of Gabor and
noise across trials to map contrast threshold as a function of external noise, location and SF
(**Fig. 2B**). We estimated three parameters—gain, additive internal noise, and nonlinearity—
by fitting the perceptual template model to threshold-versus-noise curves at each location
(**Fig. 2C**). Finally, we assessed how each parameter varies across eccentricity and around
polar angle, and contributes to location differences of performance.

Figure 2. Stimulus, procedure, and model fitting. **(A)** Orientation discrimination task.
Observers saw a streak of dynamic white noise presented at nine locations (fovea, four polar
angle at 4º and 8º eccentricity) both before and after seeing a Gabor signal presented at one
location. They reported whether the Gabor was tilted clockwise or counterclockwise and
received feedback. The signal location, noise contrast and signal contrast varied across
trials. The Gabor SF varied across sessions (4 and 6 cpd, counterbalanced). Eye fixation
was enforced by an eye tracker. **(B)** To derive contrast thresholds at each location and each
noise level, we fitted a Weibull function to accuracy as a function of Gabor contrast (binned
into 10 bins) for each noise level (indicated by lines of different shades) and estimated
threshold (indicated by the dots of different shades at x-axis). **(C)** We fitted the perceptual
template model to the thresholds across external noise to derive the three computations:
gain, additive internal noise and nonlinearity. The solid dots indicate the threshold estimated
in **(B)**.

Visual field heterogeneities in the absence of external noise

We first verified the visual field heterogeneity—the eccentricity effect and polar angle
asymmetries—in absence of external noise and examined how SF modulated their
magnitudes. **Fig. 3** shows threshold-versus-noise curves for each location group at 4 and 6
cpd. Only thresholds estimated at 75% accuracy are shown, for illustration purposes. We
focused on thresholds estimated at external noise equal to 0, referred to as *performance*.
We fit linear mixed-effects models to performance and conducted Type III ANOVAs to
examine interactions between location (eccentricity/meridian/specific location) and SF (4 and
6 cpd) and their main effects (Online Methods, equation (9)).

Figure 3. Threshold-vs-noise curves and performance throughout the visual field. Each panel plots thresh-vs-noise curve for each location group to visualize one of the three visual field heterogeneities: eccentricity effect along the horizontal **(A)** and vertical meridian **(B)**, HVA **(C)** and VMA **(D)**. Dots and error bars represent group-averaged contrast thresholds (±1 SEM) across nine levels of external noise. Many error bars are smaller than the symbols. Shaded bands are PTM model predictions (±1 SEM). Colors denote visual field locations, indicated by icons in each panel, and corresponding R^2 values for PTM fitting are shown alongside. Dot shapes indicate SF (circles=4 cpd, triangles=6 cpd), plotted separately in the left and right panels within each dashed cell. **(A, B)** Threshold-vs-noise curves at the fovea, and at 4º and 8º eccentricity along the horizontal **(A)** and vertical **(B)** meridian. To examine eccentricity effects, we compared *performance*—defined as the absolute value of the log contrast threshold at zero external noise—between each pair of eccentricities (fovea vs. 4º, fovea vs. 8º, and 4º vs. 8º). Vertical comparison lines indicate the pairs tested; asterisks denote Bonferroni-corrected p-values (***p < .001, **p < .01, *p < .05, ns, p>.1). **(C)** Threshold-vs-noise curves at the horizontal and vertical meridians at both 4º and 8º eccentricity. Subpanel (a) plots performance as a function of meridian × eccentricity, separately for 4 and 6 cpd, illustrating how both eccentricity and SF modulate the HVA. Group means (±1 SEM) are plotted as colored dots; individual subjects' performance (median of 1,000 bootstraps) is shown in grey lines. We focused on comparisons between horizontal and vertical meridian, indicated by horizontal comparison lines. Asterisks denote significance levels from post-hoc pairwise tests. **(D)** Threshold-vs-noise curves at the lower and upper vertical meridians at 4º and 8º eccentricity. Subpanel **(a)** shows performance collapsed across SF as a function of meridian × eccentricity, revealing eccentricity-dependent modulation of the VMA. Formatting follows that of **(C)**.

We observed an interaction between eccentricity and SF along both the horizontal meridian ($F(2, 45.7) = 7.277$ [1.623, 9.959], $p_{obs} = 0.002$, eta2p = 0.241 [0.067, 0.303]; **Fig. 3A**) and the vertical meridian ($F(2, 45.7) = 29.089$ [10.615, 30.430], $p_{obs} < 0.001$, eta2p = 0.560 [0.318, 0.570]; **Fig. 3B**). Post-hoc pairwise comparisons revealed that performance decreased with eccentricity along both meridians and both SFs (all eccentricity pairs had $p_{emp} < 0.001$). This decrement was more pronounced for 6 than 4 cpd and for vertical than horizontal meridian (averaged Cohen's *d*: Vertical-6 cpd > Vertical-4 cpd > Horizontal-6 cpd > Horizontal-4 cpd). These results confirm the well-established eccentricity effect, in which a decline in performance is more pronounced at higher SFs and along the vertical than the horizontal meridian.

We then compared performance between the horizontal and vertical meridian (**Fig. 3C-a**)
and examined the modulation of eccentricity (4º and 8º) and SF (4 and 6 cpd) (Online
methods, equation (10)). We found a three-way interaction among meridian, eccentricity and
SF ($F(1,64.7) = 6.728$ [0.178, 10.399], $p_{obs} = 0.012$, eta2p = 0.094 [0.003, 0.139]). Post-hoc
analyses revealed that performance was higher at the horizontal than the vertical meridian
for all four SF x eccentricity combinations, confirming a HVA in performance. The largest
HVA was observed at 6 cpd x 8º eccentricity ($t(65.0) = 14.030$ [7.869, 14.100], $p_{emp} < 0.001$,
Cohen's $d = 17.702$), followed by 4 cpd x 8º (4 cpd; $t(65.0) = 5.890$ [3.060, 7.730], $p_{emp} <$
0.001, Cohen's $d = 8.122$). The asymmetries at 4º eccentricity were smaller but still
statistically significant for both 4 cpd ($t(65.0) = 3.010$ [1.410, 10.399], $p_{emp} < 0.001$, Cohen's
$d = 5.907$) and 6 cpd ($t(65.0) = 5.090$ [3.060, 5.390], $p_{emp} < 0.001$, Cohen's $d = 5.225$). In
sum, the performance HVA was robust and stronger at further eccentricity and higher SF.

Along the vertical meridian (**Fig. 3D-a**), the three-way interaction (location x eccentricity x
SF) was not significant ($p_{obs} > 0.1$). Location interacted with eccentricity ($F(1,64.6) = 4.400$
[0.945, 9.634], $p_{obs} = 0.040$, eta2p = 0.064 [0.014, 0.130]) but not with SF ($p_{obs} > 0.1$),
indicating that performance variation along the vertical meridian was modulated by
eccentricity, but not by SF. Post-hoc analysis showed that performance was better at lower
than upper vertical meridian at both 4º ($t(65.0) = 1.300$ [0.139, 1.890], $p_{emp} = 0.044$, Cohen's
$d = 2.258$) and 8º eccentricity ($t(65.0) = 4.270$ [2.469, 5.060], $p_{emp} < 0.001$, Cohen's $d =$
5.806), confirming a VMA in performance.

Together, we observed robust eccentricity effects and polar angle asymmetries in
performance. Effects are more pronounced at the higher SF and farther eccentricities.

Gain, additive noise and nonlinearity vary throughout the visual field

To examine how computations underlie the observed visual field heterogeneity, we first
assessed whether and how each parameter varied throughout the visual field. We fitted
linear mixed-effects models (Online Methods, equation (11-12)) to each parameter and
conducted Type III ANOVAs to test interactions between locations and SF and their main
effects. Matching between a parameter's variation pattern with typical visual field
performance patterns—eccentricity effect, HVA and VMA—suggests that the corresponding
computation underlies these visual field heterogeneities.

Gain, additive noise and nonlinearity vary across eccentricity

When eccentricity increases from fovea to 4º, at 4 cpd the curve differs at lower external
noise levels but converges at higher levels (**Fig. 3A**), indicating increased internal noise at
parafoveal locations (**Fig. 1D**, *middle*). As eccentricity increases further to 8º eccentricity,
performance decreased across all levels of external noise, suggesting a reduction in gain at
peripheral locations (**Fig. 1D**, *left*). Based on these qualitative differences, we quantitatively
compared gain and internal noise across the visual field. For gain, along the horizontal
meridian (**Fig. 4A**), eccentricity interacted with SF ($F(2, 57.0) = 6.032$ [0.509, 9.016], $p_{obs} =$
0.004, eta2p = 0.175 [0.018, 0.264]). Post-hoc pairwise comparisons showed that for 6 cpd,
8º eccentricity had lower gain than both fovea ($t(46.7) = 3.220$ [0.799, 3.830], $p_{emp} = 0.048$,
Cohen's $d = 3.248$) and 4º eccentricity ($t(46.7) = 4.680$ [1.430, 4.620], $p_{emp} = 0.036$, Cohen's
$d = 4.400$); but for 4 cpd, gain was similar across eccentricity. For additive internal noise

(**Fig. 4B**), there was a main effect of eccentricity ($F(2, 45.1) = 34.842$ [10.783, 38.066], $p_{obs} <$
0.001, eta2p = 0.607 [0.307, 0.623]): fovea was lower than 4º ($t(46.3) = -5.310$ [-5.991, -
2.350], $p_{emp} = 0.006$, Cohen's $d = -4.380$), which in turn was lower than 8º eccentricity
($t(46.3) = -2.920$ [-3.540, -1.110], $p_{emp} < 0.001$, Cohen's $d = -3.611$). Lastly, nonlinearity (**Fig.
4C**) also varied with eccentricity along the horizontal meridian ($F(2, 46.6) = 11.916$ [1.653,
14.295], $p_{obs} < 0.001$, eta2p = 0.338 [0.065, 0.379]), being higher at the fovea than 8º
eccentricity ($t(46.3) = 4.870$ [1.640, 5.221], $p_{emp} = 0.006$, Cohen's $d = 3.523$).

Figure 4. Gain, additive noise and nonlinearity vary across eccentricity. Each panel shows
group means (±1 SEM) of estimated model parameters—gain **(A, D)**, additive noise **(B, E)**,
and nonlinearity **(C, F)**—as a function of eccentricity along the horizontal (top row) and
vertical (bottom row) meridian. Grey lines represent individual subject data (median of 1,000
bootstraps). Asterisks above horizontal comparison lines indicate Bonferroni-corrected post-
hoc significance levels (***p < .001, *p < .05, ns p > .1). Parameters that showed a
significant interaction between eccentricity and SF (e.g., gain in panels **A** and **D**) are plotted
separately for 4 and 6 cpd. Parameters showing only a main effect of eccentricity without
such interaction are collapsed across SFs (additive noise in panels B and E, and nonlinearity
in panels **C** and **F**).

A similar pattern emerged at the vertical meridian (**Fig. 4D-F**). For gain, eccentricity
interacted with SF ($F(2, 57.0) = 11.098$ [0.657, 10.210], $p_{obs} < 0.001$, eta2p = 0.280 [0.028,
0.297]; **Fig. 4D)**. At 6 cpd, Gain was lowest at 8º eccentricity and lower than at 4º ($t(46.7) =$
8.720 [3.229, 8.011], $p_{emp} < 0.001$, Cohen's $d = 7.123$) and at the fovea ($t(46.7) = 9.870$
[3.449, 8.752], $p_{emp} < 0.001$, Cohen's $d = 8.542$). Moreover, at 4 cpd, gain was lower at 8º
than 4º eccentricity ($t(46.7) = 5.990$ [1.339, 7.720], $p_{emp} = 0.024$, Cohen's $d = 3.418$).
Additive noise increased with eccentricity (**Fig. 4E**), ($F(2, 45.3) = 48.393$ [13.872, 46.819],
$p_{obs} < 0.001$, eta2p = 0.681 [0.361, 0.673]) without interacting with SF ($p_{obs} > 0.1$). Additive
noise was lower at the fovea than at 4º ($t(46.2) = -6.130$ [-6.780, -2.569], $p_{emp} = 0.024$,
Cohen's $d = -4.343$), which in turn was lower than at 8º eccentricity ($t(46.2) = -3.590$ [-4.120,
-1.000], $p_{emp} = 0.006$, Cohen's $d = -3.215$). Similarly, nonlinearity varied across eccentricity
($F(2, 47.0) = 6.787$ [0.745, 11.888], $p_{obs} = 0.023$, eta2p = 0.224 [0.030, 0.337]; **Fig. 4F**),

being higher at the fovea than 8^o eccentricity (t(46.4) = 3.600[0.930, 4.580], p_{emp} = 0.024,
Cohen's d = 2.883).

In sum, gain decreased with eccentricity from fovea to 8^o, and more so from 4^o to 8^o—with
this effect being the most pronounced along the vertical meridian and for 6 cpd. Additive
noise increased with eccentricity, and more so along the vertical than horizontal meridian.
Nonlinearity was higher at fovea than 8^o eccentricity similarly along both meridians.

Gain, but not internal noise or nonlinearity, varies around polar angle

We next assessed whether gain differed between horizontal and vertical meridian and
whether this was modulated by eccentricity and SF (**Fig. 5A**). There was no three-way
interaction (p_{obs} > 0.1), but meridian interacted with eccentricity (F(1, 56.9) = 13.960 [1.503,
25.598], p_{obs} < 0.001, eta2p = 0.197 [0.022, 0.290]; **Fig. 5A-a**) and SF (F(1, 56.9) = 4.430
[0.006, 8.976], p_{obs} = 0.040, eta2p = 0.072 [0, 0.123] **Fig. 5A-b**). Post-hoc analyses showed
that gain was higher at the horizontal than the vertical meridian at 8º eccentricity (t(65.5) =
6.130 [1.870, 6.560], p_{emp} < 0.001, Cohen's d = 4.297) but not at 4º (p_{emp} > 0.1), consistent
with the HVA pattern for performance. Similarly, gain reflected the HVA pattern at 6 cpd
(t(65.5) = 5.370 [1.560, 4.650], p_{emp} = 0.008, Cohen's d = 4.649), but not at 4 cpd (p_{emp} >
0.1). In sum, gain showed a pattern consistent with HVA, with stronger effects at greater
eccentricities and higher SF. Neither additive noise (**Fig. 5B**) nor nonlinearity (**Fig. 5C**)
differed between horizontal and vertical meridian (p_{obs} > 0.1) at any eccentricity or SF. Note
that the lower additive noise at 4º than 8º eccentricity (**Fig. 5B**) was already captured by the
eccentricity difference (**Fig. 4B**). These findings suggest that HVA arises specifically from
variation in gain.

Figure 5. Gain **(A and D)**, but not internal noise **(B and E)** or nonlinearity **(C and F)**, varies
around polar angle. We plot estimated parameters as a function of location × eccentricity
pairs, separately for 4 and 6 cpd. To highlight post hoc comparisons from significant two-way
interactions for gain **(A and D)**, we collapsed across SF to illustrate the modulation effect by
eccentricity (panels **A-a** and **B-a**), given a significant meridian/location × eccentricity
interaction. Similarly, we collapsed across eccentricity to illustrate the modulation effect by
SF (panels **A-b** and **B-b**), given a significant meridian/location × SF interaction. Figure
format follows that of **Fig. 4**.

We next examined whether gain differed between the lower and upper vertical meridian (**Fig.**
5D). Location interacted with SF (F(1, 76.0) = 6.469 [0.071, 12.967], p_{obs} = 0.013, eta2p =
0.078 [0.001, 0.165] **Fig. 5D-a**), and gain was higher at lower than upper vertical meridian at
4 cpd (t(65.5) = 4.410 [1.280, 5.580], p_{emp} < 0.001, Cohen's d = 3.374), but not at 6 cpd (p_{emp}
> 0.1). This result means that VMA for gain was stronger at 4 than 6 cpd. The lack of VMA at
a higher SF is because for some observers, gain at 6 cpd (particularly at 8º eccentricity) was

near the floor level (close to 1), limiting any differences. There was no three-way interaction
or interaction between location and eccentricity ($p_{obs} > 0.1$). Neither additive noise (**Fig. 5E**)
nor nonlinearity (**Fig. 5F**) differed between the upper and lower vertical meridian ($p_{obs} > 0.1$).
Together, these findings indicate that differences in gain, rather than in internal noise or
nonlinearity, may underlie VMA.

In sum, gain alone consistently varies around polar angle, indicating that it is related to both
HVA and VMA. These location-based effects are more pronounced at higher SFs. Distinct
from the eccentricity effect, additive noise and nonlinearity did not vary around polar angle.

Contribution: Gain, but not additive noise or nonlinearity, contributes to eccentricity
effect

We examined the extent to which each system-level parameter—gain, additive internal
noise, and nonlinearity—contributed to visual field asymmetries. First, we identified the
direction in which each parameter modulated performance. Using linear mixed-effects
models (Online Methods, equation (13)), we assessed the relation between performance
and each parameter across seven locations in the visual field: the fovea, the horizontal
meridian and the upper and lower vertical meridians at both 4º and 8º eccentricities.

As shown in **Fig. 6A-C**, locations with higher performance were associated with higher gain
(slope = 0.090 [0.047, 0.130], $p_{emp} < 0.001$, conditional R-squared=0.75 [0.67, 0.80]; **Fig.**
6A) and lower additive noise (slope = -0.029 [-.042, -0.008], $p_{emp} < 0.05$, conditional R-
squared=0.77 [0.70, 0.81]; **Fig. 6B**). Nonlinearity showed no significant relation with
performance (slope = -0.019 [-0.043, 0.02], $p_{emp} = 0.082$, conditional R-squared = 0.79 [0.72,
0.82]; **Fig. 6C**). These findings confirm our hypothesis that better filtering and lower internal
noise improve performance, whereas nonlinearity has no effect.

The direction of each modulation informs the directionality of parameter contribution to
location-based performance differences: Parameters that enhance performance should vary
across locations in line with performance variation, whereas parameters that impair
performance should vary in the opposite direction. We examined whether and how these
location differences of each parameter are proportional to the heterogeneity in the expected
direction (equation (14)). Location variation was indexed by the difference between two
locations divided by their sum, providing a normalized measure. In **Fig. 6D,** we plot location
difference indices for each parameter against those for performance.

For gain, the only significant correlation was between the fovea and 8º along the horizontal
meridian (slope = 0.164 [0.020,0.412], $p_{emp, right-tailed} = 0.038$, conditional R-squared = 0.75
[0.48, 0.93], **Fig. 6D-a**). The effect was driven by the positive correlation at 6 cpd ($p_{emp, right-}$
$_{tailed} = 0.030$). No correlations were observed for the other eccentricity pairs (**Fig. 6D-b and**
c), or for internal noise and nonlinearity (all $p_{emp} > 0.1$, **Fig. 6D-e-i**), despite their variations
across eccentricities. These results suggest that gain, particularly at higher SF, is the
primary factor underlying the eccentricity effect along the horizontal meridian.

In contrast, no correlation with polar angle differences was significant ($ps_{emp}>.1$), meaning that
although gain varies around polar angle, it does not contribute to polar angle asymmetries as
much as to the eccentricity effect.

Figure 6. Gain contributes to performance differences between the fovea and 8⁰
eccentricity. **(A-C)** Each panel shows estimated gain **(A)**, additive internal noise **(B)**, and
nonlinearity **(C)** as a function of performance across locations and SFs. Each data point
represents an individual location × SF combination (color-coded by location, shape-coded by
SF). Colored regression lines show predictions from linear mixed-effects models. Medians of
bootstrapped slopes and empirical p-values (p_{emp}) are reported in the figure. Performance
improves with increasing gain **(A)** and decreasing internal noise **(B)**, while nonlinearity
shows no relationship with performance **(C)**. **(D)** Quantify the contribution of each
computation. Each panel plots the location difference index (LDI) of a parameter against the
LDI of performance. Shapes denote individual observer data at 4 cpd (circles) and 6 cpd
(triangles). Regression lines reflect linear mixed-effects model predictions for each SF.
Median of bootstrapped regression slopes and empirical one-tailed p-values are shown in
each panel. Subpanels **(a-c)** show gain results across three eccentricity pairs (fovea vs. 8⁰,
fovea vs. 4⁰ and 4⁰ vs. 8⁰) along the horizontal meridian. A significant positive correlation
appears only for the fovea vs. 8⁰ pair **(D-a)**, especially at 6 cpd, indicating that gain
differences contribute to the eccentricity effect. Subpanels **(d-f)** show additive internal noise,
which shows no significant correlations with performance for any eccentricity pair. Subpanels
(g-i) shows nonlinearity, which similarly shows no significant correlation with performance
across eccentricity pairs.

419   **Discussion**

Using the equivalent noise method and perceptual template modeling, we mapped contrast
sensitivity across the visual field and identified distinct computations that underlie
performance variations across eccentricity –which have been examined primarily along the
horizontal meridian or collapsed across polar angle (review[3])– and around polar angle. With
increasing eccentricity, all three computations varied: gain and nonlinearity decreased,
whereas internal noise increased. These variations across eccentricity were stronger along
the vertical than horizontal meridian, consistent with visual performance (e.g[5,48,50,52], and they
were also stronger at higher SFs, in line with previous studies[5,53,54]. Moreover, gain predicted
contrast sensitivity differences between fovea and 8°, indicating that gain is a key driver of
the eccentricity effect. In contrast, around polar angle, neither internal noise nor nonlinearity
varied, only gain did—being higher along the horizontal than the vertical meridian and at the
lower than the upper vertical meridian—mirroring the HVA and VMA in performance.
Together, these findings indicate that polar angle asymmetries arise from a distinct
computational profile.

We found that the eccentricity effect and the HVA were more robust at 6 cpd than at 4 cpd.
These findings are consistent with previous studies (eccentricity effect[5,53,54]; HVA[5,7,9,48,55]). A
parallel pattern emerged for gain: the eccentricity effect and HVA were stronger at 6 cpd.
This shared SF dependence reveals a qualitative link between performance and underlying
computations, which likely reflect neural constraints: peripheral neurons prefer lower SFs in
both humans[56,57] and non-human primates[39,58,59]. Higher SFs place greater demands on
spatial resolution, taxing locations with lower resolving power and thereby amplifying
performance differences across eccentricity—and potentially around polar angle as well. The
distinctions in underlying computations for eccentricity effect and polar angle asymmetries
are consistent with a recent study, which using reverse correlation and a noisy-observer
model, showed higher sensitivity to signal orientation and SFs and lower internal noise at the
fovea than the parafovea[47]. In contrast, only orientation sensitivity varies around polar angle,
mirroring performance[60]. The gain —an overall measure of how much the signal is amplified
(current study) — and featural representation —the sensitivity profile to different
features[47,60]— provide complementary descriptions of how the system enhances relevant
input. Moreover, internal noise estimates obtained using the perceptual template model in
the current study, and those derived from a noisy-observer model [47,60], quantify the same
system-level property and together offer a more complete account of internal variability
across the visual field.

These distinctions in underlying computations may reflect the interplay of four
neurophysiological factors: (1) the number of neurons co-activated by visual stimuli, (2) the
tuning properties of individual neurons (e.g., sensitivity and selectivity for visual features), (3)
the intrinsic neural noise of individual neurons (e.g., firing variability and dendritic noise[40]),
which contributes to system-level internal noise, and (4) correlations in activity across
neurons[61]. By linking behavioral results to model-derived system-level computations, our
results provide inferences about how these neurophysiological properties vary throughout
the visual field.

Across eccentricity, the decrease in gain is consistent with two neurophysiological factors:
First, as eccentricity increases, the cortical surface area stimulated by a given visual input

decreases (e.g[19–24]), and thus, assuming uniform cortical density[62], fewer neurons are
recruited. Second, fewer neurons are tuned to high SFs at peripheral than central locations[63]
and receptive fields become larger[64]. Pooling from fewer and less well-tuned neurons likely
result in weaker signal amplification and thus lower gain.

Whereas gain was substantially reduced from the fovea to the perifovea, it did not differ
between the fovea and the parafovea. This result may seem surprising given the known
decline in spatial resolution with eccentricity, but it may reflect an interaction between
broader neural tuning and increased neural correlation. The increase in neural correlation at
the parafovea can be attributed to a lower proportion of orientation-selective neurons[65] and
fewer parvocellular neurons[64], which exhibit broader selectivity for SF [66,67], than in the fovea.
Together, these factors would offset the tuning advantage at the fovea, resulting in similar
gain as at the parafovea. However, as eccentricity increases towards the perifovea, tuning
continues to decrease while correlations remain high, resulting in a substantial drop in gain,
consistent with our observations. The increase in internal noise with eccentricity may reflect
a tradeoff between neural noise and neural correlation: The more correlated the neurons'
activity, the less likely their noise is to average out[41,42]. Although direct measurements of
intrinsic noise across eccentricity are lacking, even if the noise of single neurons remains
constant, increased neural correlation from fovea to periphery will result in greater system-
level internal noise. This connection between higher neural correlation and higher internal
noise is supported by findings that attention is associated with both lower neural
correlations[66,67] and reduced internal noise (e.g[33]).

The present study also provides the first assessment of nonlinearity across eccentricity.
Here, nonlinearity —which reflects the system's sensitivity to amplified signals (modeled by
the power function in the perceptual template model)— exceeded 1 at all locations.
Importantly, models without a nonlinear component, including the linear amplifier model[21–
25,51,68] and variants of perceptual template models[33,38], failed to account for our data. These
results underscore the need to consider nonlinear computations to explain visual
performance. Further research is needed to determine whether this enhanced foveal
nonlinearity shares neural mechanisms with the higher gain and lower additive noise
observed at central locations.

Around polar angle, only gain varied and reflected both the HVA and VMA. Higher gain at
better-performing locations may arise from greater cortical surface area[69]—implying more
neurons, smaller population receptive field sizes[70], and a preference for higher SFs [56,57,60].
This observed gain variation constrains hypotheses about how neural correlations vary
around polar angle: robust gain variation around polar angle suggests that better-performing
locations not only have superior tuning but also exhibit weaker neural correlations.

This interpretation on neural correlation, in turn, informs the observed invariance in internal
noise around polar angle. As discussed in the eccentricity section, such invariance may
reflect a tradeoff between neural correlation and neural noise. Thus, better-performing
locations with weaker neural correlations —as suggested by our gain results—should have
higher intrinsic neural noise. Although it may seem counterintuitive that neurons at better-
performing locations could have higher neural noise, increased neural noise can be
advantageous, e.g., it can facilitate the detection of subthreshold signals[69]. Future studies

should map both neural correlations and neural noise across the visual field to characterize
their interplay.

In sum, the observed gain variation across both eccentricity and polar angle likely results
from an interplay of three factors at better-performing locations: a greater number of
neurons, preference for higher SFs, and stronger neural correlations. In contrast, variation in
internal noise across eccentricity —and its invariance around polar angle— may be jointly
influenced by neural correlations and intrinsic neural noise, which our results suggest
varying across the visual field. The relative impact of these neural factors remains an
empirical question. A perceptual template model with population coding showed that neural
correlations have a greater impact on performance than mere amplification or sharpening of
tuning functions[71].

Our findings may help explain why M-scaling eliminates eccentricity effects but not polar
angle asymmetries in contrast sensitivity[6]. While M-scaling normalizes the cortical surface
area stimulated, it may not equate other neurophysiological properties. For example,
increasing the number of neurons can reduce neural correlation[41] and diversify tuning
profiles, further lowering such correlations. This reduction in neural correlation in the
perifoveal region likely decreases system-level internal noise and therefore eliminates the
eccentricity effect. In contrast, because internal noise appears uniform around polar angle,
polar angle asymmetries—which are primarily driven by gain variations—may persist despite
M-scaling.

It has been proposed that asymmetries between the upper and lower hemifields may relate
to functional specialization, where the upper visual field primarily processes distant objects
and the lower visual field processes near objects[72]. Polar angle asymmetries in contrast
sensitivity[4–8] and acuity[72] decrease gradually with angular distance from the cardinal
meridians, suggesting that whereas the near-versus-far specialization contributes to
meridional asymmetries, it cannot fully account for the polar angle variations. We therefore
hypothesize that gain-related polar angle asymmetries progressively diminish along non-
cardinal meridians and disappear along the intercardinal meridians.

534
Previous studies have revealed ontogenetic and phylogenetic factors in these visual field
asymmetries. Ontogenetically, children show an HVA both behaviorally[72] and in cortical
surface area[73] but lack a behavioral[74] or cortical[24] VMA. The HVA progressively strengthens
through adolescence, whereas the VMA emerges and becomes as pronounced as that of
adults in late adolescence[73]. Phylogenetically, non-human primates show the typical HVA
but an inverted VMA in visual performance. Evolutionary differences in locomotion,
navigation, and tool use, which result in distinct visuomotor interactions at specific visual
field locations, may contribute to shaping the organization of the visual field[74].

543
In conclusion, by linking our novel behavioral data to underlying system-level computations,
this study provides insights into the neurophysiological architecture of the visual field and
generates novel, testable hypotheses. We found that visual field heterogeneities arise from
distinct system-level computations. The eccentricity effect is driven by a combination of
reduced gain and nonlinearity alongside increased additive noise, consistent with changes in
cortical magnification and receptive field properties. Polar angle asymmetries are associated
with differences in gain, but not in internal noise or nonlinearity. Together, these findings

highlight a fundamental distinction in how the visual system implements computations across
eccentricity versus and around polar angle, advancing our understanding of visual
processing throughout the visual field.
554

Online Methods

Observers

Twelve observers (9 females; age mean=26.44; SD=3.24) participated in this experiment. All
observers except two authors (S.X. and Q.C.) were naïve to the study's purpose. All
observers had normal or corrected-to-normal vision and provided written informed consent.
The experimental methods complied with the Helsinki Declaration and were approved by the
Institutional Review Board for Human observers at New York University.

Apparatus

All experimental stimuli were generated using MATLAB 2020b (MathWorks) and the
Psychophysics Toolbox[75], and displayed on a gamma-calibrated CRT monitor (40 cm x 30
cm, 1,280 × 960 screen resolution, 100 Hz, 26 cd/m^2 background luminance). Each pixel
extends 0.03 x 0.03 dva squared. The display was calibrated using a Photo Research
(Chatsworth, CA) PR650 Spectracolorimeter to generate linear lookup tables for this
experiment. The display resolution was 32 pixels per degree of visual angle. An EyeLink
1000 eye-tracker was positioned in front of observers to track gaze position. Observers were
seated in a dark room with their head stabilized using a chin rest 57 cm from the display
monitor.

Stimuli

The stimulus presented in each trial consisted of a target, sandwiched between two streaks
of dynamic noise patches. The target was a Gabor patch—a sinusoidal grating modulated by
a Gaussian envelope—with luminance ( ) defined at each pixel position (and , relative to
the patch's center) (**Fig. 2A**, *inset*):

[Equation 1]

$$L(x, y) = C \cos(2\pi f(x' + \phi)) \exp\left(-\frac{x^2 + y^2}{2\sigma_{env}^2}\right)$$

$$x' = x \sin\theta + y \cos\theta$$

and   range from -48 to 48 pixels (diameter=3 dva).   represents linear contrast, either
sampled from a staircase procedure or picked at a fixed value (Procedure).   denotes SF
(0.125 or 0.1875 cycles per pixel, equivalent to 4 and 6 cpd).   denotes orientation in
radians, either counterclockwise (/4) or clockwise (- /4) with equal probability on each trial.
denotes the phase which was randomized on each trial.   specifies the Gaussian
envelope size (=32 pixels, or 1 dva, one-third of the grating size).

The Gabor was embedded in dynamic noise, which were generated independently on each

frame   by sampling pixels from a Gaussian distribution with standard deviation (   ) and

centered at 0 (hence, the RMS contrast of the noise patch equals   ) (**Fig. 2A**, *inset*):

[Equation 2]

$$N_t(x, y) \sim \mathcal{N}(0, \sigma_{\text{ext}}^2)$$

On each trial,   could be one of nine possible values (0, 0.0275, 0.055, 0.0825, 0.11,

0.165, 0.22, 0.33, 0.44).

The possible stimulus locations were indicated by placeholder (**Fig. 2A**, *first display*): one at

the fovea, and four along the horizontal (Left: 0º; right: 180º) and vertical meridians (upper:

90º; lower: 270º) at either 4° (parafovea) or 8° (perifovea) eccentricity. Each placeholder

consisted of four corners (line length=0.75°).

Procedure

Fig. 2A shows the trial sequence. Each trial began with all placeholders and a fixation dot

(radius=0.1º of visual angle) at the center of a gray background. A streak of dynamic white

noise was then presented at each location for a total duration of 630 ms. On each trial, a

target Gabor was presented for 30 ms at one of the possible locations, 300 ms after the

onset of the dynamic white noise, while other locations were only filled with noise. The streak

of dynamic white noise persisted for 300 ms following target offset. The target location was

marked by a boldened placeholder at the onset of the target Gabor, and until the observer's

response, eliminating location uncertainty.

Target location, orientation, and external noise levels were randomly selected on each trial.

The Gabor SF (4 or 6 cpd) was constant within each experimental session. Observers

performed an orientation discrimination task, pressing one of two keys to indicate whether

the Gabor was tilted counterclockwise or clockwise (±45º). Task difficulty varied with

stimulus parameters—Gabor visibility increased, hence tasks got easier, with higher contrast

and lower noise standard deviation   . Observers received auditory feedback after

making a response: a high pitch for correct and a low pitch for incorrect responses. Their

average accuracy was displayed at the end of each block.

We collected data to estimate contrast thresholds in two steps. First, at each noise level and

location, we conducted a titration procedure using four interleaved staircases (two 1-up-2-

down and two 1-up-3-down) starting from different contrast levels, with 40 trials per

staircase. Second, based on Weibull fitting, we collected 30 to 40 additional trials at selected

Gabor contrast levels—both within the dynamic range and at anchoring levels. We

interleaved external noise and locations for all observers. Each observer completed 12,600

trials across approximately eight one-hour sessions for each SF. Nine observers were tested

with both 4 cpd and 6 cpd stimuli, three were tested only at 6 cpd. They completed 25,200

trials in approximately 16 sessions. Stimuli generation and response recording were

conducted with MATLAB. Data analysis and visualization were conducted in both MATLAB

and R (R Core Team, 2024; RStudio v2024.12.1+563).

Eye-tracking

We monitored the gaze position of the right eye in real-time during each trial using an SR
Research Eyelink 1000 Desktop Mount eye tracker at a sampling rate of 1000 Hz. Stimulus
presentation was contingent on maintaining central fixation from the onset until the offset of
the dynamic white noise. Trials were terminated and re-queued to the end of the block if
observers' gaze deviated more than 1.5⁰ of visual angle from the fixation center.

Estimate contrast thresholds

For each noise level and location, we constructed psychometric functions by fitting a Weibull
function to trials tested at varying Gabor contrast and corresponding binary responses (**Fig.**
2B). Trials collected through titration were binned into 10 bins based on percentile. Following
titration, we collected additional trials at 2 to 3 fixed signal contrast levels per noise
condition. These contrast levels were manually selected based on preliminary titration fits to
improve the robustness of the psychometric function fitting.

[Equation 3]

$$f(C) = \lambda + (1 - \lambda - \gamma)\left(1 - e^{-\left(\frac{C}{\alpha}\right)^{\beta}}\right)$$

denotes accuracy as a function of Gabor contrast for each bin. , , and
represent the function's threshold at 75% accuracy, steepness, lapse rate, and chance
performance, respectively. Model parameters were estimated using constrained nonlinear
optimization via MATLAB's fmincon() function. Parameter bounds were set as follows: thresh
(\alpha) ranging from 1% to 100%, steepness ( ) from 0.01 to 100, guess rate ( ) from 0.45
to 0.55, and lapse rate (lambda) from 0 to 0.1, in 100 steps.

To balance the trade-off between trial count and fitting quality, we added "stabilizer" trials at
low contrast levels (50 trials each at 0.01% and 0.1% Gabor contrast, enforcing 50%
accuracy). At these low contrast levels, all observers were expected to perform at chance
level, assuming no bias in reporting leftward or rightward orientation. The stabilizer trials
were used only for fitting purposes and their trial counts were not included in the TvN fitting
weights.

Perceptual template model characterizes TvN curves

For each observer, we fitted a simplified version of the perceptual template model[33,34] to
contrast thresholds ( ) as a function of external noise levels ( ) at a given performance
level ( ) (**Fig. 2C**):

[Equation 4]

$$C_{\tau,d'}^{\mathrm{pred}} = \left\{ \frac{d'^2\left(\sigma_{\mathrm{ext}}^{2\gamma} + \sigma_{\mathrm{add}}^2\right)}{\beta^{2\gamma}} \right\}^{\frac{1}{2\gamma}}$$

The model includes three key free parameters: (gain), a scaling factor reflecting how well
the observer's template matches the signal; (additive noise), the standard deviation of a
Gaussian noise component independent of the stimulus strength; (nonlinearity), describing
how the template-matched output is amplified. Note that the original full model also included
a multiplicative noise parameter ( ) which reflects how internal noise scales with signal
contrast[34]. By doing a model comparison we found that the reduced model excluding
multiplicative noise outperformed the full model ($p < 0.05$, Bonferroni-corrected) (see Model
Comparison below). Therefore, we proceeded with the more parsimonious three-parameter
model (**Fig. 2C**).

Model Comparison

We evaluated the need to include all four parameters for the perceptual template model by
conducting a nested model comparison between the full model and 15 reduced models,
each systematically excluding one or more parameters. The full model devised and tested by
Lu and Dosher is:

[Equation 5]

$$C_{\tau,d'}^{\text{pred}} = \left\{ \frac{d'^2 \left[\left(1 + N_{\text{mul}}^2\right) \sigma_{\text{ext}}^{2\gamma} + \sigma_{\text{add}}^2 \right]}{\beta^{2\gamma} - N_{\text{mul}}^2 \beta^{2\gamma} d'^2 / 2} \right\}^{\frac{1}{2\gamma}}$$

When excluded from fitting, multiplicative ( ) and additive ( ) internal noise were fixed
at 0, and gain ( ) and nonlinearity ( ) were fixed at 1. We fitted each candidate model to the
seven locations we tested (fovea, horizontal meridian, upper and lower vertical meridian at
4^0 and 8^0 eccentricity). We first excluded models that failed to account for the data (indicated
by negative R-squared), which occurred when gain and/or additive noise were omitted. For
the remaining four models that included both gain and additive noise, we quantified their
goodness of fit by the Bayesian Information Criterion (BIC):

[Equation 6]

$$\text{BIC} = n \log \left(\frac{\text{RSS}}{n} \right) + k \log(n)$$

RSS is the sum of squared residuals specified above, is the number of data, and is the
number of parameters. Lower BIC scores indicate better model fits while penalizing
overfitting. We fitted the full set of perceptual template models to each location group
independently for each observer, allowing all parameters to vary freely across locations
within each location group. Delta BIC was calculated by subtracting the minimum BIC score
from each candidate model's BIC score. Finally, we fitted linear mixed-effects models to
delta BIC values to evaluate the fixed effect of model type with random effects specified
across observers:

[Equation 7]

$$\Delta\text{BIC}_{\text{nested}} = 1 + \text{Model}_{\text{nested}} + (1|\text{Subj})$$

The reduced model excluding multiplicative noise outperformed the full model (p < 0.05,
Bonferroni-corrected). Therefore, we proceeded with the more parsimonious three-
parameter model.

When fitting the perceptual template model, additive noise was calculated in log10 units
because they are in contrast units. Model fitting was implemented by MATLAB's fmincon()
function, which minimized the sum of squared residuals (RSS) between the measured and
predicted thresholds. Prior to squaring, residuals were calculated on log contrast values and
weighted by the number of trials per noise level (   ).

[Equation 8]

701
$$\mathrm{RSS} = \sum_{N} \left(n_N \left(\log C_N^{\mathrm{meas}} - \log C_N^{\mathrm{pred}} \right) \right)^2$$

Parameter bounds were decided according to the literature[9,34]: multiplicative noise (   )
was constrained between 0 to $\overline{}$ (to ensure a positive denominator in equation (4)),
additive noise (   ) ranged from 10^-8 to 1 in log steps, gain ( ) was bounded between 0
to 5, and nonlinearity ( ) ranged from 0 to 5.

To constrain model estimation[33,34], we obtained thresholds (   ) at three performance
levels   =0.77, 1.05, 1.35, where   , corresponding to 65%, 70% and 75%
accuracy ( ), respectively. Only thresholds at 75% correct were shown as TvN functions
(**Fig. 3**). This approach was originally intended to constrain the model by ruling out
substantial reductions in internal multiplicative noise as a plausible mechanism of
performance variation[38]. Although we ultimately adopted a simpler version of the model that
does not require this constraint anymore, we retained the method for consistency with prior
literature.

To examine visual field heterogeneity (eccentricity effect, HVA, and VMA), we grouped
locations according to the effect of interest (see Statistical Analysis-Location groups below),
then fitted perceptual template models to all locations within each location group.

Statistical Analysis

Location groups

At each noise level, we derived thresholds for eccentricity and meridian by combining data
across locations. To represent the horizontal meridian at each eccentricity, we collapsed
trials from the left and right locations and fit a single psychometric function. Thresholds for
the vertical meridian were computed as the mean of the upper and lower vertical meridian.
Thresholds for each eccentricity were computed as the means across the horizontal and
vertical meridian.

We grouped visual field locations to systematically examine the eccentricity effect, HVA, and
VMA. To assess the eccentricity effect, we compared thresholds at the fovea, 4º, and 8º
along the horizontal and vertical meridians separately (i.e., three locations in each group),

allowing us to examine whether the meridian modulates eccentricity-related changes. To
evaluate polar angle asymmetries, we compared thresholds between the horizontal and
vertical meridians (for HVA) and between the lower and upper vertical meridians (for VMA)
at 4º and 8º eccentricities, with four locations included in each group. In addition, we defined
four supplementary groups to isolate each asymmetry at individual eccentricities, enabling
follow-up analyses when needed. Within each group, we fit the perceptual template model to
thresholds across external noise levels to estimate the underlying system-level
computations. Grouping was essential to isolate location-specific computations without
conflating repeated measures. Some locations appeared in multiple groups (e.g., the fovea
appeared in both eccentricity groups), leading to multiple parameter estimates per location.
One-way ANOVAs confirmed that these estimates did not differ across groups (all ps > 0.1),
and all were retained for further analysis.

Linear mixed-effects model

We used linear mixed-effects models for three sets of analyses: (1) verifying the eccentricity
effect and polar angle asymmetries, (2) determining the optimal parameter configuration of
the perceptual template model, and (3) analyzing the spatial pattern of computations. This
approach was chosen for two reasons. First, linear mixed-effects models handle unbalanced
datasets effectively, which was optimal given our mixed design: a between-observer factor
(nine observers were tested at both 4 and 6 cpd, and an addition of three people were tested
only at 6 cpd) and a within-observer factor (three thresholds were derived per location x
noise per observer). Second, linear mixed-effects models account for random effects by
including observer-specific intercepts, capturing individual variability across conditions.

All analyses were conducted in RStudio (v2024.12.1+563) using R. Linear mixed-effects
models were fitted using the lmer() function from the lmerTest package, with maximum
likelihood estimation and degrees of freedom calculated using the Satterthwaite
approximation, which improves accuracy under unequal sample sizes and the inclusion of
random effects. We used the anova() function to perform Type III ANOVA tests for assessing
main effects and interactions, and the emmeans package for post-hoc comparisons with
Bonferroni corrections. Model fit was evaluated using conditional R-squared values, which
reflect both fixed and random effects.

Visual field heterogeneity for performance

Before examining the computations, we first verified that eccentricity effects and polar angle
asymmetries were robust for contrast threshold in the absence of external noise, consistent
with previous studies. We defined "performance" as the intercept threshold—the threshold
measured without external noise— the standard metric in studies without external noise. To
align with conventional performance definitions, where higher values indicate better
performance, we took the absolute value of log-transformed thresholds. We assessed how
performance varied across visual field locations within each location group and whether SF
modulated these variations.

To verify eccentricity effects, we separately modeled performance within each location group
using linear mixed-effects models, with fixed effects for eccentricity, SF, and their interaction:

[Equation 9]

770
$$\text{Perf} = 1 + \text{Eccentricity} * \text{SF} + (1|\text{Subj})$$

To verify HVA and VMA, we fit separate linear mixed-effects models for each asymmetry.
For HVA, we tested the fixed effects of *meridian* (horizontal vs. Vertical meridian),
eccentricity (4º and 8º) and SF and their interactions. For VMA, we modeled the fixed effects
of *location* (lower vs. upper vertical meridian), eccentricity and SF, and interactions among
these factors.

[Equation 10]

777
$$\text{Perf} = 1 + \text{Location} * \text{SF} * \text{Eccentricity} + (1|\text{Subj})$$

Variation and contribution

We investigated how these parameters underlie visual field heterogeneity throughout the
visual field (eccentricity effect, HVA, and VMA) by examining the variation and contribution of
neural computations.

First, to analyze how each computation varies across locations and how these variations
interact with SF, we fitted a linear mixed-effects model to each parameter. To assess
variation across eccentricity, similar to equation (6), the model assesses the interaction
between eccentricity (4º and 8º) and SF (4 and 6 cpd):

[Equation 11]

787
$$\text{Gain} = 1 + \text{Eccentricity} * \text{SF} + (1|\text{Subj})$$

788 To assess variation around polar angle, similar to equation (7), the model assesses the
789 interaction among location (meridian for HVA and location for VMA), SF and eccentricity:

790 [Equation 12]

791
$$\text{Gain} = 1 + \text{Location} * \text{SF} * \text{Eccentricity} + (1|\text{Subj})$$

792 Both models had random intercepts for observers. Regression slopes were constrained to
793 be constant, enabling assessment of overall trends.

794 Then, we examined how each parameter modulates performance. We analyzed the fixed
795 effect of each parameter. Using gain as an example:

796 [Equation 13]

797
$$\text{Perf} = 1 + \text{Gain} + (1|\text{Subj} * \text{SF})$$

This model includes a random intercept for each location- SF combination. Hence, any
observed fixed effect cannot be attributed to differences across locations or SFs. We chose
not to include random slopes because this analysis aims to estimate the overall fixed effect

across locations and SF. The model was fitted to the seven locations that cover the
isoeccentric locations at the tested eccentricity range.

After determining whether each parameter enhanced or impaired performance, we assessed
the independent contribution of each parameter to each performance heterogeneity. First,
we quantified the extent of asymmetry for both performance and parameters using a location
difference index (LDI)—the arithmetic difference between two locations divided by their sum.
This index provides a normalized measure of location variation. By design, the first of the
two values in the LDI calculation was always taken from the location with higher
performance. For example, when comparing the fovea and 8°, both the numerator and
denominator of the LDI were computed as the difference between fovea and 8°. We then
fitted linear mixed-effects models to LDI values of each parameter (e.g., gain) and that of
performance, and the resulting fixed effects indicate contribution of each parameter.

For gain and additive noise, we computed one-tailed empirical p-values (see Bootstrapping
section for definition) because the directionality of their contribution to performance
heterogeneity could be inferred from the directionality of their modulation on performance.
Specifically, if better-performing locations have higher gain, then gain is assumed to
enhance performance. In this case, both the performance LDI and gain LDI would be
positive, and a stronger difference in performance would correspond to a larger difference in
gain—resulting in a positive correlation. Conversely, if a parameter impairs performance
(e.g., additive noise), a negative correlation would be expected. If a parameter does not
modulate performance (e.g., nonlinearity), we evaluated the two-tailed empirical p-value.

[Equation 14]

$$\mathrm{LDI_{Perf}} = 1 + \mathrm{LDI_{Gain}} + (1|\mathrm{SF}) + (1|\mathrm{Subj})$$

Bootstrapping

To derive confidence intervals and assess statistical significance, we performed
bootstrapping beginning at the level of threshold estimation. In each iteration, we resampled
trials with replacement at each location and noise level separately. Psychometric functions
were then fitted to the resampled data to estimate contrast thresholds at each noise level.
These thresholds were subsequently fitted with the perceptual template model within each
location group to estimate the three model parameters. Finally, statistical analyses were
performed across observers to assess modulation, variation, and contribution effects. Both
the resampling and the two-stage model fitting were conducted independently for each
observer. This procedure was repeated 1,000 times. We reported the observed statistics for
all tests (F-values, t-values, effect sizes, slopes) and p-values (p_{obs}) from the non-
bootstrapped data, along with 95% confidence intervals (excluding p-values) from the
bootstrap distribution. For pairwise comparisons and correlation analyses, empirical two-
tailed p-values (p_{emp}) were computed as twice the smaller proportion of bootstrapped
differences (and slopes) greater or less than zero, and empirical Cohen's d was computed
as the median of the bootstrapped differences divided by the standard deviation of those
differences. For correlation analyses assessing contribution effects, the choice between one-
tailed and two-tailed empirical p-values depended on the parameter's modulation effect: if

the parameter significantly modulated performance in a specific direction, a one-tailed p-
value was used; otherwise, a two-tailed p-value was applied.

References

1. Banks, M. S., Sekuler, A. B. & Anderson, S. J. Peripheral spatial vision: limits imposed

by optics, photoreceptors, and receptor pooling. *J. Opt. Soc. Am. A* **8**, 1775–1787

(1991).

2. Bradley, C., Abrams, J. & Geisler, W. S. Retina-V1 model of detectability across the

visual field. *J. Vis.* **14**, (2014).

3. Himmelberg, M. M., Winawer, J. & Carrasco, M. Polar angle asymmetries in visual

perception and neural architecture. *Trends Neurosci.* **46**, 445–458 (2023).

4. Abrams, J., Nizam, A. & Carrasco, M. Isoeccentric locations are not equivalent: the

extent of the vertical meridian asymmetry. *Vision Res.* **52**, 70–78 (2012).

5. Baldwin, A. S., Meese, T. S. & Baker, D. H. The attenuation surface for contrast

sensitivity has the form of a witch's hat within the central visual field. *J. Vis.* **12**, (2012).

6. Jigo, M., Tavdy, D., Himmelberg, M. M. & Carrasco, M. Cortical magnification eliminates

differences in contrast sensitivity across but not around the visual field. *Elife* **12**,

2022.04.27.489757 (2023).

7. Cameron, E. L., Tai, J. C. & Carrasco, M. Covert attention affects the psychometric

function of contrast sensitivity. *Vision Res.* **42**, 949–967 (2002).

8. Kwak, Y., Lu, Z.-L. & Carrasco, M. How the window of visibility varies around polar

angle. *J. Vis.* **24**, 4 (2024).

9. Barbot, A., Xue, S. & Carrasco, M. Asymmetries in visual acuity around the visual field.

J. Vis. **21**, 2 (2021).

10. Talgar, C. P. & Carrasco, M. Vertical meridian asymmetry in spatial resolution: visual

and attentional factors. *Psychon. Bull. Rev.* **9**, 714–722 (2002).

11. Kurzawski, J. W. *et al.* The Bouma law accounts for crowding in 50 observers. *J. Vis.*

23, 6 (2023).

12. Montaser-Kouhsari, L. & Carrasco, M. Perceptual asymmetries are preserved in short-

term memory tasks. *Atten. Percept. Psychophys.* **71**, 1782–1792 (2009).

13. Roux-Sibilon, A., Peyrin, C., Greenwood, J. A. & Goffaux, V. Radial bias in face

identification. *Proc. Biol. Sci.* **290**, 20231118 (2023).

14. Jaeken, B. & Artal, P. Optical quality of emmetropic and myopic eyes in the periphery

measured with high-angular resolution. *Invest. Ophthalmol. Vis. Sci.* **53**, 3405–3413

(2012).

15. Curcio, C. A., Sloan, K. R., Jr, Packer, O., Hendrickson, A. E. & Kalina, R. E.

Distribution of cones in human and monkey retina: individual variability and radial

asymmetry. *Science* **236**, 579–582 (1987).

16. Curcio, C. A., Sloan, K. R., Kalina, R. E. & Hendrickson, A. E. Human photoreceptor

topography. *J. Comp. Neurol.* **292**, 497–523 (1990).

17. Curcio, C. A. & Allen, K. A. Topography of ganglion cells in human retina. *J. Comp.*

Neurol. **300**, 5–25 (1990).

18. Watson, A. B. A formula for human retinal ganglion cell receptive field density as a

function of visual field location. *J. Vis.* **14**, 15 (2014).

19. Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional

magnetic resonance imaging. *Science* **268**, 889–893 (1995).

20. Duncan, R. O. & Boynton, G. M. Cortical magnification within human primary visual

cortex correlates with acuity thresholds. *Neuron* **38**, 659–671 (2003).

21. Silva, M. F. *et al.* Radial asymmetries in population receptive field size and cortical

magnification factor in early visual cortex. *Neuroimage* **167**, 41–52 (2018).

22. Benson, N. C., Kupers, E. R., Barbot, A., Carrasco, M. & Winawer, J. Cortical

magnification in human visual cortex parallels task performance around the visual field.

Elife **10**, (2021).

23. Himmelberg, M. M. *et al.* Cross-dataset reproducibility of human retinotopic maps.

Neuroimage **244**, 118609 (2021).

24. Himmelberg, M. M. *et al.* Comparing retinotopic maps of children and adults reveals a

late-stage change in how V1 samples the visual field. *Nat. Commun.* **14**, 1561 (2023).

25. Himmelberg, M. M., Winawer, J. & Carrasco, M. Linking individual differences in human

primary visual cortex to contrast sensitivity around the visual field. *Nat. Commun.* **13**,

3309 (2022).

26. Kupers, E. R., Carrasco, M. & Winawer, J. Modeling visual performance differences

"around" the visual field: A computational observer approach. *PLoS Comput. Biol.* **15**,

e1007063 (2019).

27. Kupers, E. R., Benson, N. C., Carrasco, M. & Winawer, J. Asymmetries around the

visual field: From retina to cortex to behavior. *PLoS Comput. Biol.* **18**, e1009771 (2022).

28. Virsu, V. & Rovamo, J. Visual resolution, contrast sensitivity, and the cortical

magnification factor. *Exp. Brain Res.* **37**, 475–494 (1979).

29. Watson, A. B. & Solomon, J. A. Model of visual contrast gain control and pattern

masking. *J. Opt. Soc. Am. A Opt. Image Sci. Vis.* **14**, 2379–2391 (1997).

30. Eckstein, M. P., Ahumada, A. J., Jr & Watson, A. B. Visual signal detection in structured

backgrounds. II. Effects of contrast gain control, background variations, and white noise.

J. Opt. Soc. Am. A Opt. Image Sci. Vis. **14**, 2406–2419 (1997).

31. Abbey, C. K. & Eckstein, M. P. Classification images for detection, contrast

discrimination, and identification tasks with a common ideal observer. *J. Vis.* **6**, 335–355

(2006).

32. Sebastian, S., Abrams, J. & Geisler, W. S. Constrained sampling experiments reveal

principles of detection in natural scenes. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5731–

E5740 (2017).

33. Lu, Z. L. & Dosher, B. A. External noise distinguishes attention mechanisms. *Vision*

Res. **38**, 1183–1198 (1998).

34. Lu, Z.-L. & Dosher, B. A. Characterizing observers using external noise and observer

models: assessing internal representations with external noise. *Psychol. Rev.* **115**, 44–

82 (2008).

35. Pelli, D. G. Effects of visual noise.

http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.706.822&rep=rep1&type=pdf

(1981).

36. Singh, V., Burge, J. & Brainard, D. H. Equivalent noise characterization of human

lightness constancy. *Journal of Vision* **22**, 610–610 (2020).

37. Creighton, S. E., Bennett, P. J. & Sekuler, A. B. Contribution of internal noise and

calculation efficiency to face discrimination deficits in older adults. (2023)

doi:10.2139/ssrn.4474233.

38. Dosher, B. A. & Lu, Z. L. Mechanisms of perceptual learning. *Vision Res.* **39**, 3197–

3221 (1999).

39. Movshon, J. A., Thompson, I. D. & Tolhurst, D. J. Spatial and temporal contrast
sensitivity of neurones in areas 17 and 18 of the cat's visual cortex. *J. Physiol.* **283**,
101–120 (1978).

40. Faisal, A. A., Selen, L. P. J. & Wolpert, D. M. Noise in the nervous system. *Nat. Rev.*
Neurosci. **9**, 292–303 (2008).

41. Zohary, E., Shadlen, M. & Newsome, W. Correlated neuronal discharge rate and its
implications for psychophysical performance. *Nature* **370**, 140–143 (1994).

42. Kohn, A., Coen-Cagli, R., Kanitscheider, I. & Pouget, A. Correlations and neuronal
population information. *Annu. Rev. Neurosci.* **39**, 237–256 (2016).

43. Turner, M. H. & Rieke, F. Synaptic rectification controls nonlinear spatial integration of
natural visual inputs. *Neuron* **90**, 1257–1271 (2016).

44. Carandini, M. & Heeger, D. J. Normalization as a canonical neural computation. *Nat.*
Rev. Neurosci. **13**, 51–62 (2011).

45. Mareschal, I., Bex, P. J. & Dakin, S. C. Local motion processing limits fine direction
discrimination in the periphery. *Vision Res.* **48**, 1719–1725 (2008).

46. Wardle, S. G., Bex, P. J., Cass, J. & Alais, D. Stereoacuity in the periphery is limited by
internal noise. *J. Vis.* **12**, 12 (2012).

47. Xue, S., Fernández, A. & Carrasco, M. Featural representation and internal noise
underlie the eccentricity effect in contrast sensitivity. *J. Neurosci.* **44**, (2024).

48. Rijsdijk, J. P., Kroon, J. N. & van der Wildt, G. J. Contrast sensitivity as a function of
position on the retina. *Vision Res.* **20**, 235–241 (1980).

49. Seiple, W., Holopigian, K., Szlyk, J. P. & Wu, C. Multidimensional visual field maps:

relationships among local psychophysical and local electrophysiological measures. *J.*

Rehabil. Res. Dev. **41**, 359–372 (2004).

50. Carrasco, M., Williams, P. E. & Yeshurun, Y. Covert attention increases spatial

resolution with or without masks: support for signal enhancement. *J. Vis.* **2**, 467–479

(2002).

51. Liu, T., Heeger, D. J. & Carrasco, M. Neural correlates of the visual vertical meridian

asymmetry. *J. Vis.* **6**, 1294–1306 (2006).

52. Seiple, W., Holopigian, K., Shnayder, Y. & Szlyk, J. P. Duration thresholds for target

detection and identification in the peripheral visual field. *Optom. Vis. Sci.* **78**, 169–176

(2001).

53. Carrasco, M., McLean, T. L., Katz, S. M. & Frieder, K. S. Feature asymmetries in visual

search: effects of display duration, target eccentricity, orientation and spatial frequency.

Vision Res. **38**, 347–374 (1998).

54. Jigo, M. & Carrasco, M. Differential impact of exogenous and endogenous attention on

the contrast sensitivity function across eccentricity. *J. Vis.* **20**, 11 (2020).

55. Himmelberg, M. M., Winawer, J. & Carrasco, M. Stimulus-dependent contrast sensitivity

asymmetries around the visual field. *J. Vis.* **20**, 18 (2020).

56. Aghajari, S., Vinke, L. N. & Ling, S. Population spatial frequency tuning in human early

visual cortex. *J. Neurophysiol.* **123**, 773–785 (2020).

57. Broderick, W. F., Simoncelli, E. P. & Winawer, J. Mapping spatial frequency preferences

across human primary visual cortex. *J. Vis.* **22**, 3 (2022).

978 58. Schiller, P. H., Finlay, B. L. & Volman, S. F. Quantitative studies of single-cell properties

979   in monkey striate cortex. III. Spatial frequency. *J. Neurophysiol.* **39**, 1334–1351 (1976).

980 59. De Valois, R. L., Albrecht, D. G. & Thorell, L. G. Spatial frequency selectivity of cells in

981   macaque visual cortex. *Vision Res.* **22**, 545–559 (1982).

982 60. Xue, S. & Carrasco, M. Featural representation underlies performance differences

983   around the visual field. *Journal of Vision* **23**, 4771 (2023).

984 61. Rockel, A. J., Hiorns, R. W. & Powell, T. P. The basic uniformity in structure of the

985   neocortex. *Brain* **103**, 221–244 (1980).

986 62. Hubel, D. H. & Wiesel, T. N. Receptive fields of cells in striate cortex of very young,

987   visually inexperienced kittens. *J. Neurophysiol.* **26**, 994–1002 (1963).

988 63. Yu, H.-H. & Rosa, M. G. P. Uniformity and diversity of response properties of neurons in

989   the primary visual cortex: selectivity for orientation, direction of motion, and stimulus

990   size from center to far periphery. *Vis. Neurosci.* **31**, 85–98 (2014).

991 64. Azzopardi, P., Jones, K. E. & Cowey, A. Uneven mapping of magnocellular and

992   parvocellular projections from the lateral geniculate nucleus to the striate cortex in the

993   macaque monkey. *Vision Res.* **39**, 2179–2189 (1999).

994 65. Derrington, A. M. & Lennie, P. Spatial and temporal contrast sensitivities of neurones in

995   lateral geniculate nucleus of macaque. *J. Physiol.* **357**, 219–240 (1984).

996 66. Cohen, M. R. & Maunsell, J. H. R. Attention improves performance primarily by reducing

997   interneuronal correlations. *Nat. Neurosci.* **12**, 1594–1600 (2009).

998 67. Mitchell, J. F., Sundberg, K. A. & Reynolds, J. H. Spatial attention decorrelates intrinsic

999   activity fluctuations in macaque area V4. *Neuron* **63**, 879–888 (2009).

68. Himmelberg, M. M., Kwak, Y. M., Carrasco, M. & Winawer, J. Unpacking the V1 map:

Differential covariation of preferred spatial frequency and cortical magnification across

spatial dimensions. *bioRxivorg* 2025.03.19.644195 (2025)

doi:10.1101/2025.03.19.644195.

69. McDonnell, M. D. & Ward, L. M. The benefits of noise in neural systems: bridging theory

and experiment. *Nat. Rev. Neurosci.* **12**, 415–426 (2011).

70. Bejjanki, V., Beck, J., Lu, Z.-L. & Pouget, A. Perceptual learning as improved

probabilistic inference in early sensory areas. *Nat. Neurosci.* **14**, 642–648 (2011).

71. Previc, F. H. Functional specialization in the lower and upper visual fields in humans: Its

ecological origins and neurophysiological implications. *Behav. Brain Sci.* **13**, 519–542

(1990).

72. Carrasco, M., Roberts, M., Myers, C. & Shukla, L. Visual field asymmetries vary

between children and adults. *Curr. Biol.* **32**, R509–R510 (2022).

73. Carrasco, M., Myers, C. & Roberts, M. Visual field asymmetries vary between

adolescents and adults. *bioRxivorg* (2025) doi:10.1101/2023.03.04.531124.

74. Tünçok, E., Kiorpes, L. & Carrasco, M. Opposite asymmetry in visual perception of

humans and macaques. *Curr. Biol.* (2025) doi:10.1016/j.cub.2024.12.024.

75. Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3. *Perception* **36**, 1–16

(2007).