In vitro and *in vivo* derived macrophages occupy distinct phenotypic states

Stéphane Chevrier[1], Vito R. T. Zanotelli[1,2], Daniel Schulz[1], Mark D. Robinson[1,3], Laurie Ailles[4,5], Michel A.S.Jewett[5], Craig Gedye[6], Bernhard Reis[7], and Bernd Bodenmiller[1,8]

[1] Department of Quantitative Biomedicine, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

[2] Systems Biology PhD Program, Life Science Zürich Graduate School, ETH Zürich and University of Zürich, 8057 Zürich, Switzerland

[3] SIB Swiss Institute of Bioinformatics, University of Zurich, Zurich, Switzerland

[4] Dept. of Medical Biophysics, University of Toronto, Toronto, ON, Canada

[5] Princess Margaret Cancer Center, University Health Network, Toronto ON M5G 1L7, Canada

[6] School of Biomedical Sciences and Pharmacy, University of Newcastle, Hunter Medical Research Institute, NSW, 2305, Australia

[7] Roche Pharma Research and Early Development, Pharmaceutical Sciences, Roche Innovation Center Basel, F. Hoffmann-La Roche Ltd, Grenzacherstrasse 124, 4070 Basel, Switzerland

[8] Lead contact

Correspondence should be addressed to B.B. bernd.bodenmiller@imls.uzh.ch

SUMMARY

Monocyte-derived macrophages (MDMs) are widely used to model human macrophage biology *in vitro* and standardized polarization conditions were proposed to recapitulate the different macrophage activation states. Although surface markers specific for distinct MDM populations have been identified, a systematic analysis of surface marker expression on these consensus MDM populations has not previously been reported. Here, we use mass cytometry to perform an in-depth characterization of MDM surface profiles and determine how these markers evolve with time. We also compared the phenotypes found *in vitro* with patient-derived tumor-associated macrophages (TAMs) and found that although MDMs and TAMs shared most markers investigated, the cell-surface signatures markedly differed in

terms of expression levels and combinations. Our high-dimensional, single-cell analyses

clarifies the surface expression profile of the core *in vitro* differentiated macrophage

populations and highlights some limitations of the *in vitro* system to represent the complexity

of *in vivo* polarized tumor-associated macrophage phenotypes. These findings provide new

opportunities to improve the models used to study macrophage biology and give a pathway

to improve our understanding of the *in vivo* complexity of these systems.

KEYWORDS

Monocyte-derived macrophage, M1/M2 polarization, single-cell analysis, mass cytometry.

INTRODUCTION

Macrophages are distributed throughout the body and are involved in a wide variety of tasks

including clearance of senescent cells, angiogenesis, bone resorption, elimination of

allergens, clearance of pathogens, antigen presentation, inflammation, and tissue repair [1–3].

Abnormal macrophage functions can lead to or reinforce pathological conditions, as

observed in metabolic diseases, allergic disorders, autoimmune diseases, and cancer [1,2].

Macrophages have been often segregated into classically activated M1 and alternatively

activated M2 subtypes, which are assumed to constitute two extremes of a continuum [4–6].

M1 macrophages are characterized by the secretion of pro-inflammatory cytokines, reactive

oxygen species, and nitric oxide, which mediate their microbicidal and tumoricidal activities.

M2-like macrophages secrete immunosuppressive cytokines as well as growth and

angiogenic factors and are proposed to be involved in tissue remodeling and resolution of

the inflammation [7].

In vitro models using monocyte-derived macrophages (MDMs) have been developed to

mimic the functional phenotypes found *in vivo*. The polarization stimuli used to generate

MDMs have evolved from the original M1/M2 dichotomous model induced using IFNγ and

IL4, respectively, to include glucocorticoid, immune complexes, and additional cytokines.

These differentiation conditions resulted in better defined subsets such as M2a, M2b, and

M2c [8,9]. The growing numbers of MDM populations have been extensively studied at the

transcriptional and protein levels using microarrays, deep sequenced transcriptome,

proteomics as well as multiplexed cytokine/chemokine secretion assays [10–14]. These studies

reported that macrophage polarizations cover a spectrum of activation states [5,15]. To avoid

confusion in the terminology used and to allow better comparison of experimental data

across laboratories and disciplines, standardized *in vitro* polarization conditions and

nomenclatures were proposed [16]. The core polarizations that cover the spectrum of

macrophage activation states were defined, and the authors proposed that these activation

states could serve as a framework to map *ex vivo* isolated macrophages [16]. Cell surface

marker signatures are a key tool for mapping and analyzing macrophage types from

diseased tissues within a clinical context. Although previous studies have identified surface

markers specific for distinct MDM populations, no systematic analyses have been performed

to characterize and catalogue the surface expression profile of those core MDM activation

states, thus hindering their further assessment and utility.

Unlike most immune cell types, macrophage subsets cannot be defined based on the

expression or the lack of expression of a few well-defined markers [16–18]. Due to their inherent

plasticity, a large number of markers are required to distinguish the phenotypic spaces

occupied by macrophage subsets. Mass cytometry, which makes use of metal-tagged

antibodies, enables the simultaneous detection of up to 40 protein readouts at the single-cell

level with limited signal overlap [19–21]. Moreover, this method is not affected by the auto-

fluorescence of myeloid cells, which can lead to artifacts in classic fluorescence flow

cytometry. With the ability to measure dozens of samples and millions of cells in a short

time, mass cytometry is a method-of-choice to directly profile and compare the surface

expression profiles of differentially activated macrophages.

To systematically analyze the surface expression profiles of MDMs generated under

standard conditions and in a time resolved manner [16], we employed an adapted version of a

36-antibody panel used to characterize tumor associated macrophage in the context of renal

carcinoma[22] in a mass cytometry analysis. We then directly compared the surface

phenotypes of MDMs with the phenotypes of tumor-associated macrophages (TAMs) from

ascites isolated from ovarian cancer patients. Although the surface markers identified by

analysis of MDMs were detected on TAMs, we observed little overlap in the actual surface

marker expression patterns. Our data highlight the complex phenotypic diversity of

macrophages found *in vivo*, emphasizing the limitations of the *in vitro* MDM model to

recapitulate *in vivo* macrophage states.

RESULTS

MDM characterization using mass cytometry

In order to develop an antibody panel that best discriminate blood monocytes (M0),

immature macrophages (M5), and populations obtained by stimulation with the nine MDM

conditions proposed by Murray and colleagues [16] (Figure 1A), we selected markers based

on a fluorescent surface screening previously performed on those populations [22]. The final

antibody panel (Table 1) was used to analyze simultaneously the expression of 36 markers

on the surface of eleven populations of monocytes and monocyte derived macrophages from

four healthy donors. In order to minimize technical variability during staining and to measure

all samples simultaneously, the samples were barcoded using a 60-sample mass-tag cell

barcoding method with a doublet filtering scheme [23,24].

Using the nonlinear dimensionality reduction approach based on the t-distributed stochastic

neighbor embedding (t-SNE) [25,26], the high-dimensional single-cell data were visualized in

two dimensions (Figure 1B). This approach showed that monocytes, M(IL4), M(Immune

complex (IC)), and M(IFNγ) were clearly separated from all other populations. M(LPS) and

M(LPS-IFNγ), although distinct, were localized in close proximity. M5, M(Mock), M(IL10),

M(Glucocorticoid (GC)), and M(GC-TGFβ) showed partial overlap. This configuration was

consistent for each of the four donors (Figure S1A).

Unique patterns for each marker assessed in this study were observed in the t-SNE map

(Figure 1C). The canonical macrophage markers CD68, CD14, and CD13 were expressed in

all populations, yet at different abundances. CD11b was high only in monocytes and

downregulated upon *in vitro* culture. As expected from the original optimization of the

antibody panel, we identified markers that were population specific, such as CD209 in

M(IL4), CD87 in M(IC), and CD64 in M(IFNγ). Most markers were shared across a range of

conditions, however; and this was particularly obvious for costimulatory molecules. CD40

was induced both in M(LPS) and M(LPS-IFNγ) culture conditions. CD155, Slamf7, and

CD274 (PD-L1) were expressed in the M1 subsets (LPS, LPS-IFNγ, and IFNγ) and in the

M(IC). CD273 (PD-L2) and CD86 were induced in M1 and M(IC) populations and showed

some expression in the M(IL4). Similarly, the adhesion molecules CD54 (ICAM-1) and

CD82, the cyclic ADP ribose hydrolase CD38, and the chemokine receptor CCR7 were also

strongly induced in the M1 and the M(IC) subsets. The IL3 receptor CD123 was expressed

on both M1 and M(IC) populations and was observed at the highest level in M(IL4). The

scavenger receptors CD163 and CD36, the chemokine receptor CXCR4, the Fc receptor

CD32, the VEGF co-receptor CD304, and the complement receptor CD88 had strong

expression in the M2-like populations polarized with CSF-1, IL4, IL10, GC, and GC-TGFβ.

The CD4 activation marker had an expression pattern overlapping with typical M1 and M2

subsets and showed a high induction upon IL4, IFNγ, and IL10 treatment. The data were

consistent across the four samples analyzed (Figure S1B). Despite the heterogeneity of

monocytes used to generate macrophages, all the markers used in the analysis showed a

unimodal (albeit sometimes broad) distribution on all the MDM populations investigated
(Figure S1C).

Hierarchical clustering showed that despite the biological variation, most MDM donor
replicates clustered based on the activator used. To systematically identify the marker
signature that best characterizes the different MDM populations, we used marker enrichment
modelling, a metric that takes into account the median intensity and the distribution of the
markers in each population [27]. The marker signature identified for each subset based on this
approach was largely consistent with the observation made based on the t-SNE map, as
shown in Figure 1D. Taken together, our data showed that the antibody panel used captures
MDM heterogeneity and establishes condition-specific surface expression signatures, which
can be used to compare samples across studies.

Evolution of MDM surface marker expression over time

To understand better the dynamic nature of marker expression over time, we harvested
MDMs 6h, 12h, 24h, 36h and 48h after activation with the different stimuli. Displaying the
cells from the different time points for each stimulation on a t-SNE map revealed that the *in*
vitro differentiation process is dynamic and phenotypic changes could be observed
throughout the entire kinetic (Figure 2A, Figure S2A). A systematic analysis of the log2 fold
change of each marker at the different time points upon stimulations compared to the initial
state (M5) revealed that most markers were induced upon stimulation whereas only a
minority of markers were repressed (Figure 2B). For most markers, the induction or the
repression was reinforced throughout the kinetic and the highest or lowest level was reached
after 48h. The markers that best defined the M5 signature (Figure 1D), including CD4, CD14,
CD163, but also CD32 or CD206 were among the markers that were repressed upon
stimulation (Figure 2B). The repression of those markers was occurring mostly on the M1
populations (LPS, LPS-IFNγ, and IFNγ) and the M(IC) while their expression levels were
maintained or even slightly increased on the M2-like subsets. This was consistent with the
fact that the CSF-1 cytokine used to generate the M5, triggers an anti-inflammatory M2-like
response and therefore many M2 specific markers were already induced on M5 cells. For
most markers induced during the kinetic, including CD64, CD81, CD38, CD40, CD197,
CD54, CD274, CD273, CD166 or CD82, the expression level showed a progressive increase
over at least the first 24h of activation. One notable exception was Slamf7, which was
already expressed at its maximum level only 6h after LPS or LPS-IFNγ stimulation and 12h
after IC activation (Figure 2B).

When looking specifically at cells activated with LPS, LPS-IFNγ and IFNγ alone, both the t-SNE and the fold change analysis suggested strong similarities between cells activated with LPS and with LPS-IFNγ during the first 12h (Figure 2A-B). A further analysis based on diffusion mapping, an algorithm that aligns cells along putative developmental trajectories confirmed that monocytes activated with LPS and LPS-IFNγ followed the same trajectory during the first time points before adopting different differentiation paths in the later time points (Figure 2C). Conversely, cells activated with IFNγ followed a different trajectory from the beginning (Figure 2C). This suggested a dominant role of LPS at earlier time points, while the effect of IFNγ in presence of LPS was mostly observed after 24h. The presence of IFNγ was critical to induce a high expression level of costimulatory molecules including CD86, CD273 and CD40 and to maintain a long lasting induction of Slamf7 on LPS treated cells (Figure 2C). When looking systematically at the evolution of marker expression at the different time points, we observed that different canonical macrophage markers, including CD14, CD16 or HLA-DR were indeed expressed at the same levels on cells activated with LPS and LPS-IFNγ for up to 24h before diverging at the later time points (Figure 2D, first column). This approach also revealed different regulatory mechanisms for the different markers investigated. LPS was a more potent inducer of CD82, CD166 and CD155 than IFNγ or the combination of both (Figure 2D, second column). On the other hand, IFNγ had a stronger effect on the induction of CD273, CD64 or CD81 (Figure 2D, third column) compared to LPS alone or in combination with IFNγ. Finally, for most of the costimulatory molecules, including CD274, CD40, SLAMF7, CD86, but also for the ecto-enzyme CD38 and the chemokine receptor CCR7 (CD197), the combination of both LPS and IFNγ was required for a sustained, high expression level (Figure 2D, fourth and fifth columns). Together, these analyses enabled to get a better understanding on the constant evolution of marker expression over time and demonstrated how LSP, IFNγ and the combination thereof differentially regulated the expression of key macrophage markers.

MDMs and ascites associated macrophages have distinct phenotypes

The transcriptional and phenotypic signatures of different MDM populations were used to map myeloid cells isolated from human pathological conditions [5,28]. However, understanding to which extend MDM populations can serve as a framework to map and classify macrophages isolated from patient samples is still an open question. We used our comprehensive MDM dataset to directly assess how close *in vitro* derived macrophages resemble to *in vivo* isolated tumor associated macrophages. In order to avoid tissue dissociation, a step susceptible to modify the native phenotype found in vivo, we focused this comparison on ascites from advanced ovarian cancer patients, known to contain high

amount of macrophages. Since ascites were stored as viably frozen cells, we compared the

profile of MDMs before and after freezing and we observed no significant changes (Figure

S2 A-B).

We analyzed the phenotypes of macrophages isolated from ascites of nine ovarian patients

simultaneously with the MDM populations. To allow the detection of T cells and

macrophages, the panel was complemented with antibodies to CD3 and CD45. TAMs were

identified as CD45$^+$ CD3$^-$ CD68$^+$ CD64$^+$ live cells and T cells as CD45$^+$ CD3$^+$ (Figure S2C).

The phenotypes of MDMs, TAMs, and T cells were compared based on a t-SNE analysis on

those populations. This approach showed that a small proportion of macrophages had an

antibody staining signature similar to that of M0 (blood monocytes) consistent with the fact

that tumors are vascularized and likely contain circulating monocytes (Figure 3A). However,

when we compared the signatures of ascites associated macrophages with those of the

MDMs, no overlap was observed. This indicated that macrophages differentiated *in vitro* do

not recapitulate the cell-surface protein phenotypes of macrophages found in the ascites

samples (Figure 3A).

The absence of overlap between TAMs and MDMs on the t-SNE map could be explained if

the markers chosen based on *in vitro* differentiated macrophages were not expressed on the

surface of TAMs. We therefore compared the expression levels of all the markers used in

our study, on the surface of T cells, monocytes, MDMs, and macrophages. For this

comparison, data on all the MDM populations were pooled as were data on macrophages

from the nine ascites samples. As expected, T cells were positive only for CD4, CD81,

CXCR4, and HLA-ABC (Figure 3B). The monocytes expressed high levels of CD11b, CD13,

CD14, and CD36 and intermediate levels of CD68, CD64, and HLA-DR. The pooled MDM

population showed detectable expression for all the markers investigated except CD11b.

Strikingly, the expression pattern of the different markers on the surface of the TAM

population obtained by pooling the nine samples was similar to the expression pattern

observed for MDM populations. In particular, the expression patterns of CD166, CD32,

CD38, CD81, CD204, HLA-ABC, and HLA-DR were almost indistinguishable between ex

vivo isolated macrophages and MDMs. Some markers, such as CD4 and CD64 were

expressed at higher levels on ascites associated macrophages than on MDMs. Although

some markers, including CD206, CD163, and CD169, were expressed at lower levels on ex

vivo isolated macrophages than on MDMs, the expression was still higher than on

monocytes, and a proportion of cells expressed a level close to the level observed on

MDMs. Costimulatory molecules (such as CD40, CD274, and Slamf7) and chemokine and

cytokine receptors (such as CCR7 and CD119) were significantly reduced on ex vivo

isolated macrophages compared to MDMs. These data show that the markers analyzed
were appropriate for the characterization of *in vivo* macrophage phenotypes.

As an alternative approach to better understand the absence of surface marker phenotype
overlap between TAMs and MDMs and to exclude the possibility that t-SNE leads to
misleading conclusions, we identified ex vivo isolated macrophages and MDM subsets using
the unsupervised graph-based clustering algorithm PhenoGraph [30]. Consistent with t-SNE-
based observations, the ex vivo isolated macrophages - and MDM-containing clusters were
mutually exclusive, whereas one cluster contained both TAMs and monocytes (Figure 3D,
middle panel, cluster 10). Ex vivo isolated macrophages subsets were characterized by
subtle differences in a wide variety of surface markers with only a few markers reaching a
high expression level. In contrast, the MDM subsets were characterized by clear differences
in marker expression with multiple markers expressed at high levels. Ascites associated
clusters were characterized by different combinations of markers that could not individually
be associated with specific MDM populations.
To assess the similarity between ascites associated cells and *in vitro* derived macrophages,
we developed a machine learning approach comparing the marker signature of the cells
found *in vivo* with the one generated *in vitro* based on a random forest classifier. Using a
wheel plot, we visualized each cell based on its probability of being assigned to each of the
MDM populations used to train the model (Figure 2E). Apart from the cells activated with
glucocorticoid alone or in presence of TGFB, which are difficult to distinguish, cells from
each MDM subset were located in close vicinity to the correct node. This was further
confirmed when looking at the frequency of cells assigned to the different cell types based
on a 40% probability threshold (Figure 2F). In comparison, macrophages isolated from
ascites formed a range of poorly defined cells, originating from the M0 monocyte subset, but
rapidly losing any connections with *in vitro* derived cell subsets (Figure 2E). Consistently,
most of the macrophages from ascites samples were assigned to the monocyte subset, or
left unassigned based on a 40% probability threshold (Figure 3F). Thus, our data show that
the markers identified based on MDM populations do capture ex vivo macrophage diversity
but are expressed in different combinations and often at a lower level leading to an absence
of phenotype overlap between ex vivo macrophages and MDM subsets.

DISCUSSION

Macrophages are highly plastic cells that respond to stimuli present in the local environment
by adopting a continuum of functional phenotypes [3,31]. To comprehend this complexity in
humans, numerous studies over the last two decades have relied on the MDM *in vitro*

system and the M1/M2 conceptual framework (Mills et al., 2000; Martinez and Gordon,
2014). Although this system has been used to evaluate transcriptional networks and
chemokine and cytokine profiles of MDMs, no studies have systematically investigated the
surface expression profiles of MDM populations, a necessary step to determine how the *in*
vitro system relates to macrophages *in vivo*.

We used a systematic approach to characterize the cell-surface phenotypes of the nine core
MDM populations proposed by Murray and colleagues (Murray et al., 2014) based on a
modified version of a recently developed mass cytometry antibody panel [22]. Despite
differences in culture protocols, our results were mostly consistent with previous reports, with
features such as an increased expression of CD11b and CD206 on M(IL4) and an
upregulation of CD64, Slamf7, and CD86 on M(IFNγ) as previously described [10,33]. Our
strategy extended the surface expression signature to a larger panel of markers and to more
MDM populations and was consistent across different donors.

Since most markers included in the panel have well-established biological functions, this
characterization furthered our understanding of the functional differences between the MDM
subsets. In a time course analysis, we observed that LPS has a dominant effect on IFNγ
during MDM activation and that cells activated with LPS alone or in presence of IFNγ
displayed strongly overlapping phenotypes up to 12 hours after activation. At later time
points, these two MDM subsets still share many features. Our approach also showed how
LPS or IFNγ alone or in combination differentially controls the expression of costimulatory
molecules such as CD40, Slamf7, CD273 and CD274, highlighting their complex regulation
during macrophage activation. Although MDM activation by immune complexes was
previously associated with an anti-inflammatory, M2-like phenotype [16], our data suggest that
these cells share surface features with pro-inflammatory cells including induction of the
costimulatory molecules CD155, CD274, and Slamf7 and expression of CD38, CD54, CD82,
and CD197.

We then took advantage of the newly defined MDM signatures to determine whether MDM
phenotypes could be used as a framework to classify *ex vivo* isolated TAM populations [22].
We compared the cell-surface phenotypes of MDM populations with TAMs isolated from
ccRCC tumors from 20 patients using the nonlinear dimensionality reduction algorithm t-
SNE, which enables the visualization of high-dimensional single-cell data on a 2D map that
groups cells based on their similarity, and the cluster algorithm PhenoGraph. This approach
showed that MDM populations failed to reproduce the phenotypes of TAMs found *in vivo,*
and our attempt to use machine learning to classify ascites associated macrophage into the

different MDM subsets showed that ex vivo macrophages could not be classified into any of

the populations derived *in vitro*.

Due to the strictly controlled culture conditions, involving the presence of only one or two

cytokines, it is perhaps not surprising that *in vitro* populations did not mimic the phenotypes

of macrophages found in the very complex *in vivo* environment where cells are

simultaneously exposed to a multitude of stimulations and where various feedback

mechanisms are at work. This likely explains the fact that many markers, especially

costimulatory molecules and chemokine receptors, were induced at higher levels in MDMs

compared to TAMs.

Comparing macrophages activated *in vitro* with macrophages isolated *ex vivo* is inherently

limited by the fact that samples are treated in ways that can potentially prevent a direct

comparison. To circumvent this issue, we focused here on macrophages isolated from

ascites samples, which do not require any tissue dissociation, susceptible to affect marker

expression due to the enzymatic treatment and to further stimulate macrophages due to the

presence of dying cells. A limitation of our study is that we only analyzed surface markers,

whose expression level is more likely to change compared to more hardwired markers, such

as key transcription factors or even cytokine secretion. The detection of such markers is

limited by antibody availability, the sensitivity of the system, and the need to reactivate the

cells, all of which can influence results.

Although the MDM system has proven valuable in the study of macrophage biology over the

past decades, our study highlights the limitation of this system. A better understanding of

macrophages present in pathological conditions requires a deeper characterization based on

direct analysis of *ex vivo* isolated cells. The advent of mass cytometry, which allows

simultaneous analysis of multiple functional markers, has enabled the characterization of cell

subsets on a substantial scale. Combining mass cytometry with other high-dimensional

analyses, such as RNA sequencing or multiplexed cytokine assays, has already started to

uncover biological features of macrophage subsets in complex tissues [22,34]. More effort in

this direction is critical to better understand the role of macrophages in various pathological

conditions and to facilitate the development of new therapeutic treatments.

MATERIALS AND METHODS

Monocyte isolation and macrophage generation

Buffy coats from healthy donors were obtained from the Zurich Blood Transfusion Service.

PBMCs isolation was performed by histopaque (Sigma Aldrich) density gradient

centrifugation, and monocytes were isolated by MACS using the pan monocyte isolation kit

(Miltenyi Biotech) according to manufacturer's instructions. Monocytes (M0) were

differentiated into immature macrophages (M5) by culture for 5 days in Nunc UpCell^{TM} 6-well

plates in presence of 30 ng/ml M-CSF (PeproTech). Cells were subsequently polarized for

the indicated amount of time in fresh medium complemented with the following stimuli: 20

ng/ml IL4 (PeproTech), 10 ng/ml IL10 (PeproTech), 10 µg/ml IC (Sigma Aldrich), 40 ng/ml of

the GC dexomethasone (Sigma Aldrich), 40 ng/ml GC (Sigma Aldrich) and 10 ng/ml TGFβ

(CST), 100 ng/ml LPS (Sigma Aldrich), 100 ng/ml LPS (Sigma Aldrich) and 100 ng/ml IFNγ

(PeproTech), and 100 mg/ml IFNγ (PeproTech). Mock-treated cells were cultured for the

same amount of time with fresh culture medium only. After culture, cells were harvested by

temperature reduction. For mass cytometry analysis, cells were stained for viability with 10

µM cisplatin (Enzo Life Sciences) in a 1-min pulse before quenching with 10% FBS as

previously described [35]. Cells were then fixed with 1.6% paraformaldehyde (Electron

Microscopy Sciences) for 10 min at room temperature. Cells were then stored at -80 °C.

381   Tumor sample preparation

Primary ascites samples from ovarian cancer patients were obtained from the xxx from

consenting patients under University Health Network Research Ethics Board approval.

Tumors were graded based on were histological analysis by a pathologist (reported in Table

S3). Single-cell were collected by centrifugation, resuspended in D-SDCM and counted with

a hemocytometer. Cells were then frozen in 10% DMSO in D-SDCM complemented with

10% FBS. Cryopreserved cells were resuscitated for mass cytometry analyses by rapid

thawing and slow dilution. Cells were stained for viability with 10 µM cisplatin (Enzo Life

Sciences) in a 1-min pulse before quenching with 10% FBS as previously described [35]. Cells

were then fixed with 1.6% paraformaldehyde (Electron Microscopy Sciences) for 10 min at

room temperature. Cells were then stored at -80 °C.

393   Fluorescence cell barcoding and antibody screening

Fixed cells were washed in 0.03% saponin in PBS and incubated in PBS containing 0, 0.01,

0.1, or 1 µg/ml Alexa Fluor 700 carboxylic acid, succinimidyl ester (Molecular Probes) and 0,

0.1, or 10 µg/ml of Alexa Fluor 750 carboxylic acid, succinimidyl ester (Molecular Probes) for

15 min on ice, enabling the barcoding of up to 16 samples [37]. Cells were then washed twice

with 0.03% saponin in PBS, and the samples were pooled. The combined samples were

screened with the BioLegend human cell screening kit containing 342 pre-titrated PE-

conjugated antibodies arrayed on four 96-well plates according to manufacturer's instructions. To avoid nonspecific staining, cells were first incubated for 10 min at 4 °C with human Fc-receptor (FcR) blocking reagent (Miltenyi Biotech). Cells were washed, filtered, and resuspended at a density of 4.0 x 10^6 cells/ml. Approximately 2 x 10^5 cells were added to each well of the BioLegend plates, and samples were incubated for 20 min at room temperature for antibody staining. After staining, cells were washed and fixed. Immediately after staining, data were acquired on a Canto II (BD Biosciences) using the auto sampler. Data were analyzed in Cytobank 4.10 as previously described [38].

Mass cytometry barcoding

To ensure homogenous staining, 0.5 x 10^6 cells from each sample were barcoded prior to cell staining and mass cytometry analyses. For this purpose, cells were labeled with a unique combination of four of eight barcoding reagents, enabling the simultaneous analysis of up to 60 samples as previously described [23]. Six palladium isotopes (^{102}Pd, ^{1042}Pd, ^{105}Pd, ^{106}Pd, ^{108}Pd, and ^{110}Pd, Fludigm) were conjugated to bromoacetamidobenzyl-EDTA (BABE) and two indium isotopes (^{113}In and ^{115}In, Fludigm) were conjugated to 1,4,7,10-tetraazacy-clododecane-1,4,7-tris–acetic acid 10-maleimide ethylacetamide (mDOTA) following standard procedures [39]. Mass tag barcoding reagents were titrated to ensure an equivalent staining for each reagent; final concentrations were between 50 nM and 200 nM. Cells were barcoded using the transient partial permeabilization protocol described by Behbehani and colleagues [40]. Briefly, approximately 0.5 x 10^6 fixed cells were loaded into the appropriate number of wells in a 96-well plate. Cells were washed with 0.03% saponin in PBS and incubated for 30 min with 200 µl of mass tag barcoding reagent. After staining, cells were washed twice with 0.03% saponin in PBS and twice with cell staining medium (CSM, PBS with 0.5% bovine serum albumin and 0.02% sodium azide). Samples were then pooled for subsequent cell staining.

Antibodies and antibody labeling

Provider, clone, and metal tag of each antibody used in this study are listed in Table S1. Antibodies were labeled with the indicated metal tag using the MaxPAR antibody conjugation kit (Fluidigm). After metal conjugation, the concentration of each antibody was assessed using the Nanodrop (Thermo Scientific) and adjusted to 200 µg/ml in Candor Antibody Stabilizer. Conjugated antibodies were titrated to determine the optimal concentration for use. All antibodies used in this study were managed using the cloud-based platform AirLab [41].

Antibody staining and data acquisition

After barcoding, pooled cells were incubated with FcR blocking reagent (Miltenyi Biotech) for

10 min at 4 °C. Samples were stained with 200 µl of the antibody panel per 10^7 cells for 30

min at 4 °C. Cells were washed twice in CSM and resuspended in 1 ml of nucleic acid Ir-

Intercalator (Fluidigm) in 1.6% PFA/PBS for 1 h at room temperature or overnight at 4 °C.

Cells were then washed twice in PBS and twice in water. Cells were then diluted to 0.5 x 10^6

cells/ml in water containing 10% of EQ™ Four Element Calibration Beads (Fluidigm). Data

were acquired on a CyTOF 2 mass cytometer, using instrument-based dual-count calibration

and noise reduction and randomization, and cells were selected based on event length

between 10 and 75. Exported flow cytometry standard (FCS) files were concatenated using

the FCS concatenation tool from Cytobank and normalized using the executable MATLAB

version of the Normalizer tool [42]. Individual samples were debarcoded using the executable

MATLAB version of the single cell debarcoder tool [23].

Data analysis

FCS files were uploaded into Cytobank, populations of interest were manually gated, and

events of interest were exported as FCS files. For downstream analysis, FCS files were

loaded into R [43]. Signal intensities for each channel were arcsinh transformed with a cofactor

of 5 (x_transf = asinh(x/5)). To visualize the high-dimensional data in two dimensions, the t-

SNE algorithm was applied on data from 2000 random cells from each sample. The total cell

population was used when less than 2000 cells were available. The R t-SNE package for

Barnes-Hut implementation of t-SNE was used [44]. Data were displayed using the ggplot2 R

package [45]. To visualize marker expression analyses on t-SNE map, the top percentile was

excluded and the maximum intensity was defined as the 99^{th} percentile. The data from all

samples were divided by this value leading to signal intensities ranging between 0 and 1 for

each channel. For hierarchical clustering, pairwise distances between samples were

calculated using the Spearman correlation. Dendrograms were generated using Ward's

method and arranged according to the 'order.optimal' function of the 'cba' R package using

optimal leaf ordering [46]. Heatmaps were displayed in R using the heatmap.2 function form

the gplot package. For single-cell clustering analysis, the Python implementation of

PhenoGraph was run on all samples simultaneously with the parameter *k* defining the

number of nearest neighbors set to 30 [30]. For single cell classification, a random forest

classifier (RandomForest R package) was trained on 3,000 cells of each MDM subsets using

the default parameters (500 trees and 6 variables tried at each split). The remaining cells of

the MDM populations, as well as the total populations of macrophages and T cells found in

the ascites samples were classified based on this model. Probability assignment of each cell

was visualized on a wheel plot (Radviz R package).

REFERENCES

1. Murray, P. J. & Wynn, T. a. Protective and pathogenic functions of macrophage subsets. *Nat. Rev. Immunol.* **11,** 723–37 (2011).

2. Joyce, J. a & Pollard, J. W. Microenvironmental regulation of metastasis. *Nat. Rev. Cancer* **9,** 239–52 (2009).

3. Okabe, Y. & Medzhitov, R. Tissue biology perspective on macrophages. *Nat. Immunol.* **17,** 9–17 (2015).

4. Martinez, F. O. & Gordon, S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep.* **6,** 13 (2014).

5. Xue, J. *et al.* Transcriptome-Based Network Analysis Reveals a Spectrum Model of Human Macrophage Activation. *Immunity* **40,** 274–288 (2014).

6. Biswas, S. & Mantovani, A. Macrophage plasticity and interaction with lymphocyte subsets: cancer as a paradigm. *Nat. Immunol.* **11,** 889–896 (2010).

7. Sica, A. & Mantovani, A. Macrophage plasticity and polarization: in vivo veritas. *J. Clin. Invest.* **122,** 787–795 (2012).

8. Mantovani, A. *et al.* The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol.* **25,** 677–86 (2004).

9. Stein, M., Keshav, S., Harris, N. & Gordon, S. Interleukin 4 potently enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. *J. Exp. Med.* **176,** 287–92 (1992).

10. Beyer, M. *et al.* High-resolution transcriptome of human macrophages. *PLoS One* **7,** e45466 (2012).

11. Martinez, F. O., Gordon, S., Locati, M. & Mantovani, a. Transcriptional Profiling of the Human Monocyte-to-Macrophage Differentiation and Polarization: New Molecules and Patterns of Gene Expression. *J. Immunol.* **177,** 7303–7311 (2006).

12. Eligini, S. *et al.* Human monocyte-derived macrophages are heterogenous: Proteomic profile of different phenotypes. *J. Proteomics* **124,** 112–123 (2015).

13. Lu, Y. *et al.* Highly multiplexed profiling of single-cell effector functions reveals deep functional heterogeneity in response to pathogenic ligands. *Proc. Natl. Acad. Sci.* 201416756 (2015). doi:10.1073/pnas.1416756112

14. Roussel, M. *et al.* Mass cytometry deep phenotyping of human mononuclear phagocytes and myeloid-derived suppressor cells from human blood and bone marrow. *J. Leukoc. Biol.* **102,** jlb.5MA1116-457R (2017).

15. Ginhoux, F., Schultze, J. L., Murray, P. J., Ochando, J. & Biswas, S. K. New insights into the multidimensional concept of macrophage ontogeny, activation and function. *Nat. Immunol.* **17,** 34–40 (2015).

510   16.   Murray, P. J. *et al.* Macrophage Activation and Polarization: Nomenclature and
511   Experimental Guidelines. *Immunity* **41,** 14–20 (2014).
512   17.   Noy, R. & Pollard, J. W. Tumor-Associated Macrophages: From Mechanisms to
513   Therapy. *Immunity* **41,** 49–61 (2014).
514   18.   Quatromoni, J. G. & Eruslanov, E. Tumor-associated macrophages: function,
515   phenotype, and link to prognosis in human lung cancer. *Am. J. Transl. Res.* **4,** 376–89
516   (2012).
517   19.   Ornatsky, O. *et al.* Highly multiparametric analysis by mass cytometry. *J. Immunol.*
518   *Methods* **361,** 1–20 (2010).
519   20.   Bandura, D. R. *et al.* Mass cytometry: technique for real time single cell multitarget
520   immunoassay based on inductively coupled plasma time-of-flight mass spectrometry.
521   *Anal. Chem.* **81,** 6813–22 (2009).
522   21.   Chevrier, S. *et al.* Compensation of Signal Spillover in Suspension and Imaging Mass
523   Cytometry. *Cell Syst.* (2018). doi:10.1016/j.cels.2018.02.010
524   22.   Chevrier, S. *et al.* An Immune Atlas of Clear Cell Renal Cell Carcinoma. *Cell* **169,**
525   736–749.e18 (2017).
526   23.   Zunder, E. R. *et al.* Palladium-based mass tag cell barcoding with a doublet-filtering
527   scheme and single-cell deconvolution algorithm. *Nat. Protoc.* **10,** 316–333 (2015).
528   24.   Bodenmiller, B. *et al.* Multiplexed mass cytometry profiling of cellular states perturbed
529   by small-molecule regulators. *Nat. Biotechnol.* **30,** 858–67 (2012).
530   25.   Amir, E. D. *et al.* viSNE enables visualization of high dimensional single-cell data and
531   reveals phenotypic heterogeneity of leukemia. *Nat. Biotechnol.* **31,** 545–52 (2013).
532   26.   Maaten, L. Van Der & Hinton, G. Visualizing Data using t-SNE. *J. Mach. Learn. Res.*
533   **9,** 2579–2605 (2008).
534   27.   E Diggins, K., R Greenplate, A., Leelatian, N., E Wogsland, C. & M Irish, J.
535   Characterizing cell subsets using marker enrichment modeling. *Nat. Methods* **14,** 1–9
536   (2017).
537   28.   Sander, J. *et al.* Cellular Differentiation of Human Monocytes Is Regulated by Time-
538   Dependent Interleukin-4 Signaling and the Transcriptional Regulator NCOR2.
539   *Immunity* **47,** 1051–1066 (2017).
540   29.   Hollmén, M., Roudnicky, F., Karaman, S. & Detmar, M. Characterization of
541   macrophage--cancer cell crosstalk in estrogen receptor positive and triple-negative
542   breast cancer. *Sci. Rep.* **5,** 9188 (2015).
543   30.   Levine, J. H. *et al.* Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like
544   Cells that Correlate with Prognosis. *Cell* 1–14 (2015). doi:10.1016/j.cell.2015.05.047
545   31.   Glass, C. K. & Natoli, G. Molecular control of activation and priming in macrophages.
546   *Nat. Immunol.* **17,** 26–33 (2015).

32. Mills, C. D., Kincaid, K., Alt, J. M., Heilman, M. J. & Hill, A. M. M-1/M-2 Macrophages
and the Th1/Th2 Paradigm. *J. Immunol.* **164,** 6166–6173 (2000).
33. Rey-Giraud, F., Hafner, M. & Ries, C. H. In vitro generation of monocyte-derived
macrophages under serum-free conditions improves their tumor promoting functions.
PLoS One **7,** e42656 (2012).
34. Lavin, Y. *et al.* Innate Immune Landscape in Early Lung Adenocarcinoma by Paired
Single-Cell Analyses. *Cell* **169,** 750–765.e17 (2017).
35. Fienberg, H. G., Simonds, E. F., Fantl, W. J., Nolan, G. P. & Bodenmiller, B. A
platinum-based covalent viability reagent for single-cell mass cytometry. *Cytometry. A*
81, 467–75 (2012).
36. Gedye, C. & Ailles, L. *Basic Cell Culture Protocols. Methods in molecular biology*
(Clifton, N.J.) **946,** (Humana Press, 2013).
37. Krutzik, P. O. & Nolan, G. P. Fluorescent cell barcoding in flow cytometry allows high-
throughput drug screening and signaling profiling. *Nat. Methods* **3,** 361–8 (2006).
38. Kotecha, N., Krutzik, P. O. & Irish, J. M. Web-based analysis and publication of flow
cytometry experiments. *Curr. Protoc. Cytom.* **Chapter 10,** Unit10.17 (2010).
39. Zivanovic, N., Jacobs, A. & Bodenmiller, B. A Practical Guide to Multiplexed Mass
Cytometry. *Curr. Top. Microbiol. Immunol.* (2013). doi:10.1007/82_2013_335
40. Behbehani, G. K. *et al.* Transient partial permeabilization with saponin enables
cellular barcoding prior to surface marker staining. *Cytom. Part A* n/a-n/a (2014).
doi:10.1002/cyto.a.22573
41. Catena, R., Özcan, A., Jacobs, A., Chevrier, S. & Bodenmiller, B. AirLab: a cloud-
based platform to manage and share antibody-based single-cell research. *Genome*
Biol. **17,** 142 (2016).
42. Finck, R. *et al.* Normalization of mass cytometry data with bead standards. *Cytometry.*
A **83,** 483–94 (2013).
43. R Developement Core Team. R: A Language and Environment for Statistical
Computing. *R Found. Stat. Comput.* **1,** 409 (2015).
44. Krijthe, J. H. Rtsne: T-Distributed Stochastic Neighbor Embedding using a Barnes-Hut
Implementation. (2015).
45. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag New York,
2009).
46. Bar-Joseph, Z., Gifford, D. K. & Jaakkola, T. S. Fast optimal leaf ordering for
hierarchical clustering. *Bioinformatics* **17 Suppl 1,** S22-9 (2001).
581
582

ACKNOWLEDGEMENTS

We are very grateful for the generous donation of tumor samples by patients undergoing
surgery, collected through the University Health Network Biobank and Cooperative Health
Tissue Network. The authors thank the Bodenmiller lab for fruitful discussions and the Cytof
facility (University of Zurich). B.B.'s research is funded by a SNSF R`Equip grant, a SNSF
Assistant Professorship grant, the SystemsX Transfer Project "Friends and Foes", the
SystemsX MetastasiX grant, and by the European Research Council (ERC) under the
European Union's Seventh Framework Program (FP/2007-2013)/ERC Grant Agreement n.
336921. S.C. was funded by a Roche Postdoctoral Fellowship; D.S. by an EMBO fellowship
(ALTF 970-2014) co-funded by the European Commission (LTFCOFUND2013, GA-2013-
609409); C.G. by an RACP CSL Fellowship, a CIHR/KCC SHOPP Fellowship, and an
NHMRC Early Career Fellowship.

596   AUTHOR CONTRIBUTIONS

597   S.C., B.R., and B.B. conceived the study. S.C. performed all experiments with help from D.S.

598   L.A., M.A.S.J., and C.G. S.C., V.R.T.Z., M.D.R. performed data analysis. S.C., L.A.,

599   M.A.S.J., C.G., B.R., and B.B. performed the biological analysis and interpretation. S.C. and

600   B.B. wrote the manuscript with input from all authors.

602   FIGURE LEGENDS

603   Figure 1. Characterization of MDM surface phenotypes by mass cytometry

(A) Monocyte preparation and stimulation to induce differentiation. (B) t-SNE map of M0 and

MDM populations based on the 36 antibodies used for mass cytometry analysis. Cells are

colored by sample types. (C) t-SNE map showing the normalized expression levels of the 36

surface markers across the different populations. Data are from one representative replicate.

(D) Heatmap showing the normalized expression level of the 36 antibodies for the four

replicates of the different populations included in the analysis. Hierarchical clustering was

performed by calculating first the pairwise distance between samples based on the

Spearman correlation, and by clustering based on Ward's method. For each subset, the six

proteins showing the strongest enrichment based on marker enrichment modelling is

indicated on the right. IC, immune complex; GC, glucocorticoid. See also Figure S1 and

Table S1.

616   Figure 2. Kinetic analysis of MDM polarization

(A) t-SNE maps of MDMs activated with the indicated stimuli for the indicated amount of

time. For each stimulation, the cells are colored with a gradient of red corresponding to the

stimulation time shown on a grey background corresponding to all the other cells included in

the analysis. (B) Heatmap showing the log2 fold change in marker expression across time
points and stimuli compared to M5. Markers are clustered based on Spearman correlation
and Ward's linkage. (C) Diffusion mapping based on the first three diffusion components
showing the putative trajectories followed by monocytes activated with the indicated stimuli
over 48h (top-left panel). The other panels are showing the diffusion map colored by the
expression level of three markers differentially expressed on the different trajectories. (D)
Line plot showing the log2 fold change of the indicated marker in monocyte population
activated with LPS, LPS-IFNγ and IFNγ for the indicated time points compared to M5.
628

Figure 3. MDMs and TAMs express different combinations of similar markers
(A) t-SNE maps of TAMs, T cells, M0, and MDM populations based on the CyTOF antibody
panel. Cells were colored per MDM sample types (top panel) or per ascites samples (bottom
panel). In each case, the counterpart populations are colored in grey. (B) Violin plot showing
the expression pattern of the 36 markers across T cells, monocytes, pooled MDMs, and
pooled TAMs. For pooled populations, data on 2,000 cells of each individual sample were
combined. (C) t-SNE map of the 36 markers across T cells, monocytes, pooled MDMs, and
pooled TAMs colored by PhenoGraph clusters. (D) Heatmap showing the mean expression
levels of the 36 markers assessed by mass cytometry across the 26 PhenoGraph clusters.
Three clusters with less than 100 cells were left out of the analysis. Hierarchical clustering
was performed by calculating first the pairwise distance between samples based on the
Spearman correlation and by clustering based on Ward's method. The composition (comp.)
and the frequency (freq.) of each cluster are depicted on the right. (E) Wheel plot showing
the prediction score of MDMs, TAMs and T cells, based on a random forest classifier trained
on the MDM populations. The distance between each dot, representing a cell and the
vertices of the polygons is proportional to the predication score of each cell to the different
MDM populations. (F) Heatmap showing for each cell type the frequency of cells classified
with a 40% confidence into the different population used to train the model. See also Figure
S2.
648

Figure S1. Comparison of Marker Expression across the Different MDM Populations,
Related to Figure 1
(A) The data on MDMs generated from monocytes from four healthy donors were analyzed
simultaneously using the t-SNE statistic. For each sample, the data are displayed
individually on a scatter plot. In each graph, the indicated replicate is displayed using the
indicated color code, and the three other replicates are shown in grey. (B) Median
expression levels of the indicated markers in monocytes and MDM populations across four
replicates. (C) Violin plots showing the distribution of the indicated markers in the different

MDM populations as assessed by mass cytometry. Data were normalized to the 99
percentile. Data shown are from one representative sample.

Figure S2. Effect of Cell Freezing on Marker Expression as Assessed by Mass
Cytometry, Related to Figure 2
(A) Gating strategy based on Ir, Pt, CD45, CD3, CD64, and CD68 used to identify T cells
and TAMs. (B) t-SNE maps of the different MDM populations based on the CyTOF antibody
panel in presence (right panel) or in absence (left panel) of freezing step. Cells were colored
per MDM sample types and in each case, the counterpart populations are colored in grey.
(C) Correlation between marker expression in presence or in absence of enzymatic digestion
in the indicated cell samples. For each marker, the coefficient of determination R2 and the
linear model are displayed.

TABLE

Table 1. Surface molecules included in the final panel

Metal Tag	Target	Antibody Clone	Biological function
La139	CD209	DCS-8C1	C-type lectin involved in phagocytosis
Pr141	CD64	10.1	High affinity Fc receptor
Nd142	CD38	HIT2	cyclic ADP ribose hydrolase
Nd143	CD68	Y1/82A	glycoprotein
Nd144	CD36	5-271	scavenger receptor class B
Nd145	CD71	CY1G4	Transferrin receptor I
Nd146	CD155 (PVR)	SKII.4	Costimulatory receptor
Sm147	CD206	15-2	Mannose receptor
Nd148	CD13	WM15	Alanine aminopeptidase
Sm149	CD204	351615	Scavenger receptor
Nd150	CD123	6H6	Interleukin-3 receptor
Eu151	CD11b	M1/70	Complement receptor 3
Sm152	CD40	5c3	Costimulatory receptor
Eu153	Slamf7	162.1	Costimulatory ligand
Sm154	CD197	GO43H7	Chemokine receptor
Gd155	CD273	MIH18	coinhibitory receptor
Gd156	CD304	446921	VEGF co-receptor
Gd158	CD166	3A6	Adhesion molecule
Tb159	CD82	ASL-24	Tetraspanin glycoprotein
Gd160	CD274	130021	coinhibitory ligand

Dy161	CD119	GIR-208	Interferon-γ receptor
Dy162	CXCR4	12G5	Chemokine receptor
Dy163	CD87	VIM5	Urokinase receptor
Dy164	CD120b	3G7A02	Tumor necrosis factor receptor 1B
Ho165	CD4	RPA-T4	T cell coreceptor
Er166	CD32	FUN-2	Low affinity Fc receptor
Er167	CD16	3G8	Low affinity Fc receptor
Er168	CD14	RMO52	LPS co-receptor
Tm169	CD163	GHI/61	Scavenger receptor
Er170	CD169	7-239	Syaloadhesin
Yb171	CD86	2331 (FUN-1)	Costimulatory ligand
Yb172	HLA-ABC	W6/32	Antigen presentation
Yb173	CD81	5A6	Tetraspanin glycoprotein
Yb174	CD88	S5/1	complement component 5a receptor 1
Lu175	HLA-DR	L243	Antigen presentation
Yb176	CD54	HA58	Adhesion molecule

Figure 1. Characterization of MDM surface phenotypes by mass cytometry

(A) Monocyte preparation and stimulation to induce differentiation. (B) t-SNE map of M0 and MDM populations based on the 36 antibodies used for mass cytometry analysis. Cells are colored by sample types. (C) t-SNE map showing the normalized expression levels of the 36 surface markers across the different populations. Data are from one representative replicate. (D) Heatmap showing the normalized expression level of the 36 antibodies for the four replicates of the different populations included in the analysis. Hierarchical clustering was performed by calculating first the pairwise distance between samples based on the Spearman correlation, and by clustering based on Ward's method. For each subset, the six proteins showing the strongest enrichment based on marker enrichment modelling is indicated on the right. IC, immune complex; GC, glucocorticoid. See also Figure S1 and Table S1.

Figure 2. Kinetic analysis of MDM polarization

(A) t-SNE maps of MDMs activated with the indicated stimuli for the indicated amount of time. For each stimulation, the cells are colored with a gradient of red corresponding to the stimulation time shown on a grey background corresponding to all the other cells included in the analysis. (B) Heatmap showing the log2 fold change in marker expression across time points and stimuli compared to M5. Markers are clustered based on Spearman correlation and Ward's linkage. (C) Diffusion mapping based on the first three diffusion components showing the putative trajectories followed by monocytes activated with the indicated stimuli over 48h (top-left panel). The other panels are showing the diffusion map colored by the expression level of three markers differentially expressed on the different trajectories. (D) Line plot showing the log2 fold change of the indicated marker in monocyte population activated with LPS, LPS-IFNγ and IFNγ for the indicated time points compared to M5. See also Figure S2.

Figure 3. MDMs and TAMs are phenotypically distinct

(A) t-SNE maps of TAMs, T cells, M0, and MDM populations based on the CyTOF antibody panel. Cells were colored per MDM sample types (top panel) or per ascites samples (bottom panel). In each case, the counterpart populations are colored in grey. (B) Violin plot showing the expression pattern of the 36 markers across T cells, monocytes, pooled MDMs, and pooled TAMs. For pooled populations, data on 2,000 cells of each individual sample were combined. (C) t-SNE map of the 36 markers across T cells, monocytes, pooled MDMs, and pooled TAMs colored by PhenoGraph clusters. (D) Heatmap showing the mean expression levels of the 36 markers assessed by mass cytometry across the 26 PhenoGraph clusters. Three clusters with less than 100 cells were left out of the analysis. Hierarchical clustering was performed by calculating first the pairwise distance between samples based on the Spearman correlation and by clustering based on Ward's method. The composition (comp.) and the frequency (freq.) of each cluster are depicted on the right. (E) Wheel plot showing the prediction score of MDMs, TAMs and T cells, based on a random forest classifier trained on the MDM populations. The distance between each dot, representing a cell and the vertices of the polygons is proportional to the predication score of each cell to the different MDM populations. (F) Heatmap showing for each cell type the frequency of cells classified with a 40% confidence into the different population used to train the model.