Global diversity, predictors, and predictions of AMR evolutionary pathways in *Klebsiella pneumoniae*

Olav N. L. Aga[1,2], Sabrina J. Moyo[3,4,5], Joel Manyahi[4], Upendo Kibwana[4], Iren H. Löhr[6,1], Nina Langeland[1,3,7,8], Bjørn Blomberg[1,3,7], Iain G. Johnston[9,2,*]

1. Department of Clinical Science, University of Bergen, Bergen, Norway
2. Computational Biology Unit, University of Bergen, Bergen, Norway
3. Department of Medicine, Haukeland University Hospital, Bergen, Norway
4. Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania
5. Tropical Disease Biology Department, Liverpool School of Tropical Medicine, Liverpool, UK
6. Department of Medical Microbiology, Stavanger University Hospital, Stavanger, Norway
7. National Advisory Unit for Tropical Infectious Diseases, Haukeland University Hospital, Bergen, Norway
8. The Norwegian Institute of Public Health, Oslo, Norway
9. Department of Mathematics, University of Bergen, Bergen, Norway
* correspondence to iain.johnston@uib.no

Abstract

Antimicrobial resistance (AMR) is a substantial and growing global health burden. Understanding, and predicting, its evolution in specific pathogens will help responses across scales from individual patient cases to large-scale policy. Data-driven approaches to this question often focus more on genomes and less on the evolutionary dynamics generating these genomes. Here, we use global data on AMR features in *Klebsiella pneumoniae* with hypercubic transition path sampling (HyperTraPS), a machine learning approach, for Bayesian inference of the evolutionary pathways of AMR in *K. pneumoniae* (KpAMR) in 102 different countries, territories and areas. We identify "globally consistent" evolutionary behaviours that hold across countries, and "globally divergent" behaviours including carbapenem and fluoroquinolone resistance that vary across countries. We show how these divergent dynamics covary both with public health superregion and drug use policy, and reveal competing evolutionary pathways within and between countries. Using newly-sequenced data across several decades from sub-Saharan Africa, we show that this inferred global roadmap of KpAMR evolution successfully predicts prospective evolutionary dynamics. Together, we hope that the ability to characterize and predict evolutionary dynamics of AMR acquisition, connected to socio-economic and drug policy predictors, will help strengthen our understanding of AMR evolution worldwide.

Introduction

Antimicrobial resistance (AMR), the resistance of microbial pathogens to the medicines we use to treat them, is a major threat to global health. Antibiotic resistance, the resistance of bacteria to antibacterials, is an important subset of AMR. In 2019 and 2021, 1.27m and 1.14m deaths respectively were attributable to bacterial AMR (many more were associated with AMR), with highest rates in sub-Saharan Africa (Murray et al., 2022; Naghavi et al., 2024). The evolution of genetic and phenotypic features that cause AMR in bacterial pathogens is the focus of tremendous research interest, from basic biology through epidemiology, clinical studies and health policy.

Klebsiella pneumoniae (Kp) is a gram-negative bacterium and a major cause of nosocomial infections across settings. (Related species and subspecies of Kp form the KpSC (species complex); in this report we will focus on *K. pneumoniae sensu stricto*). Kp is found in multiple environmental niches and can both cause opportunistic infections in hospitalized patients and community acquired infections, including severe infections caused by hypervirulent strains (Wyres et al., 2020). Kp has been identified as an important trafficker for antibiotic resistance genes from different ecological niches into several other clinically important pathogenic bacteria (Wyres & Holt, 2018). Kp contributes significantly to the global burden of AMR. In 2021, 45 600 and 36 000 deaths were attributed to carbapenem and fluoroquinolone resistant Kp respectively. In 2021, Kp accounted for

12.7% of all deaths attributable to AMR in the population over five years old and 19.4% of AMR-attributable deaths in children under five years (Naghavi et al., 2024).

Given the substantial health burden associated with the evolution of AMR in Kp, large-scale data-driven research has explored the genomic structure of, and AMR features in, Kp populations (Argimón et al., 2021; Hetland et al., 2025; Holt et al., 2015; Munk et al., 2022; Wyres et al., 2020). Here, we consider a complementary perspective: the evolutionary pathways by which these features are acquired (Renz et al., 2024). Many related subquestions here can broadly be unified as: from a global perspective, in what orders are AMR features acquired over different instances of pathogen evolution? For example, given a newly observed isolate in the clinic with a given set of AMR features, can we predict which drug resistance(s) will evolve next (and thereby inform treatment guidelines accordingly)? And can we identify which extrinsic features – from physical environment, demography, drug policy, or others – influence the evolutionary acquisition of AMR?

These questions require approaches that can use the expanding volumes of AMR data to make and train descriptive and predictive evolutionary models. However, the large sets of features involved in typical AMR systems are often not addressable using traditional methods from evolutionary biology. For example, an often-studied dataset of *Mycobacterium tuberculosis* isolates contains susceptibility-resistance information for a panel of 10 different drugs (Casali et al., 2014). Comparative methods like the Mk (Markov k-state) model often struggle for more than around 7 potentially coupled characters under study (Johnston & Diaz-Uriarte, 2024). An alternative class of methods for studying the evolution of multiple characters – evolutionary accumulation modelling or EvAM – has emerged jointly from the cancer and evolution literatures (Diaz-Uriarte & Herrera-Nieto, 2022; Renz et al., 2024; Schill et al., 2024), but has traditionally considered independent samples, without the phylogenetic relationships found in bacterial populations (Dewar et al., 2025). While many EvAM approaches have the ability to infer dependencies between multiple interacting characters, neglecting this phylogenetic relatedness can lead to substantial artefacts in analysis (Maddison & FitzJohn, 2015).

Hypercubic transition path sampling (HyperTraPS) is an EvAM approach designed to incorporate phylogenetic information into the (Bayesian) inference of evolutionary pathways for multiple interacting characters (Aga et al., 2024; Greenbury et al., 2020; Johnston & Williams, 2016). Other EvAM approaches including TreeMHN (Luo et al., 2023), HyperHMM (Moen & Johnston, 2023), and HyperMk (Johnston & Diaz-Uriarte, 2024), also support phylogenetically coupled data, but we focus on HyperTraPS here because of its flexibility (Aga et al., 2024; Renz et al., 2024). In this article, we harness large-scale global datasets on AMR in Kp (KpAMR) with HyperTraPS to learn the structure and variability of evolutionary pathways of drug resistance in Kp across the globe, as well as the extrinsic factors shaping these pathways.

Figure 1. **Global genomic data on KpAMR. (A)** We use 47 721 *Klebsiella pneumoniae* (Kp) isolates from around the world, with (i) country-specific counts of isolates given by colour and years of isolation given in (ii). **(B)** UpSet plot giving counts of different KpAMR characters across these isolates. **Character names**: *AGly* (resistance to aminoglycosides); *Col* (colistin); *Fcyn* (Fosfomycin); *Flq* (Fluoroquinolones); *Gly* (Glycopeptides); *MLS* (Macrolides); *Phe* (Phenicols); *Rif* (Rifampin); *Sul* (Sulfonamides); *Tet* (Tetracyclines); *Tgc* (Tigecycline); *Tmt* (Trimethoprim); *Bla_a* (β-lactam resistance without extended spectrum or inhibitor-resistance); *Bla_inhR* (resistance to β-lactam/inhibitor combinations); *Bla_ESBL* (resistance to extended-spectrum β-lactams); *Bla_ESBL_inhR* (resistance to extended-spectrum β-lactam/inhibitor combinations); *Bla_Carb* (resistance to carbapenems); *SHV* (SHV β-lactamase with expanded enzyme activity), *Bla_chr* (SHV alleles conferring resistance to ampicillin), *Omp* (outer membrane protein). Further details available via Kleborate documentation at https://github.com/klebgenomics/Kleborate. We will use *-a*: acquired (via horizontal gene transfer); *-m*: mutation; *-chr*: chromosomal (intrinsic).

Results

Local and global inferred pathways of AMR evolution in Klebsiella

To learn and compare the likely pathways by which KpAMR is acquired in different countries, we obtained a dataset of 47 721 genomes from 102 different countries, territories and areas, and extracted details of the presence or absence of genes corresponding to each of 22 AMR classes, using PathogenWatch and Kleborate (Argimón et al., 2021) (Lam et al., 2021) (Methods; Fig. 1A-B). We refer to presence/absence of resistance genes for each of these classes as binary "resistance characters" ("character" referring to a particular property of a species in evolutionary biology). The set of characters we consider, following Kleborate, is given in the Fig. 1 caption; in this report we will use the shorthand defined there, grouping genes by drug resistance class (and Lahey class for β-lactamases) (Gupta et al., 2014; Lam et al., 2021; Tsang et al., 2024).

For each country in our dataset, we next used hypercubic transition path sampling (HyperTraPS), a machine learning approach, to infer the "pathways" of KpAMR character acquisition: that is, the *ordered sequences* with which characters are acquired over time. Figs. 2A-C illustrate this pipeline for a single country example (Romania, chosen for its moderate number of samples). HyperTraPS takes presence-absence patterns of characters on a phylogeny as input (Fig. 2A) and outputs an evolutionary model describing the pathways of character acquisition most supported by the data (Aga et al., 2024; Greenbury et al., 2020; Johnston & Williams, 2016; Renz et al., 2024). This model can be summarized, for example, as the set of posterior probabilities P_{ij} with which character i is acquired at ordering j in a putative evolutionary process (Fig. 2B). We refer to a plot where P_{ij} is visualised with point size, as in Fig. 2B, a "bubble plot". The full output of the inference process is a parameterised transition network describing the probability of different sequences of transitions through the state space of possible resistance characters (Fig. 2C). Respecting the fact that many KpAMR characters may be acquired reversibly (for example, via gain and loss of plasmids) and HyperTraPS assumes irreversible dynamics, we confirmed with synthetic control studies that reversible dynamics marginally increase uncertainty in our evolutionary inference but do not lead to systematic errors (Supp. Fig. 1).

Figure 2. **Inferring evolutionary dynamics from genome data across countries. (A)** Example source data from Romania, with 118 Kp isolates connected via a putative phylogeny derived from LIN codes (see Methods) and assigned presence (dark pixel) or absence (white pixel) markers for resistance to each of 22 antibiotic groups. **(B)** Summary "bubble plot" of evolutionary pathways inferred for Romanian data with HyperTraPS. The area of a bubble gives the posterior probability that a given character (row) is acquired at a given ordering in the accumulative evolutionary process. For example, *Bla_chr* is inferred to be a likely early acquisition, while *Bla_ESBL-a* likely occurs later and *Tgc_a* is inferred likely later still (or never). **(C)** A subset of the hypercubic transition graph inferred by HyperTraPS for Romanian data. Each edge's width gives the probability of a transition involving a single character acquisition from the upper state to the lower state. The uppermost state is the ancestral state with no AMR characters. Only early transitions above a threshold probability 0.025 are plotted for clarity. **(D)** Global plot, taking mean acquisition probability over all countries in the dataset. Probabilities higher than the "null model" of uniform acquisitions are plotted in blue.

The large majority of genomes with timing information were isolated in the period 2015-2020 (Fig. 1A). Our approach here does not directly consider the specific, real-world timings of the evolutionary transitions involved – although, given reliable data on the divergence times of all pairs of isolates, HyperTraPS can readily analyse continuous-time data (Aga et al., 2024; Renz et al., 2025). Rather, our approach uses the *relative* evolutionary timings from the source data phylogeny (for example, Fig. 2A) to report the *orderings* of events: which acquisitions precede or follow which others, given that characters may influence each other, with the timescale of these events (see Discussion).

While Figs. 2A-C demonstrate the approach for a single country, Fig. 2D gives the average acquisition probabilities across all countries in our dataset, and Supp. Fig. 2 gives the corresponding summary plots for every country in the dataset. Resistance to some antibiotic groups

are consistently observed across all countries: following very common presence of *Bla-chr* and
SHV-m, resistance characters against betalactams and extended-spectrum betalactams (ESBLs)
(*Bla-a*, *Bla_ESBL-a*) are often acquired earlier, while resistance to colistin (drug of last resort for
multi-drug resistant *Kp*; *Col-a* and *Col-m*) is acquired later. Country-specific differences exist in the
evolutionary acquisition of other characters, including resistance to carbapenems (*Bla_Carb-a)*.

In several instances, bimodality in the inferred evolutionary dynamics is observed. That is, a given
character may be acquired at an earlier stage or a later stage in the evolutionary process, but not at
intermediate stages. Such bimodality is characteristic of several different evolutionary pathways
existing in a system. Examples can readily be seen in Fig. 2B for Romania and Fig. 2D for the global
mean behaviour, where *SHV-m* is likely acquired either very early or at more intermediate stages,
with little probability in between. These competing dynamics are directly visible in Fig. 2C, where
distinct, canalised pathways exist: one involving *SHV-m* as the first acquisition, one involving
Bla_chr as the first acquisition and *SHV-m* only following rather later. Across countries (Supp. Fig. 2),
bimodality is observed in several characters: some examples are in *Bla_chr* (Tanzania, Zambia); *Bla-*
a (Ireland, S Korea; *Bla_ESBL-a* (Laos); and *SHV-m* (Romania and at least 15 other countries across
GDB regions). The bimodality in the mutational and chromosomal characters reflects the
differences in evolutionary dynamics between horizontal-gene-transfer-mediated acquisitions and
the other KpAMR characters we consider.

Across subsets of countries, interactions between our KpAMR resistance characters are observed
(Supp. Fig. 3). For example, in around a quarter of cases, the acquisition of tigecycline resistance
(*Tgc_a*) is inferred to promote the acquisition of glycopeptide resistance (*Gly_a*), and the acquisition
of sulfonamide resistance (*Sul_a*) is inferred to promote the acquisition of trimethoprim resistance
(*Tmt_a*). In some cases, the acquisition of MLS resistance (*MLS_a*) is inferred to repress the
acquisition of carbapenem resistance (*Bla_carb_a*), suggesting possible interactions that could be
clinically exploited in combination therapies (see Discussion).

However, although Supp. Fig. 2 gives a detailed "roadmap" of KpAMR evolution across countries, it
is hard to immediately extract global-scale insights from this representation. We therefore next
considered how to compare these results in a reduced-dimensionality picture.

Global consistency in evolutionary acquisition of resistance to a subset of drug families

To proceed, we used principal components analysis (PCA) to embed the high-dimensional set of
posterior probabilities in a 3D space (Fig. 3). Here, the axes of highest variance in the inferred
evolutionary pathways are identified and used to naturally lay out the individual country outputs.
Comparison of country-by-country behaviour along these axes can then reveal major sources of
variability (and similarity) in inferred KpAMR evolution across countries.

Inferred evolutionary pathways for Kp across countries consistently involve early acquisition of
characters conferring resistance to aminoglycosides, sulfonamides, trimethoprim, and betalactams
(*Agly*, *Sul*, *Tmt* and *Bla* characters). Aminoglycosides and sulfonamides were both discovered and
entered clinical use early in the mid 19[th] century. Resistance properties inferred to be acquired at
later orderings include extended spectrum beta-lactam and phenicol resistance, acquired via
mobile genetic elements (*Bla_ESBL-a*, *Phe-a*). Acquisitions inferred to be acquired late across
countries include inhR-driven resistance (*Bla_inhR-a*, *Bla_ESBL_inhR-a*), and resistance to later-
and final-line drugs including colistin, fosfomycin, tigecyclin, and glycopeptides (*Col-a*, *Fcyn-a*, *Tgc-*
a, *Gly-a*). These form a set of "globally consistent" evolutionary observations.

Figure 3. **Global structure of inferred pathways of AMR evolution. (A-B)** Principal components analysis (PCA) of country-specific "bubble plots" as in Fig. 2B, reflecting inferred evolutionary dynamics of KpAMR in different countries. Individual countries are plotted on PCAs 1-2 (A) and 2-3 (B), individually and with ellipses demarcating their corresponding GBD region (acronyms below). **(C-D)** Example "bubble plots" from countries reflecting the extremes of PCA1 (C) and PCA2 (D). In (C), Venezuela reflects poorly characterised evolutionary dynamics, with relatively uniform probabilities everywhere; Nigeria reflects precisely characterised evolutionary dynamics, with individual characters having highly likely, precisely specified orderings. In (D), country-specific differences in ordering emerge, including in those characters highlighted with boxes. For example, KpAMR evolution in Argentina is likely to involve *Flq-m* acquisition earlier than other characters; evolution in Senegal is likely to involve *Flq-m* acquisition later than other characters. GBD regions: CEEECA, Central Europe, Eastern Europe, Central Asia; HI, High-income; LAC, Latin American and Caribbean; NAME, North Africa and Middle East; SA, South Asia; SEEAO, South-east and East Asia and Oceania; SSAf, Sub-Saharan Africa. Zanzibar (dark outline) is from data newly sequenced in this study. Character names given in Fig. 1 caption.

Country-to-country differences in evolutionary acquisition of resistance to carbapenems, fluoroquinolones, tetracycline, and others

The general patterns above appear across evolutionary dynamics across countries and regions, with the first principal component of inferred Kp variability (accounting for 23% of the overall variance) simply reporting the precision with which these patterns can be characterised (Fig. 3C, Fig. 4). In countries supporting less precise inference (for example, with limited datasets), the expected ordering of these characters is more uniform and closer to the average null hypothesis of all characters behaving equally; in countries with more precise output the expected orderings diverge to early and late values (Fig. 3C, Fig. 4).

Figure 4. **KpAMR characters determining aspects of global variability in evolutionary dynamics.** Each country has an expected acquisition ordering for each KpAMR character. The plots show how these patterns of expected ordering are linked to the principal components of global KpAMR evolution variability in Fig. 3: PCA1 (top, grey); PCA2 (centre, red); and PCA3 (bottom, blue). **(A)** "Globally consistent" characters: those that covary tightly with PCA1. All characters here have either early or late expected orderings across all countries (either left or right of the central vertical line). Their position on PCA1 reflects the precision of the inference output. Less precise outputs (low PCA1 values, like Venezuela in Fig. 3C) will have orderings close to the central "null"; more precise outputs (high PCA1 values, like Nigeria in Fig. 3C) will have more divergent "early" and "late" orderings towards the edges of the plot. **(B-C)** "Globally divergent" characters: those that covary more tightly with PCA2 (B) or PCA3 (C). Expected orderings in these character sets span a wider range, showing sources of variability in the evolutionary dynamics that are independent of PCA1 (and hence less influenced by observational uncertainty). An alternative visualisation is given in Supp. Fig. 4. Character names given in Fig. 1 caption.

Once variability in this precision is accounted for (by considering the first principal component), variability corresponding to other differences in the inferred dynamics can be extracted. Several characters covary comparatively little with the (precision-aligned) PCA1 and instead show stronger covariance with PCA2 or PCA3, suggesting that there is genuine country-to-country variability in these characters after controlling for differences in observational noise (Fig. 3D, Fig. 4, Supp. Fig. 4). PCA2 (accounting for 15% of overall variance) displays strong covariance with carbapenem, fluoroquinolone, MLS, and tetracycline resistance and OMP mutation (*Bla_Carb-a*, *Flq-m*, *MLS-a*, *Tet-a, Omp-m,*). PCA3 (accounting for 8% of overall variance) displays covariance with *Flq-a*, *Rif-a*, *SHV-m*. None of these "globally divergent" characters display the "null-to-extreme" divergence seen in those characters most closely related to PCA1, suggesting genuine country-to-country differences in evolutionary dynamics, rather than observational differences, are responsible for this variability.

Figure 5. **Geographical variation in inferred AMR evolutionary pathways. (A-B)** Distributions of country-specific PCA projection values across different GBD regions for (A) PCA2 and (B) PCA3. In (A), sub-Saharan African countries show systematically higher positions in PCA2 than other regions. **(C)** Inferred acquisition orderings of KpAMR characters by GBD region. The left five characters covary with PCA2; the right three covary with PCA3. Carbapenem resistance and OMP mutations are inferred to be acquired substantially later in sub-Saharan Africa; rifampicin and fluoroquinolone resistance are inferred to be acquired substantially earlier in S, SE, E Asia and Oceania. New Zanzibar data from this study is shown by large circles. Character names given in Fig. 1 caption.

Geographical correlates with Kp evolutionary dynamics

To explore the sources of this evolutionary variability in more detail, we asked whether geographical region was linked to different variants of the inferred evolutionary dynamics (after accounting for observational differences). We first asked whether countries from different regions showed systematic differences in their PCA2 and PCA3 projections – corresponding to different orderings of the "globally divergent" country-specific characters above (Fig. 5). We found that sub-Saharan Africa was a dramatic outlier in PCA2, involving notably later acquisition of carbapenem and fluoroquinolone resistance and OMP mutation than the other regions (Fig. 5A, C). This ordering result agrees with the continuous-time history of treatment in the region, where carbapenem and fluoroquinolone use was established later (especially for children) than in other regions. Other regions were broadly consistent in PCA2 but showed substantial differences in PCA3. These differences correspond mainly to acquired resistance to fluoroquinolone and rifampicin, with substantially earlier inferred acquisition in South-Southeast-East Asia and Oceania, and substantially later in Central and Eastern Europe, Central Asia, and sub-Saharan Africa (Fig. 5B, C).

Figure 6. **Connecting AMR character acquisition propensity with drug treatment regimens across countries.** Median (across observed years 2016-2021) per-capita drug use in different drug classes (J codes), with expected acquisition ordering of associated resistance characters. "Globally consistent" characters *Tet-a*, *Bla_inhR-a*, *Sul-a, AGly-a* were associated with PCA1 in the global analysis and display little correlation between evolutionary ordering and drug use levels. "Globally divergent" characters *Bla_Carb-a* and *MLS-a* were associated with PCA2 in the global analysis and display stronger correlations, with earlier evolution of resistance linked to higher levels of drug use. Character names given in Fig. 1 caption.

Policy predictors of variance in AMR evolutionary dynamics

We next asked whether known policy of antibiotic use influence the inferred evolutionary dynamics from our analysis. We gathered statistics on drug use from 2016-2021 in countries available from the WHO GLASS surveillance programme (*Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report 2022*, 2022) and looked at correlations between per-capita drug use statistics and the inferred evolutionary dynamics of resistance to those drugs (Fig. 6). We found that for those drug types associated with PCA1 (where variability in inferred dynamics is largely determined by observation uncertainty) there was little correlation between drug use and the ordering of resistance acquisition. However, for carbapenems, associated with PCA2 (more non-observational country-to-country variability), there was a link between drug usage statistics and inferred resistance evolution, with resistance evolving earlier in countries with high drug use rates. The direction of this correlation was also observed in MLS, also associated with PCA2, though this relationship did not show statistical support at the $p < 0.05$ level. This pattern of correlations is consistent with the picture above: acquisition of some (PCA1) KpAMR characters is relatively constant across countries and circumstances, while other (PCA2) characters are more plastic and depend more tightly on country- and region-specific drivers.

Testing predictions of AMR evolutionary dynamics with newly sequenced, cross-decadal data

One motivating application of evolutionary inference to AMR problems is the ability to predict the next steps in evolutionary pathways (Renz et al., 2024). A question of potentially direct clinical relevance is: given a clinically observed isolate with a given set of AMR characters, which character(s) will likely evolve next – and can treatment guidelines be adapted accordingly?

Previous studies of AMR with HyperTraPS have demonstrated its ability to predict properties of evolutionary transitions in a withheld subset of the overall dataset, following the common training-test machine learning paradigm (Aga et al., 2024). Here, we are in the unique position of using independently obtained and Kp genome data to test the predictions of the evolutionary models trained above. We sequenced Kp isolates from patients with bloodstream infections in 2001-2 and 2017-8 in Tanzania and 2015-6 in Zanzibar, collected in previous studies (Blomberg et al., 2007; Moyo et al., 2020; Onken et al., 2015, 2024) (Fig. 7; see Methods). Bioinformatic analysis placed these new genomes in a phylogenetic tree with a subset of existing genomes from the training data, including their AMR profiles (Fig. 7A, Supp. Fig. 5). Using only transitions independent of those in the training data (see Methods), we tested the predictions of the HyperTraPS model trained on data from Tanzania. Specifically, for each new transition, we queried the trained model about likely next steps from the ancestral state, and recorded the predicted ranks of the next steps actually present in the transition (Fig. 7B). By far the most common outcome was that the true next step was ranked first (predicted to be most likely); compared to predictions from an untrained model, the trained model was systematically much better at predicting the true next steps.

This analysis demonstrates that within-country predictions of future evolutionary dynamics are possible given a trained HyperTraPS model. We also asked whether the model could predict evolutionary dynamics at the regional scale. We use Zanzibar as a test case here – a region of Tanzania that was not explicitly represented in the original Kleborate training dataset. Our findings would predict KpAMR in Zanzibar to fall into the sub-Saharan Africa set of behaviours, with high values on PCA2 and associated signatures of late carbapenem resistance and others (Fig. 5). To this end, we applied HyperTraPS inference to newly sequenced isolates from Zanzibar. The resulting inferred dynamics clearly place Zanzibar within the set of sub-Saharan Africa behaviours (Fig. 3A-B), with high values on PCA2 arising from later acquisition of carbapenem resistance, and high values on PCA3 arising from later acquisition of rifampicin resistance (Fig. 5).

We also explored the representation of different KpAMR characters in the snapshot genome sequences alone, without connecting with evolutionary dynamics. Fig. 7C shows the relative prevalence of acquired KpAMR characters in existing and newly-sequenced data, not accounting for phylogenetic relatedness (and without representation of an explicit non-AMR control group). With this sampling, most characters showed an increase from 2001-2 to 2015-6, particularly in mobile genetic acquisition of fluoroquinolone resistance, while the relative prevalence of acquired phenicol resistance and fluoroquinolone resistance mutations decreased. Proportions of these characters were in broad agreement with known coarser-grained observations from Tanzania (Mapunjo et al., 2025; Sangeda et al., 2021) and existing Tanzanian genomes from Pathogenwatch, but we observed more acquired rifampicin resistance and lower acquired phenicol resistance than in existing samples. Our samples from Zanzibar showed lower relative proportions of aminoglycoside and β-lactam resistance, and higher prevalence of outer membrane protein mutations.

Figure 7. **Prediction of new evolutionary transitions in KpAMR. (A)** Pathogenwatch (grey) and newly sequenced 2001-2 (blue) and 2017-8 (red) genome properties from Tanzania, connected by a phylogeny estimated from average nucleotide identity (ANI). **(B)** For each independent transition in the set of newly sequenced observations, we used HyperTraPS trained on existing data to predict which characters would likely evolve next given the ancestral state. The bars show the predicted rankings of the characters that were truly observed to evolve in the data. The trained model (blue-green) overwhelmingly predicted top rankings for the characters that were truly acquired; an untrained model (red) predicts intermediate rankings. Character names given in Fig. 1 caption. **(C)** Proportion of Tanzanian isolates in Kleborate and new datasets (uncorrected for relatedness) where different KpAMR characters are present.

Discussion

Powerful data-driven approaches to study the evolution of AMR are emerging in parallel with large genomic datasets (Argimón et al., 2021; Hetland et al., 2025; Holt et al., 2015; Munk et al., 2022; Wyres et al., 2020). Most of these studies focus on the current genomic structure of pathogen populations. In this research we have attempted to take a complementary perspective, using large-scale genome data while focusing on the (inferred) evolutionary dynamics that produce these genomic patterns. We believe that this new way of working has substantial power to support other data-driven approaches, particularly in the prediction of unseen and future evolutionary behaviours of pathogens (Renz et al., 2025).

The core technology we use, HyperTraPS, is an instance of evolutionary accumulation modelling or EvAM (Diaz-Uriarte & Herrera-Nieto, 2022; Diaz-Uriarte & Johnston, 2024). EvAM has previously been used to learn AMR pathways, including in HIV (Beerenwinkel et al., 2005) and multidrug resistance in tuberculosis (Aga et al., 2024; Greenbury et al., 2020; Moen & Johnston, 2023; Renz et

al., 2024). One question that has remained is to what extent such results reflect universal, general behaviors, versus a response to the specific selective pressures arising from a given country's drug use (for example). Our global comparison suggests a form of answer to this question. A subset of the KpAMR characters we consider appear to behave similarly across countries regardless of specific selective pressure, drug regimes, and other differences (those aligned with PCA1). Another set of characters (those aligned with PCA2-3) evolves in a way that is more dependent on country- and region-specific influences, which we have shown include public health superregions and drug use levels – although certainly other factors, from population change to the prevalence of other diseases, will also influence these dynamics. While these results are specific to *Kp*, and the characters we consider, this combination of global and country-specific behaviors may reasonably hold across other pathogens.

Inferring evolutionary dynamics of multiple, coupled traits across groups is a challenging problem. There will be countless, multiscale selective influences that impact the evolution of specific lineages of *Kp* around the globe. Our perspective attempts to "coarse-grain" away from these (unobservable) specific pressures and consider their aggregated effect on large-scale genomic properties in populations. This perspective is not without challenges. Some of these are: (a) The acquisition of many of the KpAMR characters we consider is reversible (for example, mediated by mobile genetic elements), which challenges an assumption of our modelling. (b) We assume that genomes from a country can be reasonably placed on their own phylogeny. (c) We use datasets which likely exhibit bias towards AMR presence: in most circumstances, due to timing and/or human research focus, genome databases will be enriched for cases with AMR characters acquired. And (d), in most of this analysis, we use relative evolutionary timings rather than the absolute real-world (day-month-year) timings associated with observed genomes. For (a), we have shown with a synthetic control study that violating the irreversibility assumption does not dramatically challenge our findings. (We note, in parallel to this point, that simultaneous acquisition of multiple features is very compatible with our approach; the resulting output simply places equal probabilities over all compatible ordering pathways.) For (b), coupling between countries (through, for example, host or pathogen migration) will not affect the interpretation of our results, which make statements conditioned on the set of pathogens reported within a country. And likewise for (c), the enrichment of AMR characters in our source data does not introduce systematic bias in our results, which report on accumulation of AMR characters, controlling for the existing AMR profile of an ancestral state in each case. Our phylogenetic reconstruction bridges the gap between observed, AMR-enriched genomes and their unobserved, less-enriched ancestors, ensuring that the whole spectrum of character profiles is captured in the underlying model.

For (d), the precise timescales of the evolutionary process are in principle addressable using a version of our approach called HyperTraPS-CT (continuous time) (Aga et al., 2024), but this requires a consistent estimation of divergence times between all isolates in the dataset. Importantly, our approach here does not completely neglect a timescale – it is implicit in the phylogenetic relationships included in the source data, and this inclusion both accounts for the temporal ordering of events and "pseudoreplication" due to the relatedness of sampled genomes (Boyko & Beaulieu, 2023; Dewar et al., 2025; Renz et al., 2025). However, this approach based on relative evolutionary timings means that we cannot directly tie differences in inferred evolutionary behaviour to known real-world historical events shaping AMR profiles: shifts in drug policy, for example, or outbreaks or other events changing the effective size of the evolving population. While our results, focusing on the ordering of acquisition events, already demonstrate descriptive and predictive power in KpAMR evolution across countries, a continuous time picture on a smaller scale in the future (perhaps a single country or small set) would enable finer-grained linking between model prediction and the real-world details of policy and disease history.

In learning estimates for evolutionary dynamics, HyperTraPS also infers potential interactions between characters, including whether the acquisition of one character makes acquiring another character less likely. Such "repressive interactions", if truly present, could form the basis of combination therapies: resistance to drug X makes resistance to drug Y less likely. These

interactions correspond to the phenomenon of *collateral sensitivity* in AMR, where resistance to one
drug induces sensitivity to another (Pál et al., 2015; Szybalski & Bryson, 1952): for example, in
Escherichia coli, beta-lactamase expression is linked to heightened sensitivity to colistin (among
other) (Herencias et al., 2024). While not observed consistently throughout all countries in our
dataset, a subset of countries showed statistical support for some such "repressive interactions"
(Supp. Fig. 3). These include, for example, MLS resistance repressing the acquisition of carbapenem
resistance. Such indications could suggest more detailed followup investigation to verify and
explore mechanisms behind these possibly exploitable collateral interactions.

In conclusion, we hope to show that evolutionary accumulation modelling – inferring the historic
evolutionary pathways of AMR acquisition, in addition to considering contemporary properties – can
complement and enhance established ways of working in AMR genomic analysis. By considering
the dynamics by which global KpAMR patterns have become established, a collection of research
directions are opened or expanded, including the similarities and differences between countries,
links with social and environmental covariates, and predictions of future dynamics. We anticipate
that these methods may readily be applied to provide insight into AMR evolution in other pathogens
in future.

Methods

Existing Klebsiella data acquisition

We obtained all 47 721 *Klebsiella pneumoniae sensu stricto* records from pathogenwatch as of
March 2024 (Argimón et al., 2021). The most common isolation years were 2015-2020, with a peak
at 2018 (Fig. 1A). We used AMR features (genes and mutations) reported by drug class as reported
by Kleborate version 2.3 (Lam et al., 2021). Kleborate groups genes or mutations known to confer
resistance to clinically relevant antibiotic groups and related phenotypes. Betalactamases are
further grouped by enzyme activity (Lam et al., 2021). We consider a set of L = 22 drug classes for
each genome, dichotomizing Kleborate output by presence of any gene or mutation related to each
group. That is, if there is any gene or mutation present for a resistance phenotype, the isolate is
considered resistant, otherwise it is considered susceptible. This assumption holds for most
resistance phenotypes considered in this study. However, fluoroquinolone resistance (*Flq-m*) often
require multiple mutations to cause resistance. Thus, the presence of these characters cannot be
interpreted as a resistance phenotype, but rather an increased MIC (minimum inhibitory
concentration). The isolates were divided by country based on metadata, and countries were
grouped into superregions as defined by the Global Burden of Disease regional classification
system (Rudd et al., 2020). A coarse-grained phylogeny was generated via LIN-codes (Life
identification numbers) using lincoding.py supplied by (Hennart et al., 2022). 2 102 genomes were
excluded due to missing metadata. Any country with less than two genomes available were
excluded due to the inability to construct a tree. The resistance profiles and phylogenetic tree were
combined into a phylogenetic tree annotated with resistance to drug classes.

Evolutionary pathway inference with hypercubic transition path sampling (HyperTraPS)

HyperTraPS (Aga et al., 2024) models an "evolutionary space" containing every possible state of a
system involving L binary characters, then infers the probabilities of (Markovian) transitions between
states in this space (here, involving the ordered accumulation of different AMR characters) that is
most compatible with observed data (Greenbury et al., 2020; Johnston & Williams, 2016). A recent
overview of the method in an AMR context is given in (Renz et al., 2024). The source data is a
collection of length-L "barcodes" labelling presence or absence of our characters -- resistance to
drug classes as defined in Kleborate -- on the tips of a phylogeny. Ancestral state reconstruction
(here, assuming that AMR resistance character accumulation is rare and irreversible, so that an
ancestor possesses a character if and only if all descendants possess it) is used to infer ancestral
states, and thereby construct a set of transitions from the data. These transitions are the input data
for the HyperTraPS algorithm. The target of inference is an L x L matrix θ, where θ_{ii} is the base rate
for acquiring character i, and θ_{ij} is the influence that acquisition of character j has on the base rate

of acquisition of character *i* (as in mutual hazard networks (Schill et al., 2020)). In this way, positive and negative interactions between the acquisition of different AMR characters is supported, so that (for example) acquisition of resistance to drug X can make acquisition of resistance to drug Y more (or less) likely. The output of the inference process is, most broadly, a transition matrix λ, with elements $\lambda_{a \to b}$ describing the probability with which an isolate in state *a* will next transition to state *b*. For each of the 102 countries, territories and areas we ran HyperTraPS on three different random seeds to ensure that the results were consistent across runs (Supp. Fig. 2). Simulations were run for 10^6 steps with 10^3 random walkers for sampling, employing a penalised likelihood. The US genomes were sub-sampled to a subset of 1000 genomes; three different subsamples clustered closer together than with any other countries in the PCA plot.

Synthetic control study for inference of reversible characters

To determine the effect of our assumption of irreversible accumulation of AMR characters (given that the loss of plasmids is not uncommon in Kp evolution (Holt et al., 2015)), we constructed a synthetic control study. A random phylogeny over 128 tips was constructed using a birth-death process with birth rate 1 and death rate 0.5 (chosen empirically to roughly match the phylogenies from our case studies) using *phangorn* (Schliep, 2011). A synthetic accumulation process reflecting a single evolutionary pathway was simulated on this phylogeny. Reversibility was captured by modelling random loss of the first *L*/2 characters as a Poisson process with a given rate parameter. The same HyperTraPS pipeline (assuming irreversibility), including ancestral state reconstruction and pathway inference, was run on both the pre-loss and post-loss datasets, and the outputs compared (Supp. Fig. 1).

Global structure in inferred pathways

The outputs of HyperTraPS can be summarized as a matrix *P*, where element P_{ij} gives the posterior probability that character *i* is acquired at ordering *j* in a putative evolutionary process from an ancestor with no AMR characters towards a final state with all AMR characters. For example, P_{32} gives the probability that the third character is the second to be acquired in such an evolutionary process. We obtained this matrix for each country in our dataset, then performed principal components analysis (PCA) on the set of matrices (Williams et al., 2013). We observed the structure of matrices with "bubble plots", where an array of points is plotted with areas proportional to P_{ij}.

Drug usage data

We collected antibiotic consumption data from 57 countries, territories and areas enrolled in the WHO Glass surveillance program in the period 2016 to 2021 (*Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report 2022*, 2022). Overlaying the consumption data with the countries that had 3 or more available genomes reduced the number to 53. We use the median consumption statistics over the available time points for each country.

New Klebsiella data acquisition

The 79 newly sequenced Kp isolates in this article are derived from a collection of studies performed in Tanzania and Zanzibar over the last three decades. These comprise bacterial isolates from (a) blood samples from pediatric patients with bloodstream infections in Dar-es-Salaam, Tanzania in 2001-2002; (b) blood samples from adult patients with bloodstream infections in Dar-es-Salaam, Tanzania in 2017-2018; (c) fecal samples from the patients in (b); and (d) blood samples from adult and pediatric patients with bloodstream infections in Mnazi Mmoja hospital, Zanzibar in 2015-2016. These cultures were sequenced by MicrobesNG (MicrobesNG, Birmingham, UK) using Illumina HiSeq technology. Assembled contigs were provided by the sequencing service, which formed the initial step for the bioinformatics pipeline below.

New Klebsiella data analysis

The analysis pipeline begins with the contigs from the initial sequencing and assembly process. We first used *dnadiff*, which wraps *nucmer* from the MUMmer 3.0 package (Kurtz et al., 2004), to

scaffold the contigs from a given isolate on the *Klebsiella* reference genome
GCA_000240185.2_ASM24018v2 (Liu et al., 2012) and report statistics of the alignment. We
retained only isolates with <20% unaligned bases as instances of *Klebsiella*. Combining the newly
sequenced isolates with the existing Tanzanian genomes from Pathogenwatch, we then ran pairwise
dnadiff across the combined set. We extracted the average identity across the aligned regions for
each pair and recorded this as the ANI (average nucleotide identity). We then used d = 1-ANI as a
distance measure for phylogenetic tree estimation using the unweighted pair group method with
arithmetic mean (UPGMA) (Sokal & Michener, 1958) implemented in the *phangorn* (Schliep, 2011)
package in R (R Core Team, 2022). We confirmed that an alternative method, neighbourhood joining
(Saitou & Nei, 1987), did not give qualitatively different results for the predictions of AMR evolution.
We used AMR profiles of existing and new isolates to reconstruct ancestral states across this
complete tree, then identified the set of subtrees that had only newly-sequenced isolates at their
tips. As transitions within such subtrees are independent of transitions in the (Pathogenwatch)
training data set, we used this set of transitions as our independent test set.

Testing predicted dynamics from the trained model on new data

We queried the model fitted from the training data to predict the likely next states from each
precursor state in this independent set of transitions (Aga et al., 2024; Renz et al., 2024). We ranked
the possible next steps from each state from most likely to least likely under the fitted model. We
then recorded the ranks corresponding to the actual step(s) corresponding to the observed
transition, and compared these to a null model where each possible transition could occur with
equal probability.

Data and code availability

The code base for this study is publically available at https://github.com/StochasticBiology/kp-evolution-inference. In addition to the software referenced above, we use R (R Core Team, 2022)
with libraries *hypertrapsct* for evolutionary inference (Aga et al., 2024), *dplyr* (Wickham et al., 2023),
tidyverse (Wickham et al., 2019), *countrycode* (Arel-Bundock et al., 2018) for data manipulation,
lme4 (Bates et al., 2015) for statistics, *phytools* (Revell, 2012) for phylogenetic analysis, and *ggplot2*
(Wickham, 2016), *ggpubr* (Kassambara, 2020), *ggbeeswarm* (Clarke et al., 2016), *ggrepel*
(Slowikowski, 2021), *ggupset* (Ahlmann-Eltze, 2025) for visualisation. We also use a Python script for
LINcoding by Melanie Hennart, which uses *numpy* (Harris et al., 2020). **Authors' note:** we are in the
process of uploading the new Kp sequences to a repository; a link will be provided when this
process is complete. In the meantime the raw FASTA files (which are the input for the pipeline) can
be downloaded from https://osf.io/36r45 .

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European
Union's Horizon 2020 Research and Innovation Programme [Grant agreement No. 805046
(EvoConBiO) to I.G.J.]. This project was supported by the Trond Mohn Foundation [project HyperEvol
under grant agreement No. TMS2021TMT09], through the Centre for Antimicrobial Resistance in
Western Norway (CAMRIA) [TMS2020TMT11], This project has received support from ERA-Net: JPI-AMR STRESST project (NFR333432). The authors are grateful to the CAMRIA collaboration and the
Stochastic Biology Group at UiB for useful discussions.

References

Aga, O. N., Brun, M., Giannakis, K., Dauda, K. A., Diaz-Uriarte, R., & Johnston, I. G. (2024). HyperTraPS-CT:
Inference and prediction for accumulation pathways with flexible data and model structures. *PLOS
Computational Biology*, *20*(9), e1012393.

Ahlmann-Eltze, C. (2025). *ggupset: Combination Matrix Axis for 'ggplot2' to Create 'UpSet' Plots* [R package].

Arel-Bundock, V., Enevoldsen, N., & Yetman, C. (2018). countrycode: An R package to convert country names
and country codes. *Journal of Open Source Software*, *3*(28), 848. https://doi.org/10.21105/joss.00848

Argimón, S., David, S., Underwood, A., Abrudan, M., Wheeler, N. E., Kekre, M., Abudahab, K., Yeats, C. A., Goater, R., Taylor, B., Harste, H., Muddyman, D., Feil, E. J., Brisse, S., Holt, K., Donado-Godoy, P., Ravikumar, K. L., Okeke, I. N., Carlos, C., ... NIHR Global Health Research Unit on Genomic Surveillance of Antimicrobial Resistance. (2021). Rapid Genomic Characterization and Global Surveillance of Klebsiella Using Pathogenwatch. *Clinical Infectious Diseases*, *73*(Supplement_4), S325–S335. https://doi.org/10.1093/cid/ciab784

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Beerenwinkel, N., Däumer, M., Sing, T., Rahnenführer, J., Lengauer, T., Selbig, J., Hoffmann, D., & Kaiser, R. (2005). Estimating HIV evolutionary pathways and the genetic barrier to drug resistance. *The Journal of Infectious Diseases*, *191*(11), 1953–1960.

Blomberg, B., Manji, K. P., Urassa, W. K., Tamim, B. S., Mwakagile, D. S. M., Jureen, R., Msangi, V., Tellevik, M. G., Holberg-Petersen, M., Harthug, S., Maselle, S. Y., & Langeland, N. (2007). Antimicrobial resistance predicts death in Tanzanian children with bloodstream infections: A prospective cohort study. *BMC Infectious Diseases*, *7*, 43. https://doi.org/10.1186/1471-2334-7-43

Boyko, J. D., & Beaulieu, J. M. (2023). Reducing the biases in false correlations between discrete characters. *Systematic Biology*, *72*(2), 476–488.

Casali, N., Nikolayevskyy, V., Balabanova, Y., Harris, S. R., Ignatyeva, O., Kontsevaya, I., Corander, J., Bryant, J., Parkhill, J., Nejentsev, S., Horstmann, R. D., Brown, T., & Drobniewski, F. (2014). Evolution and transmission of drug-resistant tuberculosis in a Russian population. *Nature Genetics*, *46*(3), Article 3. https://doi.org/10.1038/ng.2878

Clarke, E., Sherrill-Mix, S., & Dawson, C. (2016). ggbeeswarm: Categorical scatter (violin point) plots. *CRAN: Contributed Packages*. https://cir.nii.ac.jp/crid/1360019690351052672

Dewar, A. E., Belcher, L. J., & West, S. A. (2025). A phylogenetic approach to comparative genomics. *Nature Reviews Genetics*, *26*(6), 395–405. https://doi.org/10.1038/s41576-024-00803-0

Diaz-Uriarte, R., & Herrera-Nieto, P. (2022). EvAM-Tools: Tools for evolutionary accumulation and cancer progression models. *Bioinformatics*, *38*(24), 5457–5459.

Diaz-Uriarte, R., & Johnston, I. G. (2024). *A picture guide to cancer progression and monotonic accumulation models: Evolutionary assumptions, plausible interpretations, and alternative uses* (arXiv:2312.06824). arXiv. https://doi.org/10.48550/arXiv.2312.06824

Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report 2022 (1st ed). (2022). World Health Organization.

Greenbury, S. F., Barahona, M., & Johnston, I. G. (2020). HyperTraPS: Inferring probabilistic patterns of trait acquisition in evolutionary and disease progression pathways. *Cell Systems*, *10*(1), 39–51.

Gupta, S. K., Padmanabhan, B. R., Diene, S. M., Lopez-Rojas, R., Kempf, M., Landraud, L., & Rolain, J.-M. (2014). ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrobial Agents and Chemotherapy*, *58*(1), 212–220. https://doi.org/10.1128/AAC.01310-13

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Hennart, M., Guglielmini, J., Bridel, S., Maiden, M. C. J., Jolley, K. A., Criscuolo, A., & Brisse, S. (2022). A Dual Barcoding Approach to Bacterial Strain Nomenclature: Genomic Taxonomy of Klebsiella pneumoniae Strains. *Molecular Biology and Evolution*, *39*(7), msac135. https://doi.org/10.1093/molbev/msac135

Herencias, C., Álvaro-Llorente, L., Ramiro-Martínez, P., Fernández-Calvet, A., Muñoz-Cazalla, A., DelaFuente, J., Graf, F. E., Jaraba-Soto, L., Castillo-Polo, J. A., Cantón, R., San Millán, Á., & Rodríguez-Beltrán, J. (2024). β-lactamase expression induces collateral sensitivity in Escherichia coli. *Nature Communications*, *15*(1), 4731. https://doi.org/10.1038/s41467-024-49122-2

Hetland, M. A. K., Winkler, M. A., Kaspersen, H. P., Håkonsholm, F., Bakksjø, R.-J., Bernhoff, E., Delgado-Blas, J. F., Brisse, S., Correia, A., Fostervold, A., Lam, M. M. C., Lunestad, B.-T., Marathe, N. P., Raffelsberger, N., Samuelsen, Ø., Sunde, M., Sundsfjord, A., Urdahl, A. M., Wick, R. R., ... Holt, K. E. (2025). A genome-wide One Health study of Klebsiella pneumoniae in Norway reveals overlapping populations but few recent transmission events across reservoirs. *Genome Medicine*, *17*(1), 42. https://doi.org/10.1186/s13073-025-01466-0

Holt, K. E., Wertheim, H., Zadoks, R. N., Baker, S., Whitehouse, C. A., Dance, D., Jenney, A., Connor, T. R., Hsu, L. Y., Severin, J., Brisse, S., Cao, H., Wilksch, J., Gorrie, C., Schultz, M. B., Edwards, D. J., Nguyen, K. V., Nguyen, T. V., Dao, T. T., ... Thomson, N. R. (2015). Genomic analysis of diversity, population structure, virulence, and antimicrobial resistance in Klebsiella pneumoniae, an urgent threat to public health.

Proceedings of the National Academy of Sciences, *112*(27), E3574–E3581.
https://doi.org/10.1073/pnas.1501049112

Johnston, I. G., & Diaz-Uriarte, R. (2024). *A hypercubic Mk model framework for capturing reversibility in disease, cancer, and evolutionary accumulation modelling* (p. 2024.06.27.600959). bioRxiv.
https://doi.org/10.1101/2024.06.27.600959

Johnston, I. G., & Williams, B. P. (2016). Evolutionary Inference across Eukaryotes Identifies Specific Pressures Favoring Mitochondrial Gene Retention. *Cell Systems*, *2*(2), 101–111.
https://doi.org/10.1016/j.cels.2016.01.013

Kassambara, A. (2020). Ggpubr:"ggplot2" based publication ready plots. *R Package Version 0.4. 0, 438*.

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., & Salzberg, S. L. (2004). Versatile and open software for comparing large genomes. *Genome Biology*, *5*(2), R12.
https://doi.org/10.1186/gb-2004-5-2-r12

Lam, M. M. C., Wick, R. R., Watts, S. C., Cerdeira, L. T., Wyres, K. L., & Holt, K. E. (2021). A genomic surveillance framework and genotyping tool for Klebsiella pneumoniae and its related species complex. *Nature Communications*, *12*(1), 4188. https://doi.org/10.1038/s41467-021-24448-3

Liu, P., Li, P., Jiang, X., Bi, D., Xie, Y., Tai, C., Deng, Z., Rajakumar, K., & Ou, H.-Y. (2012). Complete genome sequence of Klebsiella pneumoniae subsp. Pneumoniae HS11286, a multidrug-resistant strain isolated from human sputum. *Journal of Bacteriology*, *194*(7), 1841–1842.
https://doi.org/10.1128/JB.00043-12

Luo, X. G., Kuipers, J., & Beerenwinkel, N. (2023). Joint inference of exclusivity patterns and recurrent trajectories from tumor mutation trees. *Nature Communications*, *14*(1), Article 1.
https://doi.org/10.1038/s41467-023-39400-w

Maddison, W. P., & FitzJohn, R. G. (2015). The Unsolved Challenge to Phylogenetic Correlation Tests for Categorical Characters. *Systematic Biology*, *64*(1), 127–136. https://doi.org/10.1093/sysbio/syu070

Mapunjo, S., Mbwasi, R., Nkiligi, E. A., Wilbroad, A., Francis, E. N., Msovela, K., Yahya, T., Mpembeni, R., Masunga, E., Nkungu, K., Saitoti, S., Lusaya, E., & Konduri, N. (2025). National consumption of antimicrobials in Tanzania: 2020–2022. *JAC-Antimicrobial Resistance*, *7*(2), dlaf026.
https://doi.org/10.1093/jacamr/dlaf026

Moen, M. T., & Johnston, I. G. (2023). HyperHMM: Efficient inference of evolutionary and progressive dynamics on hypercubic transition graphs. *Bioinformatics*, *39*(1), btac803.

Moyo, S. J., Manyahi, J., Blomberg, B., Tellevik, M. G., Masoud, N. S., Aboud, S., Manji, K., Roberts, A. P., Hanevik, K., Mørch, K., & Langeland, N. (2020). Bacteraemia, Malaria, and Case Fatality Among Children Hospitalized With Fever in Dar es Salaam, Tanzania. *Frontiers in Microbiology*, *11*, 2118.
https://doi.org/10.3389/fmicb.2020.02118

Munk, P., Brinch, C., Møller, F. D., Petersen, T. N., Hendriksen, R. S., Seyfarth, A. M., Kjeldgaard, J. S., Svendsen, C. A., van Bunnik, B., Berglund, F., Larsson, D. G. J., Koopmans, M., Woolhouse, M., & Aarestrup, F. M. (2022). Genomic analysis of sewage from 101 countries reveals global landscape of antimicrobial resistance. *Nature Communications*, *13*(1), 7251. https://doi.org/10.1038/s41467-022-34312-7

Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Aguilar, G. R., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-Woodhouse, G., Hamadani, B. H. K., Kumaran, E. A. P., McManigal, B., ... Naghavi, M. (2022). Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *The Lancet*, *399*(10325), 629–655.
https://doi.org/10.1016/S0140-6736(21)02724-0

Naghavi, M., Vollset, S. E., Ikuta, K. S., Swetschinski, L. R., Gray, A. P., Wool, E. E., Aguilar, G. R., Mestrovic, T., Smith, G., Han, C., Hsu, R. L., Chalek, J., Araki, D. T., Chung, E., Raggi, C., Hayoon, A. G., Weaver, N. D., Lindstedt, P. A., Smith, A. E., ... Murray, C. J. L. (2024). Global burden of bacterial antimicrobial resistance 1990–2021: A systematic analysis with forecasts to 2050. *The Lancet*, *404*(10459), 1199–1226. https://doi.org/10.1016/S0140-6736(24)01867-1

Onken, A., Moyo, S., Miraji, M. K., Bohlin, J., Marijani, M., Manyahi, J., Kibwana, K. O., Müller, F., Jenum, P. A., Abeid, K. A., Reimers, M., Langeland, N., Mørch, K., & Blomberg, B. (2024). Predominance of multidrug-resistant Salmonella Typhi genotype 4.3.1 with low-level ciprofloxacin resistance in Zanzibar. *PLOS Neglected Tropical Diseases*, *18*(4), e0012132.
https://doi.org/10.1371/journal.pntd.0012132

Onken, A., Said, A. K., Jørstad, M., Jenum, P. A., & Blomberg, B. (2015). Prevalence and Antimicrobial Resistance of Microbes Causing Bloodstream Infections in Unguja, Zanzibar. *PLoS ONE*, *10*(12), e0145632. https://doi.org/10.1371/journal.pone.0145632

Pál, C., Papp, B., & Lázár, V. (2015). Collateral sensitivity of antibiotic-resistant microbes. *Trends in Microbiology*, *23*(7), 401–407. https://doi.org/10.1016/j.tim.2015.02.009

R Core Team. (2022). *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2012*.

Renz, J., Dauda, K. A., Aga, O. N. L., Diaz-Uriarte, R., Löhr, I. H., Blomberg, B., & Johnston, I. G. (2024). *Evolutionary accumulation modelling in AMR: Machine learning to infer and predict evolutionary dynamics of multi-drug resistance* (arXiv:2411.00219). arXiv. https://doi.org/10.48550/arXiv.2411.00219

Renz, J., Dauda, K. A., Aga, O. N. L., Diaz-Uriarte, R., Löhr, I. H., Blomberg, B., & Johnston, I. G. (2025). Evolutionary accumulation modeling in AMR: Machine learning to infer and predict evolutionary dynamics of multi-drug resistance. *mBio*, *16*(6), e00488-25. https://doi.org/10.1128/mbio.00488-25

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, *2*, 217–223.

Rudd, K. E., Johnson, S. C., Agesa, K. M., Shackelford, K. A., Tsoi, D., Kievlan, D. R., Colombara, D. V., Ikuta, K. S., Kissoon, N., Finfer, S., Fleischmann-Struzek, C., Machado, F. R., Reinhart, K. K., Rowan, K., Seymour, C. W., Watson, R. S., West, T. E., Marinho, F., Hay, S. I., ... Naghavi, M. (2020). Global, regional, and national sepsis incidence and mortality, 1990–2017: Analysis for the Global Burden of Disease Study. *The Lancet*, *395*(10219), 200–211. https://doi.org/10.1016/S0140-6736(19)32989-7

Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, *4*(4), 406–425.

Sangeda, R. Z., Saburi, H. A., Masatu, F. C., Aiko, B. G., Mboya, E. A., Mkumbwa, S., Bitegeko, A., Mwalwisi, Y. H., Nkiligi, E. A., Chambuso, M., Sillo, H. B., Fimbo, A. M., & Horumpende, P. G. (2021). National Antibiotics Utilization Trends for Human Use in Tanzania from 2010 to 2016 Inferred from Tanzania Medicines and Medical Devices Authority Importation Data. *Antibiotics*, *10*(10), 1249. https://doi.org/10.3390/antibiotics10101249

Schill, R., Klever, M., Rupp, K., Hu, Y. L., Lösch, A., Georg, P., Pfahler, S., Vocht, S., Hansch, S., Wettig, T., Grasedyck, L., & Spang, R. (2024). Reconstructing Disease Histories in Huge Discrete State Spaces. *KI - Künstliche Intelligenz*. https://doi.org/10.1007/s13218-023-00822-9

Schill, R., Solbrig, S., Wettig, T., & Spang, R. (2020). Modelling cancer progression using mutual hazard networks. *Bioinformatics*, *36*(1), 241–249.

Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics*, *27*(4), 592–593. https://doi.org/10.1093/bioinformatics/btq706

Slowikowski, K. (2021). *ggrepel: Automatically Position Non-Overlapping Text Labels with'ggplot2'. R package version 0.9. 1, 2021*.

Sokal, R. R., & Michener, C. D. (1958). A statistical method for evaluating systematic relationships. *University of Kansas Science Bulletin*, *38*, 1409.

Szybalski, W., & Bryson, V. (1952). Genetic studies on microbial cross resistance to toxic agents i. *Journal of Bacteriology*, *64*(4), 489–499. https://doi.org/10.1128/jb.64.4.489-499.1952

Tsang, K. K., Lam, M. M. C., Wick, R. R., Wyres, K. L., Bachman, M., Baker, S., Barry, K., Brisse, S., Campino, S., Chiaverini, A., Cirillo, D. M., Clark, T., Corander, J., Corbella, M., Cornacchia, A., Cuénod, A., D'Alterio, N., Di Marco, F., Donado-Godoy, P., ... KlebNET-GSP AMR Genotype-Phenotype Group. (2024). Diversity, functional classification and genotyping of SHV β-lactamases in Klebsiella pneumoniae. *Microbial Genomics*, *10*(10), 001294. https://doi.org/10.1099/mgen.0.001294

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag. https://link.springer.com/book/10.1007/978-0-387-98141-3

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., ... Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Wickham, H., François, R., Henry, L., Müller, K., & Vaughan, D. (2023). Dplyr: A grammar of data manipulation. R package version 1.1. 2. *Computer Software*.

Williams, B. P., Johnston, I. G., Covshoff, S., & Hibberd, J. M. (2013). Phenotypic landscape inference reveals multiple evolutionary paths to C4 photosynthesis. *eLife*, *2*, e00961. https://doi.org/10.7554/eLife.00961

Wyres, K. L., & Holt, K. E. (2018). Klebsiella pneumoniae as a key trafficker of drug resistance genes from environmental to clinically important bacteria. *Current Opinion in Microbiology*, *45*, 131–139. https://doi.org/10.1016/j.mib.2018.04.004

Wyres, K. L., Lam, M. M. C., & Holt, K. E. (2020). Population genomics of Klebsiella pneumoniae. *Nature Reviews. Microbiology*, *18*(6), 344–359. https://doi.org/10.1038/s41579-019-0315-1

Supplementary Information

Supplementary Figure 1. **Synthetic control study on reversibility.** (top) A synthetic dataset involving the progressive accumulation of characters X1 to X10. The output inferred from HyperTraPS accurately reports the accumulation ordering of characters, with ambiguity about the first three characters which are observed across all samples. (bottom) A synthetic dataset involving reversibility. The same accumulation process is simulated, but with the potential for acquired characters to be lost reversibly. The output from HyperTraPS (which assumes irreversibility) is not dramatically different, except for a moderate amplification of the uncertainty in the early characters. The effect of reversibility here is to increase uncertainty rather than to introduce any systematic bias in the outputs.

Supplementary Figure 2. **All posterior orderings of AMR character acquisitions.** This "global roadmap"
summarises "bubble plots" describing inferred evolutionary dynamics of KpAMR characters in different
countries. Each row is a country, ordered vertically by descending number of Kp samples. Each column is a
KpAMR character; the horizontal axis within columns gives evolutionary orderings (from early to late). The size
of a point gives the probability that that character is acquired at that ordering in that country. For example,
across the vast majority of datasets (including the top row, USA, for example), *Bla_chr* has a high probability of
early acquisition and *Gly_acquired* has a high probability of late acquisition. In Cameroon, all characters have
similar inferred evolutionary orderings, reflecting the limited data available to make more precise statements.
Different colour points give the output of runs with different random seeds, to demonstrate consistency of the
estimates. Character names given in Fig. 1 caption.

Supplementary Figure 3. **Inferred interactions between KpAMR characters.** An arrow from character X to character Y reports an inferred interaction between the evolutionary acquisition of those characters. Blue arrow: acquiring X makes acquiring Y more likely ("promoting"). Red arrow: acquiring X makes acquiring Y less likely ("repressing"). The number on each arrow gives the proportion of countries in which that interaction was inferred. Character names given in Fig. 1 caption.

767
Supplementary Figure 4. **KpAMR characters determining aspects of global variability in evolutionary**
dynamics (alternative visualisation). Connected to Fig. 4. KpAMR characters (horizontal axis) are plotted by
their expected acquisition ordering (vertical axis) for each country (points). The country's projection on PCA1,
PCA2, and PCA3, for those characters that most strongly covary with each PCA axis, are given by the colour of
a point. As in Fig. 4, those characters that covary with PCA1 form a consistent spectrum linked to precision:
low values of PCA1 correspond to less precise, more uniform timing, while higher values lead to more precise
earlier or later timings. Characters covarying with the other PCAs display wider ranges connected to PCA1-
independent variability. Character names given in Fig. 1 caption.

Supplementary Figure 5. **New KpAMR data from Zanzibar and Dar es Salaam, Tanzania.** Phylogenies and KpAMR profiles from new datasets: **(A)** Zanzibar 2015-6. **(B)** Tanzania 2001-2. **(C)** Tanzania 2017-8. **(D)** Pathogenwatch Tanzania. **(E)** Circular projection of the phylogeny of Tanzanian isolates in Fig. 7A. Colours: green, 2001-2; blue, 2017-8; red, existing Pathogenwatch data.