**Main Manuscript for**

# The tissue specificity of cancer genes

Jiayi Chen[1-3], Ryan L Collins[3-5,*], Kevin M Haigis[2-3,5,*]

1. Departments of Biostatistics and Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA
2. Department of Cancer Biology, Dana-Farber Cancer Institute, Boston, MA
3. Cancer Program, Broad Institute of M.I.T. and Harvard, Cambridge, MA
4. Division of Population Sciences, Dana-Farber Cancer Institute, Boston, MA
5. Department of Medicine, Harvard Medical School, Boston, MA

* Correspondence to K.M.H. (Kevin_Haigis@dfci.harvard.edu) or R.L.C. (Ryan_Collins@dfci.harvard.edu)

**Author Contributions:** J.C. analyzed data; J.C. and R.L.C. wrote the paper; R.L.C. designed analyses; K.M.H. designed research; and K.M.H. revised and finalized the manuscript.
**Competing Interest Statement:** The authors have no conflicts to disclose.

**Classification:** Biological Sciences

**Keywords:** Cancer Biology, Somatic mutations, Cancer genes, Tissue specificity

**This PDF file includes:**
　　Main Text
　　Figures 1 to 3
　　Table 1

## Abstract

Defining genes that are somatically mutated in different cancers is a central goal of cancer genetics. Nevertheless, traditional definitions of "driver" genes based on pooled mutation frequency are biased toward common cancer types and tend to overlook genes that might be specific to rarer subtypes. As such, our understanding of the compendium of cancer genes is incomplete. In this study, we developed a statistical framework that defines genes enriched for functional somatic mutations in primary cancers to quantify incidence and tissue specificity for each gene. By applying this framework to the AACR GENIE v18.0 dataset, which comprised over 1.15 million mutations identified in 146,394 patients spanning 265 histologic subtypes, we identified a total of 95 genes significantly mutated in at least one subtype. We mined this dataset to derive tissue specificity scores for all 95 genes, demonstrating that tissue specificity in cancer is the norm, not the exception, for nearly all genes. We interrogated these new tissue specificity scores to reveal that oncogenes with restricted expression across normal tissues tend to exhibit higher tissue-specific mutation patterns in cancer. In contrast, tumor suppressors were often ubiquitously expressed irrespective of their underlying tissue specificity, which was partially correlated with differentially expressed compensatory genes and pathways in mutationally permissive tissues.

## Significance Statement

We present a statistically robust, subtype-aware framework to identify tissue-specific cancer driver genes and uncover their underlying biology. Our findings reveal that tissue-selective oncogenicity arises from expression constraints and tissue-restricted compensatory pathways. These findings provide a systematic, well-powered resource on mutational significance, tissue tropism, and biological consequences for the cancer research community.

**Main Text**

**Introduction**

Identifying cancer driver genes, defined as genes wherein somatic alterations confer a selective growth advantage to tumor cells, is fundamental to understanding tumor biology and guiding therapeutic development. Large-scale efforts such as The Cancer Genome Atlas (TCGA) have profiled thousands of tumors with a variety of genomic technologies and many sophisticated methods have been proposed for nominating mutational driver genes (1, 2). As a general rule, most approaches typically define driver genes based on mutational recurrence across a pooled set of tumor samples grouped by histology or tissue-of-origin. However, this pooled approach introduces a key limitation: genes that are commonly mutated in the common cancer types that are well represented in these studies, such as lung or breast cancer, tend to dominate the resulting lists of cancer driver genes. This bias is driven by relatively modest sample sizes for less common cancers in most pan-cancer cohorts, leading to low statistical power and a high false negative rate for driver gene detection in rarer cancers. As a result, genes that play important roles in less common cancer subtypes may be underappreciated, limiting our understanding of context-specific oncogenic mechanisms, as well as the extent of tissue selectivity for each mutational driver gene across all human cancer types, irrespective of disease prevalence.

Regardless of this limitation, numerous studies have proposed that tissue specificity is in fact quite common across cancer genetics. While a few genes such as *TP53* are broadly mutated, most oncogenes and tumor suppressors show restricted distributions of somatic mutations across tissues, shaped by underlying transcriptional, epigenetic, and proteomic states (3). For example, the flagship TCGA pan-cancer driver catalog reported that 142 of 258 driver genes (55%) were associated with just one cancer type, whereas only a handful of genes, such as *TP53*, *PIK3CA*, *KRAS*, *PTEN*, and *ARID1A,* exhibited broad distributions across a majority of cancer types (2).

Building on this perspective, we aimed to better understand the extent of tissue specificity among functional somatic mutations across as broad a spectrum of human cancers as possible. To accomplish this, we developed a subtype-aware statistical framework that evaluates somatic mutation enrichment within each histologic subtype and applied this framework to the largest publicly available pan-cancer clinicogenomic dataset, AACR GENIE v18.0 (4). After rigorous curation and quality control, our final analysis included somatic mutation data for 590 genes in 146,394 primary tumor samples spanning 265 histologic subtypes. We identified 95 genes with study-wide significant enrichment for functional mutations in at least one cancer subtype and subsequently defined a new metric to quantify how selectively each gene is mutated across cancer types while accounting for subtype-specific differential power, mutation frequencies, and other technical confounders. This approach enabled a more balanced detection of cancer drivers across the full spectrum of human malignancies, which represents a foundation for studying tissue-restricted oncogenic processes, and especially for processes relevant to less common cancers.

**Results**

**Statistically robust definitions of somatic mutation frequencies across histologic subtypes**

The first goal of our study was to quantify subtype-specific somatic mutation frequencies across a broad range of genes and histologic subtypes. To maximize our power for subtype-aware analyses, we curated somatic mutation data from AACR Project GENIE (v18.0). We excluded all metastatic samples due to their distinct mutation profiles at secondary sites. We also excluded subtypes with extremely limited power even at the large overall sample size of GENIE; specifically, we did not consider subtypes represented by fewer than 0.1% of the cohort (≤151 samples). Lastly, we identified and excluded likely misannotated samples (Fig. S1). After filtering, the final cohort included 146,394 non-metastatic tumor samples—of which 651 (0.44%) were benign—spanning

82   265 distinct histologic subtypes and 31 tissue types according to the established OncoTree
83   ontology (5) (Fig. 1A; Dataset S01).
84
Essentially all tumor samples in the AACR Project GENIE dataset were profiled using targeted
somatic DNA sequencing assays (*e.g.*, gene panel tests). Across all 146,394 samples, a total of
19,324 genes were targeted for sequencing in at least one sample; however, very few genes
(364/19,324; 1.88%) were sequenced in a majority (>50%) of the overall GENIE cohort, leading to
widespread differential missingness per gene across patient recruitment centers, sequencing
assays, and cancer types. To account for this considerable technical heterogeneity, we curated a
subset of 590 genes that had mutation frequencies high enough to rule out near-zero frequencies
with 90% confidence in at least one cancer subtype (Dataset S02). To capture functional
heterogeneity in somatic alterations, we assessed mutation enrichment separately for loss-of-
function (LoF), missense, and combined LoF or missense consequences.
95
We first wanted to establish an initial assessment of discrepancies between pan-cancer and tissue-
specific mutational frequencies in this dataset. To achieve this, we focused on a set of 112 genes
reported as cancer drivers in the IntOGen database that also met strict coverage criteria in GENIE
(≥80% of patients sequenced, ≥1000 times in at least two centers, and present in ≥10% of cancer
types) (6). For each gene, mutation frequencies were calculated using the combined frequency of
LoF and missense mutations for two viewpoints: (1) the mean mutation frequency across all
histologic subtypes, and (2) the mutation frequency for each gene's most permissive histologic
subtype, defined as the subtype with the highest lower bound of the 90% confidence interval (CI)
for that gene's mutation frequency, which was necessary to account for variable sample sizes
between subtypes (Fig. 1B). These analyses revealed the average mutation frequency for most
genes (109/112; 97.3%) was below 10% in the pan-cancer setting across the pooled cohort, with
15.2% (17/112) being mutated in fewer than 1% of all samples, underscoring the sparsity of
mutations in most genes when aggregated naively across all cancer types. In contrast, all genes
reached mutation frequencies greater than 5% within at least one specific subtype. For example,
*GNAQ* had an overall pooled mutation frequency of just 0.59% across all samples in GENIE but
reached a mutation frequency of 46.1% (90% CI = 40.5–51.8%) in uveal melanoma, reflecting
strong tissue specificity. Similarly, *VHL* was mutated in 1.06% of all samples but reached 67.8%
(90% CI = 65.4–70.0%) in clear cell renal cell carcinoma, where it is a well-established subtype-
specific canonical driver gene, while its mutation frequency remained low in non-clear cell renal
cancers (5.7%; 90% CI = 4.5–7.2%). These results emphasize the importance of applying a
subtype-aware viewpoint in analyses of somatic mutational enrichments, which can unmask
potential subtype-restricted drivers that might be obscured by pooled pan-cancer or gross organ-
level summaries.
119
120
121  **Estimating population-level gene mutation frequencies by combining molecular profiling**
122  **and epidemiologic data**
123
The overall frequency of functional somatic mutations in each gene across all human cancers is a
critical statistic for prioritizing diagnostic, preclinical research, and therapeutic development efforts.
In practice, such global mutation frequencies have been reported directly from large-scale cancer
genomic datasets like TCGA and GENIE, where certain rare cancer types are overrepresented
relative to their true population prevalence (*e.g.*, sarcomas, adrenal tumors) while common,
typically older-onset cancers are sometimes underrepresented (*e.g.*, prostate). These
discrepancies largely reflect clinical sequencing practices, as patients with more aggressive,
treatment-resistant, or recurrent tumors are more likely to undergo clinical genomic profiling.
132
We aimed to correct for these biases and more accurately approximate the real-world impact of
gene mutations across cancer types. To accomplish this, we projected the observed mutation
frequencies per gene in GENIE through population-level cancer incidence estimates from NCI
SEER (Dataset S03) (7). Overall, pan-cancer mutation frequencies per gene were broadly

concordant between SEER-adjusted and unadjusted models ($R^2$=0.7929; P<$10^{-20}$; Pearson
correlation test), but notable discrepancies were observed for certain genes enriched in subtypes
with disproportionate representation in sequencing datasets (Fig. 1C). For example, *KRAS*
exhibited an unadjusted mutation frequency of 16.2% across all primary tumor samples in GENIE
v18.0 but dropped to 10.2% after SEER adjustment, reflecting its relative enrichment among
aggressive cancers that are more likely to undergo genomic profiling as a part of their clinical
management. Similarly, *APC* decreased from 11.2% to 8.2%, and *TP53* decreased from 38.5% to
30.6%. These shifts illustrate how raw mutation frequencies derived from pan-cancer sequencing
projects may overstate the abundance of mutations prevalent in frequently sequenced tumor types,
while SEER-adjusted frequencies offer a more representative approximation of the true population-
level mutational burdens across all unselected human cancers.
148
We also noted a handful of genes that exhibited the opposite trend to genes like *VHL*, *KRAS*, and
*APC* (Fig. S2). For example, *AHNAK2* exhibited a high pan-cancer mutation frequency in GENIE
(16.05%) but substantially lower SEER-adjusted frequency (2.74%), despite being enriched in a
high-incidence cancer, breast cancer. Closer inspection revealed that *AHNAK2* was sequenced
almost exclusively (96.6%) in breast tumors in GENIE. Such cases, where a gene is sequenced
only in select tissues, can inflate unadjusted mutation frequencies and deflate incidence-adjusted
estimates. These effects highlight how gene panel design, often informed by prior knowledge or
assumptions about tissue relevance, can introduce ascertainment bias.
157
Although our study was focused on AACR Project GENIE, we also wanted to explore these similar
biases in gene mutation frequencies in TCGA, the most widely referenced pan-cancer
clinicogenomic dataset (Dataset S03). As with GENIE, the mutation frequencies for most genes
were well-correlated with their projected "true" population-wide frequencies ($R^2$=0.3142; P=5.2x$10^{-20}$
; Pearson test). However, a handful of genes, like *RNF213* and *LRP1B*, stood out as conspicuous
outliers. Both are exceptionally long genes (97th and >99th percentiles, respectively, for gene
length across all protein-coding human genes), making them more susceptible to inflated mutation
frequencies due to elevated passenger background mutations (Fig. S3). Thus, observed mutation
frequencies in large-scale sequencing efforts may reflect both biological factors (*e.g.*, gene length)
and clinical sequencing biases, which can be overcome by careful normalization of raw sequencing
data and population-level adjustment to better contextualize mutation frequency estimates across
subtypes.
170
171 **Tissue specificity is nearly ubiquitous across cancer gene mutations**
172
Having quantified the pan-cancer and subtype-specific mutation frequencies for 590 genes across
265 cancer types, we next wanted to statistically determine which gene-subtype pairs exhibited
enrichments of mutations significantly above the overall background rate of functional mutations
within each subtype. We designed and implemented a statistical framework to perform these
analyses and assessed all results at study-wide Bonferroni-corrected significance (P <1.60x$10^{-7}$).
An important aspect of this statistical framework was a *post hoc* empirical Bayes strategy to
determine a minimum fold-enrichment of somatic mutations (>10-fold; see Methods) that
distinguished established, biologically meaningful driver genes from genes that were subtly
enriched for somatic mutations but only achieved statistical significance due to very large sample
sizes for highly abundant cancer types (*e.g.*, lung, breast, colorectal). This *post hoc* fold-change-
based filter dramatically improved comparability between rare and common subtypes by
accounting for the greater power to detect significantly elevated mutational enrichments in well-
represented subtypes vs. rare subtypes; for example, the interquartile range of counts of significant
genes per subtype reduced from 17 to 5 after this *post hoc* adjustment (Fig. S4)
187
In total, we detected 487 significant gene-subtype pairs involving 95 distinct genes enriched >10-
fold in at least one subtype (Fig. 2A). Reciprocally, we identified at least one significant gene for
most subtypes (146/265, 55.1%; Fig. S5). Only two cancer subtypes, dedifferentiated liposarcoma
and ependymoma, showed no mutational enrichments despite both having >100 patient samples
and sufficient sequencing depth in ≥30% of genes (177/590) to reach a retrospective power >0.5,

based on the observed number of mutation calls. This lack of enrichment is consistent with their
known low mutational burden (8). To confirm that our framework recapitulated the known patterns
of cancer gene function, we next examined mutation consequences by genes' annotated roles in
cancer, finding that tumor suppressors were more frequently enriched for LoF mutations compared
to oncogenes ($P<10^{-20}$), consistent with their respective inactivation- and activation-driven roles in
cancer (Fig. 2B). Other than *TP53*, which was a pronounced outlier that displayed widespread
enrichment across virtually all tissues, most other cancer genes were enriched for mutations in only
a limited number of cancer subtypes. Furthermore, even within the same tissue, mutation
frequencies for many genes varied substantially across histologic subtypes (Fig. 2C). As mentioned
above, *VHL* was highly mutated in clear cell renal cell carcinoma but nearly absent in other kidney
cancers. Similarly, *RB1* was enriched exclusively in small cell lung cancers (mutation frequency of
42.1%; 90% CI = 38.8–45.4%), but no other lung histologies. In contrast, some genes, like *KRAS*,
exhibited comparatively broad enrichment across subtypes within a given tissue or organ but
nonetheless did exhibit notable variation in mutation frequencies among subtypes. *KRAS* was
mutated in >39% of samples across five of seven subtypes of bowel cancer, reaching 47.4% in
mucinous adenocarcinomas of the colon, but was rarely mutated in two subtypes: anal squamous
cell carcinomas (1.5%) and well-differentiated small bowel neuroendocrine tumors (~0%). These
examples underscore that many genes exhibit sharp mutation specificity not only across—but also
within—tissues.
212
We next wanted to systematically quantify this observed tissue specificity across all mutationally
enriched cancer genes. We therefore defined a "tissue specificity score" for each gene based on
the fraction of sufficiently powered cancer types in which the gene was significantly enriched
(Dataset S04. gene list ordered by tissue specificity score). We computed these scores separately
by mutational consequence (*e.g.*, missense, LoF) and subsequently scaled these scores from 0 to
100, with higher values indicating greater tissue specificity. Contrary to some conventional
viewpoints that many "hallmark" cancer genes are mutationally relevant across many tissues and
malignancies, our analysis underscored that a surprising degree of tissue selectivity of cancer gene
mutations is in fact the norm, with *TP53* being a singular rare exception as perhaps the only true
"pan-cancer" driver gene (Fig. 2D). Most genes, such as *VHL* and *KRAS*, exhibited significant
enrichment in a small subset of cancer types, emphasizing that colloquial discussions of a gene as
a "cancer gene" must be grounded in specific tissue or histologic context, even for very deeply
studied genes like *KRAS*.
226
227    **Limited evidence for race/ethnicity-specific driver genes in most cancer types**
228
We next wanted to understand the extent to which mutational enrichments within a cancer type
varied by other patient-specific factors, such as race or ethnicity. We therefore stratified our
mutational enrichment analyses by patient-reported race/ethnicity labels as recorded in GENIE,
which included the groups "White" (N=76,509), "Spanish/Hispanic" (N=9,472), "Asian" (N=6,502),
and "Black" (N=6,365). Importantly, as germline genetic data was not available for GENIE patients,
we emphasize that these groups were taken directly from the GENIE metadata and do not reflect
true underlying genetic ancestry; nonetheless, they provided an initial viewpoint for exploring biases
of mutation frequencies between these population.
237
Across all 1,579 genes with at least one somatic mutation reported in GENIE, we identified just five
gene-cancer pairs that were significantly enriched for somatic mutations in non-White populations,
with a significantly weaker (<10-fold) and/or non-significant (P>0.05) enrichment in White patients
(Table 1). These included some known ancestry-biased genes, such as *EGFR* missense mutations
in non-small cell lung cancer (NSCLC) enriched in Asian patients and *FOXA1* missense mutations
in prostate adenocarcinoma (PRAD) enriched in Asian and Black patients (9–11), as well as less
well-characterized examples, such as *ARID5B* LoF mutations in uterine endometrioid cancer (UEC)
in Asian patients (12), *JAK1* LoF in UEC in Hispanic patients, and *PTEN* missense mutations in
colon adenocarcinoma (COAD) in Black patients (13) (Fig. 2E). Of these five examples, we noted
that both *EGFR* in NSCLC and *PTEN* in COAD were nominally enriched in White patients but with
<10-fold enrichment. We therefore applied a *post hoc* binomial test to directly compare mutation

frequencies between White and non-White groups. Four of the five genes—*EGFR* in NSCLC,
*FOXA1* in PRAD, *ARID5B* in UEC, and *JAK1* in UEC—passed this *post hoc* binomial test for
significant enrichment in at least one non-White group after Benjamini-Hochberg correction,
confirming that the mutation frequencies in these four genes was significantly elevated in non-White
patients. The fifth gene, *PTEN* in COAD, showed a slightly elevated mutation frequency in Black
patients (5.8%) versus White patients (4.2%), but this difference did not reach statistical
significance after FDR correction.
256
We further confirmed that each gene-cancer association was reported by at least three independent
sequencing centers, and none of these genes were significantly enriched in White patients in any
parent or sister/cousin cancer type in OncoTree (*e.g.*, *EGFR* was not significantly mutated with >10-
fold enrichment in any other lung cancer subtype among White patients). These checks guarded
against biases that could arise if certain race/ethnicity groups were disproportionately sequenced
at centers that differed in the level of detailed sample annotation they submitted to GENIE. Thus,
while these analyses were constrained by comparatively limited statistical power in non-White
populations, even the smallest group included thousands of patients. The fact that only five genes
met our significance criteria therefore represents a small number of ancestry-specific differences.
Overall, these results suggested that many driver genes discovered in predominantly White
populations are likely also relevant in other racial and ethnic groups, though further studies in more
diverse cohorts remain essential, especially for rare cancers where statistical power is an even
greater concern.
270
271
272  **Gene expression in normal tissues partially explains tissue specificity of cancer mutations**
273
Having quantified tissue specificity scores for the 95 significantly mutated genes, we next sought
to investigate the biological factors that might underlie their tissue-selective mutational patterns.
Specifically, we considered whether baseline RNA expression levels across normal tissues might
help explain why certain genes incur somatic mutations observed in a restricted subset of cancers.
We hypothesized that genes broadly expressed across many normal tissues would be permissive
to somatic mutations in a wide range of cancer types, leading to lower tissue specificity scores,
whereas genes with narrower expression profiles might exhibit more selective patterns of mutation
enrichment across cancer subtypes.
282
To test this hypothesis, we obtained median gene-level RNA expression quantifications from 23
normal, non-cancerous tissues from postmortem donors from the Genotype-Tissue Expression
(GTEx) Project (v10) and computed an "expression breadth" statistic for each gene, defined as the
number of tissues in which expression exceeded a modest level of detectable expression (>5
transcripts per million [TPM]; Fig. S6) (14). We then assessed the relationship between expression
breadth and our cancer subtype specificity scores using Spearman's rank correlation (Fig. 3A-B).
We further stratified this analysis by oncogenic mechanism, as we reasoned that proto-oncogenes
and tumor suppressor genes might have different RNA expression patterns in normal tissues. For
our tumor suppressor-focused analyses, we included all 54 significantly LoF-enriched genes given
that most such genes (85.4%) corresponded to established tumor suppressors annotated in
COSMIC (15). However, as missense mutations likely represented a mixture of damaging, LoF-like
alleles and oncogenic, gain-of-function alleles, we restricted our analyses of missense mutations
to known oncogenes according to COSMIC (n = 25).
296
Missense-enriched oncogenes exhibited a negative correlation between tissue specificity and
expression breadth (Spearman $\rho$ = -0.294, P=0.0309), consistent with the hypothesis that
mutations preferentially conferred selective advantages in tissues where the proto-oncogene was
endogenously expressed, allowing the oncogenic mutant allele to be expressed in precancerous
cells. We noted that three genes—*ALK*, *RET*, and *FLT3*—were extreme examples of this trend,
exhibiting high missense subtype specificity and an extremely narrow spectrum of normal tissue
expression; curiously, all three of these genes are also known to be involved in recurrent oncogenic
fusion rearrangements in various cancer types (16–19). To confirm this overall correlation was not

driven by these three genes, we re-fitted a linear model after removing *ALK*, *RET*, and *FLT3,* which
revealed no significant differences in this overall trend (Welch's two-sample t-test of regression
coefficients, t=-0.651, P=0.518). In contrast, LoF-enriched genes showed a significant positive
correlation between expression breadth and tissue specificity (Spearman ρ=0.324, P=0.0168). This
pattern likely reflects the central roles of many tumor suppressor genes in cancer prevention across
myriad tissue contexts, as tumor suppressors tend to be ubiquitously expressed and involved in
essential cellular processes such as genome maintenance, cell cycle regulation, and apoptosis.
Collectively, these analyses provided support for our overarching hypothesis that the tissue-
specificity of somatic mutations in cancer is partially dependent on endogenous gene expression
levels in each cancer's tissue-of-origin, and that this relationship markedly differs between
oncogenes and tumor suppressors.

**Tissue-specific compensatory pathways contribute to extreme examples of subtype-**
**specificity**

Another common feature of some tumor suppressor genes is functional redundancy with other
tumor-suppressive pathways expressed in the same tissues (20). However, in specific tissue
contexts where these compensatory pathways are not utilized, LoF mutations in tumor suppressor
genes can be selectively advantageous. As a result, even broadly expressed tumor suppressors
may exhibit strong mutational specificity. Accordingly, we noted that a handful of genes—including
both oncogenes and tumor suppressors—stood out as particularly subtype-specific mutational
outliers despite being nearly ubiquitously expressed in normal tissues. We hypothesized that these
outliers may stem from common properties of the biological networks surrounding these genes and
therefore sought to investigate the role of compensatory pathways in shaping their selective
landscape across tissues. We focused on a set of six exemplar genes that exhibited broad
expression across normal tissues but had strong subtype-specific mutation enrichment. For
uniformity, we selected three LoF-enriched tumor suppressors (*KMT2D*, *MEN1*, and *PPM1D*) and
three missense-enriched oncogenes (*PDGFRA*, *FGFR1*, and *MAP2K1*), which we refer to as
"anchor" genes in subsequent analyses.

We reasoned that the mutationally permissive tissues for anchor tumor suppressors may lack
compensatory mechanisms that are otherwise active in non-permissive tissues. Similarly, we
hypothesized that the tissues permissive to oncogenic mutation in anchor oncogenes may have
signaling networks involving these oncogenes that are uniquely primed in certain tissues to be
sensitive to the molecular consequences of the oncogenic mutation. To test these hypotheses, we
first identified "peripheral" genes that were differentially expressed in permissive vs. non-permissive
normal tissues for each anchor gene at a false discovery rate of 5% (Fig. 3C). We reasoned that
the set of all differentially expressed peripheral genes would likely represent a mixture of the
compensatory pathways that could modulate the phenotypic impact of anchor gene mutations, as
well as any tissue-specific processes that were functionally irrelevant for the cancer gene in
question. Thus, to further place these peripheral genes in the context of known networks involving
each anchor gene, we queried the ENSEMBL IntAct Molecular Interaction Database to identify
physical protein-protein interactions (PPIs) between each anchor gene and its peripheral partners
(21). We constructed PPI networks up to four degrees of separation from each anchor gene and
performed Fisher's exact tests to determine whether peripheral genes were significantly enriched
within each anchor's network.

This analysis revealed significant enrichment of differentially expressed peripheral genes at higher
degrees of PPI connectivity (Fig. 3D). Among tumor suppressors, *MEN1* exhibited strong peripheral
gene enrichment among third-degree interactors (OR=2.03, P=0.0021) and fourth-degree
interactors (OR=1.36, P=0.0245), while *KMT2D* showed enrichment for third-degree interactors
(OR=1.51, P=0.0158). Among oncogenes, *MAP2K1* demonstrated peripheral gene enrichment for
third- and fourth-degree interactors (third-degree OR=1.17, P=0.0334; fourth-degree OR=1.23,
P=$6.5 \times 10^{-5}$). No significant enrichment was observed among first- or second-degree interactors for
any anchor gene, likely reflecting limited power due to the small number of direct interactors at

lower degrees: for example, these anchor genes had a median of only 4.5 first-degree PPIs (range
= 3–16). These findings support a possible model in which tissue-specific protein networks can
modulate the mutational selectivity of broadly expressed genes, irrespective of their functional
class.
365
Lastly, to characterize the specific biology implicated in tissue permissivity for each anchor gene,
we performed Gene Ontology (GO) enrichment analysis of all differentially expressed peripheral
genes within four degrees in each anchor gene's PPI network. For example, *MEN1*-associated
peripheral genes within the four-degree PPI showed significant enrichment for pathways related to
the regulation of mTOR signaling (Fig. 3E). These peripheral genes were downregulated in
pancreas, the tissue permissive to *MEN1* mutations in our analyses; indeed, *MEN1* is well-known
as a canonical driver specifically in pancreatic neuroendocrine tumors (22). Within four PPI
degrees, *MEN1* was connected to seven peripheral genes annotated as negative regulators of
mTOR, all of which coverage on the *RELA–SIRT1*–mTOR axis, consistent with prior reports in
pancreatic neuroendocrine tumors (23). This pattern provides a potential explanation for *MEN1*'s
mutational tissue-specificity: if inhibitory components of the mTOR pathway are already repressed
in pancreas relative to other tissues, then LoF mutations in *MEN1* may further weaken this
regulatory axis, thereby enhancing mTOR activity and promoting tumorigenesis. Collectively, these
analyses highlighted that tissue-selective oncogenesis is shaped by both intrinsic gene expression
levels activity and extrinsic cellular context, each acting as a gatekeeper for mutational impact.
381
382
383
384 **Discussion**
385
In this study, we conducted a systematic statistical survey of nonsynonymous mutation enrichments
in hundreds of putatively cancer-relevant genes across most histologic subtypes of human cancers.
Our analyses demonstrated that tissue selectivity is the prevailing rule among virtually all cancer
mutational driver genes, with only one exception—*TP53*—showing broad, pan-cancer enrichment
(3). Our work builds upon seminal prior studies from TCGA and others, which arrived at initial
coarse estimates of subtype specificity based on thousands of samples and dozens of cancer types
(24). For example, the TCGA consortium estimated that most cancer genes were mutated in only
2-20% of tumors (24). In our analysis of 146,394 samples spanning 265 histologic subtypes, we
similarly observed that few genes are mutated at high frequency across all tumors. However, by
evaluating each subtype separately, we found that many of these genes show clear enrichment
within specific tissue contexts, suggesting that the intermediate mutation frequencies in pan-cancer
datasets can mask prominent subtype-specific patterns. By implementing this subtype-aware
statistical framework, which adjusted for many biases endemic to large, aggregated clinicogenomic
cancer cohorts, we were able to document a more uniform landscape of genes with uniformly strong
enrichments of nonsynonymous mutations in both common and rare cancer subtypes, including
genes that would likely be overlooked in conventional pooled analyses: 11.6% of the genes we
identified as significantly enriched in at least one subtype were not catalogued as drivers in intOGen
(e.g., *TNFAIP3*, *NPM1*, *DNA2*, *H3F3A*), 7.4% were absent from COSMIC (e.g., *SOX9*, *ACD*, *TLR4*,
*ALB*), and 6.3% were absent from both. These findings suggest that our framework may expand
the current catalog of candidate driver genes by capturing subtype-restricted signals that are
missed in many pan-cancer analyses.
407
In contrast, we were surprised to detect only five candidate gene-cancer pairs that exhibited
significant mutational enrichments in self-reported non-White vs. White patients, implying that there
is apparently limited variability in strong-effect mutational drivers of cancer across major
race/ethnicity groups. As germline genetic ancestry is not available for the GENIE dataset, these
findings need to be confirmed in future well-powered studies with paired germline and somatic data.
If this observation holds in future studies, the lack of inter-ancestry variability encouragingly implies
that diagnostic and therapeutic efforts directed towards oncogenic genes may be applicable for all
populations.
416

As with most large-scale genomic studies of pooled patient cohorts, our analyses have several
important limitations that present opportunities for future studies. First, we focused on point
mutations and small indels, leaving copy number, structural, and epigenomic alterations
unexplored. Integrating these data could provide a more complete portrait of tissue-specific
oncogenesis, especially for tumor suppressor genes (25). Second, we relied on GTEx bulk RNA-
seq profiles to estimate baseline gene expression, which reflect heterogeneous mixtures of cell
types and may not capture the precise cell(s) of origin; integrating single-cell or lineage-resolved
transcriptomic data could provide more accurate insights (26). Third, despite the large cohort size
of AACR Project GENIE, statistical power remained limited for ultra-rare cancers and for analyses
stratified by race or ethnicity, particularly among non-White populations. Expanding to ancestrally
diverse cohorts will be essential for identifying any existing population-specific drivers of rarer
cancers. Fourth, our gene-level analysis did not account for allelic heterogeneity among variants
within the same gene—for example, different oncogenic alleles of *KRAS* (*e.g.*, G12D, G12V) are
known to have distinct signaling consequences and tissue-dependent activity (27). Together, these
considerations underscore the conservatism of our current findings while highlighting clear paths
toward a more mechanistic and clinically actionable understanding of tissue-specific oncogenesis.
433
434  In summary, by systematically quantifying tissue-restricted patterns of functional somatic
435  mutations, our study challenges the implicit multi-cancer framing of some driver gene catalogs and
436  emphasizes the value of a subtype-resolved perspective. This approach expands the list of
437  potential drivers across the cancer spectrum and also provides a quantitative scaffold for
438  investigating the biological and clinical consequences of tissue-specific oncogenesis.
439
440
441  **Materials and Methods**
442
443  **Data curation**
444
We analyzed somatic mutation data from the American Association for Cancer Research (AACR)
Project GENIE v18.0 (4), which comprised 250,018 tumor samples from 211,526 patients across
835 cancer subtypes. To avoid ambiguity in tissue-of-origin and to reduce patient duplication due
to serial biopsies, we first excluded all metastatic samples before retaining one representative
sample per patient (prioritizing primary tumors; see Supplementary Methods).We further collapsed
all OncoTree subtypes to level 3 before discarding tissues/subtypes that were underpowered;
specifically, we excluded all subtypes that represented <0.1% of the overall GENIE cohort, <1% of
all samples within their parent tissue, or had <1,000 samples overall. A site bias–flagging step (see
Supplementary Methods) removed 1,900 likely misannotated samples, yielding a final cohort of
146,394 primary tumors spanning 265 subtypes (Dataset S01).
455
456  **Definition of subtype-resolved gene mutation frequencies**
457
We first grouped all somatic mutations by predicted functional consequence (*e.g.*, missense, LoF,
silent). For each gene in each cancer pair, raw mutation frequencies were defined for each gene
in each cancer type as the fraction of all samples with at least one mutation reported divided by all
samples in that cancer type that had any sequencing coverage for the gene. We subsequently
normalized these frequencies for cancer-type-specific mutational burden and gene-level mutability,
the latter corrected using gnomAD-derived mutation rates adjusted for coding sequence length
(28). Statistical enrichment was quantified by chi-squared tests, in which the observed counts for
each gene-cancer pair were compared to expected counts derived from a gene length-adjusted
background within the same cancer subtype accounting for subtype-specific exome-wide
mutational density. We assessed all chi-squared tests at Bonferroni-adjusted significance and
further imposed a fold-change cutoff (≥10-fold enriched vs. baseline), ensuring that detected events
represented potent, major mutational drivers of each subtype rather than genes with weaker effects
situationally driven to significance by variation in sample size across subtypes. Refer to
Supplementary Methods for details of secondary analyses based on this dataset, such as

epidemiologic incidence adjustment of mutation frequencies or comparison of frequencies between
self-reported race/ethnicity groups.

474
475    **Computation of tissue specificity scores**

476
To quantify how selectively each gene is mutated across cancer types, we defined a tissue
specificity score for each gene by comparing the number of significantly enriched cancer types to
the number of adequately powered cancer types for that gene. Statistical power for each gene-
cancer pair was estimated using a noncentral chi-squared model parameterized by expected effect
sizes from the IntOGen driver gene compendium. Expected effect sizes were assigned as OR=9.7
(combined), OR=16.4 (LoF), and OR=13.2 (missense) relative to baseline. (see Supplementary
Methods) (6). In this framework, power reflects whether the number of sequenced samples for a
given gene-cancer pair is sufficient to detect enrichment, assuming a plausible driver-like effect
size. Cancer types with power >0.5 were considered adequately powered. We then computed a
tissue specificity score for each gene as the number of well-powered genes divided by the number
of significantly mutated genes after pruning direct parent-child OncoTree relationships to avoid
double-counting closely related cancer types. Finally, we scaled these scores from 0 to 100, with
higher values reflecting greater tissue specificity.

490
491    **Integration of normal tissue expression and interaction networks**

492
To assess whether tissue specificity of mutations related to baseline gene expression, we
integrated GTEx v10 median TPM values across 23 tissues mapped to the top-level OncoTree
nodes that also were included in our filtered version of the GENIE dataset (14). Expression breadth
was defined as the number of tissues in which a gene was expressed above 5 TPM (Fig. S6). We
correlated expression breadth with tissue specificity scores using Spearman and linear regression,
stratifying missense genes by oncogene/TSG status based on COSMIC annotations (15). We next
identified "peripheral" genes for each significantly mutated gene in these analyses. Peripheral
genes were defined as those that were differentially expressed between mutationally permissive
vs. non-permissive tissues for each cancer gene in question. Differential expression was tested
using either t-tests or z-tests across 18,772 protein-coding genes defined by GENCODE v48 that
were also quantified in GTEx (see Supplementary Methods) (29). To test whether these peripheral
genes were functionally linked to anchor genes, we assessed their enrichment in PPI networks
derived from IntAct (21), testing enrichment across 1-4 degrees of network proximity using Fisher's
exact test.

507
508
**Acknowledgments**

We thank Sasha Gusev and Eli Van Allen for their input and guidance. We would also like to
acknowledge the American Association for Cancer Research and its financial and material support
in the development of the AACR Project GENIE registry, as well as members of the consortium for
their commitment to data sharing. Interpretations are the responsibility of study authors. This work
was supported by an award from the Cancer Research UK Grand Challenge and the Mark
Foundation to the SPECIFICANCER team and NCI K99 CA286805 to RLC. KMH is also supported
by the Dana-Farber Cancer Institute Hale Center for Pancreatic Cancer Research and the Project
P fund. RLC received additional support from the Rossy Foundation Fund at Myriad Canada
(formerly KBF Canada).

520
521
**References**

1.    Cancer Genome Atlas Research Network, *et al.*, The Cancer Genome Atlas Pan-Cancer
analysis project. *Nat Genet* **45**, 1113–1120 (2013).

2.   M. H. Bailey, *et al.*, Comprehensive Characterization of Cancer Driver Genes and Mutations.
*Cell* **173**, 371-385.e18 (2018).

3.   K. M. Haigis, K. Cichowski, S. J. Elledge, Tissue-specificity in cancer: The rule, not the
exception. *Science* **363**, 1150–1151 (2019).

4.   AACR Project GENIE Consortium, AACR Project GENIE: Powering Precision Medicine
through an International Consortium. *Cancer Discov* **7**, 818–831 (2017). Data version 18.0.

5.   R. Kundra, *et al.*, OncoTree: A Cancer Classification System for Precision Oncology. *JCO*
*Clin Cancer Inform* **5**, 221–230 (2021).

6.   A. Gonzalez-Perez, *et al.*, IntOGen-mutations identifies cancer drivers across tumor types.
*Nat Methods* **10**, 1081–1082 (2013).

7.   Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov)
SEER*Stat Database: Incidence - SEER Research Data, 8 Registries, Nov 2024 Sub (1975-
2022) - Linked To County Attributes - Time Dependent (1990-2023) Income/Rurality, 1969-
2023 Counties, National Cancer Institute, DCCPS, Surveillance Research Program, released
April 2025, based on the November 2024 submission.

8.   N. Somaiah, *et al.*, Targeted next generation sequencing of well-
differentiated/dedifferentiated liposarcoma reveals novel gene amplifications and mutations.
*Oncotarget* **9**, 19891–19899 (2018).

9.   E.-E. -Ke, Y.-L. Wu, EGFR as a Pharmacological Target in EGFR-Mutant Non-Small-Cell
Lung Cancer: Where Do We Stand Now? *Trends Pharmacol Sci* **37**, 887–903 (2016).

10.   S. P. D'Angelo, *et al.*, Incidence of EGFR exon 19 deletions and L858R in tumor specimens
from men and cigarette smokers with lung adenocarcinomas. *J Clin Oncol* **29**, 2066–2070
(2011).

11.   J. Hwang, *et al.*, Structurally Oriented Classification of FOXA1 Alterations Identifies Prostate
Cancers with Opposing Clinical Outcomes and Distinct Molecular and Immunologic
Subtypes. *Clin Cancer Res* **31**, 936–948 (2025).

12.   D. S. Guttery, *et al.*, Racial differences in endometrial cancer molecular portraits in The
Cancer Genome Atlas. *Oncotarget* **9**, 17093–17103 (2018).

13.   M. Muquith, D. Hsiehchen, Landscape and Saturation Analysis of Mutations Associated With
Race in Cancer Genomes by Clinical Sequencing. *Oncologist* **29**, 219–226 (2024).

14.   GTEx Consortium, The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**, 580–585
(2013).

15.   J. G. Tate, *et al.*, COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids*
*Res* **47**, D941–D947 (2019).

16.   H. Mano, Non-solid oncogenes in solid tumors: EML4-ALK fusion genes in lung cancer.
*Cancer Sci* **99**, 2349–2355 (2008).

17.   A. Y. Li, *et al.*, RET fusions in solid tumors. *Cancer Treatment Reviews* **81**, 101911 (2019).

18.   G. Tang, *et al.*, Myeloid/lymphoid neoplasms with FLT3 rearrangement. *Mod Pathol* **34**,
1673–1685 (2021).

19. X. Zhao, *et al.*, Epigenetic activation of the FLT3 gene by ZNF384 fusion confers a
therapeutic susceptibility in acute lymphoblastic leukemia. *Nature communications* **13** (2022).

20. C. J. Sherr, Principles of Tumor Suppression. *Cell* **116**, 235–246 (2004).

21. N. Del Toro, *et al.*, The IntAct database: efficient access to fine-grained molecular interaction
data. *Nucleic Acids Res* **50**, D648–D653 (2022).

22. F. Marini, F. Giusti, F. Tonelli, M. L. Brandi, Pancreatic Neuroendocrine Neoplasms in
Multiple Endocrine Neoplasia Type 1. *Int J Mol Sci* **22**, 4041 (2021).

23. Y. Jiao, *et al.*, DAXX/ATRX, MEN1, and mTOR pathway genes are frequently altered in
pancreatic neuroendocrine tumors. *Science* **331**, 1199–1203 (2011).

24. M. S. Lawrence, *et al.*, Discovery and saturation analysis of cancer genes across 21 tumour
types. *Nature* **505**, 495–501 (2014).

25. O. Pös, *et al.*, DNA copy number variation: Main characteristics, evolutionary significance,
and pathological aspects. *Biomed J* **44**, 548–559 (2021).

26. K. A. Hoadley, *et al.*, Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000
Tumors from 33 Types of Cancer. *Cell* **173**, 291-304.e6 (2018).

27. C. Johnson, D. L. Burkhart, K. M. Haigis, Classification of KRAS-Activating Mutations and the
Implications for Therapeutic Intervention. *Cancer Discov* **12**, 913–923 (2022).

28. K. J. Karczewski, *et al.*, The mutational constraint spectrum quantified from variation in
141,456 humans. *Nature* **581**, 434–443 (2020).

29. J. M. Mudge, *et al.*, GENCODE 2025: reference gene annotation for human and mouse.
*Nucleic Acids Res* **53**, D966–D975 (2025).

586
587

**Figures and Tables**

**Figure 1.** Mutational properties of the AACR GENIE v18.0 dataset after our *post hoc* curation and
quality control steps. (A) Count of unique primary tumor samples included in our final analysis,
stratified by major tissue (top-level OncoTree layer). (B) Many genes exhibited large discrepancies
between their overall pan-cancer mutation frequency versus their highest subtype-restricted
mutation frequency. Shown here are the 112 IntOGen-annotated driver genes with adequate
sequencing coverage in GENIE (≥80% of patients sequenced, sequenced ≥1000 times in at least
two centers, and present in ≥10% of cancer types). For each gene, a gray line segment connects
its average pan-cancer mutation frequency to the mutation frequency observed in its most
permissive tissue subtype. Genes are ordered along the x-axis by increasing pan-cancer rate.
Genes exhibiting an exceptionally large discrepancy between their pan-cancer and subtype-
restricted mutation frequencies (absolute difference >0.30) are annotated with labels on the axis.
(C-D) SEER-adjusted GENIE mutation frequencies (y-axis) plotted against pan-cancer unadjusted
mutation frequencies from (C) GENIE and (D) TCGA. See Methods for adjustment procedure.

605
**Figure 2.** Landscape and specificity of gene mutation enrichments across cancers. (A) Mutation
enrichment landscape across the 31 top-level tissues in OncoTree. Each tile represents a single
gene–tissue pair; colored tiles indicate pairs with statistically significant enrichment (P<1.60x10⁻⁷).
Lower-right orange triangles denote enrichments of missense mutations; upper-left red triangles
denote enrichments of LoF mutations. Opacity reflects the proportion of mutationally enriched
histologic subtypes within each tissue. Genes are ordered by increasing tissue specificity; tissues
are clustered by mutational profile similarity. No fold-change filter was applied for this visualization.
(B) Virtually all (94.8%) of the mutational enrichments we observed among established oncogenes
corresponded to missense mutations, whereas most associations involving established tumor
suppressor genes (TSGs) were composed of loss-of-function (LoF) mutations. (C) Subtype-level
mutation frequencies for representative genes in permissive tissues: *KRAS* in bowel, *VHL* in kidney,
and *RB1* in lung. These examples illustrate how subtype-level resolution can reveal important inter-
subtype differences within overall tissue-level enrichments. (D) Relationship between our tissue
specificity scores and overall pan-cancer mutation frequencies per gene. Each point represents a
gene that was significantly enriched for mutations in at least one cancer. (E) We identified five
genes that were significantly enriched for mutations in self-reported non-White patients as
compared to White patients within the same histologic cancer subtype.
623

624
**Figure 3.** Expression breadth, tissue specificity, and pathway context of gene mutation
enrichments. (A-B) Relationship between expression breadth in normal tissues and tissue
specificity scores for (A) genes enriched in LoF mutations and (B) oncogenes with missense
mutations. Each point represents a gene, with a dashed red line indicating the linear regression fit.
Genes of interest referenced later in the text are highlighted in green. (C) Numbers of peripheral
genes for three LoF mutation enriched tumor suppressors (*KMT2D*, *MEN1*, *PPM1D*; red) and three
missense mutation enriched oncogenes (*PDGFRA*, *FGFR1*, *MAP2K1*; orange). Bars to the left
represent downregulated peripheral genes and bars to the right represent upregulated peripheral
genes. (D) PPI network centered on *MEN1*, displaying selected first- through fourth-degree
branches that terminate in peripheral genes or include key pathway nodes (*MTOR*, *AKT*). Nodes
are annotated by differential expression status: significant FDR < 5% differentially expressed genes
are shown in red, and all others in blue. (E) GO enrichment analysis of *MEN1*-associated peripheral
genes, showing the top 10 biological process terms with the lowest FDR-adjusted p-values. These
terms converge on terms related to regulation of mTOR signaling.
639
640
641

| Gene | OncoTree cancer type | Mutation consequence | Enriched non-White race/ethnicity | White frequency (%) | non-White frequency (%) | Benjamini-Hochberg Q-value |
|---|---|---|---|---|---|---|
| *EGFR* | NSCLC | Missense | Asian | 10.3 | 29.6 | $7 \times 10^{-6}$ |
| *FOXA1* | PRAD | Missense | Asian | 10.5 | 27 | $<10^{-10}$ |
| *FOXA1* | PRAD | Missense | Black | 10.5 | 13.2 | $<10^{-10}$ |
| *ARID5B* | UEC | LoF | Asian | 7.9 | 13.7 | 0.0118 |
| *JAK1* | UEC | LoF | Hispanic | 11 | 12.3 | $<10^{-10}$ |
| *PTEN* | COAD | Missense | Black | 4.2 | 5.8 | 0.251 |

643
644 **Table 1.** Genes with greater enrichments of mutations in self-reported non-White vs. White
645 patients.

## A

**Bulk Expression vs. LoF Specificity Score**

## B

**Bulk Expression vs. Missense Specificity Score (Oncogenes)**

## C

## D

## E

*MEN1*

**Peripheral genes enrichment**