Fork Reversal Safeguards Epigenetic Inheritance during DNA Replication Under Stress

Qiong Wu[1,5], Caixian Zhou[1,5], Yue Dou[2], Nadia I. Martin[2], Maria Gwit[2], Tae Hee Lee[4], Mark
Hedglin[4], Roséa Chen[1], Ke Zhu[1], Tianpeng Zhang[1,3], Shangming Tang[2,3], Tian Zhang[2,3], Tyler M.
Weaver[2,3], Wenpeng Liu[1,2,3,6]*

[1]Department of Radiation Oncology, University of Virginia School of Medicine, Charlottesville,
VA 22908, USA.

[2]Department of Biochemistry and Molecular Genetics, University of Virginia School of Medicine,
Charlottesville, VA 22908, USA.

[3]University of Virginia Comprehensive Cancer Center, Charlottesville, VA 22908, USA.

[4]Department of Chemistry, The Pennsylvania State University, University Park, PA 16802

[5]These authors contributed equally

[6]Lead Contact

*Correspondence: W. L. xbg5zt@virginia.edu

Abstract

During DNA replication, epigenetic information carried by histone modifications is faithfully
propagated and re-established on sister chromatids, ensuring cell identity. Chromatin reassembly
is tightly coupled to DNA replication, therefore whether and how perturbations to DNA replication
affects the fidelity of epigenetic inheritance remain poorly understood. In this study, we reveal a
critical role for replication fork reversal in maintaining the transmission of epigenetic information
under replication stress. We identify that cells defective in fork reversal exhibit reduced
nucleosome density at replication forks, accompanied by the loss of parental histones during their
transfer onto nascent DNA. Mechanistically, we demonstrate that PrimPol activation leads to
single-stranded DNA gaps in fork reversal deficient cells, and that subsequent PARylation (poly
ADP-ribosylation) and DNA–protein crosslinking on these gaps evicts nucleosomes. Our findings
demonstrate that replication fork reversal, a widespread physiological process, is not only essential
for preserving genome integrity but also for safeguarding epigenetic stability.

Introduction
In eukaryotic cells, DNA replication is not merely the duplication of genetic material; it is also a
critical process for the faithful transmission of epigenetic information[1]. During each S phase of the
cell cycle, DNA must be precisely duplicated, and chromatin structure must be accurately
reassembled to ensure cell identity, proper gene expression, and overall genome stability[2,3]. This
chromatin-based inheritance, referred to as epigenetics, encompasses histone modifications, DNA
methylation, and regulation by non-coding RNAs[4]. Programmed regulation of epigenetic states is
essential for cell differentiation and organismal development[5], whereas aberrant epigenetic
changes can drive diseases and tumorigenesis[6-10]. In cancer therapy, epigenetic exacerbates tumor
cell heterogeneity and promotes drug resistance[11]. Therefore, understanding how epigenetic
information is transferred during DNA replication is of fundamental importance.

During DNA replication, nucleosomes ahead of the fork are disassembled by the histone chaperone
FACT (FAcilitates Chromatin Transcription) complex into one hexamer, which contains a H3–H4
tetramer and a H2A–H2B dimer. Next, the hexamer is recycled to its original positions on one of

the sister chromatid by multiple histone chaperones [1,12-20]. With the assistance of histone
chaperones, like ASF1 and CAF1, recycled parental histones are incorporated together with newly
synthesized histones to assemble nucleosomes on nascent DNA[1,21-32]. Post-translational
modifications (PTMs) on parental histones serve as templates that are copied onto adjacent new
histones through a "read–write" mechanism, thereby completing the replication of epigenetic
information[1,33-37]. Replisome proteins such as MCM2, PCNA, DNA polymerases ε/α, Tof1, Mrc1,
Csm3, and Ctf4 contribute to parental histone transfer[25-27,35,38-48]. As a result, DNA replication
exerts a profound influence on epigenetic stability. For example, depletion of Pol E3/E4, subunits
of leading-strand histone chaperone polymerase ε, disrupt H3K9me3 marked parental histones
recycling, thereby activating expression of LINE-1 (Long Interspersed Element-1)
retrotransposon[49]. Conversely, mutations in lagging-strand histone chaperones, such as MCM2-
2A, lead to parental histone accumulation on the leading strand[38,39]. Simultaneous disruption of
both leading- and lagging-strand histone chaperones results in the loss of parental histones and
histone PTMs, impairing stem cell differentiation and viability[50].

DNA replication frequently encounters obstacles such as transcription–replication conflicts and
repetitive sequences[3,51]. Replication stress response pathways are activated to mitigate the
replication obstacles[3,51]. Among these pathways, the generation of single-stranded DNA (ssDNA)
and fork reversal are the two most common structural intermediates, occurring in diverse cell types
in response to various forms of replication stress[52,53]. We and others previously identified that
ssDNA can be processed by RAD51-mediated strand invasion together with multiple fork
remodelers, including SMARCAL1, ZRANB3, and HLTF, to form reversed forks[53-64]. This
process stabilizes replication intermediates and prevents excessive ssDNA accumulation to
promote proper DNA replication. When fork reversal is impaired, cells activate PrimPol to
reinitiate DNA synthesis downstream of the leading strand in a process known as repriming[65,66].
Although replication can resume, this generates abundant ssDNA gaps on nascent DNA that pose
a threat to chromatin structure and genome stability[67-69].

Under stressed conditions, DNA replication progress slowly compared to non-stressed conditions,
rather than completely stop[70]. It is unclear how fork reversal or ssDNA gaps influence parental
histone recycling and nucleosome assembly. In this study, we demonstrate that replication fork
reversal promotes parental histone recycling and nucleosome assembly on sister chromatids by
suppressing ssDNA formation, thereby preserving epigenetic stability. In cells lacking the ability
to undergo fork reversal, activation of the PrimPol primase leads to repriming and the generation
of ssDNA gaps, which ultimately impairs recycling of parental histones and histone PTMs. We
further show that impaired histone recycling results from the accumulation of ssDNA that triggers
PARP1-mediated PARylation and DNA–protein crosslink formation, both of which inhibit
nucleosome assembly. Together, our findings reveal a critical role for replication fork reversal in
safeguarding epigenetic stability under replication stress.

Results

Replication fork reversal deficiency reduces nucleosome density on nascent DNA under
replication stress

To investigate the impact of replication fork reversal on replication-associated nucleosome
assembly at replication forks, we performed SILAC-iPOND-MS (Stable Isotope Labeling by
Amino acids in Cell Culture - Isolation of Proteins on Nascent DNA - Mass Spectrometry) analysis

in wild-type (WT) U2OS cells and U2OS cells lacking the three major fork reversal enzymes,
including SMARCAL1, HLTF, ZRANB3 (termed as FRKO) to examine protein composition on
nascent DNA, upon hydroxyurea (HU)-induced replication stress (Figure 1A). The SILAC-
iPOND-MS analysis identified that FRKO cells have a markedly decreased abundance of nearly
all histones on nascent DNA compared to WT cells (Figure 1A). Notably, the levels of the MCM2–
7 helicase complex remained unchanged between WT and FRKO cells, indicating that histone
reduction was not due to a decrease in replication fork number (Figure 1A). In addition, compared
to WT cells, FRKO cells showed mild increased levels of EdU (5-Ethynyl-2'-deoxyuridine)
incorporation (Figure S1A), indicating that histone reduction in FRKO cells is not the result of
reduced replication progression. Immunofluorescence analysis of total chromatin bound histone
H3 showed no difference between fork reversal–deficient and wild-type cells (Figure S1B–E),
identifying that histone loss was restricted to nascent DNA.

To validate the mass spectrometry results, we pulse labeled nascent DNA with EdU, induced
replication stress with HU, and quantified histone H3 on nascent DNA using a proximity ligation
assay (PLA)[71] in WT and two clones of FRKO U2OS cells. Since FRKO cells incorporate EdU
more rapidly than wild-type cells (Figure S1A), direct comparison of PLA signals would not
accurately reflect histone abundance on nascent DNA (Figure 1B, 1C). Therefore, we normalized
PLA signals to EdU intensity, yielding a value representing histone abundance per unit length of
replicated nascent DNA (Figure 1D). Using this approach, we observed a significant decrease in
H3 levels on nascent DNA in two FRKO clones. Additional PLA analyses identified a marked
decrease in H2A, H2B, and H4 abundance, consistent with the observations for H3 (Figure 1B-D),
indicating the FRKO cells have a decreased abundance of all four core histones rather than loss of
individual histone types (Figure S1F-H). To further validate these results, we used ChOR
(Chromatin Occupancy after Replication)-seq, a high-throughput sequencing technique to
profiling the occupancy of histone (or modifications) and chromatin-associated proteins on newly
replicated DNA[72]. The ChOR-seq analysis also identified a significant reduction in histone H3
abundance on nascent DNA in the FRKO cells (Figure 1E-F, Figure S1I-J). Together, the PLA
and ChOR-seq analysis is consistent with the replication fork proteomics experiments that histone
abundance is reduced, suggesting low nucleosome density on nascent DNA in FRKO cells (Figure
1A).

These experiments were performed in U2OS cells with co-depletion of SMARCAL1, ZRANB3,
and HLTF. To determine which enzyme loss dominates the phenotype, we performed PLA
analysis in two clones for every single knockout of SMARCAL1, ZRANB3, and HLTF in U2OS
cells (Figure S2A–C). Notably, among these single knockout clones, both clones for HLTF
knockout consistently show significantly lower H3 level on nascent DNA (Figure S2A and S2C),
suggesting HLTF depletion dominates the phenotype in FRKO U2OS cells. Additionally,
SMARCAL1 and ZRANB3 single KO also shows a mild decrease in H3 level on nascent DNA
(Figure S2A-C). FRKO summarizes all three single depletions (Figure S2C). Consistent with
observations in fork reversal enzyme knockout cells, knockdown of RAD51, which is also an
essential factor for fork reversal, led to low nucleosome density on nascent DNA (Figure 1G and
Figure S2D).

Notably, PLA and ChOR-seq showed no reduction in H3 levels in FRKO cells without treatment
with HU (Figure S2E-G), supporting that histone loss arises as a result of replication stress.
Together, these data indicate that fork reversal machinery maintains nucleosome density on
nascent DNA during conditions of replication stress (Figure S2H).

Replication fork reversal deficiency leads to parental histone loss

During DNA replication, parental histones positioned ahead of the fork are disassembled and subsequently recycled onto daughter DNA through highly coordinated mechanisms, where they combine with newly synthesized histones to form nucleosomes. Under conditions of persistent replication stress, the reduced nucleosome density observed on nascent DNA in FRKO cells may arise from either impaired parental histone recycling or defective incorporation of new histones, both of which threaten faithful epigenetic inheritance. Importantly, the deposition of certain histone variants, such as CENP-A, macroH2A, and H3.3, occurs independently of DNA replication [73-75]. Thus, detection of these variants on nascent DNA reflects inheritance of parental histones, and their loss can be attributed to defective parental histone recycling (Figure 1A).

Prior work identified that specific histone post-translational modifications (PTMs), such as di- and trimethylation of H3K9, H3K79, and H4K20, are established on new histones many hours after replication or during the subsequent cell cycle, making them markers of parental histones during replication [31,39,49,50,76-78]. To assess whether parental histone recycling was impaired in FRKO cells under replication stress, we combined iPOND-MS with chemical derivatization of histone treatment to examine histone modifications at replication forks (Figure 2A) [79]. Upon HU-induced replication stress, parental histone PTMs such as H4K20me2, H3K79me2, H3K27me3, and H3K9me3 were markedly reduced on nascent DNA in fork reversal–deficient cells (Figure 2B, Figure S3A-D). In contrast, histone PTMs characteristic of newly synthesized histone modifications [49,77], such as H4K12ac, H4K5ac, H3K56ac,H3K9ac were either increased or remained unchanged (Figure 2B, Figure S3E-H). To validate H3K9me3 reduction in iPOND-MS results, we performed PLA assays to examine H3K9me3 levels on nascent DNA. We found that H3K9me3 levels on nascent DNA are reduced, consistent with iPOND-MS results (Figure 2C).

H3K9me3 is a parental histone PTMs that are associated with constitutive heterochromatin, which is lost upon replication stress in FRKO cells. To distinguish whether the reduction of H3K9me3 arises from replication stress or from preexisting global changes in FRKO cells, we performed immunofluorescence analysis. A slight decrease in cellular H3K9me3 level was observed only in S phase FRKO cells following HU treatment, rather than unperturbed condition (Figure 2D–E), indicating that the loss of this repressive mark in fork reversal–deficient cells is specifically attributable to replication stress. These results indicate the loss of parental histone during DNA replication in FRKO cells under stressed conditions.

H3K9me3 and H3K27me3 are parental histone–specific PTMs that also indicate compact chromatin. We also detected a decrease in the abundance of another compaction marker, H3K9me2 on nascent DNA in fork reversal–deficient cells (Figure 2B, 2F, Figure S3I), while the cellular H3K9me2 levels decrease also depends on HU treatment (Figure S3J-K). These findings are consistent with our model in which nucleosome density is reduced on nascent DNA in FRKO cells upon replication stress.

PrimPol-mediated repriming causes nucleosome density reduction

Histone chaperones are responsible for the deposition of newly synthesized histones and the recycling of parental histones behind the replication fork. Impaired recruitment of histone chaperones to the replication fork could explain the robust decrease in nucleosome density and parental histone recycling observed in FRKO cells during replication stress. Analysis of histone chaperone localization from our SILAC-iPOND-MS analysis revealed that almost all known

histone chaperones properly localize to the replication fork in both WT and FRKO cells (Figure
3A), suggesting that the loss of parental histones and reduced nucleosome density are not due to
defects of histone chaperone recruitment.

We noted that PrimPol (Primase And DNA Directed Polymerase) levels at replication forks were
elevated after HU treatment in FRKO cells (Figure 1A), consistent with previous reports of
PrimPol activation under fork reversal defects[65,66,80]. We hypothesized that PrimPol activation may
contribute to loss of nucleosome density. Following HU treatment, FRKO cells exhibited elevated
ssDNA gaps on nascent DNA (Figure 3B). PrimPol depletion can rescue the ssDNA gaps,
suggesting PrimPol repriming is activated in FRKO cells and these gaps are PrimPol dependent
(Figure 3C). To test whether PrimPol activation was responsible for reduced nucleosome density
under replication stress, we depleted PrimPol in FRKO cells by siRNA or CRISPR-Cas9 and
measured nucleosome density on nascent DNA by PLA assay (Figure 3D-E). PrimPol depletion
in both conditions rescued low nucleosome density in FRKO cells. To determine if the reduction
in nucleosome density requires PrimPol catalytic activities, we reintroduced either wild-type
PrimPol or a catalytically inactive point mutant (AxA (catalytic inactive), CH (primase dead,
polymerase competent) mutants) into FR/PrimPol-knockout (FR/PPKO) cells. Only wild-type
PrimPol was sufficient to restore nucleosome density, demonstrating that PrimPol repriming
activity drives the loss of nucleosome density (Figure 3F-G).

Next, to determine if the parental-specific histone modification loss in FRKO cells results from
PrimPol repriming, we performed iPOND-MS for histone modifications in FR/PPKO cells (Figure
S4A-D). Our results demonstrate that PrimPol depletion restores H4K20me2 and H3K79me2
levels (Figure S4A-B) and reverses the increase of H4K5Ac and H3K56Ac in FRKO cells (Figure
S4C-D). While we did not detect H3K9me2/3 peptides in iPOND-MS, PLA assays for H3K9me2/3
on nascent DNA confirmed restoration of H3K9me2 and H3K9me3 levels (Figure 3H-I).

Collectively, these results demonstrate that when fork reversal is absent, PrimPol-mediated
repriming leads to loss of parental histones and a reduction in nucleosome density on nascent DNA
(Figure 3J).

Single-stranded DNA gaps cause low nucleosome density in cells

To determine if the low nucleosome density attribute to the elevated levels of ssDNA gaps that
mediated by PrimPol, or enriched PrimPol levels per se, we examined nucleosome density in other
previously defined ssDNA gaps contexts, such as BRCA1 or BRCA2 deficient cells. BRCA1 and
BRCA2 are key fork protection factors, and BRCA1- or BRCA2-deficiency cells combined with
HU treatment leads to nucleolytic degradation of nascent DNA and ssDNA gap formation; PrimPol
repriming is also activated in the absence of BRCA2[67,81-85]. We performed PLA assay for EdU and
H3 in BRCA1 deficient UWB1.289 cells, and UWB1.289 cells complement with wild-type
BRCA1. Compared to complementation cells, we observed low nucleosome density in UWB1.289
cells (Figure 4A, Figure S5A). Similarly, we also observed low nucleosome density in other
BRCA1- or BRCA2-deficient cells lines (Figure 4B–D, Figure S5B–D). These findings are
consistent with prior work showing that histone H3 levels are decreased on nascent DNA in
BRCA2 knockout HeLa cells[85].

Defects in Okazaki fragment maturation, such as knockdown of DNA ligase 1 (LIG1) or FEN1
(Flap Endonuclease 1), combining HU treatment, also result in ssDNA gaps on the lagging strand
in previous study[85]. We depleted LIG1 and FEN1 with siRNA in U2OS WT cells, and PLA signals

of H3 and EdU also reduced (Figure 4E-F, Figure S5E–F). Inhibition of RAD51 triggers
nucleolytic degradation of nascent DNA by nucleases such as MRE11, producing ssDNA gaps[84,86].
Therefore, we treated WT U2OS cells with RAD51 inhibitor B02 and observed low nucleosome
density, and this effect was rescued by co-treatment with MRE11 nuclease inhibitor (Figure 4G,
Figure S5G).

In summary, we conclude that ssDNA gaps present in post-replicative nascent DNA reduce the
nucleosome density on nascent DNA.

Single strand DNA gaps do not block nucleosome assembly in the in vitro assay

We hypothesized that single-stranded gaps in nascent DNA impair the ability to form nucleosomes.
The ability of nucleosomes to form on ssDNA remains controversial. During DSB resection, the
generation of long ssDNA tails/overhangs have been shown to evict nucleosomes in some
studies[87-93], whereas others have demonstrated nucleosome formation on ssDNA in vitro[94,95].
Additionally, it remains unclear whether nucleosomes can assemble on short ssDNA gaps like
those found at the replication fork and what impact these short ssDNA gaps have on nucleosome
stability. To determine whether nucleosomes can be assembled on DNA in the presence of ssDNA
gaps, we designed a series of Widom 601 strong positioning oligonucleotides containing ssDNA
gaps of various lengths (from 1–31 nts) positioned at the nucleosome dyed or the nucleosome
entry/exit site (Supplemental Table 1). We then reconstituted recombinant nucleosomes containing
human histones on these oligonucleotides containing ssDNA gaps using an established salt-
dialysis method[96-98] (Figure 5A). Strikingly, we observed robust nucleosome formation regardless
of the ssDNA gap size or ssDNA gap position within the nucleosome (Figure 5B-D), indicating
that small ssDNA gaps do not inhibit nucleosome formation *in vitro*.

Having established that short ssDNA gaps (1–31 nt) at either the entry/exit or dyad regions do not
markedly impair nucleosome assembly (Fig. 5B–D), we next asked whether longer gaps
approaching the footprint of a full nucleosome would influence histone wrapping or stability. To
address this, we generated a 147-nt ssDNA gap substrate flanked by a 25bp duplex regions with
Cy3 label, and a 15bp duplex regions with Cy5 label (Figure 5E). We performed a high-resolution
Cy3-Cy5 FRET (Förster Resonance Energy Transfer) assay to monitor nucleosome formation and
conformational dynamics in real time (Figure 5E). When a nucleosome assembles onto a 147-nt
ssDNA gap, the Cy3 and Cy5 fluorophores are brought into close proximity, resulting in a FRET
signal (Figure 5F-G). Thus, the FRET signal serves as a sensitive readout for nucleosome assembly
on ssDNA gaps. The steady-state FRET efficiency ($E_{FRET} \approx 0.27$) confirmed the proximity of the
labeled duplex arms (Figure 5F-G), consistent with a compact, fully wrapped configuration. These
results demonstrate that even extensive single-stranded gaps are compatible with nucleosome
formation, revealing that the histone octamer has an intrinsic ability to engage and stabilize flexible
ssDNA regions. Together, these experiments (Figure 5), consistent with previous reports[95],
demonstrate that ssDNA gaps do not inherently preclude nucleosome formation in vitro. The
discrepancy between the in vivo (Figure 3 and Figure 4) and in vitro (Figure 5) results suggests
that the reduced nucleosome density observed in cells is likely caused by ssDNA-binding proteins,
rather than by the presence of ssDNA itself (Figure S6A).

RPA depletion does not rescue the low nucleosome density in cells

Nucleosomes reconstituted on ssDNA (ssDNA Nucs) are similar in size and morphology to
those assembled on dsDNA (dsDNA Nucs) and the histone octamer core engages the wrapped

DNA with similar affinities in both complexes at physiological ionic strength[94,95,99-102]. However,
ssDNA Nucleosomes are more disordered and dynamic[95,100], suggesting that regions of the ssDNA
wrapped around a histone octamer core are dynamically exposed. The major ssDNA-binding
protein complex, Replication Protein A (RPA), is highly abundant (10 μM range) in the
nucleoplasm [103-105] and has the highest ssDNA affinity ($K_D \sim$ pM to nM range) of any ssDNA-
binding protein/protein complex that has been identified in human cells. Each RPA engages ~20-
33 nt of ssDNA, extending the bound sequence into a linear conformation and increasing its
bending 2 to 3-fold [106-117].

To determine whether RPA can substitute for nucleosomes on ssDNA, we performed parallel
in vitro and in vivo experiments. In vitro, we compared the FRET signal of RPA binding to the
ssDNA gaps only, and confirmed that RPA binding on ssDNA gaps can't cause FRET signal
(ssDNA gaps with RPA E_{FRET}=0.075±0.001, ssDNA gaps only E_{FRET}=0.08±0.001, Figure S6B).
Next, we incubate pre-assembled ssDNA nucleosomes with physiological concentrations of RPA,
and we observed a time-dependent decline in FRET efficiency (E_{FRET} decreasing from 0.25±0.001
to 0.119±0.001; Figure S6C).

Furthermore, E_{FRET} values decrease upon the addition of RPA down to an equilibrium value
(E_{FRET} = 0.119 + 0.001, Figure S6C) that is noticeably higher than the equilibrium E_{FRET} values
observed for the ssDNA Gap DNA substrate in the absence (0.080 + 0.001) and presence of RPA
(0.075 + 0.001) (Figure S6B). It cannot be discerned whether the elevated equilibrium value
(Compare the E_{FRET} in Figure S6C to E_{FRET} in Figure S6B) is due to indirect effects of the fully
unwrapped histone octamer on the RPA-ssDNA filament or the presence of a minor population of
stable, fully wrapped ssDNA nucleosomes or partially wrapped ssDNA Nucleosomes.
Nonetheless, the clear decrease in E_{FRET} observed upon the addition of RPA suggests that ssDNA
readily unwraps from around histone octamer cores and the exposed sequences are engaged and
stabilized by RPA.

We performed immunofluorescence staining for RPA in FRKO cells to corroborate the in vitro
findings, since we observed increased levels of RPA2 and its phosphorylation, consistent with
elevated ssDNA accumulation in FRKO cells upon HU treatment (Figure S5D and S6E). To test
whether elevated RPA contributes to the reduced nucleosome density, we conducted a PLA assay
for H3 and EdU in RPA1/2-depleted U2OS cells (Figure S6F). However, depletion of RPA1 and/or
RPA2 did not rescue the low nucleosome density in fork-reversal-deficient cells. These results
suggest that RPA does not directly evict nucleosomes from ssDNA in cells, or that other ssDNA-
binding factors may perform this function when RPA is depleted.

Transcription reduces nucleosome density in cells

Since we observed reduced chromatin compaction marks, such as H3K9me2/3 (Figure 2) and
lower nucleosome density in FRKO cells (Figure 1), we hypothesized that chromatin
decompaction may increase DNA accessibility to enzymes such as RNA polymerase II (Pol II).
Conversely, transcription elongation and R-loop formation could further disrupt nucleosomes on
nascent DNA. Thus, we proposed that RNA Pol II–mediated transcription on nascent DNA and
low nucleosome density may reinforce each other.

To test this hypothesis, we first examined RNA Pol II occupancy on nascent DNA using a PLA
assay targeting EdU and the major RNA Pol II subunit RPB1 (Figure S7A). We observed a
significant increase in PLA signal in FRKO cells, indicating elevated Pol II association with

nascent DNA. As a control, treatment with Triptolide (TPL), a potent RNA Pol II inhibitor that
induces proteasome-dependent degradation of Pol II, dramatically reduced the RPB1 PLA signal,
confirming the signal specificity for RNA Pol II (Figure S7A).

To determine whether increased transcription contributes to the reduced nucleosome density, we
performed a PLA for H3 and EdU in TPL-treated WT and FRKO cells. TPL treatment significantly
increased nucleosome density in both WT and FRKO cells, suggesting that Pol II activity on
nascent DNA can reduce nucleosome density (Figure S7B). However, TPL treatment did not fully
restore nucleosome density in FRKO cells to the levels in WT U2OS cells, indicating that
transcription alone cannot account for the nucleosome loss in FRKO cells, and additional fork
reversal dependent mechanisms likely contribute to this phenotype.

PARylation suppresses nucleosome assembly on ssDNA gaps in fork reversal deficient cells

PARylation is known to alter chromatin structure and orchestrate DNA repair[118-124]. We observed
the increase in PARylation in FRKO cells upon HU treatment (Figure 6A), suggesting increased
ssDNA gaps in FRKO cells. To determine if PARP1-mediated PARylation impacts nucleosome
density, we depleted PARP1 in FRKO cells via siRNA. PARP1 depletion reduced total level of
PARylation and partially rescued nucleosome density despite persistent ssDNA gaps (Figure 6B-
D). Inhibition of PARG, the enzyme that hydrolyzes PAR chains, increased PARylation and
decreased nucleosome density in WT cells, further supporting the conclusion that PARylation
reduces nucleosome density in cells (Figure 6E and Figure 6F). Interestingly, treatment with the
PARP1/2 inhibitor Olaparib reduced PARylation in WT cells after HU treatment (Figure 6G), but
nucleosome density still decreased (Figure 6H). This indicates that PARP inhibition alone can
lower nucleosome density on nascent DNA. In FRKO cells, Olaparib similarly reduced
PARylation but nucleosome density remains low, reinforcing that PARP1 inhibition itself causes
this effect. Together, these findings indicate that ssDNA activates PARP1-mediated PARylation,
which in turn reduces nucleosome density on nascent strand DNA. PARP1 trapping by Olaparib
can also reduce nucleosome density on nascent DNA.

HMCES–DPC inhibits nucleosome assembly on ssDNA

We sought additional factors that might bind ssDNA and inhibit nucleosome assembly. From our
iPOND-MS data, we noticed that in fork reversal–deficient cells, APOBEC3B levels at replication
forks increased by ~1.5-fold (Figure 7A), chromatin-bound APOBEC3B increased by 3–5-fold
(Figure S6A). APOBEC3B can deaminate cytosine residues, which are subsequently processed by
glycosylases to generate AP sites[125]. HMCES, a PCNA-binding protein, has been shown to form
covalent DNA–protein crosslinks (DPCs) at AP sites on ssDNA[126-132]. We observed a modest
increase in levels of HMCES in iPOND-MS (Figure 7A), and a significant increase in HMCES-
DNA crosslink by RADAR assay (Figure 7B).

We hypothesized that in FRKO cells, the high abundance of ssDNA on nascent DNA permits
excess APOBEC3B activity, leading to the generation of AP sites that subsequently form
HMCES–DPCs. These DPCs could represent a steric hindrance that blocks nucleosome assembly.
To test this hypothesis, we depleted HMCES or APOBEC3B using siRNA and measured
nucleosome density on nascent DNA by PLA (Figure 7C, Figure S8B–C). HMCES depletion
restored nucleosome density in fork reversal–deficient cells, whereas APOBEC3B depletion only
mild rescue the low nucleosome density, but not statistically significant (Figure 7C), likely because
AP sites can also form spontaneously on ssDNA independent of APOBEC3B. Co-deletion of

PARP1 and HMCES additively rescued nucleosome density in fork reversal–deficient cells
(Figure 7D). HMCES loss is also known to activate TLS (translesion synthesis) for post-replication
repair[127], which in turn repair ssDNA gaps[67,80]. We next asked whether the rescue of nucleosome
density upon HMCES depletion was mediated by TLS-dependent gap repair. Inhibition of TLS
using the REV1 inhibitor JH-RE-06 in HMCES-deficient cells did not reduce nucleosome density
(Figure 7E), demonstrating that HMCES–DPCs themselves, rather than TLS activation, block
nucleosome assembly.

HMCES-DPC at dyad position inhibits nucleosome formation *in vitro*

To determine if a HMCES-DPC can block nucleosome assembly *in vitro*, we designed two Widom
601 strong positioning oligonucleotides containing a 31 nt ssDNA gap positioned at the
nucleosome dyed or the nucleosome entry/exit site, with an uracil located in the ssDNA region
(Figure S8D-E). These DNA substrates were treated with uracil DNA glycosylase to generate an
AP-site, and the HMCES-AP-site DPC stabilized using sodium borohydride (Figure S8F-
G)[127,128,133]. We then attempted to reconstitute recombinant human nucleosomes on these
oligonucleotides containing the HEMCES-DPC using a salt-dialysis method (Figure 5A)[96-98].
Notably, we observed robust nucleosome formation when the HMCES-DPC was positioned at the
nucleosome entry/exit site (Figure 7F). In contrast, we observed minimal nucleosome formation
when the HMCES-DPC was positioned in the DNA near the nucleosome dyad, indicating that
HMCES represents a steric hindrance to nucleosome formation (Figure 7G). We hypothesize that
elevated nucleosomal DNA dynamics at the entry/exit site compared to the dyad enables the
nucleosome to accommodate the HMCES-DPC at this location[134]. Together, this data shows that
a HMCES-DPC can directly impair nucleosome formation, though the ability to accommodate a
HMCES-DPC will likely be dictated by the location of the HMCES-DPC in relation to the histone
octamer during histone deposition on ssDNA gaps.

Together, these findings demonstrate that in FRKO cells, PrimPol-mediated repriming generates
abundant ssDNA gaps, which activate PARP1-dependent PARylation and promote HMCES–DPC
formation. At least three mechanisms: Transcription, PARylation and HMCES–DPC formation
act in concert to inhibit nucleosome assembly on ssDNA gaps, thereby reducing parental histone
recycling and lowering nucleosome density on nascent DNA (Figure 7H).

Discussion

Fork reversal is essential for both genome stability and epigenome stability

In this study, we demonstrate that fork reversal is critical not only for genome stability, but also
for epigenome stability during DNA replication under stress. We propose a mechanistic model in
which fork reversal suppresses ssDNA gap formation, thereby promoting parental histone
recycling and nucleosome assembly on nascent DNA to preserve chromatin states. In the absence
of fork reversal, PrimPol-mediated repriming generates abundant ssDNA gaps, which trigger
PARP1-dependent PARylation and HMCES–DPC formation (Figure 6 and Figure 7). These events
hinder nucleosome deposition and lead to the loss of parental histone modifications (Figure 2 and
Figure 3). Together, our work establishes a mechanistic paradigm linking replication fork
intermediates (fork reversal and ssDNA) to faithful epigenetic inheritance, reframing replication
stress from a purely DNA-centric problem to a driver of epigenetic instability with broad
implications for human disease.

Loss of parental-specific histone marks on nascent DNA in FRKO cells reflects impaired parental histone recycling

A previous elegant study showed that HU treatment promotes accumulation of H3K9me2/3 at stalled forks, thereby helping protect stalled replication structures{Gaggioli, 2023 #157}. An important question is whether, in fork-reversal–deficient (FRKO) cells, the reduced abundance of parental-specific histone modifications on nascent DNA (e.g., H3K9me2/3) primarily reflects loss of parental histones, a failure to induce/enrich these marks at stalled forks, or a combination of both. Our data support the conclusion that the decrease at least partially reflects impaired parental histone recycling for two reasons. First, HU-induced enrichment of H3K9me2/3 at stalled forks is ATR-dependent and has been reported to occur independently of fork reversal{Gaggioli, 2023 #157}. Consistent with this, we observed even stronger ATR activation in FRKO cells (Figure S6E), suggesting that an inability to activate ATR is unlikely to explain the reduced parental-mark signal in our system. Second, we detected a broad reduction across multiple parental histone–associated signatures, not limited to H3K9me2/3, including H4K20me2, H3K79me2, and the enrichment of histone variants such as CENP-A and macroH2A.

In summary, these findings indicate that fork reversal deficiency leads to a loss of parental histone information on nascent DNA, consistent with defective parental histone recycling. However, we cannot fully exclude the possibility that FRKO cells also exhibit a reduced capacity to establish or enrich specific marks under HU compared with WT cells. These mechanisms are not mutually exclusive, and both impaired parental histone recycling and altered mark induction may contribute to the phenotype observed under sustained HU treatment.

How parental histone loss in FRKO cells

Our results suggest that PrimPol repriming–mediated ssDNA gaps contribute to reduced nucleosome density on nascent DNA. However, our data do not fully exclude an alternative possibility: that FRKO cells experience impaired histone delivery during DNA synthesis due to altered usage of histone-chaperone–coupled replicative polymerases. Current evidence does not support a direct histone chaperone activity for PrimPol. Thus, one speculative model is that, in FRKO cells, PrimPol extends nascent DNA in contexts where replicative polymerases with tighter histone-coupled synthesis (e.g., Pol ε) would normally operate, resulting in DNA extension that is relatively uncoupled from histone deposition. Although we did not detect an obvious change in Pol ε abundance by iPOND-MS (Figure 3A), we can't fully exclude this possibility.

Based on our findings together with prior studies, we conclude that ssDNA is not intrinsically incompatible with nucleosome assembly: from a biophysical standpoint, ssDNA can support the formation of ssNUCs. The critical unresolved issue is the temporal and spatial order of ssDNA formation relative to ssDNA-associated adducts, particularly DPCs and PARylation. Because we cannot determine when and where DPCs and PARylation arise with respect to ssDNA gaps, our data are consistent with two alternative, non–mutually exclusive models: 1) Pre-existing ssDNA regions are present in the genome prior to fork passage and are already decorated with DPCs and/or PARylation. In this scenario, as replication proceeds through these regions, parental histones would be unable to efficiently recycle onto the ssDNA substrate, resulting in an immediate deficit in nucleosome density on nascent DNA. 2) Replication itself generates abundant ssDNA gaps; ssNUCs can initially form on or near these ssDNA regions, but subsequent accumulation of DPCs and/or PARylation destabilizes these structures and promotes their eviction, thereby reducing nucleosome density.

Although our experiments show that HU treatment is associated with decreased nucleosome
density, they do not definitively exclude the first model. Notably, HU exposure in our assays is
sustained for ~4 hours, which could allow the establishment of pre-existing ssDNA lesions and
associated adducts before or during replication, making it difficult to unambiguously assign
causality and order of events.

Previous work has elegantly shown that under replication stress, ASF1 can sequester parental
histones during fork stalling and facilitate their recycling upon fork restart[136]. Here, we focused on
parental histone recycling during ongoing replication under stress and observed a pronounced loss
of parental histones in FRKO cells. One possibility is that excessive ssDNA gaps in FRKO cells
increase the frequency of parental histone disassembly, which in turn may overwhelm the
buffering capacity of ASF1, ultimately resulting in defective parental histone recycling and
reduced nucleosome density on nascent DNA.

Parental histones are lost during replication under stress in FRKO cells

These parental marks and variants (H4K20me2, H3K9me3, MacroH2A.1/2, CENP-A) are
generally repressive features that promote chromatin compaction and transcriptional silencing.
H3K9me3-marked heterochromatin, in particular, is enriched at repetitive genomic regions, where
it suppresses transcription[137]. Thus, the loss of parental histones in FRKO cells likely results in the
derepression of repetitive elements. Prior studies in mouse embryonic stem cells have shown that
disrupting parental-histone recycling via depletion of histone chaperones activates transposon
elements, thereby triggering inflammatory and immune signaling and impairing differentiation and
cell viability[49,50]. SMARCAL1 loss has been implicated increasing the response to immunotherapy
through activating cGAS-STING pathway, and suppressing PD-L1 express[138]. Our study provided
extra rationale that fork reversal can be targeted for immunotherapy since its loss may activate
repeat genes, therefore producing antigens for activating cytokines.

ssDNA gaps reduce the nucleosome density on nascent DNA

Notably, all our cellular assays are EdU labeling-based, as we use EdU label nascent DNA, such
as PLA, iPOND, and ChOR-seq. Therefore, these assays may not resolve ssDNA-nucleosomes
beyond the resolution distance of these assays. This assumption may explain the inconsistency
between in vivo and in vitro assays. However, the spatial resolution of PLA is approximately 40
nm (~100 bp/nt), while the DNA fragments analyzed by iPOND and ChOR-seq are typically 100–
400 bp/nt in length[72,139]. Previously published electron microscopy images showed that most
ssDNA gap sizes on nascent DNA are less than 300bp[53,66,67,80,140], in very rare cases can reach up
to 1000 nt. The ssDNA-nucleosomes are even closer to the EdU in dsDNA spatially after
nucleosome assembly (Figure 5E). Therefore, we hypothesize that cellular assays such as PLA,
ChOR-seq, and iPOND should have enough resolution to detect most of the ssDNA-nucleosome
occupancy, as long as it exists. However, developing higher-resolution assays will be necessary to
further confirm whether nucleosomes bind ssDNA directly in cells in the future.

ssDNA gap events that inhibit histone assembly

We conclude that PARylation, transcription, and DPC are three events that reduce nucleosome
density. PARP1-mediated PARylation promotes chromatin relaxation via multiple mechanisms: 1)
the negatively charged poly(ADP-ribose) (PAR) chains neutralize histone positive charges and
weaken histone–DNA interactions[122]; and 2) PAR chains act as docking platforms to recruit DNA

repair enzymes and chromatin remodelers, such as ALC1 (CHD1L)[119,120,141-143], which remodels
nucleosomes. Together, these mechanisms enhance chromatin accessibility, exposing ssDNA
regions to repair factors. Surprisedly, we found that PARP inhibitor Olaparib treatment reduce the
nucleosome density on nascent DNA, which is distinct from PARP1 depletion. We hypothesize
that, PARP1 depletion removes PARylation and leaves ssDNA gaps relatively "clean," so the
nucleosome can form on the ssDNA gaps. However, PARP inhibitor blocks PARylation and trap
PARP1 on DNA, since PARylation normally promotes PARP1 release[145,146]. We therefore propose
that trapped PARP1 directly displaces nucleosomes. This idea is consistent with recent findings
showing that PARP1 inhibition leads to histone eviction from replication fork [147].

We examined the role of HMCES-DPCs, which were significantly increased in FRKO cells
(Figure 7B). HMCES depletion rescued the low nucleosome density in FRKO cells, as shown by
PLA (Figure 7C), and this was further confirmed by in vitro reconstitution assays (Figure 7F-G).
Surprisingly, HMCES-DPCs impaired nucleosome assembly only when positioned near the
nucleosome dyad (Figure 7G), not at the entry or exit sites (Figure 7H). However, we cannot rule
out inhibitory effects of entry/exit DPCs in cells. This is because the discrepancy between in vivo
and in vitro system: (1) our reconstitution assays involve single nucleosomes, where ssDNA gaps
at the entry/exit sites are flexible and may minimize DPC effects; (2) the Widom 601 sequence
was specifically selected for nucleosome formation assay in previous studies, which has high
histone-binding affinity, which could overcome HMCES-DPC–mediated inhibition; and (3) in
cells, HMCES-DPCs can undergo additional processing, such as ubiquitination or proteolytic
degradation, triggering downstream repair cascades that may collectively reduce nucleosome
density[127,144]. These complex in vivo events are not captured by simplified in vitro systems.

We cannot exclude other ssDNA binding proteins, or any ssDNA-related events that also
compromise nucleosome formation. Therefore, we conclude that PARylation, RNA Pol II, and
DPC are the least events that inhibit histone assembly.

Materials and Methods

Cell lines

U2OS and MDA-MB-436 cells were cultured in DMEM with 9% fetal bovine serum (FBS) (Bio-
techne). PEO4, PEO1, and DLD-1 cells were cultured in RPMI-1640 with 9% FBS. UWB1.289
cells were cultured in a 1:1 mixture of RPMI-1640 medium and Mammary Epithelial Growth
Medium (MEGM), supplemented with FBS and antibiotics. For iPOND-MS, U2OS cells were
cultured in SILAC-compatible DMEM (Thermo Fisher Scientific A33822) with 10% dialyzed
FBS (R&D Systems S12850H) and either isotopically light or heavy 13C, 15N lysine, and
13C,15N arginine (Cambridge Isotope Laboratories)[70]. Cells were cultured at 37°C and 5% CO2
with humidity. All cell lines were regularly tested for mycoplasma and verified using short tandem
repeat profiling.

Genome editing

PrimPol knockout was generated using CRISPR-Cas9 with the SpCas9(BB)-2A-Puro (PX459)
V2.0 plasmid (Addgene #62988) as described[148]. The single guide RNA (sgRNA) sequence was
5'- CTGCCAAGCAAGTCAAGAGC-3'. Transfected cells were selected with puromycin, and
single-cell clones were isolated and validated by sequencing and immunoblotting.

SILAC and Label free iPOND-mass spectrometry

iPOND-SILAC MS was performed as described previously[70]. Briefly, EdU-labeled and indicated
drug-treated cells were crosslinked with 1% formaldehyde, quenched with glycine, permeabilized
in 0.25% Triton X-100. Heavy-labeled and light-labeled cells were combined 1:1 prior to the click
reaction. DNA-protein complexes were captured with Dynabeads MyOne Streptavidin C1
(Thermo Fisher 65002), washed, and eluted with elution buffer (25mM biotin, 0.4% SDS, 50mM
Tris-HCl (pH 8.5)) at 65°C. Supernatant, containing eluted fractions, were incubated at 95°C to
reverse crosslinking. For histone modification analysis, lysine were derivatized with 2.5%
propionic anhydride (Sigma-Aldrich 8.00608) in 50 mM ammonium bicarbonate (pH 7.5).
Subsequently, proteins were digested with 200 ng trypsin (Thermo Fisher 90057) in 50 mM
ammonium bicarbonate (pH7.5) overnight. Peptides were desalted using C18 Stagetips (reversed-
phase resin) and carbon Top-Tips (Glygen TT1CAR), speed-vacuumed dried, and stored at −20°C
until MS analysis.

PrimPol complementation

PrimPol WT, AxA, and CH mutants were subcloned from pcDNA3.1/nV5-DEST to pLNCX2
using forward primer: 5'-CAGATCTCGAGATGAATAGAAAATGGGAAGCAA-3' and reverse
primer: 5'-CTGCGGCCGCCTCTTGTAATACTTCTATAATTAG-3'.   pLNCX2-Primpol
constructs and pVSV-G were co-transfected into GP2-293 cells using Polyethylenimine (PEI).
Twenty-four hours post-transfection, the culture medium was replaced with fresh medium, and
viral supernatants were harvested at 48 and 72 hours. The pooled viral supernatants were
subsequently used to infect U2OS Primpol knockout #206 cells. Stable cell lines were established
by selection with 100 µg/ml G418.

DNA combing assays

536 Cells were labeled with the nucleoside analogs as indicated in the figures and figure legends. For
537 ssDNA gap detection, S1 nuclease digestion assay was performed as previously described[66]. DNA
538 combing was performed using a DNA combing instrument as previously described[54].

Immunofluorescence

For analysis of proteins in the insoluble/chromatin fraction, cells were pre-extracted with 0.5%
Triton X-100 and subsequently fixed in 4% paraformaldehyde. Slides were blocked with 1% goat
serum in PBST (0.1% triton-X100) and incubated with primary antibodies. EdU incorporation was
visualized by click chemistry with an Alexa Fluor 488-conjugated azide. Immunofluorescent
images were obtained and analyzed by ImageXpress microscopy and software (Molecular
Devices).

PLA assay

Proximity ligation assays (PLAs) were performed using the Duolink kit (Sigma DUO92002,
DUO92004, DUO92008) according to the manufacturer's instructions, with the following
modifications as described previously [71]. Cells were pre-extracted and fixed as described in the
immunofluorescence method, then blocked with 3% BSA and processed by click chemistry with
biotin-azide and Alexa Fluor 488-conjugated azide (9:1 ratio). The following primary antibody
pairs were used: rabbit H2A (CST 12349) (1:100 dilution), H3 (Abcam ab1791) (1:200 dilution),
H3K9ac (Millipore 07-352) (1:100 dilution), H3K9me3 (CST 13969) (1:100 dilution), H4 (CST
13919) (1:100 dilution); mouse biotin (Millipore B7653) (1:200 dilution) and mouse H2B (Abcam
ab52484) (1:100 dilution), H3K9me2 (Abcam ab1220) (1:100dilution) and rabbit biotin (CST

5597) (1:100 dilution). Images were obtained by ImageXpress Pico and analyzed by MetaXpress
(Molecular Devices).

CHOR-Seq

ChoR-seq was performed as previous described[72]. Briefly, WT and FRKO U2OS cells (n = 3
biological replicates each) were labeled with 10 µM EdU for 15 min for no HU sample. For HU
treated sample, the cells are treated with 10uM EdU and 1 mM HU (Sigma H8627) for 4 h before
fixation with the truChIP kit (Covaris 520127). Chromatin was sonicated to 100–500 bp, and EdU-
labeled Drosophila S2 chromatin was added at 0.05% (w/w) as a spike-in for normalization. H3
ChIP was performed with anti-H3 (10 µg, Abcam ab1791). Immunoprecipitated DNA was
processed with the NEBNext Ultra II DNA Library Prep Kit (NEB E7645S), indexed, and
sequenced on an Illumina NovaSeq platform.

Data processing

All sequencing reads were mapped to the H. sapiens reference genome (hg38) and D. melanogaster
reference genome (BDGP6) using Bowtie2 (v2.5.4) with the default parameters[149]. Only uniquely
mapping reads were kept, and PCR duplicates were removed using Markup from Samtools (v1.17)
(https://www.htslib.org/doc/samtools-markdup.html). Genome occupancy was calculated using
bamCoverage from Deeptools (v3.5.5)[150] with standard parameters in 100 bp intervals, normalized
to relative reads per million (RRPM) using D. melanogaster reads as described previously[151], and
visualized by the Integrative Genomics Viewer (IGV) (v2.18)[152]. To generate a common H3 peak
dataset, peak calling in each sample were preformed using MACS2 (v2.2.7)[153] with parameters of
"--broad-cutoff 0.1" and merged into a consensus region set using the Intersect function in
Bedtools (v2.26.0)[154]. Subsequently, for H3 overlapped peaks in RRPM-normalized datasets were
exported with the computeMatrix tool from Deeptools, and the enrichment patterns were
visualized using the plotProfile and plotHeatmap function in DeepTools. Lastly, using the Intersect
function in bedtools to get the H3 distribution on genome repeat elements and visualize in R. All
datasets were deposited in the NCBI GEO: GSE000000.

Preparation of oligonucleotides for recombinant nucleosomes

All oligonucleotides were synthesized by Integrated DNA Technologies (Coralville, IA). A
complete list of oligonucleotides used to make recombinant nucleosomes can be found in
Supplementary Table 1. Each oligonucleotide was resuspended to a final concentration of 100 µM
in a buffer containing 10 mM Tris (pH-7.5) and 1 mM EDTA. Complimentary oligonucleotides
(see Supplementary Table 1) were mixed to a final concentration of 10 µM and annealed by heating
to 95°C for 2 minutes and cooling to 10 °C at a rate of -5 °C/min. The annealed oligonucleotides
were stored short-term at 4 °C prior to nucleosome reconstitution.

Radar Assay

Cells were lysed in Radar buffer (4M Guanidine thiocyanate, 1% Sarkosyl, 2% Triton X-100, 1%
1,4-dithioerythritol, 100 mM Sodium Acetate pH5.0, 20 mM Tris pH 8.0, 20 mM EDTA pH8.0;
adjusted to pH 6.5). Genomic DNA was sonicated and precipitated with ethanol, resuspended in 8
mM NaOH, and quantified using Qbit fluorometry. Samples were then subjected to slot-blot onto
nitrocellulose membrane, which were subsequently immunoblotted with HMCES primary
antibody (Cell Signaling #17636) and corresponding secondary antibody.)

RNA interference

All siRNA transfections were performed using DharmaFECT reagents (Horizon Discovery T-2001-03) according to the manufacturer's instructions. Experiments were performed 2 days after transfection. Qiagen AllStars Negative Control Nontargeting (NT) siRNA was used as a non-targeting control. Gene-targeting siRNA sequences are provided in the supplementary table

Purification of recombinant human histones

The genes encoding human histone H2A (UniProt identifier: P0C0S8), H2B (UniProt identifier: P62807), H3 C110A (UniProt identifier Q71DI3), and H4 (Uniprot identifier: P62805) were cloned into a pET3a expression vector. Histone H2A, H3, and H4 were individually transformed and expressed in BL21 (DE3) pLysS competent *E. coli* cells (New England BioLabs). Histone H2B was transformed and expressed in BL21-CodonPlus (DE3)-RIPL *E. coli* cells (Agilent). The histones were grown in M9 minimal media (33.5 mM $Na_2HPO_4 \bullet 7H_2O$, 220 mM KH_2PO_4, 100 mM NaCl, 200mM NH_4Cl, pH 7.2) supplemented with 0.4% glucose (w/v), 2 mM $MgSO_4$, 0.2 mM $CaCl_2$ and a 1% vitamin cocktail. The histones were grown at 37 °C to an OD600 of 0.4, and histone expression induced with a 0.4 mM (histone H2A, H2B, and H3) or 0.3 mM (histone H4) isopropyl- β-D-thiogalactoside (IPTG) for 3 hours (histone H3 and H4) or 4 hours (histone H2A and H2B). The cells were harvested via centrifugation and cell pellets stored long term at −80 °C. The individual histones were purified from inclusion bodies using well-established methods[96-98]. In brief, the histone cell pellets were lysed via sonication in a buffer containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM benzamidine, 1 mM DTT, and 1 mM EDTA and the cell lysate clarified via centrifugation. The resulting pellet was washed three times with a buffer containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM benzamidine, 1 mM DTT, 1 mM EDTA, and 1% Triton X-100 and washed a final time with a buffer containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM benzamidine, 1 mM DTT, and 1 mM EDTA. The histones were extracted from inclusion bodies under denaturing condition (6 M Guanidinium-HCl), dialyzed into 8 M Urea, and purified using anion-exchange and cation-exchange chromatography under gravity flow. The purified histones were dialyzed five times against H_2O, lyophilized, and stored long-term at −20 °C.

Preparation of H2A/H2B Dimer and H3/H4 Tetramer

Prior to nucleosome assembly, H2A/H2B dimers and H3/H4 tetramers were prepared using established methods [96-98]. Each individual lyophilized histone was resuspended in a buffer containing 20 mM Tris-HCl (pH 7.5), 6 M Guanidinium-HCl, and 10 mM DTT. To refold the H2A/H2B dimer, H2A and H2B were mixed in a 1:1 molar ratio and dialyzed three times against a buffer containing 20 mM Tris-HCl (pH 7.5), 2 M NaCl, and 1 mM EDTA at 4 °C. To refold the H3/H4 tetramer, H3 and H4 were mixed in a 1:1 molar ratio and dialyzed three times against a buffer containing 20 mM Tris-HCl (pH 7.5), 2 M NaCl, and 1 mM EDTA at 4 °C. The refolded H2A/H2B dimer and H3/H4 tetramer were subsequently purified using a HiPrep Sephacryl S-200 16/60 HR gel filtration column (Cytiva) in a buffer containing 20 mM Tris-HCl (pH 7.5), 2 M NaCl, and 1 mM EDTA. Pure fractions containing H2A/H2B dimers and H3/H4 tetramer were combined and stored long term at −20 °C in a 50% glycerol slurry.

Nucleosome assembly and purification

All nucleosomes were prepared using a well-established salt-dialysis method [96-98], with minor modifications. In brief, DNA, H2A/H2B dimer, and H3/H4 tetramer were mixed at a ratio of

1:2.2:1:1, respectively, in a buffer containing 20 mM Tris-HCl (pH 7.5), 2 M NaCl, 1 mM EDTA,
and 1mM DTT. Nucleosomes were assembled via a stepwise decrease in NaCl concentration over
a 24-hour period - from 2 M NaCl to 1.5 M NaCl, to 1.0 M NaCl, to 0.5 M NaCl, to 0.25 M NaCl,
and to 0 M NaCl. The assembled nucleosomes were then purified via ultracentrifugation for 42 to
44 hours at 4 °C using a 10% - 40% sucrose gradient. Final nucleosome purity and homogeneity
was assessed by native PAGE (5%, 59:1 acrylamide:bis-acrylamide ratio). The purified
nucleosomes were stored in a buffer containing 10 mM Tris (pH-7.5) and 1 mM EDTA at 4 °C.

Purification of recombinant UDG and HMCES

A pET-His-GFP vector (N-terminal His-GFP tag) with the human uracil DNA glycosylase (UDG)
gene was obtained from GenScript. The pET-His-GFP-UDG vector transformed into BL21-
CodonPlus (DE3)-RIPL *E. coli* cells (Agilent). The transformed cells were grown at 37 °C to an
OD600 of 0.6 and UDG expression induced using 1.0 mM IPTG for 18 h at 18 °C. The cells were
lysed via sonication in a buffer containing 25 mM HEPES (pH 7.5), 200 mM NaCl, 0.1 mM TCEP,
20 mM Imidazole, and a cocktail of protease inhibitors (AEBSF, leupeptin, benzamidine, pepstatin
A). The lysate was clarified via centrifugation and loaded onto a HisTrap HP column (Cytiva)
equilibrated with 25 mM HEPES (pH 7.5), 200 mM NaCl, 0.1 mM TCEP, and 20 mM Imidazole,
and eluted in a buffer 25 mM HEPES (pH 7.5), 200 mM NaCl, 0.1 mM TCEP, and 400 mM
Imidazole. The UDG protein was then liberates from the His-GFP tag via TEV protease and further
purified using a HiPrep SP HP column (Cytiva) equilibrated with 25 mM HEPES (pH 7.5), 50 mM
NaCl, and 0.1 mM TCEP, and eluted in a buffer 25 mM HEPES (pH 7.5), 1 M NaCl, and 0.1 mM
TCEP. The resulting UDG protein was combined and purified via size exclusion chromatography
using a HiPrep Sephacryl S-200 16/60 HR (Cytiva) in a buffer containing 20 mM HEPES (pH 7.5),
150 mM NaCl, and 0.1 mM TCEP. The purity of the UDG proteins were confirmed via denaturing
SDS-PAGE and the purified UDG protein stored long term at −80 °C.

Recombinant human HMCES protein was purified using a previously described protocol[155].
In brief, a pNIC-HMCES vector was transformed in *E. coli* cells. Cells were grown in terrific broth
(TB) medium at 37 °C to an OD_{600} of 0.7 and HMCES expression induced by the addition of
0.5 mM IPTG for 4 hrs. Cell pellets were collected via centrifugation and resuspended in a buffer
containing 20 mM HEPES/KOH (pH 7.5), 500 mM KCl, 5 mM $MgCl_2$, 30 mM imidazole, 10%
glycerol, 0.1% IGEPAL, 0.04 mg/ml Pefabloc SC, 1 mM TCEP and cOmplete EDTA-free
protease inhibitor cocktail. The lysate was loaded onto a Strep-Tactin® XT Superflow® high-
capacity cartridge, washed with a buffer containing 20 mM HEPES/KOH pH 7.5, 500 mM KCl, 5
mM $MgCl_2$, 1 mM TCEP, and eluted in a buffer containing 20 mM HEPES/KOH pH 7.5, 500 mM
KCl, 5 mM $MgCl_2$, 1 mM TCEP, and 50 mM biotin. HMCES was further purified using a HiTrap
Heparin HP columns and eluted with buffer containing 20 mM HEPES/KOH pH 7.5, 1 M KCl, 5
mM $MgCl_2$, 1 mM TCEP. The HMCES protein was combined and further purified using a HiPrep
Sephacryl S-200 16/60 HR (Cytiva) in a buffer containing 20 mM HEPES/KOH pH 7.8, 150 mM
KCl, 5 mM $MgCl_2$, 10% glycerol, 1 mM TCEP. The purity of the HMCES protein was confirmed
via denaturing SDS-PAGE and purified HMCES protein stored long term at −80 °C.

Generation of HEMCES-DPC and HEMCES-DPC nucleosome formation

AP-site DNA was generated via enzymatic digestion of dU-containing oligos (see Supplementary
Table 1. The dU-containing oligonucleotides (10 µM) were incubated with recombinant UDG (10
µM) for 2 hours at 25 ºC to excise the uracil and generate an AP-site. The oligonucleotides were
then incubated with recombinant HMCES (20 µM) for 2 hours at 25 ºC, and the HMCES-DPC

stabilized via sodium borohydride (NaBH$_4$) crosslinking for an additional 2 hours at 25 °C. The
HEMCES-DPC-DNA was loaded onto a HiPrep Q HP column (Cytiva) equilibrated with 50 mM
HEPES (pH 7.5) and 400 mM NaCl, and eluted in a buffer 50 mM HEPES (pH 7.5) and 2 M NaCl.
Fractions containing the HMCES-DPC-DNA were combined, exchanged into a buffer containing
20 mM Tris-HCl (pH 7.5), 1 mM EDTA, 1 M NaCl, and 1mM DTT, and stored short-term at 4 °C
prior to nucleosome reconstitution.

To generate HMCES-DPC-DNA nucleosomes, HMCES-DNA, H2A/H2B dimer, and H3/H4
tetramer were mixed at a ratio of 1:2.2:1, respectively, in a buffer containing 20 mM Tris-HCl (pH
7.5), 1 M NaCl, 1 mM EDTA, and 1mM DTT. Nucleosomes were assembly via a stepwise decrease
in NaCl concentration over a 24-hour period - from 1 M NaCl to 0.8 M NaCl, to 0.6 M NaCl, to
0.4 M NaCl, to 0.2 M NaCl, and to 0 M NaCl. The formation of the HMCES-DPC nucleosomes
was assessed via native PAGE gels (5%, 59:1 acrylamide:bis-acrylamide ratio).

FRET assay

Oligonucleotides in FRET assay

Oligonucleotides comprising the single strand DNA (ssDNA) gap substrate (**Table 1**) were
synthesized by Integrated DNA Technologies (Coralville, IA) or Bio-Synthesis (Lewisville, TX)
and purified on denaturing polyacrylamide gels. The concentrations of unlabeled DNAs were
determined from the absorbance at 260 nm using the provided extinction coefficients.
Concentrations of Cyanine-labeled DNAs were determined from the extinction coefficient of their
respective label (Cy3 at 550 nm, $\varepsilon_{550} = 150{,}000$ M^{-1}cm^{-1}, Cy5 at 650 nM, $\varepsilon_{650} = 250{,}000$ M^{-1}cm^{-1}).
For annealing the ssDNA gap substrate, the template strand and the cyanine-labeled primer strands
were mixed in equimolar amounts in 1X annealing buffer (10 mM TrisHCl, pH 8.0, 100 mM NaCl,
1 mM EDTA), heated to 95 °C for 5 minutes, and allowed to slowly cool to room temperature.

Recombinant Human Proteins in FRET assay

Human RPA was obtained as previously described [156]. The concentration of active RPA was
determined via a FRET-based activity assay as described previously [109]. Human recombinant
histone H3.1 – H4 tetramers (Cat. # 14-1056) and histone H2A-H2B dimers (Cat. # 10-1052) were
purchased from EMD Millipore (Temecula, CA) and utilized to reconstitute nucleosomes on the
ssDNA gap substrate via stepwise salt dilution dialysis method, as described previously [157].
Assembled nucleosomes were purified and verified by native PAGE, as described previously [157].

FRET Experiments

All experiments were performed in a 16.100F-Q-10/Z15 sub-micro fluorometer cell (Starna
Cells) at room temperature (23 ± 2 °C) in 1X Replication Buffer (25 mM TrisHCl, pH 7.5, 125
mM KOAc, 10 mM Mg(OAc)$_2$) supplemented with 1 mM DTT, 0.1 mg/mL BSA, and the ionic
strength adjusted to physiological (200 mM) by addition of KOAc. Fluorescence intensities were
monitored in a Horiba Scientific Duetta-Bio fluorescence/absorbance spectrometer with excitation
and emission slit widths set at 10 nm. For all experiments described below the final concentrations
of all reaction components are indicated. The concentration of RPA (6.76 μM) utilized in the
experiments described below agrees with the concentration of RPA in the nucleoplasm of a human
cell (10 μM range)[103-105].

For steady state (equilibrium) FRET assays in Figure S5B, a DNA substrate (100 nM ssDNA
Gap or ssDNA Gap Nuc) is pre-incubated for 5 minutes in the absence or presence of RPA (6.76
μM heterotrimer). Under these conditions, the concentration of RPA is ~11-fold higher than the
concentration of RPA-binding sites within the the ssDNA Gap DNA substrate. The solution is then
transferred to a fluorometer cell and the cell is placed in the instrument. Next, the solution is
excited at 514 nm and the fluorescence emission intensities (I) are monitored essentially
simultaneously ($\Delta t = 0.118$ ms) at the peak emission wavelengths for Cy3 (563 nm, I_{563}) and Cy5
(665 nm, I_{665}) over time until both stabilize for at least 1 min. The I_{665} and I_{563} values within this
stable region are utilized to calculate the approximate FRET efficiencies (E_{FRET}) from the equation
$E_{FRET} = \dfrac{I_{665}}{I_{665}+I_{563}}$ and the resultant values are averaged to obtain the E_{FRET} value observed for a
given condition at equilibrium. Finally, the solution is excited at 514 nM and the fluorescence
emission spectra is recorded from 530 to 750 nm.

For pre-steady state FRET assays (Figure S5C) ssDNA Gap Nuc (100 nM) is transferred to a
fluorometer cell and the cell is placed in the instrument. The solution is excited at 514 nm and I_{563}
and I_{665} are monitored over time (as described above) until both stabilize for at least 1 min. The
I_{665} and I_{563} values within this stable region are utilized to E_{FRET} and the resultant values are
averaged to obtain the E_{FRET} value observed prior to the addition of RPA. Next, RPA (6.76 μM
heterotrimer) is added, the resultant solution is mixed, and I_{665} and I_{563} are monitored beginning
30 s after the addition of RPA. All recorded fluorescence emission intensities are corrected by a
respective dilution factor. For all recordings of the fluorescence emission intensities (I_{665} and I_{563}),
E_{FRET} values are calculated as described above and plotted as a function of time after RPA is added.

For all plots in all figures, each data point/column/spectra represent the average ± standard
error of the mean (SEM). Error bars are present for all data points on all plots in all figures but
may be smaller than the data point.

Quantification and statistical analysis

The statistical test was performed using GraphPad Prism and R software. The statistics for the
high-content microscopy experiments was performed on the mean of individual replicates.
Statistical analyses were completed using Prism. A Kruskal-Wallis test was used for experiments
with more than two samples, and P values were calculated by Prism for the multiple comparisons.
A two-tailed t test was used to compare two samples with normally distributed data. No statistical
methods or criteria were used to estimate sample size or to include/exclude samples. Statistical
details of individual experiments can be found in the figure legends and in Results. Experiments
shown are representative of at least two biological replicates unless otherwise indicated in the
figure legend.

Reference

1   Stewart-Morgan, K. R., Petryk, N. & Groth, A. Chromatin replication and epigenetic cell
memory. *Nat Cell Biol* **22**, 361-371 (2020). https://doi.org/10.1038/s41556-020-0487-y

2   Alabert, C. & Groth, A. Chromatin replication and epigenome maintenance. *Nature reviews*
Molecular cell biology **13**, 153-167 (2012).

3   Berti, M., Cortez, D. & Lopes, M. The plasticity of DNA replication forks in response to
clinically relevant genotoxic stress. *Nature reviews Molecular cell biology* **21**, 633-651
(2020).

4   Allis, C. D. & Jenuwein, T. The molecular hallmarks of epigenetic control. *Nature reviews*
genetics **17**, 487-500 (2016).

5   Cantone, I. & Fisher, A. G. Epigenetic programming and reprogramming during
development. *Nat Struct Mol Biol* **20**, 282-289 (2013). https://doi.org/10.1038/nsmb.2489

6   Suva, M. L., Riggi, N. & Bernstein, B. E. Epigenetic reprogramming in cancer. *Science*
339, 1567-1570 (2013). https://doi.org/10.1126/science.1230184

7   Iacobuzio-Donahue, C. A. Epigenetic changes in cancer. *Annual Review of Pathology:*
Mechanisms of Disease **4**, 229-249 (2009).

8   Feinberg, A. P., Ohlsson, R. & Henikoff, S. The epigenetic progenitor origin of human
cancer. *Nature Reviews Genetics* **7**, 21-33 (2006). https://doi.org/10.1038/nrg1748

9   Feinberg, A. P., Koldobskiy, M. A. & Göndör, A. Epigenetic modulators, modifiers and
mediators in cancer aetiology and progression. *Nature Reviews Genetics* **17**, 284-299
(2016). https://doi.org/10.1038/nrg.2016.13

10   Parreno, V. *et al.* Transient loss of Polycomb components induces an epigenetic cancer fate.
Nature **629**, 688-696 (2024). https://doi.org/10.1038/s41586-024-07328-w

11   Easwaran, H., Tsai, H.-C. & Baylin, S. B. Cancer epigenetics: tumor heterogeneity,
plasticity of stem-like states, and drug resistance. *Molecular cell* **54**, 716-727 (2014).

12   Li, N. *et al.* Parental histone transfer caught at the replication fork. *Nature* **627**, 890-897
(2024).

13   Zhou, K., Gaullier, G. & Luger, K. Nucleosome structure and dynamics are coming of age.
Nature structural & molecular biology **26**, 3-13 (2019).

14   Sekine, S. I., Ehara, H., Kujirai, T. & Kurumizaka, H. Structural perspectives on
transcription in chromatin. *Trends Cell Biol* **34**, 211-224 (2024).
https://doi.org/10.1016/j.tcb.2023.07.011

15   Liu, Y. *et al.* FACT caught in the act of manipulating the nucleosome. *Nature* **577**, 426-431
(2020). https://doi.org/10.1038/s41586-019-1820-0

16   Groth, A., Rocha, W., Verreault, A. & Almouzni, G. Chromatin challenges during DNA
replication and repair. *Cell* **128**, 721-733 (2007). https://doi.org/10.1016/j.cell.2007.01.030

17   Abe, T. *et al.* The histone chaperone facilitates chromatin transcription (FACT) protein
795   maintains normal replication fork rates. *Journal of Biological Chemistry* **286**, 30504-30512
796   (2011).

797   18   Tsunaka, Y., Fujiwara, Y., Oyama, T., Hirose, S. & Morikawa, K. Integrated molecular
798   mechanism directing nucleosome reorganization by human FACT. *Genes & development*
799   **30**, 673-686 (2016).

800   19   Wang, T. *et al.* The histone chaperone FACT modulates nucleosome structure by tethering
801   its components. *Life science alliance* **1** (2018).

802   20   Wang, X. *et al.* The N-terminus of Spt16 anchors FACT to MCM2–7 for parental histone
803   recycling. *Nucleic acids research* **51**, 11549-11567 (2023).

804   21   Escobar, T. M. *et al.* Active and repressed chromatin domains exhibit distinct nucleosome
805   segregation during DNA replication. *Cell* **179**, 953-963. e911 (2019).

806   22   Madamba, E. V., Berthet, E. B. & Francis, N. J. Inheritance of histones H3 and H4 during
807   DNA replication in vitro. *Cell reports* **21**, 1361-1374 (2017).

808   23   Schlissel, G. & Rine, J. The nucleosome core particle remembers its position through DNA
809   replication and RNA transcription. *Proceedings of the National Academy of Sciences* **116**,
810   20605-20611 (2019).

811   24   Sauer, P. V. *et al.* Mechanistic insights into histone deposition and nucleosome assembly
812   by the chromatin assembly factor-1. *Nucleic Acids Res* **46**, 9907-9917 (2018).
813   https://doi.org/10.1093/nar/gky823

814   25   Stillman, B. Chromatin assembly during SV40 DNA replication in vitro. *Cell* **45**, 555-565
815   (1986).

816   26   Smith, S. & Stillman, B. Purification and characterization of CAF-I, a human cell factor
817   required for chromatin assembly during DNA replication in vitro. *Cell* **58**, 15-25 (1989).

818   27   Shibahara, K.-i. & Stillman, B. Replication-dependent marking of DNA by PCNA
819   facilitates CAF-1-coupled inheritance of chromatin. *Cell* **96**, 575-585 (1999).

820   28   Groth, A. *et al.* Regulation of replication fork progression through histone supply and
821   demand. *science* **318**, 1928-1931 (2007).

822   29   Richet, N. *et al.* Structural insight into how the human helicase subunit MCM2 may act as
823   a histone chaperone together with ASF1 at the replication fork. *Nucleic acids research* **43**,
824   1905-1917 (2015).

825   30   Wang, H., Wang, M., Yang, N. & Xu, R.-M. Structure of the quaternary complex of histone
826   H3-H4 heterodimer with chaperone ASF1 and the replicative helicase subunit MCM2.
827   *Protein & cell* **6**, 693-697 (2015).

828   31   Alabert, C. *et al.* Two distinct modes for propagation of histone PTMs across the cell cycle.
829   *Genes & development* **29**, 585-590 (2015).

830   32   Reverón-Gómez, N. *et al.* Accurate recycling of parental histones reproduces the histone
831   modification landscape during DNA replication. *Molecular cell* **72**, 239-249. e235 (2018).

33 Reinberg, D. & Vales, L. D. Chromatin domains rich in inheritance. *Science* **361**, 33-34
(2018).

34 Oksuz, O. *et al.* Capturing the onset of PRC2-mediated repressive domain formation.
Molecular cell **70**, 1149-1162. e1145 (2018).

35 Alabert, C. *et al.* Domain model explains propagation dynamics and stability of histone
H3K27 and H3K36 methylation landscapes. *Cell reports* **30**, 1223-1234. e1228 (2020).

36 López, V. G. *et al.* Read–write mechanisms of H2A ubiquitination by Polycomb repressive
complex 1. *Nature* **636**, 755-761 (2024).

37 Hathaway, N. A. *et al.* Dynamics and memory of heterochromatin in living cells. *Cell* **149**,
1447-1460 (2012).

38 Gan, H. *et al.* The Mcm2-Ctf4-Polalpha Axis Facilitates Parental Histone H3-H4 Transfer
to Lagging Strands. *Mol Cell* **72**, 140-151 e143 (2018).
https://doi.org/10.1016/j.molcel.2018.09.001

39 Petryk, N. *et al.* MCM2 promotes symmetric inheritance of modified histones during DNA
replication. *Science* **361**, 1389-1392 (2018). https://doi.org/10.1126/science.aau0294

40 Huang, H. *et al.* A unique binding mode enables MCM2 to chaperone histones H3–H4 at
replication forks. *Nature structural & molecular biology* **22**, 618-626 (2015).

41 Tian, C. *et al.* DNA polymerase delta governs parental histone transfer to DNA replication
lagging strand. *Proc Natl Acad Sci U S A* **121**, e2400610121 (2024).
https://doi.org/10.1073/pnas.2400610121

42 Serra-Cardona, A. *et al.* The PCNA-Pol delta complex couples lagging strand DNA
synthesis to parental histone transfer for epigenetic inheritance. *Sci Adv* **10**, eadn5175
(2024). https://doi.org/10.1126/sciadv.adn5175

43 Charlton, S. J. *et al.* The fork protection complex promotes parental histone recycling and
epigenetic memory. *Cell* **187**, 5029-5047 e5021 (2024).
https://doi.org/10.1016/j.cell.2024.07.017

44 Gambus, A. *et al.* A key role for Ctf4 in coupling the MCM2-7 helicase to DNA polymerase
α within the eukaryotic replisome. *The EMBO journal* **28**, 2992-3004 (2009).

45 Simon, A. C. *et al.* A Ctf4 trimer couples the CMG helicase to DNA polymerase α in the
eukaryotic replisome. *Nature* **510**, 293-297 (2014).

46 Evrin, C., Maman, J. D., Diamante, A., Pellegrini, L. & Labib, K. Histone H2A-H2B
binding by Pol α in the eukaryotic replisome contributes to the maintenance of repressive
chromatin. *The EMBO journal* **37**, e99021 (2018).

47 Katou, Y. *et al.* S-phase checkpoint proteins Tof1 and Mrc1 form a stable replication-
pausing complex. *Nature* **424**, 1078-1083 (2003).

48 Toda, T. *et al.* Mrc1 regulates parental histone segregation and heterochromatin inheritance.
Molecular Cell **84**, 3223-3236. e3224 (2024).

Li, Z. *et al.* Asymmetric distribution of parental H3K9me3 in S phase silences L1 elements. *Nature* **623**, 643-651 (2023). https://doi.org/10.1038/s41586-023-06711-3

Kollenstart, L. *et al.* Disabling leading and lagging strand histone transmission results in parental histones loss and reduced cell plasticity and viability. *Science Advances* **11**, eadr1453 (2025). https://doi.org/doi:10.1126/sciadv.adr1453

Saxena, S. & Zou, L. Hallmarks of DNA replication stress. *Mol Cell* **82**, 2298-2314 (2022). https://doi.org/10.1016/j.molcel.2022.05.004

Lopes, M., Foiani, M. & Sogo, J. M. Multiple mechanisms control chromosome integrity after replication fork uncoupling and restart at irreparable UV lesions. *Molecular cell* **21**, 15-27 (2006).

Zellweger, R. *et al.* Rad51-mediated replication fork reversal is a global response to genotoxic treatments in human cells. *J Cell Biol* **208**, 563-579 (2015). https://doi.org/10.1083/jcb.201406099

Liu, W. *et al.* RAD51 bypasses the CMG helicase to promote replication fork reversal. *Science* **380**, 382-387 (2023). https://doi.org/doi:10.1126/science.add7328

Poole, L. A. & Cortez, D. Functions of SMARCAL1, ZRANB3, and HLTF in maintaining genome stability. *Critical reviews in biochemistry and molecular biology* **52**, 696-714 (2017).

Adolph, M. B. *et al.* Strand dependent bypass of DNA lesions during fork reversal by ATP-dependent translocases SMARCAL1, ZRANB3, and HLTF. *bioRxiv*, 2024.2009. 2017.613558 (2024).

Couch, F. B. *et al.* ATR phosphorylates SMARCAL1 to prevent replication fork collapse. *Genes & development* **27**, 1610-1623 (2013).

Bétous, R. *et al.* SMARCAL1 catalyzes fork regression and Holliday junction migration to maintain genome stability during DNA replication. *Genes & development* **26**, 151-162 (2012).

Ciccia, A. *et al.* Polyubiquitinated PCNA recruits the ZRANB3 translocase to maintain genomic integrity after replication stress. *Molecular cell* **47**, 396-409 (2012).

Vujanovic, M. *et al.* Replication fork slowing and reversal upon DNA damage require PCNA polyubiquitination and ZRANB3 DNA translocase activity. *Molecular cell* **67**, 882-890. e885 (2017).

Kile, A. C. *et al.* HLTF's ancient HIRAN domain binds 3′ DNA ends to drive replication fork reversal. *Molecular cell* **58**, 1090-1100 (2015).

Achar, Y. J., Balogh, D. & Haracska, L. Coordinated protein and DNA remodeling by human HLTF on stalled replication fork. *Proceedings of the National Academy of Sciences* **108**, 14073-14078 (2011).

Blastyák, A. *et al.* Yeast Rad5 protein required for postreplication repair has a DNA helicase activity specific for replication fork regression. *Molecular cell* **28**, 167-175 (2007).

Blastyák, A., Hajdú, I., Unk, I. & Haracska, L. Role of double-stranded DNA translocase activity of human HLTF in replication of damaged DNA. *Molecular and cellular biology* **30**, 684-693 (2010).

Bai, G. *et al.* HLTF promotes fork reversal, limiting replication stress resistance and preventing multiple mechanisms of unrestrained DNA synthesis. *Molecular cell* **78**, 1237-1251. e1237 (2020).

Quinet, A. *et al.* PRIMPOL-mediated adaptive response suppresses replication fork reversal in BRCA-deficient cells. *Molecular cell* **77**, 461-474. e469 (2020).

Taglialatela, A. *et al.* REV1-Polζ maintains the viability of homologous recombination-deficient cancer cells through mutagenic repair of PRIMPOL-dependent ssDNA gaps. *Mol Cell* **81**, 4008-4025.e4007 (2021). https://doi.org/10.1016/j.molcel.2021.08.016

Cong, K. *et al.* Replication gaps are a key determinant of PARP inhibitor synthetic lethality with BRCA deficiency. *Mol Cell* **81**, 3128-3144.e3127 (2021). https://doi.org/10.1016/j.molcel.2021.06.011

Cong, K. & Cantor, S. B. Exploiting replication gaps for cancer therapy. *Mol Cell* **82**, 2363-2369 (2022). https://doi.org/10.1016/j.molcel.2022.04.023

Dungrawala, H. *et al.* The Replication Checkpoint Prevents Two Types of Fork Collapse without Regulating Replisome Stability. *Mol Cell* **59**, 998-1010 (2015). https://doi.org/10.1016/j.molcel.2015.07.030

Roy, S. & Schlacher, K. SIRF: a single-cell assay for in situ protein interaction with nascent DNA replication forks. *Bio-protocol* **9**, e3377-e3377 (2019).

Petryk, N. *et al.* Genome-wide and sister chromatid-resolved profiling of protein occupancy in replicated chromatin with ChOR-seq and SCAR-seq. *Nature Protocols* **16**, 4446-4493 (2021).

Sun, Z. *et al.* Transcription-associated histone pruning demarcates macroH2A chromatin domains. *Nature Structural & Molecular Biology* **25**, 958-970 (2018). https://doi.org/10.1038/s41594-018-0134-5

Tagami, H., Ray-Gallet, D., Almouzni, G. & Nakatani, Y. Histone H3. 1 and H3. 3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell* **116**, 51-61 (2004).

Jansen , L. E. T., Black , B. E., Foltz , D. R. & Cleveland , D. W. Propagation of centromeric chromatin requires exit from mitosis. *Journal of Cell Biology* **176**, 795-805 (2007). https://doi.org/10.1083/jcb.200701066

Rouillon, C. *et al.* CAF-1 deposits newly synthesized histones during DNA replication using distinct mechanisms on the leading and lagging strands. *Nucleic Acids Research* **51**, 3770-3792 (2023). https://doi.org/10.1093/nar/gkad171

Li, Z. *et al.* DNA polymerase α interacts with H3-H4 and facilitates the transfer of parental histones to lagging strands. *Science advances* **6**, eabb5820 (2020).

945 946 947	78	Wenger, A. *et al.* Symmetric inheritance of parental histones governs epigenome maintenance and embryonic stem cell identity. *Nature Genetics* **55**, 1567-1578 (2023). https://doi.org/10.1038/s41588-023-01476-x
948 949	79	Garcia, B. A. *et al.* Chemical derivatization of histones for facilitated analysis by mass spectrometry. *Nat Protoc* **2**, 933-938 (2007). https://doi.org/10.1038/nprot.2007.106
950 951 952	80	Tirman, S. *et al.* Temporally distinct post-replicative repair mechanisms fill PRIMPOL-dependent ssDNA gaps in human cells. *Mol Cell* **81**, 4026-4040.e4028 (2021). https://doi.org/10.1016/j.molcel.2021.09.013
953 954 955	81	Schlacher, K., Wu, H. & Jasin, M. A Distinct Replication Fork Protection Pathway Connects Fanconi Anemia Tumor Suppressors to RAD51-BRCA1/2. *Cancer Cell* **22**, 106-116 (2012). https://doi.org/https://doi.org/10.1016/j.ccr.2012.05.015
956 957 958	82	Schlacher, K. *et al.* Double-Strand Break Repair-Independent Role for BRCA2 in Blocking Stalled Replication Fork Degradation by MRE11. *Cell* **145**, 529-542 (2011). https://doi.org/https://doi.org/10.1016/j.cell.2011.03.041
959 960 961	83	Kang, Z. *et al.* BRCA2 associates with MCM10 to suppress PRIMPOL-mediated repriming and single-stranded gap formation after DNA damage. *Nature Communications* **12**, 5966 (2021). https://doi.org/10.1038/s41467-021-26227-6
962 963 964	84	Taglialatela, A. *et al.* Restoration of Replication Fork Stability in BRCA1- and BRCA2-Deficient Cells by Inactivation of SNF2-Family Fork Remodelers. *Molecular Cell* **68**, 414-430.e418 (2017). https://doi.org/https://doi.org/10.1016/j.molcel.2017.09.036
965 966 967	85	Thakar, T. *et al.* Lagging strand gap suppression connects BRCA-mediated fork protection to nucleosome assembly through PCNA-dependent CAF-1 recycling. *Nature Communications* **13**, 5323 (2022). https://doi.org/10.1038/s41467-022-33028-y
968 969 970	86	Hashimoto, Y., Ray Chaudhuri, A., Lopes, M. & Costanzo, V. Rad51 protects nascent DNA from Mre11-dependent degradation and promotes continuous DNA synthesis. *Nature Structural & Molecular Biology* **17**, 1305-1311 (2010). https://doi.org/10.1038/nsmb.1927
971 972 973	87	Bantele, S. C. S., Lisby, M. & Pfander, B. Quantitative sensing and signalling of single-stranded DNA during the DNA damage response. *Nat Commun* **10**, 944 (2019). https://doi.org/10.1038/s41467-019-08889-5
974 975 976	88	Chen, C. C. *et al.* Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. *Cell* **134**, 231-243 (2008). https://doi.org/10.1016/j.cell.2008.06.035
977 978	89	Mimitou, E. P., Yamada, S. & Keeney, S. A global view of meiotic double-strand break end resection. *Science* **355**, 40-45 (2017). https://doi.org/10.1126/science.aak9704
979 980 981	90	Tsukuda, T., Fleming, A. B., Nickoloff, J. A. & Osley, M. A. Chromatin remodelling at a DNA double-strand break site in Saccharomyces cerevisiae. *Nature* **438**, 379-383 (2005). https://doi.org/10.1038/nature04148

982 983 984	91	Tsukuda, T. *et al.* INO80-dependent chromatin remodeling regulates early and late stages of mitotic homologous recombination. *DNA Repair (Amst)* **8**, 360-369 (2009). https://doi.org/10.1016/j.dnarep.2008.11.014
985 986 987	92	van Attikum, H., Fritsch, O. & Gasser, S. M. Distinct roles for SWR1 and INO80 chromatin remodeling complexes at chromosomal double-strand breaks. *Embo j* **26**, 4113-4125 (2007). https://doi.org/10.1038/sj.emboj.7601835
988 989 990 991	93	Peritore, M., Reusswig, K. U., Bantele, S. C. S., Straub, T. & Pfander, B. Strand-specific ChIP-seq at DNA breaks distinguishes ssDNA versus dsDNA binding and refutes single-stranded nucleosomes. *Mol Cell* **81**, 1841-1853.e1844 (2021). https://doi.org/10.1016/j.molcel.2021.02.005
992 993 994	94	Palter, K. B., Foe, V. E. & Alberts, B. M. Evidence for the formation of nucleosome-like histone complexes on single-stranded DNA. *Cell* **18**, 451-467 (1979). https://doi.org/10.1016/0092-8674(79)90064-3
995 996 997	95	Adkins, N. L. *et al.* Nucleosome-like, Single-stranded DNA (ssDNA)-Histone Octamer Complexes and the Implication for DNA Double Strand Break Repair. *J Biol Chem* **292**, 5271-5281 (2017). https://doi.org/10.1074/jbc.M117.776369
998 999 1000	96	Ryan, B. J., Weaver, T. M., Spencer, J. J. & Freudenthal, B. D. Generation of Recombinant Nucleosomes Containing Site-Specific DNA Damage. *Methods Mol Biol* **2701**, 55-76 (2023). https://doi.org/10.1007/978-1-0716-3373-1_4
1001 1002 1003	97	Dyer, P. N. *et al.* Reconstitution of nucleosome core particles from recombinant histones and DNA. *Methods Enzymol* **375**, 23-44 (2004). https://doi.org/10.1016/s0076-6879(03)75002-2
1004 1005 1006	98	Luger, K., Rechsteiner, T. J. & Richmond, T. J. Expression and purification of recombinant histones and nucleosome reconstitution. *Methods Mol Biol* **119**, 1-16 (1999). https://doi.org/10.1385/1-59259-681-9:1
1007 1008 1009	99	Palter, K. B. & Alberts, B. M. The use of DNA-cellulose for analyzing histone-DNA interactions. Discovery of nucleosome-like histone binding to single-stranded DNA. *J Biol Chem* **254**, 11160-11169 (1979).
1010 1011 1012 1013	100	Caffarelli, E., De Santis, P., Leoni, L., Savino, M. & Trotta, E. Interactions of the histone octamer with single-stranded DNA. Sedimentation analysis and low-angle X-ray diffraction. *Biochim Biophys Acta* **739**, 235-243 (1983). https://doi.org/10.1016/0167-4781(83)90034-9
1014 1015	101	Caffarelli, E., Leoni, L. & Savino, M. Folding of single-stranded DNA on the histone octamer. *FEBS Lett* **181**, 69-73 (1985). https://doi.org/10.1016/0014-5793(85)81115-7
1016 1017 1018	102	Wang, Y. *et al.* Biophysical characterization of the association of histones with single-stranded DNA. *Biochim Biophys Acta Gen Subj* **1861**, 2739-2749 (2017). https://doi.org/10.1016/j.bbagen.2017.07.018
1019 1020 1021	103	Braet, C. *et al.* Mobility and distribution of replication protein A in living cells using fluorescence correlation spectroscopy. *Exp Mol Pathol* **82**, 156-162 (2007). https://doi.org/10.1016/j.yexmp.2006.12.008

Spegg, V. *et al.* Phase separation properties of RPA combine high-affinity ssDNA binding with dynamic condensate functions at telomeres. *Nat Struct Mol Biol* **30**, 451-462 (2023). https://doi.org/10.1038/s41594-023-00932-w

Takai, K. K., Kibe, T., Donigian, J. R., Frescas, D. & de Lange, T. Telomere protection by TPP1/POT1 requires tethering to TIN2. *Mol Cell* **44**, 647-659 (2011). https://doi.org/10.1016/j.molcel.2011.08.043

Chen, J., Le, S., Basu, A., Chazin, W. J. & Yan, J. Mechanochemical regulations of RPA's binding to ssDNA. *Sci Rep* **5**, 9296 (2015). https://doi.org/10.1038/srep09296

Yates, L. A. *et al.* A structural and dynamic model for the assembly of Replication Protein A on single-stranded DNA. *Nat Commun* **9**, 5447 (2018). https://doi.org/10.1038/s41467-018-07883-7

Chen, R. & Wold, M. S. Replication protein A: single-stranded DNA's first responder: dynamic DNA-interactions allow replication protein A to direct single-strand DNA intermediates into different pathways for synthesis or repair. *Bioessays* **36**, 1156-1161 (2014). https://doi.org/10.1002/bies.201400107

Li, M., Sengupta, B., Benkovic, S. J., Lee, T. H. & Hedglin, M. PCNA Monoubiquitination Is Regulated by Diffusion of Rad6/Rad18 Complexes along RPA Filaments. *Biochemistry* (2020). https://doi.org/10.1021/acs.biochem.0c00849

Kim, C., Paulus, B. F. & Wold, M. S. Interactions of human replication protein A with oligonucleotides. *Biochemistry* **33**, 14197-14206 (1994).

Kim, C., Snyder, R. O. & Wold, M. S. Binding properties of replication protein A from human and yeast cells. *Mol Cell Biol* **12**, 3050-3059 (1992).

Kim, C. & Wold, M. S. Recombinant human replication protein A binds to polynucleotides with low cooperativity. *Biochemistry* **34**, 2058-2064 (1995).

Kolpashchikov, D. M. *et al.* Polarity of human replication protein A binding to DNA. *Nucleic Acids Res* **29**, 373-379 (2001). https://doi.org/10.1093/nar/29.2.373

Pestryakov, P. E., Khlimankov, D. Y., Bochkareva, E., Bochkarev, A. & Lavrik, O. I. Human replication protein A (RPA) binds a primer-template junction in the absence of its major ssDNA-binding domains. *Nucleic Acids Res* **32**, 1894-1903 (2004). https://doi.org/10.1093/nar/gkh346

Pestryakov, P. E. *et al.* Human replication protein A. The C-terminal RPA70 and the central RPA32 domains are involved in the interactions with the 3'-end of a primer-template DNA. *J Biol Chem* **278**, 17515-17524 (2003). https://doi.org/10.1074/jbc.M301265200

Kolpashchikov, D. M. *et al.* Interaction of the p70 subunit of RPA with a DNA template directs p32 to the 3'-end of nascent DNA. *FEBS Lett* **450**, 131-134 (1999). https://doi.org/10.1016/s0014-5793(99)00484-6

Kolinjivadi, A. M. *et al.* Smarcal1-mediated fork reversal triggers Mre11-dependent degradation of nascent DNA in the absence of Brca2 and stable Rad51 nucleofilaments. *Molecular cell* **67**, 867-881. e867 (2017).

118 Ray Chaudhuri, A. & Nussenzweig, A. The multifaceted roles of PARP1 in DNA repair
and chromatin remodelling. *Nat Rev Mol Cell Biol* **18**, 610-621 (2017).
https://doi.org/10.1038/nrm.2017.53

119 Bacic, L. *et al.* Structure and dynamics of the chromatin remodeler ALC1 bound to a
PARylated nucleosome. *Elife* **10** (2021). https://doi.org/10.7554/eLife.71420

120 Lehmann, L. C. *et al.* Mechanistic Insights into Autoinhibition of the Oncogenic Chromatin
Remodeler ALC1. *Mol Cell* **68**, 847-859.e847 (2017).
https://doi.org/10.1016/j.molcel.2017.10.017

121 Yang, G. *et al.* Poly(ADP-ribosyl)ation mediates early phase histone eviction at DNA
lesions. *Nucleic Acids Res* **48**, 3001-3013 (2020). https://doi.org/10.1093/nar/gkaa022

122 Nosella, M. L. *et al.* Poly(ADP-ribosyl)ation enhances nucleosome dynamics and
organizes DNA damage repair components within biomolecular condensates. *Mol Cell* **84**,
429-446.e417 (2024). https://doi.org/10.1016/j.molcel.2023.12.019

123 Poirier, G. G., de Murcia, G., Jongstra-Bilen, J., Niedergang, C. & Mandel, P. Poly(ADP-
ribosyl)ation of polynucleosomes causes relaxation of chromatin structure. *Proc Natl Acad
Sci U S A* **79**, 3423-3427 (1982). https://doi.org/10.1073/pnas.79.11.3423

124 Strickfaden, H. *et al.* Poly(ADP-ribosyl)ation-dependent Transient Chromatin
Decondensation and Histone Displacement following Laser Microirradiation. *J Biol Chem*
291, 1789-1802 (2016). https://doi.org/10.1074/jbc.M115.694992

125 Pecori, R., Di Giorgio, S., Paulo Lorenzo, J. & Nina Papavasiliou, F. Functions and
consequences of AID/APOBEC-mediated DNA and RNA deamination. *Nature Reviews
Genetics* **23**, 505-518 (2022). https://doi.org/10.1038/s41576-022-00459-8

126 Ruiz, P. D. & Smogorzewska, A. Temporary HMCES-DNA cross-link prevents permanent
DNA damage. *Cell Rep* **43**, 113594 (2024). https://doi.org/10.1016/j.celrep.2023.113594

127 Mohni, K. N. *et al.* HMCES Maintains Genome Integrity by Shielding Abasic Sites in
Single-Strand DNA. *Cell* **176**, 144-153.e113 (2019).
https://doi.org/10.1016/j.cell.2018.10.055

128 Thompson, P. S., Amidon, K. M., Mohni, K. N., Cortez, D. & Eichman, B. F. Protection of
abasic sites during DNA replication by a stable thiazolidine protein-DNA cross-link. *Nat
Struct Mol Biol* **26**, 613-618 (2019). https://doi.org/10.1038/s41594-019-0255-5

129 Rua-Fernandez, J. *et al.* Self-reversal facilitates the resolution of HMCES DNA-protein
crosslinks in cells. *Cell Rep* **42**, 113427 (2023).
https://doi.org/10.1016/j.celrep.2023.113427

130 Mehta, K. P. M., Lovejoy, C. A., Zhao, R., Heintzman, D. R. & Cortez, D. HMCES
Maintains Replication Fork Progression and Prevents Double-Strand Breaks in Response
to APOBEC Deamination and Abasic Site Formation. *Cell Rep* **31**, 107705 (2020).
https://doi.org/10.1016/j.celrep.2020.107705

131 Thompson, P. S. & Cortez, D. New insights into abasic site repair and tolerance. *DNA
Repair (Amst)* **90**, 102866 (2020). https://doi.org/10.1016/j.dnarep.2020.102866

132   Srivastava, M. *et al.* HMCES safeguards replication from oxidative stress and ensures
error-free repair. *EMBO Rep* **21**, e49123 (2020). https://doi.org/10.15252/embr.201949123

133   Prasad, R. *et al.* Suicidal cross-linking of PARP-1 to AP site intermediates in cells
undergoing base excision repair. *Nucleic acids research* **42**, 6337-6351 (2014).

134   Li, G., Levitus, M., Bustamante, C. & Widom, J. Rapid spontaneous accessibility of
nucleosomal DNA. *Nat Struct Mol Biol* **12**, 46-53 (2005). https://doi.org/10.1038/nsmb869

135   Gaggioli, V. *et al.* Dynamic de novo heterochromatin assembly and disassembly at
replication forks ensures fork stability. *Nature Cell Biology* **25**, 1017-1032 (2023).
https://doi.org/10.1038/s41556-023-01167-z

136   Jasencakova, Z. *et al.* Replication Stress Interferes with Histone Recycling and
Predeposition Marking of New Histones. *Molecular Cell* **37**, 736-743 (2010).
https://doi.org/10.1016/j.molcel.2010.01.033

137   Padeken, J., Methot, S. P. & Gasser, S. M. Establishment of H3K9-methylated
heterochromatin and its functions in tissue differentiation and maintenance. *Nat Rev Mol*
Cell Biol **23**, 623-640 (2022). https://doi.org/10.1038/s41580-022-00483-w

138   Leuzzi, G. *et al.* SMARCAL1 is a dual regulator of innate immune signaling and PD-L1
expression that promotes tumor immune evasion. *Cell* **187**, 861-881.e832 (2024).
https://doi.org/10.1016/j.cell.2024.01.008

139   Sirbu, B. M., Couch, F. B. & Cortez, D. Monitoring the spatiotemporal dynamics of
proteins at replication forks and in assembled chromatin using isolation of proteins on
nascent DNA. *Nature protocols* **7**, 594-605 (2012).

140   Seppa, I. M. *et al.* MRN–CtIP, EXO1, and DNA2–WRN/BLM act bidirectionally to
process DNA gaps in PARPi-treated cells without strand cleavage. *Genes & Development*
39, 582-602 (2025).

141   Hewitt, G. *et al.* Defective ALC1 nucleosome remodeling confers PARPi sensitization and
synthetic lethality with HRD. *Molecular Cell* **81**, 767-783.e711 (2021).
https://doi.org/10.1016/j.molcel.2020.12.006

142   Verma, P. *et al.* ALC1 links chromatin accessibility to PARP inhibitor response in
homologous recombination-deficient cells. *Nature Cell Biology* **23**, 160-171 (2021).
https://doi.org/10.1038/s41556-020-00624-3

143   Bacic, L. *et al.* Asymmetric nucleosome PARylation at DNA breaks mediates directional
nucleosome sliding by ALC1. *Nature Communications* **15**, 1000 (2024).
https://doi.org/10.1038/s41467-024-45237-8

144   Semlow, D. R., MacKrell, V. A. & Walter, J. C. The HMCES DNA-protein cross-link
functions as an intermediate in DNA interstrand cross-link repair. *Nat Struct Mol Biol* **29**,
451-462 (2022). https://doi.org/10.1038/s41594-022-00764-0

145   Murai, J. *et al.* Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res*
72, 5588-5599 (2012). https://doi.org/10.1158/0008-5472.Can-12-2753

146   Satoh, M. S. & Lindahl, T. Role of poly(ADP-ribose) formation in DNA repair. *Nature* **356**,
356-358 (1992). https://doi.org/10.1038/356356a0

147   Moser, S. C. *et al.* NASP modulates histone turnover to drive PARP inhibitor resistance.
Nature (2025). https://doi.org/10.1038/s41586-025-09414-z

148   Liu, W., Krishnamoorthy, A., Zhao, R. & Cortez, D. Two replication fork remodeling
pathways generate nuclease substrates for distinct fork protection factors. *Science*
Advances **6**, eabc3598 (2020). https://doi.org/doi:10.1126/sciadv.abc3598

149   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome biology* **10**, R25 (2009).

150   Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible
platform for exploring deep-sequencing data. *Nucleic acids research* **42**, W187-W191
(2014).

151   Orlando, D. A. *et al.* Quantitative ChIP-Seq normalization reveals global modulation of the
epigenome. *Cell reports* **9**, 1163-1170 (2014).

152   Robinson, J. T. *et al.* Integrative genomics viewer. *Nature biotechnology* **29**, 24-26 (2011).

153   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, R137
(2008).

154   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841-842 (2010).

155   Donsbach, M. *et al.* A non-proteolytic release mechanism for HMCES-DNA-protein
crosslinks. *The EMBO Journal* **42**, e113360 (2023).

156   Henricksen, L. A., Umbricht, C. B. & Wold, M. S. Recombinant replication protein A:
expression, complex formation, and functional characterization. *J Biol Chem* **269**, 11121-
11132 (1994).

157   Das, S. K. *et al.* Methods to investigate nucleosome structure and dynamics with single-
molecule FRET. *Methods* **215**, 17-27 (2023). https://doi.org/10.1016/j.ymeth.2023.05.003

Acknowledgement:

We would like to thank David Cortez Lab for the SMARCAL1, HLTF, ZRANB3 single and triple knockout U2OS cell lines; Alessandro Vindigni lab for the pcDNA3.1/nV5-DEST PrimPol WT and Mutants vectors; Weixing Zhao lab for the MDA-MB-436 and BRCA1 complemental cell lines, DLD-1 WT and BRCA2 knockout cell lines. Part of this work was supported by funding to M. H. from the National Institutes of Health to M.H. (R35 GM147238-03).

Author contribution:

W.L. conceptualized the project. Q.W. and C.Z. performed PLA, IF, iPOND, western blot assays. C.Z. performed data analysis for genomic and proteomic experiments. Y.D. and T.Z. performed Mass spectrometry experiments. N.I.M. and T.M.W. performed in vitro nucleosome assembly assays. M.G. and S.T. purified HMCES proteins and performed RADAR assay. T.H.L. and M.H.

1175 performed the FRET assay. W.L., Q.W., T.M.W., and M.H. wrote the manuscript, T.M.W and TP.Z.
1176 proofread the manuscript with contributions from all the authors.

1177 **Competing Interests statement**

1178 Authors declare no financial or other relationships that may lead to a conflict of interest in this
1179 study.

1180

Figure Legends

Figure 1. Fork reversal deficiency reduces nucleosome density on nascent DNA

A) iPOND-SILAC-MS analysis of proteins bound to nascent DNA in HU-treated wild-type (WT)
and fork reversal enzyme knockout (FRKO) U2OS cells. WT and FRKO cells were labeled with
light and heavy isotopes for ≥4 weeks, pulse-labeled with EdU and treated with 1mM HU for 4
h prior to iPOND. N = 4 biological replicates; error bars represent SD.

B–D) PLA for EdU and H3 in WT or FRKO cells labeled with 10 μM EdU and treated with 1 mM
HU. B) Representative images (Texas Red represents PLA signal; green represents EdU signal).
C) PLA foci number per cell; each dot represents PLA foci number from a single cell. D) PLA
signal normalized to EdU intensity. ≥1,000 cells/sample were analyzed with MetaXpress software.
N ≥ 10. p-values from Kruskal–Wallis test.

E) Heatmaps of H3 ChOR-seq signal in WT and FRKO cells treated with HU, normalized to spike-
in chromatin from D. melanogaster S2 cells. N = 2.

F) Genome browser views of H3 ChOR-seq from the same experiment in (E), normalized to reads
per million using exogenous spike-in. N = 2.

G) PLA for EdU and H3 in WT or FRKO cells with siRAD51 knockdown. N = 3. p-values from
Kruskal–Wallis test.

1198

Figure 2 Fork reversal deficiency leads to parental histone loss on nascent DNA

A) Schematic of iPOND–propionylation-MS workflow to detect histone modifications on nascent
DNA in WT and FRKO U2OS cells treated with EdU together with 1mM HU for 4 h.

B) $\log_2$ fold-change of histone modifications in FRKO vs. WT cells. Modification-specific peptide
counts were normalized to unique peptide numbers, and then FRKO/WT ratios plotted on $\log_2$
scale. N = 3.

C) PLA for EdU and H3K9me3 in WT and FRKO cells treated as in (A). PLA signals normalized
to EdU intensity. ⩾1,000 cells/sample. N = 3. p-values from two-tailed unpaired t-test.

D–E) Immunofluorescence for H3K9me3 in S-phase WT and FRKO cells. D) Cells pulse-labeled
with EdU (10 min). E) Cells labeled with EdU and treated with HU (4 h). Quantification by QIBC.
N = 3. p-values from two-tailed unpaired t-test.

F) PLA for EdU and H3K9me2 in WT and FRKO cells treated as in (A). PLA signals normalized
to EdU intensity. ⩾1,000 cells/sample. N = 3. p-values from two-tailed unpaired t-test.

1212

1213

1214

Figure 3 PrimPol deletion rescues low nucleosome density

A) iPOND-SILAC-MS analysis of histone chaperones on nascent DNA (data from Figure 1A). N
= 4, error bars generated by Standard deviation.

B-C) S1 nuclease-treated DNA fiber assay to detect ssDNA gaps. N = 3. p-values from Kruskal–
Wallis test.

D-F) PLA for EdU and histone H3 in indicated conditions. PLA signals normalized to EdU
intensity; ⩾1,000 cells/sample. N ⩾ 3. p-values from Kruskal–Wallis test.

G) Western blot for PrimPol in KO and complemented cells.

H-I) PLA for EdU and histone modifications in indicated conditions. H) H3K9me2; I) H3K9me3.
PLA signals normalized to EdU intensity; ⩾1,000 cells/sample. N ⩾ 3. p-values from Kruskal–
Wallis test.

J) Model schematic.

1227

Figure 4. Single-stranded DNA gaps reduce nucleosome density on nascent DNA in cells

A–G) PLA for EdU and H3 in the indicated cell lines and treatments. PLA signals normalized to
EdU intensity; $\geqslant$1,000 cells/sample. N $\geqslant$ 3. p-values from two-tailed unpaired t-test for A-E),
Kruskal-Wallis test for F-G). UWB1.289 and MDA-MB-436 are BRCA1 deficient cells, WT
BRCA1 was complemented in these cell lines. PEO1 is BRCA2 truncate mutation cell line, PEO4
is reversed back that express WT BRCA2. B02: RAD51 inhibitor, 25μM was used in this assay.
Mirin: MRE11 inhibitor, 10μM used in this assay.

Figure 5 Nucleosomes can form on ssDNA gaps in vitro

A) Schematic of nucleosome formation procedure.

B) Native PAGE gel image for nucleosome formation on dsDNA substrate.

C) Nucleosome formation assay for dyad gap substrates. Gap size is 1-31bp as indicated.

D) Nucleosome formation assay for entry/exit gap substrates. Gap size is 1-31bp as indicated.

E) Schematic of the procedure to assemble a nucleosome on the 147nt ssDNA gap substrate
(ssDNA Gap). Wrapping of the poly(dT)147 ssDNA around the histone octamer juxtaposes the
two duplex regions and, hence, the cyanine labels, yielding a FRET signal.

F) Fluorescence emission spectra obtained by excitation at 514 nm. Each spectra is the average of
at least two independent spectra with the SEM shown in grey. The fluorescence emission
intensities at 665 nm (Cy5 FRET acceptor fluorescence emission maximum, I_{665}) and 561 nm (Cy3
FRET donor fluorescence emission maximum, I_{561}) are indicated. Cy5 can be excited through
FRET from Cy3 only when the two dyes are in close proximity of each other (less than ~10 nm).
This is indicated by an increase in I_{665} and a concomitant decrease in I_{561}.

G) Steady state FRET. The approximate FRET efficiency (E_{FRET}) for the ssDNA Gap DNA
substrate alone (ssDNA Gap) or pre-assembled in a nucleosome (ssDNA Gap Nuc) was measured
at equilibrium. The height of each column (E_{FRET}) is the average of at least 4 independent
measurements with the SEM shown. FRET is only observed when a nucleosome is pre-assembled
on the ssDNA Gap DNA substrate.

1254

1255

1256

1257

1258

Figure 6. PARylation reduces nucleosome density in fork reversal-deficient cells

A) Immunofluorescence of PARylation in WT and FRKO cells. Cells were labeled with EdU for
10 min in no HU samples, or EdU with HU for 4 h. Integrated PARylation intensity quantified. N
= 3. p-values from Kruskal-Wallis test.

B) PLA for EdU and H3 under indicated conditions. PLA normalized to EdU intensity; $\geqslant 1,000$
cells/sample. N = 9. p-values from Kruskal-Wallis test.

C) Immunofluorescence of PARylation in indicated conditions. N = 3.

D) S1 nuclease DNA fiber assay to detect ssDNA gaps. N = 3. p-values from Kruskal-Wallis test.

E) PLA for EdU and H3 in indicated conditions, quantified as in (B). N $\geqslant$ 7.

F-G) Immunofluorescence for PARylation in the indicated conditions. N = 3.

H) PLA for EdU and H3 in indicated conditions, quantified as in (B). N $\geqslant$ 7.

1270

1271

Figure 7. HMCES-DPC at ssDNA gaps reduces nucleosome density in fork reversal-deficient cells

A) APOBEC3B and HMCES levels on nascent DNA measured by iPOND-SILAC-MS. N = 4.

B) RADAR assay detecting HMCES-DPC in WT and FRKO cells. N = 3.

C–E) PLA for EdU and H3 in indicated conditions (as in Figure 1B). N ⩾ 7. p-values from
Kruskal–Wallis test.

F–G) Nucleosome reconstitution assays showing nucleosomes fail to form on dyad ssDNA gaps
but can form at side gaps. Gels show NCP bands. AP-DNA represents abasic-site DNA; AP-NCP
represents nucleosomes assembled on abasic DNA. L represents ladder, D is DNA, N is
nucleosome.

H) Model schematic of how ssDNA adducts interfere with nucleosome formation at gaps.

Supplementary Figure Legends

Figure S1. Fork reversal deficiency reduces nucleosome density on nascent DNA (related to Figure 1).

FRKO clone #29 is used in figure S1A-J.

A) EdU incorporation in WT and FRKO cells treated with EdU + HU (as in Fig. 1B). EdU detected by azide-488 click reaction.

B–E) Immunofluorescence for H3 in WT and FRKO cells under indicated conditions. N = 3; p-values from two-tailed unpaired t-test.

F–H) PLA for EdU and H2A (G), H2B (H), and H4 (I). Left: PLA foci/cell; middle: EdU intensity; right: normalized PLA/EdU. ≥1,000 cells/sample. N ≥ 3.

I) Genome browser views of H3 ChOR-seq from the same experiment in Figure 1E, normalized to reads per million using exogenous spike-in. N = 2.

J) Quantitative ChOR-seq for H3. Average profiles across H3 peaks in WT and FRKO cells, normalized by reference-adjusted RPM using D. melanogaster spike-in. N = 2; Wilcoxon signed-rank test.

Figure S2. Low nucleosome density phenotype in individual fork reversal enzymes knockout
clones of U2OS cell lines (related to Figure 1).

A–C) PLA for EdU and H3 in individual fork reversal enzyme KO clones. N = 3. p-values from
unpaired t-test in B) and Kruskal–Wallis test in A) and C).

D) Western blot showing siRAD51 knockdown efficiency.

E) PLA for EdU and H3 in WT and FRKO cells without HU. Cells labeled with EdU (10 min) and
harvested for PLA. N = 3; p-values from unpaired t-test.

F) ChOR-seq analysis of H3 occupancy in WT and FRKO cells without HU, normalized to spike-
in chromatin. The cells are treated same as in D).

G) Heatmaps of H3 ChOR-seq signal (same data from F) in unperturbed condition. N = 2.

G) Model schematic: fork reversal prevents low nucleosome density under replication stress.

1309

1310

Figure S3. Fork reversal deficiency leads to parental histone loss (related to Figure 2).

A-H) Histone modifications quantified in WT and FRKO cells by iPOND–Propionylation-MS.
Modification level equals modified peptide count/unique peptide count. N = 3; error bars is
generated from standard deviation.

I-J) Immunofluorescence for H3K9me2 in indicated conditions. N = 3; unpaired t-test.

Figure S4. PrimPol depletion rescues parental histone loss and new histone increase, related to
Figure 3.

A-D) Histone modifications quantified in WT, FRKO, and FR/PPKO cells by iPOND–
Propionylation-MS. Modification level equals modified peptide count/unique peptide count. N =
3; error bars is generated from standard deviation.

1321

1322 Figure S5. ssDNA gaps loosen the nucleosome assembly on nascent DNA in cells (related to Figure
1323 4).

1324 A–G) PLA for EdU and H3 in indicated treatments. PLA foci number for each cell and EdU
1325 intensity as shown (from the same data as Fig. 4). N = 3; p-values from unpaired t-test.

1326

Figure S6. RPA evicts nucleosome from ssDNA in vitro but doesn't not rescue the low nucleosome
density in cells

A) Model schematic.

B) FRET signal for ssDNA gap and RPA bound ssDNA gap. Saturating the ssDNA within the
ssDNA Gap DNA substrate with RPA does not significantly affect the E_{FRET} observed at
equilibrium. Fluorescence emission spectra obtained by excitation at 514 nm is shown. Each
spectra is the average of at least two independent spectra with the SEM shown in grey. The average
E_{FRET} values observed at equilibrium for each condition are shown in the inset. Each reported value
is the average of at least 4 independent measurements (±SEM).

C) Schematic representation of the FRET pair and assay to monitor unwrapping of ssDNA from
around histone octamers within ssDNA Gap nucleosomes. In ssDNA Gap nucleosomes conditions
as described in Figure 5E, RPA is added and E_{FRET} values are monitored over time. Trace is the
mean of at least three independent traces.

D-E) Immunofluorescence for RPA2 and phosphorylation on RPA2-S8 in WT and FRKO U2OS
cells. The cells are treated same as in Figure 6A. N = 3.

F) PLA for EdU and H3 in siRPA1 and/or siRPA2 cells (treated as in Fig. 1B). N ⩾ 5; Kruskal–
Wallis test. Western blot showed siRNA knockdown efficiency.

1344

1345

Figure S7. Transcription reduces nucleosome density in both WT and FRKO cells.

A) PLA for EdU and RBP1 in indicated treatments. Left, PLA foci number for each cell and EdU
intensity as shown. Middle, EdU integrity intensity for each cell in the indicated treatment. Right,
normalized PLA signal in each cell in the indicated treatment. TPL concentration is 10μM. N = 3;
p-values from Kruskal-Wallis test.

B) PLA for EdU and H3 in indicated treatments. N = 3; TPL concentration is 10μM. p-values from
Kruskal–Wallis test.

1353

1354

Figure S8. HMCES-DPC at ssDNA gaps reduces nucleosome density in FRKO cells (related to
Figure 7).

A) Western blot for chromatin-bound APOBEC3B in WT and FRKO cells with or without HU
treatment.

B-C) Western blot for HMCES (B) and APOBEC3B (C) to confirm knockdown efficiency in WT
and FRKO cells.

D-E) Structural models of ssDNA-gapped nucleosomes predicted by AlphaFold. D) ssDNA gap at
entry/exit. E) ssDNA gap at dyad.

F-G) HMCES-ssDNA crosslink preparation. Gels show HMCES–DNA crosslink products. AP-
DNA represents DNA that contains abasic sites; L represents ladder.

Figure 1 Fork reversal deficiency reduces the nucleosome density on nascent DNA

Figure 2 Fork reversal deficiency leads to parental histone loss on nascent DNA

Figure 3 PrimPol deletion rescues low nucleosome density

Figure 4 Single-Stranded DNA Gaps Decrease Nucleosome Density on Nascent DNA in Cells

Figure 5 Nucleosomes can form on ssDNA gaps in vitro

Figure 6 PARylation reduces nucleosome density in fork reversal deficient cells

Figure 7 HMCES-DPC at ssDNA gaps reduces nucleosome density in fork reversal deficient cells

Figure S2 Low density of nucleosome is due to fork reversal deficiency rather than individual fork reversal enzyme depletion, related to Figure 1

Figure S3 Fork reversal deficiency leads to parental histone loss, related to Figure 2

Figure S4 PrimPol deletion rescues parental histone loss and new histone increase, related to Figure 3

Figure S5 Single-stranded DNA gaps loosen the nucleosome assembly on nascent DNA in cells, related to Figure 4

Figure S6 RPA evicts nucleosomes from ssDNA in vitro but does not rescue the low nucleosome density in cells, related to Figure 5

Figure S7 Transcription reduces nucleosome density in cells

A

B

1379

1380

Figure S8 HMCES-DPC at ssDNA gaps reduces nucleosome density in FRKO cells, related to Figure 7

1382

1383   Table 1. Oligo sequences in this study.

Oligo	Sequence (5′ – 3′)
ND-NCP	
Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTTAAAACGCGGGGGACAGCGCGTACGTGCGTTTAAGCGGTGCTAGAGCTGTCTA CGACCAATTGAGCGGCCTCGGCACCGGGATTCTCGAT
Gap1x-NCPdyad	
Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTTAAAACGCGGGGGACA
Oligo 3	**/5Phos/**CGCGTACGTGCGTTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGG CCTCGGCACCGGGATTCTCGAT
Gap11x-NCPdyad	
Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTTAAAACGCGGG
Oligo 3	**/5Phos/**ACGTGCGTTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCG GCACCGGGATTCTCGAT
Gap21x-NCPdyad	
Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTTAAAAC
Oligo 3	**/5Phos/**CGTTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGGCACC GGGATTCTCGAT
Gap31x-NCPdyad	

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTT
Oligo 3	**/5Phos/**AAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGGCACCGGGA TTCTCGAT

1384

Gap1x-NCPend

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	TCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCGG TTAAAACGCGGGGGACAGCGCGTACGTGCGTTTAAGCGGTGCTAGAGCTGTCTAC GACCAATTGAGCGGCCTCGGCACCGGGATTCTCGAT

Gap11x-NCPend

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCGGTTAAAACGCG GGGGACAGCGCGTACGTGCGTTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGA GCGGCCTCGGCACCGGGATTCTCGAT

Gap21x-NCPend

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	CGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCGGTTAAAACGCGGGGGACAGCG CGTACGTGCGTTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGG CACCGGGATTCTCGAT

Gap31x-NCPend

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	GACTAGGGAGTAATCCCCTTGGCGGTTAAAACGCGGGGGACAGCGCGTACGTGCG TTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGGCACCGGGATT CTCGAT

dU-Gap31x-NCPdyad

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGC/**dU**/GTCCCCCGCGTTTTAACCGCCAAGGGGATTACTC CCTAGTCTCCAGGCACGTGTCAGATATATACATCCGAT
Oligo 2	ATCGGATGTATATATCTGACACGTGCCTGGAGACTAGGGAGTAATCCCCTTGGCG GTT
Oligo 3	/**5Phos**/AAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGGCACCGGGA TTCTCGAT

dU-Gap31x-NCPend

Oligo 1	ATCGAGAATCCCGGTGCCGAGGCCGCTCAATTGGTCGTAGACAGCTCTAGCACCG CTTAAACGCACGTACGCGCTGTCCCCCGCGTTTTAACCGCCAAGGGGATTACTCCC TAGTCTCCAGGCACG/**dU**/GTCAGATATATACATCCGAT

| Oligo 2 | GACTAGGGAGTAATCCCCTTGGCGGTTAAAACGCGGGGGACAGCGCGTACGTGCG TTTAAGCGGTGCTAGAGCTGTCTACGACCAATTGAGCGGCCTCGGCACCGGGATT CTCGAT |

1385

1386 Oligos for FRET assay

1387 5'-ACTCCTTCCGCACGTAATTTTTGAC-Cy5-3 '

1388 5'-Cy3-CTGGTCTCGCGATGG-3 '

1389 3'-TGAGGAAGGCGTGCATTAAAAACTG-poly (dT)$_{147}$-GACCAGAGCGCTACC-5'

1390

1391 <mark>siRNA sequences</mark>

1392

1393