Evaluating the quality of brainstem ROI registration using structural and diffusion MRI

Yi-An A. Chen*[1,2], Lars Kasper[1,3], Clement T. Chow[3,4], Alexandre Boutet[3,5], Jürgen Germann[3],
Andres M. Lozano[3,4,5], Kâmil Uludağ[3,6,7], Andreea O. Diaconescu[1,2,8], Sriranga Kashyap[3]

[1]Department of Psychology, University of Toronto, Toronto, Ontario, Canada

[2]Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto, Ontario,
Canada

[3]Krembil Brain Institute, University Health Network, Toronto, Canada

[4]Division of Neurosurgery, Toronto Western Hospital, Toronto, Canada

[5]Joint Department of Medical Imaging, University of Toronto, Toronto, Canada

[6]Department of Medical Biophysics, University of Toronto, Toronto, Canada

[7] Physical Sciences, Sunnybrook Research Institute, Toronto, ON, Canada

[8]Department of Psychiatry, University of Toronto, Toronto, Canada

* Correspondence:
Yi-An A. Chen
agnes.chen@mail.utoronto.ca

Keywords: Brainstem, Alignment, Diffusion-weighted imaging (DWI), Functional MRI
reliability, ROI registration

Number of words: 5632

Number of figures: 6

Number of tables: 0

Abstract

Accurate registration of regions of interest (ROIs) from standard atlases to participants' native
spaces is a critical step in fMRI studies, as it directly affects the reliability of sampled BOLD signals.
While T1-weighted (T1w) image-based ROI registration is well validated and widely adopted in
cortical fMRI, its performance degrades in brainstem studies due to the small size, dense packing,
and poor visibility of brainstem nuclei on T1w contrast. We hypothesized that incorporating diffusion
MR images, containing more information about internal brainstem architecture, should improve ROI
registration accuracy. To test this, we developed four registration pipelines that either included or

excluded diffusion-based alignment components and evaluated their performance using data from
n=10 healthy participants. Registration accuracy was assessed using Dice coefficients for the red
nucleus (RN) and mis-registration fractions for the dorsal raphe nucleus (DRN). The results showed
that diffusion-based pipelines, using fractional anisotropy (FA) images, non-diffusion-weighted (b0)
images, and multivariate combination, outperformed the T1w-only baseline. Probabilistic maps
derived from inverse-transformed native ROIs further supported improved sensitivity to inter-
individual anatomical variability in the diffusion-augmented pipelines. In addition, analysis of
gradient magnitude maps from the Jacobian determinants revealed associations between localized
deformation and image modality-specific landmarks. These findings demonstrate the potential of
diffusion-augmented pipelines for improving brainstem ROI registration, which could enhance the
robustness of fMRI studies on brainstem disorders characterized by functional dysregulation.

1   Introduction

The brainstem is a crucial hub of the central nervous system. It serves as a conduit for
numerous ascending and descending axonal tracts, actively linking the brain with the rest of the body
to coordinate physiological functions (Fernández-Gil et al., 2010). As one of the most evolutionarily
conserved structures (Pessoa et al., 2019), the brainstem harbors several highly specialized, small
neuronal clusters, known as brainstem nuclei, that support autonomic and neuromodulatory functions
essential to vital bodily processes. Despite its importance, the brainstem has been relatively
underexplored in functional MRI (fMRI) research due to challenges arising from a combination of its
anatomical location and constraints from image acquisition. These include increased sensitivity to
motion artifacts from pulsating surface arteries (e.g., basilar and vertebral arteries), signal dropouts
and image distortion at air-tissue interfaces (e.g. sphenoid sinus, pharynx), and reduced signal-to-
noise ratio (SNR) resulting from the small size of brainstem nuclei (Beissner, 2015; Sclocco et al.,
2018). Collectively, these factors amplify the impact of co-registration errors between functional and
anatomical data, compromising the validity and reliability of brainstem fMRI. To address these
brainstem-specific limitations and enhance the feasibility of brainstem fMRI as a research tool, the
fMRI community has been actively developing tailored data acquisition and preprocessing strategies
to improve the fidelity of brainstem imaging (Beissner, 2015; Sclocco et al., 2018; Matt et al, 2019;
Cauzzo et al., 2022; Singh et al., 2022; Hansen et al., 2024, Zhang et al., 2025).

In conventional fMRI analysis, cortical and large subcortical (e.g. basal ganglia, thalamus)
ROIs, defined using anatomical data and atlases, are used to extract BOLD time series for statistical
analysis. This process typically follows a two-step approach involving spatial transformation-that is,

a mathematical mapping between coordinate systems-between the participant's "native space,"
defined by participant-specific anatomical data, and a "standard space," defined by population-based
brain templates (e.g., MNI152 or IIT templates) constructed by averaging data from dozens to
hundreds of individuals (Fonov et al. 2009). In the first step, a nonlinear spatial transform from
standard space to native space is calculated by optimizing alignment between a standard T1-weighted
(T1w) image template and the participant's high-resolution T1w image. In the second step, this
transform is applied to map ROIs from an atlas defined in standard space into the participant's native
space. This T1w-based approach generally enables accurate registration of cortical and forebrain
subcortical ROIs due to the excellent gray matter-white matter contrast provided by T1w images. As
this method is considered standard practice and is implemented in the default preprocessing pipelines
of many fMRI analysis toolboxes, it is also widely used in brainstem fMRI studies, except in rare
cases where certain brainstem nuclei can be directly visualized in native space due to their distinct
biochemical properties and where additional acquisition time is acceptable for the studied population
(Chen et al., 2014; Priovoulos et al., 2018; Mazancieux et al., 2023). Unlike the cortex, however, the
brainstem lacks well-defined macroscopic gray matter-white matter boundaries, with nuclei blending
into a dense network of interwoven axonal tracts. Consequently, the brainstem often appears in T1w
images as a relatively homogeneous structure, with clear external boundaries yet unclear internal
architecture. As a result, when applied to brainstem fMRI, the first-step standard-to-native
registration relies primarily on gross anatomical features. This can lead to suboptimal registration of
brainstem ROIs because the spatial locations of brainstem nuclei are governed more by internal
microstructural organization than by gross surface contours and there is known inter-individual
variability of the internal organization (Shepherd et al., 2020; Bianciardi et al., 2018; García-Gomar
et al., 2019; Singh et al., 2020). As such, a registration pipeline factoring in the internal organization
of the brainstem would improve the validity of BOLD signal sampling at the individual level.

In principle, the quality of conventional standard-to-native, T1w-based registration could be
enhanced by incorporating an additional image modality that provides spatial references to the
internal architecture of the brainstem. Diffusion-weighted imaging (DWI) is one such modality that is
well suited for this purpose. Unlike susceptibility-weighted imaging or turbo spin echo imaging,
which are primarily used to visualize iron-rich structures localized to the midbrain or neuromelanin-
laden nuclei distributed in a patchy manner, DWI-derived images can reveal the densely interwoven
axonal tracts that span nearly the entire brainstem. Aside from its well-established role in identifying
anatomical substrates for functional connectivity (van den Heuvel & Sporns, 2013), the
microstructural features of axonal tracts revealed by DWI also form an internal spatial reference

frame that could be used to improve ROI registration accuracy for the brainstem nuclei embedded within it. Typically, DWI enables the derivation of various scalar and vector maps-such as the fractional anisotropy (FA), the non-diffusion-weighted (b=0), the mean diffusivity (MD), and the fiber direction maps, which offer complementary microstructural information relevant for ROI delineation. These DWI-derived scalar maps are inherently co-registered to each other, thereby avoiding registration errors that can arise when aligning images obtained from different MR sequences.

In this study, we evaluated the benefit of incorporating DWI-derived images in addition to the T1w for brainstem nuclei ROI registration. Specifically, standard-to-native transforms were generated through a two-step process that first computed a whole-brain T1w-based transform, as in the conventional approach, and then locally refined the transform within the brainstem by co-registering (i) the T1w image with a T1w template, (ii) the FA image with an FA template, (iii) the b0 image with a b0 template, (iv) all three image-template pairs in (i)-(iii) through an iterative multi-channel approach. These transforms were subsequently used to map brainstem ROIs from standard space into the native spaces of individual participants. The performance of the four types of transforms-T1w-based, FA-based, b0-based, and multivariate-was evaluated by quantifying the spatial overlap between the registered ROIs and manually delineated brainstem ROIs, which served as the ground truth. We hypothesized that incorporating DWI-derived images, by introducing internal structural features of the brainstem, would improve the accuracy of brainstem ROI registration over conventional T1w-based registration.

2   Materials and methods

2.1   Participants

Data were acquired from ten healthy participants (age: 38.6 ± 16.5 years; three females) using a Siemens Magnetom Prisma 3 T scanner (Siemens Healthineers, Erlangen, Germany) at Toronto Western Hospital, equipped with a 20-channel head and neck receive coil. The study was approved by the University Health Network Research Ethics Board (REB 21-6105), and signed informed consent was obtained from all participants.

2.2   Data acquisition

T1-weighted (T1w) images were acquired using a 3D magnetization-prepared rapid acquisition gradient echo (MPRAGE) sequence with the following parameters (Mugler & Brookeman, 1990): matrix size = 320×320; slice thickness = 1 mm; echo time (TE) = 2.6 ms;

repetition time (TR) = 2200 ms; inversion time (TI) = 944 ms; flip angle = 8°; GRAPPA = 2;
bandwidth = 252 Hz/pixel. Diffusion-weighted images (DWI) were acquired using a spin-echo SMS
2D-EPI sequence in two anterior-posterior (AP) runs: one with 64 diffusion directions (b = 2000
s/mm²) and one with 30 directions (b = 1000 s/mm²). Each run included five b0 images. An
additional five b0 images were acquired with posterior-anterior (PA) phase encoding for distortion
and eddy corrections. Acquisition parameters were: GRAPPA=2, SMS=2, resolution = 2 × 2 × 2
mm³; matrix size = 110 × 110; TE = 75 ms; TR = 4100 ms; flip angle = 90°. Susceptibility-weighted
images were acquired using a multi-echo gradient-echo sequence: matrix size = 384 × 384 × 52;
resolution = 0.5 × 0.5 × 1 mm³; TR = 53 ms; four echoes (TE = 7.3, 20, 32, 44 ms); flip angle = 20°;
bandwidth = 100 Hz/pixel.

2.3 Data preprocessing

DWI data were denoised using the MP-PCA algorithm implemented in the `dwidenoise`
tool in MRtrix3. Ten AP b0 images and five PA b0 images were used to estimate susceptibility-
induced off-resonance fields via FSL's `topup` tool (Andersson et al., 2003). These outputs were
passed to the `eddy` tool along with diffusion parameters to correct for eddy current-induced
distortions, magnetic field inhomogeneities, and motion (Andersson et al., 2016) in a single-step. The
corrected DWIs and rotated b-vectors were used with FSL's `dtifit` to compute the diffusion tensor
and scalar maps, including the fractional anisotropy (FA) and averaged b0 images. SWI data were
processed using CLEAR-SWI, which optimizes multi-echo combination of phase and magnitude
images to yield optimal susceptibility-based tissue contrast (Eckstein et al., 2021).

2.4 Image co-registration

The Brainstem Navigator Atlas v0.9 (https://www.nitrc.org/projects/brainstemnavig/) was
selected for this study as it includes both structural and diffusion templates in the IIT space. All three
IIT templates used in the study-namely, IIT-T1w, IIT-FA, and IIT-b0-were brain-extracted using a
whole-brain mask derived from the IIT-T1w template to remove extra-brain structures that could
interfere with registration (e.g., the bright basilar artery in IIT-T1w).
Within-participant alignment of the T1w, FA, and b0 images was performed in several steps.
First, each participant's T1w image was brain-extracted using FreeSurfer's `mri_synthstrip` and
then bias-field corrected using the `N4BiasFieldCorrection` tool in ANTs (Avants et al., 2009).
The T1w, FA, and b0 images were resampled to a common resolution (0.7 × 0.7 × 0.7 mm³) to
prevent resolution mismatches from affecting image alignment. Given that b0 images more closely

resemble T1w images in terms of gross anatomical contrast (albeit inverted), the affine transform

aligning diffusion and structural images was estimated between the b0 and T1w images using ANTs'

`antsRegistration` tool, with mutual information (MI) as the optimization metric. This

transform was subsequently applied to the FA and b0 images to align them with the T1w anatomical

space.

Brainstem ROIs were mapped from IIT space to each participant's native space using

compound transforms derived from a two-step registration pipeline illustrated in Figure 1A. In the

first step, a T1w-based whole-brain standard-to-native transform (T_{T1w-WB}) was estimated using

ANTs' `antsRegistration`, constrained to the participant's whole-brain mask. This registration

included sequential optimization of (1) a four-level rigid-body transformation using MI, (2) a four-

level affine transformation using MI, and (3) a five-level symmetric normalization (SyN)

transformation using cross-correlation (CC) as the metric (Avants et al., 2007). The T_{T1w-WB}

transform was then applied to the IIT-T1w, IIT-FA, and IIT-b0 templates to create three warped

templates in approximate alignment with the participant's native space. Quality control was carried

out using visual inspection for all stages of the registration process.

In the second step, ANTs' `antsRegistration` was used to estimate three additional

nonlinear transforms by registering each warped IIT template (T1w, FA, b0) to its corresponding

native image within the brainstem. A multivariate transform was also computed by jointly optimizing

alignment across all three image types with equal weighting. All second-step registrations were

constrained to a slightly dilated brainstem mask generated using FreeSurfer's `recon-all` to ensure

inclusion of relevant surrounding contours. The registrations were optimized using a CC-based, two-

level SyN algorithm with the following parameters: convergence = 50×20; shrink factors = 2×1;

smoothing sigmas = 2×0. The resulting nonlinear transforms-T_{T1w-BS}, T_{FA-BS}, T_{b0-BS}, and the

multivariate T_{mv-BS}-were each concatenated with T_{T1w-WB} to form compound transforms for mapping

brainstem ROIs from IIT space to each participant's native space (Figure 1B).

2.5   Brainstem ROI delineation

Two sets of brainstem ROIs were selected to evaluate registration accuracy: the red nucleus

(RN) and the dorsal raphe nucleus (DRN). These ROIs were chosen based on the visibility of either

their own anatomical boundaries or those of neighboring structures in the native space, enabling

reliable comparison between registered ROIs and manually delineated ground-truth references. The

RN appears as a pair of clearly identifiable, low-intensity, ellipsoidal structures in the midbrain,

visible in both susceptibility-weighted and T2-weighted b0 images. The well-circumscribed nature of

the RN makes it suitable for Dice coefficient-based assessment of registration accuracy. In contrast,

the DRN is not directly visualizable in any available MRI contrast and is unsuitable for Dice

coefficient-based evaluation. However, the DRN lies directly ventral to and parallel with the cerebral

aqueduct and the fourth ventricle, both of which are clearly delineated in T1w images. Accordingly,

we developed an alternative metric-the mis-registration fraction-to assess DRN registration accuracy.

The metric quantifies the proportion of registered DRN ROI voxels that erroneously overlap with the

voxels of the cerebral aqueduct and the fourth ventricle.

Ground-truth RN ROIs were manually delineated in ITK-SNAP based on susceptibility-

weighted and b0 images that had been co-registered to the T1w images. Ground-truth ROIs for the

cerebral aqueduct and fourth ventricle (CA-4thVen) were generated using FreeSurfer's `recon-all`

tool, with manual editing applied as needed to correct missing or incomplete segmentation of the

narrow cerebral aqueduct.

2.6   Data analysis

The quality of brainstem ROI registration achieved by the four compound transforms (T1w-

based, FA-based, b0-based, and multivariate) was evaluated using two metrics. First, the Dice

coefficient between each registered RN ROI and the corresponding ground-truth ROI was calculated

as:

$$DiceCoefficient = \frac{2 \times (RN_{GT} \cap RN_R)}{RN_{GT} + RN_R}, \quad (1)$$

where RN_{GT} denotes the ground-truth RN ROI and RN_R denotes the registered RN ROI.

Second, registration error for DRN was evaluated using the mis-registration fraction, defined

as:

$$Mis\text{-}registrationFraction = \frac{CA\text{-}4_{th}Ven_{GT} \cap DRN_R}{DRN_R}, \quad (2)$$

where DRN_R is the registered DRN ROI and $CA\text{-}4_{th}Ven_{GT}$ is the ground-truth $CA\text{-}4_{th}Ven$ ROI. For

each participant, four Dice coefficients and four mis-registration fractions were determined,

reflecting registration performance under the four different registration strategies. All analyses were

conducted using Python 3 with `nibabel` for image processing.

2.7   Data visualization

To qualitatively compare the registration approaches, two types of visualizations were

generated. First, four sets of probabilistic RN and DRN maps were created in IIT space by inverse-

transforming the native-space ROIs obtained via the two-step compound transforms, using each

participant's inverse $T_{T1w\text{-}WB}$ transform. These inverse-transformed ROIs were averaged across
participants to generate probabilistic maps, which were compared with the original Brainstem
Navigator atlas. It was hypothesized that diffusion-based second-step transforms (i.e., $T_{FA\text{-}BS}$, $T_{b0\text{-}BS}$, and
the multivariate $T_{mv\text{-}BS}$), by incorporating internal brainstem architecture, would result in greater spatial
dispersion of the probabilistic maps due to the known inter-individual variability in brainstem ROI
locations, as reported in previous studies with a similarly sized sample (Bianciardi et al., 2018;
García-Gomar et al., 2019; Singh et al., 2020).

Second, Jacobian determinant maps (JDet) were computed for each second-step transform
using ANTs' `CreateJacobianDeterminantImage` tool. These maps were normalized to the
range [0, 1], and gradient magnitude maps ($|\nabla JDet|$) were derived by calculating the spatial gradients
and taking the magnitude of each voxel's gradient vector. The resulting $|\nabla JDet|$ maps were then
averaged across participants to generate mean $|\nabla JDet|$ maps, which were used to visualize regions of
deformation targeted by the optimization algorithm. Since deformation tends to occur near modality-
specific anatomical landmarks, such as the brainstem surface in T1w images and tract boundaries in
FA images, these regions were expected to show higher gradient magnitudes in the corresponding
$|\nabla JDet|$ maps. The mean $|\nabla JDet|$ map for the multivariate transform was expected to exhibit a more
spatially uniform gradient distribution, reflecting the integration of salient anatomical features from
all three image modalities.

3   Results

Figure 3A shows the Dice coefficients between the ground-truth RN ROIs and the four RN
ROIs obtained via the two-step compound transforms for each of the ten participants. A one-way
repeated-measures ANOVA was conducted to examine the effect of compound transform type on
Dice coefficient values. The analysis revealed a statistically significant difference among the four
groups ($F(3, 27) = 9.34$, $p < 0.001$). Post hoc pairwise t-tests with Bonferroni correction (adjusted $\alpha =$
0.0083, 6 comparisons) indicated that Dice coefficients derived from the T1w-based compound
transforms were significantly lower than those from the other three transform types ($t(9) = -5.32$, $p <$
0.001 for T1w versus b0; $t(9) = -3.76$, $p = 0.004$ for T1w versus FA; $t(9) = -5.21$, $p < 0.001$ for T1w
versus multivariate), suggesting improved registration accuracy when diffusion-based images were
incorporated into the registration process.

Figure 3B shows the mis-registration fractions computed using the ground-truth CA-4_{th}Ven
ROIs and the four DRN ROIs obtained via the compound transforms for each of the ten participants.
A one-way repeated-measures ANOVA was conducted to examine the effect of compound transform

type on mis-registration fractions. The analysis revealed a statistically significant difference among
the four groups ($F(3, 27) = 14.23$, $p < 0.001$). Post hoc pairwise t-tests were performed between all
pairs of transform types. After Bonferroni correction, only the FA-based compound transforms,
which had the lowest average mis-registration fraction, showed statistically significant differences
compared to each of the other three transform types ($t(9) = 7.52$, $p < 0.001$ for T1w versus FA; $t(9) =$
4.15, $p = 0.002$ for b0 versus FA; $t(9) = -5.79$, $p < 0.001$ for FA versus multivariate).

Figure 4 presents probabilistic maps of the DRN (top row) and RN (bottom row), generated
by inverse-transforming the two-step-transformed native-space ROIs of each participant back into IIT
space and averaging across the ten participants. For reference, the unthresholded Brainstem
Navigator probabilistic maps for each nucleus are shown in the leftmost column. In every panel, red
contours mark the thresholded Brainstem Navigator ROIs (threshold = 0.35), which served as the
standard ROIs in IIT space throughout the registration pipelines.

As shown in the Brainstem Navigator maps, only 68.1% of voxels in the DRN probabilistic
map and 51.2% in the RN map fall within the thresholded ROI boundaries. These relatively low
proportions reflect the anatomical variability among individuals whose brainstem data were used to
construct the original probabilistic atlas, as shown in the original studies (Bianciardi et al., 2018;
García-Gomar et al., 2019; Singh et al., 2020). In contrast, the T1w-based probabilistic ROIs showed
much higher containment: 92.1% of DRN and 91.4% of RN voxels fell within the thresholded
boundaries. This suggests that augmenting the whole-brain T1w-based transform (T_{T1w-WB}) with a
local T1w-based refinement (T_{T1w-BS}) contributes little additional individual specificity; the resulting
two-step transformed ROIs remain largely consistent with the conventional one-step method.

By comparison, the DWI-augmented probabilistic ROIs show a broader spatial distribution,
with smaller proportions of their voxels falling within the thresholded boundaries, namely 61.9% for
b0-augmented DRN, 65.7% for FA-augmented DRN, and 72.7% for multivariate-augmented DRN;
and 86.3%, 70.0%, and 83.7% for the corresponding RN maps. Visually, these ROIs appear less
tightly clustered and more dispersed across subjects. This increased dispersion suggests that the
DWI-augmented registration approaches better capture individual anatomical variability and produce
more personalized ROI mappings.

Figure 5 presents the mean gradient magnitude maps of the Jacobian determinants ($|\nabla JDet|$)
derived from the four second-step transforms, displayed in transverse, coronal, and sagittal planes.
Corresponding IIT templates (T1w, b0, and FA) are shown on the right for anatomical reference. As
indicated by the yellow arrows in the transverse plane, the T1w-based $|\nabla JDet|$ map exhibits a high-
intensity rim that closely matches the outer contour of the brainstem visible in the IIT-T1w template.

This pattern suggests that alignment during the T1w-based registration process was primarily driven
by the brainstem surface. The interior regions of the T1w-based $|\nabla \text{JDet}|$ map display relatively low
gradient magnitudes, though a faint symmetrical pattern implies some internal alignment.

In contrast, the b0- and FA-based $|\nabla \text{JDet}|$ maps show less emphasis on the outer boundary, as
marked in the corresponding locations. Instead, both maps exhibit multiple symmetric, high-
magnitude clusters within the brainstem, indicating enhanced sensitivity to internal anatomical
features. This is particularly evident at the pontomedullary junction in the sagittal plane, where the
T1w-based map remains dark while the FA-based map shows localized bright gradients. This
contrast likely reflects underlying template differences: the IIT-T1w template is relatively
homogeneous in this region, whereas the IIT-FA template reveals intersecting fiber tracts at the
pontomedullary junction.

The multivariate $|\nabla \text{JDet}|$ map combines features observed in the unimodal maps, showing
gradient magnitudes that are more uniformly distributed across the brainstem. This pattern suggests
that the multivariate transform leveraged both external and internal anatomical features from the
three image modalities.

4   Discussion

In this study, we evaluated several DWI-augmented ROI registration pipelines for their
effectiveness in mapping standard brainstem ROIs from IIT space to individual participants' native
spaces, with the design of the pipelines and associated evaluation metrics summarized in Figures 1
and 2. As shown in Figure 3, DWI-derived images improved registration accuracy in a manner that
varied by both ROI and image modality. The greater spatial dispersion observed in the probabilistic
maps of inverse-transformed ROIs, as shown in Figure 4, suggests that DWI-augmented pipelines
may better capture inter-individual anatomical variability. The $|\nabla \text{JDet}|$ maps in Figure 5 further reveal
how different pipelines distribute registration effort across brainstem subregions, offering a
qualitative explanation for the observed ROI- and modality-dependent effects. These findings have
important implications for the brainstem fMRI community, which continues to face technical
challenges, particularly the unreliable sampling of brainstem signals resulting from inaccurate
localization of target ROIs. The findings could also benefit research areas that rely heavily on
accurate data sampling from small brainstem ROIs, such as brainstem histopathology.

There are, however, some limitations and issues in this study that would warrant further
investigation. One key issue concerns the interpretation of the increased spatial dispersion observed

in the DWI-augmented probabilistic ROIs shown in Figure 4. This dispersion may reflect enhanced
sensitivity of the DWI-augmented pipelines to individual anatomical variability. However, it could
also originate from reduced registration accuracy due to the low SNR and greater geometric
distortion of diffusion images. This latter explanation is less likely for RN registration, as the Dice
coefficients shown in Figure 3 indicate improved, rather than diminished, accuracy with the DWI-
augmented pipelines. For DRN registration, however, this possibility cannot be entirely excluded.
The mis-registration fraction metric captures voxel misplacement only in the posterior direction
relative to the DRN, while overlooking displacement in other directions. In this sense, it is less
comprehensive than the Dice coefficient, which, although also a global summary metric, accounts for
voxel misplacement across the entire ROI surface. As a result, it remains less clear whether the
increased dispersion observed in the DRN probabilistic maps reflects genuine anatomical variability
or greater random error in registration, especially compared to the results from the RN.

This uncertainty stems from a broader limitation: the unavailability of MR sequences capable
of reliably delineating the full boundaries of many brainstem nuclei at the individual level. As a
result, alternative evaluation metrics that do not require complete ground-truth segmentation, such as
the mis-registration fraction, must be used to assess registration accuracy for these structures.
Although the Dice coefficient results support the interpretation that DWI-augmented pipelines yield
more anatomically specific registrations, it would be preferable to validate this claim using nuclei
that can be directly visualized and manually segmented in native space, once advanced MR
sequences and ultra-high-field MRI scanners become more widely available. It is worth noting that
even as such sequences are developed, the optimization of atlas-to-subject ROI registration pipelines
will remain important, as incorporating multiple specialized scans to visualize different brainstem
nuclei may not always be practical in research or clinical settings due to limitations in scan time,
participant motion, and subject tolerance (Greene et al., 2016).

Another issue warranting further systematic exploration-though beyond the scope of the present
work-is the refinement of the multivariate registration pipeline. As described in the Materials and
Methods section, the T_{mv-BS} transforms were estimated using equally weighted image modalities,
which is the default setting in ANTs' `antsRegistration` tool for multivariate optimization.
However, the optimal weighting may vary across subregions or specific brainstem nuclei, given
differences in how well individual structures and their surrounding landmarks are visualized in each
modality. For example, Figure 3 shows that incorporating b0-based optimization consistently

improves the registration accuracy of the RN but appears to introduce greater variability in the mis-
registration fraction of the DRN. This discrepancy likely stems from how well each structure can be
visualized: while the b0 image provides direct contrast for the RN, it offers poor visibility in the
dorsal midbrain, where the DRN is located. In contrast, the impact of FA-based optimization on DRN
registration appears more consistent across participants, likely due to the presence of nearby salient
white matter tracts in FA images. These observations suggest that tailoring modality weights based
on prior anatomical knowledge could improve registration performance in an ROI-specific manner.
Further investigation in this direction may enhance the flexibility and accuracy of the proposed
multivariate pipeline across different brainstem subregions.

The significance of our findings is best appreciated in the context of the development and
application of the Brainstem Navigator-the standard-space atlas used to evaluate the DWI-augmented
registration pipelines in this study. As described in a series of publications (Bianciardi et al., 2015;
Bianciardi et al., 2018; García-Gomar et al., 2019; Singh et al., 2020; Singh et al., 2021), the
development of the Brainstem Navigator was guided by the same theoretical assumption as the
present work: that DWI-derived images provide detailed microstructural information suitable for
delineating brainstem nuclei. The Brainstem Navigator team segmented these nuclei using 7-Tesla b0
and FA images from twelve participants-a sample size comparable to that of the current study-via a
semi-automated method combining k-means clustering and manual labeling. These DWI-derived
images and segmented nuclei ROIs were then registered to the IIT diffusion imaging templates to
construct a probabilistic atlas in IIT space. Additionally, the regions of interest (ROIs) were
transformed into MNI152 space using a T1w-based IIT-to-MNI152 registration, with the IIT version
intended for the diffusion imaging community and the MNI152 version for the fMRI community
(Bianciardi et al., 2015). Accordingly, we observed a divergence in ROI registration strategies across
studies utilizing the Brainstem Navigator. While DWI-based studies tend to use the IIT-space version
for diffusion image-based ROI registration (García-Gomar et al., 2021), fMRI studies typically rely
on the T1w-based ROI registration pipelines included in standard fMRI preprocessing toolkits,
applying them to the MNI152-space version of the atlas (Zhang et al., 2025; Hansen et al., 2024).

Although the present study focuses on the application rather than the development of a
brainstem atlas, its methodology partially overlaps with that of the Brainstem Navigator, allowing for
informative comparisons. First, the Brainstem Navigator team delineated a greater number of
brainstem nuclei using only b0 and FA images. Second, although not formally evaluated, their

native-to-IIT registration appeared satisfactory using FA images alone (Bianciardi et al., 2015).
These differences may be attributable to the higher image quality achieved with a 7-Tesla scanner
compared to the 3-Tesla scanner used in the present study. Due to the lower spatial resolution and
signal-to-noise ratio in 3-Tesla DWI data, we were only able to manually delineate the boundaries of
the RN ROI with sufficient confidence, and we needed to employ a metric that did not require full
boundary delineation to assess the registration accuracy of DRN ROI. Additionally, direct
registration of our 3-Tesla DWI-derived images to the IIT diffusion templates yielded suboptimal
results. Therefore, we adopted a two-step registration approach: initial global alignment was
performed using T1w-based whole-brain images (T_{T1w-WB}), followed by local refinement using DWI-
based images through cross-correlation optimization (T_{FA-BS}, T_{b0-BS} , and T_{mv-BS}).

An intriguing aspect of the Brainstem Navigator's approach is that, although the developers
acknowledged T1w images lacked sufficient contrast for delineating brainstem nuclei-and therefore
excluded them during ROI segmentation-they nonetheless employed a T1w-based transformation to
map the atlas from IIT space to MNI152 space (Bianciardi et al., 2015; Bianciardi et al., 2018;
García-Gomar et al., 2019). While this decision was not explicitly justified, it likely reflects the
absence, at the time, of a widely accepted diffusion imaging template in MNI152 space-the default
normalization space used in popular fMRI preprocessing toolkits like SPM and FSL (Zhang &
Arfanakis, 2018; Ashburner et al., 2014; Jenkinson et al., 2012). However, registration in
neuroimaging is an inherently ill-posed problem, where the resulting transformation depends strongly
on the chosen image modality and the cost function used during optimization (Lange et al., 2024).
Because T1w images lack adequate contrast for brainstem nuclei, performing native-to-standard or
standard-to-standard (e.g., IIT-to-MNI152) ROI registration using only T1w images may
compromise the anatomical precision afforded by an atlas developed through DWI-based delineation.
This concern is supported by our findings that DWI-augmented pipelines outperformed the T1w-only
pipeline in the context of IIT-to-native registration. To extend this evaluation to standard-to-standard
registration, we generated IIT-to-OMM mappings by registering the IIT-T1w, IIT-b0, and IIT-FA
templates to the recently released OMM-1 (Oxford-MultiModal-1) T1w, T2w-FLAIR, and FA
templates (Arthofer et al., 2024). These mappings were computed using the same pipelines applied to
IIT-to-native registration in the current study, with the sole modification that the CC cost function for
the T_{b0-BS} pipeline was replaced by MI to accommodate the contrast differences between b0 and
T2w-FLAIR images. As shown in Figure 6A, the $|\nabla JDet|$ maps of the four brainstem-masked
transforms varied substantially, despite all being IIT-to-OMM mappings. Consistent with this, the

transformed RN and DRN ROIs also showed spatial discrepancies across the different registration
pipelines (Figure 6B), mirroring patterns observed in our IIT-to-native results. These differences
underscore the impact of image modality on registration accuracy-a factor that appears to have been
underappreciated in the development of brainstem atlases and brainstem fMRI preprocessing toolkits.
Although preliminary, our findings suggest that incorporating DWI-derived images at every stage of
the registration process could mitigate the loss of accuracy associated with contrast limitations in
certain modalities. For example, normalizing participant data to IIT space-an approach commonly
used by DWI researchers employing the Brainstem Navigator-avoids the intermediate T1w-only IIT-
to-MNI152 registration and may improve registration accuracy of brainstem ROIs. This strategy
would require additional diffusion acquisitions in an fMRI session; however, with appropriate EPI
distortion matching, these data may also be utilized to enhance functional-to-structural co-
registration, as previously proposed and tested with T1w images in selected studies (Renvall et al.,
2016; Zhang et al., 2025).

An important question regarding the incorporation of DWI is how best to integrate it to
maximize its benefits while minimizing potential deleterious effects, if such effects exist. The
Brainstem Navigator atlas was constructed using a sequential approach, in which only one imaging
modality was used at a time. For instance, the DRN was identified by first localizing the
periaqueductal gray (PAG)-DRN complex on the b0 image, followed by delineation of DRN
boundaries using the FA image. Similarly, b0 and FA images were used sequentially to delineate the
outer boundaries and inner subregions of the RN, respectively (Bianciardi et al., 2015). The choice of
modality for delineating specific structures was guided by prior knowledge of which image type
provided superior contrast for each brainstem nucleus. Interestingly, our findings suggest that b0-
based and FA-based registration produced more consistent improvements in registration accuracy for
the RN and DRN respectively (Figure 3), paralleling the Brainstem Navigator team's modality
selection for ROI delineation. This suggests that the registration algorithms were able to leverage
nucleus-specific structural information embedded in the corresponding DWI images to optimize the
transformations. The Brainstem Navigator team also discussed the possibility of adopting a multi-
channel approach, as opposed to a sequential one, for brainstem nuclei segmentation (Bianciardi et
al., 2015). Although our study focused on registration rather than segmentation, informative
comparisons can still be drawn from our results involving single-step (T_{T1w-WB}), sequential (e.g., T_{b0-BS}
$_{BS}$ and T_{FA-BS} following T_{T1w-WB}), and multi-channel (T_{mv-BS} following T_{T1w-WB}) registration pipelines,
as illustrated in Figure 6B. Notably, the second-step T_{b0-BS} transform actually worsened DRN

registration: more DRN ROI voxels were misplaced into the cerebral aqueduct and fourth ventricle
compared to those registered using $T_{T1w\text{-}WB}$ alone. This degradation was not observed with the multi-
channel pipeline, in which FA, b0, and T1w images were simultaneously incorporated and iterated
over within a single registration step. The deleterious effect observed with the $T_{b0\text{-}BS}$ pipeline could
be partly attributed to the use of MI rather than CC as the optimization metric due to subtle contrast
mismatch between IIT-b0 and OMM-1 T2w-FLAIR. However, the fact that the degradation occurred
specifically for the DRN suggests that it may also stem from the relatively poor contrast of these T2w
images at the dorsal midbrain, where the DRN is located. In this case, the multi-channel registration
approach, using default equal weighting across modalities, appears to effectively mitigate the adverse
impact of the lower-contrast modality. As discussed earlier, optimizing the weighting of each
modality may further enhance registration performance, but doing so would require prior knowledge
of nucleus-specific contrast characteristics and systematic empirical evaluation. Importantly, once
such optimized weighting schemes are established for brainstem ROI registration, the same
principles could be translated to brainstem ROI segmentation. Since segmentation also relies on
detecting subtle contrast differences between adjacent nuclei, tailoring the contribution of each
modality based on its ability to resolve specific anatomical boundaries may improve the precision of
segmentation algorithms, particularly in complex or low-contrast regions of the brainstem.

In summary, we proposed and evaluated a theoretically grounded approach to improving the
registration accuracy of brainstem nuclei ROIs, addressing the persistent challenge posed by the
limited availability of MRI sequences that enable direct visualization of these structures. Through a
detailed methodological comparison with studies using Brainstem Navigator, we also offered
cautious recommendations on how the anatomical precision provided by a brainstem atlas could be
better preserved using a DWI-augmented registration pipeline tailored to specific brainstem regions.
Despite certain limitations, most notably, the scarcity of ground-truth nuclei for rigorous validation,
the results demonstrate both the feasibility of the approach and its potential for ROI-specific
optimization. Although this study focused on two scalar images derived from DWI (b0 and FA),
other diffusion-based images, such as the diffusion tensor image (DTI), contain rich, high-
dimensional information that could support more precise local alignment and further augmentation
with global anatomical features, such as tract labels from cortical-subcortical tractography. In the
context of fMRI, cortical-subcortical tractography also provides a structural basis for investigating
the mechanisms underlying functional dynamics aside from serving as a spatial reference frame for
ROI registration, underscoring the value of incorporating DWI into fMRI protocols for enhancing

fidelity and biological plausibility. As ultra-high field MRI becomes more widely available and methods for correcting geometric distortion continue to advance, the integration of diffusion-weighted data into ROI registration pipelines is likely to become increasingly sophisticated. Future research could explore these technical extensions and incorporate functional-to-structural co-registration for a more holistic enhancement of brainstem fMRI reliability.

5   References

Andersson, J. L., Skare, S., & Ashburner, J. (2003). How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage*, *20*(2), 870-888.

Andersson, J. L., & Sotiropoulos, S. N. (2016). An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage*, *125*, 1063-1078.

Arthofer, C., Smith, S. M., Douaud, G., Bartsch, A., Alfaro-Almagro, F., Andersson, J., & Lange, F. J. (2024). Internally consistent and fully unbiased multimodal MRI brain template construction from UK Biobank: Oxford-MM. *Imaging Neuroscience*, *2*, 1-27.

Ashburner, J., Barnes, G., Chen, C. C., Daunizeau, J., Flandin, G., Friston, K., ... & Penny, W. (2014). SPM12 manual. *Wellcome Trust Centre for Neuroimaging, London, UK*, *2464*(4), 53.

Avants, B. B., Tustison, N., & Song, G. (2009). Advanced normalization tools (ANTS). *Insight j*, *2*(365), 1-35.

Beissner, F. (2015). Functional MRI of the brainstem: common problems and their solutions. *Clinical neuroradiology*, *25*(Suppl 2), 251-257.

Bianciardi M, Toschi N, Edlow BL, Eichner C, Setsompop K, Polimeni JR, Brown EN, Kinney HC, Rosen BR, Wald LL. (2015). *Toward an In Vivo Neuroimaging Template of Human Brainstem Nuclei of the Ascending Arousal, Autonomic, and Motor Systems.* Brain Connect. Dec;5(10):597-607

Bianciardi M, Strong C, Toschi N, Edlow BL, Fischl B, Brown EN, Rosen BR, Wald LL. (2018) *A probabilistic template of human mesopontine tegmental nuclei from in vivo 7T MRI.* Neuroimage. Apr 15;170:222-230

Cauzzo, S., Singh, K., Stauder, M., García-Gomar, M. G., Vanello, N., Passino, C., ... & Bianciardi,
M. (2022). Functional connectome of brainstem nuclei involved in autonomic, limbic, pain and
sensory processing in living humans from 7 Tesla resting state fMRI. *Neuroimage*, *250*, 118925.

Chen, X., Huddleston, D. E., Langley, J., Ahn, S., Barnum, C. J., Factor, S. A., ... & Hu, X. (2014).
Simultaneous imaging of locus coeruleus and substantia nigra with a quantitative neuromelanin MRI
approach. *Magnetic resonance imaging*, *32*(10), 1301-1306.

Eckstein, K., Bachrata, B., Hangel, G., Widhalm, G., Enzinger, C., Barth, M., ... & Robinson, S. D.
(2021). Improved susceptibility weighted imaging at ultra-high field using bipolar multi-echo
acquisition and optimized image processing: CLEAR-SWI. *Neuroimage*, *237*, 118175.

Fernández-Gil, M. A., Palacios-Bote, R., Leo-Barahona, M., & Mora-Encinas, J. P. (2010, June).
Anatomy of the brainstem: a gaze into the stem of life. In *Seminars in Ultrasound, CT and MRI* (Vol.
31, No. 3, pp. 196-219). WB Saunders.

Fonov, V. S., Evans, A. C., McKinstry, R. C., Almli, C. R., & Collins, D. L. (2009). Unbiased
nonlinear average age-appropriate brain templates from birth to adulthood. *NeuroImage*, *47*, S102.

García-Gomar MG, Strong C, Toschi N, Singh K, Rosen BR, Wald LL, Bianciardi M. (2019) *In vivo*
Probabilistic Structural Atlas of the Inferior and Superior Colliculi, Medial and Lateral Geniculate
Nuclei and Superior Olivary Complex in Humans Based on 7 Tesla MRI. Front Neurosci. Aug
7;13:764

García-Gomar MG, Videnovic A, Singh K, Stauder M, Lewis LD, Wald LL, Rosen BR, Bianciardi
M. (2021) *Disruption of brainstem structural connectivity in RBD using 7 Tesla MRI.* Mov Disord.
Dec 29.

Greene, D. J., Black, K. J., & Schlaggar, B. L. (2016). Considerations for MRI study design and
implementation in pediatric and clinical populations. *Developmental cognitive neuroscience*, *18*, 101-
112.

Hansen, J. Y., Cauzzo, S., Singh, K., García-Gomar, M. G., Shine, J. M., Bianciardi, M., & Misic, B.
(2024). Integrating brainstem and cortical functional architectures. *Nature Neuroscience*, 1-12.

Hoopes, A., Mora, J. S., Dalca, A. V., Fischl, B., & Hoffmann, M. (2022). SynthStrip: skull-stripping
for any brain image. *NeuroImage*, *260*, 119474.

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012).
Fsl. *Neuroimage*, *62*(2), 782-790.

Lange, F. J., Arthofer, C., Bartsch, A., Douaud, G., McCarthy, P., Smith, S. M., & Andersson, J. L.
(2024). MMORF-FSL's MultiMOdal registration framework. *Imaging Neuroscience*, *2*, 1-30.

Matt, E., Fischmeister, F. P. S., Amini, A., Robinson, S. D., Weber, A., Foki, T., Gizewski, E. R.,
Beisteiner, R. (2019). Improving sensitivity, specificity, and reproducibility of individual brainstem
activation. *Brain Structure and Function*, *224*(8), 2823-2838.

Mazancieux, A., Mauconduit, F., Amadon, A., de Gee, J. W., Donner, T. H., & Meyniel, F. (2023).
Brainstem fMRI signaling of surprise across different types of deviant stimuli. *Cell Reports*, *42*(11).

Mugler III, J. P., & Brookeman, J. R. (1990). Three‐dimensional magnetization‐prepared rapid
gradient‐echo imaging (3D MP RAGE). *Magnetic resonance in medicine*, *15*(1), 152-157.

Pessoa, L., Medina, L., Hof, P. R., & Desfilis, E. (2019). Neural architecture of the vertebrate brain:
implications for the interaction between emotion and cognition. *Neuroscience & Biobehavioral*
Reviews, *107*, 296-312.

Priovoulos, N., Jacobs, H. I., Ivanov, D., Uludağ, K., Verhey, F. R., & Poser, B. A. (2018). High-
resolution in vivo imaging of human locus coeruleus by magnetization transfer MRI at 3T and
7T. *Neuroimage*, *168*, 427-436.

Renvall, V., Witzel, T., Wald, L. L., & Polimeni, J. R. (2016). Automatic cortical surface
reconstruction of high-resolution T1 echo planar imaging data. *Neuroimage*, *134*, 338-354.

Sclocco, R., Beissner, F., Bianciardi, M., Polimeni, J. R., & Napadow, V. (2018). Challenges and
opportunities for brainstem neuroimaging with ultrahigh field MRI. *Neuroimage*, *168*, 412-426.

Shepherd, T. M., Ades-Aron, B., Bruno, M., Schambra, H. M., & Hoch, M. J. (2020). Direct in vivo
MRI discrimination of brain stem nuclei and pathways. *American Journal of Neuroradiology*, *41*(5),
777-784.

Singh, K., Cauzzo, S., García-Gomar, M. G., Stauder, M., Vanello, N., Passino, C., & Bianciardi, M.
(2022). Functional connectome of arousal and motor brainstem nuclei in living humans by 7 Tesla
resting-state fMRI. *NeuroImage*, *249*, 118865.

Singh K, García-Gomar MG, Bianciardi M. (2021) *Probabilistic Atlas of the Mesencephalic*
Reticular Formation, Isthmic Reticular Formation, Microcellular Tegmental Nucleus, Ventral
Tegmental Area Nucleus Complex, and Caudal-Rostral Linear Raphe Nucleus Complex in Living
Humans from 7 Tesla Magnetic Resonance Imaging. Brain Connect. Oct;11(8):613-623

Singh K, Indovina I, Augustinack JC, Nestor K, García-Gomar MG, Staab JP, Bianciardi M. (2020)
Probabilistic Template of the Lateral Parabrachial Nucleus, Medial Parabrachial Nucleus,
Vestibular Nuclei Complex, and Medullary Viscero-Sensory-Motor Nuclei Complex in Living
Humans From 7 Tesla MRI. Front Neurosci. Jan 23;13:1425.

Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee, J. C.
(2010). N4ITK: improved N3 bias correction. *IEEE transactions on medical imaging*, *29*(6), 1310-
1320.

Tustison, N. J., Cook, P. A., Holbrook, A. J., Johnson, H. J., Muschelli, J., Devenyi, G. A., ... &
Avants, B. B. (2021). The ANTsX ecosystem for quantitative biological and medical
imaging. *Scientific reports*, *11*(1), 9068.

Van den Heuvel, M. P., & Sporns, O. (2013) An anatomical substrate for integration among
functional networks in human cortex. *Journal of Neuroscience*, *33*(36), 14489-14500.

Zhang, J., Chen, D., Deming, P., Srirangarajan, T., Theriault, J., Kragel, P. A., ... & Bianciardi, M.
(2025). Cortical and subcortical mapping of the allostatic-interoceptive system in the human brain
using 7 Tesla fMRI. *BiorXiv*, 2023-07.

Zhang S, Arfanakis K. (2018) Evaluation of standardized and study-specific diffusion tensor imaging
templates of the adult human brain: Template characteristics, spatial normalization accuracy, and
detection of small inter-group FA differences. Neuroimage. May 15;172:40-50.

Figure Captions

Figure 1. ROI registration procedure. (A) Estimation of transforms for the two-step registration pipelines. The whole-brain transform (TT1w-WB) was estimated via sequential linear (mutual information, MI) and nonlinear (cross-correlation, CC) alignment between the IIT-T1w template and the participant's native T1w image. TT1w-WB was then applied to the IIT-FA and IIT-b0 templates to generate warped intermediate templates. Brainstem-specific transforms (TFA-BS, TT1w-BS, and Tb0-BS) were subsequently estimated by CC-based nonlinear registration between each warped template and its corresponding participant image. (B) ROI registration using compound transforms. The transforms obtained in (A) were concatenated to form four compound transforms-either including or excluding diffusion-based alignment-which were then applied to register brainstem ROIs from IIT space to native space.

Figure 2. Evaluation metrics for assessing registration accuracy of the red nucleus (RN) and dorsal raphe nucleus (DRN). The Dice coefficient was used to evaluate RN registration accuracy by quantifying the spatial overlap between the registered ROI and the manually delineated ground-truth ROI. It is calculated as the ratio of twice the intersection volume to the sum of the two ROI volumes. For DRN, registration accuracy was assessed using the mis-registration fraction, defined as the proportion of voxels in the transformed DRN ROI that overlapped with the ground-truth ROI of the cerebral aqueduct and fourth ventricle.

Figure 3. Registration accuracy of the four ROI registration pipelines. (A) Dice coefficients for the red nucleus (RN) across ten participants, with each participant represented by a separate colored line. Mean Dice coefficients ± standard deviation: T1w = 0.72 ± 0.02; b0 = 0.75 ± 0.02; FA = 0.76 ± 0.05; multivariate (mv) = 0.75 ± 0.03. (B) Mis-registration fractions for the dorsal raphe nucleus (DRN) across the same participants. Mean values ± standard deviation: T1w = 0.28 ± 0.06; b0 = 0.35 ± 0.13; FA = 0.18 ± 0.08; mv = 0.26 ± 0.08.

Figure 4. Probabilistic ROIs of the red nucleus (RN) and dorsal raphe nucleus (DRN). The leftmost column shows the unthresholded probabilistic ROIs from the Brainstem Navigator atlas (blue), overlaid with the boundaries of the thresholded binary ROIs at probability = 0.35 (red). Columns from second-to-left through rightmost display the probabilistic ROIs constructed by averaging inverse-transformed native-space ROIs obtained from each of the four two-step registration pipelines (yellow), overlaid with the same thresholded ROI boundaries (red).

Figure 5. Gradient magnitude maps of brainstem-specific transforms and corresponding templates.
Left: Mean gradient magnitude maps of the Jacobian determinants ($|\nabla$JDet$|$) computed from each
brainstem-specific transform, shown in transverse, coronal, and sagittal planes. Right: Corresponding
IIT templates (T1w, b0, and FA) displayed in the same anatomical planes for reference. Yellow
arrows highlight subregional differences in deformation magnitude, which may reflect variation in
anatomical landmark visibility across image modalities.

Figure 6. Gradient magnitude maps of brainstem-specific IIT-to-OMM transforms. (A) Mean
gradient magnitude maps of the Jacobian determinants ($|\nabla$JDet$|$) computed from each brainstem-
specific transform, shown in transverse, coronal, and sagittal planes. (B) Brainstem ROIs transformed
from the IIT space to the OMM-1 space, overlaid on the OMM-1 T1w template. Upper: RN. Lower:
DRN. b0: concatenated T_{T1w-WB} and T_{T1w-BS}. FA: concatenated T_{T1w-WB} and T_{FA-BS}. T1w:
concatenated T_{T1w-WB} and T_{T1w-BS}. mv: concatenated T_{T1w-WB} and T_{mv-BS}. T1w (1^{st} step): T_{T1w-WB}
only.

Acknowledgements

620 The authors thank Asma Naheed, MRT for the scanning support at the Slaight Family Centre for
621 Advanced MRI, Toronto Western Hospital, University Health Network, Toronto, Canada.

Conflict of Interest

623 AML serves as a consultant for Boston Scientific and Abbott. All other authors declare no conflicts
624 of interest.

Author Contributions

YAC: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software,
Visualization, Writing-original draft.

LK: Data curation, Writing-review & editing.

CTC: Data curation, Writing-review & editing.

AB: Data curation, Writing-review & editing.

JG: Data curation, Writing-review & editing.

AML: Funding acquisition, Resources, Writing-review & editing.

KU: Funding acquisition, Project administration, Resources, Supervision, Writing-review & editing.

AOD: Funding acquisition, Supervision, Writing-review & editing.

SK: Conceptualization, Data curation, Project administration, Resources, Software, Supervision,
Writing-review & editing.

Funding

638   This work was supported by funding from the Alan and Susan Hudson Chair in Neurosurgery,
639   University Health Network and University of Toronto (AML and KU), and by the NSERC Discovery
640   Grant 214566 (AOD).

Red Nuclei

Dorsal Raphe Nucleus

$$\text{Dice Coefficient} = \frac{2 \times (\bullet \cap \bullet)}{\bullet + \bullet}$$

$$\text{Mis-registration Fraction} = \frac{\cap}{}$$

A Red Nuclei

B Dorsal Raphe Nucleus

Jacobian Determinant Gradient Maps

IIT Templates