Title: A connexin 43 targeting peptide prevents blood vessel neointima formation

Running Title: CycliCx peptide disrupts neointima

Authors: Meghan W. Sedovy[1,2#], Mark C. Renton[1#], Xinyan Leng[1], Farwah Iqbal[3], Melissa R.
Leaf[3], Kailynn Roberts[1,2], Ramya C. Joshi[1], Arya Malek[1], Clare L. Dennison[1], Angela K Best[4],
Paul D. Lampe[5], Joseph W. Baker[3,6], Mark Joseph[3,6], George S. Baillie[7], Brant E. Isakson[4,8] and
Scott R. Johnstone[1,3,6,9*]

Affiliations:

[1] The Fralin Biomedical Research Institute at Virginia Tech Carilion, Center for Vascular and Heart
Research, 4 Riverside Circle, Roanoke, Virginia, VA 24016 USA

[2] Translational Biology, Medicine, and Health Graduate Program, Virginia Tech, Blacksburg, VA
24061, USA

[3] Virginia Tech Carilion School of Medicine, Roanoke, VA 24016 USA

[4] Robert M. Berne Cardiovascular Research Center, University of Virginia School of Medicine,
Virginia, VA 22908, USA

[5] Fred Hutchinson Cancer Research Center, Seattle, WA 98109 USA.

[6] Department of Surgery, Virginia Tech Carilion School of Medicine, Roanoke, VA 24016 USA

[7] School of Cardiovascular and Metabolic Health, University of Glasgow, Glasgow, UK G128QQ,
UK

[8] Department of Molecular Physiology and Biophysics, University of Virginia School of Medicine,
Virginia, VA 22908, USA

[9] Department of Biological Sciences, College of Science, Virginia Tech, Blacksburg, VA 24061,
USA

Coauthors: Meghan W. Sedovy and Mark C. Renton share the position of first author.

* **Corresponding Author:**

Scott R. Johnstone, The Fralin Biomedical Research Institute at Virginia Tech Carilion, Roanoke,
VA 24016, USA. Email: scottrj@vt.edu Phone: (+44) 540-526-2296

Conflict-of-Interest Statement:

The authors have declared no conflict of interest exists. Peptides generated and used in this
manuscript are covered by patent #WO2021021716 (SRJ, BEI).

Abstract:

The gap junction protein connexin 43 (Cx43) is associated with human pathological vascular smooth muscle cell (SMC) proliferation and neointima formation. We previously identified mitogen-activated protein kinase (MAPK) phosphorylation of Cx43 results in binding with the cell cycle protein cyclin E, facilitating neointima formation in mice. However, the specific nature of these interactions and their relevance to human disease have not been elucidated. Using an *ex vivo* human saphenous vein model of neointima formation, we identified increased MAPK-phosphorylated Cx43 and cyclin E in explant tissues. We used peptide arrays to define a cyclin E-Cx43 binding region and generated 'CycliCx', a stearate-linked Cx43 phospho-mimetic peptide. In human coronary artery SMC, CycliCx inhibits platelet-derived growth factor-BB (PDGF-β)-induced changes in Cx43 trafficking and interactions with cyclin E, and stimulation of proliferation. RNAseq analysis identified CycliCx significantly inhibits PDGF-β-induced proliferative pathways in SMC by limiting PDGF-induced early G1/S phase cell cycle progression transcripts. Finally, we show CycliCx limits neointima formation in mice *in vivo* and in *ex vivo* human saphenous vein explants. Our data provide strong evidence for selective targeting Cx43 as a viable therapeutic strategy for preventing neointimal formation in humans.

Introduction:

Atherosclerosis in the coronary artery narrows the vessel lumen, restricts arterial blood flow, affecting more than 20.5 million Americans, making it the primary underlying cause of cardiovascular disease[1]. Surgical interventions, such as coronary artery bypass and percutaneous coronary interventions (angioplasty/stent placement) can be employed to revascularize and alleviate symptoms and risks of coronary artery disease. Of those who undergo surgical revascularization, 16.4% will experience a serious adverse event, including a heart attack or death[2]. This highlights a need for novel therapeutics that can enhance vessel patency following revascularization surgeries.

The leading cause of long-term patency issues after coronary artery bypass grafting (CABG) or percutaneous coronary intervention is neointimal hyperplasia[3], which occurs when vascular smooth muscle cells (SMC) in the medial layer of arteries/ vascular conduits dedifferentiate, take on a proliferative and migratory phenotype, and block the vessel intima[3]. Drug-eluting stents are often used to limit neointimal growth, yet these stents use broad antiproliferative agents, which delay vessel healing and necessitate the prolonged use of dual antiplatelet therapy, increasing the risk for clinically significant bleeding[3-5]. As such, identifying mechanisms of neointima development specific to SMC proliferation is essential in developing novel therapeutics.

Connexins form hexamers on the plasma membrane, which dock with connexin hexamers on adjacent cells, forming gap junctions, facilitating cell-cell communication[6]. Increasing evidence suggests connexins also regulate cell function in a gap junction-independent manner through protein-protein interactions[7-14]. In SMC, altered Cx43 expression has been associated with neointimal hyperplasia, but the specific role Cx43 plays is currently unresolved, with conflicting studies showing both increased and decreased proliferative states in response to altered Cx43 expression[15-19]. Additionally, decreased expression and gap junction signaling are associated with both enhanced and delayed vascular disease progression[15,16,20-22].

Nuclear magnetic resonance analysis of the cytoplasmic Cx43 C-terminus reveals two α-helical regions with a randomly coiled structure for the remaining length, forming a flexible protein primed for phosphorylation and protein-protein interactions[23]. In addition, Cx43 contains a long intracellular C-terminus with a high percentage of amino acids (a.a.) that can be post-translationally modified[10,24,25]. The phosphorylation states influence the conformation of the Cx43 C-terminus, altering the accessibility of other Cx43 phosphorylation sites and regulating the Cx43 protein-protein binding affinity[26]. Our previous study showed in response to platelet-derived growth factor-BB (PDGF-β), Cx43 is phosphorylated by mitogen-activated protein kinases

(MAPK) at four of its C-terminal serines[24]. Phosphorylation by MAPK promotes Cx43 binding to

the cyclin E protein, which binds with CDK2 to promote G1/S phase cell cycle transition[24,27], and

results in SMC proliferation in mice[24]. Our previous data suggested disrupting Cx43-MAPK

activation does not compromise the viability of healthy SMC *in vivo,* making it a suitable target for

therapeutic intervention[24]. In transgenic Cx43-MAPK phosphorylation-null mice, Cyclin E-Cx43

interactions were inhibited, and ligation-induced carotid artery neointima was reduced[24]. Despite

this, the binding sites for Cx43 and cyclin E have not been elucidated yet.

In the present study, we aimed to define the importance of cyclin E-Cx43 interactions in human

vascular tissues during neointima formation, and hypothesized disrupting cyclin E-Cx43 protein-

protein interactions could prevent neointima formation. We developed models of *ex vivo* human

neointima formation using excess human saphenous veins intended for coronary artery bypass

grafts and show cyclin E-Cx43 interactions form during PDGF-β-induced SMC proliferation. We

used peptide array techniques to define a 14 a.a., phospho-dependent cyclin E binding site on

Cx43 and developed a stearate-linked linear peptide ('CycliCx') to disrupt cyclin E-Cx43

interactions. *In vitro* testing demonstrated CycliCx inhibits PDGF-β-induced changes in Cx43

localization and phosphorylation state, limits cyclin E-Cx43 interactions, and reduces human SMC

proliferation. Transcriptomics data highlighted the CycliCx peptide disrupts PDGF-β-induced gene

pathways regulating G1/S phase cell cycle progression in cultured human SMC. Importantly, our

data show CycliCx is a potent inhibitor of neointima in mouse and human models. This study is

the first to demonstrate specific binding domains for cyclin E-Cx43, show relevance in human

disease, and identify a specific molecule targeted to SMC that limits neointima formation.

Results

A human saphenous vein model of neointima formation shows Cx43 MAPK-phosphorylation in new neointima cells. Previous *ex vivo* models of human saphenous vein neointima developed by George et al.[28,29] demonstrated high serum levels promote SMC proliferation and migration to the intimal surface of tissue explants, forming a "new intima" on the surface of the explants. We aimed to adapt this model to examine the effects of PDGF-β-induced proliferation using low serum media (**Figure 1A**). Human saphenous vein tissue explants had no treatment (NT; no PDGF-β) or 14 day treatment (PDGF-β, 50 ng/mL). Histological staining demonstrated an increase in cell layers at the surface of the explants in PDGF-β-treated tissue (**Figure 1B, Supplemental Figure 1**). The appearance of the cells is consistent with the new intima in previously published models for *ex vivo* HSV intimal growth[28,29]. Immunofluorescent staining demonstrated cells in the upper layers of the tissue are transgelin (TGLN) positive SMC, with significant increases in PDGF-β-treated tissues (**Figure 1B, Supplemental Figure 2**). Staining also identified high levels of expression for total Cx43 and cyclin E within these cells, and antibodies specific to MAPK-phosphorylated sites on Cx43[24] highlight PDGF-β promotes Cx43-MAPK phosphorylation (**Figure 1B, Supplemental Figures 3-5)**.

Identification of a binding region between Cx43 and cyclin E and development of the CycliCx peptide. Our previously published data suggest phosphorylation at Cx43-MAPK C-terminal serines (S) regulates protein interactions between cyclin E and the Cx43 C-terminus[24]. However, specific sites of interaction are unknown, and structural modeling of Cx43 is limited. Therefore, we used a peptide array to define the critical binding sites of the Cx43 C-terminus required for binding with cyclin E. Peptide arrays of the full Cx43 C-terminus (a.a. 236-382) were designed with 25 a.a. peptide sequences arranged sequentially, with a five a.a. overlap between each sequence (**Figure 2A**). Arrays were arranged to test a requirement of the Cx43-MAPK serines located at S255/S262/S279/S282 (MK4) on Cx43, using either: non-mutated serines (Cx43^{wt}, non-phosphorylated state), mutated alanine (Cx43^{MK4A}, non-phosphorylatable state), mutated aspartate (Cx43^{MK4D}, phosphorylated state mimic), or native phosphoserine (Cx43^{MK4pS}, phosphorylated state). The full-length cyclin E protein was synthesized, as we previously described[24] (**Figure 2B**). Isolated cyclin E proteins were incubated with Cx43 peptide arrays, and binding was detected using antibodies targeting cyclin E (**Figure 2C-D**). Initial peptide arrays identified three potential binding regions in each of the Cx43^{MK4D} and Cx43^{MK4pS} sequences (**Figure 2C-D**). Given consensus between D/pS binding regions, we focused on a.a. 236-301, which were divided into three separate, but overlapping, 25 a.a. peptide sequences: Seq1 a.a.

235-260 (gray), Seq 2 a.a. 254-279 (green), Seq 3 a.a. 276-301 (black) (**Figure 2 E-F**). To test the requirement of single a.a., we prepared arrays using alanine substitutions along the length of each sequence, incorporating phosphorylation of Cx43-MAPK serines where they occurred. These data indicated there was consistent binding when phosphorylation was present in Seq 1 and Seq 2, but no evidence of binding in Seq 3 (**Figure 2G**). Further, single a.a. substitutions did not significantly impact overall binding in Seq 1-2 (**Supplemental Figure 6A**).

Using a truncation mutation peptide array approach, which reduces the sequence length by a single a.a. at a time, we tested variants within the identified short region GVKDRVKGRSD (Peptide-1, 11 a.a.) and longer region LDPSKDCGDPKYAYFNGCSSPT (Peptide-2, 22 a.a.) that showed cyclin E binding (**Supplemental Figure 6B-C**). We synthesized peptides for these sequences conjugated with an N-terminal stearyl group to aid in peptide cellular internalization. Peptides (30 µM) were incubated with cell cycle-stalled primary human coronary artery smooth muscle cells (CASMC) treated with PDGF-β (50 ng/mL) to promote proliferation. We used 5-ethynyl 2´-deoxyuridine (EDU) to label newly synthesized DNA and measured proliferation by flow cytometry. At 48 hours, PDGF-β increased proliferation in CASMC, which was reduced by Peptide 2 without notable cell loss (**Supplemental Figure 6D**). However, Peptide 1 treatment resulted in significant cell loss and was not tested further (**Supplemental Figure 6D**).

Peptide arrays indicated a minimum binding sequence within Peptide 2, between a.a. 254-268, but a loss of signal after removal of Tyrosine 267 (**Figure 2H, Supplemental Figure 6C).** Because of this, we created two peptides. The first peptide, named 'CycliCx', is a 14 a.a. sequence, LDPSKDCGDPKYAY, which showed cyclin E-Cx43 binding. The second control off-target peptide (OT-peptide) is a 15 a.a. peptide, AYFNGCSSPTAPLDP, which represents the immediately adjacent a.a. sequence unable to bind cyclin E. Both peptides were synthesized with an N-terminal stearate to promote cell entry, giving a final design of stearate-LDPSKDCGDPKYAY (CycliCx), and stearate-AYFNGCSSPTAPLDP (OT peptide), which was used as a control treatment throughout this study (**Fig 2H-I**). Because the CycliCx sequence was initially based on the mouse Cx43 sequence, we created a humanized CycliCx peptide, stearate-LDPAKDCGDQKYAY, for testing (**Fig 2I**). We saw no differences between mouse and human CycliCx peptides in effect in cells (not shown). The human CycliCx peptide was used for all future experiments. The two peptides have approximately the same molecular weight and charge. The CycliCx peptide has a sequence predicted MW of 1571 kDa, (1853.22 kDa with stearate group) and a charge of -1, and the OT peptide has a predicted MW of 1586 kDa (1806.01 kDa with stearate group) and a charge of -1 (**Supplemental Figure 6E**).

CycliCx peptide alters Cx43 cellular functions in human CASMC. Our previous data showed

PDGF-β-induced ERK phosphorylation is a key factor in regulating cyclin E-Cx43 protein

interactions in mouse primary SMC[24]. To assess ERK activation, human CASMC were treated

with PDGF-β (50 ng/mL) to induce proliferation. Treatment with PDGF-β significantly upregulated

ERK phosphorylation (pERK) in CASMC (**Figure 3A, Supplemental Figure 7**). Treatment of

CASMC with either CycliCx or OT-peptide (30 µM each) did not significantly alter levels of PDGF-

β-induced pERK in CASMC (**Figure 3A, Supplemental Figure 7**). In addition, total Cx43

expression was not affected by PDGF-β in CASMC; however, CycliCx altered Cx43 western blot

protein banding, while OT-peptide had no effect (**Figure 3B, Supplemental Figure 8**). Based on

previous reports of phosphorylation states present in each migrating band, we assessed Cx43

protein kinase-C (PKC) phosphorylation at Cx43-S368 (Cx43-pS368). These data highlight

CycliCx significantly increases the phosphorylation at Cx43-S368 in CASMC (**Figure 3B,**

Supplemental Figure 8). While most experiments in cells were performed at 16 hours, we found

CycliCx produced alterations in Cx43 as early as 3 hours (**Supplemental Figure 9**).

To assess Cx43 expression and localization, we performed immunofluorescence of cultured

CASMC and identified punctate staining characteristic of Cx43 which was not significantly altered

by treatment with PDGF-β or PDGF-β+OT peptide (**Figure 3C, Supplemental Figure 10**). In

PDGF-β + CycliCx-treated cells, Cx43 staining punctates were significantly reduced and staining

appeared more diffuse (**Figure 3C, Supplemental Figure 10**). Membrane biotinylation

experiments demonstrated PDGF-β and PDGF-β +OT peptide treatments reduced Cx43

expression at the plasma membrane. CycliCx inhibited the PDGF-B-induced removal of Cx43

from the plasma membrane (**Figure 3D, Supplemental Figure 11**). To test if CycliCx altered gap

junction signaling, we performed lucifer yellow scrape load dye transfer assays. We found CycliCx

inhibited Cx43 gap junction communication compared to controls (**Figure 3E, Supplemental**

Figure 12). Finally, we tested if CycliCx could inhibit interactions between Cx43 and cyclin E. Co-

immunoprecipitation assays demonstrate PDGF-β treatments increased cyclin E-Cx43 complex

formation in human CASMC ([24]), but were significantly reduced by treatment with CycliCx (**Figure**

3F, Supplemental Figure 13).

CycliCx peptide inhibits PDGF-β-induced proliferation by blocking G1/S phase cell cycle

transition. To test if the peptide alters the proliferation of SMC in a human model, we assessed

PDGF-β-induced proliferation in human CASMC by EDU flow cytometry. We found PDGF-β

significantly upregulated CASMC proliferation, which was inhibited by treatment with CycliCx but

not OT-peptide. (**Figure 4A**, **Supplemental Figure 14**). Bulk RNA sequencing of CASMC treated

revealed PDGF-β+CycliCx altered the transcript expression of more genes than PDGF-β+OT
peptide when compared with PDGF-β alone (**Figures 4D-E**). Pathway enrichment analysis using
the Gene Ontology (GO) database demonstrated a significant upregulation of pathways involved
in cell cycle progression and mitotic nuclear division in PDGF-β-treated compared to control
CASMC, indicating CASMC differentiation into a pro-proliferative phenotype (**Figure 4F**). The
addition of the CycliCx peptide resulted in significantly decreased enrichment of these same
proliferative pathways when compared to PDGF-β treatment alone and to PDGF-β+OT peptide-
treated CASMC (**Figure 4F, Supplemental Figure 16**).

Furthermore, CycliCx treatment significantly blunted the PDGF-β-induced transcript expression
of large arrays of genes involved in 'cell cycle G1/S phase transition' (GO:0000082; **Figure 4G**),
'cell cycle G2/M phase transition' (GO:0044839), and 'mitotic nuclear division' (GO:0140014;
Supplemental Figure 16). Importantly, PDGF-β+OT peptide treatment did not significantly blunt
the PDGF-β-induced expression of these genes. We next interrogated the expression of key
regulatory genes at each phase of the cell cycle (**Figure 4G**; **Supplemental Figure 17**) and found
CycliCx treatment had no effect on the PDGF-β-induced expression of Ccnd1 (Cyclin D1) and
CDK4, which regulate early G1 progression, but partially blunted CDK6 and Cdkn2c (p18)
expression (**Supplemental Figure 17**). Although CycliCx impairs binding of Cyclin E, it had no
effect on Ccne1 (cyclin E1) gene expression (**Figure 4H)**. However, CycliCx significantly blunted
the PDGF-β-induced increase in the expression of G1/S phase transition promoters CDK2 and
E2f1 and partially reversed the PDGF-β-induced decrease in expression of G1/S phase transition
inhibitors Tp53 (p53) and Cdkn1a (p21) (**Figure 4H**; **Supplemental Figure 17**). Finally, we
assessed late G2/M phase regulatory genes and found CycliCx inhibited the PDGF-β-induced
increase in Ccna2 (Cyclin A2) and Cdk1, while OT peptide treatment had no effect (**Figure 4H**;
Supplemental Figure 17). Last, CycliCx did not alter the expression of Cx43, Timp1 (**Figure 4H**),
Timp3, and Il1b, or the downstream effector kinases Mapk3 (Erk1) and Mapk1 (Erk2;
Supplemental Figure 18). Smooth muscle contractile markers were decreased by treatment with
PDGF-β, but these do not appear to be directly altered by treatments with CycliCx or OT peptide
(**Supplemental Figure 19**).

CycliCx peptide disrupts neointima formation in mice and human blood vessels. To test
CycliCx *in vivo*, we combined carotid ligation surgeries in mice with the application of peptides
mixed in pluronic gel to the outside of carotids (**Figure 6A**). We found significant neointima
formation in pluronic controls and OT peptide-treated mice compared with sham ligation mice
(**Figure 6A**). Treatment with CycliCx peptide significantly inhibited ligation-induced neointima in

mice and was not significantly different from sham control mice (**Figure 6A, Supplemental**

Figure 20). We found no sex differences in mice in terms of neointima formation or the effect of

the peptide, and data shown represent both male and female mice (**Supplemental Figure 20**).

We then tested the effect of CycliCx in PDGF-β-induced neointima formation in ex vivo human

saphenous veins. As shown in **Figure 1B and Supplemental Figures 1-5**, PDGF-β promotes

the formation of SMC-derived new intima regions at the surface of tissue explants. This is

significantly increased when compared with no treatment at 14 days and is unaffected by OT-

peptide treatments (**Figure 6B, Supplemental Figures 1-5)**. In *ex vivo* human saphenous veins,

CycliCx significantly reduces the presence of SMC in the intimal region of explants, suggesting

inhibition of neointima formation (**Figure 6B, Supplemental Figures 1-5)**.

Discussion

Targeted therapies to address the mechanisms of neointima formation are needed to reduce the risk of stent/bypass failure. Our previously published data identified phosphorylation of Cx43 at its C-terminus MAPK residues regulates SMC proliferation and neointima formation in mouse models[24]. This control of SMC proliferation is dependent on interactions between the Cx43 C-terminus and cell cycle-regulating protein cyclin E[24]. In the present study, we defined for the first time the binding region of cyclin E on Cx43 and demonstrated MAPK phosphorylation sites within this binding region are required for direct cyclin E-Cx43 interactions. We developed the CycliCx peptide, which mimics a short 14 a.a phosphorylated region of the Cx43 C-terminus within the MAPK region, and was designed to inhibit cyclin E-Cx43 binding. Our data highlight CycliCx alters Cx43 localization, signaling, and trafficking while limiting cyclin E binding with Cx43. Using a transcriptomic approach, we found CycliCx significantly blunts PDGF-β-induced cell cycle progression, leading to reduced proliferation in cultured human CASMC. Finally, we showed CycliCx significantly alters neointima formation in models of disease in mouse and human tissues. This work establishes critical binding sites on Cx43 required for SMC proliferation and identifies Cx43 as a viable therapeutic target for the prevention of human neointima formation.

Current strategies for mitigating neointima formation after stenting procedures primarily include the use of non-cell-specific drugs such as the Rapamycin analogs, Everolimus and Zotarolimus. These drugs work by targeting mTOR signaling, a pathway essential for protein, nucleotide, and lipid synthesis, ultimately regulating cell growth and division[30]. Although mTOR inhibitors can stop the pathological proliferation and migration of SMC, which form neointima, they present a risk to other cell types in the vasculature, particularly endothelial cells that protect the vessel from platelet aggregation and ultimately thrombus formation[4]. Our recent investigation revealed significant endothelial damage caused by standard preservation techniques during coronary bypass surgery, and highlighted the need for endothelial healing[31]. Targeting cyclin E-Cx43 interactions disrupts a key signal required for pathological SMC growth without impacting cell viability or global mTOR signaling. As such, this is a promising strategy for targeting neointima formation/restenosis without impacting healthy cellular healing functions and adversely impacting patient health.

Emerging research reveals Cx43 proteins can act independent of forming cell-cell gap junction channels to control cellular functions[7-13]. We previously found Cx43 interacts with the cell cycle control protein cyclin E in mouse SMC *in vitro* and *in vivo*. Cyclin E tightly controls G1/S phase cell cycle transition, which is critical for pathological SMC proliferation[27,32]. Our data in human saphenous vein explants showed both Cx43 and cyclin E are expressed in the new intima layers

after PDGF-β stimulation, suggesting a role for Cx43 in the SMC proliferation in these tissues.

Given our previous evidence that blocking these interactions is important in mouse neointima

formation, we set out to define the role of cyclinE-Cx43 interactions in human neointima and SMC

proliferation.

This is the first study to comprehensively define the location of cyclin E-Cx43 binding. We

identified two regions on the Cx43 C-terminus with consistent cyclin E binding activity and used

these to generate Cx43 mimetic peptides. The first region, used to generate peptide-1, spans

Cx43 a.a. 235-245, which does not encompass any reported Cx43-MAPK phosphorylation sites.

This peptide is highly similar to the Cx43 mimetic peptide juxtamembrane 2 (JM2)[33]. The JM2

peptide sequence is designed to cover the binding region between Cx43 and microtubule subunit

β-tubulin[33]. JM2 enhances the Cx43 β-tubulin interactions, reduces Cx43 trafficking to the plasma

membrane, and reduces Cx43 hemichannel ATP release[33,34]. In our *in vitro* model of PDGF-β-

induced CASMC proliferation, the overlapping peptide 1 resulted in cell death, and further testing

of this peptide was not performed, as our goal is to limit CASMC from entering a proliferative

phenotype while maintaining vessel health. This is in line with previous work in 4T1 tumor cells,

in which JM2 was found to induce ROS-mediated apoptosis by disrupting mitochondrial

trafficking[35]. However, no JM2 cytotoxicity was reported in healthy cultured human microvascular

endothelial cells[33]. Conflicting cellular responses to the JM2/peptide 1 sequences could arise from

differences in cell types, with CASMC more susceptible than the endothelium. Duration of

treatment and dose may also be responsible for the differential effects reported.

The second cyclin E-Cx43 binding region, used to generate peptide 2 and the final CycliCx

peptide, spans Cx43 a.a. 254-275, with phosphomimetic aspartate mutations at the known Cx43-

MAPK phosphorylation sites, S255 and S262. Cyclin-dependent kinase 1 (CDK1) and ERK5 are

implicated in the phosphorylation of Cx43 at one or both of the S255 and S262 sites[36-38], which

correlate with gap junction internalization and cell division[24,36-39]. To include an appropriate control

condition when testing the CycliCx peptide, we developed the OT peptide based on the sequence

immediately adjacent to CycliCx (Cx43 a.a. 266-280). The OT peptide contains the known Cx43-

MAPK phosphorylation site Cx43 S279 (but not the S282 site), and in line with the CycliCx

peptide, the OT peptide contains a phosphomimetic aspartate at its corresponding residue and

an identical N-terminal stearate group. The OT peptide sequence was selected as a control due

to its proximity to CycliCx and lack of cyclin E binding activity. The OT peptide is similar to the first

15 a.a. of the 18 a.a. TAT-Cx43266-283 peptide published by Gangoso et al., with the exception

of the a.a. 279 serine to aspartate mutation[7,40]. TAT-Cx43266-283 is a Cx43 mimetic based on

the cSRC binding domain. cSRC perpetuates cell proliferation in some cancers, and evidence
suggests binding with Cx43 inactivates cSRC and limits cancer growth. It is thought TAT-
Cx43266-283 binds cSRC directly, mimicking the effect of full-length Cx43 and limiting
glioblastoma growth in culture and *in vivo*[7,40]. In the present study, the OT peptide did not inhibit
SMC proliferation or neointima formation. Due to the similarity between TAT-Cx43266-283 and
our OT peptide, it is possible our OT peptide is lacking critical a.a. for cSRC binding, the aspartate
mutation within the OT peptide prevents cSRC binding, or cSRC does not play a significant role
in injury-induced proliferation in SMC.

Proposed mechanisms of action for previously published Cx43 mimetic peptides include binding
between the Cx43 mimetic and a physically interacting protein, or direct binding between the Cx43
mimetic and Cx43 itself[7,40,41]. Our peptide array data demonstrate sequences for CycliCx and the
cyclin E protein can bind directly. Although we do not present direct evidence of CycliCx-Cx43
interactions, CycliCx induced a shift in Cx43 molecular weight attributable to an increase in Cx43-
pS368, a known PKC phosphorylation site. Cx43-pS368 phosphorylation is associated with
control over gap junction open/closed states and has been observed with other Cx43 mimetic
peptides, a phenomenon we summarized in our previously published review[42]. For example, the
Cx43 mimetic peptide αCT1, developed by the Gourdie lab to disrupt Cx43/ZO-1 interactions,
binds directly to the Cx43 C-terminus, causing an increase in Cx43-PKC phosphorylation[41].
Mutating αCT1 to interact only with ZO-1 and not directly with Cx43 did not produce increases in
Cx43-pS368, indicating direct Cx43 binding is required for αCT1-induced Cx43-pS368 [41]. As
CycliCx induces a similar phosphorylation shift, it is possible CycliCx is also capable of directly
binding the Cx43 protein.

Cyclin E exerts control over the cell cycle via binding with its kinase CDK2. Although the complete
mechanism of action of cyclin E-CDK2 is uncertain, it is known this complex translocates from the
cytoplasm to the cell nucleus to phosphorylate the retinoblastoma protein and exert its effect on
cell division[43-46]. In our previous studies, cyclin E-Cx43 complexes are initially present in the cell
membrane but bind Cx43 and are removed after prolonged PDGF-β treatment, demonstrating
Cx43 can accompany cyclin E-CDK2 during translocation[24]. In the present study, CycliCx
treatment retained Cx43 at the cell membrane after PDGF-β treatment, likely attributable to the
shift toward Cx43-pS368 phosphorylation, which is associated with stabilizing Cx43 at the
membrane[47,48]. Therefore, it is possible Cx43 is required to chaperone cyclin E to the cell nucleus
to progress the cell cycle and induce proliferation, and disrupting the cyclin E-Cx43 complex
interaction and trafficking prevents this.

In line with this, our transcriptomic data suggest CycliCx treatment blunts the PDGF-β-induced
increase in transcript expression of critical cell cycle promoter genes downstream of the cyclin E-
CDK2 complex. Cyclin E-CDK2 primarily regulates the G1/S phase transition of the cell cycle[27].
Indeed, there was little change in the expression of genes involved in the regulation of the early
G1 phase of cell cycle progression with CycliCx treatment, while CycliCx significantly blunted the
PDGF-β-induced regulation of G1/S phase transition and subsequent G2/M phase transition and
M phase genes. This suggests CycliCx targets cyclin E to arrest the cell cycle during the G1/S
phase transition.

In addition to halting pathological SMC proliferation and neointima formation, it is important to
examine potential off-target effects of the CycliCx peptide. We found many PDGF-β downstream
transcription targets unrelated to cell cycle, such as tissue inhibitors of metalloproteases (TIMPs)
and interleukins, were not affected by the CycliCx peptide. Our investigation also revealed a
reduction in gap junction communication following CycliCx treatment. This may be attributable to
the observed increase in Cx43-pS368, which is associated with gap junction closure[49,50].
Interestingly, PDGF-β also induced gene expression of Cx40, but this increase in expression was
reduced after treatment with CycliCx. Together, these data demonstrate CycliCx treatment
specifically inhibits PDGF-β-induced pathological SMC cell cycle progression without significant
disruption to other cell signaling pathways.

A caveat to this study is the use of veins over arteries in testing CycliCx treatment in human
tissues. In modern clinical practice, the internal mammary or radial arteries are preferred for
coronary bypass surgeries over the saphenous vein due to lower levels of neointima formation[51-
53]. Even so, the saphenous vein is still used more often than any other vessel for coronary bypass
when additional bypass conduits are required[54]. As such, testing CycliCx in venous tissue is highly
relevant to surgical practice. Currently, we are unable to trace the half-life of the CycliCx peptide
due to a lack of available antibodies to CycliCx and the mechanism of synthesizing the peptide.
The CycliCx peptide is synthesized with an N-terminal stearate group, limiting our ability to add a
reporter such as GFP to the C-terminus. Attempts to add 6xHis tags to the peptide resulted in
insoluble peptides and were not viable for testing. However, our data show clear effects in vivo in
mice at 14 days, and our in vitro studies show CycliCx effects appear around 3 hours and remain
at >16 hours, suggesting a peptide with good stability, although this remains to be fully elucidated.
Our data also shows CycliCx limits gap junction signaling; thus, we are unable to distinguish gap
junction and connexin-mediated effects directly in this study. However, our previous study found
that Cx43-gap junction signaling did not play a significant role in PDGF-β-induced SMC

proliferation in mice[24]. The CycliCx peptide may therefore also act as a promising tool for the
specific regulation of gap junction signaling, which is not currently available.

In conclusion, we show targeting Cx43 and cyclin E interactions with the novel Cx43 mimetic
peptide CycliCx limits SMC proliferation and neointima formation. The findings from this study
highlight new pathways important in neointima formation, which could be targeted to improve
patency in human revascularization strategies. While our study focuses on pathological
proliferation in SMC in blood vessels, Cx43 has been implicated in the progression of various
proliferative diseases[55]. As our data suggests, CycliCx targets a Cx43-dependent pathological
proliferative mechanism. Future research will also focus on exploring the potential of CycliCx
treatment in other proliferative diseases.

Methods

Sex as a biological variable

Male and female mice were used to assess sex as a biological variable in animal studies. Our data do not show differences in cell pathology or neointima formation in male vs female mice, and the results report combined data for both male and female mice. The sex of the individual mice is indicated in **Supplemental Figure 20**.

Key Resources: Reagents, materials, and antibodies are all listed in **Supplemental Table 1**

***Ex vivo* human saphenous vein model of neointima formation:** We adapted a previously described *ex vivo* model of human saphenous vein neointima formation developed by George et al.[28,29]. At the end of surgical procedures for coronary artery bypass graft, excess, normally discarded saphenous vein tissue was collected. De-identified specimens from Carilion Clinic cardiovascular patients were processed within 1 hr of the end of surgery. Tissues greater than 2 inches in length, free from visible valves and branch regions, were used in the study. Tissues were placed into a container of harvest media (20 mM HEPES-buffered RPMI-1640, 0.225 mg/mL papaverine hydrochloride, 5 µg/ml amphotericin B, 20 IU/mL sodium heparin). Samples were transferred to a wash media (20 mM HEPES Buffered RPMI-1640 with 2 mM L-Glut, 8 µg/mL gentamicin, 1 IU/mL penicillin, and 1 µg/mL streptomycin). Tissues were cut into four consecutive explant sections (corresponding to treatments) and pinned on 6-well plates containing Sylgard 184 elastomer (**Figure 1**). Explants were maintained in a low-serum media (bicarbonate buffered RPMI-1640, 2 mM L-Glut, 8 ug/mL gentamicin, 1 IU/mL penicillin, 1 µg/mL streptomycin, and 2% FBS). The four treatment groups corresponding to four sections were: 1) no treatment; 2) PDGF-β (50 ng/mL, Millipore #01-305); 3) PDGF-β (50 ng/mL) plus OT peptide (30 µM); and 4) PDGF-β (50 ng/mL) plus CycliCx peptide (30 µM). Media were changed each day for 14 days. At the end of the experiments, tissue explants were fixed *in situ* in 4% PFA at 4°C overnight. To preserve integrity, tissues were encapsulated in 3% agarose before tissue processing and paraffin embedding by the FBRI Histology Core. Comparisons were made to PDGF-β treatment or to PDGF-β+OT peptide and PDGF-β+CycliCx treatments for changes in new intima formation.

Cx43 peptide arrays for cyclin E binding: SPOT synthesis of peptides and overlay experiments were carried out as described[56]. Full-length cyclin E-GST protein was synthesized as previously described by us[24]. Cyclin E-GST proteins for arrays were isolated using anti-GST beads, spin concentrated (Amicon, 10K MWCO, Millipore #UFC901008), and buffer exchanged to PBS 125 mM NaCl, pH 6.5 on PD10 columns (Millipore #GE17-0851-01). Peptide arrays were initially bathed in 100% ethanol, equilibrated in PBS-Tween (PBST, 125 mM NaCl, 0.1% tween, pH 6.5)

for 10 min, then blocked in PBST-5% block (PBST with 5% milk) at room temperature for 2 hr.

Arrays were washed in PBST, then incubated with cyclin E proteins (2.86 mg/mL) in PBST-1%

block (1% Milk). Binding on arrays was detected using (cyclin E primary antibodies) in PBST-1%

block, followed by secondary detection using rabbit HRP (Thermo Fisher #31458) in PBS-T-1%

block for 1 hr at room temp, then developed by chemiluminescent substrate (SuperSignal, Thermo

Fisher # 34579) on x-ray film in a dark room. Developed films were scanned for image analysis

(cyclin E, Odyssey, LiCor). Arrays were imaged using UV to provide imaging of total peptides

(Odyssey, LiCor). Sequences for the truncation mutation array can be found in **Supplemental**

List 1.

Peptides: Peptides used in the studies were designed based on the peptide array minimal binding

sequences shown in **Figure 2**. Peptides were synthesized with an N-terminal stearate group to

improve cellular uptake[57]. Peptide final sequences used in the study are CycliCx: stearate-

LDPSKDCGDPKYAY and Off-Target (OT) peptide: stearate-AYFNGCSSPTAPLDP. Peptide

properties are described in **Supplemental Figure 6**. Patent #WO2021021716: "Compositions

and methods for inhibiting neointimal formation" covers the discovery and use of peptide (CycliCx)

and control peptides. Peptides were commercially synthesized (ThermoFisher custom peptide

synthesis services) to >80% purity. Peptides were reconstituted at room temperature in sterile

PBS (for cell culture) or saline (for *in vivo*) to a concentration of 2 mM and stored at -80°C for use.

Peptides used in cell culture and explant culture experiments were prepared to a final

concentration of 30 μM in media. Peptides for *in vivo* applications in mice were prepared to a

concentration of 200 μM in a pluronic gel (35% final concentration), with 50 μL of the pluronic gel

applied around the ligated carotids.

Animals: All mice were cared for under the provisions of the Virginia Tech Animal Care and Use

Committee and followed the National Institute of Health guidelines for the care and use of

laboratory animals. All studies were performed in C57BL6/J mice; strain number #000664,

purchased from Jackson Labs. Mice were bred in-house and used between 12-14 weeks of age.

Carotid ligations and carotid Pluronic gel applications in mice: Permanent 14-day carotid

ligations were used to promote neointima in mice, as we previously described[24]. Mice were,

anesthetized using isoflurane, placed on a surgically sterile heating pad with isoflurane, and the

neck area cleared of hair and sterilized. Mice received analgesic (Ethiqa 0.5 mL) proximal to the

surgical site. An incision was made in the neckline of the mice, carotid exposed, and a 6/0 suture

was secured to the carotid immediately before the bifurcation. In sham control mice, a 6/0 suture

was passed under the exposed carotid and then removed. In ligated and sham mice, carotids

were elevated, and cold Pluronic gel (50 μL) was applied evenly around the carotid. Carotids were

held in place for around 10 seconds until the Pluronic gel solidified (as we previously described)[58].

The surgical site was then closed, and the skin sutured, followed by post-operative recovery.

Comparisons were made to surgical controls (sham) and OT peptide controls.

Pluronic gel preparation and application in mice. A 50% Pluronic gel stock was made by

adding 3 g of Pluronic powder to 3 mL of ice-cold sterile saline and mixing under cold conditions

until dissolved. Pluronic gels were sterilized by autoclaving. For surgeries, Pluronic stocks (50%)

were mixed with sterile saline or with peptides in sterile saline to achieve a final concentration of

35% Pluronic. Samples were maintained on ice to maintain a liquid form until ligations were placed

in the mouse carotid, then 50 μL of Pluronic was applied using a positive displacement pipette to

the surgical site. The 35% Pluronic gels solidify around the carotid at body temperature in mice

prior to closing the surgical site.

Cell culture. Primary human CASMC were sourced from American Type Culture Collection

(ATCC, #PCS-100-02), ThermoFisher (#C0175C) and Lonza (#CC-2583). Initial stocks of cells

were thawed and maintained in vascular smooth muscle cell basal media (ThermoFisher #M231-

500) containing smooth muscle growth serum (SMGS, ThermoFisher # S00725). Cells were

maintained between 70-90% confluency in culture for up to 16 cell divisions. For experiments,

5×10^4 CASMC per mL were plated in each well of a six-well plate. CASMC were serum starved

to stall cell cycle by washing in PBS and then incubating with vascular smooth muscle cell basal

media containing 2% FBS for 72 hr prior to treatments. Media were changed every 48 hours.

Stalled CASMC were induced with PDGF-β (50 ng/mL) and harvested at 16 hr for western blot or

48 hr for flow cytometry of proliferation. For peptide treatments, cells were treated simultaneously

with PDGF-β (50 ng/mL) and peptide (30 μM).

Flow cytometry: For studies of proliferation, CASMC were treated with 5-ethynyl-2'deoxyuridine

(EDU, Thermo Fisher, #C10425) at the same time as PDGF-β and peptide treatments. At the end

of experiments, CASMC were trypsinized, centrifuged (700 rpm, 5 min), then resuspended in 0.5

mL PBS. Cells were fixed by adding 70% ethanol while vortexing. EDU detection was performed

as per the manufacturer's instructions. Flow cytometry data were collected using a BD Accuri C6

flow cytometer, and data analyzed using FlowJo software. Proliferation was calculated as a

percentage of EDU-positive cells to total cells.

Imaging and analysis: Histological imaging was completed on a Nikon Ci-L histology microscope

using tile scan functions to capture whole tissues. Data for the quantification of neointima was

performed by a researcher blinded to sample identification. All immunofluorescent images were

captured using Nikon NIE confocal microscope (A1, with resonance scanning). Single images,

maximum intensity Z-stack projection, along with tile scanning, was used to visualize samples. All

quantitative analysis was performed using NIS Elements Advanced Research software (Nikon).

To analyze new intima in TGLN stained tissues, a line (16pt) was drawn at the luminal edge of

the tissues using NIE software using the signal intensity tool. Background signal was set for

individual images based on tissue autofluorescence, and the area under the curve (AUC) for the

TGLN signal was divided by the length of the tissue. 5-6 images were taken across each tissue,

and the AUC was averaged for these values. For each HSV experiment, consecutive samples

are used, and a fold change to the individual patient NT sample TGLN signal was made to show

changes.

Immunofluorescence: Tissues fixed in 4% PFA were embedded in paraffin and sectioned.

Antigen retrieval was performed (Universal R, antigen retrieval solution, Electron Microscopy

Solutions) on deparaffined slides containing tissue sections and tissues blocked in a solution of

PBS, 0.1% NP-40 substitute, 1% BSA, where the appropriate blocking solution contained anti-

goat serum. Primary antibody incubations were performed overnight at 4°C, samples washed,

and secondary antibodies applied.

Scrape load dye transfer (SLDT) assays: CASMC were grown as described in wound healing.

CASMC were treated with PDGF-β and peptides for either 6 hr or 16 hr. Cells were washed in

PBS with calcium and magnesium, and scratch wounds were created using a #10 scalpel. Cells

were then incubated with the gap junction permeable Lucifer Yellow (457 mw, final concentration

0.05%, Thermo #2790937) and gap junction impermeable Rhodamine Dextran 647 (10,000 mw,

final concentration 100 µg/mL ThermoFisher #2860812) for 5 min. Cells were then washed in

PBS three times, fixed in 4% PFA for 20 min at room temperature, washed in PBS, and imaged

for dye transfer (Nikon A1 confocal). Analysis of dye transfer area for each condition was

performed using NI Elements Advanced Research (**Supplemental Figure 12**). Normalized data

(to length) was presented as Area/Length of dye transfer.

Western blotting and membrane protein biotinylation and coimmunoprecipitation: Western

blotting, cell surface biotinylation, and co-immunoprecipitation studies were performed as we

previously described[24,59]. Proteins for western blotting and cell surface protein biotinylation were

harvested in PBS lysis buffer containing 5 mM EDTA, 0.5% sodium deoxycholate, 0.5% NP-40

substitute, 1 mM sodium orthovanadate, 10 mM sodium fluoride protease inhibitor cocktail (1:100,

Millipore #P8340), phosphatase inhibitor cocktail 2 (1:100, Millipore #P5726), phosphatase

inhibitor cocktail 3 (1:100, Sigma #P0044). For co-immunoprecipitation, 10 mM n-

ethylmethylamine (Millipore, #291145) was added to the lysis buffer. Protein samples were

quantified by BCA assay before loading, and equal loading was confirmed using β-tubulin and

total protein assay (Total protein stain, LiCor). Membranes were developed using LiCor secondary

antibodies, and imaged on a LiCor Odyssey scanner. Expression analysis was performed using

Image Studio (LiCor). Values were normalized to total protein stain or β-tubulin, and changes

were calculated as a fold change compared to non-treated controls.

Bulk RNA sequencing. CASMC were plated, stalled for 72 hr and treated with PDGF-β and

peptides overnight. Human CASMC were washed 2x in ice-cold PBS and then lysed in 350 µl of

RLT lysis buffer. RNA was harvested (RNeasy Micro Kit, Qiagen #74004) according to the

manufacturer's instructions, including the optional DNA removal step. Extracted RNA was

quantified using a NanoDrop One spectrophotometer (Thermo Fisher, USA). Aliquots of RNA (1.5

µg) were then diluted to equal concentrations (75 ng/µl) in RNase-free water.

RNA integrity analysis, library preparation, sequencing, raw read count generation, and

differential gene expression analysis were performed commercially by Novogene (Sacremento,

CA, USA) using their Human mRNA Sequencing pipeline. Full parameters, protocols, and quality

control reports are available on request, but are explained briefly below:

RNA integrity was assessed using the 5400 Fragment Analyzer System (Agilent Technologies,

CA, USA). mRNA library preparation was performed with poly A enrichment and library quality

was verified using Qubit assay for quantification, Agilent Bioanalyzer for size distribution, and

qPCR for molarity using adaptor-specific primers. Sequencing was performed on the NovaSeq X

Plus (Illumina, USA) platform with 150 bp paired end reads and a minimum of 6 G raw data. Raw

sequencing data in FASTQ format were cleaned using custom in-house (Novogene) Perl scripts.

Reads with adapter contamination, reads containing more than 10% uncertain nucleotides, and

reads in which more than 50% of bases had a quality (Q) score below 5 were removed. At this

point, an outlying sample in the PDGF-β group was removed from all further analysis due to

significant AT/GC content separation (**Supplemental Figure 15).** Reads were then aligned to the

human genome using 'HISAT2' (v2.0.05). Raw read counts were generated using 'featureCounts'

(v1.5.0-p3). Fragments Per Kilobase of transcript per Million mapped reads (FPKM) was derived

from raw data by Novogene and used to calculate Pearson's correlation coefficient to assess intra

group and between group variability, and hierarchical clustering. However, for all downstream

analysis raw count normalization was performed using 'DESeq2' (v1.44.0) in the R statistical

analysis program (v4.4.1). Gene symbols were extracted based on provided Ensembl ID using

'clusterProfiler' (v4.12.0) and 'org.Hs.eg.db' (v3.19.1). 'DESeq2' normalized counts were $\log_2$

transformed to ensure a normal distribution. Euclidean distances were then calculated and
principal component analysis (PCA) was performed using 'dist()' and 'prcomp()', respectively, in
base R.

Differential gene expression analysis was performed by Novogene for every possible combination
of sample groups using 'DESeq2' (v.1.20.0) with Benjamini-Hochberg multiple comparison
adjustment. An adjusted p-value (padj) < 0.05 and an $|log_2(FC)| > 0$ was set as the threshold for
differentially expressed genes (DEGs). Volcano plots displaying $-log_{10}(padj)$ and $log_2(FC)$ were
generated in R.

Pathway enrichment analysis was performed on DEGs with a padj < 0.05 with 'clusterProfiler' in
R using the Gene Ontology (GO) database. GO terms with a padj < 0.05 following Benjamini-
Hochberg multiple comparison adjustment were considered statistically significantly enriched.
Separate pathway enrichment analysis was performed for upregulated DEGs ($log_2(FC) > 0$) and
downregulated DEGs ($log_2(FC) < 0$). Individual pathway heatmaps were generated by extracting
'DESeq2' normalized counts for all DEGs (when comparing PDGF-β+CycliCx to PDGF-β+OT
peptide) in indicated statistically significantly enriched pathways and calculating a gene
expression z-score for each sample. Data were then visualized using 'pheatmap' (v1.0.12).

To interrogate changes in expression of single genes, 'DESeq2' normalized counts were
extracted, and the mean normalized counts were plotted in GraphPad Prism 10. A 1-way ANOVA
followed by Tukey's multiple comparison testing was performed to assess changes in gene
expression between groups. Unless otherwise stated, RNA sequencing data were visualized
using 'ggplot2' (v3.5.1).

Statistical analysis

Statistical analysis was performed using Prism version 10.4.2. Samples were assessed for normal
distribution using a Shapiro-Wilk test, and outliers were identified using the Prism robust
regression and outlier removal (ROUT) test (Q =1%). 1-way or 2-way multiple comparison ANOVA
followed by Tukey or Dunnett post-test was used for comparisons between >2 groups. The t-test
was used for comparisons of 2 treatment groups. Kruskal-Wallis and Mann-Whitney non-
parametric tests were used for groups that did not display normal distribution. A minimum of n=3
was used for all statistical analyses, and n values for each experiment are stated in the figure
legends. P values are shown for all statistical analyses, with a P value of less than 0.05 considered
significant. Data represent mean ± SD.

Approval for the use of human excess vascular tissues. The Institutional Review Board (IRB) at Carilion Roanoke Memorial Hospital reviewed and approved the use of de-identified human saphenous vein tissues normally discarded at the end of coronary artery bypass graft surgeries. The Carilion Clinic IRB determined the research does not meet the definition of human subject research as outlined in 45 CFR 46.102(d). Patient data, health data, and identifiable information were not included with specimens.

Data availability

All data generated and used for analysis and original data sets are freely available for further analysis

Author contributions

MWS, MCR, MJ, GSB, BEI, and SRJ designed research studies; MWS, MCR, XL, FI, MRL, KR, RCJ, AM, CLD, AKB, MJ, GSB, and SRJ conducted experiments and acquired data; MWS, MCR, XL, KR, RJ, AM, CLD, GSB, BEI and SRJ analyzed data; PDL provided reagents; MWS, MCR, BEI and SRJ wrote the manuscript; All authors approved the final version of the manuscript. A coin flip decided the listed order of co-first authors MWS and MCR; both authors contributed equally.

Acknowledgments

We gratefully acknowledge the resources and services provided by Dr. Clare Dennison, at the FBRI Histology Core Facility at VTC, and Ms Ruth Macleod at the University of Glasgow for the production of peptide arrays. Servier medical art, licensed under CC BY 4.0 https://creativecommons.org/licenses/by/4.0/, was used to create figure schematics. Funding for this research includes AHA 19CDA34630036 (SJ), AHA 23PRE1010870 (MS), AHA 25POST1410066 (MR), NIH-F31HL170721 (MS), NIH-R215R21HL168614-02 (SJ), VT-Proof of Concept (SJ), and the Seale Innovation Award (SJ). Medical Research Council grant MR/y00364/1 (GSB). NIH HL120840, NIH HL137112.

References

1. Nedkoff L, Briffa T, Zemedikun D, Herrington S, Wright FL. Global Trends in Atherosclerotic Cardiovascular Disease. *Clin Ther*. Nov 2023;45(11):1087-1091. doi:10.1016/j.clinthera.2023.09.020

2. Maron DJ, Hochman JS, Reynolds HR, et al. Initial Invasive or Conservative Strategy for Stable Coronary Disease. *N Engl J Med*. Apr 09 2020;382(15):1395-1407. doi:10.1056/NEJMoa1915922

3. Déglise S, Bechelli C, Allagnat F. Vascular smooth muscle cells in intimal hyperplasia, an update. *Front Physiol*. 2022;13:1081881. doi:10.3389/fphys.2022.1081881

4. Cornelissen A, Guo L, Fernandez R, et al. Endothelial Recovery in Bare Metal Stents and Drug-Eluting Stents on a Single-Cell Level. *Arterioscler Thromb Vasc Biol*. Aug 2021;41(8):2277-2292. doi:10.1161/ATVBAHA.121.316472

5. Bønaa KH, Mannsverk J, Wiseth R, et al. Drug-Eluting or Bare-Metal Stents for Coronary Artery Disease. *N Engl J Med*. Sep 29 2016;375(13):1242-52. doi:10.1056/NEJMoa1607991

6. Goodenough DA, Goliger JA, Paul DL. Connexins, connexons, and intercellular communication. *Annu Rev Biochem*. 1996;65:475-502. doi:10.1146/annurev.bi.65.070196.002355

7. Gangoso E, Thirant C, Chneiweiss H, Medina JM, Tabernero A. A cell-penetrating peptide based on the interaction between c-Src and connexin43 reverses glioma stem cell phenotype. *Cell Death Dis*. Jan 23 2014;5:e1023. doi:10.1038/cddis.2013.560

8. Moorer MC, Hebert C, Tomlinson RE, Iyer SR, Chason M, Stains JP. Defective signaling, osteoblastogenesis and bone remodeling in a mouse model of connexin 43 C-terminal truncation. *J Cell Sci*. Feb 01 2017;130(3):531-540. doi:10.1242/jcs.197285

9. Langlois S, Cowan KN, Shao Q, Cowan BJ, Laird DW. The tumor-suppressive function of Connexin43 in keratinocytes is mediated in part via interaction with caveolin-1. *Cancer Res*. May 15 2010;70(10):4222-32. doi:10.1158/0008-5472.CAN-09-3281

10. Gellhaus A, Dong X, Propson S, et al. Connexin43 interacts with NOV: a possible mechanism for negative regulation of cell growth in choriocarcinoma cells. *J Biol Chem*. Aug 27 2004;279(35):36931-42. doi:10.1074/jbc.M404073200

11. Fu CT, Bechberger JF, Ozog MA, Perbal B, Naus CC. CCN3 (NOV) interacts with connexin43 in C6 glioma cells: possible mechanism of connexin-mediated growth suppression. *J Biol Chem*. Aug 27 2004;279(35):36943-50. doi:10.1074/jbc.M403952200

12. Strauss RE, Mezache L, Veeraraghavan R, Gourdie RG. The Cx43 Carboxyl-Terminal Mimetic Peptide αCT1 Protects Endothelial Barrier Function in a ZO1 Binding-Competent Manner. *Biomolecules*. Aug 12 2021;11(8)doi:10.3390/biom11081192

13. Smyth JW, Guo S, O'Rourke L, et al. Increased interaction between connexin43 and microtubules is critical for glioblastoma stem-like cell maintenance and tumorigenicity. *bioRxiv*. Jan 27 2024;doi:10.1101/2024.01.26.576347

14. Hong T, Hill CE. Restricted expression of the gap junctional protein connexin 43 in the arterial system of the rat. *J Anat*. May 1998;192 (Pt 4)(Pt 4):583-93. doi:10.1046/j.1469-7580.1998.19240583.x

15. Liao Y, Regan CP, Manabe I, et al. Smooth muscle-targeted knockout of connexin43 enhances neointimal formation in response to vascular injury. *Arterioscler Thromb Vasc Biol*. May 2007;27(5):1037-42. doi:10.1161/ATVBAHA.106.137182

16. Chadjichristos CE, Matter CM, Roth I, et al. Reduced connexin43 expression limits neointima formation after balloon distension injury in hypercholesterolemic mice. *Circulation*. Jun 20 2006;113(24):2835-43. doi:10.1161/CIRCULATIONAHA.106.627703

17. Zhang A, Han Y, Wang B, Li S, Gan W. Beyond Gap Junction Channel Function: the Expression of Cx43 Contributes to Aldosterone-Induced Mesangial Cell Proliferation via the ERK1/2 and PKC Pathways. *Cell Physiol Biochem*. 2015;36(3):1210-22. doi:10.1159/000430291

18.   Tittarelli A, Guerrero I, Tempio F, et al. Overexpression of connexin 43 reduces melanoma
proliferative and metastatic capacity. *Br J Cancer*. Jul 14 2015;113(2):259-67.
doi:10.1038/bjc.2015.162

19.   Arshad M, Conzelmann C, Riaz MA, Noll T, Gündüz D. Inhibition of Cx43 attenuates
ERK1/2 activation, enhances the expression of Cav-1 and suppresses cell proliferation. *Int J Mol*
Med. Nov 2018;42(5):2811-2818. doi:10.3892/ijmm.2018.3828

20.   Chadjichristos CE, Morel S, Derouette JP, et al. Targeting connexin 43 prevents platelet-
derived growth factor-BB-induced phenotypic change in porcine coronary artery smooth muscle
cells. *Circ Res*. Mar 28 2008;102(6):653-60. doi:10.1161/CIRCRESAHA.107.170472

21.   Plenz G, Ko YS, Yeh HI, et al. Upregulation of connexin43 gap junctions between
neointimal smooth muscle cells. *Eur J Cell Biol*. Oct 2004;83(10):521-30. doi:10.1078/0171-9335-
00417

22.   Yeh HI, Lupu F, Dupont E, Severs NJ. Upregulation of connexin43 gap junctions between
smooth muscle cells after balloon catheter injury in the rat carotid artery. *Arterioscler Thromb*
Vasc Biol. Nov 1997;17(11):3174-84. doi:10.1161/01.atv.17.11.3174

23.   Sorgen PL, Duffy HS, Sahoo P, Coombs W, Delmar M, Spray DC. Structural changes in
the carboxyl terminus of the gap junction protein connexin43 indicates signaling between binding
domains for c-Src and zonula occludens-1. *J Biol Chem*. Dec 24 2004;279(52):54695-701.
doi:10.1074/jbc.M409552200

24.   Johnstone SR, Kroncke BM, Straub AC, et al. MAPK phosphorylation of connexin 43
promotes binding of cyclin E and smooth muscle cell proliferation. *Circ Res*. Jul 6
2012;111(2):201-11. doi:10.1161/circresaha.112.272302

25.   González-Sánchez A, Jaraíz-Rodríguez M, Domínguez-Prieto M, Herrero-González S,
Medina JM, Tabernero A. Connexin43 recruits PTEN and Csk to inhibit c-Src activity in glioma
cells and astrocytes. *Oncotarget*. Aug 02 2016;7(31):49819-49833.
doi:10.18632/oncotarget.10454

26.   Solan JL, Marquez-Rosado L, Sorgen PL, Thornton PJ, Gafken PR, Lampe PD.
Phosphorylation at S365 is a gatekeeper event that changes the structure of Cx43 and prevents
down-regulation by PKC. *J Cell Biol*. Dec 17 2007;179(6):1301-9. doi:10.1083/jcb.200707060

27.   Ohtsubo M, Theodoras AM, Schumacher J, Roberts JM, Pagano M. Human cyclin E, a
nuclear protein essential for the G1-to-S phase transition. *Mol Cell Biol*. May 1995;15(5):2612-24.
doi:10.1128/MCB.15.5.2612

28.   George SJ, Wan S, Hu J, MacDonald R, Johnson JL, Baker AH. Sustained reduction of
vein graft neointima formation by ex vivo TIMP-3 gene therapy. *Circulation*. Sep 13 2011;124(11
Suppl):S135-42. doi:10.1161/CIRCULATIONAHA.110.012732

29.   George SJ, Williams A, Newby AC. An essential role for platelet-derived growth factor in
neointima formation in human saphenous vein in vitro. *Atherosclerosis*. Feb 1996;120(1-2):227-
40. doi:10.1016/0021-9150(95)05717-x

30.   Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. *Cell*. Mar
09 2017;168(6):960-976. doi:10.1016/j.cell.2017.02.004

31.   Sedovy MW, Leng X, Iqbal F, et al. Preserving Endothelial Integrity in Human Saphenous
Veins during Preparation for Coronary Bypass Surgery. *J Vasc Res*. Jan 12 2024:1-9.
doi:10.1159/000535843

32.   O'Sullivan M, Scott SD, McCarthy N, et al. Differential cyclin E expression in human in-
stent stenosis smooth muscle cells identifies targets for selective anti-restenosis therapy.
Cardiovasc Res. Dec 01 2003;60(3):673-83. doi:10.1016/j.cardiores.2003.09.018

33.   Rhett JM, Calder BW, Fann SA, Bainbridge H, Gourdie RG, Yost MJ. Mechanism of action
of the anti-inflammatory connexin43 mimetic peptide JM2. *Am J Physiol Cell Physiol*. Sep 01
2017;313(3):C314-C326. doi:10.1152/ajpcell.00229.2016

34.   Calder BW, Matthew Rhett J, Bainbridge H, Fann SA, Gourdie RG, Yost MJ. Inhibition of
connexin 43 hemichannel-mediated ATP release attenuates early inflammation during the foreign
body response. *Tissue Eng Part A*. Jun 2015;21(11-12):1752-62. doi:10.1089/ten.TEA.2014.0651
35.   He D, Ma Z, Xue K, Li H. Juxtamembrane 2 mimic peptide competitively inhibits
mitochondrial trafficking and activates ROS-mediated apoptosis pathway to exert anti-tumor
effects. *Cell Death Dis*. Mar 24 2022;13(3):264. doi:10.1038/s41419-022-04639-6
36.   Lampe PD, Kurata WE, Warn-Cramer BJ, Lau AF. Formation of a distinct connexin43
phosphoisoform in mitotic cells is dependent upon p34cdc2 kinase. *J Cell Sci*. Mar 1998;111 (Pt
6):833-41. doi:10.1242/jcs.111.6.833
37.   Kanemitsu MY, Jiang W, Eckhart W. Cdc2-mediated phosphorylation of the gap junction
protein, connexin43, during mitosis. *Cell Growth Differ*. Jan 1998;9(1):13-21.
38.   Cameron SJ, Malik S, Akaike M, et al. Regulation of epidermal growth factor-induced
connexin 43 gap junction communication by big mitogen-activated protein kinase1/ERK5 but not
ERK1/2   kinase   activation.   *J   Biol   Chem*.   May   16   2003;278(20):18682-8.
doi:10.1074/jbc.M213283200
39.   Doble BW, Dang X, Ping P, et al. Phosphorylation of serine 262 in the gap junction protein
connexin-43 regulates DNA synthesis in cell-cell contact forming cardiomyocytes. *J Cell Sci*. Jan
26 2004;117(Pt 3):507-14. doi:10.1242/jcs.00889
40.   Jaraíz-Rodríguez M, Talaverón R, García-Vicente L, et al. Connexin43 peptide, TAT-
Cx43266-283, selectively targets glioma cells, impairs malignant growth, and enhances survival
in mouse models in vivo. *Neuro Oncol*. Apr 15 2020;22(4):493-504. doi:10.1093/neuonc/noz243
41.   Jiang J, Hoagland D, Palatinus JA, et al. Interaction of α Carboxyl Terminus 1 Peptide
With the Connexin 43 Carboxyl Terminus Preserves Left Ventricular Function After Ischemia-
Reperfusion   Injury.   *J   Am   Heart   Assoc*.   Aug   20   2019;8(16):e012385.
doi:10.1161/JAHA.119.012385
42.   King DR, Sedovy MW, Leng X, et al. Mechanisms of Connexin Regulating Peptides.
International Journal of Molecular Sciences. 2021;22(19)doi:10.3390/ijms
43.   Jackman M, Kubota Y, den Elzen N, Hagting A, Pines J. Cyclin A- and cyclin E-Cdk
complexes shuttle between the nucleus and the cytoplasm. *Mol Biol Cell*. Mar 2002;13(3):1030-
45. doi:10.1091/mbc.01-07-0361
44.   Moore JD, Yang J, Truant R, Kornbluth S. Nuclear import of Cdk/cyclin complexes:
identification of distinct mechanisms for import of Cdk2/cyclin E and Cdc2/cyclin B1. *J Cell Biol*.
Jan 25 1999;144(2):213-24. doi:10.1083/jcb.144.2.213
45.   Gaulin JF, Fiset A, Fortier S, Faure RL. Characterization of Cdk2-cyclin E complexes in
plasma membrane and endosomes of liver parenchyma. Insulin-dependent regulation. *J Biol
Chem*. Jun 02 2000;275(22):16658-65. doi:10.1074/jbc.275.22.16658
46.   Akiyama T, Ohuchi T, Sumida S, Matsumoto K, Toyoshima K. Phosphorylation of the
retinoblastoma protein by cdk2. *Proc Natl Acad Sci U S A*. Sep 01 1992;89(17):7900-4.
doi:10.1073/pnas.89.17.7900
47.   O'Quinn MP, Palatinus JA, Harris BS, Hewett KW, Gourdie RG. A peptide mimetic of the
connexin43 carboxyl terminus reduces gap junction remodeling and induced arrhythmia following
ventricular injury. Research Support, N.I.H., Extramural Research Support, Non-U.S. Gov't.
Circulation Research. Mar 18 2011;108(6):704-15. doi:10.1161/circresaha.110.235747
48.   Rhett JM, Jourdan J, Gourdie RG. Connexin 43 connexon to gap junction transition is
regulated by zonula occludens-1. *Mol Biol Cell*. May 2011;22(9):1516-28. doi:10.1091/mbc.E10-
06-0548
49.   Ek-Vitorin JF, King TJ, Heyman NS, Lampe PD, Burt JM. Selectivity of connexin 43
channels is regulated through protein kinase C-dependent phosphorylation. *Circ Res*. Jun 23
2006;98(12):1498-505. doi:10.1161/01.RES.0000227572.45891.2c

50. Lampe PD, TenBroek EM, Burt JM, Kurata WE, Johnson RG, Lau AF. Phosphorylation of
connexin43 on serine368 by protein kinase C regulates gap junctional communication. *J Cell Biol*.
Jun 26 2000;149(7):1503-12. doi:10.1083/jcb.149.7.1503
51. Gaudino M, Benedetto U, Fremes S, et al. Radial-Artery or Saphenous-Vein Grafts in
Coronary-Artery Bypass Surgery. *N Engl J Med*. May 31 2018;378(22):2069-2077.
doi:10.1056/NEJMoa1716026
52. van der Meer J, Hillege HL, van Gilst WH, et al. A comparison of internal mammary artery
and saphenous vein grafts after coronary artery bypass surgery. No difference in 1-year occlusion
rates and clinical outcome. CABADAS Research Group of the Interuniversity Cardiology Institute
of The Netherlands. *Circulation*. Nov 1994;90(5):2367-74. doi:10.1161/01.cir.90.5.2367
53. Tector AJ, Schmahl TM, Janson B, Kallies JR, Johnson G. The internal mammary artery
graft. Its longevity after coronary bypass. *JAMA*. Nov 13 1981;246(19):2181-3.
54. Gaudino M, Bakaeen FG, Sandner S, et al. Expert Systematic Review on the Choice of
Conduits for Coronary Artery Bypass Grafting: Endorsed by the European Association for Cardio-
Thoracic Surgery (EACTS) and The Society of Thoracic Surgeons (STS). *Ann Thorac Surg*. Oct
2023;116(4):659-674. doi:10.1016/j.athoracsur.2023.06.010
55. Bonacquisti EE, Nguyen J. Connexin 43 (Cx43) in cancer: Implications for therapeutic
approaches via gap junctions. *Cancer Lett*. Feb 01 2019;442:439-444.
doi:10.1016/j.canlet.2018.10.043
56. Bolger GB, Baillie GS, Li X, et al. Scanning peptide array analyses identify overlapping
binding sites for the signalling scaffold proteins, beta-arrestin and RACK1, in cAMP-specific
phosphodiesterase PDE4D5. *Biochem J*. Aug 15 2006;398(1):23-36. doi:10.1042/BJ20060423
57. Futaki S, Ohashi W, Suzuki T, et al. Stearylated arginine-rich peptides: a new class of
transfection systems. *Bioconjug Chem*. Nov-Dec 2001;12(6):1005-11. doi:10.1021/bc015508l
58. Johnstone SR, Ross J, Rizzo MJ, et al. Oxidized phospholipid species promote in vivo
differential cx43 phosphorylation and vascular smooth muscle cell proliferation. *Am J Pathol*. Aug
2009;175(2):916-24. doi:ajpath.2009.090160 [pii]
10.2353/ajpath.2009.090160
59. Yang Y, Delalio LJ, Best AK, et al. Endothelial Pannexin 1 Channels Control Inflammation
by Regulating Intracellular Calcium. *J Immunol*. Jun 2020;204(11):2995-3007.
doi:10.4049/jimmunol.1901089

Figure Legends:

Figure 1: PDGF-β induces new intima in human saphenous veins in vitro, associated with an increase in Cx43 and proliferation markers. Schematic of human saphenous vein tissue harvesting. Excess saphenous veins, used for coronary artery bypass graft, were dissected open with the luminal side up, cut free from valves and sutured vessels, and sequential sections were plated for testing. Media containing 2% serum was compared with samples treated with 50 ng PDGF, which was changed daily (**A**). Tissues were sectioned and stained for H&E, Tgln (orange), Cx43 (magenta), Cx43pS279 (red), and cyclin E (white) (**B**). n=3-5 samples used for staining. * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 μM.

Figure 2: Design of the CycliCx peptide using peptide arrays demonstrates cyclin E binding regions on the phosphorylated Cx43CT. Schematic showing the design for peptide arrays for using a sequence corresponding to the 180 a.a. Cx43-CT (**A**). Sequential 25 a.a. sequences were printed to nitrocellulose; consecutive sequences were overlapped by 5 a.a. (**A**). Eluted fractions from GST columns for cyclin E-GST proteins expressed in bacteria, expression shown by western blotting (anti-cyclin E, upper blot) and coomassie staining (lower gel, **B**). Peptide arrays with Cx43CT^{WT} (red box) or MAPK phosphorylated Cx43-S255/262/279/282 (MK4) substitutions with either alanine (Cx43^{MK4A}, blue), aspartate (Cx43^{MK4D}, green), or native phosphor-serine (Cx43^{MK4pS}, orange) were bound by cyclin E proteins and detect with anti-cyclin E (**C**). Potential binding sites highlighted by red circles (**C**) and cyclin E arrays aligned with UV imaging of total peptide (**D**). Three potential binding sequence regions in Cx43^{MK4D} and Cx43^{MK4pS} are underlined (**E**). Analysis of the first three 25 a.a. binding sequences in Cx43 CT spanning Cx43 a.a. 236-300 shows phospho-specific binding only within Cx43 a.a. 236-285 (**F**), locations shown in schematic (**G**). Truncation mutation peptide array across Cx43 a.a. 255-272 used to define the minimum binding region (**H**). A schematic showing the location of the binding region and selected adjacent region used as an Off-Target (OT) peptide (**I**). The final design of the CycliCx peptide incorporating an N-terminal stearyl group (**J**)

Figure 3: CycliCx Peptide alters Cx43 expression and interactions with cyclin E.

CASMC treated with PDGF (PD, 50 ng/ mL) alone or in combination with OT peptide or CycliCx (30 μM) for 16 hr. Western blots were performed for total-ERK (tERK), phosphorylated-ERK (pERK phosphor ERK Thr202/Tyr204), and ratio of pERK: pERK calculated (n=6) (**A**), for total Cx43 and PKC-phosphorylated Cx43 (Cx43-PKC, pS368) (n=6), and alterations in expression from not treated (NT) calculated (**B**). Distribution and frequency of Cx43 GJ punctates (magenta) in treated CASMC (TGLN, Green) were captured by confocal microscopy, and differences in gap junction punctates/ cell were measured using NIS-elements software (n=3) (**C**). Cell plasma membrane protein biotinylation western blots for membrane-bound Cx43 (streptavidin eluted) compared to cytosolic (non-biotin bound) proteins (n=3) (**D**). Cell-to-cell communication through gap junctions was assessed by lucifer yellow dye transfer (green) in scrape-wounded monolayers of CASMC after treatment with PDGF or PDGF plus OT/ CycliCx peptide (n=6). Scrape load dye transfer (SLDT) of lucifer yellow (green) permeability was used to measure gap junction dye transfer, area of dye normalized to length, with co-loading of high molecular weight dextran (magenta) used to determine non-gap junction dye area (n=6) (**E**) Co-immunoprecipitation western blots were performed to assess Cx43-cyclin E binding in CASMC following peptide

treatment (IP cyclin E, WB Cx43) (n=4), with loading control western blot for Cx43 or cyclin E proteins (**F**). Sample loading for western blots was normalized to total protein or β-tubulin expression, and comparisons were made to NT cells (**A, B, D, F**). One-way ANOVA with multiple comparison test performed, P<0.05 is significant.

Figure 4: Transcriptomics reveals that the CycliCx peptide arrests cell cycle progression at G1/S phase transition. 5-ethynyl-2'-deoxyuridine (EDU) proliferation assay in human coronary artery smooth muscle cells (CASMC; **A**). Schematic showing the design for bulk RNA sequencing and group combinations for analysis (**B**). Principal component analysis (PCA) of 'DESeq2' normalized and $\log_2$ transformed counts (**C**). Volcano plots of 'DESeq2' differentially expressed genes (DEGs) identified with $|\log_2(Fold Change (FC))| > 0$ and an adjusted p-value (padj) < 0.05 for indicated group comparisons (**D**). Venn diagram of the total number of identified DEGs unique to and shared by the PDGF-β (PD) + Off Target Peptide (OT) and PD + CycliCx peptide treatment groups when compared to PD treatment alone (**E**). Top 10 pathways by padj from Gene Ontology (GO) Biological Process pathway analysis ranked by gene ratio (left panel shows top 10 upregulated pathways in PD vs no treatment (NT) and right panel shows top 10 downregulated pathways in PD + CycliCx vs PD + OT; **F**). Heatmap of between replicate 'DESeq2' normalized gene expression z-scores for PD + CycliCx vs PD + OT DEGs from the significantly enriched GO pathway GO:0000082: 'cell cycle G1/S phase transition (**G**). 'DESeq2' normalized gene counts in selected genes (**H**). N = 3-4 per group. Data displayed as mean ± SEM. P-values calculated by Tukey's multiple comparison testing after one-way ANOVA.

Figure 5: CycliCx peptide limits neointima formation in mouse and human blood vessels

H&E staining of mouse carotids following 14 days of sham surgery (no ligation) or ligation with either pluronic gel, pluronic+OT peptide or pluronic+CycliCx pep (200 µM). Measure of media area, neointima area, and ratio of neointima to media. Kruskal-Wallis multiple comparison test, N=12 (9 male, 3 female). H&E staining of human saphenous veins treated with PDGF+OT pep or PDGF+CycliCx for 14 days. Tgln staining of human saphenous veins treated with PDGF+OT pep or PDGF+CycliCx for 14 days. T-test, n=5.

Supplemental Figure Legends:

Supplemental Figure 1: Human saphenous vein H&E staining comparison. Representative images of H&E-stained human saphenous veins for no treatment (NT), PDGF, PDGF+ OT peptide, and PDGF+ CycliCx. * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 µM. N=5.

Supplemental Figure 2: Human saphenous vein smooth muscle cell transgelin staining. Representative images of anti-TGLN stained human saphenous veins for no treatment (NT), PDGF, PDGF+ OT peptide, PDGF+ CycliCx. Measurement of the leading edge (16pt) TGLN staining, averaged n=5-6 measures per tissue. * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 µM. N=5. Mann-Whitney tests were used for NT:PDGF, PD+OT peptide:PDGF+CycliCx, and Kruskal-Wallis multiple comparison test of all values. Exact P values shown.

Supplemental Figure 3: Human saphenous vein Cx43 staining comparison. Representative images of anti-Cx43 stained human saphenous veins for no treatment (NT), PDGF, PDGF+ OT peptide, and PDGF+ CycliCx. * indicates luminal side of the

vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 µM. N=3.

Supplemental Figure 4: Human saphenous vein Cx43-pS279 staining comparison. Representative images of anti-Cx43-pS279 stained human saphenous veins for no treatment (NT), PDGF, PDGF+ OT peptide, and PDGF+ CycliCx. * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 µM. N=3.

Supplemental Figure 5: Human saphenous vein cyclin E staining comparison. Representative images of anti-cyclin E1-stained human saphenous veins for no treatment (NT), PDGF, PDGF+ OT peptide, and PDGF+ CycliCx. * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 50 µM. N=3.

Supplemental Figure 6: Identification of active cyclin E binding sites on Cx43.

In **A**, Single Alanine amino acid substitutions for Cx43 sequences Seq1 Cx43 a.a. 235-260 (white), Seq 2 Cx43 a.a. 254-279 (green), Seq 3 Cx43 a.a. 276-301 (black). Full length sequences contain either native phosphoserine (ps) or aspartate (D). The first four spots contain a full sequence with either serines (S), alanines (A), pS, or D, and were used as representative images for **Figure 2**. In **A**, arrays were imaged with UV (total peptides, left), then probed with anti-cyclin E, and detected with a secondary antibody before development by chemiluminescence onto x-ray film (right). In **B**, truncation mutation arrays were performed on identified sequences (see **supplemental list** for sequences), detected with anti-cyclin E antibodies (top), and imaged with UV and Coomassie (total peptides). In **C**, alignments show the minimum binding sites identified across the Cx43-MAPK binding region (top, Sequence 2) and at a distant site (bottom, Sequence 1). Flow cytometry of stearate-linked peptides targeting Sequence 1 and Sequence 2 were generated and incubated with SMC (30 µM) in the presence of PDGF and EDU for proliferation (**D**). Sequence 1 reduced cell numbers, indicating cell death. Final design of peptide sequences based on binding characteristics seen by truncation arrays for CycliCx peptide (binding), and Off-Target peptide (OT peptide, non-binding), approximate charge, and molecular weight not including stearate group (**E**).

Supplemental Figure 7: Full blots for ERK measurements. CASMC were treated with PDGF-β (50 ng/ mL) alone or in combination with OT peptide or CycliCx (30 µM) for 16 hr. Western blots were performed for total ERK (tERK, p42/p44), phosphorylated ERK (pERK phosphor- p42/p44 Thr202/Tyr204) and ratio of pERK: pERK calculated (n=6). Total protein blots highlight equal loading of samples. Sample loading was normalized to total protein expression, and differences in expression were measured against non-treated cells. One-way ANOVA with multiple comparison test was performed, $P < 0.05$ is significant. Orange box indicates blots used in **Figure 3**.

Supplemental Figure 8: Full blots for Cx43 measurements: CASMC were treated with PDG-β (50 ng/ mL) alone or in combination with OT peptide or CycliCx (30 µM) for 16 hr. Western blots were performed for Total Cx43 and Cx43 (n=3), Cx43-PKC Cx43-pS368) (n=6). Sample loading was normalized to total protein expression, and differences in expression measured against NT cells. One way ANOVA with multiple comparison test performed, $P < 0.05$ is significant. Orange box indicates blots used in **Figure 3**.

Supplemental Figure 9: CycliCx alters Cx43 phosphorylation expression from 3 hours post-treatment. CASMC were treated with CycliCx (30 µM) and harvested every hour up until 5 hr. Expression of total Cx43 and Cx43-PKC (Cx43-pS368) was measured. Data show a trend towards an increase in Cx43-PKC from approximately 3 hr post-treatment (n=2).

Supplemental Figure 10: IF of Cx43 expression in SMC following PDGF and peptide treatments. Cells were treated for 16 hr with PDGF-β or PDGF-β with OT pep or CycliCx (30 µm). An n=3 images were taken per experiment (n=3), and results were averaged in the analysis. Highlighted boxes indicate images used in **Figure 3**.

Supplemental Figure 11: Full blots for plasma membrane protein biotinylation in CASMC. CASMC were treated with PDGF-β (50 ng/ mL) alone or in combination with OT peptide or CycliCx (30 µM) for 16 hr. Western blots for Cx43 show plasma membrane-bound proteins (streptavidin eluted, upper blots) compared to cytosolic (non-biotin bound, Lower) proteins (n=3). Orange boxes indicate blots used in **Figure 3**.

Supplemental Figure 12: IF of scrape load dye transfer in SMC following PDGF-β and peptide treatments. CASMC were scraped using a scalpel, followed by the addition of lucifer yellow (green) and Dextran (magenta). For each treatment group (NT, PDGF-β, PDGF-β + OT peptide, and PDGF-β + CycliCx) n=6 biological replicas were used (**A**). NIE image processing software was used to define the dye transfer area (**B**) and calculate the area of lucifer yellow gap junction dye spread minus the area of dextran used to detect scrape-injured cells (**C**). Both total dye area and dye area per scrape length were graphed (**D**). One way ANOVA with multiple comparison test performed, P<0.05 is significant.

Supplemental Figure 13: Full Co-immunoprecipitation western blots for Cx43 and cyclin E. CASMC were treated with PDGF-β (50 ng/ mL) alone or in combination with OT peptide or CycliCx (30 µM) for 16 hr. Blots show cyclin E co-immunoprecipitation with Cx43 detected by western blot (n=3). Control loading blots for Cx43, cyclin E, β- β-tubulin, and total protein. N=4, One-way ANOVA with multiple comparison test performed, P<0.05 is significant. (Orange boxes indicate blots used in Figure 3)

Supplemental Figure 14: SMC EDU proliferation flow cytometry histogram data. Histogram replicates for SMC EDU fluorescence gating for cells treated with PDGF-β, PDGF-β+OT pep, and PDGF-β+CycliCx. N=6 replicates

Supplemental Figure 15: Bulk RNA sequencing quality control, correlation, and clustering. G/C and A/T content distribution along sequence reads for individual replicates in the PDGF-β (PD) group (**A**). Pearson's correlation coefficient matrix of FPKM normalized gene counts before (top panel) and after (bottom panel) removal of the outlying PD replicate (**B**). Heatmap of between replicate FPKM normalized gene expression z-scores with hierarchical clustering (**C**). N = 3-4 per group. Con = no treatment control group; Pep4 = off-target (OT) control peptide; Pep5 = CycliCx peptide.

Supplemental Figure 16: CycliCx peptide blunts mitotic gene pathways. Heatmaps of between replicate 'DESeq2' normalized gene expression z-scores for PDGF-β (PD) + CycliCx vs PD + Off Target Peptide (OT Pep) differentially expressed genes (DEGs) from the significantly enriched GO pathways GO:0044839: 'cell cycle G2/M phase transition (**A**), and GO:0140014 'mitotic nuclear division' (**B**). N = 3-4 per group. NT = No Treatment control group.

Supplemental Figure 17: CycliCx peptide alters the expression of critical cell cycle regulatory genes. 'DESeq2' normalized gene counts in selected genes that regulate early G1 phase progression (**A**), G1/S phase transition (**B**), and G2/M phase transition (**C**). Separated based on CycliCx regulation. N = 3-4 per group. Data displayed as mean ± SEM. NT = No Treatment control group; PD = PDGF-β; OT = Off Target peptide.

Supplemental Figure 18: Gene expression of gap junction proteins and PDGF regulators and transcription targets. 'DESeq2' normalized gene counts in selected gap junction genes (**A**), G1/S phase transition (**B**), and in PDGF regulatory kinases (Erk1/2) and non-cell cycle PDGF transcription targets (**C**). N = 3-4 per group. Data displayed as mean ± SEM. NT = No Treatment control group; PD = PDGF-β; OT = Off Target peptide.

Supplemental Figure 19: CycliCx peptide blunts the expression of smooth muscle contractile genes. 'DESeq2' normalized gene counts in selected vascular smooth muscle cell contractile genes. N = 3-4 per group. Data displayed as mean ± SEM. NT = No Treatment control group; PD = PDGF-β; OT = Off Target peptide.

Supplemental Figure 20: Mouse ligation neointima measurements. H&E staining of mouse carotids following 14 days of sham surgery (no ligation) or ligation with either pluronic gel, pluronic plus OT peptide or pluronic CycliCx peptide. Measure of media area, neointima area and ration of neointima to media. Highlighted images are used as a representative in **Figure 5**. Kruskal-Wallis multiple comparison test, N=12 (9 male, 3 female). * indicates luminal side of the vessel, arrow indicates regions of new intima that were detected in tissues, scale bar is 100 μM.

Supplemental Table 1: Key Resource Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies (IF/ WB/ CO-IP)		
Cx43 (WB/ CO-IP)	Millipore Sigma	C6219 RRID:AB_476857
Cx43 (IF, Rabbit)	Millipore Sigma	HPA035097 UNSPSC:12352203
Cx43 (WB/ CO-IP)	Millipore Sigma	C8093 RRID:AB_259085
Phospho-Cx43 (pSer368) (WB)	Millipore Sigma	SAB4504371
Phospho-Cx43 (pSer 279/282) 0.19 g/mL (IF)	custom antibody Source Dr. Paul Lampe,	Reference PMID: 22652908
Cyclin E1, CCNE1-HE12 (IF, Mouse)	Cell Signaling Technologies	4129
Cyclin E (CCNE1) (IF, WB, Rabbit)	Millipore Sigma	SAB1400044
p44/42 MAPK (Erk1/2) (137F5, Rabbit)	Cell Signaling Technologies	4695
Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) (D13.14.4E) XP®, Rabbit	Cell Signaling Technologies	4370
β-Tubulin (WB, Mouse)	DSHB	E7 RRID:AB_2315513
Transgelin (IF, Goat)	Abcam	ab10135
IRDye® 800CW, Donkey Anti-Rabbit	LI-COR Biosciences	926-32213 RRID:AB_621848
IRDye® 800CW, Donkey Anti-Mouse	LI-COR Biosciences	926-32212, RRID:AB_621847

IRDy(R) 680RD Donkey anti-Rabbit	LI-COR Biosciences	926-68073 RRID:AB_10954442
IRDye 680RD Donkey anti-Mouse	LI-COR Biosciences	926-68072 RRID:AB_10953628
IRDye®800CW Streptavidin	LI-COR Biosciences	926-32230

Chemicals and Reagents

Gibco™ M231 media	ThermoFisher Scientific	M-231-500
VWR Life Science Seradigm Premium Grade Fetal Bovine Serum	VWR Life Science	97068-085
Smooth muscle growth supplement (SMGS)	ThermoFisher Scientific	S00725
Trypsin-EDTA (0.5%), no phenol red	ThermoFisher Scientific	15400-054
Gibco™ M231 media	ThermoFisher Scientific	M-231-500
DPBS (1X), no calcium, no magnesium	ATCC	30-2200
Recombinant PDGF-bb	Millipore	01-305
Ethylene glycol-bis(2-aminoethylether)-N,N,N′,N′-tetraacetic acid	Sigma-Aldrich	E4378
Protease Inhibitor Cocktail	Millipore Sigma	P8340
phosphatase Inhibitor Cocktail 2	Millipore Sigma	P5726
phosphatase Inhibitor Cocktail 3	Millipore Sigma	P0044
RNeasy kit	Qiagen	74004
Odyssey Blocking Buffer (PBS)	LI-COR Biosciences	927-40000

Dynabeads™ M-280 Streptavidin	ThermoFisher Scientific	11205D
EZ-Link™ Sulfo-NHS-LC-Biotin	ThermoFisher Scientific	21335
5-ethynyl-2'deoxyuridine (EDU) Flow cytometry kitt-488 (	ThermoFisher Scientific	C10425
Ethiqa XR 1.3 mg/mL	Covetrus	072117
Isoflurane	Covetrus	029405
Pluronic F127	Millipore Sigma	P2443
CycliCx Peptide: stearate-LDPSKDCGDPKYAY	ThermoFisher custom peptide synthesis services	Custom
OT Peptide: stearate-AYFNGCSSPTAPLDP	ThermoFisher custom peptide synthesis services	Custom

Commercial Assays

Pierce™ BCA Protein Assay Kit	ThermoFisher Scientific	23225
REVERT™ Total Protein Stain for immunoblot Normalization	LI-COR Biosciences	926-11010

Experimental Models: Cell lines and Mice

CASMC: primary human SMC	ATCC	PCS-100-02
CASMC: primary human SMC	Lonza	CC-2583
CASMC: primary human SMC	Thermo Fisher Scientific	C0175C
C57BL/6J mice	Jackson Laboratory	000664

Key:
IF – Immunofluorescence

WB – Western blot
Co-IP co-immunoprecipitation

Supplemental List Cx43CT Truncation Mutation array
Full sequence
QGPLGVKDRVKGRSDPYHATTGPLSPSKDCGSPKYAYFNGCSSPTAPLSPMSPPGYKLVTGDRNNSSCRN

Arrays

(1)
1. Q G P L G V K D R V K G R S D P Y H A T T G P L pS
2. Q G P L G V K D R V K G R S D P Y H A T T G P L S
3. Q G P L G V K D R V K G R S D P Y H A T T G P L A
4. G P L G V K D R V K G R S D P Y H A T T G P L pS
5. P L G V K D R V K G R S D P Y H A T T G P L pS
6. L G V K D R V K G R S D P Y H A T T G P L pS
7. G V K D R V K G R S D P Y H A T T G P L pS
8. V K D R V K G R S D P Y H A T T G P L pS
9. K D R V K G R S D P Y H A T T G P L pS
10. D R V K G R S D P Y H A T T G P L pS
11. R V K G R S D P Y H A T T G P L pS
12. V K G R S D P Y H A T T G P L pS
13. K G R S D P Y H A T T G P L pS
14. G R S D P Y H A T T G P L pS
15. R S D P Y H A T T G P L pS
16. S D P Y H A T T G P L pS
17. D P Y H A T T G P L pS
18. P Y H A T T G P L pS
19. Y H A T T G P L pS
20. H A T T G P L pS
21. A T T G P L pS
22. T T G P L pS
23. T G P L pS
24. P L pS
25. L pS
26. pS
27. .space
(2)
1. Q G P L G V K D R V K G R S D P Y H A T T G P L pS
2. Q G P L G V K D R V K G R S D P Y H A T T G P L S
3. Q G P L G V K D R V K G R S D P Y H A T T G P L A
4. Q G P L G V K D R V K G R S D P Y H A T T G P L
5. Q G P L G V K D R V K G R S D P Y H A T T G P
6. Q G P L G V K D R V K G R S D P Y H A T T G
7. Q G P L G V K D R V K G R S D P Y H A T T
8. Q G P L G V K D R V K G R S D P Y H A T
9. Q G P L G V K D R V K G R S D P Y H A
10. Q G P L G V K D R V K G R S D P Y H
11. Q G P L G V K D R V K G R S D P Y
12. Q G P L G V K D R V K G R S D P
13. Q G P L G V K D R V K G R S D
14. Q G P L G V K D R V K G R S
15. Q G P L G V K D R V K G R
16. Q G P L G V K D R V K G
17. Q G P L G V K D R V K
18. Q G P L G V K D R V
19. Q G P L G V K D R
20. Q G P L G V K D
21. Q G P L G V K
22. Q G P L G V
23. Q G P L G
24. Q G P L
25. Q G P
26. Q G
27. Q
28. .space
(3)pS
1. P S K D C G pS P K Y A Y F N G C S S P T A P L pS P
2. P S K D C G S P K Y A Y F N G C S S P T A P L S P
3. P S K D C G A P K Y A Y F N G C S S P T A P L A P
4. S K D C G pS P K Y A Y F N G C S S P T A P L pS P
5. K D C G pS P K Y A Y F N G C S S P T A P L pS P
6. D C G pS P K Y A Y F N G C S S P T A P L pS P
7. C G pS P K Y A Y F N G C S S P T A P L pS P

8. G pS P K Y A Y F N G C S S P T A P L pS P
9. pS P K Y A Y F N G C S S P T A P L pS P
10. P K Y A Y F N G C S S P T A P L pS P
11. K Y A Y F N G C S S P T A P L pS P
12. Y A Y F N G C S S P T A P L pS P
13. A Y F N G C S S P T A P L pS P
14. Y F N G C S S P T A P L pS P
15. F N G C S S P T A P L pS P
16. N G C S S P T A P L pS P
17. G C S S P T A P L pS P
18. C S S P T A P L pS P
19. S S P T A P L pS P
20. S P T A P L pS P
21. P T A P L pS P
22. A P L pS P
23. P L pS P
24. L pS P
25. pS P
26. P
27. .space

(4)pS
1. P S K D C G pS P K Y A Y F N G C S S P T A P L pS P
2. P S K D C G S P K Y A Y F N G C S S P T A P L S P
3. P S K D C G A P K Y A Y F N G C S S P T A P L A P
4. P S K D C G pS P K Y A Y F N G C S S P T A P L pS
5. P S K D C G pS P K Y A Y F N G C S S P T A P L
6. P S K D C G pS P K Y A Y F N G C S S P T A P
7. P S K D C G pS P K Y A Y F N G C S S P T A
8. P S K D C G pS P K Y A Y F N G C S S P T
9. P S K D C G pS P K Y A Y F N G C S S P
10. P S K D C G pS P K Y A Y F N G C S S
11. P S K D C G pS P K Y A Y F N G C S
12. P S K D C G pS P K Y A Y F N G C
13. P S K D C G pS P K Y A Y F N G
14. P S K D C G pS P K Y A Y F N
15. P S K D C G pS P K Y A Y F
16. P S K D C G pS P K Y A Y
17. P S K D C G pS P K Y A
18. P S K D C G pS P K Y
19. P S K D C G pS P K
20. P S K D C G pS P
21. P S K D C G pS
22. P S K D C G
23. P S K D C
24. P S K D
25. P S K
26. P S
27. P
28. .space

(5)
1. A P L pS P M pS P P G Y K L V T G D R N N S S C R N
2. A P L S P M S P P G Y K L V T G D R N N S S C R N
3. A P L A P M A P P G Y K L V T G D R N N S S C R N
4. P L pS P M pS P P G Y K L V T G D R N N S S C R N
5. P L pS P M pS P P G Y K L V T G D R N N S S C R N
6. pS P M pS P P G Y K L V T G D R N N S S C R N
7. P M pS P P G Y K L V T G D R N N S S C R N
8. M pS P P G Y K L V T G D R N N S S C R N
9. pS P P G Y K L V T G D R N N S S C R N
10. P P G Y K L V T G D R N N S S C R N
11. P G Y K L V T G D R N N S S C R N
12. G Y K L V T G D R N N S S C R N
13. Y K L V T G D R N N S S C R N
14. K L V T G D R N N S S C R N
15. L V T G D R N N S S C R N
16. V T G D R N N S S C R N
17. T G D R N N S S C R N
18. G D R N N S S C R N
19. R N N S S C R N
20. N N S S C R N
21. N S S C R N

22. S S C R N
23. S C R N
24. C R N
25. R N
26. N
27. .space

(6) pS
1. A P L pS P M pS P P G Y K L V T G D R N N S S C R N
2. A P L S P M S P P G Y K L V T G D R N N S S C R N
3. A P L A P M A P P G Y K L V T G D R N N S S C R N
4. A P L pS P M pS P P G Y K L V T G D R N N S S C R
5. A P L pS P M pS P P G Y K L V T G D R N N S S C
6. A P L pS P M pS P P G Y K L V T G D R N N S S
7. A P L pS P M pS P P G Y K L V T G D R N N S
8. A P L pS P M pS P P G Y K L V T G D R N N
9. A P L pS P M pS P P G Y K L V T G D R N
10. A P L pS P M pS P P G Y K L V T G D R
11. A P L pS P M pS P P G Y K L V T G D
12. A P L pS P M pS P P G Y K L V T G
13. A P L pS P M pS P P G Y K L V T
14. A P L pS P M pS P P G Y K L V
15. A P L pS P M pS P P G Y K L
16. A P L pS P M pS P P G Y K
17. A P L pS P M pS P P G Y
18. A P L pS P M pS P P G
19. A P L pS P M pS P P
20. A P L pS P M pS P
21. A P L pS P M pS
22. A P L pS P M
23. A P L pS P
24. A P L pS
25. A P L
26. A P
27. A
28. .space

(7) pS
1. pS P S K D C G pS P K Y A Y F N G C S S P T A P L pS P M pS
2. S P S K D C G S P K Y A Y F N G C S S P T A P L S P M S
3. A P S K D C G A P K Y A Y F N G C S S P T A P L A P M A
4. P S K D C G pS P K Y A Y F N G C S S P T A P L pS P M pS
5. S K D C G pS P K Y A Y F N G C S S P T A P L pS P M pS
6. K D C G pS P K Y A Y F N G C S S P T A P L pS P M pS
7. D C G pS P K Y A Y F N G C S S P T A P L pS P M pS
8. C G pS P K Y A Y F N G C S S P T A P L pS P M pS
9. G pS P K Y A Y F N G C S S P T A P L pS P M pS
10. pS P K Y A Y F N G C S S P T A P L pS P M pS
11. P K Y A Y F N G C S S P T A P L pS P M pS
12. K Y A Y F N G C S S P T A P L pS P M pS
13. Y A Y F N G C S S P T A P L pS P M pS
14. A Y F N G C S S P T A P L pS P M pS
15. Y F N G C S S P T A P L pS P M pS
16. F N G C S S P T A P L pS P M pS
17. N G C S S P T A P L pS P M pS
18. G C S S P T A P L pS P M pS
19. C S S P T A P L pS P M pS
20. S S P T A P L pS P M pS
21. S P T A P L pS P M pS
22. P T A P L pS P M pS
23. T A P L pS P M pS
24. A P L pS P M pS
25. P L pS P M pS
26. L pS P M pS
27. pS P M pS
28. P M pS
29. M pS
30. pS
31. .space

(8) pS
1. pS P S K D C G pS P K Y A Y F N G C S S P T A P L pS P M pS

2. S P S K D C G S P K Y A Y F N G C S S P T A P L S P M S
3. A P S K D C G A P K Y A Y F N G C S S P T A P L A P M A
4. pS P S K D C G pS P K Y A Y F N G C S S P T A P L pS P M
5. pS P S K D C G pS P K Y A Y F N G C S S P T A P L pS P
6. pS P S K D C G pS P K Y A Y F N G C S S P T A P L pS
7. pS P S K D C G pS P K Y A Y F N G C S S P T A P L
8. pS P S K D C G pS P K Y A Y F N G C S S P T A P
9. pS P S K D C G pS P K Y A Y F N G C S S P T A
10. pS P S K D C G pS P K Y A Y F N G C S S P T
11. pS P S K D C G pS P K Y A Y F N G C S S P
12. pS P S K D C G pS P K Y A Y F N G C S S
13. pS P S K D C G pS P K Y A Y F N G C S
14. pS P S K D C G pS P K Y A Y F N G C
15. pS P S K D C G pS P K Y A Y F N G
16. pS P S K D C G pS P K Y A Y F N
17. pS P S K D C G pS P K Y A Y F
18. pS P S K D C G pS P K Y A Y
19. pS P S K D C G pS P K Y A
20. pS P S K D C G pS P K Y
21. pS P S K D C G pS P K
22. pS P S K D C G pS P
23. pS P S K D C G pS
24. pS P S K D C G
25. pS P S K D C
26. pS P S K D
27. pS P S K
28. pS P S
29. pS P
30. pS
31. sp

(9) D
1. Q G P L G V K D R V K G R S D P Y H A T T G P L D
2. Q G P L G V K D R V K G R S D P Y H A T T G P L S
3. Q G P L G V K D R V K G R S D P Y H A T T G P L A
4. G P L G V K D R V K G R S D P Y H A T T G P L D
5. P L G V K D R V K G R S D P Y H A T T G P L D
6. L G V K D R V K G R S D P Y H A T T G P L D
7. G V K D R V K G R S D P Y H A T T G P L D
8. V K D R V K G R S D P Y H A T T G P L D
9. K D R V K G R S D P Y H A T T G P L D
10. D R V K G R S D P Y H A T T G P L D
11. R V K G R S D P Y H A T T G P L D
12. V K G R S D P Y H A T T G P L D
13. K G R S D P Y H A T T G P L D
14. G R S D P Y H A T T G P L D
15. R S D P Y H A T T G P L D
16. S D P Y H A T T G P L D
17. D P Y H A T T G P L D
18. P Y H A T T G P L D
19. Y H A T T G P L D
20. H A T T G P L D
21. A T T G P L D
22. T T G P L D
23. T G P L D
24. P L D
25. L D
26. D
27. .space

(10)
1. Q G P L G V K D R V K G R S D P Y H A T T G P L D
2. Q G P L G V K D R V K G R S D P Y H A T T G P L S
3. Q G P L G V K D R V K G R S D P Y H A T T G P L A
4. Q G P L G V K D R V K G R S D P Y H A T T G P L
5. Q G P L G V K D R V K G R S D P Y H A T T G P
6. Q G P L G V K D R V K G R S D P Y H A T T G
7. Q G P L G V K D R V K G R S D P Y H A T T
8. Q G P L G V K D R V K G R S D P Y H A T
9. Q G P L G V K D R V K G R S D P Y H A
10. Q G P L G V K D R V K G R S D P Y H

11. Q G P L G V K D R V K G R S D P Y
12. Q G P L G V K D R V K G R S D P
13. Q G P L G V K D R V K G R S D
14. Q G P L G V K D R V K G R S
15. Q G P L G V K D R V K G R
16. Q G P L G V K D R V K G
17. Q G P L G V K D R V K
18. Q G P L G V K D R V
19. Q G P L G V K D R
20. Q G P L G V K D
21. Q G P L G V K
22. Q G P L G V
23. Q G P L G
24. Q G P L
25. Q G P
26. Q G
27. Q
28. .space;

(11) D
1. P S K D C G D P K Y A Y F N G C S S P T A P L D P
2. P S K D C G S P K Y A Y F N G C S S P T A P L S P
3. P S K D C G A P K Y A Y F N G C S S P T A P L A P
4. S K D C G D P K Y A Y F N G C S S P T A P L D P
5. K D C G D P K Y A Y F N G C S S P T A P L D P
6. D C G D P K Y A Y F N G C S S P T A P L D P
7. C G D P K Y A Y F N G C S S P T A P L D P
8. G D P K Y A Y F N G C S S P T A P L D P
9. D P K Y A Y F N G C S S P T A P L D P
10. P K Y A Y F N G C S S P T A P L D P
11. K Y A Y F N G C S S P T A P L D P
12. Y A Y F N G C S S P T A P L D P
13. A Y F N G C S S P T A P L D P
14. Y F N G C S S P T A P L D P
15. F N G C S S P T A P L D P
16. N G C S S P T A P L D P
17. G C S S P T A P L D P
18. C S S P T A P L D P
19. S S P T A P L D P
20. S P T A P L D P
21. P T A P L D P
22. A P L D P
23. P L D P
24. L D P
25. D P
26. P
27. .space
(12) D
1. P S K D C G D P K Y A Y F N G C S S P T A P L D P
2. P S K D C G S P K Y A Y F N G C S S P T A P L S P
3. P S K D C G A P K Y A Y F N G C S S P T A P L A P
4. P S K D C G D P K Y A Y F N G C S S P T A P L D
5. P S K D C G D P K Y A Y F N G C S S P T A P L
6. P S K D C G D P K Y A Y F N G C S S P T A P
7. P S K D C G D P K Y A Y F N G C S S P T A
8. P S K D C G D P K Y A Y F N G C S S P T
9. P S K D C G D P K Y A Y F N G C S S P
10. P S K D C G D P K Y A Y F N G C S S
11. P S K D C G D P K Y A Y F N G C S
12. P S K D C G D P K Y A Y F N G C
13. P S K D C G D P K Y A Y F N G
14. P S K D C G D P K Y A Y F N
15. P S K D C G D P K Y A Y F
16. P S K D C G D P K Y A Y
17. P S K D C G D P K Y A
18. P S K D C G D P K Y
19. P S K D C G D P K
20. P S K D C G D P
21. P S K D C G D
22. P S K D C G
23. P S K D C

24. P S K D
25. P S K
26. P S
27. P
28. .space
1.
(13) D
1. D P S K D C G D P K Y A Y F N G C S S P T A P L D P M D
2. S P S K D C G S P K Y A Y F N G C S S P T A P L S P M S
3. A P S K D C G A P K Y A Y F N G C S S P T A P L A P M A
4. P S K D C G D P K Y A Y F N G C S S P T A P L D P M D
5. S K D C G D P K Y A Y F N G C S S P T A P L D P M D
6. K D C G D P K Y A Y F N G C S S P T A P L D P M D
7. D C G D P K Y A Y F N G C S S P T A P L D P M D
8. C G D P K Y A Y F N G C S S P T A P L D P M D
9. G D P K Y A Y F N G C S S P T A P L D P M D
10. D P K Y A Y F N G C S S P T A P L D P M D
11. P K Y A Y F N G C S S P T A P L D P M D
12. K Y A Y F N G C S S P T A P L D P M D
13. Y A Y F N G C S S P T A P L D P M D
14. A Y F N G C S S P T A P L D P M D
15. Y F N G C S S P T A P L D P M D
16. F N G C S S P T A P L D P M D
17. N G C S S P T A P L D P M D
18. G C S S P T A P L D P M D
19. C S S P T A P L D P M D
20. S S P T A P L D P M D
21. S P T A P L D P M D
22. P T A P L D P M D
23. T A P L D P M D
24. A P L D P M D
25. P L D P M D
26. L D P M D
27. D P M D
28. P M D
29. M D
30. D
31. .space
(14) D
1. D P S K D C G D P K Y A Y F N G C S S P T A P L D P M D
2. S P S K D C G S P K Y A Y F N G C S S P T A P L S P M S
3. A P S K D C G A P K Y A Y F N G C S S P T A P L A P M A
4. D P S K D C G D P K Y A Y F N G C S S P T A P L D P M
5. D P S K D C G D P K Y A Y F N G C S S P T A P L D P
6. D P S K D C G D P K Y A Y F N G C S S P T A P L D
7. D P S K D C G D P K Y A Y F N G C S S P T A P L
8. D P S K D C G D P K Y A Y F N G C S S P T A P
9. D P S K D C G D P K Y A Y F N G C S S P T A
10. D P S K D C G D P K Y A Y F N G C S S P T
11. D P S K D C G D P K Y A Y F N G C S S P
12. D P S K D C G D P K Y A Y F N G C S S
13. D P S K D C G D P K Y A Y F N G C S
14. D P S K D C G D P K Y A Y F N G C
15. D P S K D C G D P K Y A Y F N G
16. D P S K D C G D P K Y A Y F N
17. D P S K D C G D P K Y A Y F
18. D P S K D C G D P K Y A Y
19. D P S K D C G D P K Y A
20. D P S K D C G D P K Y
21. D P S K D C G D P K
22. D P S K D C G D P
23. D P S K D C G D
24. D P S K D C G
25. D P S K D C
26. D P S K D
27. D P S K
28. D P S
29. D P
30. D
31. .space

(13) pS
1. pSPSKDCGpSPKYAYFNGCSSPT
2. pSPSKDCGpSPKYAYFNGCSSP
3. pSPSKDCGpSPKYAYFNGCSS
4. pSPSKDCGpSPKYAYFNGCS
5. pSPSKDCGpSPKYAYFNGC
6. pSPSKDCGpSPKYAYFNG
7. pSPSKDCGpSPKYAYFNG
8. pSPSKDCGpSPKYAYFN
9. pSPSKDCGpSPKYAYF
10. pSPSKDCGpSPKYAY
11. pSPSKDCGpSPKYA
12. pSPSKDCGpSPKY
13. pSPSKDCGpSPK
14. pSPSKDCGpSP
15. pSPSKDCGpS
16. pSPSKDCG
17. pSPSKDCGpSPKYAYFNG
18. PSKDCGpSPKYAYFNG
19. SKDCGpSPKYAYFNG
20. KDCGpSPKYAYFNG
21. DCGpSPKYAYFNG
22. CGpSPKYAYFNG
23. GpSPKYAYFNG
24. pSPKYAYFNG
25. PKYAYFNG
26. KYAYFNG
27. AYFNG
28. YFNG

(14)
1. dSPSKDCGdSPKYAYFNGCSSPT
2. dSPSKDCGdSPKYAYFNG
3. FNG
4. .space;

(13) D
1. A P L D P M D P P G Y K L V T G D R N N S S C R N
2. A P L S P M S P P G Y K L V T G D R N N S S C R N
3. A P L A P M A P P G Y K L V T G D R N N S S C R N
4. P L D P M D P P G Y K L V T G D R N N S S C R N
5. P L D P M D P P G Y K L V T G D R N N S S C R N
6. D P M D P P G Y K L V T G D R N N S S C R N
7. P M D P P G Y K L V T G D R N N S S C R N
8. M D P P G Y K L V T G D R N N S S C R N
9. D P P G Y K L V T G D R N N S S C R N
10. P P G Y K L V T G D R N N S S C R N
11. P G Y K L V T G D R N N S S C R N
12. G Y K L V T G D R N N S S C R N
13. Y K L V T G D R N N S S C R N
14. K L V T G D R N N S S C R N
15. L V T G D R N N S S C R N
16. V T G D R N N S S C R N
17. T G D R N N S S C R N
18. G D R N N S S C R N
19. R N N S S C R N
20. N N S S C R N
21. N S S C R N
22. S S C R N
23. S C R N
24. C R N
25. R N
26. N
27. .space
(14) D
2. A P L D P M D P P G Y K L V T G D R N N S S C R N
3. A P L S P M S P P G Y K L V T G D R N N S S C R N
4. A P L A P M A P P G Y K L V T G D R N N S S C R N
5. A P L D P M D P P G Y K L V T G D R N N S S C R
6. A P L D P M D P P G Y K L V T G D R N N S S C
7. A P L D P M D P P G Y K L V T G D R N N S S

8. A P L D P M D P P G Y K L V T G D R N N S
9. A P L D P M D P P G Y K L V T G D R N N
10. A P L D P M D P P G Y K L V T G D R N
11. A P L D P M D P P G Y K L V T G D R
12. A P L D P M D P P G Y K L V T G D
13. A P L D P M D P P G Y K L V T G
14. A P L D P M D P P G Y K L V T
15. A P L D P M D P P G Y K L V
16. A P L D P M D P P G Y K L
17. A P L D P M D P P G Y K
18. A P L D P M D P P G Y
19. A P L D P M D P P G
20. A P L D P M D P P
21. A P L D P M D P
22. A P L D P M D
23. A P L D P M
24. A P L D P
25. A P L D
26. A P L
27. A P
28. A

Figure 1: PDGF-β induces new intima in human saphenous veins in vitro, associated with an increase in Cx43 and proliferation markers

Figure 2: Design of the CycliCx peptide using peptide arrays demonstrates cyclin E binding regions on the phosphorylated Cx43CT

Figure 3: CycliCx Peptide alters Cx43 expression and interactions with cyclin E

Figure 4: Transcriptomics reveals that the CycliCx peptide arrests cell cycle progression at G1/S phase transition

Figure 5: CycliCx peptide limits neointima formation in mouse and human blood vessels

A
14 Day mouse ligation peptide

ligation
+
Pluronic
+
Peptide

Neointima: Media Ratio

Ratio (neointima: media)

>0.9999
<0.0001
<0.0001

Sham | Lig | Lig+OT | Lig+CycliCx

Sham

Lig + Pluronic

Lig + OT Peptide

Lig + CycliCx Peptide

B
Human Saphenous Vein Peptide

TGLN (Fold Change)

0.0079

PD+OT Pep | PD+CycliCx

H&E

TGLN

PDGF + OT Peptide

PDGF + CycliCx Peptide