Directing egg traffic: Internal mechanosensory feedback modulates rhythmic motor activity to coordinate ovulation in *Drosophila*

Shao-Cong Su (苏少聪)[1], Ning Zhang (张宁)[1], Chen-Yu Li (李晨雨)[1], Ji-Yang Xing (邢霁阳)[1], Dick R. Nässel[3], Cong-Fen Gao (高聪芬)[1], Shun-Fan Wu (吴顺凡)[1,2*]

Affiliations:
[1] State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China
[2] College of Sciences, Nanjing Agricultural University, Nanjing 210095, China
[3] Department of Zoology, Stockholm University, Stockholm, Sweden
*Correspondence: wusf@njau.edu.cn

D. R. Nässel Orcid ID: 0000-0002-1147-7766
D. R. Nässel e-mail: dnassel@zoologi.su.se
S. F. Wu Orcid ID: 0000-0003-0096-147X
S. F. Wu e-mail: wusf@njau.edu.cn

Significance

Ovulation requires precisely coordinated contractions of the lateral oviducts to propel eggs into the common oviduct without jamming. Yet the sensory cells and mechanotransduction channels that coordinate movement through the paired lateral oviducts remain unclear. Here, we identify a novel pair of lateral-oviduct multidendritic neurons (mdn-LO) that express the conserved mechanosensory channels TMC (transmembrane channel-like protein), PPK (pickpocket), and Piezo, and show that TMC is uniquely required to prevent egg jamming at the junction between the lateral and common oviducts. The mdn-LO neurons are activated by oviduct contractions and relay signals to female-specific glutamatergic motoneurons in the abdominal neuromere. These paired oviduct motoneurons are rhythmically active and regulate contractions of the lateral oviducts. Together, our findings demonstrate that TMC-mediated mechanosensory feedback fine-tunes motor output to ensure coordinated ovulation and successful reproduction.

Abstract

Ovulation is a prerequisite for successful reproduction and requires precise coordination of muscle contractions, especially to propel eggs from the paired lateral oviducts into the common oviduct. Although internal mechanosensation is expected to provide critical feedback for this process, the sensorimotor circuitry and mechanisms by which sensory feedback shapes motoneuron activity and coordinates oviduct contractions are still poorly known. Here, in *Drosophila,* we identify a novel pair of multidendritic mechanosensory neurons (mdn-LO) in the lateral oviducts that express the mechanosensitive channels TMC (transmembrane channel-like protein), PPK (pickpocket), and Piezo. We show that TMC is essential for coordinated ovulation; *tmc* mutation or neuron-specific knockdown increases egg-jamming at the junction between the lateral and the common oviduct. In contrast, *ppk* or *Piezo* mutants display reduced egg-laying but no jamming. Calcium imaging reveals that the mdn-LO neurons are rhythmically activated by oviduct contractions, and chemogenetic stimulation of these neurons triggers muscle contractions. Genetic manipulations of the mdn-LO neurons disrupt egg passage and induce egg-jamming. The axons of the mdn-LO neurons form functional contacts with female-specific oviduct motoneurons in the abdominal neuromeres that express insulin-like peptide 7 (ILP7) and glutamate. Activation of mdn-LO neurons triggers Ca^{2+} activity in ILP7 motoneurons and elicits rhythmic oviduct contractions. Importantly, the ILP7 motoneurons exhibit spontaneous rhythmic Ca^{2+} activity that is modulated by oviduct contractions and egg movement. Furthermore, triggering activity in the ILP7 neurons induces egg-jamming. Finally, knockdown of ILP7 peptide affects the rate of ovulation, but plays no role in egg jamming, which instead is likely induced by colocalized glutamate. In summary, our study identifies a novel circuit, including the TMC-expressing mdn-LO neurons and a small set of ILP7 motoneurons, that organizes an innate behavior by coupling strategic sensory information and rhythmical motor output to ensure bilateral coordination of oviduct contractions to prevent egg jamming during oviposition.

Introduction

Successful reproduction in oviparous organisms relies on the precise coordination of muscle contractions during ovulation and oviposition, a process orchestrated by neural circuits that generate rhythmic motor outputs and sensory feedback. Studies of the genetically tractable insect *Drosophila* have shown that successful reproduction requires strict and precise regulation of events to occur in coordination, including mechanisms and behaviors associated with oogenesis, follicle cell rupture, motor control of reproductive oviduct muscles to ensure ovulation, egg activation, the choice of egg-laying site, and finally egg deposit [1, 2, 3, 4, 5, 6]. In *Drosophila,* the neuronal circuitry involved in mating and post-mating is well understood [7, 8, 9, 10, 11, 12, 13, 14, 15, 16], and several aspects of the motor control of ovulation and oviposition have been

studied [4, 8, 17, 18]. One critical obstacle during ovulation is the passage of eggs from
the paired ovaries into the common oviduct. To avoid jamming of eggs at the junction
between the two lateral oviducts, critical mechanosensory feedback is required to
ensure sequential activity in motoneurons to generate alternating contractions of left
and right oviduct muscle. The internal sensory neurons providing feedback and motor
circuitry responsible for this directing of egg traffic into the common oviduct have not
been identified. Here, we untangle the neuronal components responsible for this
coordinated egg transport, starting with sensory neurons in the oviduct and
proceeding to motoneurons in the abdominal neuromeres of the ventral nerve cord
(VNC).

Previous studies have identified some sensory neurons in the oviduct. The
contractions of the common oviduct can be sensed by mechanosensory neurons in
the common oviduct that express pickpocket (PPK) [8]. Further down the common
oviduct, a pair of posterior uterine sensory neurons (PU neurons) sense the passage
of eggs into the ovipositor [1]. However, it remains unclear which neurons are
responsible for sensing the contractions of the lateral oviducts. Here, we identify a
novel pair of multidendritic mechanosensory neurons in the lateral oviducts
(designated mdn-LO) that express three evolutionarily conserved mechanosensitive
channels: TMC (transmembrane channel-like protein), PPK (pickpocket), and Piezo,
along with the sex-determination gene *doublesex* (distinct from the adjacent
fruitless-expressing multidendritic neurons). We show that TMC is uniquely required
for ovulation coordination, in contrast to PPK and Piezo. TMC mutation or
neuron-specific knockdown causes severe egg-jamming. Calcium imaging confirms
that mdn-LO neurons are rhythmically activated by lateral oviduct contractions,
consistent with a mechanosensory role.

What about motoneurons regulating contractions of the lateral oviduct muscle?
Two types of efferent neurons that supply axons to the oviduct muscle have been
identified. The first is a set of efferent octopaminergic neurons identified in the
abdominal neuromeres of the VNC. These release octopamine (OA) that acts on two
different receptors (OAMB and Octβ2R) to cause a series of contractions and
relaxations of the muscle in the oviducts to move the egg forward [17, 18, 19, 20, 21]. More
specifically, OA acts on the OAMB receptor to allow entrance of the egg into one of
the lateral oviducts by the rupture of the posterior mature follicle cells surrounding the
oocyte [5]. Further movement through the lateral and common oviducts is caused by
OA modulation of contractions in the myogenic muscle in the reproductive tract [17, 18,
19, 20].

In addition to OA regulation of oviduct contractions, a group of female-specific
glutamatergic motor neurons located in the abdominal neuromeres has been
identified. These bilateral pairs of neurons co-express insulin-like peptide 7 (ILP7) and

were shown to induce contractions of the lateral and the common oviducts [4, 8, 22]. In the following, we call these neurons ILP7-neurons. We show here that silencing of these ILP7-neurons causes jamming of eggs at the junction between the common and lateral oviducts. Using trans-Tango and GRASP analyses together with calcium imaging, we demonstrate that mdn-LO neurons form functional connections with the ILP7 neurons in the abdominal neuromeres. Activation of either mdn-LO or ILP7 neurons induces egg jamming. Moreover, ILP7 neurons exhibit spontaneous rhythmic calcium oscillations that are modulated by mechanosensory input from the lateral oviducts. Thus, acting in concert, the left and right mechanosensory mdn-LO cells and ILP7 motoneurons generate the alternating contractions in the lateral oviducts necessary for coordinated egg entrance into the common oviduct. Although ILP7 neuropeptide is produced by the ILP7-neurons, knockdown of the peptide does not affect egg jamming, but reduces the number of eggs laid, suggesting that its action may be on the common oviduct rather than the lateral ones. In summary, we outline a circuit consisting of a pair of TMC-dependent mechanosensory neurons and a set of rhythmically active ILP7 oviduct motoneurons that ensures smooth passage of eggs through the lateral oviducts and thereby successful oviposition and reproduction. We furthermore describe a novel function of the evolutionarily conserved TMC channels.

Results

Loss of the transmembrane channel-like (*tmc*) gene leads to egg-jamming in *Drosophila*

The paired oviducts connect the ovaries to the uterus and are essential for successful egg transport during ovulation. In *Drosophila*, typically only one egg traverses the oviduct at a time, indicating that internal sensory mechanisms coordinate egg movement to prevent egg jamming. Regarding internal sensation within the female reproductive system, existing reports are limited to *ppk*-expressing neurons in the common oviduct that enhance *Drosophila*'s preference for acetic acid by sensing oviduct contractions [8], as well as posterior uterine (PU) sensory neurons that are activated by egg progression [1]. However, how internal sensory mechanisms coordinate the process of egg transport from the two lateral into the unpaired uterus to avoid egg jamming remains unclear.

In addition to Pickpocket (PPK), PIEZO, and Transmembrane Channel-like (TMC) proteins have also been identified in the female reproductive system of *Drosophila*, though their functional roles remain unclear [8, 23]. PPK, PIEZO, and TMC are evolutionarily conserved mechanosensitive ion channels [8, 23, 24, 25, 26, 27, 28]. In *Drosophila* these channels have been implicated in the regulation of sexual behavior [29, 30], oviposition preference [28], and feeding behavior [23, 31].

Given our previous finding that *Nltmc3*, a TMC ortholog in the brown planthopper (BPH) *Nilaparvata lugens*, is essential for normal ovary development and oviposition

[32](#), we sought to investigate whether the *Drosophila* ortholog *tmc* (CG46121) plays a
similar role. Using *tmc* mutant flies, *tmc[1]* [23](#), we conducted oviposition assays and
found that egg-laying was significantly reduced compared to controls (Fig. 1A).

Since reduced egg-laying can result either from defects in ovarian development
or disruptions in ovulation, we first examined whether developmental effects
contributed to the phenotype. To rule out effects from mating, we initially used virgin
female flies (for other experiments, unless noted otherwise, mated females were
used). We assessed ovarian development in virgin *tmc[1]* mutants by measuring ovary
size and found no developmental abnormalities, in contrast to BPHs [32](#). Instead, we
observed significantly enlarged ovaries in *tmc[1]* flies at both day 2 and day 7
post-eclosion compared to w^{1118} controls (Fig. 1B, D). The enlarged ovaries were due
to the accumulation of mature eggs, suggesting a severe ovulation defect in these
unmated flies (Fig. 1C, D), which was not attributable to abnormalities in egg
morphology (Fig. 1—figure supplement 1).

To further confirm the role of TMC in ovulation, we reintroduced the *tmc* gene into
tmc[1] mutants using the *tmc-GAL4* driver line [23](#). The reduced oviposition, enlarged
ovaries, and accumulation of mature eggs observed in *tmc[1]* mutants were all rescued
by expressing *UAS-tmc* under the *tmc* promoter (Fig. 1A–D).

We next dissected mated *tmc* mutant females to assess ovulation. Interestingly,
we occasionally observed egg-jamming at the junction where the two lateral oviducts
merge into the common oviduct (Fig. 1E and Fig. 1F, respectively). The frequency of
this blockage was significantly higher in *tmc* mutants than in controls. Importantly, the
reintroduction of the *tmc* gene in *tmc[1]* mutants eliminated this egg-jamming phenotype
(Fig.1G). Similarly, RNAi-mediated knockdown of *tmc* in *tmc-GAL4-*expressing cells
also resulted in egg-jamming (Fig. 1H–I).

Finally, we explored whether the other two channels might contribute to
oviposition and ovulation regulation in *Drosophila* (Figure 1—figure supplement 2).
Using *ppk* mutants (*ppk[ESB]*) and *Piezo* mutants (*Piezo[KO]*), we observed a significant
reduction in egg-laying in both strains after mating (Figure 1—figure supplement 2A).
However, we did not observe increased egg-jamming in these mutants or in flies with
RNAi-mediated knockdown of *ppk* or *Piezo* in TMC-GAL4-expressing cells (Figure
1—figure supplement 2B–E).

These findings show that TMC, specifically in *tmc-GAL4-*expressing cells, plays a
key role in coordinating normal ovulation in the lateral oviducts of *Drosophila*, whereas
Piezo and PPK do not. The disruption of TMC leads to a significantly elevated
incidence of egg-jamming, whereas the other channels may affect
ovulation/oviposition via the common oviduct or uterus. The roles of Piezo and PPK in
tmc-positive sensory neurons remain unresolved, underscoring the need for future
studies to reveal the distinct contributions of these two mechanosensory channels and
how the three channels together affect oviduct function.

Figure 1. The *Tmc* channel is essential for ovulation and oviposition in
Drosophila. *Tmc* mutants lay fewer eggs and display ovulation defects. All experiments,
except those in Fig. 1B-D, used mated flies. Groups that share at least one letter are not
statistically distinguishable ($P > 0.05$), whereas groups with different letters are statistically
distinguishable ($P < 0.05$).
(**A**) The number of eggs laid per female of the w^{1118}, tmc^1 mutant, and tmc-GAL4/UAS-tmc;
tmc^1 (mutant with tmc selectively rescued in tmc-GAL4-expressing cells) for 14 h at 25 °C.
One-way ANOVA with Tukey's Multiple Range Test. (**B-C**) Quantification was performed on the
ovary size and the number of mature eggs retained in each pair of ovaries from w^{1118}, tmc^1,
and tmc-GAL4/UAS-tmc;tmc^1 flies. The results show that unmated tmc^1 mutant females
retained more mature eggs in their ovaries, resulting in larger ovaries. This phenotype can be
rescued by expressing tmc in neurons specifically marked by tmc-GAL4. Representative
images of the ovaries are shown in Figure 1D. One-way ANOVA with Tukey's Multiple Range
Test. (**D**) Representative images of w^{1118}, tmc^1, tmc-GAL4/UAS-tmc; tmc^1 ovaries on d2AAE (2
d after eclosion) and d7AEE (7 d after eclosion), from unmated flies. (**E**) Representative
images of w^{1118}, tmc^1, tmc-GAL4/UAS-tmc; tmc^1/tmc^1 ovaries. White arrows mark the egg in
the lateral oviduct. (**F**) Illustration of egg-jamming in the lateral oviduct. (**G**) The egg-jamming
ratio (in %). Genotypes as in Fig 1E. Fisher's exact test. (**H**) Representative images of
w-;;tmc-GAL4/+, tmc-RNAi/+;tmc-GAL4/+ ovaries. White arrows mark the egg in the lateral
oviduct. (I) The egg-jamming ratio (in %). Genotypes as in Fig. 1H. Fisher's exact test.

Figure 1 — figure supplement 1. The *tmc* gene does not affect egg shape.

(**A**) Representative images of eggs of *w1118* flies and the *tmcGAL4* mutant. (**B - C**) The egg length and width of the *w1118* and *tmcGAL4* mutants. Groups that share at least one letter are not statistically distinguishable ($P > 0.05$), whereas groups with completely different letters are statistically distinguishable ($P < 0.05$); Student's t-test.

Figure 1 - figure supplement 2. Egg jamming is not affected by the manipulation of other mechanosensory channel genes.

(**A**) Number of eggs laid per female of *w1118*, *Piezoko*, *ppkESB* flies for 14 h at 25 °C. One-way ANOVA with Tukey's Multiple Range Test. In this and the panels **C and E**: groups that share at least one letter are not statistically distinguishable ($P > 0.05$), whereas groups with completely different letters are statistically distinguishable ($P < 0.05$). (**B**) Representative images of *w1118*, *Piezoko*, *ppkESB* ovaries. (**C**) The egg-jamming ratio. Genotypes as in **B**. Fisher's exact test.(**D**) Representative images of *w-;;tmc-GAL4/+, ppk-RNAi/+; tmc-GAL4/+, Piezo-RNAi/+; tmc-GAL4/+* ovaries. (**E**) The egg-jamming ratio. Genotypes as in **D**. Fisher's exact test.

Tmc is localized in multi-dendritic neurons in the lateral oviduct

Whereas PPK and PIEZO have been previously localized to sensory neurons of the lateral oviducts [8], the presence of TMC in the reproductive system was hitherto not known. Thus, we set out to examine the detailed expression of this channel using a *tmc-GAL4* and by immunohistochemistry. For the latter, we generated a TMC antiserum using a previously reported method [26]. As revealed by immunocytochemical labeling and *tmc-GAL4* expression, TMC is localized in a pair of multi-dendritic neurons in the lateral oviducts that we designate **m**ulti-**d**endritic **n**eurons in the **L**ateral **O**viduct, mdn-LO (Figure 2A-B). Double-labeling experiments with TMC antibodies and *tmc-GAL4*-driven green fluorescent protein (GFP) confirmed the expression of the channel protein in these mdn-LO neurons (Figure 2A). We did not observe anti-TMC immunoreactivity in *tmc*[1] mutant animals, confirming the specificity of the TMC antibodies (Figure 2C). To acquire high-resolution images of the processes of the *tmc-GAL4* labeled neurons, we expressed a membrane-bound GFP (*UAS-mCD8::GFP*) under the control of *tmc-GAL4.* We found that the *tmc-GAL4* specifically identified one mdn-LO neuron in each of the two bilaterally symmetrical lateral oviducts (Figure 2D). TMC is expressed in the dendritic branches, suggesting a sensory role (Figure 2D). Each of these *tmc* neurons covers the native lateral oviduct (the one with the neuronal cell body) with its dendrites (Figure 2D). We found no *tmc*-positive sensory neuron processes in the common oviduct or the uterus (Figure 2D), where branches of PPK-positive neurons have been detected [8, 33]. We also investigated the neuronal expression of two *tmc-GAL4* lines (insertions on the second and third chromosomes, respectively) and a *tmc-LexA* line in the central nervous system. Similar to previous studies [23], we found no *tmc*-positive cell bodies in the abdominal neuromeres (ABD) of the VNC in adult females [34] (Figure 2-supplemental file 1A-C), but there are many *tmc*-expressing neuronal processes projecting from peripheral tissues into the brain and posterior ABD. This part of the ABD is a region dedicated to processing female reproductive sensory input from the periphery and the origin of motor output to muscles involved in ovulation and egg-laying behavior [34] (Figure 2-supplemental file 1A-C).

We next tested whether Piezo and PPK are expressed in neurons of the lateral oviduct. Similar to previous studies [8], we observed both *Piezo-GAL4* and *PPK-GAL4* expression in multidendritic neurons in each lateral oviduct (Figure 2-supplemental file 2A and B). Then we asked whether TMC is co-expressed with PPK or PIEZO in neurons of the lateral oviduct. Indeed, double labeling of *PPK-LexA* and *tmc-GAL4* neurons revealed that the pair of TMC-expressing mdn-LO neurons is also PPK-positive (Figure 2E). We then employed an intersectional strategy [35, 36] using a combination of *tmc-GAL4*, *PPK-LexA,* and FLP recombinase to label mdn-LO neurons and observed one pair of TMC and PPK co-expressing mdn-LO neurons (Figure 2J). The axons of this pair of neurons project from the reproductive system to the abdominal neuromeres of the VNC. (Figure 2J-K)

Using the same approach, we examined whether Piezo and PPK, TMC and Fruitless (fru), as well as TMC and Piezo, are co-expressed in neurons of the lateral oviduct (Figure 2F–I). Taken together, our results indicate that among the two pairs of

286 multidendritic neurons located in the lateral oviducts, one pair (mdn-LO) expresses all
287 three channels, TMC, PPK, and Piezo, as well as Dsx (Doublesex), while the other
288 pair expresses only PPK and fru (Figure 2L) (for Dsx, see also Rezával et al., 2012).
289 Note that FruM is male-specific and absent in females. Thus, the fru expression in
290 females indicates fruP1 regulatory activity rather than FruM protein expression [37].
291

Figure 2. Expression pattern of mechanosensory channels in mdn-LO neurons of the lateral oviducts

(A) Labeling of a lateral oviduct (lat. ovi.) with antiserum to TMC and anti-GFP staining from flies expressing the *UAS-mCD8::GFP* reporter under the control of the *tmc-GAL4*. mdn-LO neurons are indicated by arrows. **(B - C)** Labeling of the lateral oviduct (lat. ovi.) using TMC antibodies in (B) w^{1118} control and (C) tmc^1 mutant. The arrow points to one labeled mdn-LO neuron. **(D)** TMC is specifically expressed in a pair of multi-dendritic sensory neuron cells in the lateral oviduct. Even when the egg enters the lateral oviduct, the mdn-LO neuron dendrites are still wrapped around the lateral oviduct. *tmc* is not expressed in any other part of the female reproductive system: the common oviduct (com.ovi.), spermatheca (sp.) uterus, or parovaria (female accessory glands). **(E)** Lateral oviduct neurons (mdn-LO neurons) that express both *tmc* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows). **(F)** Lateral oviduct neurons (mdn-LO neurons) that express both *Piezo* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows). **(G)** Lateral oviduct neurons that express tmc (green arrows) and are labeled by the fruP1 driver (fruP1+) (red arrows). **(H)** Lateral oviduct neurons that express Piezo (green arrows) and are labeled by the fruP1 driver (fruP1+) (red arrows). **(I)** Lateral oviduct neurons(mdn-LO neurons) that express both *Piezo* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows). **(J)** Expression of *tmc* and *ppk* expressing neurons in the brain, VNC, and lateral oviducts (lat. ovi.), using the intersectional technique (*GAL4/LexA*). The ANm region is targeted by mdn-LO neurons from the lateral oviducts. The genotype: *tmc-GAL4/UAS>stop>myrGFP;ppk-LexA/8xLexAop2-FlpL.* Common oviduct (com. ovi). median abdominal nerve (MAN). **(K)** Illustration of the morphology of the lateral oviduct *tmc-GAL4*-expressing neurons (mdn-LO) in the female reproductive system and ABD of the ventral nerve cord. **(L)** Summary illustration of the expression pattern of mechanosensory and other genes in lateral oviduct neurons. *Dsx*: doublesex.

Figure 2—figure supplement 1. Expression patterns of different *tmc* drivers in the brain, ventral nerve cord, and reproductive system.

(**A-B**) Expression patterns of *tmc-GAL4* (lines with insertion on different chromosomes) expressing neurons in the brain, ventral nerve cord (VNC), and ovary. (**C**) Expression of *tmc-LexA* expressing neurons (cell bodies indicated by white arrows) in the oviduct.

Figure 2—figure supplement 2. Expression of *Piezo* and *ppk* in CNS and reproductive system.

(**A-B**) Expression patterns of *ppk* and *Piezo* in the female VNC (A1, B1) and reproductive system (**A2, B2**). (**A3, A4**) Expression patterns of *Piezo* in the ovary and uterus. White arrowheads mark the lateral oviduct neurons.

A pair of *Tmc*-expressing mdn-LO neurons in the lateral oviduct is essential for ovulation, and avoidance of egg-jamming

We selectively silenced the TMC-expressing neurons by using the *tmc-GAL4* to direct expression of the inward-rectifying potassium channel *Kir2.1* [38] and found that this manipulation causes animals to retain a similar number of eggs as the *tmc^1* mutants do (Figure 3A). The binding of GAL80 to the C-terminal 30 amino acids of GAL4 can prevent GAL4-mediated transcriptional activation [39] and thus inhibit the activity of GAL4 in *Drosophila* [40]. We thus silenced *tmc-GAL4* neurons with *Kir2.1* and examined the effect of *elav-GAL80* suppression of the *tmc-GAL4*-mediated transgene expression in mdn-LO neurons. Interestingly, *tmc-GAL4 > Kir2.1* flies expressing *elav-GAL80* laid almost normal numbers of eggs compared to control flies, indicating that the oviposition phenotype arises predominantly from the *tmc*-GAL4 expressing mdn-LO neurons (Figure 3B). Taken together, these findings show that mdn-LO neurons are necessary for normal ovulation. Furthermore, we asked whether silencing *tmc*-expressing neurons also affects ovulation coordination in *Drosophila*. However, although acute silencing of *tmc-GAL4*-expressing mdn-LO neurons with Kir2.1 reduced oviposition, it did not increase the incidence of egg jamming at the junction between the lateral and common oviducts (Figure 3C-D). We interpret this difference from *tmc* loss-of-function mutants as follows: in *tmc* mutants, mdn-LO neurons are present and active but fail to properly transduce mechanical stimuli, thus producing a defective sensory drive to the motor circuit and thereby promoting

jamming. In contrast, silencing mdn-LO removes the sensory drive altogether, yielding
a state in which fewer eggs reach the junction, leading to reduced oviposition without
increased jamming.

Next, we selectively activated the TMC-expressing neurons thermo-genetically by
using the *tmc-GAL4* to direct the expression of dTrpA1 41. Activation of *tmc*-GAL4
neurons via dTrpA1 (at 30°C) severely impaired oviposition and induced a significant
egg-jamming rate (Figure 3F and G). To clarify which of the *tmc-GAL4* neurons are
functionally required for coordinated oviposition and avoidance of egg-jamming, we
suppressed the activity of GAL4 in selected tissues and cell types using the GAL80
repressor 40. For this, we used two GAL80 lines, *elav-GAL80* 42 and vesicular
glutamate transporter (*vGlut*)-*GAL80* 43. We found that *elav-GAL80* and *vGlut-GAL80*
lines both block the expression of TMC in mdn-LO neurons in the lateral oviduct
(Figure 3E, Figure 3—figure supplement 1). Given that the motor neurons controlling
muscle movements in *Drosophila* are glutamatergic 44, *vGlut-GAL80* serves to
repress any potential GAL4 miss-expression in motor neurons.

To determine whether the behavioral abnormality was due to the mdn-LO
neurons, we combined the *elav-GAL80* or *vGlut-Gal80* transgene with the *tmc-GAL4*
and *UAS-dTrpA1* to suppress activation of mdn-LO neurons. We found that animals
with these manipulations showed no difference in egg-jamming compared with control
flies, while *tmc-GAL4/UAS-dTrpA1* animals displayed increased egg-jamming at 30°C
(Figure 3F and G).

To further confirm our findings, we used the optogenetic effector CsChrimson 45,
which affords greater control over the dynamic range of neuronal activation than
dTrpA1 46, and found that optogenetic stimulation of *tmc-GAL4* expressing neurons in
UAS-CsChrimsonVenus/tmc-GAL4 females abolished oviposition, dramatically
different from all control females [see also 28]. We also observed a significant
egg-jamming rate in the activated *UAS-CsChrimsonVenus/tmc-GAL4* females (Figure
3H and I). Combining pan-neuronal expression of GAL80 with
UAS-CsChrimsonVenus/tmc-GAL4 significantly reduces the rate of egg-jamming
seen after photoactivation (Figure 3I).

Figure 3

Figure 3. A pair of Tmc-expressing mdn-LO neurons in the lateral oviduct is
essential for ovulation, and activation egg-jamming oviduct
In these graphs, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001, and ns
(non-significant). Groups that share at least one letter are not statistically
distinguishable (P > 0.05), whereas groups with completely different letters are

statistically distinguishable ($P < 0.05$).
(**A**) The number of eggs laid per female is reduced when the mdn-LO neurons
are selectively silenced by expression of the hyperpolarizing channel *Kir2.1* in the
tmc-GAL4 line. One-way ANOVA with Tukey's Multiple Range Test. In this and the
other figures: Groups that share at least one letter are not statistically distinguishable
($P > 0.05$), whereas groups with completely different letters are statistically
distinguishable ($P < 0.05$). (**B)** The number of eggs laid per female of mdn-LO
neurons selectively silenced, using Kir2.1, in the presence and absence of
suppression by *elav-GAL80*. The number of eggs laid is restored by neuron-specific
elav-GAL80 suppression. One-way ANOVA with Tukey's Multiple Range Test. (**C**) The
egg-jamming ratio of *UAS-Kir2.1/+, tmc-GAL4/+, tmc-GAL4/+;UAS-Kir2.1/+,* and
tmc-GAL4/elav-GAL80;UAS-Kir2.1/+ flies. Fisher's exact test. (**D**) Representative
images of ovaries of *UAS-Kir2.1/+, tmc-GAL4/+, tmc-GAL4/+;UAS-Kir2.1/+,* and
tmc-GAL4/elav-GAL80;UAS-Kir2.1/+ flies. (**E**) Using the nuclear-localized RFP
(UAS-Redstinger) and the membrane-localized GFP (UAS-mCD8::GFP;
UAS-mCD8::GFP) as a marker, we investigated the labeling of mdn-LO neurons in
the lateral oviducts in the presence of vGlut-GAL80 and elav-GAL80 strains. Both
vGlut-GAL80 and elav-GAL80 can effectively block the tmc-GAL4 labeling of lateral
oviduct neurons (mdn-LO) but not neurons in the CNS. Arrowheads indicate the
mdn-LO neurons in the lateral oviducts. (**F**) Representative images of ovaries from
UAS-dTrpA1/+, tmc-GAL4/UAS-dTrpA1, elav-GAL80;tmc-GAL4/ UAS-dTrpA1, and
tmc-GAL4,vGlut-GAL80/+; UAS-dTrpA1/+ flies at 22°C and 30°C. (**G**) The
egg-jamming ratio of *UAS-dTrpA1/+, tmc-GAL4/UAS-dTrpA1, elav-GAL80;tmc-GAL4/*
UAS-dTrpA1, and *tmc-GAL4,vGlut-GAL80/+; UAS-dTrpA1/+* flies. Fisher's exact test.
(**H**)   Representative   images   of   *UAS-CsChrimsonmVenus*/+   and
tmc-GAL4/+;UAS-CsChrimsonmVenus/+ ovaries after optogenetic activation. The
arrow indicates the location where the egg is jamming in the oviduct. (**I**) Optogenetic
activation of *tmc-GAL4* neurons using CsChrimson. The egg-jamming ratio after
optogenetic activation of *tmc-GAL4* neurons. Fisher's exact test.
427

429 Figure 3 —figure supplement 1

w⁻;UAS-mCD8::GFP;UAS-mCD8::GFP		
w⁻;tmc-GAL4	*w⁻;elav-GAL80;tmc-GAL4*	*w⁻;tmc-GAL4;vGlut-GAL80*

Brain

anti-GFP nc82

VNC

anti-GFP nc82 — 50µm

Figure 3—figure supplement 1. Expression of tmc-GAL4 in the presence of different

GAL80 in the brain and ventral nerve cord.

Using the membrane-localized GFP (*UAS-mCD8::GFP; UAS-mCD8::GFP*) as a marker, we

investigated the labeling of mdn-LO neurons in the CNS in the presence of *vGlut-GAL80* and

elav-GAL80 strains. Note that in the *tmc-GAL4, elav-GAL80 > GFP*, only a subset of the

GFP-labeled inputs to the subesophageal zone and the metathoracic ganglion of the VNC

are retained; in the *tmc-GAL4; vGlut-GAL80 > GFP*, by contrast, there are fewer labeled

axons in the ANm, indicating that inputs from the reproductive system are blocked.

***tmc*-expressing sensory neurons in the reproductive oviduct can sense**

contractions of the lateral oviduct

Our experiments, so far, suggest that the female flies sense the passage of eggs

through the reproductive tract by means of mechanosensory neurons. Thus, we next

asked whether the mdn-LO neurons monitor oviduct muscle contractions and

oviposition.

To investigate whether ovulation and oviposition affect activity in mdn-LO neurons,

we conducted an assay using a transcriptional reporter of intracellular Ca^{2+} (TRIC)

that monitors sustained neuronal activity 47. Indeed, we found a significant elevation

of TRIC signals, corresponding to increased intracellular Ca^{2+} levels, in mdn-LO
neurons of egg-laying virgin flies compared with virgin flies that did not lay eggs,
monitored over 5 days (Figure 4A and A'). This finding is consistent with elevated
long-term activity in mdn-LO neurons during egg laying.

Next, to further explore the requirement of mdn-LO neurons for ovulation-induced
neuronal activity, we monitored Ca^{2+} dynamics in the cell body of mdn-LO neurons in
response to ovulation. To visualize changes in Ca^{2+} levels in cell bodies of mdn-LO,
we expressed a transgene encoding the Ca^{2+} sensor GCaMP6m (*UAS-GCaMP6m*)
and Tdtomato (*UAS-Tdtomato*) [48] in mdn-LO neurons under the control of the
tmc-GAL4. We surgically exposed the lateral oviduct region of the female reproductive
area, aiming to monitor the fluorescence changes of mdn-LO neurons as the egg
passed through the lateral oviduct. However, we found that when the egg passes
through the lateral oviduct, an inevitable movement occurs along the Z-axis, causing
us to lose the signal. Even with the addition of tdTomato as a negative control, the
tissue displacement still prevented us from accurately monitoring the change in
fluorescence in the mdn-LO neurons as the egg passed through the lateral oviduct.
Therefore, we monitored the activity of mdn-LO neurons when the lateral oviducts
undergo spontaneous contractions. We observed that calcium activity in the cell body
of the mdn-LO neurons increased when the muscle contracted (Figure 4B and Movie
S1).

We also used another calcium sensor, GCaMP7s [49,] to monitor activity in axonal
termini of mdn-LO neurons in the VNC. This was done since the GCaMP7s signal in
tmc-expressing processes in the VNC is less likely to move out of the focal plane
when oviduct contraction occurs, and thus facilitates measurements. We found that
the CGaMP7s signal in the axonal termini of *tmc* neurons increased significantly
during oviduct contraction (Figure 4C-C"; Movie S2). To ensure that such GCaMP7
increase was not due to "local interaction" between *tmc*-expressing neurons and
other neurons in the VNC, we physically severed the connection (median abdominal
nerve, MAN) between the oviducts and the VNC. In these severed preparations, we
did not observe a significant GCaMP7s increase in *tmc*-expressing axons in the VNC
during oviduct contractions (Figure 4D-D"; Movie S3). This finding supports that
mdn-LO neurons convey contraction-related signals from the oviduct to the VNC
rather than the Ca^{2+} transients being the result of local neuronal interactions in the
VNC. Thus, our data strongly suggest that the *tmc*-expressing mdn-LO neurons of the
lateral oviducts are mechanosensory and can detect contractions/distentions in this
portion of the reproductive tract.

We next asked whether mdn-LO neurons can also be influenced by inputs other
than mechanosensory stimulation during ovulation. Since octopamine (OA) has been
implicated as a regulator of ovulation in insects [18], we tested the effect of exogenous
OA on mdn-LO activity. We found that OA application increased Ca^{2+} signals in the
axon terminals of GCaMP6m-expressing mdn-LO neurons both in intact animals and
after severing the connection to the oviduct (Figure 4 —figure supplement 1),
suggesting OA modulates activity in mdn-LO neurons independently of
mechanosensory input.

Finally, we used a chemical-genetic approach [50, 51] to stimulate the mdn-LO
neurons. We expressed the ATP-gated P2X2 channel, which is not expressed
endogenously in flies, and can be used for 'chemogenetic activation' of specific
neurons of interest. When stimulating the P2X2 expressing mdn-LO neurons with ATP,
a robust contraction of the oviduct muscle was induced (Movie S4). This shows that
mdn-LO mechanosensory neurons in the lateral oviduct can not only sense oviduct
contraction, but also act through downstream motoneurons to regulate contractions of
the lateral oviduct.
502

503 Figure 4

506 **Figure 4. The lateral oviduct *tmc*-expressing neurons (mdn-LO) are mechanosensory**
507 **and are activated during ovulation.**

(**A**) Monitoring sustained calcium activity in mdn-LO neurons using the TRIC method. Comparison of TRIC signals of mdn-LO sensory neurons between egg-laying and non-egg-laying females and representative images. We collected flies that were 5 days post-eclosion and dissected them, separating those that laid eggs and those that did not lay any eggs. After dissection, without staining, we immediately observed the TRIC signal of mdn-LO in real time under the same parameters (we used GFP/RFP as the TRIC signal). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$, and ns (non-significant); Two-tailed t-tests. (**B**) When the lateral oviduct contracts rhythmically, it is accompanied by rhythmic excitation of mdn-LO sensory neurons, as shown by increased Ca^{2+} levels monitored with Gcamp6m. (**C-C''**) In intact animals (median abdominal nerve, MAN, intact), lateral oviduct mdn-LO sensory neurons transmit excitation to the abdominal neuromere (ANm), and Ca^{2+} activity in axon terminations of the left and right neurons can be monitored. (**D- D''**) When the MAN is surgically severed, no Ca^{2+} activity is detected in axon terminations in the ANm during spontaneous muscle contractions. This supports that the activation in intact animals is derived from the axons of the lateral oviduct mdn-LO sensory neurons.

Figure 4—figure supplement 1. Mdn-LO neurons can be activated by octopamine in intact animals and after severing the connection to the oviduct.

(A-A'') *tmc*+ neurons can be activated by octopamine (A). Illustration of maintaining the connection via the median abdominal nerve (MAN) and cutting MAN. (A') Calcium imaging of mdn-LO neuron axons in the ANm region before and after the addition of 1 mM octopamine in intact animals and after severing the MAN. (A'') Changes in fluorescence intensity ($\triangle F/F_0$) of Gcamp6m in mdn-LO neuron axons in the ANm region before and after octopamine addition.

ILP7-expressing neurons in the abdominal ganglion receive input from *tmc*-expressing mdn-LO neurons

We next sought to identify how TMC-expressing mechanosensory mdn-LO neurons engage the central circuitry that controls oviduct contractility.

To map candidate downstream partners of mdn-LO neurons, we first used trans-Tango [52] to visualize neurons postsynaptic to *tmc*-GAL4–labeled mdn-LO neurons. Expression of trans-Tango in mdn-LO neurons labeled numerous putative postsynaptic neurons, with the strongest labeling in the ABD of the VNC (Figure 5A-C'') and in the suboesophageal zone (SEZ) (Figure 5—figure supplement 1A). We,

however, did not detect a trans-Tango signal in any local neurons in the ovary or
oviduct (Figure 5—figure supplement 1B). This finding further supports that mdn-LO
neurons act as sensory neurons that transmit signals from the oviduct muscle to the
central nervous system. Since the trans-Tango labeled numerous neurons, the
identification of individual post-synaptic neurons was not possible. It has been shown
previously 22, and we confirmed here, that some of the *Ilp7-GAL4*-expressing neurons
in the ventral portion of the posterior VNC are efferent neurons that innervate the
oviduct (Figure 5D, Figure 5—figure supplement 2H). Importantly, it was previously
shown that inhibiting ILP7-neurons consistently causes one or more eggs to be
jammed in the oviduct 4. Furthermore, efferent neurons in the posterior VNC that
produce OA, identified by a *tdc2-GAL4,* are also involved in the regulation of ovulation
19, 53, 54, 55. Thus, we next used the syb:GRASP technique 56 to address ~~the question~~
~~of~~ whether axons of *tmc*-expressing mdn-LO neurons directly contact processes of
Ilp7-expressing or *tdc2*-expressing neurons in the abdominal neuromeres of the VNC.
In this method, half of the GFP is targeted specifically to the presynaptic termini of the
mdn-LO neurons and the other to *Ilp7* or *Tdc* neurons, and GFP reconstitution occurs
at exposure to the synaptic cleft only upon synaptic vesicle fusion. With *Ilp7* neurons
and *Tdc* neurons separately as partners of mnd-LO neurons, we were able to detect
reconstituted GFP clearly in the ABD of the VNC (Figure 5E-H; Figure 5—figure
supplement 2B, D), suggesting that mdn-LO neurons can act as presynaptic partners
of both *Ilp7* neurons and *Tdc* neurons in the ABD. However, we did not find evidence
with the GRASP technique that *Ilp7* or *tdc2* neurons are coupled pre-synaptically to
mdn-LO neurons (Figure 5—figure supplement 1C and 1E). Thus, the activation of
mdn-LO neurons by applied OA, shown in Figure 5—figure supplement 1, is likely to
represent paracrine (non-synaptic) signaling from octopaminergic neurons.
Furthermore, we found that *tdc2*-GAL4 and *Ilp7*-GAL4 signals are not co-expressed in
neurons in the ABD or the brain (Figure 5—figure supplement 1I). However, we could
show by GRASP that *tdc2* neurons are presynaptically coupled to *Ilp7*-expressing
neurons, suggesting functional connections between these neurons (Figure 5—figure
supplement 1F-G).
To test whether the contacts observed between mdn-LO and ILP7 neurons using
the GRASP technique permit functional signal transmission, we next asked whether
stimulation of mdn-LO neurons affects Ca²⁺ levels in ILP7 neurons. For this, we
established an *ex vivo* "brain-less" preparation in which the full oviduct, where ovaries
remain connected to the VNC, while the connection to the brain is severed. In this
preparation, we monitored Ca²⁺ dynamics in ILP7 neurons using GCaMP6m 48. To
stimulate mdn-LO neurons, we expressed in these neurons the ATP-gated P2X2
channel 50, 51.
We found that stimulating the P2X2-expressing mdn-LO neurons with ATP
triggered a significant increase in GCaMP signal in the cell bodies of *Ilp7* neurons in
the ventral portion of the abdominal neuromeres of the VNC (Figure 5I-O),
accompanied by rhythmic contractions of the oviduct (Movie S4). We performed three
control experiments where no increase in GCaMP signals was observed in ILP7
neurons: (i) genotype controls lacking P2X2 expression in mdn-LO neurons (Figure

5I), (ii) vehicle controls, AHLS (ATP solvent) applied with the same timing/volume elicited no response (Figure 5K), and (iii) preparations in which the median abdominal nerve that connects oviduct and VNC was severed (Figure 5L).

Interestingly, we did not detect comparable elevations of the GCaMP signal in ILP7 neurons located on the dorsal side of the ABD. Thus, the Ca²⁺ increase was confined to the ventral cluster that includes the previously identified female-specific post-embryonic ILP7 oviduct motoneurons (Figure 5D, 5K, 5M-5O). Collectively, our findings demonstrate that mdn-LO axons form functional contacts with ILP7 neurons in the ABD, and that activation of the mdn-LO neurons induces increased Ca²⁺ activity in the ventral subset of these ILP7 neurons, which in turn modulates contractions in the lateral oviducts.

tmc-GAL4/UAS-myrGFP attp8,QUAS-mTdtomato (3xHA);trans-Tango/+

Type	Innervate
Embryonic *ilp7* neurons (5, 6)	Gut
Embryonic *ilp7* neurons (7-10)	unknown
Post-embryonic *ilp7* neurons
Female-specific post-embryonic *ilp7* neurons	Oviduct

(Modified from Castellanos, Development, 2013)

w-;tmc-LexA/LexAop-nsyb-GFP^{1-10},UAS-CD4-GFP^{11};ilp7-GAL4/+

mated

LexAop-P2x2/+;
ilp7-GAL4/UAS-Gcamp6m

tmc-LexA/LexAop-P2x2;ilp7-GAL4/UAS-Gcamp6m

▬ *Lexaop-P2x2;ilp7>Gcamp6m* +ATP (Ven) n=3	▬ *Tmc>P2x2;ilp7>Gcamp6m* +ATP (Ven) n=5
▬ *Tmc>P2x2;ilp7>Gcamp6m* +ATP (Dor) n=3	▬ *Tmc>P2x2;ilp7>Gcamp6m* +ATP (Ven,cut MAN) n=3
▬ *Tmc>P2x2;ilp7>Gcamp6m* +AHLS (Ven) n=5	

Embryonic *ilp7* neurons (5,6) | Embryonic *ilp7* neurons (7-10) | Female-specific post-embryonic *ilp7* neurons / Post-embryonic *ilp7* neurons

Figure 5. The mdn-LO sensory neurons regulate activity in the *Ilp7* neurons in the ANm, and their connections are visualized by trans-tango and GRASP.

(A-C'') Putative second-order neurons targeted by *tmc*+ neurons (green) in the ANm region revealed by trans-Tango (red) driven by *tmc*-GAL4. In A-B', we show the expression of presynaptic signal (GFP/dark) in mdn-LO neurons and postsynaptic signal (RFP/dark) of *tmc*+ neurons in the ANm region of VNC. In C-C, merged confocal images of trans-Tango labeling of postsynaptic neurons (GFP/green) to *tmc*+ neurons (RFP/red).

(D) Location of *Ilp7* neurons in the ventral and dorsal ANm regions. Blue circles indicate embryonic *Ilp7* neurons; pink circles indicate large female-specific post-embryonic *Ilp7* neurons; Green circles indicate the post-embryonic *Ilp7* neurons. Original data from 22. The three types of *Ilp7* -expressing cell clusters innervate different tissues. The embryonic *Ilp7* neurons innervate the intestine, and the female-specific post-embryonic *Ilp7* neurons and post-embryonic *Ilp7* neurons are glutamatergic motoneurons that innervate the oviduct muscle of the female reproductive system. (E-H) GRASP signals between mdn-LO sensory neurons and *Ilp7* neurons are observed in the brain, dorsal and ventral VNC, and ovary. (I–L) The mdn-LO sensory neurons regulate the excitation of ventral VNC Ilp7 neurons in the ANm region. In control preparations lacking P2X2 expression in mdn-LO neurons (LexAop-P2X2/+), application of 1 mM ATP did not increase Ca^{2+} activity in ventral Ilp7 neurons (I). In contrast, activating P2X2-expressing mdn-LO neurons with 1 mM ATP robustly increased Ca^{2+} activity in ventral Ilp7 neurons, whereas dorsal Ilp7 neurons were not affected (J). (K) Vehicle control: AHLS (ATP solvent) applied with the same timing/volume elicited no response. (L) MAN-severed preparation: after cutting the median abdominal nerve (MAN) connecting the reproductive tract/periphery to the abdominal ganglion, ATP application failed to evoke ventral Ilp7 responses. (M-O) Graphs showing activation of *Tmc*+ neurons via *P2X2* increased different *Ilp7*+ motor neuron activity. Fluorescence changes ($\Delta F/F_o$) of *GCaMP6s* in *Ilp7* neurons are shown following different treatments. The yellow bar indicates the addition of the solution.

Figure 5—figure supplement 1. The presumed second-order neurons of mdn-LO target the subesophageal zone (SEZ) region in the brain.

(A) Putative second-order neurons of *tmc*+ neurons (green) in the suboesophageal ganglion (SOG) region revealed by trans-Tango (red). (B) Using the trans-tango system, it was found that there are no putative second-order neurons of mdn-LO in the oviduct.

Figure 5—figure supplement 2. Probing connections between *tdc2+*, *Ilp7+*, and *tmc+*

neurons by the GRASP technique and calcium imaging.

(A) Lack of GRASP signal from expressing *tmc-LexA* alone in the brain and ventral VNC.

(B) Flies in virgin status were used to obtain the GRASP signal between mdn-LO (expressing

tmc-LexA) and *Ilp7* + neurons with *Ilp7-GAL4* in the brain, dorsal, and ventral VNC. The black

signal represents the region of putative synaptic connections (indicated by black dashed lines).

(C) No GRASP signal is seen from the *Ilp7* + neurons to *tmc*+ neurons in the brain, dorsal VNC,

and ovary. (D) GRASP signal from the *tmc*+ neurons to *tdc2*+ neurons in the dorsal and ventral VNC. The dark signal represents the region of synaptic connections (indicated by black dashed lines). (E) No GRASP signal is seen from the *tdc2*+ neurons to *tmc*+ neurons in the ventral VNC. (F) GRASP signal from the *tdc2*+ neurons to *Ilp7* + neurons in the ventral and dorsal VNC. The dark signal represents the region of putative synaptic connections (indicated by black dashed lines). (G) GRASP signal from the *Ilp7* + neurons to *tdc2*+ neurons in the ventral and dorsal VNC. The dark signal represents the region of putative synaptic connections (indicated by black dashed lines). (H) The expression of *Ilp7*-GAL4 positive neurons in the brain, VNC, and ovary. (I) *Ilp7*+ (green) and *tdc2*+ (red) neurons within the ANm and brain are distinct. (J) There is no functional connection between ILP7 neurons and Tdc2 neurons. When ILP7 neurons are activated by ATP, the calcium signal in Tdc2 neurons shows no significant change; the converse is also true.

ILP7 neuron activity disrupts egg passage and exhibits mechanically tunable rhythmic Ca²⁺ dynamics.

Previous work showed that hyperpolarizing *Ilp7-GAL4* neurons in the posterior VNC leads to egg jamming in the oviduct [57]. However, the effects of activating these neurons and the role of ILP7 peptide on ovulation and oviposition have not been investigated. We found that optogenetic activation of *Ilp7-GAL4* neurons using CsChrimson markedly increased the fraction of females exhibiting egg-jamming compared with genetic controls (Figure 6A). Furthermore, thermogenetic activation of *Ilp7-GAL4* neurons by driving dTrpA1 reduced the number of eggs laid at 30 °C compared to 22 °C controls and increased the incidence of egg jamming (Figure 6B–D).

To assess the contribution of the ILP7 neuropeptide during ILP7 neuron activation, we reduced ILP7 levels by RNAi (*Ilp7-GAL4 > UAS-Ilp7-RNAi*) while activating ILP7 neurons with dTrpA1. Under these conditions, the number of eggs laid was partially restored relative to ILP7 activation alone (Figure 6B), whereas the egg-jamming frequency was not significantly reduced (Figure 6D). Together, these data indicate that elevating neuronal activity in ILP7 neurons elicits egg-jamming, while the ILP7 peptide contributes primarily to overall oviposition rather than playing a role in the jamming phenotype (at least under our assay conditions). This finding might suggest that the ILP7 peptide acts at the level of the common oviduct rather than on the lateral ones.

We next examined the activity dynamics of ILP7 neurons using *Ilp7-GAL4 > GCaMP6m*. In *ex vivo* "brain-free" VNC preparations, *Ilp7*-GAL4–expressing female-specific post-embryonic neurons and embryonic ILP7 neurons display spontaneous rhythmic Ca²⁺ oscillations (Figure 6E). These two neuron types can be distinguished by their locations.

Following enzymatic digestion (protease/collagenase), the rhythmic oscillations were still observed, although they were slower (Figure 6F). Because digestion makes it difficult to classify ILP7 subtypes by position, we pooled the ILP7 neurons for quantification. The oscillation frequency decreased from ~0.040 Hz before digestion to ~0.019 Hz after digestion (Figure 6G), consistent with an intrinsic or locally

generated oscillatory capacity, although additional network contributions to the rhythmicity cannot be excluded.

Finally, we asked whether the rhythmic activity of the ILP7-nerons is modulated by oviduct-linked internal mechanosensory feedback. Thus, we recorded prolonged Ca^{2+} activity from multiple ILP7 neurons while scoring egg position and movement within the reproductive tract. Across such a long recording, ILP7 neurons showed distinct oscillatory modes and transitions that covaried with changes in egg movement (Figure 6H). Time points at which egg position was scored are indicated (Figure 6H), and the corresponding sequence of egg movement states is summarized schematically (Figure 6I, J). Our observations suggest an association between the movement of the egg through the oviduct and Ca^{2+} dynamics in ILP7 neurons, and furthermore provide an indication of how internal mechanosensory feedback influences excitatory motor output during egg transport.

Figure 6. ILP7 neuron activity disrupts egg passage and exhibits mechanically tunable rhythmic Ca²⁺ dynamics.

(A) Optogenetic activation of ILP7-GAL4 neurons (UAS-CsChrimsonmVenus; ILP7-GAL4) increases the fraction of females exhibiting egg jamming (left). Representative images of dissected female reproductive tracts after optogenetic stimulation are shown (right); the boxed example highlights a jammed egg in the oviduct.

(B) Thermogenetic activation of ILP7 neurons (UAS-dTrpA1; ILP7-GAL4) reduces oviposition

at 30 °C relative to 22 °C controls, and this reduction is partially alleviated by Ilp7 RNAi during
activation (UAS-Ilp7-RNAi; UAS-dTrpA1; ILP7-GAL4).
(C) Representative images of dissected female reproductive tracts from the thermogenetic
activation experiment in (B) at 22 °C and 30 °C, showing egg retention/jamming in activated
conditions (arrowheads).
(D) Egg-jamming frequency following thermogenetic activation of ILP7 neurons, with or without
Ilp7 RNAi, at 22 °C and 30 °C (genotypes as in B). Groups sharing at least one letter are not
statistically distinguishable.
(E) Spontaneous rhythmic Ca²⁺ oscillations recorded with GCaMP6m in ILP7-GAL4 neurons in
the adult VNC preparation, including female-specific post-embryonic ILP7 neurons (pink) and
embryonic ILP7 neurons (dark and light blue).
(F) Rhythmic Ca²⁺ oscillations persist in ILP7 neurons after enzymatic digestion
(protease/collagenase), although oscillations are slowed.
(G) Quantification of oscillation frequency in ILP7 neurons before and after enzymatic
digestion (AHLS, Adult Hemolymph-Like Saline).
(H) ILP7-neuron Ca²⁺ dynamics are modulated by egg position within the oviduct. Traces show
Ca²⁺ signals from eight ILP7 neurons (color-coded ROIs) across a continuous recording that
was segmented into 15 consecutive time windows (roman numerals I–XV). Red numbered
boxes indicate periods corresponding to discrete egg-position states defined in (J), whereas
gray boxes mark recording intervals during which no egg manipulation was performed.
In the initial state G1 (I–IV; right egg enters the oviduct), neuron 4 exhibits rhythmic Ca²⁺
oscillations that progressively slow and terminate by IV. After the left egg is pushed into the
oviduct (G2, V–IX), neuron 4—previously quiescent—shows a brief burst of oscillations, then
pauses again, with oscillations re-emerging by IX. After the left egg is pulled out (G3, X),
neurons 1, 6, and 8 decrease in baseline signal, neuron 4 stops oscillating, and neurons 4 and
5 show increased activity; notably, neuron 2 displays a newly emergent oscillatory pattern.
When the right egg is pulled out (G4, XI), neuron 2 activity decreases, whereas pushing the
right egg back in (G5, XII–XIV) restores and re-initiates rhythmic oscillations in neuron 2,
consistent with regulation by right-side mechanosensory feedback. Finally, when the left egg is
pushed back into the oviduct (G6, XV), oscillations in neuron 2 disappear.
(I) Schematic of the ex vivo imaging configuration used for the recording in (H), in which the
ANm region of the VNC was retained for Ca²⁺ imaging. The upper inset indicates the eight
ILP7 neurons monitored and the corresponding ROI color codes used in (H).
(J) Schematic timeline of egg-position manipulations and scored egg-location states. Numbers
indicate the sequence of events and correspond to the red numbered boxes in (H).
744
745

Female reproductive system

Working model of mechanosensory feedback control of ovulation in *Drosophila*

Figure 7. Proposed model of mechanosensory feedback control of ovulation in *Drosophila*.

(A) Graphic summary of the female reproductive tract and ventral nerve cord (VNC).

(B) We propose a working model of ovulation in which TMC-dependent lateral-oviduct mechanosensory neurons (mdn-LO; TMC/PPK/Piezo$^+$) detect contraction- and egg-movement–linked mechanical cues and relay this information to the abdominal ganglion, where it shapes motoneuron activity patterns of ILP7-expressing glutamatergic oviduct motoneurons to coordinate egg passage from the paired lateral oviducts into the common oviduct and thereby promote successful ovulation. ILP7 peptide signaling contributes to overall oviposition output but is not a determinant of egg-jamming phenotypes under our experimental conditions. The yellow ppk$^+$/fru$^+$ LO neuron (here labeled mdn-LO_fru for clarity; previously described without a specific name). The mdn-LO_fru neurons constitute a second pair of lateral-oviduct multidendritic neurons (ppk$^+$ fru$^+$) reported previously; their role in egg-traffic coordination was not directly examined here.

Discussion

In this study, we describe a novel neuronal circuit that regulates coordinated oviposition in *Drosophila*. A pair of multidendritic sensory neurons, mnd-LO, in the lateral oviducts respond to muscle contractions as the egg passes and provide feedback to bilateral sets of motoneurons (ILP7 neurons) in the VNC to ensure coordinating lateral oviduct contractions to propel eggs into the common oviduct without jamming. In this circuit, we discovered a previously unrecognized role for the mechanosensory channel protein TMC. The TMC channel is expressed in the mnd-LO neurons together with two additional mechanosensory channels, PPK and PIEZO. Mutations in all three channel genes result in defects in ovulation/oviposition. However, only mutation or cell-specific knockdown of *tmc* markedly elevates the

incidence of egg jamming at the junction between the two lateral oviducts and the common oviduct. Together, these findings highlight a key role for mechanosensory feedback in ensuring reliable egg transport in the reproductive tract, with TMC being specifically required to prevent jamming at the entrance to the common oviduct. Thus, while mdn-LO neurons co-express TMC, PPK, and PIEZO, our data suggest that these channels contribute in distinct ways to the mechanosensation of the lateral oviduct and that they function complementarily in this sensory process. Future work is required to understand in detail the specific roles of Piezo and PPK in the *Tmc*-positive sensory neurons. Interestingly, a recent report showed that PIEZO channels are required for effective uterine contractions during mammalian parturition [58], suggesting an evolutionary conservation of function.

The mdn-LO neurons display rhythmic [Ca^{2+}] activity during spontaneous and induced contractions of the oviduct muscle, consistent with a mechanosensory role. Using trans-Tango and GRASP, we found that mdn-LO neurons are presynaptic to a set of female-specific efferent neurons in the abdominal neuromere that co-express glutamate and the neuropeptide ILP7. Chemogenetic stimulation of mdn-LO neurons increases [Ca^{2+}] in ILP7 neurons and elicits rhythmic contractions of oviduct musculature. Previous studies established that activity in these female-specific ILP7 neurons is required for proper ovulation and oviposition. It is likely mediated via glutamatergic neuromuscular transmission [4, 8, 19, 22]. However, the functional contribution of the co-localized ILP7 peptide to oviduct contractions was not investigated. Our study demonstrates that blocking ILP7 peptide release (by *Ilp7*-RNAi) during activation of ILP7 neurons recovers oviposition but does not affect egg jamming. Thus, ILP7 may play a modulatory role further down the reproductive tract, such as in the common oviduct, and thereby influence the rate of egg laying, possibly by means of short-range hormonal signaling (or paracrine/non-synaptic signaling). In contrast, it is likely that the ILP7 neurons utilize glutamate for direct motor output to coordinate lateral oviduct muscle [4, 8, 19, 22].

Interestingly, the specific location of egg jamming in the oviduct depends on which component of the pathway is manipulated. Activation or inactivation of ILP7 neurons leads to egg jamming in the entrance of the common oviduct, whereas activation of mdn-LO neurons preferentially generates egg jamming in the lateral oviducts. This finding suggests that the two mdn-LO neurons provide sensory feedback from each of the lateral oviducts to modulate the two pairs of ILP7-motoneurons to generate alternating activity in the two lateral oviducts to ensure a smooth passage of eggs into the common oviduct.

Apart from the ILP7-neurons and mdn-LO neurons shown here and descending neuronal pathways from the brain [3, 16], there are additional neurons in the control of ovulation in *Drosophila*. These are a second pair of multidendritic sensory neurons in the lateral oviducts that express PPK, but not the TMC and PIEZO channels [8], and a small set of octopaminergic efferent neurons (*tdc*-neurons) in the abdominal neuromeres of the VNC that innervate the muscle of the oviduct [17, 18, 19]. We did not specifically address the function of this second set of sensory neurons, but did investigate the possible interactions between mdn-LOs, ILP7-neurons, and the

tdc2-expressing octopaminergic neurons. Using GRASP technique to reveal synaptic partners, we could not demonstrate inputs from mdn-LO neurons to *tdc*-expressing neurons or vice versa. However, we found that OA applied to the abdominal neuromere induced [Ca^{2+}] increases in the axon terminations of the mdn-LO neurons. Since GRASP analysis could not visualize any synaptic contacts from *tdc2* neurons onto mdn-LOs, we suggest that OA acts after paracrine release (non-synaptic action) to modulate these sensory cells. By severing the axons of the *tdc2* neurons by cutting the nerve from the VNC to the oviduct, we could demonstrate that this OA action on mdn-LOs occurs locally in the VNC. Furthermore, using GRASP, we found that the *tdc2*-neurons are presynaptic to the ILP7 neurons. Hence, the *tdc2*-neurons, which are known triggers of ovulation [17,18,19], may also trigger activity in the ILP7-neurons. In summary, we found no evidence that ILP7- and *tdc2*-neurons signal synaptically to mdn-LO neurons, but *tdc2*-neurons may synapse onto ILP7-neurons and also modulate mdn-LOs by paracrine signaling. Thus, while OA released from *tdc2*-neurons is a main trigger of ovulation, [17,18,19], it may also be a signal to activate ILP7-neurons and to modulate the activity of mdn-LOs. Interestingly, the mdn-LOs do not provide mechanosensory feedback to *tdc2*-neurons, only the ILP7 neurons. Finally, the mechanistic role of the octopaminergic modulation of the mdn-LOs was not revealed in our study.

While our study delineates key elements of the regulatory ovulation pathway, the complete neuronal connectivity and state-dependent recruitment of additional premotor and interneuron partners remain to be described. For the octopaminergic neurons in the VNC that initiate ovulation, a recent screen of membrane receptors has identified regulatory inputs including ILP8, GABA, and SIFamide [59]. Similarly, single-cell transcriptomic datasets could be leveraged to search for candidate receptors and ion channels expressed by ILP7 neurons and mdn-LO neurons, thereby generating testable hypotheses for future functional validation. The emergence of comprehensive female adult VNC connectomic resources will further enable circuit-level tests of this working model by identifying additional mechanosensory inputs, premotor partners, and connectivity motifs associated with ILP7 neurons.

Notably, ILP7 neurons exhibit spontaneous rhythmic [Ca^{2+}] dynamics in *ex vivo* preparations, and these dynamics are influenced by mechanical feedback linked to egg movement. Establishing how this rhythmicity relates to contractions in the oviduct muscle *in vivo* will benefit from future experiments that directly assay phase relationships and causal coupling under intact physiological conditions.

The precise role of ILP7 in the ovulation pathway remains to be clarified since we only found a possible role in sustaining oviposition. Previous studies detected no effect on ovulation of *Ilp7*-knockdown in the ILP7-neurons [4,22]. However, overexpression of *Ilp7* in ILP7-neurons, or ubiquitously, led to flies that were more susceptible to laying eggs on a high sucrose medium than a regular medium [4]. It is not clear how ILP7 mediates this effect, but it should be noted that apart from the sex-specific ILP7-neurons, there are further ILP7-expressing neurons (in both sexes) in the VNC of flies that may play roles in oviposition-related behavior and its relation to gut function and nutritional state [22,60]. Two of these are ascending interneurons that

co-express short neuropeptide F (sNPF) and supply axons to the brain [60, 61], where they act on the insulin-producing cells (IPCs) to regulate insulin-signaling when flies ingest high-carbohydrate medium [62, 63]. While we only investigated the role of the efferent oviduct-innervating ILP7-neurons (with ILP7 as a putative co-transmitter of glutamate), it may be interesting for the future to determine whether these and the ascending ILP7/sNPF neurons are functionally coupled and thereby link nutrition and oviposition.

Finally, the contribution of ovulation-related sensory input to the ramping activity of OviDN neurons that initiate egg deposition [3] warrants further investigation. A possible hypothesis is that the initial mechanosensory signals generated as an egg enters the oviduct or uterus are relayed, via as-yet-unknown pathways, to the brain, thereby prompting the fly to initiate the oviposition evaluation program and triggering the onset of calcium ramping in OviDNs. During this ramping phase, OviDNs may integrate external environmental cues and internal state signals until activity reaches a threshold, at which point the oviposition motor sequence is engaged to complete egg deposition. Therefore, elucidating the primary sensory circuits that trigger the oviposition evaluation program may enable the development of a model linking interoceptive sensory inputs to behavioral decision-making.

881 Materials and methods

882 Animals

883   Unless otherwise mentioned, all fly stocks were raised in standard
884 cornmeal–molasses–agar medium maintained at 25 ˚C, 60% humidity, and a 12 hr:12
885 hr light: dark cycle. We used w^{1118} flies as the wild-type strain.

886 Key resources table

REAGENT or RESOURCE	Source or reference	Identifiers
Antibodies		
Goat anti-Rabbit Alexa 488	Invitrogen	Cat#A11008; RRID:AB_14...
Goat anti-mouse Alexa 488	Invitrogen	Cat#A28175; RRID:AB_2536161
Goat anti-mouse Alexa 555	Invitrogen	Cat# A21428; RRID:AB_2535849
Goat anti-mouse Alexa 647	Invitrogen	Cat# A21235; RRID:AB_2535804
Mouse monoclonal anti-Brp	Developmental Studies Hybridoma Bank	Cat# nc82; RRID:AB_2314...
Rabbit polyclonal anti-GFP	Invitrogen	Cat# A11122; RRID:AB_221569
Rabbit polyclonal anti-RFP	Invitrogen	
Mouse polyclonal anti-GFP	Invitrogen	
Chemicals, Peptides, and Recombinant Proteins		
4% PFA	Leagene	Cat# DF0134
Agarose	Invitrogen	Cat# 75510-019
Yeast	Angel	N/A
Propionic acid	MACKLIN	Cat# C14876266
Adenosine 50-triphosphate (ATP)	New England Biolabs	Cat# P0756S
VECTASHIELD® Antifade Mounting Medium	Vector Laboratories	H-1000-10
DAPI	Solarbio	Cat# C0060
Phalloidine	Cytoskeleton, Inc	Cat# PHDN1
PBS	Solarbio	Cat# P1022
ReadyProbes 2.5% Normal Goat Serum	Invitrogen	Cat# R37624
Octopamine	Sigma-Aldrich	Cat# O0250

Trehalose	Biosharp	Cat# BS106B
Glutamate	Sigma- Aldrich	W328512
protease	Sigma- Aldrich	P5147
collagenase	Sigma- Aldrich	C0130
Schneider's Drosophila Medium	BasalMedia	C210908

Experimental Models: Organisms/Strains		
w^{1118}	Bloomington Drosophila Stock Center	3605
tmc^1	Zhang et al., 2016	BDSC_66556; RRID:BDSC_66556
tmc^{GAL4}	Guo et al., 2016	Flybase: FBal0321088
UAS-tmc/Cyo; tmc^1/TM6B	Zhang et al., 2016	BDSC_66560; RRID:BDSC_66560
w;tmc-GAL4	Zhang et al., 2016	
w;;tmc-GAL4		
tmc^{LexA}	This paper	N/A
tmc-LexA	This paper	N/A
$Piezo^{ko}$	Kim et al., 2012	BDSC_58770; RRID:BDSC_58770
ppk^{ESB}	Laboratory of Wei Zhang, Tsinghua Uni	N/A
ppk^{ko} and $Piezo^{ko}$	Laboratory of Wei Zhang, Tsinghua Uni	N/A
$Piezo^{ko}$; tmc^1	Laboratory of Wei Zhang, Tsinghua Uni	N/A
UAS-mCD8::GFP; UAS-mCD8::GFP		
UAS-stinger, UAS-Tdtomato	Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS>Stop>myrGFP; 8xLexAop2-FlpL	Laboratory of Yufeng Pan, Southeast Uni	N/A
fru-LexA	Laboratory of Yufeng Pan, Southeast Uni	N/A
ppk-GAL4	Laboratory of	N/A

		Wei Zhang, Tsinghua Uni	
ppk-LexA		Laboratory of Wei Zhang, Tsinghua Uni	N/A
Piezo-GAL4		Bloomington Drosophila Stock Center	BDSC_59266
UAS-Kir2.1		Bloomington Drosophila Stock Center	Flybase:FBtp0125506
UAS-dTrpA1		Hamada et al., 2008	BDSC:26263
UAS-shibire		Klapoetke et al., 2014	N/A
UAS-CsChrimson		Klapoetke et al., 2014	N/A
vGlut-GAL80/Cyo;TM2/TM6B		Bussell et al., 2014	N/A
elav-GAL80;tmc-GAL4			
tmc-GAL4;ppk-GAL80			
UAS-Redstinger		Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS>Stop>dTrpA1; 8xLexAop2-FlpL		Laboratory of Yufeng Pan, Southeast Uni	N/A
w*; 10XUAS-IVS-mCD8::RFP attP18 (13XLexAop2-mCD8::GFP)su(Hw)attP8; P(nSyb-MKII::nlsLexADBDo)attP24/Cyo; M(UAS-p65AD::CaM}ZH-86Fb		Laboratory of Yufeng Pan, Southeast Uni	N/A
20xUAS-Gcamp7s		Laboratory of Yufeng Pan, Southeast Uni	N/A
w-;UAS-tdtomato,UAS-Gcamp6m/Cyo		Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS-P2X$_2$		Lima and Miesenbo¨ ck, 2005	N/A
LexAop-Gcamp6m			
LexAop-P2X$_2$		Gou et al., 2014	BDSC_76030; RRID:BDSC_76030
tdc2-LexA;44588		Laboratory of Zhefeng Gong,	N/A

	Zhejiang University	
Sp/Cyo;ILP7-GAL4/TM6B	Laboratory of Yufeng Pan, Southeast Uni	N/A
tdc2-LexA;ILP7-GAL4	This paper	N/A
tmc-LexA;ILP7-GAL4	This paper	N/A
LexAOP-nSyb-spGFP[1-10],UAS-CD4-spGFP[11]; MKRS/TM6B	Macpherson et al., 2015	BDSC_64315; RRID:BDSC_64315
yw,UAS-myrGFP,QUAS-mtdTomato,attp8;trans-Tango,attp40,tubp-FRT-QS/Cyo	Laboratory of Yufeng Pan, Southeast Uni	N/A
w-;UAS-ppk-RNAi attp40	TsingHua Fly Center	N/A
w-;;UAS-ppk-RNAi attp2	TsingHua Fly Center	N/A
w-;UAS-piezo-RNAi attp40	TsingHua Fly Center	N/A
w-;UAS-tmc-RNAi/cyo; MKRS/TM6B		N/A
Software and Algorithms		
FIJI		https://doi.org/10.1038/nme
Prism 9	GraphPad	RRID:SCR_002798
Photoshop	Adobe	RRID:SCR_014199
Illustrator	Adobe	RRID:SCR_010279
ZEN	Zeiss	N/A
Motic Images Plus3.0	Motic	N/A
Motic Images Devices	Motic	N/A
Other		
Egg-laying device	Gou et al., 2016	N/A
Anatomical tweezers	Dumont	Cat# 0209-5-PO

Genomic DNA and cDNA amplifications.

Genomic DNA was extracted using a buffer (0.1 M Tris-HCl, pH 9.0, 0.1 M EDTA; 1% SDS). In brief, 30 fruit flies were ground with steel beads in the extraction buffer and incubated at 65°C for 1 hour. The samples were then lysed by adding potassium acetate to a final concentration of 800 mM and incubated on ice for 2 hours. Precipitation was pelleted by centrifugation at 10,000g for 15 min, and the supernatant was collected. The same volume of ethanol was added to the supernatant at -20℃ for 2h. Precipitation was pelleted by centrifugation at 10,000g for 15 min, and the sediment was collected. The pellet was washed with ice-cold 70% ethanol once, and the pellet was dried before resuspended in DEPC water.

897   Total RNA Extraction was done using the TRizol reagent (Invitrogen, Carlsbad,
898   CA, USA) according to the manufacturer's instructions. The cDNA template used for
899   cloning was synthesized using the Biotech M-MLV reverse transcription kit and the
900   synthesized cDNA template was stored at -20°C

Egg-laying assays

Unless otherwise mentioned, all egg laying assays were performed with females in
darkness with temperature and humidity controlled at 25°C and 60%, respectively.
The experimental flies were prepared as previously described (Gou et al., 2016;
Yang et al., 2015; Gou et al., 2014). Briefly, 20 to 30 females of the appropriate
genotypes and 10 to 15 males of mixed genotypes were gathered into a single food
vial that was supplied with wet yeast paste. After 5-7 days, the food in the vial became
very chewed up by the larvae, and at this point, the females were well-fed but
deprived of egg-laying. Thus, they were ready to lay eggs when placed in our
egg-laying apparatus (Yang et al., 2015; Gou et al., 2016). We usually let females lay
eggs overnight (~14 hr). Groups to be compared were always run in parallel. Note that
regardless of the assays we performed, we always assayed at the level of single
animals - each data point on a graph denotes the outcome (number of eggs laid) of a
single female.

Assessing ovulation efficiency and egg-jamming ratio

We collected some females for ovary dissection before the egg-laying females were put in
the oviposition apparatus, and the remaining females had their ovaries dissected just after
the egg-laying experiment in the oviposition apparatus.
We assessed whether there is an ovulation defect by counting mature eggs inside the
ovaries of the females before and after the egg-laying assay The ovulation efficiency =
(Retained mature eggs before egg-laying - Retained mature eggs after egg-laying) /
Retained mature eggs before egg-laying.
Before dissection, we used carbon dioxide for brief anesthesia and severed the
connection between the VNC and the brain in the flies. This procedure facilitates
dissection and removes the brain's inhibitory effect on ovulation, allowing for autonomous
ovulation behavior in the flies, which aids in subsequent dissection and frequency
statistics of egg jamming.
The frequency of egg jamming is determined by dissecting female fruit fly ovaries and
noting the position of the eggs in the oviduct. We define egg jamming as the presence of
two eggs retained in the lateral oviducts. This definition includes the egg-jamming
phenotype caused by the inactivation/activation of *ILP7*-expressing neurons, which also
leads to an extra egg being retained in the main oviduct.

Ovary development measurement

934   We dissected and made images of the ovaries of virgin flies on d2AAE and d7AAE,
935   respectively. Using Motic Images Devices and Motic Images Plus 3.0 to capture images of
936   the ovaries. By counting the number of retained mature eggs and using FIJI to calculate
937   the ovary's area and the mature egg's length and width, we could determine whether the

938   oogenesis and ovary development process was aberrant.

dTrpA1 activation and shibire inactivation

940   Experimental flies were maintained at 22℃, cold anesthetized, and allowed to recover for at
941   least 30 min at 22℃. The egg-laying apparatus containing flies was then placed at
942   experimental temperatures (30℃ or 22℃) for 14 hours to record the egg-laying numbers.

CsChrimson activation

For optogenetic stimulation, test flies were collected within 1-3 d after eclosion and
transferred into a vial that was supplied with wet yeast paste containing 0.2 mM
all-trans-retinal (116–31–4, Sigma- Aldrich). The vials were covered by aluminum foil to
protect them from light for 3–4 days before the ovulation. As previously mentioned, we
placed the flies in a transparent oviposition device for egg-laying and stimulated the
animals with red light (620 nm, 0.03 mW/mm², Vanch Technology, Shanghai, China).
Unless otherwise noted, light stimulation was presented continuously throughout the
observation period. Light intensity was measured by placing an optical power meter
(PS-310 V2, Gentec, Canada) near the location of the glass slide.

Immunostaining and confocal imaging

We generally dissected the brain, VNC, and ovary of four- to eight-day-old females,
except for trans-Tango experiments, in which flies were reared at 20℃ on days 15-20.
Dissection was in PBS buffer, and tissues were fixed in 4% PFA solution for ~25 min at
room temperature. Samples were then washed four times for 10 minutes each with PAT3
wash buffer (0.5% Triton X-100, 0.5% bovine serum albumin in PBS). The tissues were
transferred to a block buffer (ReadyProbes™ 2.5% Normal Goat Serum) and incubated at
room temperature for 60 minutes. Primary antibodies were added to the samples and
incubated for 4h at room temperature. Samples were then incubated overnight at 4°C.
Samples were washed four times for 10 minutes each and then incubated with secondary
antibodies at room temperature for 4h. Samples were washed four times, each for 10
minutes. If DAPI staining was required, the samples were incubated at room temperature
with DAPI (1:1000, Solarbio, Cat# C0060) for 30 minutes. Finally, samples were mounted
using an anti-fade mounting medium (Vector Laboratories).
Titers for primary antibodies were as follows: mouse anti-nc82 (Developmental
Studies Hybridoma Bank nc82, 1:1000), mouse anti-GFP (1:1000, A11120, Thermo Fisher
Scientific), rabbit anti-GFP (1:1000, A11122, Thermo Fisher Scientific), and rabbit
anti-RFP (1:1000, 600-401-379, Rockland), rabbit anti-TMC (1:250). Secondary
antibodies were Donkey Alexa 488 anti-rabbit, Donkey Alexa 488 anti-mouse, Donkey
Alexa 555 anti-rabbit, and Donkey Alexa 555 anti-mouse (1:500, R37118, R37114,
R37119, R37115, Thermo Fisher Scientific).
Samples were imaged at 10X，20X, or 40X magnification and 1,024 X 1,024 resolution
or more on Zeiss LSM980 confocal microscopes (Jena, Germany) at Nanjing Agricultural
University. Images were processed with FIJI or ZEN.
For GRASP, we did not stain the samples with anti-GFP and imaged the fixed and
washed samples directly.

For Tdtomato and Stinger, we did not stain the samples with anti-GFP and imaged
the fixed and washed samples directly.
For oversized images (such as Fig2J, Fig2S3), we used the Fiji plugin:
Stitching-Pixelwise Stitching of Images to process the image stitching.

Antibody generation

We selected the last 24 amino acids of the TMC protein
(C-DPRSASPEPTVNIIRIDIENEHEK) as the peptide antigen for peptide synthesis and
conjugated it to a KLH carrier for immunization. The synthesized peptide antigen was
injected into rabbits to generate antibodies (ABclonal antibodies). Drosophila TMC protein
antibodies were obtained by affinity purification of the antiserum, and the specificity of the
antibodies was verified via Dot-Blot(DB) testing against the naked peptide before
experiments. For immunolabeling, the TMC antibody was diluted at a 1:250 ratio.

TRIC

We collected virgin flies with the genotype w*,10XUAS-IVS-mCD8::RFP attP18,
{13XLexAop2-mCD8::GFP}su(Hw)attP8; P{nSyb-MKII::nlsLexADBDo}attP24/tmc-GAL4;
M{UAS-p65AD::CaM}ZH-86Fb/+ that were enclosed for 5 days and dissected them,
separating those that laid eggs and those that did not lay any eggs (number of eggs =
0). After dissection, without staining, we immediately observed the TRIC signal of
mdn-LO neurons in real time under the same parameters. We used GFP/RFP in each
cell body as the TRIC signal.

Calcium imaging

For in vitro experiments, VNC and ovaries were dissected in *Drosophila* Adult
Hemolymph-Like Saline (AHLS) buffer. The cuticle and connective tissue covering the
VNC and ovary were removed using fine forceps. The median abdominal nerve,
connecting the abdominal neuromere and the reproductive oviduct, was either kept intact
or carefully severed, depending on the experiment.
Calcium imaging was performed on a Zeiss 980 confocal microscope (Jena,
Germany) equipped with 10× or 20× objectives. Briefly, cells were scanned with 488 nm
and 555 nm lasers at 512 × 512 pixels (or lower) and an 8-bit dynamic range to acquire
baseline GCaMP or tdTomato fluorescence. Regions of interest (ROIs) were selected in
FIJI, and fluorescence changes were quantified as $\Delta F/F_0 = (F_{peak} - F_0)/F_0$, where F_0 is the
mean fluorescence over 30 frames in mdn-LO neurons expressing GCaMP. For *ilp7* or
tdc2 neurons, F_0 was defined as the average fluorescence over 30 frames before solution
application. Depending on the experiment, solutions were applied as follows: 10 µL AHLS
buffer,10 µL of 100 mM ATP, and 10 µL of 1 mM octopamine in 90 µL AHLS; and 20 µL of
50 mM glutamate in 80 µL AHLS.
For positioning the egg within the oviduct, we used an ex vivo "brain-less" preparation.
Using forceps, we rapidly displaced the egg to different locations along the oviduct and,
after each displacement, promptly refocused and began recording one or more bouts (with
brief pauses between bouts to restart acquisition). This allowed us to capture how

mechanosensory feedback arising from changes in egg position modulates the activity of
ILP7 neurons.

Single-nucleus RNA-sequencing analyses in *ilp7* neurons.

We queried the Fly Cell Atlas (FCA, https://flycellatlas.org/) and focused on the Body (10×,
Stringent) dataset of *Drosophila melanogaster* (ASAP). We performed a three-gene
co-expression screen: (1) *nSyb* was used as a neuronal marker to restrict the analysis to
VNC neurons; (2) *Ilp7* was used to identify *Ilp7*-positive cells; and (3) each candidate gene
from our curated list was queried within that set. Cells co-expressing *nSyb* and *Ilp7* were
operationally defined as *Ilp7* neurons. For each candidate gene, we recorded its
expression within these Ilp7 neurons and mapped the common co-expression to five FCA
cell coordinate codes: (−13.14, 0.14), (−13.34, 0.24), (−14.19, −1.32), (−15.20, −0.17),
(−15.37, −1.57). Gene expression levels are reported as Norm_1_asap_seurat values as
provided by the FCA portal.

Quantification and statistical analysis

We used the GraphPad Prism6 software package to generate graphs and analyze
data statistically. We tested whether the values were normally distributed using
D'Agostino–Pearson omnibus and Shapiro–Wilk normality tests before performing
statistical analysis. When data were normally distributed, we used parametric tests; when
data were not normally distributed, we used non-parametric tests. All tests were two-tailed.
All data are presented as mean ± s.e.m. We also consistently labeled the sample numbers
directly on graphs.

Sample size

1041   Our results were based on an analysis of behavior at the level of single animals.
1042 Sample sizes were determined before experimentation based on both variance and effect
1043 sizes. All the data are biological replicates.
1044

1045   Acknowledgments

1046   This research was funded by the the National Natural Science Foundation of
1047   China (32022011 & 32472542), the National Key Research and Developmental
1048   Program of China (2022YFD1700200) and the Fundamental Research Funds for
1049   Central Universities of the Central South University (KJJQ2025014).

1050

References

1. Cury KM, Axel R. Flexible neural control of transition points within the egg-laying behavioral sequence in *Drosophila*. *Nat Neurosci* **26**, 1054–1067 (2023).

2. Oliveira-Ferreira C, Gaspar M, Vasconcelos ML. Neuronal substrates of egg-laying behaviour at the abdominal ganglion of *Drosophila melanogaster*. *Scientific Reports* **13**, 21941 (2023).

3. Vijayan V*, et al.* A rise-to-threshold process for a relative-value decision. *Nature* **619**, 563–571 (2023).

4. Yang CH, Belawat P, Hafen E, Jan LY, Jan YN. Drosophila egg-laying site selection as a system to study simple decision-making processes. *Science* **319**, 1679 (2008).

5. Deady LD, Sun J. A follicle rupture assay reveals an essential role for follicular adrenergic signaling in *Drosophila* ovulation. *PLOS Genetics* **11**, e1005604 (2015).

6. Hu Q, Wolfner MF. The *Drosophila* Trpm channel mediates calcium influx during egg activation. *Proceedings of the National Academy of Sciences* **116**, 18994-19000 (2019).

7. Dickson BJ. Wired for Sex: The Neurobiology of *Drosophila* Mating Decisions. *Science* **322**, 904-909 (2008).

8. Gou B, Liu Y, Guntur Ananya R, Stern U, Yang C-H. Mechanosensitive neurons on the internal reproductive tract contribute to egg-laying-induced acetic acid attraction in *Drosophila*. *Cell Rep* **9**, 522-530 (2014).

9. Greenspan RJ, Ferveur J-F. Courtship in *Drosophila*. *Annual Review of Genetics* **34**, 205-232 (2000).

10. Hall JC. The mating of a fly. *Science* **264**, 1702 (1994).

11. Kubli E. Sex-peptides: seminal peptides of the *Drosophila* male. *Cell Mol Life Sci* **60**, 1689-1704 (2003).

12. Rezával C, Pavlou Hania J, Dornan Anthony J, Chan Y-B, Kravitz Edward A, Goodwin Stephen F. Neural circuitry underlying *Drosophila* female postmating behavioral responses. *Curr Biol* **22**, 1155-1165 (2012).

13. Yamamoto D, Koganezawa M. Genes and circuits of courtship behaviour in

1095   *Drosophila* males. *Nat Rev Neurosci* **14**, 681 (2013).

1096

1097   14.   Wang K*, et al.* Neural circuit mechanisms of sexual receptivity in *Drosophila*
1098   females. *Nature* **589**, 577–581 (2020).

1099

1100   15.   Wang F*, et al.* Neural circuitry linking mating and egg laying in *Drosophila*
1101   females. *Nature* **579**, 101-105 (2020).

1102

1103   16.   Nallasivan MP, Singh DND, Saleh MSRS, Soller M. Sex-peptide targets
1104   distinct higher order processing neurons in the brain to induce the female
1105   post-mating response. *eLife* **13**, RP98283 (2024).

1106

1107   17.   Middleton CA, Nongthomba U, Parry K, Sweeney ST, Sparrow JC, Elliott CJH.
1108   Neuromuscular organization and aminergic modulation of contractions in the
1109   *Drosophila* ovary. *BMC Biology* **4**, 17 (2006).

1110

1111   18.   White MA, Chen DS, Wolfner MF. She's got nerve: Roles of octopamine in
1112   insect female reproduction. *J Neurogenet* **35**, 132-153 (2021).

1113

1114   19.   Deshpande SA*, et al.* Regulation of *Drosophila* oviduct muscle contractility by
1115   octopamine. *iScience* **25**, 104697 (2022).

1116

1117   20.   Lee H-G, Seong C-S, Kim Y-C, Davis RL, Han K-A. Octopamine receptor
1118   OAMB is required for ovulation in *Drosophila melanogaster*. *Developmental*
1119   *Biology* **264**, 179-190 (2003).

1120

1121   21.   Rodríguez-Valentín R, López-González I, Jorquera R, Labarca P, Zurita M,
1122   Reynaud E. Oviduct contraction in *Drosophila* is modulated by a neural
1123   network that is both, octopaminergic and glutamatergic. *Journal of Cellular*
1124   *Physiology* **209**, 183-198 (2006).

1125

1126   22.   Castellanos MC, Tang JCY, Allan DW. Female-biased dimorphism underlies a
1127   female-specific role for post-embryonic Ilp7 neurons in *Drosophila* fertility.
1128   *Development* **140**, 3915 (2013).

1129

1130   23.   Zhang YV, Aikin TJ, Li Z, Montell C. The basis of food texture sensation in
1131   *Drosophila. Neuron* **91**, 863-877 (2016).

1132

1133   24.   Coste B*, et al.* Piezo1 and Piezo2 are essential components of distinct
1134   mechanically activated cation channels. *Science* **330**, 55-60 (2010).

1135

1136   25.   Coste B*, et al.* Piezo proteins are pore-forming subunits of mechanically
1137   activated channels. *Nature* **483**, 176-181 (2012).

1138

26. Guo Y*, et al.* Transmembrane channel-like (tmc) gene regulates *Drosophila* larval locomotion. *Proc Natl Acad Sci U S A* **113**, 7243-7248 (2016).

27. Hyunjin L, Hyun Woo C, Chen Z, Zee Yong P, Young Joon K. A pair of oviduct-born *pickpocket* neurons important for egg-laying in *Drosophila melanogaster. Mol Cells* **39**, 573-579 (2016).

28. Wu SF, Ja YL, Zhang YJ, Yang CH. Sweet neurons inhibit texture discrimination by signaling TMC-expressing mechanosensitive neurons in *Drosophila. eLife* **8**, e46165 (2019).

29. Lam RM*, et al.* PIEZO2 and perineal mechanosensation are essential for sexual function. *Science* **381**, 906-910 (2023).

30. Yamanouchi HM, Tanaka R, Kamikouchi A. Piezo-mediated mechanosensation contributes to stabilizing copulation posture and reproductive success in *Drosophila* males. *iScience* **26**, 106617 (2023).

31. Qin J, Yang T, Li K, Liu T, Zhang W. Pharyngeal mechanosensory neurons control food swallow in *Drosophila melanogaster. eLife*, 1-24 (2023).

32. Jia YL*, et al.* A mechanosensory receptor TMC regulates ovary development in the brown planthopper *Nilaparvata lugens. Frontiers in Genetics* **11**, 1152 (2020).

33. Häsemeyer M, Yapici N, Heberlein U, Dickson BJ. Sensory neurons in the *Drosophila* genital tract regulate female reproductive behavior. *Neuron* **61**, 511-518 (2009).

34. Court R*, et al.* A systematic nomenclature for the *Drosophila* ventral nerve cord. *Neuron* **107**, 1071-1079.e1072 (2020).

35. Kimura K-i, Hachiya T, Koganezawa M, Tazawa T, Yamamoto D. Fruitless and doublesex coordinate to generate male-specific neurons that can initiate courtship. *Neuron* **59**, 759-769 (2008).

36. Pan Y, Meissner GW, Baker BS. Joint control of *Drosophila* male courtship behavior by motion cues and activation of male-specific P1 neurons. *Proc Natl Acad Sci U S A* **109**, 10065-10070 (2012).

37. Ji X, Wang L, Li X, Li S, Liu S, Pan Y. Hormonal control of *fruitless* expression and male sexual orientation in *Drosophila. The Innovation Life* **2**, 100060 (2024).

1183   38. Sweeney ST, Broadie K, Keane J, Niemann H, O'Kane CJ. Targeted
1184   expression of tetanus toxin light chain in *Drosophila* specifically eliminates
1185   synaptic transmission and causes behavioral defects. *Neuron* **14**, 341-351
1186   (1995).
1187

1188   39. Ma J, Ptashne M. The carboxy-terminal 30 amino acids of GAL4 are
1189   recognized by GAL80. *Cell* **50**, 137-142 (1987).
1190

1191   40. Lee T, Luo L. Mosaic analysis with a repressible cell marker for studies of gene
1192   function in neuronal morphogenesis. *Neuron* **22**, 451-461 (1999).
1193

1194   41. Hamada FN*, et al.* An internal thermal sensor controlling temperature
1195   preference in *Drosophila*. *Nature* **454**, 217-220 (2008).
1196

1197   42. Yang C-H*, et al.* Control of the postmating behavioral switch in *Drosophila*
1198   females by internal sensory neurons. *Neuron* **61**, 519-526 (2009).
1199

1200   43. Bussell Jennifer J, Yapici N, Zhang Stephen X, Dickson Barry J, Vosshall
1201   Leslie B. Abdominal-B neurons control *Drosophila* virgin female receptivity.
1202   *Curr Biol* **24**, 1584-1595 (2014).
1203

1204   44. Richard D, Maria G, Catherine C, Aaron D. Visualizing glutamatergic cell
1205   bodies and synapses in *Drosophila* larval and adult CNS. *Journal of
1206   Comparative Neurology* **508**, 131-152 (2008).
1207

1208   45. Klapoetke NC*, et al.* Independent optical excitation of distinct neural
1209   populations. *Nat Meth* **11**, 338-346 (2014).
1210

1211   46. Hoopfer ED, Jung Y, Inagaki HK, Rubin GM, Anderson DJ. P1 interneurons
1212   promote a persistent internal state that enhances inter-male aggression in
1213   Drosophila. *eLife* **4**, e11346 (2015).
1214

1215   47. Gao XJ, Riabinina O, Li J, Potter CJ, Clandinin TR, Luo L. A transcriptional
1216   reporter of intracellular Ca2+ in *Drosophila*. *Nature Neuroscience* **18**, 917-925
1217   (2015).
1218

1219   48. Chen T-W*, et al.* Ultrasensitive fluorescent proteins for imaging neuronal
1220   activity. *Nature* **499**, 295-300 (2013).
1221

1222   49. Dana H*, et al.* High-performance calcium sensors for imaging activity in
1223   neuronal populations and microcompartments. *Nat Methods* **16**, 649-657
1224   (2019).
1225

1226   50. Lima SQ, Miesenböck G. Remote control of behavior through genetically

targeted photostimulation of neurons. *Cell* **121**, 141-152 (2005).

51.   Yao Z, Macara AM, Lelito KR, Minosyan TY, Shafer OT. Analysis of functional neuronal connectivity in the *Drosophila* brain. *J Neurophysiol* **108**, 684-696 (2012).

52.   Talay M*, et al.* Transsynaptic mapping of second-order taste neurons in flies by trans-Tango. *Neuron* **96**, 783-795.e784 (2017).

53.   Rezával C, Nojima T, Neville Megan C, Lin Andrew C, Goodwin Stephen F. Sexually dimorphic octopaminergic neurons modulate female postmating behaviors in *Drosophila*. *Curr Biol* **24**, 725-730 (2014).

54.   Li Y, Fink C, El-Kholy S, Roeder T. The octopamine receptor octß2R is essential for ovulation and fertilization in the fruitfly *Drosophila melanogaster*. *Arch Insect Biochem Physiol* **88**, 168-178 (2015).

55.   Lim J*, et al.* The octopamine receptor Octβ2R regulates ovulation in *Drosophila melanogaster*. *PLoS ONE* **9**, e104441 (2014).

56.   Macpherson LJ*, et al.* Dynamic labelling of neural connections in multiple colours by trans-synaptic fluorescence complementation. *Nat Commun* **6**, 10024 (2015).

57.   Baines RA, Uhler JP, Thompson A, Sweeney ST, Bate M. Altered electrical properties in *Drosophila* neurons developing without synaptic transmission. *The Journal of Neuroscience* **21**, 1523 (2001).

58.   Zhang Y*, et al.* PIEZO channels link mechanical forces to uterine contractions in parturition. *Science* **390**, eady3045 (2025).

59.   Yang M*, et al.* Female membrane proteins regulate postmating ovulation in <i>Drosophila melanogaster</i> by ovulin-dependent and -independent pathways. *Proceedings of the National Academy of Sciences* **122**, e2508783122 (2025).

60.   Miguel-Aliaga I, Thor S, Gould AP. Postmitotic specification of *Drosophila* insulinergic neurons from pioneer neurons. *PLOS Biology* **6**, e58 (2008).

61.   Nässel DR, Enell LE, Santos JG, Wegener C, Johard HAD. A large population of diverse neurons in the Drosophilacentral nervous system expresses short neuropeptide F, suggesting multiple distributed peptide functions. *BMC Neuroscience* **9**, 90 (2008).

1271 62. Cognigni P, Bailey AP, Miguel-Aliaga I. Enteric neurons and systemic signals
1272   couple nutritional and reproductive status with intestinal homeostasis. *Cell
1273   Metabolism* **13**, 92-104 (2011).

1274

1275 63. Prince E, Kretzschmar J, Trautenberg LC, Broschk S, Brankatschk M.
1276   DIlp7-producing neurons regulate insulin-producing cells in *Drosophila.
1277   Frontiers in Physiology* **12**, 630390 (2021).

1278
1279
1280

Movie legend

Movie S1. The video shows that during lateral oviduct contractions, Ca^{2+} levels in mdn-LO neurons increase in phase with muscle contraction, indicating responsiveness to lateral oviduct activity.

Movie S2. With the median abdominal nerve (MAN) intact, Ca^{2+} in mdn-LO axons within the VNC region of the VNC rises markedly during lateral oviduct contractions.

Movie S3. Following transection of the MAN, Ca^{2+} in mdn-LO axons within the VNC region shows no significant increase during lateral oviduct contractions, indicating that mdn-LO conveys signals to the VNC via the MAN.

Movie S4. ATP stimulation of P2X2-expressing mdn-LO neurons produces a pronounced elevation of Ca^{2+} in *ilp7* neurons within the VNC and triggers strong contractions of the lateral oviduct musculature.

Movie S5. The oscillation of the female-specific post-embryonic ILP7 neurons.

Movie S6. The oscillation of the embryonic ILP7 neurons.

Movie S7. The oscillation of the dissociated ILP7 neurons.