Internal mechanosensory feedback modulates central pattern generation to coordinate ovulation in *Drosophila*

Shao-Cong Su (苏少聪)[1], Ning Zhang (张宁)[1], Chen-Yu Li (李晨雨)[1], Dick R. Nässel[3], Cong-Fen Gao (高聪芬)[1], Shun-Fan Wu (吴顺凡)[1,2*]

Affiliations:

[1] State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China

[2] College of Sciences, Nanjing Agricultural University, Nanjing 210095, China

[3] Department of Zoology, Stockholm University, Stockholm, Sweden

[*] Correspondence: wusf@njau.edu.cn

Abstract

Successful reproduction requires precise coordination of muscle contractions during ovulation and has been well investigated in insects. However, the neural mechanisms of the central pattern generator (CPG) orchestrating this process remain poorly understood. Using *Drosophila*, we identify a novel pair of multidendritic mechanosensory neurons (mdn-LO) in the lateral oviducts that express the mechanosensitive channels TMC (transmembrane channel-like protein), PPK (pickpocket), and Piezo. We demonstrate that TMC is essential for ovulation coordination: *tmc* mutation or neuron-specific knockdown significantly increases egg-jamming at the junction between the lateral and the common oviduct, while *ppk* or *Piezo* single mutants reduce egg-laying without causing jamming. Calcium imaging reveals that mdn-LO neurons are rhythmically activated by oviduct contractions, and chemogenetic stimulation of these neurons triggers muscle contraction. Trans-Tango and GRASP (GFP Reconstitution Across Synaptic Partners) analyses show that mdn-LO neurons form synapses with efferent insulin-like peptide 7 (ILP7)-expressing motor neurons in the abdominal ganglion. Activating either mdn-LO or ILP7 neurons induces egg-jamming, indicating their roles within a CPG network. This circuit, including the TMC-expressing mdn-LO neurons and a set of ILP7 neurons, organizes an innate behavior by coupling strategic sensory information and a CPG to ensure bilateral coordination of oviduct contractions to prevent egg jamming during oviposition.

Introduction

Successful reproduction relies on the precise coordination of muscle contractions during ovulation and oviposition, a process orchestrated by central pattern generators (CPGs), neural circuits that generate rhythmic motor outputs. However, the CPG mechanisms governing ovulation remain poorly understood. In insects, which are commonly oviparous, the structure of the female reproductive system and the sequence of egg-laying events are conserved. It has been found that successful reproduction requires strict and precise regulation of events to occur in coordination, including mechanisms and behaviors associated with oogenesis, follicle cell rupture, motor control of reproductive oviduct muscles to ensure ovulation, egg activation, the choice of egg-laying site, and egg deposit (1-6).

Drosophila has emerged as a genetically tractable organism well suited for analysis of reproductive behavior, and the neuronal circuitry involved in mating and post-mating events has been intensely studied (7-16). *Drosophila* is also a favorable organism to model the motor control of ovulation and oviposition (4, 8, 17, 18). Although *Drosophila* virgins can ovulate, mating is commonly a trigger (18). During copulation and transfer of sperm and seminal fluid to the *Drosophila* female also sex peptide and a few other factors are also transferred and act on sensory neurons in the oviduct that signal to target neurons in the brain to induce a change in physiology and behavior of the fly, including the onset of ovulation and oviposition (11, 19, 20). A mated *Drosophila* female can lay eggs up to 80 times in a day (2), each time following a stereotyped sequence. Thus, oviposition comprises ovulation, the transfer of a mature egg from the ovary to the uterus where fertilization occurs, and then the deposition of eggs after a thorough search for an optimal egg-laying site (1, 4, 17, 21). After each egg is laid, the female goes through a grooming program and then rests before the next egg-laying cycle (1, 4).

In *Drosophila,* the mechanisms by which the mature egg is transferred from the ovary to be deposited by the ovipositor are as follows. Contractions in circular muscle in the peritoneal sheath around the ovary push the egg towards the base of the ovary and these contractions are amplified by octopamine (OA) (17, 18). The entrance of the

egg into one of the lateral oviducts is enabled by the rupture of the posterior mature follicle cells surrounding the oocyte, which is triggered by the activation of matrix metalloproteinase 2 due to octopamine acting on the OAMB receptor ([5](#)). Further movement through the lateral and common oviducts is caused by OA modulation of contractions in the myogenic muscle in the reproductive tract ([17](#), [18](#), [21](#), [22](#)). Thus, OA acts on two different OA receptors (OAMB and Octβ2R) to cause a series of contractions and relaxations of the muscle in the oviducts to move the egg forward. The source of OA acting on the oviduct muscle is a small set of efferent neurons in the abdominal ganglion (ABG) expressing tyramine decarboxylase (TDC), a biosynthetic enzyme for tyramine and OA ([17](#), [18](#), [21-23](#)), whereas the origin of OA causing follicle cell rupture is not known. When the egg enters the common oviduct, it triggers the contraction of its muscle.

In addition to the regulatory effects of OA on oviduct contractions, a group of female-specific glutamatergic motor neurons located in the ABG, which co-express insulin-like peptide 7 (ILP7), also control the contraction of the lateral oviduct and the common oviduct ([4](#), [8](#), [24](#)). Silencing of these ILP7 neurons causes jamming of eggs at the junction between the common and lateral oviducts ([4](#), [8](#), [24](#)). These papers suggest that the signal to the oviduct muscle is mediated by glutamate and found no role for the co-localized ILP7 in muscle contractions.

The contractions of the common oviduct can be sensed by mechanosensory neurons that express pickpocket (ppk) in the common oviduct, and this has been shown to influence the oviposition preference for acetic acid ([8](#)). However, it remains unclear which neurons are responsible for sensing the contraction of the lateral oviduct. Further down the common oviduct, a pair of posterior uterine sensory neurons (PU neurons) sense the egg's passage into the ovipositor ([1](#)). Another pair of motor neurons (circular muscle of the uterus; CMU neurons), possibly with feedback from PU neurons, finally activate the egg expulsion, which is followed by grooming and then a new cycle of ovulation ([1](#)).

This ovulation process relies on the highly coordinated activity of the muscles in the reproductive tract. Typically, only one egg is present in the reproductive tract at a

time, and only one egg can pass through the lateral oviduct at a time, indicating a

coordinated movement between the oviducts to ensure the smooth passage of the

egg. As mentioned above, inhibiting female-specific *ILP7* neurons leads to blockage

of the egg at the junction between the lateral and common oviducts, referred to as egg

jamming ([4](#)). However, it is still unclear how the coordinated transport of the eggs into

the common oviduct is controlled in *Drosophila*, but a central pattern generator (CPG)

is likely involved. Additionally, ovulation in insects like locusts is driven by a

brain-independent CPG in the abdominal nerve cord ([25-27](#)), but the CPG circuit for

Drosophila ovulation, including how sensory feedback integrates with motor neurons

like ILP7-expressing cells, has not been determined.

Here, we address these gaps by investigating the neural circuits and

mechanosensory feedback mechanisms underlying ovulation CPG function in

Drosophila. We identify a novel pair of multidendritic mechanosensory neurons in the

lateral oviducts (designated mdn-LO) that express three evolutionarily conserved

mechanosensitive channels: TMC (transmembrane channel-like protein), PPK

(pickpocket), and Piezo, along with the sex-determination gene doublesex (distinct

from adjacent fruitless-expressing multidendritic neurons). We show that TMC is

uniquely required for ovulation coordination, unlike PPK or Piezo. TMC mutation or

neuron-specific knockdown causes severe egg-jamming. Calcium imaging confirms

that mdn-LO neurons are rhythmically activated by lateral oviduct contractions,

consistent with a mechanosensory role. Using trans-Tango and GRASP analyses, we

demonstrate that mdn-LO neurons form functional connections with ILP7-expressing

motor neurons in the ABG. Activation of either mdn-LO or ILP7 neurons induces

egg-jamming, indicating their integration into the ovulation CPG. Taken together, our

findings outline a novel TMC-mdn-LO-ILP7 circuit that ensures bilateral coordination

of oviduct contractions to prevent egg-jamming during ovulation.

Results

Loss of the transmembrane channel-like (*tmc*) gene leads to egg-jamming in

Drosophila

The paired oviducts are the conduits between the ovaries and the uterus and are essential for ensuring optimal ovulation. Typically, only one egg can pass through the oviducts into the uterus at a time in *Drosophila*, indicating that internal sensory mechanisms coordinate this process. Regarding internal sensation within the female reproductive system, existing reports are limited to *ppk*-expressing neurons in the main oviduct that enhance *Drosophila*'s preference for acetic acid by sensing oviduct contractions ([8](#)), as well as posterior uterine (PU) sensory neurons that are activated by egg progression ([1](#)). However, how internal sensory mechanisms within the oviduct coordinate the process of egg transport from the two lateral into the unpaired uterus (to avoid egg jamming) remains largely unexplored.

In addition to Pickpocket (PPK), also PIEZO and Transmembrane Channel-like (TMC) proteins have been identified in the female reproductive system of *Drosophila*, though their functional roles remain unknown ([8](#), [28](#)). In *Drosophila*, PPK, PIEZO, and TMC are evolutionarily conserved mechanosensitive ion channels ([8](#), [28-33](#)). These channels have been implicated in the regulation of sexual behavior ([34](#), [35](#)), oviposition preference ([33](#)) (Wu et al., 2019b), and feeding behavior ([28](#), [36](#)).

Given our previous finding that *Nltmc3*, a TMC ortholog in brown planthopper (BPH), is essential for normal ovary development and oviposition ([37](#)), we sought to investigate whether the *Drosophila* ortholog *tmc* (CG46121) plays a similar role. Using

tmc mutant flies *tmc¹* ([28](#)), we conducted oviposition assays and found that egg-laying

was significantly reduced compared to controls (Fig. 1A).

Because reduced egg-laying can result either from defects in ovarian

development or disruptions in ovulation, we first examined whether developmental

effects contributed to the phenotype. To rule out effects from mating, we used virgin

female flies (for other experiments, unless noted otherwise, mated females were

used). We assessed ovarian development in virgin *tmc¹* mutants by measuring ovary

size and found no developmental abnormalities, in contrast to BPHs ([37](#)). Instead, we

observed significantly enlarged ovaries in *tmc¹* flies at both day 2 and day 7

post-eclosion (d2AAE and d7AAE, respectively) compared to w^{1118} controls (Fig. 1B,

D). The enlarged ovaries were due to the accumulation of mature eggs, suggesting a

severe ovulation defect in these unmated flies (Fig. 1C, D), which was not attributable

to abnormalities in egg morphology (Fig. 1—figure supplement 1).

To further confirm the role of TMC in ovulation, we reintroduced the *tmc* gene into

tmc¹ mutants using the *tmc-GAL4* driver line ([28](#)). The reduced oviposition, enlarged

ovaries, and accumulation of mature eggs observed in *tmc¹* mutants were all rescued

by expressing *UAS-tmc* under the *tmc* promoter (Fig. 1A–D).

We next dissected mated *tmc* mutant females to assess ovulation. Interestingly,

we occasionally observed egg-jamming at the junction where the two lateral oviducts

merge into the common oviduct (Fig. 1E and Fig. 1F, respectively). The frequency of

this blockage was significantly higher in *tmc* mutants than in controls. Importantly, the

reintroduction of the *tmc* gene in *tmc¹* mutants eliminated this egg-jamming phenotype

(Fig.1G). Similarly, RNAi-mediated knockdown of *tmc* in *TMC-GAL4*-expressing cells

also resulted in egg-jamming (Fig. 1H–I).

We next explored whether other channels might contribute to oviposition and

ovulation regulation in *Drosophila* (Figure 1—figure supplement 2). Using *ppk* mutants

(ppk^{ESB}) and *Piezo* mutants ($Piezo^{KO}$), we observed a significant reduction in

egg-laying in both strains after mating (Figure 1—figure supplement 2A). However, we

did not observe increased egg-jamming in these mutants or in flies with

RNAi-mediated knockdown of *ppk* or *Piezo* in TMC-GAL4-expressing cells (Figure

1—figure supplement 2B–E).

These findings indicate that TMC, specifically in *tmc-GAL4*-labeled cells, plays a

key role in coordinating normal ovulation in *Drosophila*, whereas *Piezo* and PPK do

not. The disruption of TMC leads to a significantly elevated incidence of egg-jamming.

Figure 1

Figure 1. The *Tmc* channel is essential for ovulation and oviposition in

***Drosophila*.**

Tmc mutants lay fewer eggs and display ovulation defects. All experiments, except

those in Fig. 1B-D, used mated flies. Groups that share at least one letter are not

statistically distinguishable (*P* > 0.05), whereas groups with completely different

letters are statistically distinguishable (*P* < 0.05).

(**A**) The number of eggs laid per female of the w^{1118}, tmc^1 mutant, and

tmc-GAL4/UAS-tmc; tmc^1 (mutant with *tmc* selectively rescued in

tmc-GAL4-expressing cells) for 14 h at 25 °C. One-way ANOVA with Tukey's Multiple

Range Test.

(**B-C**) Quantification was performed on the ovary size and the number of mature eggs

retained in each pair of ovaries from w^{1118}, tmc^1, and $tmc\text{-}GAL4/UAS\text{-}tmc;tmc^1$ flies.

The results show that unmated tmc^1 mutant females retained more mature eggs in

their ovaries, resulting in larger ovaries. This phenotype can be rescued by

expressing tmc in neurons specifically marked by $tmc\text{-}GAL4$. Representative images

of the ovaries are shown in Figure 1D. One-way ANOVA with Tukey's Multiple Range

Test.

(**D**) Representative images of w^{1118}, tmc^1, $tmc\text{-}GAL4/UAS\text{-}tmc$; tmc^1 ovaries on d2AAE

(2 d after eclosion) and d7AEE (7 d after eclosion), from unmated flies.

(**E**) Representative images of w^{1118}, tmc^1, $tmc\text{-}GAL4/UAS\text{-}tmc$; tmc^1/tmc^1 ovaries.

White arrows mark the egg in the lateral oviduct.

(**F**) Illustration of egg-jamming in the lateral oviduct.

(**G**) The egg-jamming ratio. Genotypes as in Fig 1E. Fisher's exact test.

(H) Representative images of $w\text{-};;tmc\text{-}GAL4/+$, $tmc\text{-}RNAi/+;tmc\text{-}GAL4/+$ ovaries.

White arrows mark the egg in the lateral oviduct.

(I) The egg-jamming ratio. Genotypes as in Fig. 1H. Fisher's exact test.

Figure 1—figure supplement 1

Figure 1 — figure supplement 1. The *tmc* gene does not affect egg shape.

(**A**) Representative images of eggs of w^{1118} flies and the tmc^{GAL4} mutant.

(**B- C**) The egg length and width of the w^{1118} and tmc^{GAL4} mutants. Groups that share

at least one letter are not statistically distinguishable ($P > 0.05$), whereas groups with

completely different letters are statistically distinguishable ($P < 0.05$); Student's t-test.

Figure 1—figure supplement 2

Figure 1 - figure supplement 2. Egg jamming is not affected by the manipulation

of other mechanosensory channel genes.

(**A**) Number of eggs laid per female of w^{1118}, $Piezo^{ko}$, ppk^{ESB} flies for 14 h at 25 °C.

One-way ANOVA with Tukey's Multiple Range Test. In this and the panels **C and E**:

groups that share at least one letter are not statistically distinguishable ($P > 0.05$),

whereas groups with completely different letters are statistically distinguishable ($P <$

0.05).

(**B**) Representative images of w^{1118}, $Piezo^{ko}$, ppk^{ESB} ovaries.

(**C**) The egg-jamming ratio. Genotypes as in **B**. Fisher's exact test.

(**D**)   Representative   images   of   *w-;;tmc-GAL4/+,   ppk-RNAi/+;   tmc-GAL4/+,*

Piezo-RNAi/+; tmc-GAL4/+ ovaries.

(**E**) The egg-jamming ratio. Genotypes as in **D**. Fisher's exact test.

Tmc is localized in multi-dendritic neurons in the lateral oviduct

Whereas PPK and PIEZO have been previously localized to neurons of the lateral

oviducts ([8](#)), the presence of TMC in the reproductive system was hitherto not known.

Thus, we set out to examine the detailed expression of *tmc-GAL4*.

To further determine the expression pattern of TMC, we generated a TMC

antiserum using a previously reported method ([31](#)). As revealed by

immunocytochemical labeling, TMC is localized in a pair of multi-dendritic neurons in

the lateral oviducts that we designate **m**ulti-**d**endritic **n**eurons in the **l**ateral **o**viduct,

mdn-LO (Figure 2A-B). Double-labeling experiments with TMC antibodies and

tmc-GAL4-driven *GFP* identified the mdn-LO neurons and thus confirmed the

expression of the channel protein (Figure 2A). In contrast, we did not observe

anti-TMC immunoreactivity in *tmc^1* mutant animals, confirming the specificity of the

TMC antibodies (Figure 2C). To acquire high-resolution images of the processes of

the *tmc-GAL4* labeled neurons, we expressed a membrane-bound green fluorescent

protein (*UAS-mCD8::GFP*) under the control of *tmc-GAL4.* We found that the

tmc-GAL4 specifically identified one mdn-LO neuron in each of the two bilaterally

symmetrical lateral oviducts (Figure 2D). TMC is expressed in the dendritic branches,

suggesting a sensory role (Figure 2D). Each of these *tmc* neurons covers the native

lateral oviduct (the one with the neuronal cell body) with their dendrites ([Figure 2D](#)).

We found no *tmc*-positive sensory neuron processes in the common oviduct or the

uterus (Figure 2D), where branches of PPK-positive neurons have been detected ([8](#),

[38](#)). We also investigated the neuronal expression of two *tmc-GAL4* lines (insertions

on the second and third chromosomes, respectively) and a *tmc-LexA* line in the

central nervous system. Similar to previous studies (28), we found no cell bodies in the

adult female ABG (Figure 2-supplemental file 1A-C), but there are many *tmc-GAL4*

positive neuronal processes projecting from peripheral tissues into the brain and ABG.

This is a VNC region dedicated to processing female reproductive sensory input from

the periphery and output to muscles involved in ovulation and egg-laying behavior (39)

(Figure 2-supplemental file 1A-C).

We also tested whether Piezo and PPK are expressed in neurons of the lateral

oviduct. Similar to previous studies (8), we observed both *Piezo-GAL4* and *PPK-GAL4*

expression in multidendritic neurons in each lateral oviduct (Figure 2-supplemental file

2A and B). Next, we asked whether TMC is co-expressed with PPK or PIEZO in

neurons of the lateral oviduct. Indeed, double labeling of *PPK-LexA* and *tmc-GAL4*

neurons revealed that the pair of TMC-expressing mdn-LO neurons is also

PPK-positive (Figure 2E). We then employed an intersectional strategy (40, 41) using

a combination of *tmc-GAL4*, *PPK-LexA,* and FLP recombinase to label mdn-LO

neurons and observed one pair of TMC and PPK co-expressing mdn-LO neurons

(Figure 2J). The axons of this pair of neurons project from the reproductive system to

the ABG region of the VNC (Figure 2J-K)

Using the same approach, we examined whether Piezo and PPK, TMC and Fru,

as well as TMC and Piezo, are co-expressed in neurons of the lateral oviduct (Figure

2F–I). Taken together, our results indicate that among the two pairs of multidendritic

neurons located in the lateral oviducts, one pair (mdn-LO) expresses all three

channels, TMC, PPK, and Piezo, as well as Dsx (Doublesex)—while the other pair

expresses only PPK and fru (Figure 2L) (for Dsx, see also Rezával et al., 2012).

Figure 2. Expression pattern of mechanosensory channels in mdn-LO neurons of the lateral oviducts

(A) Labeling of a lateral oviduct (lat. ovi.) with antiserum to TMC and anti-GFP staining from flies expressing the *UAS-mCD8::GFP* reporter under the control of the *tmc-GAL4*. mdn-LO neurons are indicated by arrows.

(B - C) Labeling of the lateral oviduct (lat. ovi.) using TMC antibodies in (B) w^{1118} control and (C) tmc^1 mutant. The arrow points to one labeled mdn-LO neuron.

(D) TMC is specifically expressed in a pair of multi-dendritic sensory neuron cells in the lateral oviduct. Even when the egg enters the lateral oviduct, the mdn-LO neuron dendrites are still wrapped around the lateral oviduct. *tmc* is not expressed in any other part of the female reproductive system: the common oviduct (com.ovi.), spermatheca (sp.) uterus, or parovaria (female accessory glands).

(E) Lateral oviduct neurons (mdn-LO neurons) that express both *tmc* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows).

(F) Lateral oviduct neurons (mdn-LO neurons) that express both *Piezo* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows).

(G) Lateral oviduct neurons that express *tmc* (indicated by green arrows) and *fruitless* (*fru*) (indicated by red arrows).

(H) Lateral oviduct neurons that express *Piezo* (indicated by green arrows) and *fru* (indicated by red arrows).

(I) Lateral oviduct neurons(mdn-LO neurons) that express both *Piezo* and *ppk* are revealed by a combinatorial strategy (indicated by yellow arrows).

(J) Expression of *tmc* and *ppk* expressing neurons in the brain, VNC, and lateral

oviducts (lat. ovi.), using the intersectional technique (*GAL4/LexA*). A distinct

abdominal ganglion (ABG) region is targeted by mdn-LO neurons from the lateral

oviducts. The genotype: *tmc-GAL4/UAS>stop>myrGFP;ppk-LexA/8xLexAop2-FlpL.*

Common oviduct (com. ovi). median abdominal nerve (MAN).

(K) Illustration of the morphology of the lateral oviduct *tmc-GAL4*-expressing neurons

(mdn-LO) in the female reproductive system and ABG.

(L) Summary illustration of the expression pattern of mechanosensory and other genes in

lateral oviduct neurons. *Dsx*: doublesex.

Figure 2—figure supplement 1

Figure 2—figure supplement 1. Expression patterns of different *tmc* drivers in

the brain, ventral nerve cord, and reproductive system.

(**A-B**) Expression patterns of *tmc-GAL4* (lines with insertion on different chromosomes)

expressing neurons in the brain, ventral nerve cord (VNC), and ovary.

(**C**) Expression of *tmc-LexA* expressing neurons (cell bodies indicated by white arrows)

in the oviduct.

321   Figure 2 —figure supplement 2

Figure 2—figure supplement 2. Expression of *Piezo* and *ppk* in CNS and

reproductive tract.

(**A-B**) Expression patterns of *ppk* and *Piezo* in the VNC (A1, B1) and female

reproductive system (**A2, B2**). (**A3, A4**) Expression patterns of *Piezo* in the ovary and

uterus. White arrowheads mark the lateral oviduct neurons.

The *tmc*-expressing mdn-LO neurons are required for normal ovulation

We selectively silenced the output of TMC-expressing neurons by using the

tmc-GAL4 to direct expression of the inward-rectifying potassium channel *Kir2.1* ([42])

and found that this manipulation causes animals to retain a similar number of eggs as

the *tmc[1]* mutants do (Figure 3A). The binding of GAL80 to the C-terminal 30 amino

acids of GAL4 can prevent GAL4-mediated transcriptional activation ([43]) and thus

inhibit the activity of GAL4 in *Drosophila* ([44]). We thus silenced *tmc-GAL4* neurons

with *Kir2.1* and examined the effect of *elav-GAL80* suppression of the

tmc-GAL4-mediated transgene expression in mdn-LO neurons. Interestingly,

tmc-GAL4 > Kir2.1 flies expressing *elav-GAL80* ([45]) laid almost normal numbers of

eggs compared to control flies, indicating that the *tmc-GAL4* effect on oviposition is

due to the mdn-LO neurons (Figure 3B). These results show that mdn-LO neurons are

necessary for normal ovulation. Furthermore, we asked whether silencing

tmc-expressing neurons also affects ovulation coordination in *Drosophila*. Indeed, we

did not observe eggs that were jammed in the two lateral oviducts (Figure 3C-D).

Figure 3

Figure 3. A pair of *Tmc*-expressing mechanosensory neurons (mdn-LO) in the

lateral oviduct is essential for ovulation.

(**A**) The number of eggs laid per female is reduced when the mdn-LO neurons are

selectively silenced by expression of the hyperpolarizing channel *Kir2.1* in the

tmc-GAL4 line. One-way ANOVA with Tukey's Multiple Range Test. In this and the

other figures: Groups that share at least one letter are not statistically distinguishable

($P > 0.05$), whereas groups with completely different letters are statistically

distinguishable ($P < 0.05$).

(**B)** The number of eggs laid per female of mdn-LO neurons selectively silenced, using

Kir2.1, in the presence and absence of suppression by *elav-GAL80*. The number of

eggs laid is restored by neuron-specific *elav-GAL80* suppression. One-way ANOVA

with Tukey's Multiple Range Test.

(**C**) The egg-jamming ratio of *UAS-Kir2.1/+, tmc-GAL4/+, tmc-GAL4/+;UAS-Kir2.1+,*

and *tmc-GAL4/elav-GAL80*;*UAS-Kir2.1/+* flies. Fisher's exact test.

(**D**) Representative images of ovaries of *UAS-Kir2.1/+, tmc-GAL4/+,*

362   *tmc-GAL4/+;UAS-Kir2.1/+,* and *tmc-GAL4/elav-GAL80*;*UAS-Kir2.1/+* flies.

Activation of *tmc*-expressing mdn-LO neurons inhibits oviposition and

normal ovulation

Next, we selectively activated the output of TMC-expressing neurons thermo-genetically by using the *tmc-GAL4* to direct the expression of dTrpA1 ([46]). Activation of *tmc*-GAL4 neurons via dTrpA1 severely impaired oviposition (Figure 4B) and induced a significant egg-jamming rate (Figure 4C and D). To clarify which of the *tmc-GAL4* neurons are functionally required for coordinated oviposition and avoidance of egg-jamming, we suppressed the activity of GAL4 in selected tissues and cell types using the GAL80 repressor ([44]). For this, we used two GAL80 lines, *elav-GAL80* ([45]) and vesicular glutamate transporter (*vGlut*)-*GAL80* ([47]). We found that *elav-GAL80* and *vGlut-GAL80* lines both block the expression of TMC in mdn-LO neurons in the lateral oviduct (Figure 4A, Figure 4—figure supplement 1). Given that the motor neurons controlling muscle movements in *Drosophila* are glutamatergic ([48]), *vGlut-GAL80* serves to repress any potential GAL4 miss-expression in motor neurons.

To determine if the behavioral abnormality was due to the mdn-LO neurons, we combined the *elav-GAL80* or *vGlut-Gal80* transgene with the *tmc-GAL4* and *UAS-dTrpA1* to suppress activation of mdn-LO neurons. We found that these animals showed no difference in egg-jamming compared with control flies, while *tmc-GAL4/UAS-dTrpA1* animals showed an increased egg-jamming ratio at 30℃ (Figure 4C and D).

We next used an intersectional strategy to visualize the overlapping expression of

PPK-LexA   and   *tmc-GAL4*   (using   *tmc-GAL4/UAS>stop>myrGFP;*

ppk-LexA/LexAop2-FlpL) to obtain a more precise target for our manipulations of

mdn-LO neurons. However, in these flies, we observed only one of the two mdn-LO

neurons (Figure 4—figure supplement 2A and figure supplement 3), possibly due to

the weak cleavage efficiency of the flp in the neurons. When we activated these

tmc/PPK-coexpressing neurons using the dTrpA1 we found that these females

(*tmc-GAL4/UAS>stop>dTrpA1; ppk-LexA/LexAop2-FlpL*) laid more eggs at 30°C than

at 20°C (Figure 4—figure supplement 2B). In addition, we observed no ovulation

deficiency and egg-jamming when we activated this single mdn-LO neuron (Figure

4—figure supplement 2C-D). These results indicate that a single mdn-LO neuron in

one of the lateral oviducts is sufficient for ovulation.

To further confirm our findings, we used the optogenetic effector CsChrimson (49),

which affords greater control over the dynamic range of neuronal activation than

dTrpA1 (50), and found that optogenetic stimulation of *tmc-GAL4* labeled neurons in

UAS-CsChrimsonVenus/tmc-GAL4 females abolished oviposition, dramatically

different from all control females [see also (33)]. We also observed a significant

egg-jamming rate in the activated *UAS-CsChrimsonVenus/tmc-GAL4* females (Figure

4E and F). Combining pan-neuronal expression of GAL80 with

UAS-CsChrimsonVenus/tmc-GAL4 significantly reduces the rate of egg-jamming

seen after photoactivation (Figure 4F).

Figure 4

Figure 4. Thermogenetic and optogenetic activation of the pair of

***tmc*-expressing mdn-LO neurons in the lateral oviduct jams the passage of**

eggs through the lateral oviduct.

In these graphs, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001, and ns

(non-significant). Groups that share at least one letter are not statistically

distinguishable (P > 0.05), whereas groups with completely different letters are

statistically distinguishable ($P < 0.05$).

(**A**) Using the nuclear-localized RFP (UAS-Redstinger) and the membrane-localized

GFP (UAS-mCD8::GFP; UAS-mCD8::GFP) as a marker, we investigated the labeling

of mdn-LO neurons in the lateral oviducts in the presence of vGlut-GAL80 and

elav-GAL80 strains. Both vGlut-GAL80 and elav-GAL80 can effectively block the

tmc-GAL4 labeling of lateral oviduct neurons (mdn-LO) but not neurons in the CNS.

Arrowheads indicate the mdn-LO neurons in the lateral oviducts.

(**B**) Thermogenetic activation of tmc neurons affects oviposition. The number of eggs

laid per female with *UAS-dTrpA1/+, tmc-GAL4/UAS-dTrpA1, elav-GAL80;tmc-GAL4/*

UAS-dTrpA1, and *tmc-GAL4,vGlut-GAL80/+*; *UAS-dTrpA1/+* in 22°C and 30°C.

One-way ANOVA with Tukey's Multiple Range Test.

(**C**) Representative images of ovaries from *UAS-dTrpA1/+, tmc-GAL4/UAS-dTrpA1,*

elav-GAL80;tmc-GAL4/ UAS-dTrpA1, and *tmc-GAL4,vGlut-GAL80/+; UAS-dTrpA1/+*

flies at 22°C and 30°C.

(**D**) The egg-jamming ratio of *UAS-dTrpA1/+, tmc-GAL4/UAS-dTrpA1,*

elav-GAL80;tmc-GAL4/ UAS-dTrpA1, and *tmc-GAL4,vGlut-GAL80/+; UAS-dTrpA1/+*

flies. Fisher's exact test

(**E**)   Representative   images   of   *UAS-CsChrimsonmVenus*/+   and

tmc-GAL4/+;*UAS-CsChrimsonmVenus/+* ovaries after optogenetic activation. The

arrow indicates the location where the egg is jamming in the oviduct.

(**F**) Optogenetic activation of *tmc-GAL4* neurons using CsChrimson. The egg-jamming

ratio after optogenetic activation of *tmc-GAL4* neurons. Fisher's exact test.

Figure 4 —figure supplement 1

Figure 4—figure supplement 1. Expression of tmc-GAL4 in the presence of

different GAL80 in the brain and ventral nerve cord.

Using the membrane-localized GFP (*UAS-mCD8::GFP; UAS-mCD8::GFP*) as a

marker, we investigated the labeling of mdn-LO neurons in the CNS in the presence

of *vGlut-GAL80* and *elav-GAL80* strains.

Figure 4 —figure supplement 2

Figure 4—figure supplement 2. Activating mdn-LO neurons on one side only

promotes oviposition but does not induce egg-jamming.

In these graphs, groups that share at least one letter are not statistically

distinguishable (P >0.05), whereas groups with different letters are statistically

distinguishable (P <0.05).

(**A**) Using the intersectional technique (*tmc-GAL4/UAS>stop>myrGFP; ppk-LexA/8*

× LexAop-FlpL) to selectively target mdn-LO neurons revealed the expression of a

single mdn-LO neuron in the lateral oviduct on one side only.

(**B-C**) Number of eggs laid per female before (22°C) and after thermogenetic (30°C)

activation (*TrpA1*) of a single mdn-LO neuron in the lateral oviduct, along with

images of the ovaries. Two-tailed t-tests.

(**D**) The calculated egg-jamming ratio at 22°C and 30°C of one mdn-LO neuron in

the lateral oviduct. Fisher's exact test.

Figure 4 —Figure supplement 3

tmc-GAL4/UAS>stop>GFP;ppk-LexA/8xLexAop-FlpL

Figure 4—figure supplement 3. Labeling of mdn-LO neuron on one side

Using the intersectional technique (*tmc-GAL4/UAS>stop>myrGFP; ppk-LexA/8 ✗ LexAop-FlpL*) was used to specifically target mdn-LO neurons. However, in all cases, it only labeled a single mdn-LO neuron in the lateral oviduct on one side. (n=6).

472   **_tmc_-expressing sensory neurons in the reproductive tract can sense**

473   **contractions of the lateral oviduct tract**

Our experiments, so far suggest that the female flies sense the passage of eggs

through the reproductive tract by means of mechanosensory neurons. Thus, we next

asked whether the mdn-LO neurons monitor oviduct muscle contractions and

oviposition.

To investigate whether oviposition and ovulation affect activity in mdn-LO neurons,

we conducted an assay using a transcriptional reporter of intracellular Ca^{2+} (TRIC)

that monitors sustained neuronal activity (51). Indeed, we found a significant elevation

of TRIC signals, corresponding to increased intracellular Ca^{2+} levels, in mdn-LO

neurons of egg-laying virgin flies compared with virgin flies that did not lay eggs,

monitored over 5 days (Figure 5A and A').

Next, to further explore the requirement of mdn-LO neurons for ovulation-induced

neuronal activity, we monitored Ca^{2+} dynamics in the cell body of mdn-LO neurons in

response to ovulation. To visualize changes in Ca^{2+} levels in cell bodies of mdn-LO,

we expressed a transgene encoding the Ca^{2+} sensor GCaMP6m (_UAS-GCaMP6m_)

and Tdtomato (_UAS-Tdtomato_) (52) in mdn-LO neurons under the control of the

tmc-GAL4. We surgically exposed the lateral oviduct region of the female reproductive

area, aiming to monitor the fluorescence changes of mdn-LO neurons as the egg

passed through the lateral oviduct. However, we found that when the egg passes

through the lateral oviduct, an inevitable shift occurs along the Z-axis, causing us to

lose the signal. Even with the addition of tdTomato as a negative control, the

significant displacement still prevented us from accurately monitoring the true calcium

fluorescence signal of mdn-LO under the condition of the egg being squeezed through

the oviduct. Therefore, we monitored the activity of mdn-LO neurons when the oviduct

undergoes spontaneous contractions. We observed that calcium activity in the cell

body of the mdn-LO neuron increased when the muscle contracted (Figure 5B and

Movie S1).

We also used another calcium sensor GCaMP7s (53) to monitor activity in axonal

termini of mdn-LO neurons in the VNC. This was done since the GCaMP7s signal in

tmc processes in the VNC is less likely to move out of the focal plane when tract

contraction occurs, and thus facilitates measurements. We found that the CGaMP7s

signal in the axonal termini of *tmc* neurons increased significantly during oviduct

contraction (Figure 5C-C''; Movie S2). To ensure that such GCaMP7 increase was not

due to "local interaction" between *tmc* and *other* neurons in the VNC, we physically

severed the connection (median abdominal nerve, MAN) between the reproductive

tract and the VNC. In these severed preparations, we did not observe a significant

GCaMP7s increase in *tmc* axons in the VNC during oviduct contractions (Figure

5D-D''; Movie S3), ruling out the possibility that other neurons locally activate *tmc*

axons in the VNC. Thus, our results suggest that the *tmc*-expressing mdn-LO neurons

of the oviduct are mechanosensory and can detect contraction/distention of the lateral

oviduct.

We next asked whether the mdn-LO neurons receive input from sources other

than   mechanical   stimulation   during   the   ovulation   process.   The   insect

neurotransmitter/neuromodulator octopamine is considered a trigger for ovulation in

insects (18), so we tested the effect of octopamine application on mdn-LO neurons.

We found that octopamine activates calcium in the axon terminations of GCaMP6m

expressing mdn-LO neurons both in intact animals and after surgically severing the

MAN and thus the connection to the oviduct (Figure 5 —figure supplement 1). This

suggests that octopamine modulates activity in the mdn-LO neurons locally in the

VNC.

Finally, we used a chemical-genetic approach (54, 55) to stimulate the mdn-LO

neurons. We expressed the ATP-gated P2X2 channel, which is not expressed

endogenously in flies and can be used for 'chemogenetic activation' of specific

neurons of interest. When stimulating the P2X2 expressing mdn-LO neurons with ATP,

a robust contraction of the oviduct muscle was induced (Movie S4). This shows that

mdn-LO mechanosensory neurons in the lateral oviduct can not only sense oviduct

contraction but also act through downstream motoneurons to regulate contractions of

the lateral oviduct.

Figure 5

Figure 5. The lateral oviduct *tmc*-expressing neurons (mdn-LO) are mechanosensory and are activated during ovulation.

(**A**) Monitoring sustained calcium activity in mdn-LO neurons using the TRIC method. Comparison of TRIC signals of mdn-LO sensory neurons between egg-laying and non-egg-laying females and representative images. We collected flies that were 5 days post-eclosion and dissected them, separating those that laid eggs and those that did not lay any eggs. After dissection, without staining, we immediately observed the TRIC signal of mdn-LO in real time under the same parameters (we used GFP/RFP as the TRIC signal). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$, and ns (non-significant); Two-tailed t-tests.

(**B**) When the lateral oviduct contracts rhythmically, it is accompanied by rhythmic excitation of mdn-LO sensory neurons, as shown by increased Ca^{2+} levels monitored with Gcamp6m.

(**C- C"**) In intact animals (median abdominal nerve, MAN, intact), lateral oviduct mdn-LO sensory neurons transmit excitation to the abdominal ganglion (ABG), and Ca^{2+} activity in axon terminations of the left and right neurons can be monitored.

(**D- D"**) When the MAN is surgically severed, no Ca^{2+} activity is detected in axon terminations in the ABG during spontaneous muscle contractions. This supports that the activation in intact animals is derived from the axons of the lateral oviduct mdn-LO sensory neurons.

Figure 5 —figure supplement 1

Figure 5—figure supplement 1. Mdn-LO neurons can be activated by

octopamine in intact animals and after severing the connection to the

reproductive tract.

(A-A'') *tmc*+ neurons can be activated by octopamine (A) Illustration of maintaining the

connection via the median abdominal nerve (MAN) and cutting MAN. (A') Calcium

imaging of mdn-LO neuron axons in the ABG region before and after the addition of 1

mM octopamine in intact animals and after severing the MAN. (A'') Changes in

fluorescence intensity ($\triangle F/F_0$) of Gcamp6m in mdn-LO neuron axons in the ABG

region before and after octopamine addition.

ILP7-expressing neurons in the abdominal ganglion receive synaptic input from tmc-expressing mdn-LO neurons

We next set out to identify the potential mechanism by which the TMC-expressing mechanosensitive neurons act on neuronal circuits that control oviduct muscle contractions.

First, therefore, we sought to map the postsynaptic targets of mdn-LO neurons by visualization of functionally connected neurons using trans-Tango (56). We found that expression of trans-Tango in mdn-LO neurons labeled many post-synaptic neurons, primarily in the abdominal ganglion (ABG) (Figure 6A-C'') and suboesophageal ganglion (SOG) (Figure 6—figure supplement 1A). We, however, did not observe that expression of trans-Tango in mdn-LO neurons labeled any post-synaptic local neurons in the ovary or oviduct (Figure 6—figure supplement 1B). This finding provides further evidence that mdn-LO neurons are sensory neurons that transfer signals to neurons in the central nervous system. Since the trans-Tango labeled numerous neurons, the identification of individual post-synaptic neurons was hard. It has been shown previously (24), and we confirmed here, that some of the *Ilp7-GAL4*-expressing neurons in the ventral portion of the posterior VNC are motor neurons that innervate the oviduct (Figure 6D, Figure 6—figure supplement 2H). Importantly, inhibiting *ILP7* neurons consistently causes one or more eggs to be jammed in the oviduct (4). Furthermore, efferent neurons in the VNC that produce octopamine, identified by a *tdc2-GAL4,* and applied octopamine, are also involved in the regulation of ovulation (22, 57-59). Thus, we next used the syb:GRASP technique

(60) to address the question of whether axons of *tmc*-expressing mdn-LO neurons

directly contact processes of *Ilp7*-expressing or *tdc2*-expressing neurons in the

abdominal ganglion (ABG). In this method, half of the GFP is targeted specifically to

the presynaptic termini of the mdn-LO neurons and the other to *Ilp7* or *Tdc* neurons,

and GFP reconstitution occurs at exposure to the synaptic cleft only upon synaptic

vesicle fusion. With *Ilp7* neurons and *Tdc* neurons separately as partners, we were

able to detect reconstituted GFP clearly in the ABG (Figure 6E-H; Figure 6—figure

supplement 2B, D), suggesting that mdn-LO neurons can act as presynaptic partners

of both *Ilp7* neurons and *Tdc* neurons in the ABG. However, we did not find evidence

with the GRASP technique that *Ilp7* or *tdc2* neurons are coupled pre-synaptically to

mdn-LO neurons (Figure 6—figure supplement 1C and 1E). Thus, the activation of

mdn-LO neurons by applied octopamine, shown in Figure 5—figure supplement 1, is

likely to represent paracrine (non-synaptic) signaling from octopaminergic neurons.

Furthermore, we found that *tdc2*-GAL4 and *Ilp7*-GAL4 signals are not co-expressed in

neurons in the ABG or the brain (Figure 6—figure supplement 1I). However, we could

show by GRASP that *tdc2* neurons are presynaptically coupled to *Ilp7* neurons,

suggesting functional connections between these neurons (Figure 6—figure

supplement 1F-G).

To probe whether the neuronal contacts we observed using the GRASP

technique permit the mdn-LO neurons to transmit signals to the *Ilp7*-expressing

neurons, we next assessed if we could detect a change in Ca^{2+} levels in the cell

bodies and processes of *Ilp7* neurons when we stimulated the mdn-LO neurons. To do

this, we developed an *ex vivo* "brain-less" preparation in which the entire ventral nerve

cord is dissected together with the full female reproductive tract and intact

connections of the nerve supply between the VNC and the reproductive system (but

with the connection to the brain is severed). In this *in vitro* preparation, we monitored

the Ca^{2+} levels in cell bodies and the processes of *Ilp7*-expressing neurons by using

the genetically encoded Ca^{2+} indicator GCaMP6m (52). To stimulate mdn-LO neurons,

we expressed in these neurons the ATP-gated P2X2 channel (54, 55).

We found that stimulating the P2X2-expressing mdn-LO neurons with ATP

triggered a significant increase in GCaMP signal in the cell bodies of *Ilp7* neurons in

the ventral ABG (Figure 6I-O), accompanied by rhythmic contractions of the oviduct

(Movie S4). In contrast, exposing control animals that did not express P2X2 in their

mdn-LO neurons to the same levels of ATP did not produce an increased GCaMP

signal (Figure 6J). Interestingly, we did not detect an increased GCaMP signal in *Ilp7*

neurons located on the dorsal side of the ABG region, only in the ventral cluster,

including the female-specific *Ilp7* neurons (Figure 6L). Collectively, our results show

that axons of mdn-LO neurons physically contact the *Ilp7* neurons in the ABG, and

stimulation of mdn-LO neurons increases Ca^{2+} levels in the cell bodies and processes

of ventrally located *Ilp7* neurons (Figure 6M-O).

Figure 6. The mdn-LO sensory neurons regulate activity in the *Ilp7* neurons in

the ABG, and their connections are visualized by trans-tango and GRASP.

(A-C'') Putative second-order neurons targeted by *tmc+* neurons (green) in the ABG

region revealed by trans-Tango (red) driven by *tmc*-GAL4.

In A-B', we show the expression of presynaptic signal (GFP/dark) in mdn-LO neurons

and postsynaptic signal (RFP/dark) of *tmc+* neurons in the ABG region of VNC.

In C-C'' are shown merged confocal images of trans-Tango labeling of postsynaptic

neurons (GFP/green) to *tmc+* neurons (RFP/red).

(D) Location of *Ilp7* neurons in the ventral and dorsal ABG regions. Blue circles

indicate embryonic *Ilp7* neurons; pink circles indicate large female-specific

post-embryonic *Ilp7* neurons; Green circles indicate the post-embryonic *Ilp7* neurons.

Original data from ([24](#)). The three types of *Ilp7* -expressing cell clusters innervate

different tissues. The embryonic *Ilp7* neurons innervate the intestine, and the

female-specific post-embryonic *Ilp7* neurons and post-embryonic *Ilp7* neurons are

glutamatergic motoneurons that innervate the oviduct muscle of the female

reproductive system.

(E-H) GRASP signals between mdn-LO sensory neurons and *Ilp7* neurons observed

in the brain, dorsal and ventral VNC, and ovary.

(I-L) The mdn-LO sensory neurons regulate the excitation of ventral VNC *Ilp7* neurons

in the ABG region. Activating *P2X2*-expressing mdn-LO neurons with 1 mM ATP

increased the activity of ventral *Ilp7* motor neurons in the abdominal part of the VNC.

The dorsal *Ilp7* neurons are not affected (J).

(M-O) Graph showing activation of *Tmc+* neurons via *P2X2* increased different *Ilp7+*

656   motor neuron activity. Fluorescence changes ($\Delta F/F_0$) of *GCaMP6s* in *Ilp7* neurons are

657   shown following different treatments.

660   Figure 6 —figure supplement 1

Figure 6—figure supplement 1. The presumed second-order neurons of mdn-LO

target the subesophageal ganglion (SEG) region in the brain.

(A) Putative second-order neurons of *tmc*+ neurons (green) in the suboesophageal

ganglion (SOG) region revealed by trans-Tango (red).

(B) Using the trans-tango system, it was found that there are no putative second-order

neurons of mdn-LO in the oviduct.

Figure 6 —figure supplement 2

Figure 6—figure supplement 2. Probing connections between *tdc2+*, *Ilp7+*, and

***tmc+* neurons by the GRASP technique.**

(A) Lack of GRASP signal from expressing *tmc-LexA* alone in the brain and ventral

VNC.

(B) Flies in virgin status were used to obtain the GRASP signal between mdn-LO

(expressing *tmc-LexA*) and *Ilp7* + neurons with *Ilp7 -GAL4* in the brain, dorsal, and

ventral VNC. The black signal represents the region of putative synaptic connections

(indicated by black dashed lines).

(C) No GRASP signal is seen from the *Ilp7* + neurons to *tmc*+ neurons in the brain,

dorsal VNC, and ovary.

(D) GRASP signal from the *tmc*+ neurons to *tdc2*+ neurons in the dorsal and ventral

VNC. The dark signal represents the region of synaptic connections (indicated by

black dashed lines).

(E) No GRASP signal is seen from the *tdc2*+ neurons to *tmc*+ neurons in the ventral

VNC.

(F) GRASP signal from the *tdc2*+ neurons to *Ilp7* + neurons in the ventral and dorsal

VNC. The dark signal represents the region of putative synaptic connections

(indicated by black dashed lines).

(G) GRASP signal from the *Ilp7* + neurons to *tdc2*+ neurons in the ventral and dorsal

VNC. The dark signal represents the region of putative synaptic connections

(indicated by black dashed lines).

(H) The expression pattern of *Ilp7*-GAL4 positive neurons in the brain, VNC, and

ovary.

(I) *Ilp7*+ (green) and *tdc2*+ (red) neurons within the ABG and brain are distinct.

Activation of *Ilp7* neurons induces egg-jamming, and ILP7 release affects

oviposition

Although previous studies reported that hyperpolarization of *Ilp7-GAL4* neurons in the VNC with *Kir2.1* ([61](#)) consistently causes one or more eggs to be jammed in the oviduct due to defects in ovulation coordination ([4](#), [8](#), [24](#)), the effects of depolarizing these neurons and ILP7 peptide (Insulin-like peptide 7) on egg-laying and ovulation have not been investigated.

Therefore, we aimed to determine whether depolarization of *Ilp7-GAL4* neurons, induced by mdn-LO sensory neuron activity, results in disrupted ovulation coordination. Additionally, we sought to assess whether the ILP7 neuropeptide released upon *Ilp7-GAL4* neuron activation plays any role in egg laying and ovulation.

We found that activating the output of *Ilp7*-expressing neurons thermo-genetically by using the *Ilp7-GAL4* to direct the expression of dTrpA1 lead to decreased oviposition and increased egg jamming (Fig. 7A-C). Optogenetic stimulation of *Ilp7-GAL4*-labeled neurons in *UAS-CsChrimsonvenus/ILP7-GAL4* females induced egg-jamming phenotypes in all individuals (Figure [7D](#)).

When ILP7 levels were reduced by RNAi (*ilp7-GAL4>UAS-Ilp7*-RNAi) during TrpA1 activation of *Ilp7*-neurons, the number of eggs laid was partly restored compared to *Ilp7*-neuron activation alone (Figure 7A), although the frequency of egg jamming was not significantly reduced (Figure 7[B](#)), similar to the activation of mdn-LO neurons shown in our study. These findings suggest that the ILP7 neuropeptide released during *Ilp7*-neuron activation reduces egg laying, but does not significantly

718   impact the coordination of ovulation.

719   Our results indicate that the ovulation-coordinating ILP7 motor neurons are

720   modulated by sensory feedback from the mdn-LO mechanosensory neurons in the

721   lateral oviducts, thereby ensuring the coordination of the ovulation process.

 Figure 7

Female reproductive system

Figure 7. Activation of ILP7 neurons induces egg-jamming, and ILP7 release

affects oviposition

Groups that share at least one letter are not statistically distinguishable (P > 0.05),

whereas groups with completely different letters are statistically distinguishable ($P <$

0.05). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, and ns (non-significant).

(A) Number of eggs laid per female of the in flies with ILP7 neurons thermogenetically

activated and ILP7 peptide RNAi (*ILP7*-GAL4>UAS-dTrpA1,

ILP7-GAL4>UAS-dTrpA1/UAS-*ILP7*-RNAi) for 14 h at 22 °C and 30 °C. One-way

ANOVA with Tukey's Multiple Range Test.

(B) The egg-jamming ratio after thermogenetic activation of *ILP7*-GAL4 neurons and

Ilp7 RNAi. Genotypes as in **A**. Fisher's exact test.

(**C**) Representative images of ovaries of *ILP7*-GAL4>UAS-dTrpA1,

ILP7-GAL4>UAS-dTrpA1/UAS-*ILP7*-RNAi at 22 °C and 30 °C.

(**D**) The egg-jamming ratio is increased after optogenetic activation of *ILP7*-GAL4

neurons.

(**E**)   Representative   images   of   *UAS-CsChrimsonmVenus*/+   and

UAS-CsChrimsonmVenus/ILP7-GAL4 ovaries after optogenetic activation. The arrow

indicates the location where the egg is jamming in the oviduct.

(F) Graphic summary of the female reproductive tract and ventral nerve cord (VNC).

(G) Graphic summary of the motor neurons and mechanosensory neurons as part of

the ovulation central pattern generator circuit.

Discussion

Here, we have identified a novel role for the mechanosensory channel protein,

TMC, in a pair of multidendritic sensory neurons in the lateral oviducts that provide

feedback to motor circuits that ensure coordinated ovulation in *Drosophila*. These

neurons, which we designated mdn-LO, also express two other mechanosensory

channels, PPK and PIEZO. Mutations in all three channel proteins result in defects in

ovulation and oviposition, whereas mutation of *tmc* further increases the incidence of

egg jamming. This indicates that all three mechanosensitive channels are involved in

the mechanosensation of the lateral oviduct and that they function complementarily in

this sensory process. The mdn-LO neurons display rhythmic $[Ca^{2+}]$ oscillations during

spontaneous and induced contractions of the oviduct muscle, supporting a

mechanosensory role. Our experiments show that the mdn-LOs are presynaptic to a

set of efferent neurons in the abdominal ganglion that produce glutamate and the

neuropeptide ILP7, here designated *Ilp7*-neurons. We found that chemo-genetically

stimulating activity in the mdn-LOs leads to increased $[Ca^{2+}]$ in the *ILP7*-neurons and

rhythmic contractions in the oviduct muscle. Previous research has demonstrated that

activity in these two pairs of female-specific *Ilp7*-neurons is required for proper

ovulation, likely by using glutamate as a neuromuscular neurotransmitter ([4], [8], [22], [24]).

However, the function of ILP7 was not shown. Our study demonstrates that blocking

ILP7 peptide release during activation of ILP7 neurons increases oviposition in

Drosophila, but does not affect egg jamming. Thus, this peptide may act further down

in the common oviduct to modulate contractions (and oviposition). We could show that

activation or inactivation of the *Ilp7*-neurons leads to egg-jamming in the common

oviduct, whereas activation of mdn-LO neurons results in egg-jamming in the lateral

oviducts. Thus, our findings suggest that the two mdn-LO neurons provide sensory

feedback from each of the lateral oviducts to modulate alternating activity in the two

pairs of *Ilp7*-motor neurons in order to generate a smooth passage of eggs into the

common oviduct. Our data thus suggest that these *Ilp7*-neurons are part of the

ovulation CPG network that acts in concert with octopaminergic neurons in the control

of ovulation, as discussed below.

Apart from the *Ilp7* and mdn-LO neurons shown here and descending neuronal

pathways from the brain (3, 16), there are additional neurons in the control of ovulation

in *Drosophila*. These are a second pair of multidendritic sensory neurons in the lateral

oviducts that express PPK, but not the TMC and PIEZO channels (8), and a small set

of octopaminergic efferent neurons (*tdc*-neurons) in the abdominal ganglia that

innervate muscle of the reproductive tract (17, 18, 22). We did not specifically address

the function of this second set of sensory neurons but did investigate the possible

interactions between mdn-LOs, *Ilp7*-neurons, and the *tdc2*-expressing octopaminergic

neurons.

Using the GRASP technique to reveal synaptic partners, we could not

demonstrate inputs from mdn-LO neurons to *tdc*-expressing neurons or vice versa.

However, we found that OA applied to the abdominal ganglion induced $[Ca^{2+}]$

increases in the axon terminations of the mdn-LO neurons. Since GRASP analysis

could not visualize any synaptic contacts from *tdc2* neurons onto mdn-LOs, we

suggest that OA acts after paracrine release (non-synaptic action) to modulate these

sensory cells. By severing the axons of the *tdc2* neurons by cutting the nerve from the

abdominal ganglion to the reproductive tract, we could demonstrate that this OA

action on mdn-LOs occurs locally in the ganglion. Furthermore, using GRASP, we

found that the *tdc2*-neurons are presynaptic to the *Ilp7*-neurons, but did not find any

function of OA on these neurons. Thus, the *tdc2*-neurons, which are known triggers of

ovulation (17, 18, 22), may also trigger activity in the *Ilp7*-neurons. In summary, we

found no evidence that *Ilp7-* and *tdc2*-neurons signal synaptically to mdn-LO neurons,

but *tdc2*-neurons may synapse onto *Ilp7*-neurons and also modulate mdn-LOs by

paracrine signaling. The roles of this OA signaling were not studied. Thus, OA

released from *tdc2*-neurons is a main trigger of ovulation (17, 18, 22) but also a likely

signal to activate *Ilp7*-motor neurons and to modulate the activity of mdn-LOs.

Interestingly, the mdn-LOs do not provide mechanosensory feedback to *tdc2*-neurons,

only the *Ilp7*-neurons. Furthermore, we do not yet have data to clarify the mechanistic

role for the octopaminergic modulation of the mdn-LOs.

The specific role of the ILP7 peptide in the ovulation circuit remains to be tested;

we only found a possible role in sustaining oviposition. Previous studies detected no

effect on ovulation of *Ilp7*-knockdown in the *Ilp7*_neurons (4, 24). However,

overexpression of *Ilp7* in *Ilp7* -neurons, or ubiquitously, led to flies that were more

susceptible to laying eggs on a high sucrose medium than a regular medium (4). It is

not clear how ILP7 mediates this effect, but it should be noted that there are further

Ilp7-expressing neurons in the VNC of flies (24, 62)_that may play roles in

oviposition-related behavior and its relation to nutritional state.   Some of these are

interneurons that supply axons to the brain, where they act on the insulin-producing

cells (IPCs) to regulate insulin-signaling when flies ingest high-carbohydrate medium

(63, 64). Thus, it is necessary to specifically pinpoint the role of ILP7 in the efferent

neurons as a putative co-transmitter of glutamate.

TMC proteins have been identified in a range of invertebrates and as well as

mammals, including humans (31, 37, 65-68). There are eight TMC proteins in

mammals (66, 67), two in *Caenorhabditis elegans* (65), three in brown planthoppers

(37) , and only one in *Drosophila* (28, 31). The first function of a TMC was described in

humans and mice where intact TMC1 and TMC2 activity is necessary for cochlear

hair-cell function, and mutations in these genes cause deafness (67, 69-71). The TMC

proteins are pore-forming subunits of channels required for mechano-transduction

currents in hair cells (72). In *C. elegans,* the *tmc1* gene encodes a sodium-sensitive

cation channel that functions in salt gustation and mediates behavioral avoidance at

high NaCl concentrations (65). Furthermore, TMC-1 expressed in nociceptive neurons

of *C. elegans* mediates avoidance of alkalines (73). Interestingly, TMC proteins also

modulate egg-laying in nematodes (74, 75). Another study found that TMC-1 affects *C.*

elegans development and sexual behavior in a diet-dependent context (76). In

Drosophila, the *tmc* gene has been implicated in proprioception during larval

locomotion (31), sensing of food texture (28), and discrimination of the texture of

egg-laying site (33). Finally, in the brown planthopper *Nilaparvata lugens,* the

orthologous gene *Nltmc3* was found to be highly expressed in the female reproductive

tract, including the oviduct (37). Knockdown of the *Nltmc3* resulted in reduced egg

laying and impaired ovary development. Thus, the present study adds an important

role of TMC protein as a mechanosensory channel in mdn-LOs in the lateral oviducts

that sense the movement of the eggs and thereby provide feedback to motoneurons

that control oviduct muscle contractions. This proprioceptive feedback ensures

smooth passage of eggs into the common oviduct.

Our data suggest that the two other mechanosensory channels, PPK and PIEZO,

expressed in mdn-LO neurons, are also necessary for proper ovulation. Thus, the

delicate process of ovulation and oviposition requires both "belt and suspenders,", i. e.

several types of multidendritic neurons to ensure efficient proprioceptive feedback to

motor circuits in the VNC. However, it is not not yet clear what the contribution of each

of the channels is and why all three seem necessary for optimal ovulation.

Materials and methods

Animals

Unless otherwise mentioned, all fly stocks were raised in standard

cornmeal–molasses–agar medium maintained at 25 ˚C, 60% humidity, and a 12 hr:12

hr light: dark cycle. We used w^{1118} flies as the wild-type strain.

Key resources table

REAGENT or RESOURCE	Source or reference	Identifiers
Antibodies		
Goat anti-Rabbit Alexa 488	Invitrogen	Cat#A11008; RRID:AB_143165
Goat anti-mouse Alexa 488	Invitrogen	Cat#A28175; RRID:AB_2536161
Goat anti-mouse Alexa 555	Invitrogen	Cat# A21428; RRID:AB_2535849
Goat anti-mouse Alexa 647	Invitrogen	Cat# A21235; RRID:AB_2535804
Mouse monoclonal anti-Brp	Developmental Studies Hybridoma Bank	Cat# nc82; RRID:AB_2314866
Rabbit polyclonal anti-GFP	Invitrogen	Cat# A11122; RRID:AB_221569
Rabbit polyclonal anti-RFP	Invitrogen	
Mouse polyclonal anti-GFP	Invitrogen	
Chemicals, Peptides, and Recombinant Proteins		
4% PFA	Leagene	Cat# DF0134
Agarose	Invitrogen	Cat# 75510-019
Yeast	Angel	N/A
Propionic acid	MACKLIN	Cat# C14876266
Adenosine 50-triphosphate (ATP)	New England Biolabs	Cat# P0756S
VECTASHIELD® Antifade Mounting Medium	Vector Laboratories	H-1000-10
DAPI	Solarbio	Cat# C0060

Phalloidine	Cytoskeleton, Inc	Cat# PHDN1
PBS	Solarbio	Cat# P1022
ReadyProbes 2.5% Normal Goat Serum	Invitrogen	Cat# R37624
Octopamine	Sigma-Aldrich	Cat# O0250
Trehalose	Biosharp	Cat# BS106B
Glutamate	Sigma-Aldrich	W328512

Experimental Models: Organisms/Strains		
w^{1118}	Bloomington Drosophila Stock Center	3605
tmc^1	Zhang et al., 2016	BDSC_66556; RRID:BDSC_66556
tmc^{GAL4}	Guo et al., 2016	Flybase: FBal0321088
UAS-tmc/Cyo; tmc^1/TM6B	Zhang et al., 2016	BDSC_66560; RRID:BDSC_66560
w^-;tmc-GAL4	Zhang et al., 2016	
w^-;;tmc-GAL4		
tmc^{LexA}	This paper	N/A
tmc-LexA	This paper	N/A
$Piezo^{ko}$	Kim et al., 2012	BDSC_58770; RRID:BDSC_58770
ppk^{ESB}	Laboratory of Wei Zhang, Tsinghua Uni	N/A
ppk^{ko} and $Piezo^{ko}$	Laboratory of Wei Zhang, Tsinghua Uni	N/A
$Piezo^{ko}$; tmc^1	Laboratory of Wei Zhang, Tsinghua Uni	N/A

UAS-mCD8::GFP; UAS-mCD8::GFP

UAS-stinger, UAS-Tdtomato	Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS>Stop>myrGFP; 8xLexAop2-FlpL	Laboratory of Yufeng Pan, Southeast Uni	N/A
fru-LexA	Laboratory of Yufeng Pan, Southeast Uni	N/A
ppk-GAL4	Laboratory of Wei Zhang, Tsinghua Uni	N/A
ppk-LexA	Laboratory of Wei Zhang, Tsinghua Uni	N/A
Piezo-GAL4	Bloomington Drosophila Stock Center	BDSC_59266
UAS-Kir2.1	Bloomington Drosophila Stock Center	Flybase:FBtp0125506
UAS-dTrpA1	Hamada et al., 2008	BDSC:26263
UAS-shibire	Klapoetke et al., 2014	N/A
UAS-CsChrimson	Klapoetke et al., 2014	N/A
vGlut-GAL80/Cyo;TM2/TM6B	Bussell et al., 2014	N/A

elav-GAL80;tmc-GAL4		
tmc-GAL4;ppk-GAL80		
UAS-Redstinger	Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS>Stop>dTrpA1; 8xLexAop2-FlpL	Laboratory of Yufeng Pan, Southeast Uni	N/A
w*; 10XUAS-IVS-mCD8::RFP attP18 (13XLexAop2-mCD8::GFP)su(Hw)attP8; P(nSyb-MKII::nlsLexADBDo)attP24/Cyo; M(UAS-p65AD::CaM}ZH-86Fb	Laboratory of Yufeng Pan, Southeast Uni	N/A
20xUAS-Gcamp7s	Laboratory of Yufeng Pan, Southeast Uni	N/A
w-;UAS-tdtomato,UAS-Gcamp6m/Cyo	Laboratory of Yufeng Pan, Southeast Uni	N/A
UAS-P2X$_2$	Lima and Miesenbö̈ck, 2005	N/A
LexAop-Gcamp6m		
LexAop-P2X$_2$	Gou et al., 2014	BDSC_76030; RRID:BDSC_76030
tdc2-LexA;44588	Laboratory of Zhefeng Gong, Zhejiang University	N/A
Sp/Cyo;ILP7-GAL4/TM6B	Laboratory of Yufeng Pan, Southeast Uni	N/A
tdc2-LexA;ILP7-GAL4	This paper	N/A

tmc-LexA;ILP7-GAL4	This paper	N/A
LexAOP-nSyb-spGFP[1-10],UAS-CD4-spGFP[11]; MKRS/TM6B	Macpherson et al., 2015	BDSC_64315; RRID:BDSC_64315
yw,UAS-myrGFP,QUAS-mtdTomato,attp8;trans-Tango,attp40,tubp-FRT-QS/Cyo	Laboratory of Yufeng Pan, Southeast Uni	N/A
w-;UAS-ppk-RNAi attp40	TsingHua Fly Center	N/A
w-;;UAS-ppk-RNAi attp2	TsingHua Fly Center	N/A
w-;UAS-piezo-RNAi attp40	TsingHua Fly Center	N/A
w-;UAS-tmc-RNAi/cyo; MKRS/TM6B		N/A
Software and Algorithms		
FIJI		https://doi.org/10.1038/nmeth.2019
Prism 9	GraphPad	RRID:SCR_002798
Photoshop	Adobe	RRID:SCR_014199
Illustrator	Adobe	RRID:SCR_010279
ZEN	Zeiss	N/A
Motic Images Plus3.0	Motic	N/A
Motic Images Devices	Motic	N/A

Primer		
Primer1: GGGCACGTTTTGGCTTTGAT		
Primer2: TGCCTCGCTGTCCTTTGTAG		
Primer3: GAACACCATCGTGGGGAACT		
Primer4: AACAAGCGCAGCTGAACAAG		
Other		
Egg-laying device	Gou et al., 2016	N/A
Anatomical tweezers	Dumont	Cat# 0209-5-PO

Genomic DNA and cDNA amplifications.

Genomic DNA was extracted using a buffer (0.1 M Tris-HCl, pH 9.0, 0.1 M EDTA;

1% SDS). In brief, 30 fruit flies were ground with steel beads in the extraction buffer

and incubated at 65°C for 1 hour. The samples were then lysed by adding potassium

acetate to a final concentration of 800 mM and incubated on ice for 2 hours.

Precipitation was pelleted by centrifugation at 10,000g for 15 min, and the

supernatant was collected. The same volume of ethanol was added to the

supernatant at -20℃ for 2h. Precipitation was pelleted by centrifugation at 10,000g for

15 min, and the sediment was collected. The pellet was washed with ice-cold 70%

ethanol once, and the pellet was dried before resuspended in DEPC water.

Total RNA Extraction was done using the TRizol reagent (Invitrogen, Carlsbad,

CA, USA) according to the manufacturer's instructions. The cDNA template used for

cloning was synthesized using the Biotech M-MLV reverse transcription kit and the

synthesized cDNA template was stored at -20˚C

Egg-laying assays

Unless otherwise mentioned, all egg laying assays were performed with females

in darkness with temperature and humidity controlled at 25˚C and 60%, respectively.

The experimental flies were prepared as previously described (Gou et al., 2016;

Yang et al., 2015; Gou et al., 2014). Briefly, 20 to 30 females of the appropriate

genotypes and 10 to 15 males of mixed genotypes were gathered into a single food

vial that was supplied with wet yeast paste. After 5-7 days, the food in the vial became

very chewed up by the larvae, and at this point, the females were well-fed but

deprived of egg-laying. Thus, they were ready to lay eggs when placed in our

egg-laying apparatus (Yang et al., 2015; Gou et al., 2016). We usually let females lay

eggs overnight (~14 hr). Groups to be compared were always run in parallel. Note that

regardless of the assays we performed, we always assayed at the level of single

animals - each data point on a graph denotes the outcome (number of eggs laid) of a

single female.

883 Assessing ovulation efficiency and egg-jamming ratio

We collected some females for ovary dissection before the egg-laying females were

put in the oviposition apparatus, and the remaining females had their ovaries dissected

just after the egg-laying experiment in the oviposition apparatus.

We assessed whether there is an ovulation defect by counting mature eggs inside the

ovaries of the females before and after the egg-laying assay The ovulation efficiency =

(Retained mature eggs before egg-laying - Retained mature eggs after egg-laying) /

Retained mature eggs before egg-laying.

Before dissection, we used carbon dioxide for brief anesthesia and severed the

connection between the VNC and the brain in the flies. This procedure facilitates

dissection and removes the brain's inhibitory effect on ovulation, allowing for autonomous

ovulation behavior in the flies, which aids in subsequent dissection and frequency

statistics of egg jamming.

The frequency of egg jamming is determined by dissecting female fruit fly ovaries and

noting the position of the eggs in the oviduct. We define egg jamming as the presence of

two eggs retained in the lateral oviducts. This definition includes the egg-jamming

phenotype caused by the inactivation/activation of *ILP7*-expressing neurons, which also

leads to an extra egg being retained in the main oviduct.

901 Ovary development measurement

We dissected and made images of the ovaries of virgin flies on d2AAE and d7AAE, respectively. Using Motic Images Devices and Motic Images Plus 3.0 to capture images of the ovaries. By counting the number of retained mature eggs and using FIJI to calculate the ovary's area and the mature egg's length and width, we could determine whether the oogenesis and ovary development process was aberrant.

dTrpA1 activation and shibire inactivation

Experimental flies were maintained at 22 □C, cold anesthetized, and allowed to recover for at least 30 min at 22 □C. The egg-laying apparatus containing flies was then placed at experimental temperatures (30 □C or 22 □C) for 14 hours to record the egg-laying numbers.

CsChrimson activation

For optogenetic stimulation, test flies were collected within 1-3 d after eclosion and transferred into a vial that was supplied with wet yeast paste containing 0.2 mM all-trans-retinal (116–31–4, Sigma- Aldrich). The vials were covered by aluminum foil to protect them from light for 3–4 days before the ovulation. As previously mentioned, we placed the flies in a transparent oviposition device for egg-laying and stimulated the animals with red light (620 nm, 0.03 mW/mm², Vanch Technology, Shanghai, China). Unless otherwise noted, light stimulation was presented continuously throughout the observation period. Light intensity was measured by placing an optical power meter (PS-310 V2, Gentec, Canada) near the location of the glass slide.

Immunostaining and confocal imaging

We generally dissected the brain, VNC, and ovary of four- to eight-day-old females, except for trans-Tango experiments, in which flies were reared at 20℃ on days 15-20.

Dissection was in PBS buffer, and tissues were fixed in 4% PFA solution for ~25 min at

room temperature. Samples were then washed four times for 10 minutes each with PAT3

wash buffer (0.5% Triton X-100, 0.5% bovine serum albumin in PBS). The tissues were

transferred to a block buffer (ReadyProbes™ 2.5% Normal Goat Serum) and incubated at

room temperature for 60 minutes. Primary antibodies were added to the samples and

incubated for 4h at room temperature. Samples were then incubated overnight at 4°C.

Samples were washed four times for 10 minutes each and then incubated with secondary

antibodies at room temperature for 4h. Samples were washed four times, each for 10

minutes. If DAPI staining was required, the samples were incubated at room temperature

with DAPI (1:1000, Solarbio, Cat# C0060) for 30 minutes. Finally, samples were mounted

using an anti-fade mounting medium (Vector Laboratories).

Titers for primary antibodies were as follows: mouse anti-nc82 (Developmental

Studies Hybridoma Bank nc82, 1:1000), mouse anti-GFP (1:1000, A11120, Thermo Fisher

Scientific), rabbit anti-GFP (1:1000, A11122, Thermo Fisher Scientific), and rabbit

anti-RFP (1:1000, 600-401-379, Rockland), rabbit anti-TMC (1:250). Secondary

antibodies were Donkey Alexa 488 anti-rabbit, Donkey Alexa 488 anti-mouse, Donkey

Alexa 555 anti-rabbit, and Donkey Alexa 555 anti-mouse (1:500, R37118, R37114,

R37119, R37115, Thermo Fisher Scientific).

Samples were imaged at 10X ,20X, or 40X magnification and 1,024 X 1,024 resolution

or more on Zeiss LSM980 confocal microscopes (Jena, Germany) at Nanjing Agricultural

University. Images were processed with FIJI or ZEN.

For GRASP, we did not stain the samples with anti-GFP and imaged the fixed and

washed samples directly.

For Tdtomato and Stinger, we did not stain the samples with anti-GFP and imaged

the fixed and washed samples directly.

For oversized images (such as Fig2J, Fig2S3), we used the Fiji plugin:

Stitching-Pixelwise Stitching of Images to process the image stitching.

Antibody generation

We selected the last 24 amino acids of the TMC protein

(C-DPRSASPEPTVNIIRIDIENEHEK) as the peptide antigen for peptide synthesis and

conjugated it to a KLH carrier for immunization. The synthesized peptide antigen was

injected into rabbits to generate antibodies (ABclonal antibodies). Drosophila TMC protein

antibodies were obtained by affinity purification of the antiserum, and the specificity of the

antibodies was verified via Dot⬛Blot(DB) testing against the naked peptide before

experiments. For immunolabeling, the TMC antibody was diluted at a 1:250 ratio.

TRIC

We collected virgin flies with the genotype w*,10XUAS-IVS-mCD8::RFP attP18,

{13XLexAop2-mCD8::GFP}su(Hw)attP8; P{nSyb-MKII::nlsLexADBDo}attP24/tmc-GAL4;

M{UAS-p65AD::CaM}ZH-86Fb/+ that were enclosed for 5 days and dissected them,

separating those that laid eggs and those that did not lay any eggs (number of eggs =

0). After dissection, without staining, we immediately observed the TRIC signal of

mdn-LO neurons in real time under the same parameters. We used GFP/RFP in each

cell body as the TRIC signal.

Calcium imaging

For in vitro experiments, VNC and ovaries were dissected in *Drosophila* Adult

Hemolymph-Like Saline (AHLS) buffer. The cuticle and connective tissue covering the

VNC and ovary were removed using fine forceps. The median abdominal nerve,

connecting the abdominal ganglion and the reproductive tract, was either kept intact or

carefully severed, depending on the experiment.

Calcium imaging was performed on a Zeiss 980 confocal microscope (Jena,

Germany) equipped with 10× or 20× objectives. Briefly, cells were scanned with 488 nm

and 555 nm lasers at 512 × 512 pixels (or lower) and an 8 bit dynamic range to

acquire baseline GCaMP or tdTomato fluorescence. Regions of interest (ROIs) were

selected in FIJI, and fluorescence changes were quantified as $\Delta F/F_0 = (F - F_0)/F_0$,

where F_0 is the mean fluorescence over 30 frames in mdn LO neurons expressing

GCaMP. For ILP7 or tdc2 neurons, F_0 was defined as the average fluorescence over 30

frames before solution application. Depending on the experiment, solutions were applied

as follows: 10 µL AHLS buffer,10 µL of 100 mM ATP, and 10 µL of 1 mM octopamine

in 90 µL AHLS; and 20 µL of 50 mM glutamate in 80 µL AHLS.

Quantification and statistical analysis

We used the GraphPad Prism6 software package to generate graphs and analyze

data statistically. We tested whether the values were normally distributed using

D'Agostino–Pearson omnibus and Shapiro–Wilk normality tests before performing

statistical analysis. When data were normally distributed, we used parametric tests; when

data were not normally distributed, we used non-parametric tests. All tests were two-tailed.

All data are presented as mean ± s.e.m. We also consistently labeled the sample numbers

directly on graphs.

Sample size

Our results were based on an analysis of behavior at the level of single animals.

Sample sizes were determined before experimentation based on both variance and effect

sizes. All the data are biological replicates.

Acknowledgments

This research was funded by the National Key Research and Developmental

Program of China (2022YFD1700200), Key Research and Development Project of

Xizang Autonomous Region (XZ202401ZY0031) and the National Natural Science

Foundation of China (32022011 & 32472542).

1003

References

1. K. M. Cury, R. Axel, Flexible neural control of transition points within the egg-laying behavioral sequence in *Drosophila*. *Nat. Neurosci.* **26**, 1054–1067 (2023).

2. C. Oliveira-Ferreira, M. Gaspar, M. L. Vasconcelos, Neuronal substrates of egg-laying behaviour at the abdominal ganglion of *Drosophila melanogaster*. *Scientific Reports* **13**, 21941 (2023).

3. V. Vijayan, F. Wang, K. Wang, A. Chakravorty, A. Adachi, H. Akhlaghpour, B. J. Dickson, G. Maimon, A rise-to-threshold process for a relative-value decision. *Nature* **619**, 563–571 (2023).

4. C. H. Yang, P. Belawat, E. Hafen, L. Y. Jan, Y. N. Jan, Drosophila egg-laying site selection as a system to study simple decision-making processes. *Science* **319**, 1679 (2008).

5. L. D. Deady, J. Sun, A follicle rupture assay reveals an essential role for follicular adrenergic signaling in *Drosophila* ovulation. *PLOS Genetics* **11**, e1005604 (2015).

6. Q. Hu, M. F. Wolfner, The *Drosophila* Trpm channel mediates calcium influx during egg activation. *Proceedings of the National Academy of Sciences* **116**, 18994-19000 (2019).

7. B. J. Dickson, Wired for Sex: The Neurobiology of *Drosophila* Mating Decisions. *Science* **322**, 904-909 (2008).

8. B. Gou, Y. Liu, Ananya R. Guntur, U. Stern, C.-H. Yang, Mechanosensitive neurons on the internal reproductive tract contribute to egg-laying-induced acetic acid attraction in *Drosophila*. *Cell Rep.* **9**, 522-530 (2014).

9. R. J. Greenspan, J.-F. Ferveur, Courtship in *Drosophila*. *Annual Review of Genetics* **34**, 205-232 (2000).

10. J. C. Hall, The mating of a fly. *Science* **264**, 1702 (1994).

11. E. Kubli, Sex-peptides: seminal peptides of the *Drosophila* male. *Cell. Mol. Life Sci.* **60**, 1689-1704 (2003).

12. C. Rezával, Hania J. Pavlou, Anthony J. Dornan, Y.-B. Chan, Edward A. Kravitz, Stephen F. Goodwin, Neural circuitry underlying *Drosophila* female postmating behavioral responses. *Curr. Biol.* **22**, 1155-1165 (2012).

13. D. Yamamoto, M. Koganezawa, Genes and circuits of courtship behaviour in *Drosophila* males. *Nat. Rev. Neurosci.* **14**, 681 (2013).

14. K. Wang, F. Wang, N. Forknall, T. Yang, C. Patrick, R. Parekh, B. J. Dickson, Neural circuit mechanisms of sexual receptivity in *Drosophila* females. *Nature* **589**, 577–581 (2020).

15. F. Wang, K. Wang, N. Forknall, C. Patrick, T. Yang, R. Parekh, D. Bock, B. J. Dickson, Neural circuitry linking mating and egg laying in *Drosophila* females. *Nature* **579**, 101-105 (2020).

16. M. P. Nallasivan, D. N. D. Singh, M. S. R. S. Saleh, M. Soller, Sex-peptide targets distinct higher order processing neurons in the brain to induce the female post-mating response. *eLife* **13**, RP98283 (2024).

17. C. A. Middleton, U. Nongthomba, K. Parry, S. T. Sweeney, J. C. Sparrow, C. J.
H. Elliott, Neuromuscular organization and aminergic modulation of
contractions in the *Drosophila* ovary. *BMC Biology* **4**, 17 (2006).

18. M. A. White, D. S. Chen, M. F. Wolfner, She's got nerve: Roles of octopamine
in insect female reproduction. *J. Neurogenet.* **35**, 132-153 (2021).

19. N. Yapici, Y.-J. Kim, C. Ribeiro, B. J. Dickson, A receptor that mediates the
post-mating switch in *Drosophila* reproductive behaviour. *Nature* **451**, 33-37
(2008).

20. M. F. Wolfner, The gifts that keep on giving: physiological functions and
evolutionary dynamics of male seminal proteins in *Drosophila*. *Heredity* **88**,
85-93 (2002).

21. H.-G. Lee, C.-S. Seong, Y.-C. Kim, R. L. Davis, K.-A. Han, Octopamine
receptor OAMB is required for ovulation in *Drosophila melanogaster*.
Developmental Biology **264**, 179-190 (2003).

22. S. A. Deshpande, E. W. Rohrbach, J. D. Asuncion, J. Harrigan, A. Eamani, E.
H. Schlingmann, D. J. Suto, P.-T. Lee, F. E. Schweizer, H. J. Bellen, D. E.
Krantz, Regulation of *Drosophila* oviduct muscle contractility by octopamine.
iScience **25**, 104697 (2022).

23. R. Rodríguez-Valentín, I. López-González, R. Jorquera, P. Labarca, M. Zurita,
E. Reynaud, Oviduct contraction in *Drosophila* is modulated by a neural
network that is both, octopaminergic and glutamatergic. *Journal of Cellular*
Physiology **209**, 183-198 (2006).

24. M. C. Castellanos, J. C. Y. Tang, D. W. Allan, Female-biased dimorphism
underlies a female-specific role for post-embryonic Ilp7 neurons in *Drosophila*
fertility. *Development* **140**, 3915 (2013).

25. K. J. Thompson, Oviposition digging in the grasshopper: II. descending neural
control. *Journal of Experimental Biology* **122**, 413-425 (1986).

26. M. Vyas, S. K. Parepally, P. D. Kamala Jayanthi, Is the natural instinct to
oviposit in mated female oriental fruit fly, *Bactrocera dorsalis* more of a
brain-independent act? *Frontiers in Physiology* **13**, 800441 (2022).

27. A. Ayali, A. B. Lange, Rhythmic behaviour and pattern-generating circuits in
the locust: Key concepts and recent updates. *Journal of Insect Physiology* **56**,
834-843 (2010).

28. Y. V. Zhang, T. J. Aikin, Z. Li, C. Montell, The basis of food texture sensation in
Drosophila. *Neuron* **91**, 863-877 (2016).

29. B. Coste, J. Mathur, M. Schmidt, T. J. Earley, S. Ranade, M. J. Petrus, A. E.
Dubin, A. Patapoutian, Piezo1 and Piezo2 are essential components of distinct
mechanically activated cation channels. *Science* **330**, 55-60 (2010).

30. B. Coste, B. Xiao, J. S. Santos, R. Syeda, J. Grandl, K. S. Spencer, S. E. Kim,
M. Schmidt, J. Mathur, A. E. Dubin, M. Montal, A. Patapoutian, Piezo proteins
are pore-forming subunits of mechanically activated channels. *Nature* **483**,
176-181 (2012).

31. Y. Guo, Y. Wang, W. Zhang, S. Meltzer, D. Zanini, Y. Yu, J. Li, T. Cheng, Z. Guo,
Q. Wang, J. S. Jacobs, Y. Sharma, D. F. Eberl, M. C. Göpfert, L. Y. Jan, Y. N.

Jan, Z. Wang, Transmembrane channel-like (tmc) gene regulates *Drosophila*
larval locomotion. *Proc. Natl. Acad. Sci. U S A.* **113**, 7243-7248 (2016).

32.   L. Hyunjin, C. Hyun Woo, Z. Chen, P. Zee Yong, K. Young Joon, A pair of oviduct-born *pickpocket* neurons important for egg-laying in *Drosophila melanogaster*. *Mol. Cells* **39**, 573-579 (2016).

33.   S. F. Wu, Y. L. Ja, Y. J. Zhang, C. H. Yang, Sweet neurons inhibit texture discrimination by signaling TMC-expressing mechanosensitive neurons in *Drosophila*. *eLife* **8**, e46165 (2019).

34.   R. M. Lam, L. J. von Buchholtz, M. Falgairolle, J. Osborne, E. Frangos, M. R. Servin-Vences, M. Nagel, M. Q. Nguyen, M. Jayabalan, D. Saade, A. Patapoutian, C. G. Bönnemann, N. J. P. Ryba, A. T. Chesler, PIEZO2 and perineal mechanosensation are essential for sexual function. *Science* **381**, 906-910 (2023).

35.   H. M. Yamanouchi, R. Tanaka, A. Kamikouchi, Piezo-mediated mechanosensation contributes to stabilizing copulation posture and reproductive success in *Drosophila* males. *iScience* **26**, 106617 (2023).

36.   J. Qin, T. Yang, K. Li, T. Liu, W. Zhang, Pharyngeal mechanosensory neurons control food swallow in *Drosophila melanogaster*. *eLife*, 1-24 (2023).

37.   Y. L. Jia, Y. J. Zhang, D. Guo, C. Y. Li, J. Y. Ma, C. F. Gao, S. F. Wu, A mechanosensory receptor TMC regulates ovary development in the brown planthopper *Nilaparvata lugens*. *Frontiers in Genetics* **11**, 1152 (2020).

38.   M. Häsemeyer, N. Yapici, U. Heberlein, B. J. Dickson, Sensory neurons in the *Drosophila* genital tract regulate female reproductive behavior. *Neuron* **61**, 511-518 (2009).

39.   R. Court, S. Namiki, J. D. Armstrong, J. Börner, G. Card, M. Costa, M. Dickinson, C. Duch, W. Korff, R. Mann, D. Merritt, R. K. Murphey, A. M. Seeds, T. Shirangi, J. H. Simpson, J. W. Truman, J. C. Tuthill, D. W. Williams, D. Shepherd, A systematic nomenclature for the *Drosophila* ventral nerve cord. *Neuron* **107**, 1071-1079.e1072 (2020).

40.   K.-i. Kimura, T. Hachiya, M. Koganezawa, T. Tazawa, D. Yamamoto, Fruitless and doublesex coordinate to generate male-specific neurons that can initiate courtship. *Neuron* **59**, 759-769 (2008).

41.   Y. Pan, G. W. Meissner, B. S. Baker, Joint control of *Drosophila* male courtship behavior by motion cues and activation of male-specific P1 neurons. *Proc. Natl. Acad. Sci. U S A.* **109**, 10065-10070 (2012).

42.   S. T. Sweeney, K. Broadie, J. Keane, H. Niemann, C. J. O'Kane, Targeted expression of tetanus toxin light chain in *Drosophila* specifically eliminates synaptic transmission and causes behavioral defects. *Neuron* **14**, 341-351 (1995).

43.   J. Ma, M. Ptashne, The carboxy-terminal 30 amino acids of GAL4 are recognized by GAL80. *Cell* **50**, 137-142 (1987).

44.   T. Lee, L. Luo, Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron* **22**, 451-461 (1999).

45.   C.-H. Yang, S. Rumpf, Y. Xiang, M. D. Gordon, W. Song, L. Y. Jan, Y.-N. Jan,

Control of the postmating behavioral switch in *Drosophila* females by internal sensory neurons. *Neuron* **61**, 519-526 (2009).

46. F. N. Hamada, M. Rosenzweig, K. Kang, S. R. Pulver, A. Ghezzi, T. J. Jegla, P. A. Garrity, An internal thermal sensor controlling temperature preference in *Drosophila*. *Nature* **454**, 217-220 (2008).

47. Jennifer J. Bussell, N. Yapici, Stephen X. Zhang, Barry J. Dickson, Leslie B. Vosshall, Abdominal-B neurons control *Drosophila* virgin female receptivity. *Curr. Biol.* **24**, 1584-1595 (2014).

48. D. Richard, G. Maria, C. Catherine, D. Aaron, Visualizing glutamatergic cell bodies and synapses in *Drosophila* larval and adult CNS. *Journal of Comparative Neurology* **508**, 131-152 (2008).

49. N. C. Klapoetke, Y. Murata, S. S. Kim, S. R. Pulver, A. Birdsey-Benson, Y. K. Cho, T. K. Morimoto, A. S. Chuong, E. J. Carpenter, Z. Tian, J. Wang, Y. Xie, Z. Yan, Y. Zhang, B. Y. Chow, B. Surek, M. Melkonian, V. Jayaraman, M. Constantine-Paton, G. K.-S. Wong, E. S. Boyden, Independent optical excitation of distinct neural populations. *Nat. Meth.* **11**, 338-346 (2014).

50. E. D. Hoopfer, Y. Jung, H. K. Inagaki, G. M. Rubin, D. J. Anderson, P1 interneurons promote a persistent internal state that enhances inter-male aggression in Drosophila. *eLife* **4**, e11346 (2015).

51. X. J. Gao, O. Riabinina, J. Li, C. J. Potter, T. R. Clandinin, L. Luo, A transcriptional reporter of intracellular Ca2+ in *Drosophila*. *Nature Neuroscience* **18**, 917-925 (2015).

52. T.-W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A. Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda, D. S. Kim, Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300 (2013).

53. H. Dana, Y. Sun, B. Mohar, B. K. Hulse, A. M. Kerlin, J. P. Hasseman, G. Tsegaye, A. Tsang, A. Wong, R. Patel, J. J. Macklin, Y. Chen, A. Konnerth, V. Jayaraman, L. L. Looger, E. R. Schreiter, K. Svoboda, D. S. Kim, High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nat. Methods* **16**, 649-657 (2019).

54. S. Q. Lima, G. Miesenböck, Remote control of behavior through genetically targeted photostimulation of neurons. *Cell* **121**, 141-152 (2005).

55. Z. Yao, A. M. Macara, K. R. Lelito, T. Y. Minosyan, O. T. Shafer, Analysis of functional neuronal connectivity in the *Drosophila* brain. *J. Neurophysiol.* **108**, 684-696 (2012).

56. M. Talay, E. B. Richman, N. J. Snell, G. G. Hartmann, J. D. Fisher, A. Sorkaç, J. F. Santoyo, C. Chou-Freed, N. Nair, M. Johnson, J. R. Szymanski, G. Barnea, Transsynaptic mapping of second-order taste neurons in flies by trans-Tango. *Neuron* **96**, 783-795.e784 (2017).

57. C. Rezával, T. Nojima, Megan C. Neville, Andrew C. Lin, Stephen F. Goodwin, Sexually dimorphic octopaminergic neurons modulate female postmating behaviors in *Drosophila*. *Curr. Biol.* **24**, 725-730 (2014).

58. Y. Li, C. Fink, S. El-Kholy, T. Roeder, The octopamine receptor octß2R is

essential for ovulation and fertilization in the fruitfly *Drosophila melanogaster*. *Arch. Insect Biochem. Physiol.* **88**, 168-178 (2015).

59. J. Lim, P. R. Sabandal, A. Fernandez, J. M. Sabandal, H.-G. Lee, P. Evans, K.-A. Han, The octopamine receptor Octβ2R regulates ovulation in *Drosophila melanogaster*. *PLoS ONE* **9**, e104441 (2014).

60. L. J. Macpherson, E. E. Zaharieva, P. J. Kearney, M. H. Alpert, T.-Y. Lin, Z. Turan, C.-H. Lee, M. Gallio, Dynamic labelling of neural connections in multiple colours by trans-synaptic fluorescence complementation. *Nat. Commun.* **6**, 10024 (2015).

61. R. A. Baines, J. P. Uhler, A. Thompson, S. T. Sweeney, M. Bate, Altered electrical properties in *Drosophila* neurons developing without synaptic transmission. *The Journal of Neuroscience* **21**, 1523 (2001).

62. I. Miguel-Aliaga, S. Thor, A. P. Gould, Postmitotic specification of *Drosophila* insulinergic neurons from pioneer neurons. *PLOS Biology* **6**, e58 (2008).

63. P. Cognigni, A. P. Bailey, I. Miguel-Aliaga, Enteric neurons and systemic signals couple nutritional and reproductive status with intestinal homeostasis. *Cell Metabolism* **13**, 92-104 (2011).

64. E. Prince, J. Kretzschmar, L. C. Trautenberg, S. Broschk, M. Brankatschk, DIlp7-producing neurons regulate insulin-producing cells in *Drosophila*. *Frontiers in Physiology* **12**, 630390 (2021).

65. M. Chatzigeorgiou, S. Bang, S. W. Hwang, W. R. Schafer, tmc-1 encodes a sodium-sensitive channel required for salt chemosensation in *C. elegans*. *Nature* **494**, 95-99 (2013).

66. G. Keresztes, H. Mutai, S. Heller, TMC and EVER genes belong to a larger novel family, the TMC gene family encoding transmembrane proteins. *BMC Genomics* **4**, 24 (2003).

67. K. Kurima, L. M. Peters, Y. Yang, S. Riazuddin, Z. M. Ahmed, S. Naz, D. Arnaud, S. Drury, J. Mo, T. Makishima, M. Ghosh, P. S. N. Menon, D. Deshmukh, C. Oddoux, H. Ostrer, S. Khan, S. Riazuddin, P. L. Deininger, L. L. Hampton, S. L. Sullivan, J. F. Battey Jr, B. J. B. Keats, E. R. Wilcox, T. B. Friedman, A. J. Griffith, Dominant and recessive deafness caused by mutations of a novel gene, TMC1, required for cochlear hair-cell function. *Nat. Genet.* **30**, 277-284 (2002).

68. X. Yue, Y. Sheng, L. Kang, R. Xiao, Distinct functions of TMC channels: a comparative overview. *Cellular and Molecular Life Sciences* **76**, 4221-4232 (2019).

69. Y. Kawashima, K. Kurima, B. Pan, A. J. Griffith, J. R. Holt, Transmembrane channel-like (TMC) genes are required for auditory and vestibular mechanosensation. *Pflugers Arch.* **467**, 85-94 (2015).

70. K. Kurima, L. M. Peters, Y. Yang, S. Riazuddin, Z. M. Ahmed, S. Naz, D. Arnaud, S. Drury, J. Mo, T. Makishima, M. Ghosh, P. S. N. Menon, D. Deshmukh, C. Oddoux, H. Ostrer, S. Khan, S. Riazuddin, P. L. Deininger, L. L. Hampton, S. L. Sullivan, J. F. Battey, B. J. B. Keats, E. R. Wilcox, T. B. Friedman, A. J. Griffith, Dominant and recessive deafness caused by

1223 mutations of a novel gene, TMC1, required for cochlear hair-cell function. *Nature Genetics* **30**, 277-284 (2002).

71. Y. Kawashima, K. Kurima, B. Pan, A. J. Griffith, J. R. Holt, Transmembrane channel-like (TMC) genes are required for auditory and vestibular mechanosensation. *Pflügers Archiv - European Journal of Physiology* **467**, 85-94 (2015).

72. B. Pan, Gwenaelle S. Géléoc, Y. Asai, Geoffrey C. Horwitz, K. Kurima, K. Ishikawa, Y. Kawashima, Andrew J. Griffith, Jeffrey R. Holt, TMC1 and TMC2 are components of the mechanotransduction channel in hair cells of the mammalian inner ear. *Neuron* **79**, 504-515 (2013).

73. X. Wang, G. Li, J. Liu, J. Liu, X. Z. S. Xu, TMC-1 mediates alkaline sensation in *C. elegans* through nociceptive neurons. *Neuron* **91**, 146-154 (2016).

74. X. Yue, J. Zhao, X. Li, Y. Fan, D. Duan, X. Zhang, W. Zou, Y. Sheng, T. Zhang, Q. Yang, J. Luo, S. Duan, R. Xiao, L. Kang, TMC proteins modulate egg laying and membrane excitability through a background leak conductance in *C. elegans*. *Neuron* **97**, 571-585 (2018).

75. E. Kaulich, D. S. Walker, Y. Q. Tang, W. R. Schafer, The *Caenorhabditis elegans* tmc-1 is involved in egg-laying inhibition in response to harsh touch. *microPublication biology* **2021**, (2021).

76. L. Zhang, D. G. Gualberto, X. Guo, P. Correa, C. Jee, L. R. Garcia, TMC-1 attenuates *C. elegans* development and sexual behaviour in a chemically defined food environment. *Nat. Commun.* **6**, 6345 (2015).