Title: Oligodendrocyte progenitor cell responses to inflammatory demyelination with aging in

mouse model of multiple sclerosis

Authors: Emily E. Fresenko[1], Camilla N. Bahri[1], Benjamin J. Burson[1], Noor F. Ahmed[1], Davin

Packer[1], Benjamin J. Tabor Jr[1], Bogdan Beirowski[1], Wenjing Sun[2], Michelle A. Wedemeyer[3,4],

Cole A. Harrington[1*]

Author affiliations and footnotes

[1]The Ohio State University Wexner Medical Center, Department of Neurology, The

Neuroscience Research Institute, College of Medicine, Columbus, OH, USA

[2]The Ohio State University Wexner Medical Center, Department of Neuroscience, College of

Medicine, Columbus, OH, USA

[3]The Ohio State University Wexner Medical Center, Department of Neurosurgery and Pediatrics,

College of Medicine, Columbus, OH, USA

[4]Nationwide Children's Hospital, Division of Neurosurgery, Columbus, OH, USA

Contact information

*Corresponding author:

Cole A. Harrington

460 W 12th Ave, Biomedical Research Tower 604

Columbus, OH 43210

1-614-688-1184

cole.harrington@osumc.edu

Conflict of Interest Statement

The authors have declared that no conflict of interest exists.

Abstract

Oligodendrocyte progenitor cells (OPCs) have the capacity to self-renew, differentiate, and

remyelinate the CNS. Aging is associated with a reduction in the functional capacity of OPCs

even in the absence of an autoimmune insult. To determine how aging affects the response of

oligodendroglia to a strong inflammatory insult comparable to an immune-mediated

demyelinating event in multiple sclerosis (MS), we performed adoptive transfer of young myelin-

reactive Th17 T cells into young and aged OPC lineage tracing mice. After adoptive transfer,

OPCs were enriched within spinal cord lesions of both young and aged mice. However

differentiated oligodendrocytes (OLs) were significantly reduced after adoptive transfer. Both

young and aged OPCs differentiated into mature OLs during adoptive transfer. Transmission

electron microscopy revealed thinly myelinated axons without degenerative features that likely

represent remyelinated axons in lesions of both age groups. Young and aged OPCs rise to the

challenge after a strong auto-immune attack, suggesting that compensatory strategies permit

both young and aged oligodendroglia to survive despite an inflammatory environment.

Identifying pathways that promote resilience of oligodendroglia in the face of an inflammatory

challenge will facilitate the development of remyelinating therapies for people with MS.

Introduction

In the postnatal CNS, oligodendrocyte progenitor cells (OPCs) are lineage restricted

progenitors with the capacity to proliferate, migrate, and differentiate into mature myelinating

oligodendrocytes during development and adulthood (1-3) and in response to demyelinating

injury (4,5). Multiple sclerosis (MS) is a chronic demyelinating disorder of the CNS in which an

autoimmune attack of oligodendroglia and their myelin processes results in demyelinating

lesions (6-8). OPCs and maturing oligodendrocytes have been identified in MS lesions from

patients of all ages (9,10) suggesting that OPCs may retain the capacity to differentiate and

remyelinate after a demyelinating attack despite an inhibitory inflammatory environment (8,11).

Aging is a risk factor for progressive accumulation of neurological disability in MS

(12,13). Cellular aging in MS leads to senescence of immune cells and other CNS resident cell

populations (13) which may influence the ability of aged OPC to differentiate and remyelinate.

Physiological aging of OPCs, in the absence of an immune disorder or model, is associated with

phenotypic changes in OPC ion channel expression (14), reduced proliferation and

differentiation (14,15), increased cellular senescence and upregulation of immune response

pathways (14-16). Developmental inhibitors of OPC differentiation, e.g. Wnt and hypoxia

inducible factor-1 alpha (HIF-1a) pathways, are upregulated in aged OPCs and pharmacological

inhibition of these pathways promotes differentiation of aged OPCs *in vitro* (16). In toxin-

mediated demyelination models, OPC proliferation, recruitment into lesions, differentiation and

remyelination are all slower in aged animals (17-19). Metformin plus alternative day fasting (15)

and young macrophages (20) can partially rescue the slower remyelination kinetics in aged

animals. These studies suggest that aged OPC function can be augmented to levels similar to

young OPCs, and targeting mechanisms that promote aged OPC function may be a viable

therapeutic strategy to promote remyelination in MS.

Formal investigations of the impact of aging on OPC performance after an autoimmune-

mediated inflammatory insult have not been conducted to date. While toxin-mediated models

benefit from stereotyped OPC responses, the lack of an adaptive immune response in these

models may not accurately reflect the microenvironment of human MS lesions. Inflammatory

models offer the benefit of a T cell mediated auto-immune process targeting oligodendroglia and

myelin, better simulating an acute MS demyelinating event (6, 21). Although inflammatory

models reproduce the histological features of severe disease, including early axonal

degeneration, high inflammatory infiltrate, rapid severe disease course and lack of disease

remission (22-24), the rapid progression and severity of clinical disease prevent the use of this

model for chronic timepoints. Despite these limitations, the study of murine models with a strong

autoimmune inflammatory response may uncover mechanisms that oligodendroglia utilize to

adapt to a harsh inflammatory environment. Studies in both murine models (25-31) and human

MS tissue (29,32) indicate the oligodendroglia adopt distinct transcriptional states in response to

inflammation. These "disease-associated" oligodendroglia upregulate immune signaling

pathways and may directly interact with T cells by expression of MHC class I and II molecules

(30, 33-34). The function of disease-associated oligodendroglia is unclear and may be context

and model dependent. Investigation of OPC properties in a strong inflammatory model will offer

insight into how young and aged OPCs respond to an auto-immune demyelinating insult.

We performed lineage tracing of OPCs in the Th17 adoptive transfer model to

investigate how aging affects the response of OPCs to a robust autoimmune demyelinating

event. A tamoxifen-inducible OPC Cre line (*Pdgfra-Cre^{ERT}*) (35) was crossed to an inducible

GFP reporter line (*RCE- Rosa26^{flstopflEGFP}*) (36), tamoxifen was administered to lineage trace

OPCs starting two weeks prior to adoptive transfer of young myelin-reactive Th17 T cells into

aged (10-14 months) and young (3-5 months) *Pdgfra-Cre^{ERT}*;*RCE* recipients. The Th17 adoptive

transfer model offers several advantages over other immune-mediated models including 1)

control of adoptive transfer T cell age to prevent potential confounding from aged donor T cells

(37) and 2) selective determination of oligodendroglial responses to a Th17 T cell mediated

disease as Th17 T cells play a key role in MS pathogenesis (22, 38-39).

Following myelin-reactive Th17 adoptive transfer into *Pdgfra-Cre^{ERT};RCE* mice, we found

that OPCs from both young and aged mice were enriched in spinal cord lesions. Higher OPC

densities were seen in lesions compared to non-lesion white matter suggesting that the lesion

environment promotes OPC recruitment and/or survival mechanisms. Differentiated

oligodendrocytes (OLs) were reduced in adoptive transfer in both young and aged animals. A

small minority of lineage traced OPCs differentiated into mature OLs in both ages. Transmission

electron microscopy (TEM) analysis of lesions identified similar proportions of thinly myelinated

axons without degenerative features in both young and aged lesions suggesting that

remyelination can occur even in the presence of an ongoing inflammatory response. These

findings indicate that both young and aged OPCs respond similarly to an auto-immune

inflammatory demyelinating insult and OPCs may have compensatory strategies to support

survival and physiological function in an inflammatory environment.

Results

Aged Th17 adoptive transfer animals have increased disease severity

To identify differences in the response of young and aged OPCs to an adaptive auto-

immune response, young and aged mice cohorts were used for this study (Figure 1A, age at

sacrifice/tissue analysis- young: P87-P148, 3-5 months; aged: P295-P427,10-14 months). Aged

mice age range was selected based on mouse aging functional studies correlating our aged

animal cohort to human age in 4^{th} decade (40) which is the average onset of late-onset

relapsing remitting MS (LORRMS) and progressive MS (41,42). Experiments with aged animals

beyond 14 months was not performed due to increased disease severity and early morbidity in

our aged cohort (Figure 1B-C). OPC lineage tracing was initiated by administering tamoxifen to

young and aged *Pdgfra-Cre^{ERT}*; *RCE* (*Rosa26^{flstopflEGFP}*) mice two weeks prior to adoptive

transfer (Figure 1A) to avoid tamoxifen-mediated impairment of T-cell activation and disease

induction (43). Myelin oligodendrocyte glycoprotein (MOG)-reactive T cells were generated by

immunizing young (2-3-month-old) female wild-type C57BL/6J donors with MOG$_{35-55}$ peptide

followed by *ex-vivo* polarization into Th17 phenotype and adoptive transfer into young and aged

Pdgfra-Cre^{ERT}; *RCE* OPC lineage traced mice. Young and aged *Pdgfra-Cre^{ERT}*; *RCE* mice that

were administered tamoxifen but did not receive adoptive transfer served as naïve controls. For

transmission electron microscopy analysis (TEM), two additional experimental lines were used:

wild-type C57BL/6J mice that did not receive tamoxifen and OPC Cre combined with the

Tau^{mGFP} reporter (*Pdgfra-Cre^{ERT}*; *Tau^{mGFP}*) which have been previously used to visualize newly

formed myelin sheaths (43-45). *Pdgfra-Cre^{ERT}*; *RCE* OPC lineage traced mice were sacrificed at

9-10 days post-adoptive transfer, corresponding to peak disease (Figure 1B) for

immunohistochemistry analysis of spinal cord. Wild-type and *Pdgfra-Cre^ERT*; *Tau^mGFP* OPC

remyelination reporter mice were sacrificed at 10 days post-adoptive transfer for TEM.

Extension of the analysis timepoint in aged mice beyond the acute peak disease time-point was

not possible without significant loss of aged animals as one third of aged mice reached humane

end-point criteria at day 9 (Figure 1C, score ≥4, no movement around cage, inability to feed).

Aged adoptive transfer mice had significantly higher experimental autoimmune

encephalomyelitis (EAE) clinical scores relative to young adoptive transfer mice (Figure 1B) and

faster onset to quadriplegia (Figure 1C). Increased disease severity in aged Th17 adoptive

transfer recipients in other studies is not ameliorated by young bone marrow transplantation

indicating that the aged host CNS environment drives an exacerbated disease course (46).

Pdgfra-Cre lines lineage trace hematopoietic cells in CNS inflammatory models

Lineage traced *Pdgfra-Cre^ERT;RCE* mice demonstrated high recombination efficiency

and GFP reporter induction in OPCs (Figure 1D) with ~70% of Pdgfra-positive cells expressing

GFP (Figure 1E). Lineage traced GFP reporter-positive cells demonstrated expression of pan-

oligodendroglial marker Olig2 (Figure 1F). However, this OPC Cre *Pdgfra-Cre^ERT* line

consistently demonstrated non-oligodendroglial lineage traced cells (GFP+/Olig2-) in adoptive

transfer spinal cord particularly within lesions (Figure 1G-H). GFP reporter positive cells were

positive for hematopoietic cell marker CD45 and Pdgfra (Figure 1I). Flow cytometry confirmed

the presence of a GFP+CD45+ cell population in adoptive transfer CNS tissue (Figure 1J),

indicating that peripheral immune cells that traffic into the CNS undergo recombination. These

findings highlight the need for rigorous validation of Cre drivers with a lineage tracing approach

when using CNS inflammatory disease models. Given the presence of GFP reporter positive

non-oligodendroglial cells in adoptive transfer, subsequent quantitative analysis of lineage

traced OPCs required inclusion of oligodendroglial restricted markers.

Young and aged lineage traced OPCs are enriched within adoptive transfer lesions

In the Th17 adoptive transfer model, immune infiltrates are concentrated at

leptomeningeal and perivascular spaces of the spinal cord and hindbrain (47-48). To

characterize the impact of aging on OPC density and differentiation in response to T cell

adoptive transfer, transverse spinal cord cross-sections from young and aged adoptive transfer

and naïve animals were stained for lineage traced OPCs with GFP and pan-oligodendroglial

marker Olig2 (Figure 2A). Lineage traced OPCs were quantified in total spinal cord cross-

sections, lesions (determined by DAPI hypercellularity), and non-lesional white matter. For initial

quantification all lesions present in one spinal cord cross-section were grouped together for the

total lesional density (Figure 2B). To capture the differences across lesions within a single spinal

cord section, individual lesions were quantified separately for downstream analyses (Figure 2C-

F, H). Aged lineage traced OPC (GFP+Olig2+) density was significantly lower in adoptive

transfer whole cord compared to young, however aged OPCs were present at similar densities

in adoptive transfer compared to aged naïve whole cord (Figure 2B) reflecting lower baseline

OPC densities with aging. Lineage traced OPC (GFP+Olig2+) density was not significantly

different between aged and young lesions when pooling all lesions across a spinal cord section

and not accounting for differentiation status (Figure 2B).

To better characterize the fate and differentiation status of lineage traced OPCs, spinal

cord sections were co-labeled with GFP, mature oligodendrocyte marker (Aspa), and

hematopoietic marker (CD45) followed by quantification of all GFP+CD45- cells in spinal cord

cross-sections (Figure 2C-H). In both young and aged animals, lineage traced OPCs that

remained OPCs (OPC->OPC, GFP+Aspa-CD45-) were significantly increased within lesions

compared to non-lesional white matter (Figure 2C). OPC->OPC density was significantly lower

in aged compared to young lesions (Figure 2C). To determine the enrichment of OPCs within

lesions compared to surrounding non-lesional white matter, a ratio of mean lesional density over

non-lesional white matter (nlwm) density was calculated for each spinal cord cross-section

(Figure 2D). Lineage traced OPCs that remained OPCs (GFP+Aspa-CD45-) had a greater than

2-fold enrichment in both young and aged lesions compared to surrounding nlwm suggesting

recruitment, proliferation, and/or retention of OPCs within the lesional environment.

Young and aged OPCs demonstrate the ability to differentiate in acute adoptive transfer

The density of OPCs increased in lesions of young and aged animals, suggesting that

despite the more severe disease observed in aged animals, OPCs retain the capacity to

respond to demyelination and potentially differentiate. A small minority of lineage traced OPCs

underwent differentiation and expressed Aspa (GFP+Aspa+CD45-) in both young and aged

adoptive transfer spinal cord (Figure 2E). Compared to non-lesion white matter, differentiated

OPCs were significantly higher in young lesions and significantly lower in aged lesions (Figure

2E). Young OPCs may have a higher or faster capacity for differentiation within lesions and/or

aged OPCs undergoing differentiation may be more susceptible to cell death in the aged

environment. Comparing the ratio of newly differentiated OPCs to surrounding nlwm on an

individual spinal cord section level (Figure 2F) did not indicate a significant difference between

the ratio of differentiated OLs within young and aged lesions compared to surrounding nlwm

(Figure 2F). To determine the percentage of lineage traced OPCs that remained OPCs within

lesions, all lineage traced OPCs (GFP+CD45-) were quantified within individual lesions and

assessed for Aspa expression. Similar to density analysis, lineage traced OPCs that remained

OPCs made up over 95% of lineage traced OPCs within lesions of young and aged animals

(Figure 2H). The enrichment of lineage traced OPCs within young and aged lesions and ability

of OPCs to differentiate, particularly within young lesions at this peak disease time-point,

indicates that OPCs can quickly respond even with ongoing inflammatory activity and high

disease severity. Alternative immune-mediated models with remission of clinical disease may

offer additional insight into the kinetics of OPC responses after resolution of inflammatory

activity.

Young and aged differentiated oligodendrocytes are lost in adoptive transfer

Adoptive transfer of myelin oligodendrocyte glycoprotein (MOG)-reactive Th17 T cells may result in targeted cytotoxicity to MOG-expressing mature OLs. To determine whether OLs are lost or preserved in young and aged adoptive transfer, spinal cord sections were stained for OL marker Aspa (Figure 3A). High densities of differentiated OLs in spinal cord cross-sections necessitated the development of a semi-automated cell counting method. An ImageJ macro was developed to quantify Aspa-positive cell densities (Figure 3B). Aspa+ OL densities were significantly lower at baseline in aged compared to young naïve spinal cord (Figure 3C). Adoptive transfer resulted in a significant decrease in OL density by ~1/4 of density relative to naïve controls (Figure 3C) and aged adoptive transfer spinal cord had the highest degree of OL loss. To determine if existing OLs (not derived from OPCs) were selectively lost within lesions we performed manual quantification of OLs in individual lesions and adjacent nlwm with use of GFP/Aspa/CD45 co-labeling (Figure 2G). Existing differentiated OLs (Aspa+GFP-CD45-) were significantly lower in lesions compared to non-lesions in both young and aged animals with the lowest density of OLs within aged lesions (Figure 3D). The ratio of OL lesional to non-lesional density was not significantly different between young and aged adoptive transfer (Figure 3E).

The lower density of aged OLs throughout spinal cord regions may reflect overall higher disease severity in aged animals and/or increased susceptibility of aged OLs to cell death. To investigate the impact of EAE score and CD45 infiltrate on OPC and OL densities within the spinal cord of young and aged adoptive transfer animals we performed a correlation analysis on matched cell counts from individual spinal cord sections. Variables of age, EAE score, percentage of cord occupied by CD45 infiltrate, and density of OPCs (GFP+Aspa-CD45-), OPC->OL (GFP+Aspa+CD45-), and OLs (GFP-Aspa+CD45-) across the whole cord and mean of pooled lesions were compared for each spinal cord section (Supplemental Figure 1A). Strong correlations (r values $<$-0.5 or $>$+0.5 and p values of $\leq$0.001) included the following correlations:

positive correlations- age to EAE score, OPCs in cord to OPCs in lesion, OPCs in cord to OPC-

>OL in cord, and negative correlation- age to OL lesion. EAE score was not significantly

correlated with OPC/OL densities. Degree of CD45 infiltrate was not correlated with age, EAE

score or OPC/OL densities. While pooling counts from lesions may limit the power of this

analysis, it suggests that the range of EAE scores across aged and young animals does not

markedly impact OPC/OL densities. Notably, higher levels of OPCs throughout the whole cord

was correlated with higher OPCs within lesions and OPCs that differentiated in the cord. Higher

age may also correlate with reduced OL lesional densities which was also suggested by our

macro and individual lesional quantification (Figure 3C-D).

Aging increases myelin debris accumulation after adoptive transfer

To determine the impact of aging on the degree of axonal loss, myelin debris and

demyelination after adoptive transfer, transverse spinal cord sections from young and aged

adoptive transfer mice were stained for neurofilament (NF), degraded myelin basic protein

(dMBP), and FluoroMyelin (total myelin) (Figure 4). To quantify axonal loss within lesions an

ImageJ macro was developed to quantify neurofilament-positive axonal density (Figure 4A).

Axonal density was highly variable across lesions, and no significant differences were observed

between young and aged lesions (Figure 4B). The accumulation of myelin debris was assessed

by the presence of dMBP staining (Figure 4C) and lesions in aged animals had a significantly

higher percentage of white matter with positive dMBP staining (Figure 4D). Higher levels of

myelin debris in aged animals may reflect impaired myelin debris clearance by aged

macrophages (20). Although myelin debris was higher in aged animals, there was no significant

differences in percent of white matter demyelination quantified by FluoroMyelin staining

between age groups (Figure 4E-F).

Remyelination is present to a similar extent in young and aged lesions

In the Th17 adoptive transfer model, areas of demyelination are localized in the

peripheral white matter adjacent to leptomeningeal infiltrates indicated by DAPI hypercellularity

(Figure 4E) (47-48). To determine if OPCs can remyelinate in adoptive transfer, we crossed

OPC inducible Cre line *Pdgfra-Cre^{ERT}* to inducible membrane targeted GFP within the

microtubule associated protein tau locus (*Tau^{mGFP}*) which has been used to visualize newly

generated myelin sheaths (43-45). Newly generated OL GFP+ myelin sheaths were not seen in

naïve spinal cord (Figure 5A). In adoptive transfer, GFP+ processes surrounding axons and co-

labeled with MBP were visualized at the edge of lesions (Figure 5B) suggesting the presence of

remyelination from newly formed OLs. Immunohistochemistry of a young and aged adoptive

transfer cohort of remyelination reporter mice (*Pdgfra-Cre^{ERT};Tau^{mGFP}*) was performed however

quantification of remyelination was not pursued due to GFP reporter background expression in

OPC processes in the parenchyma (Figure 5B).

To more reliably assess remyelination transmission electron microscopy (TEM) was

performed on wild-type C57BL/6J young and aged ventromedial white matter lesions at 10 days

post-adoptive transfer. Longer time-point analysis was not possible in aged animals due to

increased morbidity in this model (Figure 1C). Ventromedial white matter was chosen based on

the high rate of lesions in this region, ability to identify landmarks with the presence of a ventral

vein, and large diameter axons (49) that could facilitate the ability to detect thinner remyelinated

axons. Toxin-mediated demyelinating models have demonstrated that remyelinated axons can

be identified by thinner myelin sheaths (50-52), however thin myelin sheaths can also be a sign

of degenerating axons both in toxin-mediated and immune-mediated models (53). Perivascular

immune infiltrates lining areas of demyelination were identified in all ventromedial white matter

sections imaged as well as high concentration of oligodendrocytes within demyelinated areas

(Figure 5C). Thin myelin sheaths surrounding axons with degenerative features (electron dense

material, myelin blebbing, myelin infoldings, myelin unraveling, high mitochondrial

concentration) and axons without these degenerative features were present within lesions

(Figure 5D). TEM lesion images at 1150x magnification were used for quantification of young

and aged lesions with inverted images facilitating the visualization of myelin sheaths and axonal

components (Figure 5E). G-ratio quantification was performed on 1150x magnification images

of young and aged ventromedial white matter lesions (non-inverted) using MyelTracer (54)

combined with manual outlining for axons with a diameter $\geq 2\mu$m. Axons with thin myelin sheaths

were noted to have g-ratios of 0.8 or higher. Axons with g-ratio $\geq$ 0.8 were present in both age

groups. G-ratio slopes were significantly different between young and aged lesions (Figure 5F).

Relative frequency distribution analysis of g-ratios demonstrated a significantly higher frequency

of lower g-ratios (thicker myelin) in aged lesions (Figure 5G), indicating that the difference in g-

ratio slope was driven by thicker myelin sheaths in aged compared to young animals. This

finding is consistent with previous studies of human (55) and mouse spinal cord tracts (56)

which have demonstrated thicker myelin with aging. With thin myelin sheaths surrounding both

degenerating axons and axons without degenerative components (Figure 5D) we further

analyzed axons with thin myelin sheaths (g-ratio $\geq$0.8) for the presence or absence of

degenerative features. Thinly myelinated axons without degenerative features, likely

representing remyelinated axons, were present in similar proportions in aged and young

adoptive transfer lesions (Figure 5H). Thinly myelinated axons with degenerative features were

also present at similar proportions in young and aged lesions (Figure 5I). Relative frequency

distribution analysis of axonal cross-sectional area of thinly myelinated axons classified as

degenerating compared to non-degenerating indicated that degenerating axons had a

significantly higher frequency of large axonal areas (Figure 5J) as would be expected for

degenerating axons.

We acknowledge that thin myelin sheaths may represent remyelinated axons or axons in

the process of demyelination and our time-point analysis at 10 days-post adoptive transfer

would require fast kinetics for OPC differentiation and remyelination. Young and aged OPC

remyelination reporter mice *Pdgfra-Cre^ERT;Tau^mGFP* that received adoptive transfer were

processed for TEM with immunogold staining for GFP to attempt to further validate OPC

remyelination. However, due to GFP reporter expression in OPC processes in the parenchyma

(Figure 5B) and anti-GFP immunogold deposition in similar pattern, analysis of

TEM/immunogold was not further pursued. Despite these limitations, remyelination appears

similar between age groups, even in the presence of a robust inflammatory infiltrate and severe

disease model.

Discussion

OPCs are dynamic cells with the ability to self-renew, migrate, and differentiate into

mature, myelinating oligodendrocytes. Although the response of OPCs to demyelination has

been investigated in toxin-mediated models and genetic ablation strategies (5), the fate of OPCs

after exposure to a robust auto-immune inflammatory insult remains poorly understood.

Biological aging influences multiple sclerosis disease activity and progression (13) and the

response of OPCs in an aged environment is particularly relevant for a chronic CNS

autoimmune disease that affects individuals at all stages of life. Physiological aging in the

absence of a disease model is associated with a decreased rate of OPC proliferation (14,16)

and differentiation into mature OLs (14). Aged OPCs express higher levels of the cell cycle

inhibitors p16 and p21, which may reflect age-associated senescence (57). In response to toxin-

mediated demyelination insults, aged OPCs have slower proliferation, recruitment, and

differentiation kinetics; however, remyelination in aged animals eventually catches up to young

animals (17-19).

To determine whether aging impacts the response of OPCs to a strong adaptive immune

response akin to an acute MS attack, we performed lineage tracing of OPCs in young adult and

aged spinal cord after adoptive transfer of myelin reactive Th17 polarized T cells. Both aged and

young OPCs respond to an acute inflammatory demyelination event with a proportionally similar

increase in lesional OPC density. The highest densities of lineage traced OPCs in both aged

and young animals were within lesions, suggesting that compensatory mechanisms allow for

recruitment, survival and persistence of OPCs within a lesion environment. Both young and

aged OPCs can differentiate into mature OLs, a response that is less robust in aged OPCs.

While existing differentiated OLs sustain damage after adoptive transfer, the magnitude of aged

OL loss is more severe. Although the presence of ongoing axonal degeneration in adoptive

transfer and other models can complicate the interpretation of thin myelin sheaths by TEM

analysis (53), we quantified thin myelin sheaths by TEM and found a similar proportion of non-

degenerative and likely remyelinated axons in lesions of young and aged animals. With TEM we

were unable to distinguish whether remyelination occurs from newly generated

oligodendrocytes versus existing differentiated oligodendrocytes, both of which can occur in

toxin-mediated models (4). Further lineage tracing studies using other transgenic membrane

targeted reporter lines may offer additional insight into the origin of remyelinating

oligodendroglia in immune-mediated models.

352   Previous studies combining genetic CNS interferon gamma production with toxin-

mediated demyelination indicate that OPCs may by targets of CD8 T cell mediated killing (33).

Our findings in the Th17 adoptive transfer model indicate that an inflammatory lesion

environment stimulates OPCs responses and both young and aged OPCs may have

compensatory strategies that allow them to survive and differentiate despite the presence of T

cells. Interferon-response pathways are upregulated in disease-associated oligodendroglia (25-

32) and with physiological aging (16, 58) which may be a result of CD8 T cell accumulation

within the aged CNS (58). Oligodendrocytes upregulate MHC class I (59) and immune

checkpoint inhibitor programmed death-ligand 1 (PD-L1) (60) in CNS viral models. Upregulation

of interferon-response pathway in young and aged OPCs may in fact be beneficial for

oligodendroglial survival allowing OPCs and OLs to evade the immune attack and dampen

immune signaling. Further studies are needed to investigate the role of disease-associated

oligodendroglia in inflammatory models to determine whether immunoregulatory pathways in

oligodendroglia are beneficial or inhibitory to oligodendroglial function.

This study demonstrates that oligodendroglia in young and aged animals respond

similarly to an acute auto-immune mediated demyelinating attack mediated by MOG-reactive

Th17 T cells. The ability of both young and aged OPCs to respond to a severe autoimmune

attack and preservation of OPCs within lesions suggests that remyelination and repair strategies

that target oligodendroglial survival mechanisms may be promising strategy for people of all

ages with MS.

Methods

Sex as a biological variable

Both male and female mice transgenic and wild-type mice were used as adoptive transfer

recipients with a roughly equal even distribution between female and male mice in young and

aged cohorts. Due to the research strategy involving use of transgenic lines, timing of adoptive

transfer cohorts and need for aged cohorts, all available positive transgenic mice were used

regardless of sex and the experiments were not sufficiently powered to assess differences

between sexes. Wild-type mice used for immunization for donor T cells were female only to

avoid potential sex-based T cell differences.

Animal care and use

Young (P87-P148, 3-5-month-old) and aged (P295-P427,10-14-month-old) *Pdgfra-Cre^{ERT}*

hemizygote +/- (Jackson Laboratory 018280); *RCE- Rosa26^{flstopflEGFP}* hemizygote +/- (Jackson

Laboratory 032037-JAX, MMRRC stock #32027). Young (P131, 4-month-old) and aged (P366-

P426, 12-14-month-old) C57BL/6J (Jackson Laboratory 000664) wild-type mice and *Pdgfra-*

Cre^{ERT} hemizygote +/- (Jackson Laboratory 018280); *Tau^{mGFP}* hemizygote +/- (Jackson

Laboratory 021162) were used for transmission electron microscopy experiments. Age indicated

is animal age at sacrifice. Female and male mice were assigned to experiments based on

availability. Mice were maintained on a 12-hour light/dark cycle, housed in cages of five or less,

and provided food and water ad libitum. Adoptive transfer mice were monitored daily between

days 4-10 post-adoptive transfer for clinical score and weight. All animal experiments were

conducted in accordance with the National Institute of Health Guide for the Care and Use of

Laboratory Animals.

Adoptive transfer/Experimental autoimmune encephalomyelitis (EAE)

Transgenic mice *Pdgfra-Cre^{ERT}*; *RCE* and *Pdgfra-Cre^{ERT};Tau^{mGFP}* were administered tamoxifen

(Sigma T5648) intraperitoneally (100-150ul of 10mg/ml stock reconstituted in corn oil for

working dose of 75mg/kg/day) daily for 5 days to induce recombination starting two weeks prior

to adoptive transfer. Wild-type C57BL/6J mice used for transmission electron microscopy (TEM)

did not receive tamoxifen. T cells for adoptive transfer were generated from wild-type C57BL/6J

(Jackson Laboratory 000664) 2-3-month-old female mice which were anesthetized with

intraperitoneal injection of ketamine (90mg/kg)/ xylazine (10mg/kg) and immunized with four

subcutaneous flank injections with a total of 0.2-0.4mg of myelin oligodendrocyte glycoprotein

(MOG)$_{35-55}$ peptide (1mg/ml, biosynthesis) resuspended in Complete Freund's Adjuvant (CFA)

with concentration of 5mg/ml heat killed *M. tuberculosis*. After 10 days immunized wild-type

mice were euthanized with CO_2 and spleen and lymph nodes were dissected and placed on ice

in EAE media: RPMI (ThermoFisher 21870084), 10% FBS (GeminiBio 100-106-500), P/S

(Fisher 15-140-148), 1x GlutaMAX (Fisher 35-050-061), 1x NEAA (ThermoFisher 11140050),

1mM sodium pyruvate (Fisher 11-360-070), 55nM beta-mercaptoethanol (ThermoFisher

21985023). Spleens and lymph nodes were mashed across filter, washed with EAE media, and

centrifuged at 800g 5min. Spleens were resuspended in RBC lysis buffer (ThermoFisher 00-

4300-54) for 2 minutes at room temperature followed by quenching with EAE media, centrifuged

at 800g 5min, and washed with EAE media. Spleen and lymph nodes were resuspended in EAE

media with MOG_{35-55} peptide (50μg/ml, Biosynthesis), recombinant mouse IL-23 (8ng/ml, R&D

1887-ML), recombinant mouse IL-1a (10ng/ml, ThermoFisher 211-11A), anti-IFNg (10mg/ml,

Bio X Cell BE0055) at volume of $5x10^6$ cells/ml in T75 flasks. Cells suspensions were incubated

at 37°C in 5% CO_2 incubator for 3.5 days. After *ex-vivo* Th17 polarization cells were collected

into conical tubes, centrifuged 800g 5min, filtered and washed twice with MACs buffer (0.5%

BSA and 2mM EDTA in PBS). CD4 T cell selection was performed with mouse CD4 (L3T4)

microbeads (Miltenyi Biotec 130-117-043) with LS column per Miltenyi protocol. Eluted CD4+

cells were washed in MACs buffer twice and $4x10^6$ cells were injected intraperitoneally into

transgenic mice (tamoxifen administered) and wild-type mice (no tamoxifen). For naïve/non-

adoptive transfer animals, littermates of transgenic mice that received tamoxifen were separated

into naïve cages. Mice were monitored daily starting on day 4 post-adoptive transfer with

weights and EAE clinical scoring scale. EAE scoring scale: 0.5 reduced tail tone; 1 complete

loss of tail tone; 1.5 slight gait imbalance; 2 gait imbalance with footfalls or spinning/ataxia; 2.5

hindlimb foot drag able to move above hip; 3 partial to complete hindlimb paralysis, not able to

move either hindlimb above hip, moving around cage with forelimbs; 3.5 complete hindlimb

paralysis, hindlimbs held to one size, moving around cage with forelimbs; 4 complete hindlimb

paralysis, hindlimbs held to one size, no movement around cage but alert; 4.5 moribund not

alert; 5 death. Mice were sacrificed at days 9-10 post-adoptive transfer with most severe mice

sacrificed on day 9 time-point to avoid death and following end-point euthanasia protocols

(euthanasia at score 4 or greater within 24 hours or sustained weight loss $\geq$30% for 3 days). All

mice in this study were between clinical scores of 2-4.5 at day of sacrifice and a total of 5

separate cohorts of adoptive transfer were performed. EAE clinical score graphs from mice

used for immunohistochemistry and transmission electron microscopy including several

additional cohorts that were used to optimize TEM fixation protocol were combined for EAE

clinical score graphs. There was no significant difference between mice that received tamoxifen

and those that did not. Animals that were sacrificed due to endpoint criteria or reached score 5

(death) were included as data-points with score 5 on subsequent days on EAE clinical scoring

graphs.

Immunohistochemistry

Mice were anesthetized with intraperitoneal injection of ketamine (90mg/kg)/ xylazine (10mg/kg)

and perfused transcardially with 15ml of 0.1M phosphate-buffered saline (PBS) followed by cold

4% paraformaldehyde (PFA) in PBS (ThermoScientific J19943.K2). Spinal cords were dissected

out of the spinal column and post-fixed in 4% PFA/PBS at 4°C overnight then cryoprotected in

30% sucrose/1xPBS at 4°C for at least 2 days. Prior to embedding two transverse cuts were

performed (at the end of the cervical enlargement and beginning of lumbar enlargement) to

separate the spinal cord into cervical, thoracic and lumbar segments. Spinal cord segments

were embedded in Tissue-Tek OCT compound (Sakura) with a cross-section of the beginning of

each spinal cord segment facing the cutting edge of the block. Spinal cord blocks were

sectioned at 15μm thickness with Leica CM1950 cryostat onto Superfrost Plus microscope

slides (Fisher 12-550-15) with each 8-10 series of slides having four sets of

cervical/thoracic/lumbar transverse sections (12 total sections) representing ~600mm distance

from first to fourth section in each region to obtain representative analysis across each spinal

cord segment on one slide. Slides were stored at -80°C until use. For immunostaining slides

were permeabilized with 1xPBS +0.1% Triton-X100 (PBST) for 10 minutes followed by outlining

of sections with PAP pen (ThermoFisher 008899). To prevent non-specific binding, 5% normal

goat serum (JacksonImmuno 005-000-121) in PBST was added to sections for 1 hour at room

temperature followed by addition of primary antibodies in same blocking solution overnight at

4°C. Slides were washed in PBST three times 5 minutes each followed by incubation in

secondary antibodies in blocking buffer for 1 hour at room temperature. Slides were washed

with PBST twice for 5 minutes each, once with PBS for 5 minutes then mounted with

ProlongGold antifade (ThermoFisher P36935). For FluoroMyelin co-stained slides, FluoroMyelin

Red (Invitrogen F34652) was added at 1:300 dilution in PBS after last wash and incubated for

20 minutes followed by PBS wash and mounting. The following primary antibodies were used:

chicken anti-GFP (Abcam ab13970, 1:1000), rabbit anti-Olig2 (MilliporeSigma ab9610, 1:1000),

rabbit anti-Aspa (MilliporeSigma ABN1698, 1:250), rat anti-CD45 (MilliporeSigma 05-1416,

1:500), rabbit anti-degraded myelin basic protein dMBP (MilliporeSigma ab5864, 1:1000), rabbit

anti-Pdgfra (CellSignaling 3174, 1:500), mouse anti-neurofilament heavy NF-H (EnCor MCA-

AH1), rat ant-MBP (Biorad MCA409S, 1:1000), mouse anti-neurofilament medium NF-M

(Invitrogen 130700, 1:1000, for remyelination reporter staining). The following secondary

antibodies were used all purchased from Invitrogen and used at dilution of 1:500: Alexa Fluor

488 goat anti-chicken IgY (H+L) (A-11039), Alexa Fluor 594 goat anti-rat IgG (H+L) (A-11007),

Alexa Fluor 647 goat anti-rat IgG (H+L) (A-21247), Alexa Fluor 647 goat anti-rabbit IgG (H+L)

(A-21245), Alexa Fluor 647 goat anti-mouse IgG (H+L) (A-21236), and Alexa Fluor 594 goat

anti-mouse IgG (H+L) (A-11032).

Immunohistochemistry imaging and analysis

Images of immunohistochemistry-stained spinal cord sections were acquired using Olympus

IX83 inverted microscope with Zeiss ZEN microscopy software. Tiled images of transverse

spinal cord sections at a resolution of 650nm/pixel (10x objective) were obtained for GFP/Olig2,

GFP/Aspa, and dMBP stains. Higher resolution tiled images at 325nm/pixel (20x objective) were

taken for GFP/Aspa/CD45, neurofilament and FluoroMyelin stains. For manual-counting, image

stacks were analyzed in ImageJ/Fiji using channels tool, cell counter and ROI manager. For

macro-analyzed images (DAPI/GFP/Aspa and DAPI/GFP/NF stains), images were exported as

non-compressed TIFF grayscale stack of channels. Macros were developed using a

combination of background subtraction, moments thresholding and analyzing particle

ImageJ/Fiji functions to identify positive cells. In refinement of macros for individual stains,

quality assessment images with overlay of counted objects were analyzed for accuracy and

parameters were adjusted for each macro to obtain the highest accuracy of cell detection. After

importing the spinal cord section image stack, the macros prompt the user to outline regions of

interest (ROI)- whole cord, gray matter and lesions. Lesions were identified and outlined by the

presence of DAPI hypercellularity which correlated with CD45 infiltrate. Non-lesion white matter

counts were generated by subtracting whole cord counts from gray matter and total lesion

counts, this was due to the ease in outlining gray matter during the macro execution compared

to total non-lesion white matter. For Aspa macro counts total lesions within each spinal cord

section were outlined as one lesion. In NF macro and DAPI/GFP/Aspa/CD45 manual counts

individual lesions were outlined separately to increase power of analysis given potential

variability across lesions. For neurofilament quantification, DAPI/GFP/NF 20x tiled images were

analyzed with NF macro allowing users to outline lesions based on DAPI hypercellularity and

NF positive axonal density was calculated for individual lesions. For dMBP quantification

DAPI/GFP/dMBP 10x tiled image stacks were quantified in ImageJ/Fiji by outlining total white

matter area and white matter with positive dMBP staining for each spinal cord section. For

FluoroMyelin quantification, DAPI/GFP/Fluoromyelin 20x tiled image stacks were quantified by

outlining total white matter and demyelinated white matter without Fluoromyelin. To generate

non-biased transverse spinal cord cross-section counts throughout the cervical, thoracic and

lumbar spinal cord, all spinal cord sections without significant folds or other tissue processing

artifacts were imaged. Adoptive transfer sections without lesions were not quantified to focus on

spinal cord regions with inflammatory infiltrate. Sections quantified per animal: GFP/Olig2 6-16,

GFP/Aspa/CD45 hand counts 3-6, GFP/Aspa macro 6, GFP/NF macro 3-8, GFP/FlouroMyelin

6-9, GFP/dMBP 6-12. Sections counts were not averaged across individual animals to focus on

comparisons on an individual section/region level which allowed for analysis of enrichment

within lesions compared to non-lesion white matter. Lesion to non-lesion cell density ratios were

generated by calculating the ratio of average lesional density/average non-lesional density for

each spinal cord transverse section. For remyelination reporter assessment *Pdgfra-*

Cre^{ERT};*Tau^{mGFP+/-}* ventral medial lumbar spinal cord stained with GFP/MBP/NF were imaged on

W1 SoRA spinning disk confocal with Nikon NIS-Elements using 40x oil objective (108nm/pixel)

with z-stacks and maximum intensity projections generated for ROIs.

Transmission electron microscopy (TEM)

Young (P131, 4-month-old) and aged (P366-P426, 12-14-month-old) C57BL/6J (Jackson

Laboratory 000664) wild-type mice and *Pdgfra-Cre^{ERT}* hemizygote +/- (Jackson Laboratory

018280); *Tau^{mGFP}* hemizygote +/- (Jackson Laboratory 021162) were used for transmission

electron microscopy experiments. C57BL/6J mice were perfused with 4% PFA/5%

glutaraldehyde and *Pdgfra-Cre^{ERT}*;*Tau^{mGFP}* mice for immunogold were perfused with 4%

PFA/0.25% glutaraldehyde. On day 10 post-adoptive transfer young and aged EAE mice were

anesthetized with intraperitoneal injection of ketamine (90mg/kg)/ xylazine (10mg/kg) and

perfused transcardially with 15ml of 0.1M PBS followed by cold fixation solution: PFA (EMS

15714-S) and glutaraldehyde (EMS 16210) in 80mM sodium cacodylate and 9mM calcium

chloride dihydrate in EM grade dH$_2$O (EMS 22800-01) pH 7.4. The spinal cord was dissected

out of the column and post-fixed for 24 hours in perfusion buffer (2 hours for immunogold

followed by switch to 1% PFA overnight). Spinal cord was cut into 2mm transverse segments on

spinal cord matrix (EMS 69085-C) and 3-5 lumbar and 3-5 cervical blocks per animal were

washed with wash buffer (70mM sodium cacodylate, 70mM sodium chloride, 0.5mM calcium

chloride, 1mM magnesium chloride hexahydrate in EM grade dH$_2$O) for 30 minutes followed by

post-fixation in 1% osmium tetroxide (EMS 19152) in 100mM sodium cacodylate for 1 hour

protected from light. Blocks were washed with EM grade dH$_2$O twice 2 minutes each then

transferred to glass vials followed by dehydration (50% EtOH, 70% EtOH, 80% EtOH, 95%

EtOH, 100% EtOH x 2, 100% propylene oxide 2 times, with each wash 30 minutes). Blocks

were incubated overnight in 1:1 Embed 812 resin (EMS 14120, medium hardness

concentration):propylene oxide (EMS 20412) at room temperature. Solution was changed to 3:1

Embed 812 resin:propylene oxide for 24 hours followed by 100% Embed 812 resin for 24 hours

at room temperature. Embedding molds were filled with fresh Embed 812 resin and individual

spinal cord blocks were transferred to molds and baked at 60°C for 24 hours. Semi-thin sections

from lumbar spinal cord blocks were cut on Leica Ultracut UCT at 1μm thickness with Histo

Diamond Knife (EMS 60-HIS) and stained with 1% toluidine blue to visualize lesions. Ultrathin

sections for TEM were cut on Ultra Diamond Knife (EMS 40-UL) at 70nm thickness on

Ultramicrotome Reichert Ultracut E and mounted onto grids (Pelco 200 mesh 1GC200). Grids

were stained using Hiraoka staining kit (EMS 71560-00) with uranyl acetate alternative (EMS

22405) for 20 minutes followed by rinses in EMS grade dH$_2$O 30 seconds 3 times, then

incubation in 3mg/ml lead citrate in 0.1N NaOH in dH$_2$O for 3 minutes. Grids were washed with

20mM NaOH followed by 2 washes with dH$_2$O for 30 seconds then air dried. For immunogold

grids were stained with Hiraoka staining kit washed with 0.01% Triton TBS, blocked with 1%

NGS (JacksonImmuno 005-000-121), 1% BSA (Sigma A4161), 1% fish gelatin (Sigma G7041)

in TBST (0.01% Triton TBS) for 30 minutes then incubated with anti-GFP (Abcam ab13970,

1:100) overnight at 4°C. Grids were washed with TBST 3 times, incubated in 12nm colloidal

gold donkey anti-chicken (JacksonImmuno 703-205-155) in blocking solution for 90 minutes at

room temperature. Grids were washed with TBST, post-fixed in 1% glutaraldehyde in PBS for 5

minutes, washed in EM grade dH$_2$O followed by staining for myelin as described above. Grids

were imaged on FEI Tecnai G2 Spirit TEM and ventral side was identified by the presence of

ventral central vein and ventromedial white matter regions were imaged at 1150x magnification.

All lumbar blocks were sectioned, imaged and quantified. All lumbar blocks from aged and

young animals contained lesions with notable loss of myelin and presence of immune infiltrate in

this region. TEM lesion images acquired at 1150x resolution representing one grid box were

analyzed in MyelTracer (54) with users selecting and manually outlining myelin sheaths for all

axons with $\geq$2μm given the difficulty in outlining smaller diameter axons at this magnification and

desire to quantify remyelination which is more feasible for large diameter axons. Axon diameter

and axon plus myelin diameters and calculated g-ratios were exported and compiled in excel

and g-ratios were plotted with a cutoff of 2μm axon diameter in GraphPad Prism. For

quantification of thin myelin sheaths with or without degenerative axon components TEM lesion

images acquired at 1150x resolution were inverted in ImageJ/Fiji for highlighting myelin sheaths

and degenerative axon components. Axons with g-ratio of 0.8 or less with a high number of

mitochondria, myelin infoldings, unraveling or blebbing, and/or electron dense material were

counted as thinly myelinated axons with degenerative features. Axons with g-ratio of 0.8 or less

without these features were counted as without degenerative features and likely remyelinated

axons. The percentage of these two types of axons was quantified over total myelinated axons

with 2μm or greater axon diameter for each 1150x lesion TEM image. Total myelinated axons

with 2μm or greater axon diameter were counted by using StarDist ImageJ/Fiji plugin followed

by manual confirmation and adjustment as needed.

Flow cytometry

Transgenic *Pdgfra-Cre^{ERT}*; *RCE- Rosa26^{flstopflEGFP}* mice that received tamoxifen and Th17 T cells

or no Th17 T cells (naïve) as described above were sacrificed at 9 days post adoptive transfer

for flow cytometry analysis of GFP reporter expression. Mice were anesthetized with

intraperitoneal injection of ketamine (90mg/kg)/xylazine (10mg/kg) and perfused transcardially

with 15ml of 0.1M PBS. Brains and spinal cord were dissected and dissociated in 20U/ml papain

and 100U/ml of DNase in papain buffer (1mM EDTA, 5mM L-cystine, EBSS pH 7.4-7.6) for 30-

45 minutes in 37°C 5% CO_2 incubator with 3-4 manual trituration steps during incubation to

dissociate into single cell suspension. Dissociated cells were spun at 800g 5min 4°C and

resuspended in 30% Percoll (VWR 89428-524) in FACs buffer (2% FBS in PBS no cations) and

spin 1000g 10 minutes 4°C speed 8. Myelin layer was removed and pellet was resuspended in

FACs buffer followed by transfer over 70mm filter, PBS wash and resuspension in PBS with Fc

block (BD Biosciences rat anti-mouse CD16/CD32 552142, 1:200) and viability stain

(eBioscience 65-0865-14, 1:1000) for 10 minutes on ice. Cells were washed in FACs buffer and

resuspended in primary antibody CD45 e450 (Invitrogen 48-0451-82 1:100) in FACs and

incubated 30 minutes on ice followed by wash and resuspension in FACs. Flow cytometry of

spinal cord and brain samples was performed with BD FACS Melody 4-way sorter and analyzed

in FlowJo software.

Statistics and reproducibility

GraphPad Prism was used to generate graphs and perform statistical analysis. EAE scoring and

quantification was blinded by use of genotyping animal ID that was later linked to a sample

group in GraphPad prism analysis. A target sample size of 6 mice per group for

immunohistochemistry and 4 mice per group for TEM were set for our analysis. For IHC

analysis, average EAE score at sacrifice for young was 3.5 (minimum 2.5, maximum 4.5) and

aged 3.8 (minimum 2.5, maximum 4.5) with male and females in both groups. A minimum of 4

animals per group and 3 sections per animal were analyzed for IHC outcomes and individual

section and lesion data-points were represented to allow for more robust statistics given

potential variability on a micro and macroenvironment level within different lesions and sections.

For transmission electron microscopy on wild-type mice a total of 4 young and 6 aged wild-type

Th17 AT mice were sacrificed and processed with an average EAE score at sacrifice for young

was 2.3 (minimum 2, maximum 2.5) and aged 3.6 (minimum 2.5, maximum 4) with males and

females in both groups. All lumbar spinal cord blocks were quantified from these animals. In

each figure and legend, the number of biological replicates and regions of interest (ROI)

quantified, statistical test and significance levels are indicated. Due to complexity of variability

including recombination, EAE severity, region, lesional difference and age; datasets did not

meet normality assumptions. Statistical significance was assessed using nonparametric Mann-

Whitney (two-tailed) and 2-sample median bootstrap (100,000 resamples, two-tailed, shift

bootstrap, BCa bias-corrected & accelerated), with bootstrap preferred with better control for

Type I errors and unequal variances (61).

Study approval

All animal experiments were conducted in accordance with the National Institute of Health

Guide for the Care and Use of Laboratory Animals and were approved by The Ohio State

University Institutional Animal Care and Use Committee (IACUC) protocol 2023A00000086.

Data availability

Data presented in available in the Supporting Data Values file. All macro codes have been

made publicly available in GitHub-https://github.com/coleayden/spinalcordmacros. For

additional data inquires contact the corresponding author

Author contributions

CAH conceptualized the project. WS, MAW, and BB provided scientific guidance. EEF and CAH

conducted the experiments. EEF, CNB, NFA and BJT performed quantification of histology. BJB,

EEF, CNB, and NFA performed quantification of transmission electron microscopy. DP

developed initial macro code and CAH modified macro codes. CAH wrote the manuscript. All

authors contributed to editing and finalizing the manuscript.

Funding support

This study was funded by The Ohio State University College of Medicine Research Innovator

Career Development Award (RICDA) and American Academy of Neurology Career

Development Award. BJT was funded by the National Multiple Sclerosis Society Undergraduate

Summer Research Program.

Acknowledgements

We would like to thank the following contributors: Dr. Benjamin M. Segal for scientific

mentorship and access to shared equipment used for this study. Dr. Dana McTigue and Dr.

Anthony Brown for assistance with transmission electron microscopy protocols and equipment

usage. For equipment used for TEM sectioning and TEM imaging we acknowledge resources

from the Campus Microscopy and Imaging Facility (CMIF) and the OSU Comprehensive Cancer

Center (OSUCCC) Microscopy Shared Resource (MSR), The Ohio State University. This facility

is supported in part by grant P30 CA016058, National Cancer Institute, Bethesda, MD. We

would like to thank Jeffery Tonniges CMIF director for providing personal training for TEM

ultrathin sectioning.

References

1. Bergles DE, Richardson WD. Oligodendrocyte Development and Plasticity. *Cold Spring Harb*
   Perspect Biol. 201;8(2):a020453.

2. Zhang X,et al. Replenishing the Aged Brains: Targeting Oligodendrocytes and Myelination?
   Front Aging Neurosci. 2021;13:760200.

3. Rawji KS, Neumann B, Franklin RJM. Glial aging and its impact on central nervous system
   myelin regeneration. *Ann N Y Acad Sci*. 2023;1519(1):34–45.

4. Franklin RJM, Frisén J, Lyons DA. Revisiting remyelination: Towards a consensus on the
   regeneration of CNS myelin. *Semin Cell Dev Biol*. 2021;116:3–9.

5. Tepavčević V, Lubetzki C. Oligodendrocyte progenitor cell recruitment and remyelination in
   multiple sclerosis: the more, the merrier? *Brain*. 2022;145(12):4178–92.

6. Gold R, Linington C, Lassmann H. Understanding pathogenesis and therapy of multiple
   sclerosis via animal models: 70 years of merits and culprits in experimental autoimmune
   encephalomyelitis research. *Brain*. 2006;129(Pt 8):1953–71.

7. Lassmann H. Multiple sclerosis: lessons from molecular neuropathology. *Exp Neurol*.

2014;262 Pt A:2–7.

8. Franklin RJM, Simons M. CNS remyelination and inflammation: From basic mechanisms to

therapeutic opportunities. *Neuron*. 2022;110(21):3549–65.

9. Chang A, et al. NG2-positive oligodendrocyte progenitor cells in adult human brain and

multiple sclerosis lesions. *J Neurosci*. 2000;20(17):6404–12.

10. Chang A, et al. Premyelinating oligodendrocytes in chronic lesions of multiple sclerosis. *N

Engl J Med*. 2002;346(3):165–73.

11. Antel JP, et al. Immunology of oligodendrocyte precursor cells in vivo and in vitro. *J

Neuroimmunol*. 2019;331:28–35.

12. Stankoff B, et al. Age at onset determines the occurrence of the progressive phase of

multiple sclerosis. *Neurology*. 2007;68(10):779–81.

13. Graves JS, et al. Ageing and multiple sclerosis. *Lancet Neurol*. 2023;22(1):66–77.

14. Spitzer SO, et al. Oligodendrocyte Progenitor Cells Become Regionally Diverse and

Heterogeneous with Age. *Neuron*. 2019;101(3):459–71.e5.

15. Neumann B, et al. Metformin Restores CNS Remyelination Capacity by Rejuvenating Aged

Stem Cells. *Cell Stem Cell*. 2019;25(4):473–85.e8.

16. Heo D, et al. Transcriptional profiles of mouse oligodendrocyte precursor cells across the

lifespan. *Nat Aging*. 2025;5(4):675–90.

17. Shields SA, et al. Remyelination occurs as extensively but more slowly in old rats compared

to young rats following gliotoxin-induced CNS demyelination. *Glia*. 1999;28(1):77–83.

18. Sim FJ, et al. The age-related decrease in CNS remyelination efficiency is attributable to an

impairment of both oligodendrocyte progenitor recruitment and differentiation. *J Neurosci*.

2002;22(7):2451–9.

19. Shen S, et al. Age-dependent epigenetic control of differentiation inhibitors is critical for

remyelination efficiency. *Nat Neurosci*. 2008;11(9):1024–34.

20. Ruckh JM, et al. Rejuvenation of regeneration in the aging central nervous system. *Cell*

Stem Cell. 2012;10(1):96–103.

21. Lassmann H. The changing concepts in the neuropathology of acquired demyelinating

central nervous system disorders. *Curr Opin Neurol*. 2019;32(3):313–9.

22. Rangachari M, Kuchroo VK. Using EAE to better understand principles of immune function

and autoimmune pathology. *J Autoimmun*. 201;45:31–9.

23. Kipp M, et al. Multiple sclerosis animal models: a clinical and histopathological perspective.

Brain Pathol. 2017;27(2):123–37.

24. Packer D, Fresenko EE, Harrington EP. Remyelination in animal models of multiple

sclerosis: finding the elusive grail of regeneration. *Front Mol Neurosci*. 2023;16:1207007.

25. Shen K,et al. Multiple sclerosis risk gene Mertk is required for microglial activation and

subsequent remyelination. *Cell Rep*. 2021;34(10):108835.

26. Pandey S, et al. Disease-associated oligodendrocyte responses across neurodegenerative

diseases. *Cell Rep*. 202;40(8):111189.

27. Hou J, et al. Transcriptomic atlas and interaction networks of brain cells in mouse CNS

demyelination and remyelination. *Cell Rep*. 2023;42(4):112293.

28. Aboelnour EL, et al. A comparative transcriptomic analysis of mouse demyelination models

and Multiple Sclerosis lesions. [preprint]. http://dx.doi.org/10.1101/2025.03.24.645058.

Posted on bioRxiv. March 25, 2025.

29. Kukanja P,et al. Cellular architecture of evolving neuroinflammatory lesions and multiple

sclerosis pathology. *Cell.* 2024;187(8)1990-2009.

30. Falcão AM, et al. Disease-specific oligodendrocyte lineage cells arise in multiple sclerosis.

Nat Med. 2018;24(12):1837–44.

31. Harrington EP, et al. MHC class I and MHC class II reporter mice enable analysis of immune

oligodendroglia in mouse models of multiple sclerosis. *Elife*. 2023;12:eLife82938.

32. Jäkel S, et al. Altered human oligodendrocyte heterogeneity in multiple sclerosis. Nature.

2019;566(7745):543–7.

33. Kirby L, et al. Oligodendrocyte precursor cells present antigen and are cytotoxic targets in

inflammatory demyelination. *Nat Commun*. 2019;10(1):3887.

34. Harrington EP, Bergles DE, Calabresi PA. Immune cell modulation of oligodendrocyte

lineage cells. *Neurosci Lett*. 2020;715:134601.

35. Kang SH, et al. NG2+ CNS glial progenitors remain committed to the oligodendrocyte

lineage in postnatal life and following neurodegeneration. *Neuron*. 201;68(4):668–81.

36. Sousa VH, et al. Characterization of Nkx6-2-derived neocortical interneuron lineages. *Cereb*

Cortex. 2009;19 Suppl 1(Suppl1):i1–10.

37. Doss PMIA, et al. Male sex chromosomal complement exacerbates the pathogenicity of

Th17 cells in a chronic model of central nervous system autoimmunity. *Cell Rep*.

2021;34(10):108833.

38. Fletcher JM, et al. T cells in multiple sclerosis and experimental autoimmune

encephalomyelitis. *Clin Exp Immunol*. 2010;162(1):1-11.

39. Maddur MS, et al. Th17 cells: biology, pathogenesis of autoimmune and inflammatory

diseases, and therapeutic strategies. *Am J Pathol*. 2012;181(1):8-18.

40. Yanai S, Endo S. Functional Aging in Male C57BL/6J Mice Across the Life-Span: A

Systematic Behavioral Analysis of Motor, Emotional, and Memory Function to Define an

Aging Phenotype. *Front Aging Neurosci*. 2021;13:697621.

41. Tremlett H, et al. Natural history comparisons of primary and secondary progressive multiple

sclerosis reveals differences and similarities. *J Neurol*. 2009;256(3):374-381.

42. D'Amico E, et al. Late-onset and young-onset relapsing-remitting multiple sclerosis:

evidence from a retrospective long-term follow-up study. *Eur J Neurol*. 2018;25(12):1425-

1431.

43. Bebo BF Jr, et al. Treatment with selective estrogen receptor modulators regulates myelin

specific T-cells and suppresses experimental autoimmune encephalomyelitis. *Glia*.

2009;57(7):777-790.

43. Mei F, et al. Accelerated remyelination during inflammatory demyelination prevents axonal

loss and improves functional recovery. *Elife*. 2016 Sep 27;5:e18246.

44. Wang, F, et al*.* Myelin degeneration and diminished myelin renewal contribute to age-

related deficits in memory. *Nature Neurosci*. 2020;23(4):481-486.

45. Chen, J, et al*.* Enhancing myelin renewal reverses cognitive dysfunction in a murine model

of Alzheimer's disease. *Neuron*. 2021;109(14):2292-2307.e5

46. Atkinson JR, et al. Biological aging of CNS-resident cells alters the clinical course and

immunopathology of autoimmune demyelinating disease. JCI Insight. 2022;7(12):e158153.

47. Stromnes IM, et al. Differential regulation of central nervous system autoimmunity by T(H)1

and T(H)17 cells. *Nat Med*. 2008;14(3):337–42.

48. Jäger A, et al. Th1, Th17, and Th9 effector cells induce experimental autoimmune

encephalomyelitis with different pathological phenotypes. *J Immunol*. 2009;183(11):7169–77.

49. Zaimi A, et al. AxonDeepSeg: automatic axon and myelin segmentation from microscopy

data using convolutional neural networks. *Sci Rep*. 2018;8(1):3816.

50. Blakemore WF. Demyelination of the superior cerebellar peduncle in the mouse induced by

cuprizone. *J Neurol Sci*. 1973;20(1):63–72.

51. Blakemore WF. Pattern of remyelination in the CNS. *Nature*. 1974;249(457):577–8.

52. Neumann B, et al. Problems and Pitfalls of Identifying Remyelination in Multiple Sclerosis.

Cell Stem Cell. 2020;26(5):617–9.

53. Schäffner E, et al. Myelin insulation as a risk factor for axonal degeneration in autoimmune

demyelinating disease. *Nat Neurosci*. 2023;26(7):1218-1228.

54. Kaiser T, et al. MyelTracer: A semi-automated software for myelin g-ratio quantification.

eNeuro. 2021;8(4):ENEURO.0558–20.2021.

55. Poulen G, et al. Sex and age differences in glia and myelin in nonhuman primate and

human spinal cords: implications for pathology. *Cell Death Discov*. 2025;11(1):129.

56. Xie F, et al. Effects of normal aging on myelin sheath ultrastructures in the somatic

sensorimotor system of rats. *Mol Med Rep*. 2014;10(1):459–66.

57. Zhang P, et al. Senolytic therapy alleviates Aβ-associated oligodendrocyte progenitor cell

senescence and cognitive deficits in an Alzheimer's disease model. *Nat Neurosci*.

2019;22(5):719–28.

58. Kaya T, et al. CD8+ T cells induce interferon-responsive oligodendrocytes and microglia in

white matter aging. *Nat Neurosci*. 2022;25(11):1446–57.

59. Pan R, et al. Oligodendrocytes that survive acute coronavirus infection induce prolonged

inflammatory responses in the CNS. *Proc Natl Acad Sci U S A*. 2020;117(27):15902–10.

60. Phares TW, et al. Target-dependent B7-H1 regulation contributes to clearance of central

nervous system infection and dampens morbidity. *J Immunol*. 2009;182(9):5430–8.

61. Johnston MG, Faulkner C. A bootstrap approach is a superior statistical method for the

comparison of non-normal data with differing variances. *New Phytol*. 2021;230(1):23-26.

Figure 1- OPC lineage tracing strategy in young and aged Th17 adoptive transfer

(**A**). Experimental strategy for MOG-reactive Th17 adoptive transfer (AT) in young and aged *Pdgfra-Cre^ERT^;RCE* OPC lineage traced mice (for immunohistochemistry), wild-type C57BL/6J (for TEM), and *Pdgfra-Cre^ERT^;Tau^mGFP^* OPC remyelination mice (for TEM immunogold). No adoptive transfer young and aged *Pdgfra-Cre^ERT^;RCE* OPC lineage traced mice (for immunohistochemistry) served as naïve controls. Created in BioRender. (**B**). EAE clinical scoring of all young and aged adoptive transfer animals. Aged adoptive transfer animals have significantly higher EAE clinical scores compared to young animals. All young and aged experimental animals pooled. Two-way repeated measures ANOVA, age ** p≤0.001. Non-linear regression sigmoidal variable slope lines, error bars 95% CI. (**C**). Kaplan-Meier survival analysis of probability of EAE score of 4 or greater (quadriplegia, euthanasia endpoint) in young and aged adoptive transfer with significant difference in time to score 4 endpoint. Logrank (Mantel-Cox) test, * p≤0.05. (**D**). Representative images of *Pdgfra-Cre^ERT^;RCE* Th17 adoptive transfer spinal cord with lineage-traced OPCs labeled with GFP antibody and Pdgfra. (**E**). Quantification of OPC recombination efficiency in naïve young and aged *Pdgfra-Cre^ERT^;RCE* by percentage of GFP+Pdgfra+ of total Pdgfra+ cells in whole spinal cord section. Data points mean individual animals, n=6 animals. (**F**). Representative images of *Pdgfra-Cre^ERT^;RCE* Th17 adoptive transfer spinal cord co-labeled with pan-oligodendroglial marker Olig2. (**G**). Lineage traced GFP+ OPCs express Olig2 (arrows) in spinal cord non-lesion and lesion areas. GFP+Olig2- cells (arrowheads) are present particularly within lesions. (**H**). Quantification of the percentage of lineage traced GFP+Olig2+ of total GFP+ cells in whole cord (wc) and lesions. Percentage of GFP+ cells that are oligodendroglia (Olig2+) are significantly lower in adoptive transfer wc and lesions compared to naïve wc in both young and aged animals. Median shift Bootstrap test, ****p<0.0001. (**I**). Representative image of lineage-traced *Pdgfra-Cre^ERT^;RCE* adoptive transfer spinal cord lesion co-labeled with GFP, Pdgfra and CD45. GFP+Pdgfra+CD45- OPCs (arrows) and GFP+Pdgfra+CD45+ cells (arrowhead) are present within lesions. (**J**). Flow cytometry of lineage traced *Pdgfra-Cre^ERT^;RCE* naïve and adoptive transfer brain and spinal cord with GFP+CD45+ cells (box) in adoptive transfer tissue. Box and whiskers plots: box 25 to 75 percentile, whiskers min to max, median line, show all datapoints. Scale bars: D,F 200μm; G,I 50μm.

Figure 2- Young and aged lineage traced OPCs are enriched within adoptive transfer lesions

(**A**). Representative images of young and aged lineage traced OPC (*Pdgfra-Cre^{ERT};RCE*) naïve and adoptive transfer spinal cord cross-sections stained for pan-oligodendroglial marker (Olig2), lineage traced OPCs (GFP) and DAPI (lesions outlined). (**B**). Quantification of lineage traced OPC (Olig2+GFP+) densities in whole cord (wc), adoptive transfer (AT) non-lesion white matter (nlwm) and adoptive transfer lesions. Lesion white matter density datapoints are means of all lesions in an individual spinal cord section. Median shift Bootstrap test. (**C**). Quantification of density of lineage traced OPCs that remained OPCs (OPC->OPC, GFP+Aspa-CD45-) in young and aged adoptive transfer wc, nlwm, and individual lesions. OPC->OPC density is significantly higher in lesions compared to nlwm in young and aged AT. OPC->OPC density is significantly lower in aged compared to young AT lesions. Median shift Bootstrap test. (**D**). Quantification of the ratio of mean OPC->OPC density in lesions compared to nlwm across a whole spinal cord section. OPC->OPC density is enriched in lesions compared to surrounding nlwm in both young and aged AT and lesion/nlwm ratios are not significantly different between ages. Median shift Bootstrap test. (**E**). Quantification of density of lineage traced OPCs that differentiated into OLs (OPC->OL, GFP+Aspa+CD45-) in young and aged adoptive transfer wc, nlwm, and individual lesions. Differentiated OPCs OPC->OL density is significantly higher in lesions compared to nlwm in young and aged AT. Median shift Bootstrap test. (**F**). Quantification of the ratio of mean OPC->OL density in lesions compared to nlwm across a whole spinal cord section. OPC->OL lesion/nlwm ratios are not significantly different between ages. Median shift Bootstrap test. (**G**). Representative images of young and aged *Pdgfra-Cre^{ERT};RCE* adoptive transfer spinal cord stained for GFP, Aspa, and CD45. OPCs that have differentiated (GFP+Aspa+) are indicated with arrows. Lesions outlined by presence of DAPI hypercellularity (not shown) and CD45 infiltrate. (**H**). Quantification of percentage of lineage traced OPCs that remain OPCs (GFP+Aspa-CD45- of GFP+CD45-) within young and aged individual lesions. Median shift Bootstrap test. Box and whiskers plots: box 25 to 75 percentile, whiskers min to max, median line, show all datapoints. p values: ns p>0.05, * p≤0.05, **p≤0.01, ***p≤0.001, ****p<0.0001. Scale bars: A, 200μm; F 50μm.

Figure 3- Differentiated oligodendrocytes are reduced in adoptive transfer

(**A**). Representative images of spinal cord sections from young and aged lineage traced OPC *Pdgfra-Cre^{ERT};RCE* adoptive transfer and naïve spinal cord sections stained for differentiated oligodendrocyte (OL) marker (Aspa), lineage traced OPCs (GFP) and DAPI (lesion areas outlined). (**B**). Representative adoptive transfer spinal cord section stained for DAPI/GFP/Aspa and processed with GFP/Aspa ImageJ macro with Aspa-positive macro counted cells outlined in red overlayed on raw grayscale of Aspa staining (bottom panel). (**C**). Quantification of Aspa+ OL densities with macro in young and aged naïve and adoptive transfer (AT) whole cord (wc) cross-sections. Aspa+ OL density is significantly lower in aged naïve compared to young naïve and in adoptive transfer compared to naïve in both ages. Median shift Bootstrap test. (**D**). Manual quantification of differentiated OLs (GFP-Aspa+CD45-) in young and aged adoptive transfer non-lesion white matter (nlwm) and individual lesions. OL density is significantly lower in lesions compared to nlwm in both ages. Aged AT OL density is significantly lower than young AT in both regions. Median shift Bootstrap test. (**E**). Quantification of the ratio of mean OL density in lesions compared to nlwm across a whole spinal cord section. OL lesion/nlwm ratios are not significantly different between ages. Median shift Bootstrap test. Box and whiskers plots: box 25 to 75 percentile, whiskers min to max, median line, show all datapoints. p values: ns p>0.05, * p≤0.05, **p≤0.01, ****p<0.0001. Scale bars: A,B 200μm.

Figure 4- Axonal loss, myelin debris and demyelination in young and aged adoptive transfer
(**A**). Representative images of macro quantification of axonal density in *Pdgfra-Cre^{ERT};RCE* adoptive transfer spinal cord section stained for neurofilament-H (NF) and DAPI (left panel), NF only grayscale image (middle left panel), NF macro thresholded image (middle right panel) and NF macro processed image with NF+ axons outlined in red (right panel). Lesions outlined. (**B**). Quantification of NF axonal density by macro in individual lesions of young and aged adoptive transfer (AT). No significant difference in axonal density in lesions between ages. Median shift Bootstrap test. (**C**). Representative image of myelin debris/degraded myelin (dMBP) and GFP staining in young and aged *Pdgfra-Cre^{ERT};RCE* adoptive transfer spinal cord sections. (**D**). Quantification of the percentage of white matter area with dMBP staining in young and aged AT spinal cord with significant increase white matter with myelin debris in aged white matter. Median shift Bootstrap test. (**E**). Representative images of FluoroMyelin and GFP staining in young and aged adoptive transfer *Pdgfra-Cre^{ERT};RCE* spinal cord sections. White matter regions with DAPI hypercellularity demonstrate absence of FluoroMyelin staining. (**F**). Quantification of the percentage of demyelinated white matter with no FluoroMyelin staining in young and aged AT spinal cord with no significant difference between ages. Median shift Bootstrap test. Box and whiskers plots: box 25 to 75 percentile, whiskers min to max, median line, show all datapoints. p values: ns p>0.05, **p≤0.01. Scale bars: A,C,E 200μm.

Figure 5- Remyelination in young and aged adoptive transfer lesions

(**A**). Representative image of lumbar ventrolateral spinal cord in lineage traced OPC Cre line crossed to membrane GFP reporter (*Pdgfra-Cre^{ERT};Tau^{mGFP}*) for labeling remyelinating myelin sheaths from OPCs. Staining for myelin (MBP), lineage traced OPC processes (GFP) and axons (NF) with no GFP+ myelin sheaths in naïve/non-adoptive transfer. (**B**). Representative image of young *Pdgfra-Cre^{ERT};Tau^{mGFP}* ventral spinal cord 10 days post-adoptive transfer with lineage traced OPC GFP+ processes wrapping NF+ axons (arrows) at the lesion edge, box indicates region of higher magnification. (**C**). Representative transmission electron microscopy (TEM) images of ventromedial white matter lesions in young and aged adoptive transfer. (**D**). Higher magnification TEM images of a young adoptive transfer lesion with thin myelin sheaths around degenerating axons with electron dense material or myelin infoldings (black asterisks) and thin myelin sheaths around axons without signs of degeneration (red asterisks). (**E**). Representative TEM inverted images of young and aged adoptive transfer lesions to highlight myelin sheaths. (**F**). Quantification of g-ratios of axons ≥2μm diameter in young and aged adoptive transfer lesions. Simple linear regression analysis. (**G**). Cumulative relative frequency distribution of g-ratios indicating a significantly higher proportion of lower g-ratios (thicker myelin sheaths) in aged lesions. Kolmogorov-Shirnov test. (**H**). Quantification of percent of thin myelin sheaths (g-ratio ≥0.8) without degenerative features of total myelinated axons with diameter ≥2μm within a lesion. No significant difference between young and aged lesions. Median shift Bootstrap test. (**I**). Quantification of percent of thin myelin sheaths (g-ratio ≥0.8) with degenerative features of total myelinated axons with diameter ≥2μm within a lesion. No significant difference between young and aged lesions. Median shift Bootstrap test. (**J**). Cumulative relative frequency distribution of axonal areas of thin myelin sheaths with degenerative features (black) or no degenerative features (gray) with a significantly higher proportion of large axons in axons with degenerative features. Kolmogorov-Shirnov test. F-G, n=4 young animals with 31 lesions, n=6 aged animals with 36 lesions. H-J, n=4 young animals with 25 lesions, n=6 aged animals with 18 lesions. Box and whiskers plots: box 25 to 75 percentile, whiskers min to max, median line, show all datapoints. p value: ns, p>0.05, **p≤0.01, ***p≤0.001, ****p<0.0001.