The Arabidopsis nucleoporins 98A/B are essential for NLP7 nuclear
accumulation in response to nitrate

Zsolt Kelemen[1,#], Garry Duville[1,2,#], Virginie Bréhaut[1], and Anne Krapp[1*]

[1] University Paris-Saclay, INRAE, AgroParisTech, Institute Jean-Pierre Bourgin for Plant Sciences
(IJPB), 78000, Versailles, France
[2] University Paris-Saclay, 91405 Orsay, France
7
# These authors have equally contributed.

*Anne Krapp, Institute Jean-Pierre Bourgin for Plant Sciences (IJPB), Route de Saint Cyr, 78000,
Versailles, France

Email: anne.krapp@inrae.fr

Keywords: Nucleoporin, transcription factor, nitrate signaling, Arabidopsis

This PDF file includes:

Main Text
Figures 1 to 6
Supplemental Table S1
Supplemental Figure S1
Graphical abstract
21

Abstract

The nuclear compartment is an essential evolutionary gain in eukaryotes. Stimulus-induced nuclear accumulation of transcription factors allows rapid and reversible regulation of gene expression. Nitrate-regulated nuclear accumulation of the master transcriptional regulator NLP7, a member of the RWP-RK family of transcription factors, triggers a major gene expression cascade. Whether and how this process is regulated at the level of the nuclear pore remains elusive. Here we show that the mobile nucleoporin NUP98A and its close homolog NUP98B interact with NLP7 in the nucleus and that the nitrate-triggered nuclear accumulation of NLP7 is impaired in their absence. We also demonstrated that the simultaneous loss of NLP7 and NUP98A has a strong and specific effect on the expression of nitrate-regulated genes, resulting in a drastic reduction in plant growth. Our results indicate that plant NUP98 acts on nitrate use by plants through its interaction with the top-tier NLP7 transcription factor.

Significance Statement

Nucleoporins (NUPs) are fundamental components of all eukaryotic cells. They are increasingly recognized not only as structural components of the nuclear pore complex but also as active regulators of gene expression and signaling pathways across eukaryotes. NUP98 proteins in mammals are multifunctional proteins with key roles beyond nucleocytoplasmic transport. The adaptation of plants to nitrogen availability is crucial for their development and yield. The master transcription factor NLP7 orchestrates early responses to nitrate supply and is regulated by a rapid nitrate-regulated nucleo-cytosolic shuttling. In this study, we demonstrate that the Arabidopsis nucleoporins 98A/B enable acclimation of Arabidopsis to N status by gating the nucleocytoplasmic trafficking of the nitrate sensor NLP7 transcription factor, enlarging their functional network.

Main Text

Introduction

Nucleoporins (NUPs) are fundamental components of all eukaryotic cells. They are constituents of the nuclear pore complexes (NPCs), which are extremely large protein complexes embedded in the nuclear envelope and consist of over 1,000 copies of ~30 individual NUPs. The central channel of the NPC functions as a permeability barrier between cyto- and nucleoplasm. The FG-repeat-containing NUPs within the central channel constitute a permeability barrier that allows the passive diffusion of small molecules, while limiting the passage of macromolecules larger than ~40 kDa

and enabling nucleocytoplasmic exchange. It thus plays the role of a regulator of intracellular molecule trafficking, constituting an important gatekeeper for macromolecular transports between the nucleus and cytoplasm (Li et al., 2016).

In addition to their main function in nucleocytoplasmic transport, NPCs and their components have been shown to play important roles in many biological processes, such as kinetochore and spindle assembly, cell cycle control, regulation of gene expression, chromatin organization, DNA repair, and DNA replication (Capelson and Hetzer 2009; Capelson et al., 2010; Strambio-De-Castillia et al., 2010; Wozniak et al., 2010; Van de Vosse et al., 2011; Bukata et al., 2013). In Arabidopsis, NUPs are also involved in multiple biological processes, including flowering and immune responses (Gu et al., 2016; Cheng et al., 2020).

Changing environmental conditions necessitate rapid adaptation of cytoplasmic and nuclear protein content, and a number of key transcription factors are regulated by a rapid nuclear accumulation in response to a signal (Cartwright and Helin, 2000). These TFs must be transported from the cytoplasm to the nucleus to reach their targets and induce or repress their transcription, and exported from the nucleus to the cytoplasm.

Nitrogen (N) is a crucial macronutrient, and N availability in the soil can be a limiting factor for plant growth and development (Marschner et al., 1995). Thus, plants evolved mechanisms to sense and promptly adapt to fluctuating nitrate availability. Importantly, nitrate is not only a nutrient but also a signaling molecule (Crawford, 1995; Scheible et al., 1997). The rapid transcriptional response to nitrate supply triggers expression changes of thousands of genes.

Members of the Nin Like-Protein (NLP) family are top tier TFs of the nitrate signaling pathway through inducing a cascade of transcriptional regulation (reviewed in Mu and Luo, 2019). The Arabidopsis NLP7 TF is regulated by a very rapid nuclear accumulation within seconds after exposure to nitrate. This mechanism involves the phosphorylation of an evolutionarily conserved Serine by three CALCIUM-DEPENDENT PROTEIN KINASEs (CPK10, CPK30, CPK32, Liu et al., 2017). In addition, NLP7 activity is directly regulated by nitrate binding to its N-terminal region (Liu et al., 2022). The Arabidopsis genome encodes nine NLPs that bind in yeast the nitrate-responsive cis-element (NRE), a conserved DNA sequence motif found in the proximal regions of promoters of nitrate-responsive genes and shown to be necessary and sufficient for their nitrate-induced gene expression (Konishi and Yanagisawa, 2011, 2013). NLP7 was shown as a major TF governing the early nitrate response (Marchive et al., 2013; Konishi and Yanagisawa, 2019; Alvarez et al., 2020). NLP6, the closest homolog of NLP7, exhibits a partially redundant function to that of NLP7. However, although the loss of NLP7 is sufficient to cause a growth reduction under ample nitrate supply and impaired induction of nitrate-responsive gene, the additional loss of function of NLP6 leads to a stronger phenotype than that of the single mutant *nlp7-1* (Guan et al., 2017, Cheng et

al., 2023). Recent studies demonstrated that NLP2 orchestrates both N and carbon assimilation in
response to external nitrate availability (Konishi et al., 2021; Durand et al., 2023). Other NLPs also
contribute to the regulation of nitrate-inducible gene expression and nitrate-dependent promotion
of vegetative growth in Arabidopsis (Konishi et al., 2021; Liu et al., 2022). The functionality of NLPs
is further enriched by their ability to form homo-, heterodimer, or to interact with other proteins via
their PB1 domain (Guan et al., 2017; Konishi et al., 2021).

By exploring NLP7 interacting proteins, we discovered that nucleoporins NUP98A and NUP98B
interact directly with NLP7. Here, our goal was to elucidate how NUP98A/B bridges nitrate signaling
to the NPC function. We first confirmed the interaction of NUP98A/B with NLP7 in planta. Then,
applying genetics, we showed that the double mutant *nup98anup98b* has a higher sensitivity to
nitrate limitation and that the loss of function of NUP98A in the *nlp7* mutant background leads to a
strongly impaired early transcriptional nitrate response and impaired growth on nitrate fertilization.
Finally, the significant reduction of GFP-NLP7 nuclear accumulation in response to nitrate in the
genetic background of *nup98anup98b* strongly underscores the critical role of NUP98A/B proteins
in regulating nitrate-dependent nuclear accumulation of NLP7.

Results

NLP7 protein interacts with the nucleoporin NUP98A/B

Using a Y2H approach (performed by Hybrigenics), we found an interaction between the full-length
NLP7 protein and NUP98A. We then confirmed this interaction by pairwise Y2H assay (Figure 1A)
and in planta by co-expressing ^{C}YFP-NLP7 and ^{N}YFP-NUP98A in *Nicotiana benthamiana*. The
YFP signal indicating an interaction between the co-expressed proteins was present in the nucleus
as well as in the cytoplasm (Figure 1B-G).

We then tested whether the closely related nucleoporin NUP98B also interacts with NLP7 after
transient expression in *N. benthamiana*, using two other nucleoporins, NUP96 and CPR5, as
negative controls (Figure 2). NLP7 showed interaction with NUP98B with a localization of the
interaction similar to NUP98A (Figure 2D-F), whereas the nucleoporins NUP96 and CPR5 showed
no interaction with NLP7 (Figure 2G-L). Taken together, these results demonstrate that the
interaction between NLP7 and NUP98A/B is specific to these two NUPs and is not a general
interaction with proteins of the NPCs.

PB1 and GAF domains of NLP7 are not essential for NLP7-NUP98A interaction

NLPs contain 2 predicted protein-protein interaction domains. The C-terminal PB1 domain, which
has been shown to be involved in the formation of homo- and heterodimers of NLP proteins (Konishi

and Yanagisawa, 2019) and with proteins that do not contain PB1 domains (Guan et al., 2017).
The second, the N-terminal GAF-like domain, is reported as a protein-protein interacting domain
(Ho et al., 2000; Wu et al., 2022). To identify the NLP7 protein domains involved in the interaction
with NUP98A, we produced NLP7 constructs without either the GAF-like domain or the PB1
domain. NLP7 without either GAF or PB1 domain interacted with NUP98A in the BIFC assays,
producing positive signal both in the nucleus and in the cytoplasm (Supplemental Figure S1). Thus,
the NLP7 PB1 and GAF domains might contribute to the interaction between NLP7 and NUP98A/B
but are not essential for it.

The intracellular localization of the NUP98A-NLP7 interaction is modified by the N status of
the plants

NLP7 accumulates in the nucleus in the presence of nitrate and is localized in the cytosol in N-
starved seedlings (Marchive et al., 2013). Because NUP98A/B are key components of the NPC,
we then ask whether their association with NLP7 might depend on the N status. We studied the
interaction between NLP7 and NUP98A in *N. benthamiana* plants that have been grown under
ample nitrate supply (10 mM) and in plants that were treated with a 5-day N depletion. In addition,
leaf discs from N-depleted plants were resupplied with nitrate 10 min before observation (Figure
3). The fluorescent signal indicating the interaction between NLP7 and NUP98A was present both
in the cytoplasm and in the nucleus in all three conditions, however, in the case of the starved
leaves, the nuclear signal was less defined (Figure 3 D-F). Quantification of the fluorescence ratio
between nucleus and cytoplasm confirmed these observations (Figure 3 J). Our results suggest
that NUP98A/B interact with NLP7s when both proteins are localized in the same cellular
compartment with enhanced interaction in the nucleus compared to the cytosol in the presence of
nitrate.

Limiting nitrate supply amplifies the early senescence of nup98anup98b double mutants.
The protein interaction between NLP7 and NUP98A/B led us to evaluate the role of NUP98A/B in
the response of plants to nitrate availability. We analyzed the growth of NUP98A/B loss-of-function
mutants (*nup98a-1*, *nup98a-2*, *nup98b-1*, *nup98b-2* and *nup98a-1nup98b-1*) under contrasting
nitrate supply that had revealed nitrate-dependent growth phenotypes of *nlp7* and *nlp2* mutants
(Durand et al., 2023). We grew the plants in sand culture under short-day conditions (8 h light/16 h
dark) with either 0.5 mM nitrate or 5 mM nitrate supply for 39 days and determined rosette and root
biomass (Figure 4A - D). Under both N supply conditions, the rosette and root biomass of *nup98a*
and *nup98b* single mutants were similar to that of the wild type (WT) Col-0. Under non-limiting N
supply, WT plants displayed the expected (2.5-fold) increase in rosette fresh weight compared to
WT plants grown under 0.5 mM N supply (Figure 4A and B). Rosette biomass of the double mutant
nup98a-1nup98b-1 was strongly decreased in both N regimes (by 72 and 84 % under 0.5 and 5mM

nitrate supply, respectively). Root fresh weight of *nup98a* and *nup98b* mutants was also similar to

WT, with a trend towards slightly lower root biomass compared to WT under limiting N supply

(Figure 4 C and D). Root biomass was also decreased in the case of the *nup98a-1nup98b-1* double

mutant, and the rosette/root ratio was modified for the double mutant compared to WT (Figure 4 E

and F). Interestingly, it decreased slightly by 30% under non-limiting N supply, but increased 2-fold

under limiting N supply. The shoot/root ratio is an indicator of N limitation, suggesting that the

double mutant experienced less N limitation than the WT and the single mutants under limiting N

supply. Remarkably, the early senescence phenotype of *nup98a-1nup98b-1* double mutant that

was described before for this double mutant (Xiao et al., 2020) and that was visible by the loss of

chlorophyll, particularly in older leaves, had appeared much earlier in plants grown under nitrate-

limiting conditions than under ample nitrate supply (at 28, 35 and 38 days, respectively, Figure 4,

G and H). Under these conditions no loss of chlorophyll was observed for WT under both nitrate

regimes. Taken together, these observations suggest that NUP98A/B are necessary to prevent

early N limitation-induced senescence.

Double and triple mutant phenotypes support an interplay between NLP7 and NUP98A/B. To

study the genetic interaction of NUP98A/B and NLP7 and to further unravel the roles played by

NUP98A/B in the NLP7-dependent nitrate responses, we generated the *nup98a-1nlp7-1* and

nup98b-1nlp7-1 double mutants and the triple mutant *nup98a-1nup98b-1nlp7-1*.

We first studied the growth of single and higher order mutants in comparison to the WT in sand

culture under low and ample nitrate supply in short day conditions (Figure 5 A-C). Under limiting N

supply, the single mutants *nup98a*, *nup98b* and *nlp7-1* attained a rosette biomass comparable to

that of the WT, as expected. The rosette FW of the double mutants *nlp7-1nup98a* and *nlp7-

1nup98b* decreased by 40% and 30% respectively, compared to *nlp7-1*. The double mutant

nup98anup98b had a rosette biomass of 37 mg, corresponding to 22% of WT and the triple mutant

nup98a-1nup98b-1nlp7-1 barely attained an FW of 1 mg (Figure 5A). Under non-limiting nitrate

supply, whereas the *nup98a* and *nup98b* showed no difference to WT, the biomass of the *nlp7-1*

mutant was decreased 2-fold compared to WT, as reported previously and explained by a

constitutive N-limitation phenotype under ample nitrate supply (Castaings et al., 2009). In these

conditions, rosette biomass of the double mutants *nlp7-1nup98a* and *nlp7-1nup98b* was decreased

by 50% and 35%, respectively, compared to *nlp7-1*, thus slightly stronger than under limiting nitrate

supply. The triple mutant reached a biomass of 5.5 mg compared to 75 mg for the double mutant

nup98anup98b. Together, our data suggest that both NLP7 and NUP98A/B determine plant growth

in response to N supply. This can be due either to their participation in a shared regulatory pathway
or in two different pathways affecting growth in response to the N status of the plant.

Loss of function of both NLP7 and NUP98 amplified the altered early response to nitrate in
nlp7-1 mutants. Given the conserved role of nucleoporins and transcription factors in many
eukaryotes, the interaction between NLP7 and NUP98A/B may reflect a broader principle whereby
nuclear transport components cooperate with transcriptional regulators to fine-tune N responses.
We compared steady state levels of mRNA in seedlings either grown for 11 days on 3 mM nitrate
medium, or deprived of nitrogen for 3 days after 11 days in medium containing 3 mM nitrate, or
resupplied for 30 min by 3 mM nitrate after the 3-day deprivation period. We measured by RT-
qPCR the relative transcript levels of 18 NLP7-dependent nitrate-regulated genes in *nlp7-1nup98a*
double mutants compared to the single mutants and the WT (Figure 5 D-G). The main differences
between *nlp7-1* and the double mutant *nlp7-1nup98a* have been observed after nitrate resupply
(Figure 5 G). For all of the studied genes, the nitrate-triggered increase in steady-state transcript
level was diminished for *nlp7-1* as expected. For the double mutant *nlp7-1nup98a*, this reduced
increase was further diminished in the case of *NITRATE TRANSPORTER 2.1* (*NRT2.1*), *NRT2.2*,
NLP1, *NITRITE REDUCTASE* (*NII*), *NITRATE REDUCTASE1* (*NIA1*), *NAR2*, and *HRS1*
HOMOLOGOUS 1 (*HHO1*). A similar tendency was observed for *LOB BINDING DOMAIN 38*
(*LBD38*), *NLP2* and *BTB/POZ AND TAZ DOMAIN-CONTAINING PROTEIN 5* (*BT5*), whereas no
difference in the expression level after 30 min of nitrate resupply for the double mutant *nlp7-*
1nup98a compared to *nlp7-1* was found for *MAP KINASE KINASE KINASE 14* (*MAP3K14*),
MAP3K13, *NLP3*, *BT2*, *HYPERSENSITIVITY TO LOW PI-ELICITED PRIMARY ROOT*
SHORTENING 1 (*HRS1*), *LBD39*, *NIA2* and *NRT1.1*. Taken together, for several nitrate-regulated
genes, the combination of *nlp7* and *nup98* mutations resulted in an additive decrease in nitrate
induction, likely due to the consequences of impaired nuclear accumulation of other NLPs or other
transcription factors necessary for complete early response to nitrate.

NUP98A and NUP98B are required for the nitrate-triggered nuclear accumulation of NLP7.
To decipher the mechanism underlying the various developmental and molecular phenotypes of
the higher-order mutants, we studied the nuclear accumulation of GFP-NLP7 in response to nitrate
resupply to N-depleted seedlings in different mutant backgrounds. We hypothesized that the
interaction of NUP98A and NUP98B with NLP7 is required for the nitrate-triggered nuclear
accumulation and NLP7-dependent nitrate-regulated gene expression and growth.

We introduced the *GFP-NLP7* translation fusion driven by the *NLP7* promoter into these NUP98
loss of function mutants by crossing an *NLP7pro:GFP-NLP7* (*nlp7-1*) transgenic line (Castaings et
al., 2009) with the single mutant *nup98a-1* and the double mutant *nup98a-1nup98b-1*. In N-
depleted 10-day-old seedlings, GFP-NLP7 was localized in the cytosol of all genotypes (Figure 6

A, C, E). At 10 minutes after nitrate resupply, in the presence of NUP98A and NUP98B GFP-NLP7
accumulated rapidly in the nucleus (Figure 6 B). The loss of function of NUP98A was accompanied
by a decrease in the accumulation in the nucleus after nitrate resupply (Figure 6 D). In the absence
of both NUP98A and NUP98B, the accumulation of GFP-NLP7 in the nucleus in response to nitrate
resupply was almost abolished (Figure 6 F). Quantification of the fluorescence ratio between
nucleus and cytoplasm confirmed these observations (Figure 6 G). Thus, our results indicate a
specific role of NUP98A/B for the nitrate-triggered nuclear accumulation of NLP7.

Discussion

Nucleoporins are increasingly recognized not only as structural components of the nuclear pore
complex (NPC) but also as active regulators of gene expression and signaling pathways across
eukaryotes. NUP98 proteins in mammals are multifunctional components of the nuclear pore
complex with key roles beyond nucleocytoplasmic transport. Increasing evidence shows that
NUP98 and other nucleoporins are frequently associated with active chromatin regions, particularly
enhancer elements, where they contribute to the regulation of gene expression and chromatin
organization. NUP98 can act as a scaffold at these regulatory sites, promoting the maintenance of
transcriptionally active chromatin by stabilizing histone modifications and facilitating epigenetic
memory. Several models have been proposed to explain this chromatin-binding function of NPCs:
(i) NPCs may help maintain the active state of genes through direct interaction with histone-
modifying enzymes; (ii) the targeting of active genes and enhancers to the NPC could be
functionally linked to transport-related mechanisms, such as the nuclear shuttling of transcription
factors; and (iii) the intrinsically disordered FG-repeat domains of NUP98 may enable phase
separation at NPCs, creating specialized transcriptional compartments that enhance gene
activation at enhancers and super-enhancers (for review, see Pascual-Garcia & Capelson (2019).
In this study, we uncover a novel role for the plant nucleoporins NUP98A and NUP98B in controlling
the nuclear accumulation of the transcription factor NLP7, a central regulator of nitrogen signaling.
While NUP98 proteins are generally thought to maintain NPC integrity and mediate nucleo-cytosolic
trafficking (Tamura et al., 2010; Raveh et al., 2016), studies in yeast, animals, and plants have
highlighted their mobility and functional versatility beyond nuclear transport (Capitanio et al., 2017;
Gallemí et al., 2016). Our work now extends this paradigm to nitrogen signaling in plants.

Here, we report a novel role for the nucleoporins NUP98A and NUP98B in regulating the nuclear
accumulation of the TF NLP7. Based on their sequence homology with other species, plant
NUP98A/B are considered to be required for maintaining the NPC structure and facilitating
macromolecule nucleo-cytosolic trafficking (Tamura et al., 2010; Raveh et al., 2016). Although they
have an important structural role in the NPC, NUP98s in yeast, animals and plants are mobile

proteins that can be localized in different cellular compartments (Capitanio et al., 2017; Gallemí et
al., 2016). This has various implications for their role in other biological processes. We validated
the interaction between NUP98A and NLP7 using in planta BiFC assays, confirming the results
obtained through yeast two-hybrid screening. Interaction was detected both in the cytosol and the
nucleus. We cannot exclude that the strong transient expression of the splitYFP proteins driven by
the *CaMV 35S* promoter leads to a protein localization different from the native conditions.
However, the localization of the interacting complex corresponded to the localization of NLP7
tagged with GFP under the particular condition, as in N-depletion condition, the interaction in the
nucleus was less pronounced, as expected from the NLP7 localization change under this condition
(Marchive et al., 2013).

Here, we showed that neither the NLP7 GAF domain nor its PB1 domain is essential for the
interaction with NUP98A/B, however, it is possible that both domains act redundantly to ensure this
interaction. Found in a variety of proteins, including TFs, the GAF domain is important for protein-
protein interactions and can bind a variety of ligands, including cGMP, cAMP, and small molecules
such as oxygen and nitrate (Ho et al., 2000; Moglich et al., 2009; Niemann et al., 2014; Su and
Lagarias, 2007; Grefen et al., 2008). It has been shown that the NLP7 PB1 domain is important for
its interaction with other NLPs, but also with the transcription factor teosinte
branched1/cycloidea/proliferating cell factor20 (TCP20), and SUCROSE NON-FERMENTING 1-
RELATED KINASE 1 (SnRK1) (Konishi and Yanagisawa, 2019; Guan et al., 2017; Wang et al.,
2022). The NLP7 protein possesses short intrinsic disordered regions (IDRs), which leads to the
hypothesis that these IDRs interact with the FG motifs of NUP98. However, the nature of the
underlying molecular interactions between FG repeats that might lead to protein condensates (Ahn
et al., 2025) is largely unknown. Interestingly, it has been shown recently that protein condensation
is involved in the specificity of transport through the NPC (Wang et al., 2023).

Despite the high probability that the positive BiFC between NLP7 and NUP98A/B is due to a direct
interaction between these proteins, we cannot exclude that a third partner is necessary for the
interaction. For example, NirA, the specific transcription factor of the nitrate assimilation pathway
of Aspergillus nidulans, interacts with the nuclear export factor KapK, which bridges an interaction
with a nucleoporin-like protein. Interestingly, NirA, which has no structural homology with NLP7,
also accumulates in the nucleus upon induction by nitrate via a nuclear retention mechanism
(Bernreiter et al., 2007).

To further assess the biochemical mechanisms, further cell biological experiments are needed in
Arabidopsis thaliana transgenic plants using native promoters to examine whether NUP98A/B and
NLP7 co-localize and translocate together into the nucleus, particularly under nitrate treatment
conditions. In addition, biochemical validation of the NLP7–NUP98 interaction under native

expression levels and mapping of the interacting domains of NUP98A/B will be important to
determine whether this regulation relies on direct binding to FG-repeat regions or involves
additional transport factors. However, our genetic data demonstrate clearly that NUP98A/B are
essentiel for nitrate-triggered nuclear accumulation of NLP7. Future work will be necessary to
reveal the precise molecular mechanism and distinguish whether NUP98A/B primarily facilitate
nuclear import of NLP7, promote its nuclear retention, or modulate its export dynamics.

Several other *nup* mutants are, like *nup98anup98b*, also modified for the intracellular partitioning
of regulatory proteins, including TFs. For example, the mutation of the *HIGH EXPRESSION OF*
OSMOTICALLY RESPONSIVE GENES 1 (*HOS1*) reduces the nuclear accumulation of SnRK1a1
(Margalha et al., 2023) and of PHYTOCHROME-INTERACTING FACTOR 4 (PIF4) (Zhang et al.,
2020). The NUP62-subcomplex modulates Flowering locus C (FLC) nuclear import, and NUP85 is
a player in the nuclear-cytosolic trafficking of ACQOS, encoding a nucleotide-binding leucine-rich
repeat (NLR) protein (Mori et al., 2024). The nuclear-cytosolic partitioning of these proteins is
regulated by abiotic stresses or the transition to flowering. Here, we show that the response to
nutrient availability is also governed by the function of a NUP, NUP98A/B.

nup98anup98b double mutants display early flowering (Jiang et al., 2019) with early senescence
during flowering. NUP98A/B are also involved in the control of starch degradation (Xiao et al., 2020)
and in the circadian control (Jiang et al. 2019, 2020). Here, we found that single *nup98a* mutants
had only minor phenotypes, whereas the *nup98anup98b* double mutants showed a severe growth
phenotype associated with early senescence as reported before (Xiao et al., 2020). Interestingly,
the early senescence is exacerbated by N limitation. *nlp7* mutants display several traits of N
limitation, despite growing under non-limiting nitrate supply (Castaings et al., 2009). Whereas this
does not lead to early senescence of the *nlp7* mutants, impaired nuclear accumulation of other
NLPs in the *nup98anup98b* mutant might lead to its early senescent phenotype under N limitation.
Indeed, our study shows that in *nup98a-1nlp7-1* double mutants, a subset of nitrate-regulated
genes displayed reduced transcript abundance in response to nitrate resupply compared to the
nlp7 mutant. This might be due to the impact of the loss of function of NUP98A on the activities of
other TFs involved in nitrate signaling, including NLP1/2/6/7 (Konishi et al., 2021, Cheng et al.,
2023).

Besides the pivotal role in gating nuclear and cytoplasmic functions, the nuclear pore proteins
function in key biological processes, including chromatin tethering to the nuclear periphery, a
mechanism contributing to the regulation of gene expression (Ungricht and Kutay, 2017; Smith et
al., 2015). Thus, it will require further investigations to determine whether NUP98A/B act with NLP
to orchestrate the nitrate-triggered transcriptional regulation of nitrate-regulated genes by

facilitating their transcriptional activation by NLPs in addition to regulating nuclear accumulation of
NLPs in response to nitrate.

Interestingly, in humans, NUP98 has oncogenic roles as a genetic fusion partner (Gough et al.,
2011). Indeed, in malignant cells, NUP98 occurs as part of fusion proteins with TFs, pointing to the
involvement of NUP98 in transcriptomic dysregulation during oncogenesis. In plants, such NUP98
fusion proteins have not been reported, and it can be speculated that instead, protein-protein
interactions evolved. Further studies are required to reveal further regulatory mechanisms
governed by the NUP98-NLP7 protein complex.

In conclusion, this study demonstrates that in Arabidopsis, NUP98A/B promotes NLP7 activity
through the control of its nucleocytoplasmic trafficking and thereby impacts the rapid transcriptional
response to nitrate availability and the acclimation of plant growth to nitrate supply.

Materials and Methods

Plant materials. All Arabidopsis material is in the Col-0 background. The *Arabidopsis thaliana*
mutant *nlp7-1* (SALK_26134) derived from T-DNA–mutagenized populations of the Col-0
accession (Alonso-Blanco et al., 2003, Castaings et al., 2009) and was backcrossed twice to Col-
0. All mutant seeds were obtained from the NASC, and homozygous lines were identified by
genotyping using the primers listed in Supplemental Table S1. Homozygote lines of *nup98a1*
(SALK_103803, Jiang et al., 2019), *nup98a2* (SALK_015016, Jiang et al., 2019, also named *dra2-*
4, Gallemi et al., 2016), *nup98b1* (GABI_288A08, Jiang et al, 2019), and *nup98b2* (SALK_132527)
have been selected by genotyping. The higher-order mutants with and without the *NLP7pro:GFP-*
NLP7 construct have been obtained by crosses of *NLP7pro:GFP-NLP7* (Col-0) line A4.4 (Castaings
et al., 2009) followed by selfing of the F1 generation, genotyping the F2 generation and selecting
for homozygous transgene by hygromycin selection in F3.

Bimolecular Fluorescence Complementation (BiFC). Open reading frames (without stop codon)
flanked by AttB1 and AttB2 sites were amplified from Arabidopsis cDNA clones (Columbia
accession) using the primers as given in Supplemental Table S1, cloned into Gateway vector
pDONR207 using BP recombination (Invitrogen), and sequenced, except *NUP98A*, where the
cDNA sequence optimized for *Arabidopsis thaliana* was synthetized and cloned into pTwist ENTR
Kozak entry vector by Twist Biosciences (Supplemental Data Set 1). An LR reaction between the
entry vector and the complete set of four pBiFP vectors (Azimzadeh et al., 2008) produced the final
expression vectors, where coding sequences are cloned in fusion with the N- or C-terminal parts
of YFP, either as N-terminal or C-terminal fusions, under the control of the cauliflower mosaic virus
35S promoter.

Modified versions of NLP7 were produced by PCR. For NLP7 without PB1 domain, the region
corresponding to 1-863 amino acids was amplified by PCR and cloned into the Gateway vector
pDONR207 using BP recombination. For NLP7 without the predicted GAF domain (Chardin et al.,
2014), regions corresponding to amino acids 1-159 and 301-959 were amplified separately. The
two fragments were fused together by a second PCR, and the resulting fragment flanked by AttB1
and AttB2 sites was cloned into pDONR207. The primer sequences are given in Table S1. A
description of the pH2B2:H2B2-mTurquoise construct can be found in Bouchez et al. (2024).

For BiFC interaction assays, the vectors were transformed into *Agrobacterium tumefaciens* strain
C58C1(pMP90) (Kámán-Tóth et al., 2018) and co-infiltrated into *Nicotiana benthamiana* leaves.
Plants were grown either in fertilized soil in an air-conditioned greenhouse or on sand in a growth
chamber under long day conditions (day 22 oC / night 18 °C, 16h photoperiod, 130-160 µmolm-2s-
1 light intensity (Systion 32-42 VDC 330mA, 11V). Plants grown in sand culture were watered with
a 2 mM nitrate-containing nutrient solution for 29 days, followed by watering for 3 days with a
nutrient solution without N after washing the sand with the N-depleted solution or by watering with
a 10 mM nitrate nutrient solution. Agrobacterium carrying clones of interest were grown overnight
at 28°C in 5 ml LB medium with appropriate antibiotics. Aliquots from the overnight cultures were
resuspended in 10 mM $MgCl_2$ and 10 mM 2-(N-morpholine)-ethanesulphonic acid (MES), pH 5.6,
to obtain a final OD600nm of 0.5 for tobacco leaf infiltration. YFP fluorescence was detected 2-3
days after infiltration. During that time, plants were watered with either 0 mM or 10 mM nitrate-
containing nutrient solution. Ion equilibrium of the nutrient solution was ensured by replacing KNO3
and $Ca(NO_3)_2$ with KCl and $CaSO_4$, respectively.

Yeast Two-Hybrid Assays. Coding sequences of NLP7 and NUP98A cloned in pDONR207 were
recombined into the pGBKT7 and pGADT7 vectors (Clontech) via LR Clonase reactions and co-
transformed into the Saccharomyces cerevisiae strain MaV203 (Invitrogen). Yeast was cultured on
YPDA plates (Clontech) at 28 °C. For transformation, an overnight liquid culture at OD600nm =0.6
was harvested, resuspended in 1 mL sterile water, pelleted by centrifugation (maximum speed,
15 s), and resuspended in 1 mL of 0.1 M lithium acetate in 1X TE buffer. Transformation mixes
contained 5 µg of each plasmid, 5 µL of carrier DNA (Finnzymes, Fisher Scientific), 0.1 mL of yeast
suspension, and 0.6 mL of 40% [v/v] PEG 4000 in 0.1 M lithium acetate/1X TE. After a 30 min
incubation at 28 °C, mixtures were heat-shocked at 42 °C for 25 min. Cells were pelleted (15 s, max
speed), resuspended in 0.25 mL sterile water, and plated on SD/-Trp-Leu selective medium
(Clontech). To assess protein–protein interactions, cotransformed colonies were resuspended in
0.1 mL sterile water, and 10 µL was spotted on SD/-Trp-Leu-His plates supplemented with various
concentrations of 3-amino-1,2,4-triazole (3-AT). Growth under these conditions was interpreted as
a positive interaction.

Growth conditions of Arabidopsis plants. Seeds were stratified for 5 days at 4°C in the dark and
then grown either on non-fertilized soil (Stender) or on sand, both supplemented with nutrient
solutions (Loudet et al., 2003) containing either 0.5 mM (growth-limiting nitrate supply) or 5 mM
(non-limiting nitrate supply) nitrate, in short-day photoperiod with an 8-h light/16-h dark cycle at
21°C/18°C, respectively, 65% relative humidity, and a constant 150 µmol photons $m^{-2} s^{-1}$ irradiation
with Osram 36w light sources. For N limitation conditions, ion equilibrium of the medium was
ensured by partly replacing KNO_3 and $Ca(NO_3)_2$ with KCl and $CaSO_4$, respectively.

For nitrate response assays, both for the determination of steady state mRNA levels and confocal
analyses, seeds were stratified for 3 days at 4°C and then grown in liquid medium containing 0.07%
(w/v) MES pH 5.8, 1 mM KH_2PO_4, 0.5 mM $MgCl_2$, 0.5 mM $CaSO_4$, microelements as in Estelle and
Somerville (1987), 0.25% (w/v) sucrose and 3 mM KNO_3 for 10 days in long-day photoperiod (16-
h light/8-h dark) at 25°C with a light intensity of 80 µmol photons $\cdot m^{-2} \cdot s^{-1}$ by Philips F17T8/TL741
light sources. For gene expression analysis after 7 days of growth, the medium was changed to
fresh medium every 2 days. Eleven-day-old seedlings were then nitrogen-starved for 3 days using
fresh medium containing 3 mM KCl instead of 3 mM KNO_3. After 3 days of N starvation, 3 mM
nitrate was added for the indicated time periods, and harvests were performed at several time
points. For confocal analysis, seedlings were grown as in Marchive et al. (2013) for five days on a
3 mM KNO_3 liquid medium, then N-starved for two days by exchanging the culture medium with an
N-free medium in which KNO_3 was replaced by KCl.

Confocal Imaging. Laser scanning confocal imaging was performed using the SP8 confocal laser
microscope (Leica, Germany) equipped with an argon laser (488 nm for GFP excitation, intensity
20%) and an argon laser (514 nm for YFP excitation, intensity 20%). Emission was collected at 505
to 525 nm (GFP, gain= 500) or at 520 to 540 nm (YFP, gain= 500). The images were coded green
for both GFP and YFP. Images were processed with ImageJ. Plant samples were mounted in 40
µL of a top agar solution (0.1%, w/v). During confocal imaging, 5 µL of 0.5 M KNO_3 was added by
capillarity to the mounting medium (45 µl).

Gene expression analysis. Measurements of relative steady-state mRNA levels were performed
on seedlings (plants at 1.02 developmental growth stages (Boyes et al., 2001)) that had been
cultivated in liquid medium as described above. Each sample was shock frozen after harvest and
was composed of 26-34 seedlings. Total RNA was extracted using the RNeasy plant mini kit
(QIAGEN), including a DNase treatment according to the manufacturer's instructions, or by Trizol
method (Invitrogen). First-strand cDNAs were synthesized according to Daniel-Vedele and
Caboche (1993) using Moloney murine leukemia virus reverse transcriptase and oligo(dT)15
primers (Promega) or using the SuperScript III Fast strand synthesis system (Invitrogen).

cDNAs were first checked by PCR for contamination by genomic DNA, then relative expression
was determined in 384-well plates with a QuantStudio 5 System device (40 3-step cycles; step 1,
95°C, 5 s; step 2, 58°C, 30 s; step 3, 72°C, 30 s) using Takyon qPCR kit for SYBR assay
(Eurogentec) (2.5 μL TAKYON SYBR 2X, 0.015 μL Forward primer 100 μM, 0.015 μL Reverse
primer 100 μM, 2.5 μL cDNA 1/20 per well). Relative expression was calculated according to the
2-ΔCt method. Target gene expression was normalized to the geometric mean of the expression
of two plant genes, *ACTIN2* (At3g18780) and *TIP41* (At4g34270). All primer sequences are given
in Supplemental Table S1.

Statistical analyses. Data were analyzed using non-parametric tests due to non-normal
distributions and/or unequal variances. Comparisons between multiple groups were performed
using the Kruskal–Wallis test. When a significant difference was detected ($p < 0.05$), Dunn's post
hoc test was used for pairwise comparisons without correction for multiple testing. Statistical
analyses were conducted in Python using the scipy and scikit-posthocs libraries.

Acknowledgments

We thank Hervé Ferry, Jean-Christophe Guedet and Philippe Maréchal for plant care, Gladys
Cloarec for support for confocal analyses, and Alizée Teinturier for technical assistance. We also
thank the NASC and ABRC for supplying Arabidopsis T-DNA insertion lines. This work has
benefited from the support of IJPB's Plant Observatory technological platforms PO-Plants and PO-
Cyto and was supported by ANR NITRASENSE (ANR-16-CE20-022, ZK, AK). The IJPB benefits
from the support of the Saclay Plant Sciences (SPS) Network (ANR-17-EUR-0007). We thank
Sylvie Ferrario-Méry, Thomas Girin, Anne-Sophie Leprince and Christian Meyer for fruitful
discussions.

Author Contributions

Z.K and A.K. designed research; Z.K, G.D., V.B., and A.K. performed research and analyzed
data; and Z.K., G.D., and A.K. wrote the paper.

Competing Interest Statement

The authors declare no competing interest.

References

Ahn, J.H., Guo, Y., Lyons, H., Mackintosh, S.G., Lau, B.K., Edmondson, R.D., Byrum, S.D., Storey, A.J., Tackett, A.J., Cai, L., Sabari, B.R., Wang, G.G., 2025. The phenylalanine-and-glycine repeats of NUP98 oncofusions form condensates that selectively partition transcriptional coactivators. Mol Cell 85, 708-725.e9. https://doi.org/10.1016/j.molcel.2024.12.026

Alonso-Blanco, C., Bentsink, L., Hanhart, C.J., Blankestijn-de Vries, H., Koornneef, M., 2003. Analysis of natural allelic variation at seed dormancy loci of Arabidopsis thaliana. Genetics 164, 711–729. https://doi.org/10.1093/genetics/164.2.711

Alvarez, J.M., Schinke, A.-L., Brooks, M.D., Pasquino, A., Leonelli, L., Varala, K., Safi, A., Krouk, G., Krapp, A., Coruzzi, G.M., 2020. Transient genome-wide interactions of the master transcription factor NLP7 initiate a rapid nitrogen-response cascade. Nature Communications 11, 1157. https://doi.org/10.1038/s41467-020-14979-6

Azimzadeh, J., Nacry, P., Christodoulidou, A., Drevensek, S., Camilleri, C., Amiour, N., Parcy, F., Pastuglia, M., Bouchez, D., 2008. Arabidopsis TONNEAU1 proteins are essential for preprophase band formation and interact with centrin. The Plant Cell 20, 2146–2159. https://doi.org/10.1105/tpc.107.056812

Bernreiter, A., Ramon, A., Fernández-Martínez, J., Berger, H., Araújo-Bazan, L., Espeso, E.A., Pachlinger, R., Gallmetzer, A., Anderl, I., Scazzocchio, C., Strauss, J., 2007. Nuclear export of the transcription factor NirA is a regulatory checkpoint for nitrate induction in Aspergillus nidulans. Molecular and Cellular Biology 27, 791–802. https://doi.org/10.1128/MCB.00761-06

Bouchez, D., Uyttewaal, M., Pastuglia, M., 2024. Spatiotemporal regulation of plant cell division. Curr Opin Plant Biol 79, 102530. https://doi.org/10.1016/j.pbi.2024.102530

Boyes, D.C., Zayed, A.M., Ascenzi, R., McCaskill, A.J., Hoffman, N.E., Davis, K.R., Görlach, J., 2001. Growth stage-based phenotypic analysis of Arabidopsis: a model for high throughput functional genomics in plants. The Plant Cell 13, 1499–1510. https://doi.org/10.1105/tpc.010011

Bukata, L., Parker, S.L., D'Angelo, M.A., 2013. Nuclear pore complexes in the maintenance of genome integrity. Current Opinion in Cell Biology 25, 378–386. https://doi.org/10.1016/j.ceb.2013.03.002

Capelson, M., Hetzer, M.W., 2009. The role of nuclear pores in gene regulation, development and disease. EMBO reports 10, 697–705. https://doi.org/10.1038/embor.2009.147

Capelson, M., Liang, Y., Schulte, R., Mair, W., Wagner, U., Hetzer, M.W., 2010. Chromatin-bound nuclear pore components regulate gene expression in higher eukaryotes. Cell 140, 372–383. https://doi.org/10.1016/j.cell.2009.12.054

Capitanio, J.S., Montpetit, B., Wozniak, R.W., 2017. Human Nup98 regulates the localization and activity of DExH/D-box helicase DHX9. Elife 6, e18825. https://doi.org/10.7554/eLife.18825

Cartwright, P., Helin, K., 2000. Nucleocytoplasmic shuttling of transcription factors. Cellular and molecular life sciences: CMLS 57, 1193–1206. https://doi.org/10.1007/pl00000759

Castaings, L., Camargo, A., Pocholle, D., Gaudon, V., Texier, Y., Boutet-Mercey, S., Taconnat, L., Renou, J.-P., Daniel-Vedele, F., Fernandez, E., Meyer, C., Krapp, A., 2009. The nodule inception-like protein 7 modulates nitrate sensing and metabolism in Arabidopsis. The Plant

Journal: For Cell and Molecular Biology 57, 426–435. https://doi.org/10.1111/j.1365-313X.2008.03695.x

Chardin, C., Girin, T., Roudier, F., Meyer, C., Krapp, A., 2014. The plant RWP-RK transcription factors: key regulators of nitrogen responses and of gametophyte development. Journal of Experimental Botany 65, 5577–5587. https://doi.org/10.1093/jxb/eru261

Cheng, Y.-H., Durand, M., Brehaut, V., Hsu, F.-C., Kelemen, Z., Texier, Y., Krapp, A., Tsay, Y.-F., 2023. Interplay between NIN-LIKE PROTEINs 6 and 7 in nitrate signaling. Plant Physiology 192, 3049–3068. https://doi.org/10.1093/plphys/kiad242

Cheng, Z., Zhang, X., Huang, P., Huang, G., Zhu, J., Chen, F., Miao, Y., Liu, L., Fu, Y.-F., Wang, X., 2020. Nup96 and HOS1 Are Mutually Stabilized and Gate CONSTANS Protein Level, Conferring Long-Day Photoperiodic Flowering Regulation in Arabidopsis. The Plant Cell 32, 374–391. https://doi.org/10.1105/tpc.19.00661

Crawford, N.M., 1995. Nitrate: nutrient and signal for plant growth. The Plant Cell 7, 859–868. https://doi.org/10.1105/tpc.7.7.859

Daniel-Vedele, F., Caboche, M., 1993. A tobacco cDNA clone encoding a GATA-1 zinc finger protein homologous to regulators of nitrogen metabolism in fungi. Mol Gen Genet 240, 365–373. https://doi.org/10.1007/BF00280388

Durand, M., Brehaut, V., Clement, G., Kelemen, Z., Macé, J., Feil, R., Duville, G., Launay-Avon, A., Roux, C.P.-L., Lunn, J.E., Roudier, F., Krapp, A., 2023. The Arabidopsis transcription factor NLP2 regulates early nitrate responses and integrates nitrate assimilation with energy and carbon skeleton supply. The Plant Cell 35, 1429–1454. https://doi.org/10.1093/plcell/koad025

Estelle, M.A., Somerville, C., 1987. Auxin-resistant mutants of Arabidopsis thaliana with an altered morphology. Mol Gen Genet 206, 200–206. https://doi.org/10.1007/BF00333575

Gallemí, M., Galstyan, A., Paulišić, S., Then, C., Ferrández-Ayela, A., Lorenzo-Orts, L., Roig-Villanova, I., Wang, X., Micol, J.L., Ponce, M.R., Devlin, P.F., Martínez-García, J.F., 2016. DRACULA2 is a dynamic nucleoporin with a role in regulating the shade avoidance syndrome in Arabidopsis. Development (Cambridge, England) 143, 1623–1631. https://doi.org/10.1242/dev.130211

Gough, S.M., Slape, C.I., Aplan, P.D., 2011. NUP98 gene fusions and hematopoietic malignancies: common themes and new biologic insights. Blood 118, 6247–6257. https://doi.org/10.1182/blood-2011-07-328880

Grefen, C., Städele, K., Růzicka, K., Obrdlik, P., Harter, K., Horák, J., 2008. Subcellular localization and in vivo interactions of the Arabidopsis thaliana ethylene receptor family members. Molecular Plant 1, 308–320. https://doi.org/10.1093/mp/ssm015

Gu, Y., Zebell, S.G., Liang, Z., Wang, S., Kang, B.-H., Dong, X., 2016. Nuclear Pore Permeabilization Is a Convergent Signaling Event in Effector-Triggered Immunity. Cell 166, 1526-1538.e11. https://doi.org/10.1016/j.cell.2016.07.042

Guan, P., Ripoll, J.-J., Wang, R., Vuong, L., Bailey-Steinitz, L.J., Ye, D., Crawford, N.M., 2017. Interacting TCP and NLP transcription factors control plant responses to nitrate availability.

Proceedings of the National Academy of Sciences of the United States of America 114, 2419–2424. https://doi.org/10.1073/pnas.1615676114

Ho, Y.S., Burden, L.M., Hurley, J.H., 2000. Structure of the GAF domain, a ubiquitous signaling motif and a new class of cyclic GMP receptor. The EMBO journal 19, 5288–5299. https://doi.org/10.1093/emboj/19.20.5288

Jiang, S., Xiao, L., Huang, P., Cheng, Z., Chen, F., Miao, Y., Fu, Y.-F., Chen, Q., Zhang, X.-M., 2019. Nucleoporin Nup98 participates in flowering regulation in a CONSTANS-independent mode. Plant Cell Reports 38, 1263–1271. https://doi.org/10.1007/s00299-019-02442-w

Kámán-Tóth, E., Pogány, M., Dankó, T., Szatmári, Á., Bozsó, Z., 2018. A simplified and efficient Agrobacterium tumefaciens electroporation method. 3 Biotech 8, 148. https://doi.org/10.1007/s13205-018-1171-9

Konishi, M., Okitsu, T., Yanagisawa, S., 2021. Nitrate-responsive NIN-like protein transcription factors perform unique and redundant roles in Arabidopsis. Journal of Experimental Botany 72, 5735–5750. https://doi.org/10.1093/jxb/erab246

Konishi, M., Yanagisawa, S., 2019. The role of protein-protein interactions mediated by the PB1 domain of NLP transcription factors in nitrate-inducible gene expression. BMC plant biology 19, 90. https://doi.org/10.1186/s12870-019-1692-3

Konishi, M., Yanagisawa, S., 2013. Arabidopsis NIN-like transcription factors have a central role in nitrate signalling. Nature Communications 4, 1617. https://doi.org/10.1038/ncomms2621

Konishi, M., Yanagisawa, S., 2011. Roles of the transcriptional regulation mediated by the nitrate-responsive cis-element in higher plants. Biochemical and Biophysical Research Communications 411, 708–713. https://doi.org/10.1016/j.bbrc.2011.07.008

Li, C., Goryaynov, A., Yang, W., 2016. The selective permeability barrier in the nuclear pore complex. Nucleus 7, 430–446. https://doi.org/10.1080/19491034.2016.1238997

Liu, K.-H., Liu, M., Lin, Z., Wang, Z.-F., Chen, B., Liu, C., Guo, A., Konishi, M., Yanagisawa, S., Wagner, G., Sheen, J., 2022. NIN-like protein 7 transcription factor is a plant nitrate sensor. Science (New York, N.Y.) 377, 1419–1425. https://doi.org/10.1126/science.add1104

Liu, K.-H., Niu, Y., Konishi, M., Wu, Y., Du, H., Sun Chung, H., Li, L., Boudsocq, M., McCormack, M., Maekawa, S., Ishida, T., Zhang, C., Shokat, K., Yanagisawa, S., Sheen, J., 2017. Discovery of nitrate-CPK-NLP signalling in central nutrient-growth networks. Nature 545, 311–316. https://doi.org/10.1038/nature22077

Loudet, O., Chaillou, S., Merigout, P., Talbotec, J., Daniel-Vedele, F., 2003. Quantitative Trait Loci Analysis of Nitrogen Use Efficiency in Arabidopsis. Plant Physiol 131, 345–358. https://doi.org/10.1104/pp.102.010785

Marchive, C., Roudier, F., Castaings, L., Bréhaut, V., Blondet, E., Colot, V., Meyer, C., Krapp, A., 2013. Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. Nature Communications 4, 1713. https://doi.org/10.1038/ncomms2650

Margalha, L., Elias, A., Belda-Palazón, B., Peixoto, B., Confraria, A., Baena-González, E., 2023. HOS1 promotes plant tolerance to low-energy stress via the SnRK1 protein kinase. Plant J 115, 627–641. https://doi.org/10.1111/tpj.16250

Marschner, H., 1995. Mineral Nutrition of Higher Plants, 2nd ed. Academic Press, London, England.

Möglich, A., Ayers, R.A., Moffat, K., 2009. Structure and signaling mechanism of Per-ARNT-Sim domains. Structure (London, England: 1993) 17, 1282–1294. https://doi.org/10.1016/j.str.2009.08.011

Mori, K., Murakoshi, Y., Tamura, M., Kunitake, S., Nishimura, K., Ariga, H., Tanaka, K., Iuchi, S., Yotsui, I., Sakata, Y., Taji, T., 2024. Mutations in nuclear pore complex promote osmotolerance

in Arabidopsis by suppressing the nuclear translocation of ACQOS and its osmotically induced
immunity. Front Plant Sci 15, 1304366. https://doi.org/10.3389/fpls.2024.1304366

Mu, X., Luo, J., 2019. Evolutionary analyses of NIN-like proteins in plants and their roles in nitrate
signaling. Cellular and molecular life sciences: CMLS 76, 3753–3764.
https://doi.org/10.1007/s00018-019-03164-8

Niemann, V., Koch-Singenstreu, M., Neu, A., Nilkens, S., Götz, F., Unden, G., Stehle, T., 2014.
The NreA protein functions as a nitrate receptor in the staphylococcal nitrate regulation system.
Journal of Molecular Biology 426, 1539–1553. https://doi.org/10.1016/j.jmb.2013.12.026

Pascual-Garcia, P, Capelson M. Nuclear pores in genome architecture and enhancer function. Curr
Opin Cell Biol. 2019 Jun;58:126-133. doi: 10.1016/j.ceb.2019.04.001. Epub 2019 May 4. PMID:
31063899; PMCID: PMC7347242.

Raveh, B., Karp, J.M., Sparks, S., Dutta, K., Rout, M.P., Sali, A., Cowburn, D., 2016. Slide-and-
exchange mechanism for rapid and selective transport through the nuclear pore complex.
Proceedings of the National Academy of Sciences of the United States of America 113, E2489-
2497. https://doi.org/10.1073/pnas.1522663113

Scheible, W.-R., Lauerer, M., Schulze, E.-D., Caboche, M., Stitt, M., 1997. Accumulation of nitrate
in the shoot acts as a signal to regulate shoot-root allocation in tobacco. The Plant Journal 11,
671–691. https://doi.org/10.1046/j.1365-313X.1997.11040671.x

Smith, C., Lari, A., Derrer, C.P., Ouwehand, A., Rossouw, A., Huisman, M., Dange, T., Hopman,
M., Joseph, A., Zenklusen, D., Weis, K., Grunwald, D., Montpetit, B., 2015. In vivo single-particle
imaging of nuclear mRNA export in budding yeast demonstrates an essential role for Mex67p.
The Journal of Cell Biology 211, 1121–1130. https://doi.org/10.1083/jcb.201503135

Strambio-De-Castillia, C., Niepel, M., Rout, M.P., 2010. The nuclear pore complex: bridging nuclear
transport and gene regulation. Nature Reviews. Molecular Cell Biology 11, 490–501.
https://doi.org/10.1038/nrm2928

Su, Y., Lagarias, J.C., 2007. Light-independent phytochrome signaling mediated by dominant GAF
domain tyrosine mutants of Arabidopsis phytochromes in transgenic plants. The Plant Cell 19,
2124–2139. https://doi.org/10.1105/tpc.107.051516

Tamura, K., Fukao, Y., Iwamoto, M., Haraguchi, T., Hara-Nishimura, I., 2010. Identification and
characterization of nuclear pore complex components in Arabidopsis thaliana. The Plant Cell
22, 4084–4097. https://doi.org/10.1105/tpc.110.079947

Ungricht, R., Kutay, U., 2017. Mechanisms and functions of nuclear envelope remodelling. Nature
Reviews. Molecular Cell Biology 18, 229–245. https://doi.org/10.1038/nrm.2016.153

Van de Vosse, D.W., Wan, Y., Wozniak, R.W., Aitchison, J.D., 2011. Role of the nuclear envelope
in genome organization and gene expression. Wiley Interdisciplinary Reviews. Systems Biology
and Medicine 3, 147–166. https://doi.org/10.1002/wsbm.101

Wang, H., Han, C., Wang, J.-G., Chu, X., Shi, W., Yao, L., Chen, J., Hao, W., Deng, Z., Fan, M.,
Bai, M.-Y., 2022. Regulatory functions of cellular energy sensor SnRK1 for nitrate signalling
through NLP7 repression. Nature Plants 8, 1094–1107. https://doi.org/10.1038/s41477-022-
01236-5

Wang, J., Pei, G., Wang, Y., Wu, D., Liu, X., Li, G., He, J., Zhang, X., Shan, X., Li, P., Xie, D., 2023.
Phase separation of the nuclear pore complex facilitates selective nuclear transport to regulate

plant defense against pathogen and pest invasion. Molecular Plant 16, 1016–1030.
https://doi.org/10.1016/j.molp.2023.04.008

Wozniak, R., Burke, B., Doye, V., 2010. Nuclear transport and the mitotic apparatus: an evolving
relationship. Cellular and molecular life sciences: CMLS 67, 2215–2230.
https://doi.org/10.1007/s00018-010-0325-7

Wu, X., Han, J., Guo, C., 2022. Function of Nuclear Pore Complexes in Regulation of Plant Defense
Signaling. International Journal of Molecular Sciences 23, 3031.
https://doi.org/10.3390/ijms23063031

Xiao, L., Jiang, S., Huang, P., Chen, F., Wang, X., Cheng, Z., Miao, Y., Liu, L., Searle, I., Liu, C.,
Wu, X.-X., Fu, Y.-F., Chen, Q., Zhang, X.-M., 2020. Two Nucleoporin98 homologous genes
jointly participate in the regulation of starch degradation to repress senescence in Arabidopsis.
BMC plant biology 20, 292. https://doi.org/10.1186/s12870-020-02494-1

Zhang, A., Wang, S., Kim, J., Yan, J., Yan, X., Pang, Q., Hua, J., 2020. Nuclear pore complex
components have temperature-influenced roles in plant growth and immunity. Plant Cell Environ
43, 1452–1466. https://doi.org/10.1111/pce.13741

Figures and Tables
648
649   **Figure 1. NLP7 interacts with NUP98A.**

650
651
652   **Figure 1. NLP7 interacts with NUP98A.**
653
(**A**) Yeast two-hybrid assay using NLP7 fused with the binding domain (BD) and NUP98A fused
with the DNA activation domain (AD). Cells were grown on selective and non-selective plates.
(**B-G**) NLP7 interacts with NUP98A in Bimolecular Fluorescence Complementation (BiFC) assays
in *N. benthamiana.*
(**B, E**) YFP fluorescence (green), (**C, F**) chlorophyll autofluorescence (magenta), (**D, G**) H2B
nuclear marker - mTurcoise fluorescence (blue)
Scale bar = (**B-D**) 50 μm, (**E-F**) 5 μm
661
662
663

664 **Figure 2.**

665
666
Figure 2. NLP7 interacts with NUP98A and NUP98B.
(A-L) NLP7 interacts with NUP98A and NUP98B in BiFC assays in *N. benthamiana.* (**A-C**) NLP7-
NUP98A interaction, (**D-F**) NLP7-NUP98B interaction, **(G-I)** NLP7-CPR5 interaction (negative
control), **(J-L)** NLP7-NUP96 interaction (negative control). (**A, D, G, J**) YFP fluorescence (green),
(**B, E, H, K**) chlorophyll autofluorescence (magenta) (**C, F, I, L**) YFP fluorescence (green) was
merged with chlorophyll autofluorescence (magenta).
Scale bar = 25 µm.
674

 Figure 3.

^{C}YFP-NLP7 ^{N}YFP-NUP98A

Figure 3. **NLP7 interacts with NUP98A in the nucleus in the presence of nitrate**

(A-I) BiFC assays between NLP7 and NUP98A in *N. benthaminia*. (**A-C**) Plants were grown under ample (10 mM) nitrate nutrition (**D-F**) Plants have been N-depleted for 5-days **(G-I)** leaf disc from transitory transformed N-depleted plants were treated for 10 min with nitrate. (A, D, G) YFP fluorescence (green), (B, E, H) chlorophyll autofluorescence (magenta) (C, F, I) YFP fluorescence (green) was merged with chlorophyll autofluorescence (magenta). Scale bar = 50 µm.   **(J)** Nuclear-to-cytoplasmic fluorescence intensity ratios in plant cells across different samples. Ratios were calculated from measurements in 13 - 15 images per genotype and treatment. Data are presented as boxplots. Statistical significance was assessed using the Kruskal–Wallis test followed by Dunn's post hoc test for pairwise comparisons. *P*-values: 4.47e-02 (**), 2.7e-07 (***), 1.08e-03 (###) .

678 **Figure 4.**

679
Figure 4. Biomass and Shoot/Root ratio of *nup98a/b single and double* mutants grown
under limiting and non-limiting nitrate supply.
Plants were grown on sand under short days with a light intensity of 150 µE for 39 d (**A, B**)
Rosette fresh weight of plants grown under limiting (0.5 mM, **A**) and non-limiting (5 mM, **B**) nitrate
supply. (**C, D**) Root fresh weight of plants grown under limiting (0.5 mM, **C**) and non-limiting (5
mM, **D**) nitrate supply. (**E, F**) Shoot/root ratio of plants grown under limiting (0.5 mM, **E**) and non-
limiting (5 mM, **F**) nitrate supply. Data were obtained from six plants from two independent
experiments. Significant differences between genotypes for each nitrate supply condition were
determined by Student t-Test (* *P*-value ≤ 0.05). **(G, H)** Top view of rosettes of Col-0 and
nup98anup98b-double mutants grown under limiting (0.5 mM, **G**) and non-limiting (5 mM, **H**)
nitrate supply at different times of the culture. Scale bar = 2 cm.
691

Figure 5.

694
Figure 5. Growth and gene expression in response to nitrate are strongly impacted by the
simultaneous loss of function of Nucleoporin 98A/B and NLP7. (A-C) Plants were grown on
sand under short days with a light intensity of 150 µE for 39 days (**A, B**) Rosette fresh weight of
plants grown under limiting (0.5 mM, **A**) and non-limiting (5 mM, **B**) nitrate supply. **(C)** Top view of
rosettes of plants grown under non-limiting (5 mM) nitrate supply, Scale bar = 2 cm. **(D-F)** Gene
expression analysis. Relative transcript levels of selected genes before and after nitrate addition
to N-starved 14-day-old seedlings. Expression was measured by RT-qPCR and normalized as a
percentage of the expression of a constitutive synthetic gene composed of *TIP41* (At4g34270)
and *ACTIN2* (*ACT2*, At3g18780). **(G)** Heatmap representation of the fold-change in steady state
transcript level after 30 min of nitrate resupply in comparison to Col-0. Red, lower abundance;
blue, higher abundance (* *P*-value ≤ 0.05 in comparison to *nlp7-1*).
706

707   **Figure 6.**

N starvation   **Nitrate resupply (10 min)**

A   B

NLP7pro:GFP-NLP7 (nlp7-1)

C   D

NLP7pro:GFP-NLP7 (nlp7-1nup98a-1)

E   F

NLP7pro:GFP-NLP7 (nlp7-1nup98a-1nup98b-1)

G

Figure 6. **NUP98A *and* NUP98B loss of function abolished the nitrate-mediated nuclear accumulation of NLP7.**

Analysis of the nitrate-triggered nuclear accumulation of GFP-NLP7. **(A-H)** 10-days-old seedlings were N starved for 2 days before observation by confocal microscopy. Acquisitions were performed before nitration addition **(A, C, E)**, or 10 min after nitrate addition **(B, D, F)**.

Representative images out of 3 independent experiments are shown. Scale bar = 50 μm.

(**G**) Nuclear-to-cytoplasmic fluorescence intensity ratios in plant cells across different samples. Ratios were calculated from measurements in 7 - 15 images per genotype and treatment. Data are presented as boxplots. Statistical significance was assessed using the Kruskal–Wallis test followed by Dunn's post hoc test for pairwise comparisons. *P*-values: 1.49e-3 (**), 6.5e-6 (***). Black squares (■) and black downward triangles (▼): regular outliers; white upward triangle (△): extreme outlier.

708
709

710 **Supplemental Table S1**. Primers used for cloning, genotyping and RT-qPCR analyses.
711

			Primer sequences	
			Forward	Reverse
Cloning	NLP7 into pDONR207		GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGTGCGAGCCCGATGATAATTCCG	GGGGACCACTTTGTACAAGAAAGCTGGGTCCAATTCTCCAGTGCTCTCGCAGGAG
	NLP7 without PB1 into pDONR207		GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGTGCGAGCCCGATGATAATTCCG	GGGGACCACTTTGTACAAGAAAGCTGGGTCTCTCATTTCTGAGCCTGATGG
	NLP7 GAF-like domain deletion		GACTCAGGCGCTTCGATACTTCAAAGCGGTAAATCTGAAAAGCTCGGAAATAC	GTATTTCCGAGCTTTTCAGATTTACCGCTTTGAAGTATCGAAGCGCCTGAGTC
	NUP98B into pDONR207		GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGTTCGGTTCTTCTAATAATAATCC	GGGGACCACTTTGTACAAGAAAGCTGGGTCCACATCATATTCATCTCCAAGCTTGTAG
	CPR5 into pDONR207		GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGGAAGCCCTCCTCCTCCCTCCTTCG	GGGGACCACTTTGTACAAGAAAGCTGGGTCAGCATAGTCAGACCCACCATGAAGGTG
	NUP96 into pDONR207		GGGGACAAGTTTGTACAAAAAAGCAGGCTTCATGACGTCTCTTTCTATTCAACCATTTC	GGGGACCACTTTGTACAAGAAAGCTGGGTCAGCTGAGATTTGGCCCGTCTCTGAGAG
Genotyping	*nlp7-1* - gene primers		GTTTCCAGTTTCCGTCTCCA	GGCCAAATGATGAGCAGAA
	nlp7-1 - T-DNA primer		GTTTCCAGTTTCCGTCTCCA	LBb1: GCGTGGACCGCTTGCTGCAACT
	nup98a1 - gene primers		ATTGGTCGAAGAACGTGTGTC	GAAAATCCCAAAGGTTTCTGC
	nup98a1 - T-DNA primer		LBb1.3: ATTTTGCCGATTTCGGAAC	GAAAATCCCAAAGGTTTCTGC
	nup98b - gene primers		GTTTGTCCAGGTTCCATTCTTCAGC	TCCAATCTCTCGCCTTCCATCAC
	nup98b - T-DNA primer		Lb3sail: TAGCATCTGAATTTCATAACCAATCTCGATACAC	TCCAATCTCTCGCCTTCCATCAC
	Gene	**Accession number**	**Forward**	**Reverse**
RT-qPCR	*TIP41*	AT4G34270	GCTCATCGGTACGCTCTTTT	TCCATCAGTCAGAGGCTTCC
	ACT2	At3g18780	GCCATCCAAGCTGTTCTCTC	CCCTCGTAGATTGGCACAGT
	NRT2.1	AT1G08090	AGTCGCTTGCACGTTACCTG	ACCCTCTGACTTGGCGTTCTC
	NRT2.2	AT1G08100	AATGTCATGTTTGGTGAGGTTAAGA	GCTATGCCTTCTCGGTAGATGGTAG
	NLP1	At2g17150	TAGTTGCTGAGGATGCTGATG	GGCACCTCTAGCTCTTAAACT
	NII	AT2G15620	TGCTGATGACGTTCTTCCACTCTGC	CTGAGGGTTGACTCCGAAATAGTCTC
	NIA1	AT1G77760	ATCGTCAAAGAAACCGAAGTC	ACGGAGCATGGATGAGTT
	NAR2	At5g50200	CCAGAAGATCCTCTTTGCTTCACT	CCCAATCGAGCTTAGCGTCCA
	LBD38	AT3G49940	TGCCCTGCTTTGTTTCAGTCTT	CGTTCACCGGATTCACAGTTCT
	NLP2	At4g35270	TTGTCATCATCCAGGTACTTA	CGGAGAATAGTAGCAGGAAGT
	HHO1	AT3G25790	ACAACACCGAACAACAGAA	GCAACGAACTGACCATAGG
	BT5	AT4G37610	ACAAAGACCGAAACCCCG	CATTGTTCCGAGTCTACGCA
	NLP6	At1g64530	GAGACGCGTTTTCGGCTTCCTA	TTGAACCTGATGATGTCTTCTCTGTA
	MAP3K14	AT2G30040	ACCGTCATCGAGATGCTCAC	AGAGAACCCGATCCGACTCA
	NLP3	At4g38340	GGTCGTCGGACATACACCAA	GGACACTCTCCGACAACACT
	BT2	At3g48360	TGCAAGAGGAGGAATAACGAT	CACAACGACGTCATAGCTTTT
	HRS1	At1g13300	CCTAACAACGGAAACTCTCAA	TCCACAATAATCGACGGTCTA
	NIA2	AT1G37130	GGTTACGCATATTCCGGAG	CATGCACGAACAGCAATC
	MAP3K13	AT1G07150	CGCGGCGAAGGCAATAATTT	CACCCAACCATCTGACTCCC
	LBD39	AT4G37540	GAACTCCAACGTCCTGCTTTGT	ATACCAACCGCTCCGTTAACC
	NRT1.1	At1g12110	AGACCGAACCAAAAGAACGA	CCACGATAACCGCAGCAACC

Supplemental Figure 1

Supplemental Figure S1. NLP7 GAF and PB1 domains are not essential for the NLP7-
NUP98A interaction

NLP7ΔGAF and NLP7ΔPB1, interact with NUP98A in BiFC assays in *N. benthamiana* (**A-C**) NLP7-
NUP98A interaction, (**D-F**) NLP7DGAF-NUP98A interaction, **(G-I)** NLP7DPB1-NUP98A
interaction. (**A, D, G**) YFP fluorescence (green), (**B, E, H**) chlorophyll autofluorescence (magenta)
(**C, F, I**) YFP fluorescence (green) was merged with chlorophyll autofluorescence (magenta). Scale
bar = 50 μm.

Graphical abstract