1 TITLE: UPSLOPE DIVERSIFICATION AND GENE FLOW AMONG CONGENERIC
2 ANDEAN TREES
3

4 Ellen J. Quinlan[1*], James B. Pease[2], Jhonatan Sallo Bravo[3], Alfredo F. Fuentes[4,5], William
5 Farfan-Rios[1,6], and Miles R. Silman[1,6]
6
[1]Department of Biology, Wake Forest University, Winston-Salem, NC, USA 27109
[2]Department of Evolution, Ecology, and Organismal Biology, The Ohio State University,
Columbus, OH, USA 43210
[3]Facultad de Ciencias Biológicas, Universidad Nacional de San Antonio Abad del Cusco, Cusco,
Perú
[4]Latin America Department, Missouri Botanical Garden, St. Louis, MO, USA 63110
[5]Herbario Nacional de Bolivia, Universidad Mayor de San Andrés, La Paz, Bolivia
[6]The Sabin Center for Environment and Sustainability, Wake Forest University, Winston-Salem,
NC, USA 27109
16
*Corresponding Author: Ellen J. Quinlan, quinej18@wfu.edu, ORCID: 0000-0002-6249-649X

SUMMARY

• The exceptional biodiversity of tropical Andean forests is characterized by rapid turnover

of many closely related species across elevation and other environmental gradients. The

eco-evolutionary relationships among these congeneric species are often poorly

understood—specifically, how they diversify, sort into their elevational niches, and the

extent to which species boundaries are maintained through time.

• In this study, we examined the phylogenetic relationships and patterns of ancient and

ongoing gene flow among sympatric *Prunus* species occurring along elevation gradients

in Peru and Bolivia to explore the evolutionary processes underlying this turnover.

• Phylogenetic relationships reconstructed a pattern of upslope adaptive diversification

concurrent with Andean orogeny during the Miocene. Despite deep divergence times,

species show evidence of widespread and persistent gene flow, with little constraint

imposed by ecological or evolutionary distance.

• Together, these findings show that during adaptive radiations, observable ecological and

morphological differences may emerge despite ongoing gene flow and long before

reproductive isolation is complete.

Keywords: Andes, biodiversity, diversification, elevation gradient, gene flow, hybridization,

tropical trees

 INTRODUCTION

Clarifying how species diversify and assemble into communities, and the degree to which

species boundaries persist through time is essential for understanding the origin and maintenance

of tropical tree biodiversity and its conservation. Coexistence theories of ecology generally

assume species are discrete evolutionary units (e.g. Hubbell 2000, Chesson 2000), though

interspecific gene flow is known to have been common throughout the evolutionary history of

plants (reviewed by Stull et al. 2023). No system lends itself to tests of these theories better than

hyper-diverse tropical forests. Until recently, hybridization was considered an extremely rare

occurrence in tropical trees, as hybrids were thought to be poor competitors in high-diversity

systems and few individuals with intermediate morphologies have been identified (e.g. Ashton

1969; Ehrendorfer 1970; Gentry 1982). Mounting genomic evidence suggests hybridization has

been widespread throughout the diversification of the Neotropical flora (reviewed by Schley et

al. 2022) and may be an important evolutionary mechanism for maintaining high tree diversity

with low population densities (Cannon and Lerdau 2015; Quinlan et al. 2025). These findings

collectively point to the importance of reevaluating the impact of hybridization in our

understanding of species diversity and coexistence theories in tropical ecosystems.

The tendency for inter-specific gene flow and hybridization to occur amongst closely

related, sympatric trees has been widely documented in temperate zones (De La Torre et al.

2014, Tsuda et al. 2017, Cavender-Bares et al. 2018, Chhatre et al. 2018, Menon et al. 2018), yet

species cohesion and morphological similarity is often maintained (Grant 1971; Gailing and

Curtu 2014; Cannon and Petit 2019). This may be partially due to assortative mating and variable

mating behaviors (selfing, conspecific outcrossing, and interspecific gene flow), determined by

population densities and pollinator abundance (Hamrick and Murawski 1990; Ward et al. 2005;

Cannon and Lerdau 2015). When present, conspecific pollen can outcompete heterospecific

pollen, reinforcing assortative mating (Williams et al. 1999). However, incomplete reproductive

isolation and the ability to accept heterospecific pollen may become adaptively beneficial when

conspecific pollen is rare, as the benefit of avoiding local extinction overrides potential negative

consequences of hybridization such as reduced hybrid fitness (Barton 2001; Wu 2001; Cannon

and Lerdau 2015). Hybridization may be particularly beneficial in times of rapid environmental

change, as it can produce novel gene combinations within a single generation, increasing

adaptive potential and population resilience (Anderson and Stebbins 1954; Baskett and

Gomulkiewicz 2011; Suarez-Gonzalez 2018; Brauer et al. 2023; Kulmuni et al. 2024). Such

density-dependent mating has been reported in oaks (Lepais et al. 2009) and may be particularly

relevant to tropical trees which maintain extremely low local population densities (Pires et al.

1953; Pitman et al. 1999; ter Steege et al. 2013).

Most tropical forest trees are highly outcrossed (Bawa et al. 1985a, b; Hamrick and

Murawski 1990; Ward et al. 2005), though individuals retain variable capacities for different

mating behaviors, even within populations and across flowering events (Murawski and Hamrick

1991; Nason and Hamrick 1997). Self-fertilization has been shown to be inversely correlated

with population density (Hamrick and Murawski 1990; Ward et al. 2005), but few studies have

assessed the extent of inter-specific gene flow. Those that have (e.g. Schley et al. 2020 –

Brownea, Larson et al. 2021 – *Eschweilera,* Schley et al. 2025 – *Inga*), detected widespread

admixture. Evolutionary distance may be one important constraint on interspecific gene flow

among congeners in sympatry, as hybridization is thought to decrease with increasing divergence

due to the accumulation of pre- and postzygotic barriers through time (Mendelson et al. 2004;

Levin 2012; Comeault and Matute 2018). Linan et al. (2021) found evidence of this in the

Mascarene Island *Diospyros* clade, suggesting the processes by which species diversify and

assemble into communities may have lasting effects on their eco-evolutionary dynamics. For

example, adaptive radiations across gradients typically result in stepwise diversification patterns

with sister taxa occurring closest in niche proximity (Simpson 1953; Schluter 2000; Stroud and

Losos 2016), whereas communities assembled predominantly via the immigration and sorting of

pre-adapted clades demonstrate greater phylogenetic distance among co-occurring taxa

(Donoghue 2008; Linan et al. 2021).

In tropical montane forests, genera commonly comprise many rare species that turn over

rapidly with elevation (Terborgh 1971; Gentry 1982). Turnover in the abiotic environment and

biotic interactions leads to narrow, shoestring-like distributions and the packing of many closely

related species across elevation gradients (e.g. Janzen 1967, Jankowski et el. 2009, Hillyer and

Silman 2010, Freeman et al. 2022). In the tropical Andes, this is consistent across taxa and is

especially true for forest trees (e.g. Terborgh 1977, Meier et al. 2010, Jankowski et al. 2012).

While many tree species have elevation ranges of < 500 m, the distributions of entire genera are

often much broader, with congeneric species occurring in partial sympatry, especially at range

margins (Feeley et al. 2011; Griffiths et al. 2020). At the genus level, immigration and sorting of

pre-adapted clades has been shown to be the primary mechanism structuring tree diversity across

elevation gradients in the Andes (Griffiths et al. 2020; Segovia et al. 2020; Linan et al. 2021).

However, many examples of adaptive diversification have been found within individual plant

clades (e.g. Nörk et al. 2013, Lagomarsino et al. 2016, Nevado et al. 2016, among many others).

Still, few studies have explored how congeners within individual tree clades have diversified and

assembled into their elevation niches, and characterizing these evolutionary relationships is

essential for understanding potential gene flow and its ecological implications.

Prunus L. (Rosaceae) is one such genus that exhibits high congeneric turnover with

elevation in the Andes (Figure 1). Globally, it has diversified into a range of environments, and

though typically considered a temperate group (Wen et al. 2008; Chin et al. 2014), a recent

revision found its highest diversity occurs in the tropics and primarily the Neotropics (Pérez-

Zabala 2022). *Prunus* is one of the only tree genera with species spanning the entire Andes-

Amazon gradient, from treeline (~3800 m) into the lowlands (<500 m), though individual species

have more narrow ranges, typically 500–1000 m or less (e.g. Achá 2013). It is a keystone clade,

whose fruits are important in the diets of frugivorous birds, primates, Andean bears, and humans

(Achá 2013). The genus also contains many globally important crop species (peach, plum,

almond, cherries) and others used as ornamentals, timber, and medicine (Andro and Riffaud

1995; Bortiri et al. 2001; Lee and Wen 2001; Wen et al. 2008).

The genus *Prunus* is divided into three major clades that correspond to inflorescence

type: solitary, corymbose, and racemose (Rehder 1940, Su et al. 2023). All species endemic to

the Americas and tropical regions are racemose (Perez-Zabala 2022). Though historically

understudied, the exclusively South American species form a strongly supported clade nested

within the tropical Racemose group (Hodel et al. 2021, 2023). Studies have placed the origin of

the genus in either North America (Chin et al. 2014) or East Asia (Hodel et al. 2023) during the

Paleocene (ca. 68–55 Ma), with the three major clades diverging by the early-mid Eocene as part

of the Boreotropical flora (Chin et al. 2014; Su et al. 2023; Hodel et al. 2023). Divergence in the

tropical Racemose group occurred in several regions including Asia, Africa, North America, and

Europe ca. 55–40 Ma (Hodel et al. 2023). Chin et al. (2014) hypothesized migration into South

America occurred via the dispersal of a North American relative ca. 34 Ma through Central

America and the Caribbean. Hodel et al. (2023) found evidence of a long period of stasis in the

Neotropical lineages followed by rapid diversification during the Miocene.

While most species in the solitary-flower and corymbose groups are diploids, the

racemose lineage is known for higher ploidy levels across temperate and tropical species and the

lineage is thought to be the result of a single or multiple ancient hybridization events during the

Eocene (Chin et al. 2104, Zhao et al. 2016, Su et al. 2023, Hodel et al. 2023). Domesticated

Prunus (particularly plums) readily hybridize in agricultural settings (Cici and Acker 2010),

resulting in the creation of some crops (e.g. *P. armeniaca* L. × *P. cerasifera* Ehrh. – 'black

apricots,' *P. salicina* × *P. armeniaca* Lindl. – plumcots, pluots, and apriums; Guerrero et al.

2022), as well as among cultivated varieties and wild conspecifics (*P. yedoensis* - Baek et al.

2018). Species in this genus are highly outcrossing and typically self-incompatible with

generalist insect pollinators (Cici and Acker 2010), though we are unaware of any study that has

specifically looked at either breeding systems or pollination in Neotropical *Prunus*.

In this study, we investigate the patterns, timing, and extent of divergence among

partially sympatric tree species which turn over across elevation gradients in the tropical Andes,

and the degree to which species boundaries are maintained through time. Specifically, we ask:

(1) how and when co-occurring species diversified into their elevational niches, (2) whether

species are genetically isolated, and, if not, (3) is introgressive gene flow related to evolutionary

or ecological distance. We reconstructed the phylogenetic relationships among sympatric *Prunus*

species, estimated the timing of divergence, and tested for historical introgression and recent

admixture. The results of these analyses expand current understanding of the origin and

maintenance of tropical Andean tree diversity and provide important context for predicting how

these species may respond to rapid environmental change.

 MATERIALS AND METHODS

Study site and sample collection

Sampling for this study primarily focused on species which co-occur across the Andes

Biodiversity and Ecosystem Research Group (ABERG; andesbiodiversity.org) plot network in

Manu National Park, Peru (Figure 1). This site is located on the eastern slope of the tropical

Andes in southeastern Peru and is a 3.5 km elevational gradient from the Andean tree line to the

Amazonian lowlands. With 23-1 ha and 33-0.1 ha forest plots along a series of descending

ridges, this is the most continuous transect in the tropical Andes. To date, ~19,000 individual

trees have been documented in these plots from 100 families, 292 genera, and 1255 species

(Farfan-Rios et al. 2025).

Intensive field sampling was conducted for *Prunus* species along this transect, with

botanists searching in lateral transects centered on the 1-ha plots (approximately every 250 m)

for up to 6 hrs. Leaf tissue was field collected and stored in silica following the Smithsonian

Institution guidelines for collecting vouchers and tissues intended for genomic work (Funk et al.

2017). Samples were also included for species known to co-occur across an additional plot

network spanning 4 km of elevation in Madidi National Park, Bolivia, and from at least one

external population when possible. These samples were collected from herbarium specimens

previously not treated with alcohol or fresh tissue stored in silica collected from individuals in

the Cotapata region of Bolivia (Figure 1, Table S1). In total, we sampled 158 individuals from 13

species including two outgroups (*P. myrtifolia* and *P. caroliniana*) and a reference (Table S1).

ddRAD-seq data

High molecular weight DNA was extracted from ~50 mg of dried leaf tissue using Aboul-

Maaty and Oraby's (2019) modified CTAB protocol for non-model plants with a few

modifications: samples were ground with a bead-beater, incubated at -20 ℃ overnight (and up to

24 hrs), and the DNA pellet was washed twice with 500 µl 70% ethanol. DNA was checked for

quality and quantity on a 1.5% agarose gel and Qubit before being sent to Floragenex

(Beaverton, Oregon, USA) for ddRAD library prep and sequencing. Samples were ligated with 6

bp barcodes and digested with the PstI/MseI+0 enzyme pair. Genomic data were demultiplexed

and assembled to sample using ipyrad v. 0.9.95 (Eaton and Overcast 2020) with an average of

$2,215,444 \pm 1,419,356$ million reads (mean $\pm$ s.d.) recovered per sample.

Reads were assembled into orthologous loci using reference-based assembly in ipyrad.

Default settings were used to trim and filter reads and the clustering threshold was set to 0.85 as

recommended by the ipyrad manual and McCartney-Melstad et al. (2019) for phylogenetic

analyses. For the reference-based assembly, reads were mapped to the *Prunus persica* (peach)

genome (GCF_000346465.2; The International Peach Genome Initiative et al. 2013, Verde et al.

2017). The proportion of mapped reads varied across samples (mean = 67%, s.d. = 13%), but

variation in the quality of sample preservation and DNA extractions likely explain much of the

variation. When calling consensus alleles, we allowed for up to 4 alleles per sample following

Donoghue et al. (2022), as all *Prunus* species in the tropical Racemose group studied to date are

tetraploids (2n=32; Hodel et al. 2023), but ploidy is not known for most of the species in our

study. The final reference-mapped dataset comprised 35,099 loci, representing non-overlapping

genomic regions where data from at least 4 samples passed filtering. Further downstream

analyses were performed using vcftools and the ipyrad-analysis toolkit, which allows for filtering

and subsampling loci, genomic windows, and/or SNPs for phylogenetic inference and D-

statistics.

Individual-level phylogeny

A maximum likelihood (ML) phylogeny was reconstructed from a subset of

representative samples with the most data present per species (up to 10 individuals per species, at

least 1000 loci per sample; n=47), including conspecifics from multiple localities when possible.

The ipa.window_extracter module in the ipyrad-analysis toolkit was used to extract data from the

8 largest scaffolds (the chromosomes) and filter sites to only maintain those shared by at least

25% of individuals. We selected the 25% threshold after testing various other filtering thresholds

(10%, 30%, 40%, 50%), as 25% provided for the best phylogenetic resolution while also

including as much information as possible. The filtered and concatenated sequence matrix is 2.8

Mb in length, containing 107,708 SNPs and 65.8% missing data. This alignment was used in

RAxML (Stamatakis 2014) under the GTR+GAMMA evolutionary model to infer a best

phylogeny under the rapid hill-climbing algorithm, with support values from 100 bootstrap

replicates. The phylogeny was rooted to the reference genome *P. persica* (a native of China and

member of the Solitary-flower group and Amygdalus subgenus) and included two additional

Racemose outgroup species – *P. caroliniana* (native of the eastern United States and Carribbean)

and *P. myrtifolia* (native to the Caribbean, Central America, and lowland South America).

Species-level phylogeny

To construct the species-level phylogeny, we selected one representative specimen for

each species in the individual-level phylogeny (n=13). The samples selected grouped with other

members of their species on the individual-level phylogeny and had the most data present. Again,

we used the ipa.window_extracter module in the ipyrad-analysis toolkit to extract data from the

chromosomes and filter sites, selecting those shared by at least 25% of all species (after testing

multiple filtering thresholds). The resulting sequence matrix is 4.2 Mb with 62,262 SNPs and

63.6% missing data. To reconstruct the phylogenetic relationships, we used both maximum

likelihood (RAxML) and Bayesian (ExaBayes) analyses. We ran RAxML using GTR+GAMMA

to infer a best phylogeny under the rapid hill-climbing algorithm with support values from 100

bootstrap replicates. For ExaBayes, we ran two Metropolis-coupling replicates with four

coupled-chains (each with three heated chains) for 1×10^6 MCMC generations, sampled every

500 generations. Estimated sample size (ESS) for all parameters and branch lengths were

summarized with the ProcParam tool and confirmed as sufficiently sampled. The majority-rule

consensus phylogeny was generated with the ExaBayes 'consense' tool with a 25% burn-in.

To assess the robustness of the resolved relationships and distinguish between conflicted

and poorly informed branches, we used Quartet Sampling (Pease et al. 2018;

https://www.github.com/ fephyfofum/quartetsampling). We performed Quartet Sampling on the

resulting RAxML tree with the full concatenated alignment, allowing 100 replicates per branch

(Figure S2).

Divergence time dating

To assess the timing of divergence among species, we generated a time-calibrated

phylogeny using BEAST v. 2.7.7 (Bouckaert et al., 2019) with the same alignment used for the

species-level phylogenies. We used an optimized relaxed molecular clock and Yule speciation

process with the GTR+GAMMA substitution model. A nodal calibration was set for the crown

age of *Prunus* using a normal prior with an offset of 61.5 Mya and a standard deviation of 3.0

Mya (Chin et al. 2014; Su et al. 2023). The crown age of the racemose group was calibrated from

a staminate flower of *Prunus hirsutipetala* D.D. Sokoloff, Remizowa et Nuraliev preserved in

the late Eocene (Priabonian) Rovno amber from northwestern Ukraine (Sokoloff et al. 2018),

using a normal prior with an offset of 35.0 Mya and standard deviation of 2.0 Mya (following Su

et al. 2023). To ensure a more rapid convergence of MCMC runs, we used our RAxML topology

as our starting tree. The Markov chain Monte Carlo (MCMC) run was conducted for 20,000,000

generations and trees sampled every 2000 generations producing a total of 10,000 trees. The

resulting log files were checked in Tracer v.1.5 (Rambaut et al. 2018) and we used TreeAnnotator

v.1.10.4 to generate the maximum clade credibility tree discarding 25% of trees as burn-in before

visualizing the results in FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Tests of historical introgression

To test for historical introgression, we used the four-taxon Patterson's D-statistic (ABBA-

BABA), which quantifies allele frequency patterns that are incongruent with the hypothesized

species tree topology (Green et al. 2010). We ran the Patterson's D tests using the ipa.baba tool as

implemented in ipyrad, which is specifically designed for analysis of RAD-seq datasets. We used

the species-level RAxML tree as our starting tree hypothesis and constrained the tests setting the

outgroup to *P. myrtifolia*, so all tests would focus on ingroup species. To assess significance, we

performed 1000 bootstrap replicates for each test with loci resampled with replacement. Z-scores

were then calculated as the number of bootstrap standard deviations from the null $D=0$ (Table 1).

Z-scores > 2.5 were considered significant.

247   *Tests of recent admixture*

To assess genetic structure and test for recent admixture, we used the Bayesian clustering

algorithm STRUCTURE v. 2.3.4 (Pritchard et al. 2000) as implemented in ipyrad with the set of

samples included in the individual-level phylogeny (minus the outgroups). SNPs were filtered

conservatively to only include those with at least 50% coverage across individuals and then

randomly sampled to one SNP per locus to avoid linkage. This resulted in a total of 1493 SNPs

included in the analysis. The model was run with a burn-in of 125,000 and run length of 500,000

generations, testing K=1–10 with 10 replicates per K. The assignment of samples into K distinct

genetic groups was assessed using DeltaK, which has been shown to be more robust to

accurately estimating the correct number of clusters in STRUCTURE, particularly with high

amounts of missing data as is common in RADseq (Evanno et al. 2005; Figure S3).

Estimates of genomic diversity

Genetic diversity was estimated within individuals and species to compare to the

admixture results. In VCFtools v.1.16 (Danecek et al. 2011), sequences were filtered using two

different missing data thresholds, retaining loci present within at least 50% or 70% of the

samples for each species. Sites were also filtered if they had a minor allele frequency less than

0.05. We calculated nucleotide diversity (π) for each species – the average pairwise nucleotide

differences between sequences in a sample at a site as well as observed (H_o) and expected

heterozygosity (H_e) – the proportion of loci where an individual is heterozygous vs. the

proportion expected by Hardy-Weinberg equilibrium – within samples and averaged across the

species. For these calculations, *P. debilis* and *P. detrita* samples were grouped together as well as

P. huantensis and *P. brittoniana,* given that STRUCTURE (Figure 4) did not detect any distinct

genetic clusters among these species' pairs. Species with < 3 samples were excluded from these

analyses given the lack of statistical power from such a small sample size.

281   RESULTS

Phylogenomic relationships and elevational niche diversification

Both the species-level maximum likelihood (RAxML) and Bayesian (ExaBayes)

phylogenetic analyses supported the same evolutionary relationships among the focal species and

their outgroups (Figure 2). Subtropical *P. caroliniana* diverged before the tropical species and *P.*

myrtifolia was sister to the Andean species. Among these partially sympatric Andean species,

diversification generally occurred from the lowlands to the highlands, with the lineage including

P. debilis and *P. detrita* diverging first, followed by the mid-elevation cloud-forest species (*P.*

herthae, P. pleiantha, P. stipulata, P. integrifolia) and finally the high-elevation species as the

most recent diversification events (*P. huantensis, P. brittoniana, and P. antioquensis*). Statistical

support for the inferred relationships was assessed using bootstrap support (RAxML), posterior

probabilities (ExaBayes), and Quartet Sampling. While the tree nodes are generally well-

supported (all posterior probabilities = 1, most bootstrap values > 75; Figure 2), Quartet

Sampling indicated that low bootstrap support values are not likely the result of insufficient

phylogenetic signal, as the proportion of informative quartets (QI) ranged from 80–100%. The

lack of a strong skew in the proportions of discordant phylogenetic trees at most nodes (QD >

0.5) suggests that incomplete lineage sorting (ILS) is more likely than introgression. The branch

ancestral to *P. huantensis, P. brittoniana*, and *P. antioquensis* had moderate concordance

(QC=0.21) and discordant skew (QD=0.22), indicating a possible introgressive event (Figure

S2). Together, these phylogenetic analyses point towards a history of diversification from the

lowlands into the highlands within Andean *Prunus*, with some evidence of incomplete lineage

sorting (ILS) and/or introgression.

Divergence time estimation

The time-calibrated phylogeny generated in BEAST recovered the same tree topology as

RAxML and ExaBayes with similar branch support (Figure 3). The estimated divergence time

for P. myrtifolia and the ancestor of the Amazonian and Andean species is ~27.3 Ma (95%

highest posterior density [HPD] 22.4–31.7). The Andean species began diverging at the start of

the Miocene, with the low-elevation *P. debilis+detrita* lineage branching off ~23.5 Ma (95%

HPD 19.4-28.1), and from each other ~17.6 Ma (95% HPD 12.8-22.6). The cloud forest species

followed, with *P. pleiantha* diverging ~20.4 Ma (95% HPD 16.3-24.4), and *P. integrifolia* ~18.0

Ma (95% HPD 14.2-21.7). The remaining lineages emerged in the mid to late-Miocene (15.9–

9.3) with the highest elevation clade diverging last, with *P. huantensis* ~11.6 Ma (95% HPD 8.6-

14.9) and *P. antioquensis+ brittoniana* ~9.3 Ma (95% HPD 6.2-12.3). In sum, these estimated

divergence times indicate that Andean *Prunus* diversified from the Amazonian lowlands into the

Andean highlands over the Miocene.

Tests of historical introgression and recent admixture

D-statistics indicated significant historical introgression events between *P. pleiantha* and

both the ancestors *P. debilis+detrita*, with more limited evidence suggesting introgression with *P.*

debilis and *P. detrita* separately (Figure 2, Table 1). A second significant introgression event was

detected between *P. pleiantha* and the *P. huantensis+antioquensis* ancestor, which is further

supported by Quartet Sampling (Figure S2). Finally, significant *D*-statistics supporting

introgression between *P. debilis* and *P. antioquensis* were also detected. Overall, introgression is

widespread and does not seem to be correlated with ecological or evolutionary distance.

STRUCTURE analysis found K=9 to be the most likely number of distinct genetic

clusters within the dataset, determined by mean log probability and DeltaK values (Figures 5,

S3). Individuals largely clustered according to their morphologically assigned species; however,

the number of detected clusters was fewer than the expected number of named taxonomic species

in the analysis (*n*=10). *P. integrifolia* and *P. amplifolia* are largely reproductively isolated, though

low levels of admixture were detected in *P. integrifolia* and only one sample was included for *P.*

amplifolia. Admixture is widespread amongst all other species, particularly *P. pleiantha*, *P.*

stipulata, *P. herthae*, *P. debilis, P. huantensis, P. brittoniana and P. antioquensis,* and like the

introgression analyses, is unrelated to evolutionary or ecological distance.

Estimates of genomic diversity

Within each species, the individuals with the highest observed heterozygosity (H_o) were

also the most admixed in the STRUCTURE results (Tables 3 and 4, Figure 4), and this was

consistent with different amounts of missing data. For example, in *P. pleiantha*, individuals 163,

164, and 165 showed high levels of admixture from multiple groups and had the highest H_o

estimates (50%: 0.18 – 0.22, 70%: 0.19 – 0.25; Table 3). Across species and filtering thresholds,

mean pairwise nucleotide diversity (π) was high (50%: 0.41 – 0.64, 70%: 0.37 – 0.51) compared

to H_o, π tracked with H_e as expected, and H_o (50%: 0.06 – 0.15, 70%: 0.06 – 0.17) was far below

H_e (50%: 0.41 – 0.64; 70%: 0.37 – 0.51; Table 2). However, the estimates for *P. herthae* and the

P. debilis+detrita group should be interpreted cautiously, given the low sample sizes for each.

These statistics corroborate the findings of the STRUCTURE analysis and further indicate a

history of interspecific gene flow among co-occurring *Prunus* species.

345   DISCUSSION

In this study, we aimed to investigate the patterns, timing, and extent of divergence

among partially sympatric congeneric trees in the tropical Andes. The phylogenomic data

presented here suggest that *Prunus* species known to co-occur as elevational replacements

diversified upslope from the Amazonian lowlands into the Andean highlands during the

Miocene, in a pattern consistent with adaptive diversification (lower montane lineage ~23.5 Ma,

cloud forest species ~20.4–13.4, high-elevation clade ~11.6–9.3 Ma). However, despite these

ancient diversification events, species do not appear to be reproductively isolated, since

substantial introgressive gene flow is detectable among species through time and is not

correlated with evolutionary or ecological distance. Together, these findings suggest that during

adaptive radiations, observable ecological and morphological differences may emerge despite

ongoing gene flow and long before reproductive isolation is complete.

Upslope diversification and biogeographic implications

Phylogenomic reconstructions and divergence time dating support the hypothesis that *Prunus*

diversified upslope, likely in response to novel habitat opportunities as has been observed in

other plant clades (e.g. Melastomataceae – Dudley 1978, *Espletia* – Pouchon et al. 2018, lupines

- Hughes and Eastwood 2006, Nevado et al. 2016, bellflowers – Lagomarsino et al. 2016,

Hypericums - Nürk et al. 2013, wild tomato – Pease et al. 2016, orchids - Pérez-Escobar et al.

2017, and bromeliads – Givnish et al. 2011, 2014, Silvestro et al. 2014, Males 2018). Though

this does not exclude the possibility that some species may have adapted elsewhere and

immigrated in once their preferred habitat became available, the predominant pattern

reconstructed from the phylogenomic analyses is one of stepwise, upslope adaptive

diversification. In agreement with prior studies, we found *P. myrtifolia* (endemic to the

Caribbean, Central America, and lowland South America) to be ancestral to the rest of the

Andean species included in this analysis, supporting the hypothesis that this lineage or its

ancestor colonized South America via migration through the Caribbean and Central America

(Chin et al. 2014; Hodel et al. 2023). While our divergence time estimate for *P. myrtifolia* of ~27

Ma is slightly later than the Neotropical colonization of ~34 Ma predicted by Chin et al. (2014),

it is still within the climatic shift from warmhouse to transitional state that coincided with an

onset of glaciation in Antarctica (~33.9 – 27.29 Ma; Judd et al. 2024) and the estimated end of

the boreotropical forests in the Northern Hemisphere.

In our analyses, diversification of the Andean species occurred throughout the Miocene,

beginning with the lineage that includes two of the lowest elevation species (*P. debilis* and *P.*

detrita) ~23.5 Ma and ending with the high-elevation clade ~11.6 Ma (*P. huantensis*), ~9.3 Ma

(*P. antioquensis* and *P. brittoniana*). This is consistent with the timing of major uplift events and

estimated paleo-elevations in the North-Central and Central Andes (Boschman 2021). It also

supports the findings of Hodel et al. (2023), which detected a long period of stasis followed by

rapid diversification of Neotropical *Prunus* during the Miocene. Although the exact timing, pace,

and sequence of Andean orogeny is a subject of ongoing study and is spatially heterogeneous, the

late Oligocene and Miocene (~25-5 Ma) are known to have been a period of intense and

accelerated uplift in the Eastern Cordillera of the North-Central and Central Andes (Boschman et

al. 2021; Pérez-Escobar et al. 2022).

The lower montane cloud forest species in our analysis (median elevation ranges 1200–2500

m; *P. pleiantha, P. integrifolia, P. herthae, P. stipulata*) diverged during the early to mid-

Miocene. This aligns with the predicted origination of cloud forests (Sempere et al. 2005) and the

estimated divergence times for other cloud forest lineages (Luebert and Weigend 2014). Andean

cloud forests are distinct habitats characterized by persistent cloud cover and immersion that

occurs from ~1200 m to the upper forest limits at ~3200–3800 m (depending on latitude).

Although they are the most speciose habitats in the Andes, persistence in these environments

often requires unique adaptations to cope with the reduced light availability, high precipitation,

and humidity such as thick, pachyphyllic leaves, low growth rates, and ability to withstand heavy

accumulations of epiphytes (Grubb 1977; Rapp and Silman 2012; Fahey et al. 2016; Pérez-

Escobar 2022). The highest elevation species in our study (median elevation ranges ≥3000 m; *P.*

huantensis, P. antioquensis, P. brittoniana) diverged during the late Miocene, as the final uplift

phase began in the North-Central and Central Andes (Boschman et al. 2021). Species in this

elevation zone are exposed to unique high-elevation stressors such as night frost, with

temperatures dipping below 0ºC, requiring further physiological specialization (Pérez-Escobar

2022).

Gene flow through time

Our genomic analyses show that Andean *Prunus* species remain porous lineages. Despite

deep Miocene divergences, we detect both ancient introgression and recent admixture among

partially sympatric taxa, with gene flow not obviously constrained by either evolutionary or

elevational distance. *D*-statistics detected at least five significant ancient introgression events.

Two of these events occurred among lineages separated by great evolutionary and ecological

distance (*P. pleiantha* × *P. huantensis*+ *antioquensis*, and *P. debilis* × *P. antioquensis).*

Introgression was also detected between *P. pleiantha* and the ancestor of *P. debilis*+ *detrita* (*P.*

pleiantha × *P. debilis*+ *detrita*) as well as among *P. pleiantha* × *P. debilis* and *P. pleiantha* × *P.*

detrita individually. Although difficult to disentangle, especially with limited sampling and

reduced-representation sequencing data, these patterns are most consistent with repeated

interspecific hybridization that occurred among the ancestors of these lineages and has continued

as elevationally segregated lineages diverged.

Historical introgression among species separated in ecological space across the elevation

gradient is plausible for several reasons. While the medians of their elevation ranges may be

quite different, they are found in close geographic proximity, and their elevational ranges overlap

(or are much closer) at the range margins. *Prunus* species are thought to be obligate outcrossers

with generalist pollinators, as is common in tropical trees (Ashton 1969; Bawa 1974, 1979). The

extent of self-incompatibility across the genus and potential for breakdown after polyploidization

are still unclear (Hauck et al. 2006, Tao et al. 2010, Aguiar et al. 2015). Generalist pollinators can

forage over kilometers, providing opportunity for introgression, especially given that many

tropical species flower semi-synchronously during the dry season (Bawa et al. 1985; Van Schaik

et al. 1993).

Even where known species ranges are separate at present, they have not necessarily

always been so. Pollen records from Lago Consuelo (1360 m a.s.l., located in southeastern Peru

adjacent to the Bolivian border) show that during the last full glacial period (43.5–22 kyr BP),

individual cloud forest species migrated downslope at least 1000–1700 vertical m in response to

lower cloud base and cooler temperatures (-5°C; Urrego et al. 2005). Today, these species ranges

are separated as far apart as they have likely been during the Quaternary, and historical

introgression events may be signatures of past ecological overlap.

Tests of recent admixture also found gene flow to be widespread among Andean *Prunus*

species. *P. integrifolia* and *P. amplifolia* are largely genomically isolated, but gene flow appears

rampant among all other species. The isolation of *P. integrifolia* is particularly interesting as this

species has the widest elevational range and is one of the most abundant species in the tropical

Andes. Gene flow was especially high among *P. pleiantha, P. stipulata*, and *P. herthae*, all cloud

forest species with largely overlapping elevation ranges. However, individuals of these species

mostly form monophyletic groups, indicating they have developed specific genetic

distinctiveness despite gene flow (Figure S1). As *P. pleiantha* was identified in most of the

historical introgression events we can infer that this species readily hybridizes when in contact

with close relatives. Further investigation will be needed to test this hypothesis, identify the

directionality of such gene flow, and determine whether physiological, ecological, or genetic

factors drive this weaker reproductive isolation. Ploidy variation may also underlie some of these

ambiguous genomic patterns, and an important next step will be to characterize ploidy across

these species and assess its influence on inferred gene flow and lineage distinctiveness.

STRUCTURE did not identify *P. debilis* and *P. detrita* as distinct genetic clusters, possibly due

to how closely related these species are, introgression, and the low number of individuals for

both species in the dataset. Our genomic data also support the most recent taxonomic revision by

Pérez-Zabala (2022) which considered *P. brittoniana* and *P. huantensis* to be the same species

given their morphological and ecological similarities (Figure S1). Taken together, these results

indicate that gene flow has been a persistent and complex feature of Andean *Prunus*

diversification, occurring across both evolutionary time and ecological space despite ancient

divergence times and emergence of species-level distinctiveness.

Conclusions

Overall, these results suggest that adaptive diversification following the emergence of

novel gradients has been important for within-genus diversification of Andean trees, particularly

in the genus *Prunus*, while immigration and sorting of pre-adapted clades may be the dominant

mechanism shaping community assembly at higher-level taxonomic levels (Segovia et al. 2020,

Griffiths et al. 2020, Linan et al. 2021). For most of the Neogene, *Prunus* diversified with the

rise of the Andes, developing sequential elevational replacements with narrow ranges, specially

adapted to live in radically different environmental conditions from the Amazonian lowlands to

Andean tree line. This has important implications in the context of climate change, as species are

now shifting their ranges to track their preferred temperatures (Feeley et al. 2011; Fadrique et al.

2018). Though most species will not be able to migrate fast enough to keep pace with climatic

shifts and are instead experiencing range retractions at the warmer-end, any upslope range shift

may increasingly result in range overlap among congeneric species and potential gene flow

(Duque et al. 2015; Farfan-Rios et al. 2025; Quinlan et al. 2025)

Here, we found evidence of both historical introgression, possibly in response to

Quaternary range shifts, and widespread admixture among co-occurring *Prunus* species. Yet

despite gene flow, these species have maintained distinct evolutionary lineages with identifiable

ecological and morphological differences since the Miocene. These results align with the

syngameon model, wherein species maintain incomplete reproductive barriers but primarily

experience divergent selection (Lotsy 1925; Grant 1971). Though oaks are the classic example

(e.g. Cannon and Petit 2020), such genomic mutualism may be an important yet under-

appreciated mechanism in the maintenance of diversity in hyper-diverse tropical forests, where

individuals predominantly outcross but retain some capacity for self-fertilization and

interspecific hybridization at low population densities (Wu 2001; Cannon and Lerdau 2015).

Further work is required to understand the prevalence of syngameons in tropical forests and their

role in species' responses to rapid environmental change, namely resilience and avoidance of

local extinction. Incorporating these complex dynamics into demographic models may be

important for accurately predicting the futures of Andean forests and effective conservation and

management under current climate change.

485   ACKNOWLEDGEMENTS

We thank Paul Fine, Michael Anderson, and Regina Joice Cordy for their helpful comments on

study design and earlier versions of this manuscript. We are also grateful to Herbario Vargas

(CUZ) and Herbario Nacional de Bolivia (LPB) for their assistance with coordination, field

work, and access to herbarium material, as well as Christopher Dick and Fairchild Botanical

Garden for their donation of samples to this project. Additional thanks to the Andes Biodiversity

Ecosystem Research Group (ABERG), and the many students and field technicians involved

with this project (especially Andrea Palomino Cardenas, Jesus Manuel Bañon Sanchez, Daire

Luciana Lafuente Palie, Paukar Alejandro Vedia Villca, and Felipe Aramburo Jaramillo). This

work was supported by NSF LTREB (1754647) and Garden Club of America's Tropical Botany

Fellowship. The material used in this study was collected under the research permits No 18-

2019-SERNANP-JEF, No 14-2022-SERNANP-JEF, No 527-2019-MINAGRI-SERFOR-

DGGSPFFS, and No D000170-2023-MIDAGRI-SERFOR-DGGSPFFS-DGSPF granted by

SERNANP and SERFOR.

499   COMPETING INTERESTS

500   None declared.

501   AUTHOR CONTRIBUTIONS

EJQ, MRS, and JBP initially conceived of the study. EJQ acquired funding, led research design,

data collection, data analysis and interpretation, and wrote the manuscript. JSB, AFF, and WFR

assisted with data collection and WFR additionally contributed to manuscript editing and

translated the summary. JBP assisted with data analysis and manuscript review and editing. MRS

provided supervision, acquired funding, and contributed to research design and manuscript

writing.

508   DATA AVAILABILITY

The genomic data generated for this study are available on NCBI's GenBank under submission

ID SUB15831409 and corresponding herbarium vouchers are deposited in Herbario Vargas

(CUZ) at the Universidad Nacional de San Antonio Abad del Cusco (UNSAAC) and Herbario

Nacional de Bolivia (LPB).

REFERENCES

Aboul-Maaty, N. A. F., & Oraby, H. A. S. (2019). Extraction of high-quality genomic DNA from
different plant orders applying a modified CTAB-based method. *Bulletin of the National*
Research Centre, *43*(1), 1-10.

Achá, S. (2013). El género Prunus (Rosaceae) en Bolivia: Novedades en su sistemática.

Aguiar, B., Vieira, J., Cunha, A. E., Fonseca, N. A., Iezzoni, A., van Nocker, S., & Vieira, C. P.
(2015). Convergent evolution at the gametophytic self-incompatibility system in Malus and
Prunus. *PloS one*, *10*(5), e0126138.

Anderson, E., & Stebbins Jr, G. L. (1954). Hybridization as an evolutionary stimulus. *Evolution*,
378-388.

Andro, M. C., & Riffaud, J. P. (1995). Pygeum africanum extract for the treatment of patients
with benign prostatic hyperplasia: a review of 25 years of published experience. *Current*
therapeutic research, *56*(8), 796-817.

Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of
recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

Baek, S., Choi, K., Kim, G. B., Yu, H. J., Cho, A., Jang, H., ... & Mun, J. H. (2018). Draft
genome sequence of wild Prunus yedoensis reveals massive inter-specific hybridization between
sympatric flowering cherries. *Genome biology*, *19*, 1-17.

Barton, N. H. (2001). The role of hybridization in evolution. *Molecular ecology*, *10*(3), 551-568.

Baskett, M. L., & Gomulkiewicz, R. (2011). Introgressive hybridization as a mechanism for
species rescue. *Theoretical Ecology*, *4*, 223-239.

Bawa, K. S. (1974). Breeding systems of tree species of a lowland tropical community.
Evolution, 85-92.

Bawa, K. S. (1979). Breeding systems of trees in a tropical wet forest. *New Zealand Journal of*
Botany, *17*(4), 521-524.

Bawa, K. S., Perry, D. R., & Beach, J. H. (1985). Reproductive biology of tropical lowland rain
forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of*
botany, *72*(3), 331-345.

Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H. (1985). Reproductive
biology of tropical lowland rain forest trees. II. Pollination systems. *American journal of*
botany, *72*(3), 346-356.

Boschman, L. M. (2021). Andean mountain building since the Late Cretaceous: A paleoelevation
reconstruction. *Earth-Science Reviews*, *220*, 103640.

Bortiri, E., Oh, S. H., Jiang, J., Baggett, S., Granger, A., Weeks, C., ... & Parfitt, D. E. (2001).
Phylogeny and systematics of Prunus (Rosaceae) as determined by sequence analysis of ITS and
the chloroplast trnL-trnF spacer DNA. *Systematic Botany*, *26*(4), 797-807.

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M., Gavryushkina, A.,
... & Drummond, A. J. (2019). BEAST 2.5: An advanced software platform for Bayesian
evolutionary analysis. *PLoS computational biology*, *15*(4), e1006650.

Brauer, C. J., Sandoval-Castillo, J., Gates, K., Hammer, M. P., Unmack, P. J., Bernatchez, L., &
Beheregaray, L. B. (2023). Natural hybridization reduces vulnerability to climate change. *Nature*
Climate Change, *13*(3), 282-289.

Cannon, C. H., & Lerdau, M. (2015). Variable mating behaviors and the maintenance of tropical
biodiversity. *Frontiers in Genetics*, *6*, 183.

Cannon, C. H., & Petit, R. J. (2020). The oak syngameon: more than the sum of its parts. *New*
Phytologist, *226*(4), 978-983.

Cavender-Bares, J., Kothari, S., Meireles, J. E., Kaproth, M. A., Manos, P. S., & Hipp, A. L.
(2018). The role of diversification in community assembly of the oaks (Quercus L.) across the
continental US. *American Journal of Botany*, *105*(3), 565-586.

Chakraborty, R., & Nei, M. (1977). Bottleneck effects on average heterozygosity and genetic
distance with the stepwise mutation model. *Evolution*, 347-356.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology*
and Systematics, *31*(1), 343-366.

Chhatre, V. E., Evans, L. M., DiFazio, S. P., & Keller, S. R. (2018). Adaptive introgression and
maintenance of a trispecies hybrid complex in range-edge populations of Populus. *Molecular*
ecology, *27*(23), 4820-4838.

Chin, S. W., Shaw, J., Haberle, R., Wen, J., & Potter, D. (2014). Diversification of almonds,
peaches, plums and cherries–molecular systematics and biogeographic history of Prunus
(Rosaceae). Molecular phylogenetics and evolution, 76, 34-48.

Cici, S. Z. H., & Van Acker, R. C. (2010). Gene flow in Prunus species in the context of novel
trait risk assessment. *Environmental biosafety research*, *9*(2), 75-85.

Comeault, A. A., & Matute, D. R. (2018). Genetic divergence and the number of hybridizing
species affect the path to homoploid hybrid speciation. *Proceedings of the National Academy of*
Sciences, *115*(39), 9761-9766.

Danecek, Petr, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A.
DePristo, Robert E. Handsaker et al. "The variant call format and VCFtools." *Bioinformatics* 27,
no. 15 (2011): 2156-2158.

De La Torre, A. R., Roberts, D. R., & Aitken, S. N. (2014). Genome-wide admixture and ecological niche modelling reveal the maintenance of species boundaries despite long history of interspecific gene flow. Molecular Ecology, 23(8), 2046-2059.

Donoghue, M. J. (2008). A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences*, *105*(supplement_1), 11549-11555.

Donoghue, M. J., Eaton, D. A., Maya-Lastra, C. A., Landis, M. J., Sweeney, P. W., Olson, M. E., ... & Edwards, E. J. (2022). Replicated radiation of a plant clade along a cloud forest archipelago. *Nature Ecology & Evolution*, *6*(9), 1318-1329.

Duque, A., Stevenson, P. R., & Feeley, K. J. (2015). Thermophilization of adult and juvenile tree communities in the northern tropical Andes. *Proceedings of the National Academy of Sciences*, *112*(34), 10744-10749.

Dudley, E. C. (1978). Adaptive radiation in the Melastomataceae along an altitudinal gradient in Peru. *Biotropica*, 134-143.

Durand, E. Y., Patterson, N., Reich, D., & Slatkin, M. (2011). Testing for ancient admixture between closely related populations. *Molecular biology and evolution*, *28*(8), 2239-2252.

Eaton, D. A., & Overcast, I. (2020). ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics*, *36*(8), 2592-2594.

Ehrendorfer, F. (1970). Evolutionary patterns and strategies in seed plants. *Taxon*, *19*(2), 185-195.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular ecology*, *14*(8), 2611-2620.

Fahey, T. J., Sherman, R. E., & Tanner, E. V. (2016). Tropical montane cloud forest: environmental drivers of vegetation structure and ecosystem function. *Journal of Tropical Ecology*, *32*(5), 355-367.

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., ... & Feeley, K. J. (2018). Widespread but heterogeneous responses of Andean forests to climate change. *Nature*, *564*(7735), 207-212.

Farfan-Rios, W. (2019). *Forest responses to climate change along an Andes-to-Amazon elevational gradient*. Wake Forest University.

Farfan-Rios, W., Feeley, K. J., Myers, J. A., Tello, S., Sallo-Bravo, J., Malhi, Y., ... & Silman, M. R. (2025). Amazonian and Andean tree communities are not tracking current climate warming. Proceedings of the National Academy of Sciences, 122(34), e2425619122.

Feeley, K. J., Silman, M. R., Bush, M. B., Farfan, W., Cabrera, K. G., Malhi, Y., ... & Saatchi, S. (2011). Upslope migration of Andean trees. *Journal of Biogeography*, *38*(4), 783-791.

Freeman, B. G., Strimas-Mackey, M., & Miller, E. T. (2022). Interspecific competition limits bird species' ranges in tropical mountains. *Science*, *377*(6604), 416-420.

Funk, V. A., Gostel, M., Devine, A., Kelloff, C. L., Wurdack, K., Tuccinardi, C., ... & Coddington, J. (2017). Guidelines for collecting vouchers and tissues intended for genomic work (Smithsonian Institution): Botany Best Practices. *Biodiversity Data Journal*, (5), e11625.

Gailing, O., & Curtu, A. L. (2014). Interspecific gene flow and maintenance of species integrity in oaks. *Annals of Forest Research*, *57*(1), 5-18.

Gentry, A. H. (1982). Neotropical floristic diversity: phytogeographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny?. *Annals of the Missouri Botanical Garden*, *69*(3), 557-593.

Givnish, T. J., Barfuss, M. H., Van Ee, B., Riina, R., Schulte, K., Horres, R., ... & Sytsma, K. J. (2011). Phylogeny, adaptive radiation, and historical biogeography in Bromeliaceae: insights from an eight-locus plastid phylogeny. *American journal of botany*, *98*(5), 872-895.

Grant, V. (1971). Plant Speciation. New York: Columbia University Press.

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., ... & Pääbo, S. (2010). A draft sequence of the Neandertal genome. science, 328(5979), 710-722.

Griffiths, A. R., Silman, M. R., Farfan-Rios, W., Feeley, K. J., Cabrera, K. G., Meir, P., ... & Dexter, K. G. (2021). Evolutionary diversity peaks at mid-elevations along an Amazon-to-Andes elevation gradient. *Frontiers in ecology and evolution*, *9*, 680041.

Grubb, P. J. (1977). Control of forest growth and distribution on wet tropical mountains: with special reference to mineral nutrition. *Annual Review of Ecology and Systematics*, 83-107.

Guerrero, B. I., Guerra, M. E., & Rodrigo, J. (2022). Simple sequence repeat (SSR)-based genetic diversity in interspecific Plumcot-type (Prunus salicina× Prunus armeniaca) hybrids. *Plants*, *11*(9), 1241.

Hamrick, J. L., & Murawski, D. A. (1990). The breeding structure of tropical tree populations. *Plant species biology*, *5*(1), 157-165.

Hauck, N. R., Yamane, H., Tao, R., & Iezzoni, A. F. (2006). Accumulation of nonfunctional S-haplotypes results in the breakdown of gametophytic self-incompatibility in tetraploid Prunus. *Genetics*, *172*(2), 1191-1198.

Hibbins, M. S., & Hahn, M. W. (2022). Phylogenomic approaches to detecting and characterizing introgression. *Genetics*, *220*(2), iyab173.

Hillyer, R., & Silman, M. R. (2010). Changes in species interactions across a 2.5 km elevation gradient: effects on plant migration in response to climate change. *Global Change Biology*, *16*(12), 3205-3214.

Hodel, R. G., Zimmer, E., & Wen, J. (2021). A phylogenomic approach resolves the backbone of Prunus (Rosaceae) and identifies signals of hybridization and allopolyploidy. *Molecular Phylogenetics and Evolution*, *160*, 107118.

Hodel, R. G., Winslow, S. K., Liu, B. B., Johnson, G., Trizna, M., White, A. E., ... & Wen, J. (2023). A phylogenomic approach, combined with morphological characters gleaned via machine learning, uncovers the hybrid origin and biogeographic diversification of the plum genus. *bioRxiv*, 2023-09.

Hubbell, S. P. (2005). Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional ecology*, *19*(1), 166-172.

Hughes, C., & Eastwood, R. (2006). Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the Andes. *Proceedings of the National Academy of Sciences*, *103*(27), 10334-10339.

International Peach Genome Initiative, Verde, I., Abbott, A. G., Scalabrin, S., Jung, S., Shu, S., ... & Rokhsar, D. S. (2013). The high-quality draft genome of peach (Prunus persica) identifies unique patterns of genetic diversity, domestication, and genome evolution. *Nature genetics*, *45*(5), 487-494.

Janzen, D. H. (1967). Why mountain passes are higher in the tropics. *The American Naturalist*, *101*(919), 233-249.

Jankowski, J. E., Ciecka, A. L., Meyer, N. Y., & Rabenold, K. N. (2009). Beta diversity along environmental gradients: implications of habitat specialization in tropical montane landscapes. *Journal of Animal ecology*, *78*(2), 315-327.

Jankowski, J. E., Graham, C. H., Parra, J. L., Robinson, S. K., Seddon, N., Touchton, J. M., & Tobias, J. A. (2012). The role of competition in structuring tropical bird communities. *Ornitologia Neotropical*, *23*, 115-124.

Judd, E. J., Tierney, J. E., Lunt, D. J., Montañez, I. P., Huber, B. T., Wing, S. L., & Valdes, P. J. (2024). A 485-million-year history of Earth's surface temperature. *science*, *385*(6715), eadk3705.

Kulmuni, J., Wiley, B., & Otto, S. P. (2024). On the fast track: hybrids adapt more rapidly than parental populations in a novel environment. *Evolution Letters*, *8*(1), 128-136.

Lagomarsino, L. P., Condamine, F. L., Antonelli, A., Mulch, A., & Davis, C. C. (2016). The abiotic and biotic drivers of rapid diversification in A ndean bellflowers (Campanulaceae). *New Phytologist*, *210*(4), 1430-1442.

Larson, D. A., Vargas, O. M., Vicentini, A., & Dick, C. W. (2021). Admixture may be extensive among hyperdominant Amazon rainforest tree species. *New Phytologist*, *232*(6), 2520-2534.

Lee, S., & Wen, J. (2001). A phylogenetic analysis of Prunus and the Amygdaloideae (Rosaceae) using ITS sequences of nuclear ribosomal DNA. *American Journal of Botany*, *88*(1), 150-160.

Lepais, O., Petit, R. J., Guichoux, E., Lavabre, J. E., Alberto, F., Kremer, A., & Gerber, S. (2009). Species relative abundance and direction of introgression in oaks. *Molecular Ecology*, *18*(10), 2228-2242.

Levin, D. A. (2012). Mating system shifts on the trailing edge. *Annals of Botany*, *109*(3), 613-620.

Linan, A. G., Myers, J. A., Edwards, C. E., Zanne, A. E., Smith, S. A., Arellano, G., ... & Tello, J. S. (2021). The evolutionary assembly of forest communities along environmental gradients: recent diversification or sorting of pre-adapted clades?. *New Phytologist*, *232*(6), 2506-2519.

Lotsy, J. P. (1925). *Evolution considered in the light of hybridization*. Canterbury College.

Luebert, F., & Weigend, M. (2014). Phylogenetic insights into Andean plant diversification. *Frontiers in Ecology and Evolution*, *2*, 27.

McCartney-Melstad, E., Gidiş, M., & Shaffer, H. B. (2019). An empirical pipeline for choosing the optimal clustering threshold in RADseq studies. *Molecular Ecology Resources*, *19*(5), 1195-1204.

Meier, C. L., Rapp, J., Bowers, R. M., Silman, M., & Fierer, N. (2010). Fungal growth on a common wood substrate across a tropical elevation gradient: temperature sensitivity, community composition, and potential for above-ground decomposition. *Soil Biology and Biochemistry*, *42*(7), 1083-1090.

Mendelson, T. C., Inouye, B. D., & Rausher, M. D. (2004). Quantifying patterns in the evolution of reproductive isolation. *Evolution*, *58*(7), 1424-1433.

Menon, M., Bagley, J. C., Friedline, C. J., Whipple, A. V., Schoettle, A. W., Leal-Sàenz, A., ... & Eckert, A. J. (2018). The role of hybridization during ecological divergence of southwestern white pine (Pinus strobiformis) and limber pine (P. flexilis). *Molecular Ecology*, *27*(5), 1245-1260.

Murawski, D. A., & Hamrick, J. L. (1991). The effect of the density of flowering individuals on the mating systems of nine tropical tree species. *Heredity*, *67*(2), 167-174.

Nason, J. D., & Hamrick, J. L. (1997). Reproductive and genetic consequences of forest fragmentation: Two case studies of neotropical... *Journal of Heredity*, *88*(4).

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The bottleneck effect and genetic variability in populations. *Evolution*, 1-10.

Nevado, B., Atchison, G. W., Hughes, C. E., & Filatov, D. A. (2016). Widespread adaptive evolution during repeated evolutionary radiations in New World lupins. *Nature communications*, *7*(1), 12384.

Nürk, N. M., Madriñán, S., Carine, M. A., Chase, M. W., & Blattner, F. R. (2013). Molecular phylogenetics and morphological evolution of St. John's wort (Hypericum; Hypericaceae). *Molecular Phylogenetics and Evolution*, *66*(1), 1-16.

Ottenburghs, J. (2020). Ghost introgression: spooky gene flow in the distant past. *Bioessays*, *42*(6), 2000012.

Pease, J. B., Haak, D. C., Hahn, M. W., & Moyle, L. C. (2016). Phylogenomics reveals three sources of adaptive variation during a rapid radiation. PLoS biology, 14(2), e1002379.

Pease, J. B., Brown, J. W., Walker, J. F., Hinchliff, C. E., & Smith, S. A. (2018). Quartet Sampling distinguishes lack of support from conflicting support in the green plant tree of life. *American journal of botany*, *105*(3), 385-403.

Pérez-Escobar, O. A., Chomicki, G., Condamine, F. L., Karremans, A. P., Bogarín, D., Matzke, N. J., ... & Antonelli, A. (2017). Recent origin and rapid speciation of Neotropical orchids in the world's richest plant biodiversity hotspot. *New Phytologist*, *215*(2), 891-905.

Pérez-Escobar, O. A., Zizka, A., Bermúdez, M. A., Meseguer, A. S., Condamine, F. L., Hoorn, C., ... & Chomicki, G. (2022). The Andes through time: evolution and distribution of Andean floras. *Trends in Plant Science*, *27*(4), 364-378.

Perez Zabala, J. A. (2022). *Taxonomic Diversity, Phylogeny, and Diversification of the Environmental Niche of the Genus Prunus L. with emphasis on the New World Tropics* (Doctoral dissertation, UC Davis).

Pires, J. M., Dobzhansky, T. H., & Black, G. A. (1953). An estimate of the number of species of trees in an Amazonian forest community. *Botanical Gazette*, *114*(4), 467-477.

Pitman, N. C., Terborgh, J., Silman, M. R., & Nuñez V, P. (1999). Tree species distributions in an upper Amazonian forest. *Ecology*, *80*(8), 2651-2661.

Pouchon, C., Fernández, A., Nassar, J. M., Boyer, F., Aubert, S., Lavergne, S., & Mavárez, J. (2018). Phylogenomic analysis of the explosive adaptive radiation of the Espeletia complex (Asteraceae) in the tropical Andes. *Systematic biology*, *67*(6), 1041-1060.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945-959.

Quinlan, E. J., Layman, C. A., & Silman, M. R. (2025). Climate-Mediated Hybridisation and the Future of Andean Forests. *Journal of Biogeography*, *52*(6), e15113.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior
summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic biology*, *67*(5), 901-904.

Rapp, J. M., & Silman, M. R. (2012). Diurnal, seasonal, and altitudinal trends in microclimate
across a tropical montane cloud forest. *Climate Research*, *55*(1), 17-32.

Rehder A (1940) Manual of cultivated trees and shrubs hardy in North America exclusive of the
subtropical and warmer temperate regions. Dioscorides, Portland, OR.

Schley, R. J., Pennington, R. T., Pérez-Escobar, O. A., Helmstetter, A. J., de la Estrella, M.,
Larridon, I., ... & Klitgård, B. (2020). Introgression across evolutionary scales suggests
reticulation contributes to Amazonian tree diversity. *Molecular Ecology*, *29*(21), 4170-4185.

Schley, R. J., Twyford, A. D., & Pennington, R. T. (2022). Hybridization: a 'double-edged
sword' for Neotropical plant diversity. *Botanical Journal of the Linnean Society*, *199*(1), 331-
356.

Schley, R. J., Piñeiro, R., Nicholls, J. A., Pezzini, F. F., Farbos, A., Lewis, G. P., ... & Pennington,
R. T. (2025). Rampant Reticulation in a Rapid Radiation of Tropical Trees-Insights from Inga
(Fabaceae). *Systematic biology*, syaf027.

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *the american
naturalist*, *156*(S4), S4-S16.

Segovia, R. A., Pennington, R. T., Baker, T. R., Coelho de Souza, F., Neves, D. M., Davis, C. C.,
... & Dexter, K. G. (2020). Freezing and water availability structure the evolutionary diversity of
trees across the Americas. *Science Advances*, *6*(19), eaaz5373.

Sempere, T., Picard, D., & Plantard, O. (2005). Assessing and dating Andean uplift by
phylogeography and phylochronology: early Miocene emergence of Andean cloud forests. In *6th
International Symposium on Andean Geodynamics (ISAG 2005, Barcelona), Extended
Abstracts* (pp. 663-665). IRD Éditions.

Silvestro, D., Schnitzler, J., Liow, L. H., Antonelli, A., & Salamin, N. (2014). Bayesian
estimation of speciation and extinction from incomplete fossil occurrence data. *Systematic
biology*, *63*(3), 349-367.

Simpson, G. G. (1953). *The major features of evolution*. Columbia University Press.

Sokoloff, D. D., Ignatov, M. S., Remizowa, M. V., Nuraliev, M. S., Blagoderov, V., Garbout, A.,
& Perkovsky, E. E. (2018). Staminate flower of Prunus s. l.(Rosaceae) from Eocene Rovno
amber (Ukraine). *Journal of Plant Research*, *131*, 925-943.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.

Stroud, J. T., & Losos, J. B. (2016). Ecological opportunity and adaptive radiation. *Annual Review of Ecology, Evolution, and Systematics*, *47*(1), 507-532.

Stull, G. W., Pham, K. K., Soltis, P. S., & Soltis, D. E. (2023). Deep reticulation: the long legacy of hybridization in vascular plant evolution. *The Plant Journal*, *114*(4), 743-766.

Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. (2018). Adaptive introgression: a plant perspective. *Biology letters*, *14*(3), 20170688.

Su, N., Hodel, R. G., Wang, X., Wang, J. R., Xie, S. Y., Gui, C. X., ... & Wen, J. (2023). Molecular phylogeny and inflorescence evolution of Prunus (Rosaceae) based on RAD-seq and genome skimming analyses. *Plant diversity*, *45*(4), 397-408.

Tao, R., & Iezzoni, A. F. (2010). The S-RNase-based gametophytic self-incompatibility system in Prunus exhibits distinct genetic and molecular features. *Scientia Horticulturae*, *124*(4), 423-433.

Ter Steege, H., Pitman, N. C., Sabatier, D., Baraloto, C., Salomão, R. P., Guevara, J. E., ... & Silman, M. R. (2013). Hyperdominance in the Amazonian tree flora. *Science*, *342*(6156), 1243092.

Terborgh, J. (1971). Distribution on environmental gradients: theory and a preliminary interpretation of distributional patterns in the avifauna of the Cordillera Vilcabamba, Peru. *Ecology*, *52*(1), 23-40.

Terborgh, J. (1977). Bird species diversity on an Andean elevational gradient. *Ecology*, *58*(5), 1007-1019.

Tricou, T., Tannier, E., & de Vienne, D. M. (2022). Ghost lineages highly influence the interpretation of introgression tests. *Systematic Biology*, *71*(5), 1147-1158.

Tsuda, Y., Semerikov, V., Sebastiani, F., Vendramin, G. G., & Lascoux, M. (2017). Multispecies genetic structure and hybridization in the Betula genus across Eurasia. *Molecular Ecology*, *26*(2), 589-605.

Urrego, D. H., Silman, M. R., & Bush, M. B. (2005). The Last Glacial Maximum: stability and change in a western Amazonian cloud forest. *Journal of Quaternary Science: Published for the Quaternary Research Association*, *20*(7-8), 693-701.

Van Schaik, C. P., Terborgh, J. W., & Wright, S. J. (1993). The phenology of tropical forests: adaptive significance and consequences for primary consumers. *Annual Review of ecology and Systematics*, 353-377.

Verde, I., Jenkins, J., Dondini, L., Micali, S., Pagliarani, G., Vendramin, E., ... & Schmutz, J. (2017). The Peach v2. 0 release: high-resolution linkage mapping and deep resequencing improve chromosome-scale assembly and contiguity. *BMC genomics*, *18*, 1-18.

Ward, M., Dick, C. W., Gribel, R., & Lowe, A. J. (2005). To self, or not to self… A review of
outcrossing and pollen-mediated gene flow in neotropical trees. Heredity, 95(4), 246-254.

Wen, J., Berggren, S. T., Lee, C. H., Ickert-Bond, S., Yi, T. S., Yoo, K. O., ... & Potter, D. (2008).
Phylogenetic inferences in Prunus (Rosaceae) using chloroplast ndhF and nuclear ribosomal ITS
sequences.

Williams Jr, J. H., Friedman, W. E., & Arnold, M. L. (1999). Developmental selection within the
angiosperm style: using gamete DNA to visualize interspecific pollen competition. *Proceedings*
of the National Academy of Sciences, *96*(16), 9201-9206.

Wu, C. I. (2001). The genic view of the process of speciation. *Journal of evolutionary*
biology, *14*(6), 851-865.

Zhao, L., Jiang, X. W., Zuo, Y. J., Liu, X. L., Chin, S. W., Haberle, R., ... & Wen, J. (2016).
Multiple events of allopolyploidy in the evolution of the racemose lineages in Prunus (Rosaceae)
based on integrated evidence from nuclear and plastid data. *PLoS One*, *11*(6), e0157123.

Table 1. Summary of significant D-statistic tests, with *P. myrtifolia* set to P4.

P1	P2	P3	dstat	Z	ABBA	BABA	nloci
P. antioquensis	*P. stipulata*	*P. debilis*	-0.69	4.75	8.38	45.13	152
P. stipulata	*P. antioquensis*	*P. debilis*	0.69	4.60	45.13	8.38	152
P. amplifolia, P. stipulata	*P. antioquensis*	*P. debilis*	0.66	4.16	43.19	8.94	163
P. antioquensis	*P. amplifolia, P. stipulata*	*P. debilis*	-0.66	4.01	8.94	43.19	163
P. stipulata	*P. pleiantha*	*P. detrita, P. debilis*	0.13	2.94	793	605.44	3968
P. pleiantha	*P. stipulata*	*P. detrita, P. debilis*	-0.13	2.87	605.44	793	3968
P. herthae, P. huantensis	*P. pleiantha*	*P. detrita, P. debilis*	0.11	2.81	864.08	686.51	4485
P. stipulata	*P. huantensis, P. antioquensis*	*P. pleiantha*	0.09	2.71	1190.20	989.74	6218
P. huantensis, P. antioquensis	*P. stipulata*	*P. pleiantha*	-0.09	2.71	989.74	1190.2	6218
P. pleiantha	*P. herthae, P. huantensis*	*P. detrita, P. debilis*	-0.11	2.67	686.51	864.08	4485
P. stipulata	*P. pleiantha*	*P. detrita*	0.13	2.66	744.38	578.5	3725
P. pleiantha	*P. antioquensis*	*P. debilis*	0.51	2.63	20.25	6.5	142
P. pleiantha	*P. stipulata*	*P. detrita*	-0.13	2.62	578.5	744.38	3725
P. antioquensis	*P. pleiantha*	*P. debilis*	-0.51	2.58	6.5	20.25	142
P. pleiantha	*P. stipulata*	*P. debilis*	-0.27	2.53	68.5	119.25	576
P. herthae, P. huantensis, P. antioquensis, P. brittoniana	*P. pleiantha*	*P. detrita*	0.11	2.52	813.45	654.78	4203

830   Table 2. Summary of within species genomic diversity statistics.

Species	# of samples	Filter	# of loci	Average π	Average H_o	Average H_e
P. integrifolia	10	50	53574	0.42 ± 0.13	0.15 ± 0.064	0.42 ± 0.0025
		70	17592	0.41 ± 0.14	0.17 ± 0.073	0.40 ± 0.0015
P. huantensis/ brittoniana	12	50	24457	0.41 ± 0.13	0.12 ± 0.078	0.41 ± 0.0012
		70	1960	0.38 ± 0.15	0.16 ± 0.10	0.37 ± 0.0031
P. stipulata	7	50	24457	0.45 ± 0.11	0.095 ± 0.096	0.45 ± 0.0024
				0.42 ± 0.13	0.11 ± 0.11	0.42 ± 0.0064
P. pleiantha	7	50	23299	0.45 ± 0.11	0.12 ± 0.081	0.44 ± 0.0015
		70	7295	0.43 ± 0.12	0.13 ± 0.091	0.42 ± 0.0013
P. herthae	3	50	6251	0.62 ± 0.077	0.085 ± 0.03	0.61 ± 0.0067
		70	699	0.51 ± 0.073	0.064 ± 0.0072	0.51 ± 0
P. debilis/detrita	3	50	1747	0.64 ± 0.061	0.064 ± 0.058	0.064 ± 0.0009

832   Table 3. Summary of within-individual heterozygosity estimates.

Species	Individual	H_o (50% filter)	H_e (50% filter)	H_o (70% filter)	H_e (70% filter)
P. integrifolia	139_integrifolia	0.11	0.42	0.12	0.41
	147_integrifolia	0.31	0.42	0.36	0.4
	148_integrifolia	0.14	0.42	0.16	0.4
	149_integrifolia	0.18	0.42	0.21	0.4
	151_integrifolia	0.10	0.42	0.12	0.4
	155_integrifolia	0.16	0.42	0.19	0.4
	156_integrifolia	0.12	0.42	0.14	0.4
	157_integrifolia	0.17	0.42	0.2	0.4
	212_integrifolia	0.09	0.41	0.1	0.4
	253_integrifolia	0.12	0.42	0.14	0.41
P. huantensis /brittoniana	108_huantensis	0.16	0.41	0.23	0.37
	115_huantensis	0.1	0.41	0.13	0.38
	116_huantensis	0.22	0.41	0.29	0.37
	117_huantensis	0.044	0.4	0.07	0.37
	119_huantensis	0.072	0.41	0.093	0.37
	123_huantensis	0.069	0.4	0.12	0.37
	124_huantensis	0.2	0.4	0.25	0.37
	130_huantensis	0.05	0.41	0.079	0.38
	131_huantensis	0.15	0.41	0.22	0.37
	132_huantensis	0.26	0.41	0.36	0.37
	179_brittoniana	0.064	0.41	0.08	0.37
	210_brittoniana	0.03	0.4	0.05	0.38
P. stipulata	171_stipulata	0.28	0.45	0.32	0.42
	173_stipulata	0.17	0.45	0.18	0.42
	175_stipulata	0.07	0.45	0.07	0.41
	285_stipulata	0.058	0.45	0.1	0.42
	286_stipulata	0.03	0.45	0.041	0.41
	312_stipulata	0.037	0.45	0.032	0.42
	332_stipulata	0.02	0.45	0.021	0.43
P. pleiantha	162_pleiantha	0.12	0.44	0.13	0.43
	163_pleiantha	0.22	0.44	0.25	0.43
	164_pleiantha	0.18	0.44	0.19	0.42
	165_pleiantha	0.2	0.44	0.22	0.42
	166_pleiantha	0.023	0.44	0.021	0.42
	167_pleiantha	0.05	0.44	0.052	0.42
	169_pleiantha	0.051	0.44	0.056	0.42
P. herthae	226_herthae	0.087	0.62	0.057	0.51
	236_herthae	0.054	0.61	0.063	0.51
	237_herthae	0.11	0.62	0.072	0.51
P. debilis/ detrita	180_debilis	0.13	0.64	NA	NA
	225_debilis	0.036	0.64	NA	NA
	107_detrita	0.026	0.64	NA	NA

Figure 1. (a) Map of samples included in this study. Black dot indicates the location of the
ABERG transect in Manu National Park, Peru where most samples (n=33) were collected, also
shown in picture to the right. Red points indicate locations of additional samples included in the
study. Further details on individual samples provided in Table S1. (b) Box plots showing
elevational distributions of the in-group Andean species across three different sources. "Aberg"
includes all known individuals along the ABERG transect >10 dbh over 20 years. "Acha"
includes all Bolivian material reviewed by Achá (2013) and "Zabala" includes all material

reviewed by Perez-Zabala (2022) from Venezuela to Bolivia. Perez-Zabala considered *P.*
huantensis to be *P. brittoniana* so those samples are grouped together on the figure under *P.*
brittoniana. Figure does not include new "varieties" of species proposed by either Achá or Perez-
Zabala.

Figure 2. Species-level topology resulting from RAxML and ExaBayes phylogenetic analyses.
Nodes show RAxML bootstrap support values where black circles indicate values > 75 and
values < 75 are shown on white nodes. All Bayesian posterior probabilities were equal to 1. The
tree is rooted to the reference (*P. persica*) and all three outgroup species are shown in grey.
Dashed arrows summarize patterns of historical admixture inferred from D-statistics.

Figure 3. Time-calibrated phylogeny generated from BEAST analysis. Posterior probabilities are shown for nodes when < 1. Node
bars represent the 95% highest probability density (HPD) interval for node ages. Red stars indicate the fossil calibration points.

Figure 4. Patterns of genetic structure among individuals of sympatric species. Each vertical bar
represents a single individual and the y-axis indicates the proportion of genetic ancestry of that
individual assigned to different inferred genetic clusters, designated with different colors.
STRUCTURE detected K=9 distinct genetic clusters as the most likely grouping (Figure S3).

Table S1. Summary of samples included in individual-level phylogeny and genetic structure
analyses. Samples in bold were used to construct the species-level phylogeny.
865

Species determination	Sample ID	Voucher (Herbarium)	Material	Sampling locality
P. detrita	**107_detrita**	**EJQ82 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. debilis	**180_debilis**	**Loza et al. 388 (LPB)**	**Herbarium**	**Madidi NP, Bolivia**
P. debilis	225_debilis	JSB6352A (CUZ)	Fresh	Manu NP, Peru
P. stipulata	**171_stipulata**	**EJQ114 (LPB)**	**Fresh**	**Cotapata, Bolivia**
P. stipulata	173_stipulata	EJQ116 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	175_stipulata	EJQ118 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	285_stipulata	JSB6320 (CUZ)	Fresh	Manu NP, Peru
P. stipulata	286_stipulata	JSB6321(CUZ)	Fresh	Manu, NP, Peru
P. stipulata	312_stipulata	Cayola 3259	Herbarium	La Paz, Bolivia
P. stipulata	332_stipulata	Villalobosos 81	Herbarium	
P. integrifolia	139_integrifolia	JSB6366 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	**147_integrifolia**	**JSB6402 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. integrifolia	148_integrifolia	JSB6396 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	149_integrifolia	JSB6403 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	151_integrifolia	JSB6400 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	155_integrifolia	JSB6397 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	156_integrifolia	JSB6391 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	157_integrifolia	JSB6392 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	212_integrifolia	Meneses 7535 (LPB)	Herbarium	Nor Yungas, Cotapata, Bolivia
P. integrifolia	253_integrifolia	JSB7255 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	162_pleiantha	JSB6382 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	**163_pleiantha**	**JSB6388 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. pleiantha	164_pleiantha	JSB6386 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	165_pleiantha	JSB6387 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	166_pleiantha	JSB6385 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	167_pleiantha	JSB6380 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	169_pleiantha	JSB6390 (CUZ)	Fresh	Manu NP, Peru
P. herthae	226_herthae	JSB6343 (CUZ)	Fresh	Manu NP, Peru
P. herthae	236_herthae	JSB6333 (CUZ)	Fresh	Manu NP, Peru
P. herthae	**237_herthae**	**JSB6334 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. amplifolia	**206_amplifolia**	**Beck 33550 (LPB)**	**Herbarium**	**La Paz, Bolivia**
P. antioquensis	**177_antioquensis**	**Araujo & Canqui 348 (LPB)**	**Herbarium**	**Madidi, Bolivia**
P. huantensis	108_huantensis	JSB6316 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	115_huantensis	JSB6310 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	**116_huantensis**	**JSB6358 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. huantensis	117_huantensis	JSB6360 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	119_huantensis	JSB6362 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	123_huantensis	JSB6353 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	124_huantensis	JSB6357 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	130_huantensis	JSB6375 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	131_huantensis	JSB6311 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	132_huantensis	JSB6312 (CUZ)	Fresh	Manu NP, Peru

P. brittoniana	**179_brittoniana**	**Paniagua et al. 5782A (LPB)**	**Herbarium**	**Pelechuco-Mojo, Bolivia**
P. brittoniana	210_brittoniana	Beck 35044 (LPB)	Herbarium	Sud Yungas, Cotapata, Bolivia
Outgroups:				
P. myrtifolia	**1_myrtifolia**	**Silica sample only**	**Fresh**	**Fairchild Botanical Garden, Miami, Florida, USA**
P. caroliniana	**2_caroliniana**	**Silica sample only**	**Fresh**	**Weymouth Woods, North Carolina, USA**

Figure S1. Maximum likelihood individual-level phylogeny reconstructed in RAxML.
Individuals were selected to include those with at least 1000 loci per sample and up to 10
individuals per species. Sites were filtered to only include those with at least 25% coverage
across samples. Tree was rooted to the reference (*P. persica*) and nodes show bootstrap support
value.

Figure S2. Quartet sampling results with quartet concordance (QC)/quartet differential
(QD)/quartet informativeness (QI) indicated next to nodes. Nodes are colored according to QC
scores, indicating how often the concordant quartet is inferred over both discordant quartets.
Outgroup samples *P. persica* and *P. caroliniana* are removed for clarity. Figure created with
https://github.com/ShuiyinLIU/QS_visualization.

Figure S3. Plot of estimators used to select the optimal number of genetic clusters (K) in the
STRUCTURE analysis. Estimated log mean probability assesses how well the model with a
particular K explains the genetic data. As the log-likelihood values are negative, the best-fit K-
value is the least negative. Delta K estimates the rate of change in the log probability of data
between successive K values. The K value corresponding to the highest delta K and estimated log
mean probability (K=9) was considered the best estimate of the number of subpopulations in this
dataset.