1 TITLE: DIVERSIFICATION AND COEXISTENCE AMID WIDESPREAD AND
2 PERSISTENT GENE FLOW IN ANDEAN TREES
3

4 Ellen J. Quinlan[1, 6*], James B. Pease[2], Jhonatan Sallo Bravo[3], Alfredo F. Fuentes[4,5], William
5 Farfan-Rios[1,6], and Miles R. Silman[1,6]
6
[1]Department of Biology, Wake Forest University, Winston-Salem, NC, USA 27109
[2]Department of Evolution, Ecology, and Organismal Biology, The Ohio State University,
Columbus, OH, USA 43210
[3]Facultad de Ciencias Biológicas, Universidad Nacional de San Antonio Abad del Cusco, Cusco,
Perú
[4]Latin America Department, Missouri Botanical Garden, St. Louis, MO, USA 63110
[5]Herbario Nacional de Bolivia, Universidad Mayor de San Andrés, La Paz, Bolivia
[6]The Sabin Center for Environment and Sustainability, Wake Forest University, Winston-Salem,
NC, USA 27109
16
17 Corresponding Author: Ellen J. Quinlan, quinej18@wfu.edu
18
ORCID
Ellen J. Quinlan 0000-0002-6249-649X
James B. Pease 0000-0003-0125-9156
Jhonatan Sallo Bravo 0000-0001-9007-4959
Alfredo F. Fuentes 0000-0003-4848-4182
William Farfan-Rios 0000-0002-3196-0317
Miles R. Silman 0000-0003-4152-2844
26

Total word count:	6509	Acknowledgements:	149
Summary:	159	Author contributions:	67
Introduction:	1613	No. of figures:	4 (main text), 3 (supplemental)
Materials and Methods:	1841	No. of tables:	3 (main text), 1 (supplemental)
Results:	869	No. of supporting information files:	1 (translated abstract)
Discussion	2181		

SUMMARY

•   Hybridization and introgression are increasingly recognized as important in plant evolution,

yet their prevalence and consequences remain poorly understood in tropical Andean trees,

where species coexist at low densities and turn over rapidly across elevation gradients.

•   Here, we investigate whether species boundaries are maintained among congeneric Andean

Prunus distributed across the Andes-Amazon elevational gradient by reconstructing

phylogenetic relationships, divergence times, and patterns of historical and recent gene flow.

•   We find that Andean *Prunus* diversified during the Miocene, likely in response to novel

habitat opportunities resulting from mountain uplift, combined with allopatric processes.

Despite this evolutionary divergence, these species show extensive ancient introgression and

recent admixture, indicating that species boundaries remain permeable across both ecological

and phylogenetic distance.

•   Together, these results suggest that Andean *Prunus* comprise distinct but evolutionarily

connected lineages that retain morphological identity while potentially functioning as a long-

lived syngameon, with implications for understanding the origin and maintenance of tropical

tree diversity and future response to global change.

Keywords: Andes, biodiversity, climate change, diversification, elevation gradient, gene flow,

hybridization, tropical trees

                              INTRODUCTION

Coexistence theories of ecology generally assume species are discrete evolutionary units

(e.g. Chesson 2000, Hubbell 2005), though interspecific gene flow is known to have been

common throughout the evolutionary history of plants (reviewed by Stull et al. 2023). No system

lends itself to tests of these theories better than species-rich tropical forests. Until recently,

hybridization was considered an extremely rare occurrence in tropical trees, as hybrids were

thought to be poor competitors in diverse systems and few individuals with intermediate

morphologies have been identified (e.g. Ashton 1969; Ehrendorfer 1970; Gentry 1982).

Mounting genomic evidence suggests hybridization has been widespread throughout the

diversification of the Neotropical flora (reviewed by Schley et al. 2022) and may be an important

evolutionary mechanism for maintaining high tree diversity with low population densities

(Cannon and Lerdau 2015, Cervantes et al. 2025, Quinlan et al. 2025). These findings

collectively point to the importance of reevaluating the impact of hybridization in our

understanding of species diversity and coexistence theories in tropical ecosystems.

The tendency for inter-specific gene flow and hybridization to occur amongst closely

related, sympatric trees has been widely documented in temperate zones (De La Torre et al.

2014, Tsuda et al. 2017, Cavender-Bares et al. 2018, Chhatre et al. 2018, Menon et al. 2018), yet

species cohesion and morphological similarity is often maintained (Grant 1971; Gailing and

Curtu 2014; Cannon and Petit 2020). This may be partially due to assortative (non-random)

mating and variable mating behaviors (selfing, conspecific outcrossing, and interspecific gene

flow), determined by population densities and pollinator abundance (Hamrick and Murawski

1990; Ward et al. 2005; Cannon and Lerdau 2015). When present, conspecific pollen can

outcompete heterospecific pollen, reinforcing assortative mating (Williams Jr. et al. 1999).

However, incomplete reproductive isolation and the ability to accept heterospecific pollen may

become adaptively beneficial when conspecific pollen is rare, as the benefit of avoiding local

extinction overrides potential negative consequences of hybridization such as reduced hybrid

fitness (Barton 2001, Wu 2001, Cannon and Lerdau 2015). Hybridization with introgression may

be particularly beneficial in times of rapid environmental change, as it can produce novel gene

combinations within a single generation, increasing adaptive potential and population resilience

(Anderson and Stebbins 1954, Baskett and Gomulkiewicz 2011, Suarez-Gonzalez 2018, Brauer

et al. 2023, Kulmuni et al. 2024). Such density-dependent mating has been reported in oaks

(Lepais et al. 2009) and may be particularly relevant to tropical trees which maintain extremely

low local population densities (Pires et al. 1953, Pitman et al. 1999, ter Steege et al. 2013).

Most tropical forest trees that have been studied are highly outcrossed (Bawa et al. 1985a,

b, Hamrick and Murawski 1990, Ward et al. 2005), though individuals retain variable capacities

for different mating behaviors, even within populations and across flowering events (Murawski

and Hamrick 1991, Nason and Hamrick 1997). Self-fertilization has been shown to be inversely

correlated with population density (Hamrick and Murawski 1990, Ward et al. 2005), but few

studies have assessed the extent of inter-specific gene flow. Those that have (e.g. Schley et al.

2020 – *Brownea*, Larson et al. 2021 – *Eschweilera,* Schley et al. 2025 – *Inga*), detected

widespread admixture. Evolutionary distance may be one important constraint on interspecific

gene flow among congeners in sympatry, as hybridization is thought to decrease with increasing

divergence due to the accumulation of pre- and postzygotic barriers through time (Mendelson et

al. 2004, Levin 2012, Comeault and Matute 2018). Linan et al. (2021) found evidence of this in

the Mascarene Island *Diospyros* clade, suggesting the processes by which species diversify and

assemble into communities may have lasting effects on their eco-evolutionary dynamics. For

example, adaptive radiations across gradients typically result in stepwise diversification patterns

with sister taxa occurring closest in niche proximity (Simpson 1953, Schluter 2000, Stroud and

Losos 2016), whereas communities assembled predominantly via the immigration and sorting of

pre-adapted clades demonstrate greater phylogenetic distance among co-occurring taxa

(Donoghue 2008, Dexter et al. 2017, Linan et al. 2021).

In tropical montane forests, genera commonly comprise many rare species that turn over

rapidly with elevation (Terborgh 1971, Gentry 1982). Turnover in the abiotic environment and

biotic interactions leads to narrow, shoestring-like distributions and the packing of many closely

related species across elevation gradients (e.g. Janzen 1967, Jankowski et al. 2009, Hillyer and

Silman 2010, Freeman et al. 2022). In the tropical Andes, this is consistent across taxa and is

especially true for forest trees (e.g. Terborgh 1977, Meier et al. 2010, Jankowski et al. 2012).

While many tree species have elevation ranges of <500 m, the distributions of entire genera are

often much broader, with congeneric species occurring in partial sympatry, especially at range

margins (Feeley et al. 2011, Griffiths et al. 2021). At the genus level, immigration and sorting of

pre-adapted clades has been shown to be the primary mechanism structuring tree diversity across

elevation gradients in the Andes (Griffiths et al. 2021, Segovia et al. 2020, Linan et al. 2021). However, many examples of adaptive diversification have been found within individual plant clades (e.g. Nürk et al. 2013, Lagomarsino et al. 2016, Nevado et al. 2016, among many others). Still, few studies have explored the evolutionary relationships among congeners within individual tree clades that co-occur across the gradient, and characterizing these evolutionary relationships is essential for understanding potential gene flow and its ecological implications.

Prunus L. (Rosaceae) is one such genus that exhibits high species turnover with elevation in the Andes (Figure 1). Globally, it has diversified into a range of environments, and though typically considered a temperate group (Wen et al. 2008, Chin et al. 2014), a recent revision found its highest diversity occurs in the tropics and primarily the Neotropics (~228 Neotropical species; Pérez-Zabala 2022). The Andes contain the highest number of species, with 41 total taxa including 39 endemics identified from the Central Andes alone. A total of 33 taxa, including 10 endemics are known from the Amazon, Guianas and northern South America (Pérez-Zabala 2022). *Prunus* is one of the only tree genera with species spanning the entire Andes-Amazon gradient, from treeline (~3800 m) into the lowlands (<500 m), though individual species have more narrow ranges, typically 500–1000 m or less (e.g. Achá 2013). It is a keystone clade, whose fruits are important in the diets of frugivorous birds, primates, Andean bears, and humans (Achá 2013). The genus also contains many globally important crop species (peach, plum, almond, cherries) and others used as ornamentals, timber, and medicine (Andro and Riffaud 1995, Bortiri et al. 2001, Lee and Wen 2001, Wen et al. 2008).

The genus *Prunus* is divided into three major clades that correspond to inflorescence type: solitary, corymbose, and racemose (Rehder 1940, Su et al. 2023). All species endemic to the Americas and tropical regions are racemose (Pérez-Zabala 2022). Though historically understudied, the exclusively South American species studied to date form a strongly supported clade nested within the tropical racemose group (Hodel et al. 2023—11species, Hyb-Seq and full plastome sequencing; Pérez-Zabala 2022—30 species, PCR). Studies have placed the origin of the genus in either North America (Chin et al. 2014) or East Asia (Hodel et al. 2023) during the Paleocene (ca. 68–55 Ma), with the three major clades diverging by the early-mid Eocene as part of the Boreotropical flora (Chin et al. 2014, Hodel et al. 2023, Su et al. 2023). Divergence in the tropical Racemose group occurred in several regions including Asia, Africa, North America, and

Europe ca. 55–40 Ma (Hodel et al. 2023). Chin et al. (2014) hypothesized migration into South

America occurred via the dispersal of a North American relative ca. 34 Ma through Central

America and the Caribbean. Hodel et al. (2023) found evidence of a long period of stasis in the

Neotropical lineages followed by rapid diversification during the Miocene.

While most species in the solitary-flower and corymbose groups are diploids

(2n=2x=16), the racemose lineage is known for higher ploidy levels, with allopolyploidy and/or

hybridization thought to have driven early diversification in the genus (Chin et al. 2014, Zhao et

al. 2016, Hodel et al. 2021, Su et al. 2023). Hodel et al. (2023) concluded that the majority, if not

all racemose species are polyploid, and all tropical species studied to date have been tetraploid

(2n=4x=32). The two exceptions being *P. laurocerasus* (2n=22x=176) and *P. gongshanensis*

(2n=8x=64)—species native to old world Europe and Southeast Asia (Hodel et al. 2023, Xie et

al. 2025). Domesticated *Prunus* (particularly plums) readily hybridize in agricultural settings

(Cici and Acker 2010), resulting in the creation of some crops (e.g. *P. armeniaca* L. × *P.*

cerasifera Ehrh. – 'black apricots,' *P. salicina* × *P. armeniaca* Lindl. – plumcots, pluots, and

apriums; Guerrero et al. 2022), as well as among cultivated varieties and wild conspecifics (*P.*

yedoensis - Baek et al. 2018). Species in this genus are highly outcrossing and typically self-

incompatible with generalist insect pollinators (Cici and Acker 2010), though we are unaware of

any study that has specifically looked at either breeding systems or pollination in Neotropical

Prunus.

Here, we investigate the extent to which species boundaries are maintained among

congeneric Andean trees, across elevation gradients and through time. Specifically, we ask: (1)

what are the evolutionary relationships among species that turnover with elevation, (2) are these

species are genetically isolated and (3) is genetic isolation related to evolutionary or ecological

distance, with ecological distance defined as elevational range overlap. To do this, we

reconstructed the phylogenetic relationships among sympatric *Prunus* species, estimated the

timing of divergence, and tested for historical introgression and recent admixture using RADseq

data. The results of these analyses expand current understanding of the origin and maintenance of

tropical Andean tree diversity and provide important context for predicting how these species

may respond to rapid environmental change.

167   MATERIALS AND METHODS

 Study site and sample collection

Sampling for this study primarily focused on species which co-occur across the Andes

Biodiversity and Ecosystem Research Group (ABERG; andesbiodiversity.org) plot network in

Manu National Park, Peru (Figure 1). This site is located on the eastern slope of the tropical

Andes in southeastern Peru and is a 3.5 km elevational gradient from the Andean tree line to the

Amazonian lowlands. With 23-1 ha and 33-0.1 ha forest plots along a series of descending

ridges, this is the most continuous transect in the tropical Andes. To date, ~19,000 individual

trees have been documented in these plots from 100 families, 292 genera, and 1255 species

(Farfan-Rios et al. 2025).

Intensive field sampling was conducted for *Prunus* species along this transect, with

botanists searching in lateral transects centered on the 1-ha plots (approximately every 250 m)

for up to 6 hrs. Leaf tissue was field collected and stored in silica following the Smithsonian

Institution guidelines for collecting vouchers and tissues intended for genomic work (Funk et al.

2017). Samples were also included for species known to co-occur across an additional plot

network spanning 4 km of elevation in Madidi National Park, Bolivia, and from at least one

external population when possible. These samples were collected from herbarium specimens

previously not treated with alcohol or fresh tissue stored in silica collected from individuals in

the Cotapata region of Bolivia (Figure 1, Table S1). In total, we sampled 158 individuals from 13

species including two non-Andean racemose species (*P. myrtifolia* and *P. caroliniana*) and a

reference (Table S1). At least four of these species have not been included in any previous

phylogeny.

ddRAD-seq data

High molecular weight DNA was extracted from ~50 mg of dried leaf tissue using Aboul-

Maaty and Oraby's (2019) modified CTAB protocol for non-model plants with a few

modifications: samples were ground with a bead-beater, incubated at -20℃ overnight (and up to

24 hrs), and the DNA pellet was washed twice with 500 µl 70% ethanol. DNA was checked for

quality and quantity on a 1.5% agarose gel and Qubit before being sent to Floragenex

(Beaverton, Oregon, USA) for ddRAD library prep and sequencing. Samples were ligated with 6

bp barcodes and digested with the PstI/MseI+0 enzyme pair. Genomic data were demultiplexed

and assembled to sample using ipyrad v. 0.9.95 (Eaton and Overcast 2020) with an average of

2,215,444 ± 1,419,356 million reads (mean ± s.d.) recovered per sample.

Reads were assembled into orthologous loci using reference-based assembly in ipyrad.

Default settings were used to trim and filter reads and the clustering threshold was set to 0.85 as

recommended by the ipyrad manual and McCartney-Melstad et al. (2019) for phylogenetic

analyses. For the reference-based assembly, reads were mapped to the *Prunus persica* (peach)

genome (GCF_000346465.2; The International Peach Genome Initiative et al. 2013, Verde et al.

2017). The proportion of mapped reads varied across samples (mean = 67%, s.d. = 13%), but

variation in the quality of sample preservation and DNA extractions likely explain much of the

variation. When calling consensus alleles, we allowed for up to 4 alleles per sample following

Donoghue et al. (2022), as all tropical racemose *Prunus* species studied to date are tetraploids

(2n=4x=32; Zhao et al. 2016, Hodel et al. 2023, Xie et al. 2025). Cytogenetic data is not

available for most species included here, but the clade is thought to derive from an ancient

allopolyploid event (Hodel et al. 2023, Xie et al. 2025). Therefore, residual uncertainty in allele

dosage and homeologous variation may remain. To evaluate the sensitivity of downstream results

to allele representation, we also generated an artificially diploidized dataset by randomly

downsampling to a maximum of two alleles per site per sample, but the qualitative patterns were

like those represented by the full polyploid-aware assembly. The final reference-mapped dataset

comprised 35,099 loci, representing non-overlapping genomic regions where data from at least 4

samples passed filtering. Further downstream analyses were performed using vcftools and the

ipyrad-analysis toolkit, which allows for filtering and subsampling loci, genomic windows,

and/or SNPs for phylogenetic inference and D-statistics.

Individual-level phylogeny

A maximum likelihood (ML) phylogeny was reconstructed from a subset of

representative samples with the most data present per species (up to 10 individuals per species, at

least 1000 loci per sample; n=47), including conspecifics from multiple localities when possible.

The ipa.window_extracter module in the ipyrad-analysis toolkit was used to extract data from the

8 largest scaffolds (the chromosomes) and filter sites to only maintain those shared by at least

25% of individuals. We selected the 25% threshold after testing various other filtering thresholds

(10%, 30%, 40%, 50%), as 25% provided for the best phylogenetic resolution while also

including as much information as possible. The filtered and concatenated sequence matrix is 2.8

Mb in length, containing 107,708 SNPs and 65.8% missing data. This alignment was used in

RAxML (Stamatakis 2014) under the GTR+GAMMA evolutionary model to infer a best

phylogeny under the rapid hill-climbing algorithm, with support values from 100 bootstrap

replicates. The phylogeny was rooted to the reference genome *P. persica* (a native of China and

member of the Solitary-flower group and Amygdalus subgenus) and included two additional

non-Andean racemose species—*P. caroliniana* (native of the eastern United States and

Carribbean) and *P. myrtifolia* (native to the Caribbean, Central America, and lowland South

America).

Species-level phylogeny

To construct the species-level phylogeny, we selected one representative specimen for

each species in the individual-level phylogeny (n=13). The samples selected grouped with other

members of their species on the individual-level phylogeny and had the most data present. For

the species that were reconstructed as non-monophyletic on the individual-level tree, individual

selection did result in some movement on the species-level phylogeny. However, we found this

was largely linked to amount of missing data and therefore relied on the individual with the

highest genomic coverage as the representative. Again, we used the ipa.window_extracter

module in the ipyrad-analysis toolkit to extract data from the chromosomes and filter sites,

selecting those shared by at least 25% of all species (after testing multiple filtering thresholds).

The resulting sequence matrix is 4.2 Mb with 62,262 SNPs and 63.6% missing data. To

reconstruct the phylogenetic relationships, we used both maximum likelihood (RAxML) and

Bayesian (ExaBayes) analyses. We ran RAxML using GTR+GAMMA to infer a best phylogeny

under the rapid hill-climbing algorithm with support values from 100 bootstrap replicates. For

ExaBayes, we ran two Metropolis-coupling replicates with four coupled-chains (each with three

heated chains) for 1×10^6 MCMC generations, sampled every 500 generations. Estimated sample

size (ESS) for all parameters and branch lengths were summarized with the ProcParam tool and

confirmed as sufficiently sampled. The majority-rule consensus phylogeny was generated with

the ExaBayes 'consense' tool with a 25% burn-in.

To assess the robustness of the resolved relationships and distinguish between conflicted

and poorly informed branches, we used Quartet Sampling (Pease et al. 2018;

https://www.github.com/ fephyfofum/quartetsampling). We performed Quartet Sampling on the

resulting RAxML tree with the full concatenated alignment, allowing 100 replicates per branch

(Figure S2).

Divergence time dating

261   To assess the timing of divergence among species, we generated a time-calibrated

phylogeny using BEAST v. 2.7.7 (Bouckaert et al., 2019) with the same alignment used for the

species-level phylogenies. We used an optimized relaxed molecular clock and Yule speciation

process with the GTR+GAMMA substitution model. A nodal calibration was set for the crown

age of *Prunus* using a normal prior with an offset of 61.5 Mya and a standard deviation of 3.0

Mya (Chin et al. 2014; Su et al. 2023). The crown age of the racemose group was calibrated from

a staminate flower of *Prunus hirsutipetala* D.D. Sokoloff, Remizowa et Nuraliev preserved in

the late Eocene (Priabonian) Rovno amber from northwestern Ukraine (Sokoloff et al. 2018),

using a normal prior with an offset of 35.0 Mya and standard deviation of 2.0 Mya (following Su

et al. 2023). To ensure a more rapid convergence of MCMC runs, we used our RAxML topology

as our starting tree. The Markov chain Monte Carlo (MCMC) run was conducted for 20,000,000

generations and trees sampled every 2000 generations producing a total of 10,000 trees. The

resulting log files were checked in Tracer v.1.5 (Rambaut et al. 2018) and we used TreeAnnotator

v.1.10.4 to generate the maximum clade credibility tree discarding 25% of trees as burn-in before

visualizing the results in FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Tests of historical introgression

277   To test for historical introgression, we used the four-taxon Patterson's *D*-statistic (ABBA-

BABA), which quantifies allele frequency patterns that are incongruent with the hypothesized

species tree topology (Green et al. 2010). We ran the Patterson's *D*-tests using the ipa.baba tool

as implemented in ipyrad, which is specifically designed for analysis of RAD-seq datasets. We

used the species-level RAxML tree as our starting tree hypothesis and constrained the tests

setting the outgroup to *P. myrtifolia*, so all tests would focus on ingroup Andean species. To

assess significance, we performed 1000 bootstrap replicates for each test with loci resampled

with replacement. Z-scores were then calculated as the number of bootstrap standard deviations

from the null *D*=0 (Table 1). Z-scores > 2.5 were considered significant.

 Tests of recent admixture

To assess genetic structure and test for recent admixture, we used the Bayesian clustering

algorithm STRUCTURE v. 2.3.4 (Pritchard et al. 2000) as implemented in ipyrad with the set of

samples included in the individual-level phylogeny (minus the reference and non-Andean

species). SNPs were filtered conservatively to only include those with at least 50% coverage

across individuals and then randomly sampled to one SNP per locus to avoid linkage. This

resulted in a total of 1493 SNPs included in the analysis. The model was run with a burn-in of

125,000 and run length of 500,000 generations, testing K=1–10 with 10 replicates per K. The

assignment of samples into K distinct genetic groups was assessed using DeltaK, which has been

shown to be more robust to accurately estimating the correct number of clusters in

STRUCTURE, particularly with high amounts of missing data as is common in RADseq

(Evanno et al. 2005; Figure S3).

Estimates of genomic diversity

Genetic diversity was estimated within individuals and species to compare to the

admixture results. In VCFtools v.1.16 (Danecek et al. 2011), sequences were filtered using two

different missing data thresholds, retaining loci present within at least 50% or 70% of the

samples for each species. Sites were also filtered if they had a minor allele frequency less than

0.05. We calculated nucleotide diversity (π) for each species – the average pairwise nucleotide

differences between sequences in a sample at a site as well as observed (H_o) and expected

heterozygosity (H_e) – the proportion of loci where an individual is heterozygous vs. the

proportion expected by Hardy-Weinberg equilibrium – within samples and averaged across the

species. For these calculations, *P. debilis* and *P. detrita* samples were grouped together as well as

P. huantensis and *P. brittoniana,* given that STRUCTURE (Figure 4) did not detect any distinct

genetic clusters among these species' pairs. Species with <3 samples were excluded from these

analyses given the lack of statistical power from such a small sample size.

<div align="center">RESULTS</div>

Phylogenomic relationships and divergence time estimation

Both the species-level maximum likelihood (RAxML) and Bayesian (ExaBayes)

phylogenetic analyses supported the same evolutionary relationships among the focal Andean

species and the non-Andean lineages (Figure 2). Subtropical *P. caroliniana* was reconstructed as

sister to the tropical species and *P. myrtifolia* was sister to the Andean species. Among the partially sympatric Andean species, the lineage including pre-montane species *P. debilis* and *P. detrita* were sister to the remaining Andean species, followed by the mid-elevation cloud-forest species (*P. herthae, P. pleiantha, P. stipulata, P. integrifolia*) with the high-elevation species being the most recently derived (*P. huantensis, P. brittoniana, and P. antioquensis*). Statistical support for the inferred relationships was assessed using bootstrap support (RAxML), posterior probabilities (ExaBayes), and Quartet Sampling.

Quartet Sampling (QS; Pease et al. 2018) complements conventional node-support metrics by distinguishing low support caused by limited phylogenetic information from low support caused by conflicting signal. It evaluates each node using the relative frequency of alternative relationships among sets of four taxa. In this framework, QI (quartet informativeness) measures the proportion of sampled quartets that are informative for a given node, QC (quartet concordance) measures the degree to which informative quartets support the focal relationship, and QD (quartet differential) measures the skew between the two alternative discordant topologies, helping distinguish balanced discordance (as expected under incomplete lineage sorting) from a strong preference for one alternative (consistent with introgression). While the tree nodes are generally well-supported (all posterior probabilities = 1, most bootstrap values >75; Figure 2), Quartet Sampling indicated that low bootstrap support values are not likely the result of insufficient phylogenetic signal, as QI ranged from 80–100%. The lack of a strong skew in discordant quartets at most nodes (QD>0.5) suggests that incomplete lineage sorting (ILS) is more likely than introgression. In contrast, the branch subtending *P. huantensis, P. brittoniana*, and *P. antioquensis* showed moderate concordance (QC=0.21) and strong discordant skew (QD=0.22), signal consistent with a possible introgressive event (Figure S2).

The time-calibrated phylogeny generated in BEAST recovered the same tree topology as RAxML and ExaBayes with similar branch support (Figure 3). The estimated divergence time *for P. myrtifolia* and the ancestor of the Amazonian and Andean species is ~27.3 Ma (95% highest posterior density [HPD] 22.4–31.7). The Andean species began diverging at the start of the Miocene, with the pre-montane *P. debilis+detrita* lineage branching off ~23.5 Ma (95% HPD 19.4-28.1), and from each other ~17.6 Ma (95% HPD 12.8-22.6). The cloud forest species followed, with the *P. pleiantha* lineage diverging ~20.4 Ma (95% HPD 16.3-24.4), and *P.*

integrifolia ~18.0 Ma (95% HPD 14.2-21.7). The remaining lineages emerged in the mid to late-

Miocene (15.9–9.3) with *P. huantensis* ~11.6 Ma (95% HPD 8.6-14.9) and *P. antioquensis+*

brittoniana ~9.3 Ma (95% HPD 6.2-12.3). Together, the phylogenetic analyses indicate that these

lineages are ancient, with potential sympatry since the Miocene.

Tests of historical introgression and recent admixture

D-statistics indicated significant historical introgression events between *P. pleiantha* and

the ancestor of *P. debilis+detrita*, with more limited evidence suggesting introgression with *P.*

debilis and *P. detrita* separately (Figure 2, Table 1). A second significant introgression event was

detected between *P. pleiantha* and the *P. huantensis+antioquensis* ancestor, which is further

supported by Quartet Sampling (Figure S2). Finally, significant *D*-statistics supporting

introgression between *P. debilis* and *P. antioquensis* were also detected. Overall, introgression is

widespread and does not seem to be limited by ecological or evolutionary distance, as evidenced

by their elevational niche overlap (Figure 1) and hypothesized phylogenomic relationships

(Figure 2).

STRUCTURE analysis found K=9 to be the most likely number of distinct genetic

clusters within the dataset, determined by mean log probability and DeltaK values (Figures 5,

S3). Individuals largely clustered according to their morphologically assigned species; however,

the number of detected clusters was fewer than the expected number of named taxonomic species

in the analysis (*n*=10). *P. integrifolia* and *P. amplifolia* are largely reproductively isolated, though

low levels of admixture were detected in *P. integrifolia* and only one sample was included for *P.*

amplifolia. Admixture is widespread amongst all other species, particularly *P. pleiantha*, *P.*

stipulata, *P. herthae*, *P. debilis, P. huantensis, P. brittoniana* and *P. antioquensis,* and like the

introgression analyses, appears unrelated to ecological or evolutionary distance.

Estimates of genomic diversity

Within each species, the individuals with the highest observed heterozygosity (H_o) were

also the most admixed in the STRUCTURE results (Tables 3 and 4, Figure 4), and this was

consistent with different amounts of missing data. For example, in *P. pleiantha*, individuals 163,

164, and 165 showed high levels of admixture from multiple groups and had the highest H_o

estimates (50%: 0.18 – 0.22, 70%: 0.19 – 0.25; Table 3). Across species and filtering thresholds,

mean pairwise nucleotide diversity (π) was high (50%: 0.41 – 0.64, 70%: 0.37 – 0.51) compared

to H_o, π tracked with H_e as expected, and H_o (50%: 0.06 – 0.15, 70%: 0.06 – 0.17) was far below

H_e (50%: 0.41 – 0.64; 70%: 0.37 – 0.51; Table 2). However, the estimates for *P. herthae* and the

P. debilis+detrita group should be interpreted cautiously, given the low sample sizes for each.

These statistics corroborate the findings of the STRUCTURE analysis and further indicate a

history of interspecific gene flow among co-occurring *Prunus* species.

381   DISCUSSION

In this study, we aimed to characterize the evolutionary relationships among congeneric

tree species that turn over with elevation in the tropical Andes, assess whether they are

genetically isolated, and the extent to which gene flow is related to ecological or evolutionary

distance. The phylogenomic data presented here suggest that *Prunus* species known to co-occur

as elevational replacements diversified across the Miocene (~23.5-9.3 Ma). However, despite

these lineages being separated by considerable evolutionary time, species are not reproductively

isolated, as substantial gene flow is detectable among species through time and does not decrease

with increasing evolutionary distance or elevational separation. Together, these findings suggest

that during diversification, observable ecological and morphological differences may emerge

despite ongoing gene flow. Further, the maintenance of incomplete reproductive barriers over

millions of years may be an important and overlooked mechanism for maintaining high tree

diversity, particularly with the low population densities and high environmental variability

characteristic of tropical montane forests.

Gene flow through time

Our genomic analyses show that Andean *Prunus* species remain distinct but porous

evolutionary lineages. Despite deep Miocene divergences, we detect both ancient introgression

and recent admixture among partially sympatric taxa. *D*-statistics detected at least five

significant ancient introgression events. Two of these events occurred among lineages separated

by great evolutionary and elevational distance (*P. pleiantha* × *P. huantensis+ antioquensis*, and *P.*

debilis × *P. antioquensis;* Figure 2*)*. Introgression was also detected between *P. pleiantha* and the

ancestor of *P. debilis+ detrita* (*P. pleiantha* × *P. debilis+ detrita*) as well as among *P. pleiantha* ×

P. debilis and *P. pleiantha* × *P. detrita* individually. These patterns are most consistent with

repeated interspecific introgression among the ancestors of these lineages, with gene flow

continuing as elevationally separated lineages diverged. However, because D-statistics do not
resolve the direction of introgression, a stronger test of this interpretation will require methods
that can distinguish donor and recipient lineages (such as D-FOIL; Pease and Hahn 2015),
together with denser taxonomic and wider genomic sampling.

Historical introgression among species separated in ecological space across the elevation
gradient is plausible for several reasons. While the medians of their elevation ranges may be
quite different, they are found in close geographic proximity, and their elevational ranges overlap
(or are much closer) at the range margins. *Prunus* species are thought to be obligate outcrossers
with generalist pollinators, as is common in tropical trees (Ashton 1969, Bawa 1974, 1979). The
extent of self-incompatibility across the genus and potential for breakdown after polyploidization
are still unclear (Hauck et al. 2006, Tao et al. 2010, Aguiar et al. 2015). Generalist pollinators can
forage over kilometers, providing opportunity for gene flow, especially given that many tropical
species flower semi-synchronously during the dry season (Bawa et al. 1985, Van Schaik et al.
1993).

Even where known species ranges are separate at present, they have not necessarily
always been so. Pollen records from Lago Consuelo (1360 m a.s.l., located in southeastern Peru
adjacent to the Bolivian border) show that during the last full glacial period (43.5–22 kyr BP),
individual cloud forest species migrated downslope at least 1000–1700 vertical m in response to
lower cloud base and cooler temperatures (-5°C; Urrego et al. 2005). Similar pulses of
connectivity likely occurred as many as ten times in response to glacial-interglacial cycles over
just the last million years and may have helped maintain these incomplete reproductive
boundaries. Today, these species ranges are separated as far apart as they have likely been during
the Quaternary, and historical introgression events may be signatures of past ecological overlap.

Tests of recent admixture also found gene flow to be widespread among Andean *Prunus*
species. *P. integrifolia* and *P. amplifolia* are largely genomically isolated, but gene flow appears
rampant among all other species. The isolation of *P. integrifolia* is particularly interesting as this
species has the widest elevational range and is one of the most abundant species in the tropical
Andes. The single sample of *P. amplifolia* included in the analysis prevents us from determining
whether this pattern of isolation is widespread in the species (as observed in *P. integrifolia*) or
whether rates of mixed genetic ancestry are highly variable (as observed in *P. pleiantha*). Gene

flow was especially high among *P. pleiantha, P. stipulata*, and *P. herthae*, all cloud forest species

with largely overlapping elevation ranges. However, individuals of these species mostly form

monophyletic groups, indicating they have developed specific genetic distinctiveness as well as

morphological similarity that allows for field identification despite gene flow (Figure S1). As *P.*

pleiantha was identified in most of the historical introgression events and had high proportions

of mixed genetic ancestry, we can infer that this species readily hybridizes when in contact with

close relatives. Yet, previous morphological studies of herbarium specimens (Achá 2013, Pérez-

Zabala 2022) have not recovered high morphological variation in *P. pleiantha* or *P. herthae*, as

would be expected if morphological variation increased with gene flow. Further investigation

will be needed to test these hypotheses, identify the directionality of such gene flow, and

determine whether physiological, ecological, or genetic factors drive this weaker reproductive

isolation and its implications.

These inferences should also be interpreted considering the likely shared ancient

allopolyploid origin of the clade. Residual homeologous variation or differential sorting of

ancestral subgenomes, may increase apparent allele sharing among taxa and complicate the

interpretation of admixture and D-statistic results, particularly in analyses that reduce polyploid

variation to diploid-like summaries. However, because this polyploid background is likely shared

across the clade, any resulting bias should be widespread rather than specific to particular taxa.

Polyploid history may therefore contribute to overall ambiguity in the genomic signal, but is

unlikely to fully explain the repeated, lineage-specific patterns of excess allele sharing recovered

here.

STRUCTURE did not identify *P. debilis* and *P. detrita* as distinct genetic clusters,

possibly due to how closely related these species are, gene flow, and the low number of

individuals for both species in the dataset. Our genomic data also support the most recent

taxonomic revision by Pérez-Zabala (2022) which considered *P. brittoniana* and *P. huantensis* to

be the same species given their morphological and ecological similarities (Figure S1). Taken

together, these results indicate that gene flow has been a persistent and complex feature of

Andean *Prunus* diversification, occurring across both evolutionary time and ecological space,

despite ancient divergence times and emergence of species-level distinctiveness.

Diversification patterns and biogeographic implications

Phylogenomic reconstructions and divergence time dating indicate that *Prunus* diversified in the tropical Andes over the Miocene. In agreement with prior studies, we found *P. myrtifolia* (endemic to the Caribbean, Central America, and lowland South America) to be sister to the Andean species included in this analysis, supporting the hypothesis that this lineage or its ancestor colonized South America via migration through the Caribbean and Central America (Chin et al. 2014, Hodel et al. 2023). While our divergence time estimate for *P. myrtifolia* of ~27 Ma is slightly later than the Neotropical colonization of ~34 Ma predicted by Chin et al. (2014), it is still within the climatic shift from warmhouse to transitional state that coincided with an onset of glaciation in Antarctica (~33.9 – 27.29 Ma; Judd et al. 2024) and the estimated end of the boreotropical forests in the Northern Hemisphere.

Among the Andean species included in our analyses, the two lowest elevation species (*P. debilis* and *P. detrita*) were sister to the rest of the group (diverging ~23.5 Ma) and the high-elevation clade was the most recently derived (~11.6 Ma—*P. huantensis*, ~9.3 Ma—*P. antioquensis* and *P. brittoniana*). This is consistent with the timing of major uplift events and estimated paleo-elevations in the North-Central and Central Andes (Boschman 2021). It also supports the findings of Hodel et al. (2023), which detected a long period of stasis followed by rapid diversification of Neotropical *Prunus* during the Miocene. Although the exact timing, pace, and sequence of Andean orogeny is a subject of ongoing study and is spatially heterogeneous, the late Oligocene and Miocene (~25-5 Ma) are known to have been a period of intense and accelerated uplift in the Eastern Cordillera of the North-Central and Central Andes (Boschman et al. 2021, Pérez-Escobar et al. 2022).

The lower montane cloud forest species in our analysis (median elevation ranges 1200–2500 m; *P. pleiantha, P. integrifolia, P. herthae, P. stipulata*) diverged during the early to mid-Miocene. This aligns with the predicted origination of cloud forests (Sempere et al. 2005) and the estimated divergence times for other cloud forest lineages (Luebert and Weigend 2014). Andean cloud forests are distinct habitats characterized by persistent cloud cover and immersion that occurs from ~1200 m to the upper forest limits at ~3200–3800 m (depending on latitude). Although they are the most speciose habitats in the Andes, persistence in these environments often requires unique adaptations to cope with the reduced light availability, high precipitation, and humidity such as thick, pachyphyllic leaves, low growth rates, and ability to withstand heavy

accumulations of epiphytes (Grubb 1977, Rapp and Silman 2012, Fahey et al. 2016, Pérez-

Escobar 2022). The highest elevation species in our study (median elevation ranges ≥3000 m; *P.*

huantensis/ P. brittoniana, P. antioquensis) diverged during the late Miocene, as the final uplift

phase began in the North-Central and Central Andes (Boschman et al. 2021). Species in this

elevation zone are exposed to unique high-elevation stressors such as night frost, with

temperatures dipping below 0ºC, requiring further physiological specialization (Pérez-Escobar

2022).

Given the phylogenomic reconstructions and divergence time estimates, two hypotheses for

understanding diversification among Andean *Prunus* emerge. First, these species may be the

result of adaptive diversification following Andean uplift. Similar diversifications have been

documented in other Andean plant clades in response to novel mountain habitat opportunities

(e.g. Melastomataceae—Dudley 1978, wild tomato—Pease et al. 2016). Dating analysis from

this and previous studies (e.g. Chin et al. 2014, Hodel et al. 2023) suggest *Prunus* arrived in

South America during the early to middle Oligocene (~34-27 Mya), long before Miocene

mountain building brought the central Andes to their modern elevations. However, an alternate

and often dominant pattern of diversification in the Andes is one in which extratropical genera

emigrate in and colonize elevation bands for which they were pre-adapted (Griffiths et al. 2020,

Segovia et al. 2020, Linan et al. 2021). These elevation bands then become fragmented by

topography, climatic oscillations, and geological processes, promoting allopatric speciation.

Examples of such latitudinal allopatric speciation have been documented across plant and bird

genera (e.g. *Ceroxylon*—Sanín et al. 2016, *Gunnera*—Bacon et al. 2018, *Scytalopus*—Cadena et

al. 2020).

Importantly, these modes of speciation (adaptive vs. allopatric) are not mutually exclusive.

Post-colonization, diversification may occur both in response to novel habitat opportunities

(across elevation) and following isolation events (across latitude), with both mechanisms

responsible for generating the immense rates of endemism and diversity characteristic of tropical

Andes (e.g. lupines—Hughes and Eastwood 2006, *Hypericum*—Nürk et al. 2013, bellflowers—

Lagomarsino et al. 2016, orchids—Pérez-Escobar et al. 2017). Disentangling when and how

these mechanisms gave rise to the >100 species of South American *Prunus* (Pérez-Zabala 2022)

will require denser species sampling than available in this study as well as higher-resolution

species distribution data. Within those limitations, our results are most consistent with

diversification in Andean *Prunus* being driven by the joint effects of adaptive diversification in

response to uplift-driven ecological opportunity and the immigration, sorting, and isolation of

lineages adapted to different elevational environments.

Conclusions

Here, we found evidence of historical introgression, possibly in response to Quaternary

range shifts, and widespread admixture among co-occurring *Prunus* species. Yet despite gene

flow, these species have maintained distinct evolutionary lineages with identifiable ecological

and morphological differences since the Miocene. These results align with the syngameon

model, wherein species maintain incomplete reproductive barriers but primarily experience

divergent selection (Lotsy 1925, Grant 1971, Cannon and Lerdau 2015, Cervantes et al. 2025,

Schley et al. 2025). Though oaks are the classic example (e.g. Cannon and Petit 2020), such

genomic mutualism may be an important yet under-appreciated mechanism in the maintenance

of diversity in species-rich tropical forests, where individuals predominantly outcross but retain

some capacity for self-fertilization and interspecific hybridization at low population densities

(Wu 2001, Cannon and Lerdau 2015). This has important implications in the context of climate

change, as species are now shifting their ranges in response to changes in temperature and

moisture (Feeley et al. 2011, Fadrique et al. 2018). Though most tree species will not be able to

migrate fast enough to track climatic shifts and are instead experiencing range retractions, any

upslope range shift may increasingly bring congeneric species into contact, especially at sparsely

populated range margins where low conspecific density could elevate the likelihood of

heterospecific mating and gene flow (Duque et al 2015, Cannon and Lerdau 2015, Farfan-Rios et

al. 2025, Quinlan et al. 2025). Further work is required to understand the prevalence of

syngameons in tropical forests and their role in species' responses to rapid environmental

change, namely resilience and avoidance of local extinction. Incorporating these complex

evolutionary dynamics into demographic models is essential for accurately predicting the futures

of Andean forests and effective conservation and management under current climate change.

ACKNOWLEDGEMENTS

We thank Paul Fine, Michael Anderson, and Regina Joice Cordy for their helpful comments on study design and earlier versions of this manuscript. We are also grateful to Herbario Vargas (CUZ) and Herbario Nacional de Bolivia (LPB) for their assistance with coordination, field work, and access to herbarium material, as well as Christopher Dick and Fairchild Botanical Garden for their donation of samples to this project. Additional thanks to the Andes Biodiversity Ecosystem Research Group (ABERG), and the many students and field technicians involved with this project (especially Andrea Palomino Cardenas, Jesus Manuel Bañon Sanchez, Daire Luciana Lafuente Palie, Paukar Alejandro Vedia Villca, and Felipe Aramburo Jaramillo). This work was supported by NSF LTREB (1754647) and Garden Club of America's Tropical Botany Fellowship. The material used in this study was collected under the research permits No 18-2019-SERNANP-JEF, No 14-2022-SERNANP-JEF, No 527-2019-MINAGRI-SERFOR-DGGSPFFS, and No D000170-2023-MIDAGRI-SERFOR-DGGSPFFS-DGSPF granted by SERNANP and SERFOR.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

EJQ, MRS, and JBP initially conceived of the study. EJQ acquired funding, led research design, data collection, data analysis and interpretation, and wrote the manuscript. JSB, AFF, and WFR assisted with data collection and WFR additionally contributed to manuscript editing and translated the summary. JBP assisted with data analysis and manuscript review and editing. MRS provided supervision, acquired funding, and contributed to research design and manuscript writing.

DATA AVAILABILITY

The genomic data generated for this study are available on NCBI's GenBank under submission ID SUB15831409 and corresponding herbarium vouchers are deposited in Herbario Vargas (CUZ) at the Universidad Nacional de San Antonio Abad del Cusco (UNSAAC) and Herbario Nacional de Bolivia (LPB).

581   REFERENCES

Aboul-Maaty, N. A. F., & Oraby, H. A. S. (2019). Extraction of high-quality genomic DNA from
different plant orders applying a modified CTAB-based method. *Bulletin of the National*
Research Centre, *43*(1), 1-10.

Achá, S. (2013). El género Prunus (Rosaceae) en Bolivia: Novedades en su sistemática.

Aguiar, B., Vieira, J., Cunha, A. E., Fonseca, N. A., Iezzoni, A., van Nocker, S., & Vieira, C. P.
(2015). Convergent evolution at the gametophytic self-incompatibility system in Malus and
Prunus. *PloS one*, *10*(5), e0126138.

Anderson, E., & Stebbins Jr, G. L. (1954). Hybridization as an evolutionary stimulus. *Evolution*,
378-388.

Andro, M. C., & Riffaud, J. P. (1995). Pygeum africanum extract for the treatment of patients
with benign prostatic hyperplasia: a review of 25 years of published experience. *Current*
therapeutic research, *56*(8), 796-817.

Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of
recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

Bacon, C. D., Velásquez-Puentes, F. J., Hinojosa, L. F., Schwartz, T., Oxelman, B., Pfeil, B., ... &
Antonelli, A. (2018). Evolutionary persistence in Gunnera and the contribution of southern plant
groups to the tropical Andes biodiversity hotspot. PeerJ, 6, e4388.

Baek, S., Choi, K., Kim, G. B., Yu, H. J., Cho, A., Jang, H., ... & Mun, J. H. (2018). Draft
genome sequence of wild Prunus yedoensis reveals massive inter-specific hybridization between
sympatric flowering cherries. *Genome biology*, *19*, 1-17.

Barton, N. H. (2001). The role of hybridization in evolution. *Molecular ecology*, *10*(3), 551-568.

Baskett, M. L., & Gomulkiewicz, R. (2011). Introgressive hybridization as a mechanism for
species rescue. *Theoretical Ecology*, *4*, 223-239.

Bawa, K. S. (1974). Breeding systems of tree species of a lowland tropical community.
Evolution, 85-92.

Bawa, K. S. (1979). Breeding systems of trees in a tropical wet forest. *New Zealand Journal of*
Botany, *17*(4), 521-524.

Bawa, K. S., Perry, D. R., & Beach, J. H. (1985a). Reproductive biology of tropical lowland rain
forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of*
botany, *72*(3), 331-345.

Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H. (1985b). Reproductive biology of tropical lowland rain forest trees. II. Pollination systems. *American journal of botany*, *72*(3), 346-356.

Boschman, L. M. (2021). Andean mountain building since the Late Cretaceous: A paleoelevation reconstruction. *Earth-Science Reviews*, *220*, 103640.

Bortiri, E., Oh, S. H., Jiang, J., Baggett, S., Granger, A., Weeks, C., ... & Parfitt, D. E. (2001). Phylogeny and systematics of Prunus (Rosaceae) as determined by sequence analysis of ITS and the chloroplast trnL-trnF spacer DNA. *Systematic Botany*, *26*(4), 797-807.

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M., Gavryushkina, A., ... & Drummond, A. J. (2019). BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS computational biology*, *15*(4), e1006650.

Brauer, C. J., Sandoval-Castillo, J., Gates, K., Hammer, M. P., Unmack, P. J., Bernatchez, L., & Beheregaray, L. B. (2023). Natural hybridization reduces vulnerability to climate change. *Nature Climate Change*, *13*(3), 282-289.

Cadena, C. D., & Céspedes, L. N. (2020). Origin of elevational replacements in a clade of nearly flightless birds: most diversity in tropical mountains accumulates via secondary contact following allopatric speciation. In *Neotropical diversification: Patterns and processes* (pp. 635-659). Cham: Springer International Publishing.

Cannon, C. H., & Lerdau, M. (2015). Variable mating behaviors and the maintenance of tropical biodiversity. *Frontiers in Genetics*, *6*, 183.

Cannon, C. H., & Petit, R. J. (2020). The oak syngameon: more than the sum of its parts. *New Phytologist*, *226*(4), 978-983.

Cavender-Bares, J., Kothari, S., Meireles, J. E., Kaproth, M. A., Manos, P. S., & Hipp, A. L. (2018). The role of diversification in community assembly of the oaks (Quercus L.) across the continental US. *American Journal of Botany*, *105*(3), 565-586.

Cervantes, S., Schley, R., Hardy, O. J., & Ojeda, D. (2025). Do syngameons exist in tropical trees? Challenges to determine their existence and estimate their frequency. Biology Letters, 21(11). https://doi.org/10.1098/rsbl.2025.0444.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology and Systematics*, *31*(1), 343-366.

Chhatre, V. E., Evans, L. M., DiFazio, S. P., & Keller, S. R. (2018). Adaptive introgression and maintenance of a trispecies hybrid complex in range-edge populations of Populus. *Molecular ecology*, *27*(23), 4820-4838.

Chin, S. W., Shaw, J., Haberle, R., Wen, J., & Potter, D. (2014). Diversification of almonds, peaches, plums and cherries–molecular systematics and biogeographic history of Prunus (Rosaceae). Molecular phylogenetics and evolution, 76, 34-48.

Cici, S. Z. H., & Van Acker, R. C. (2010). Gene flow in Prunus species in the context of novel trait risk assessment. *Environmental biosafety research*, *9*(2), 75-85.

Comeault, A. A., & Matute, D. R. (2018). Genetic divergence and the number of hybridizing species affect the path to homoploid hybrid speciation. *Proceedings of the National Academy of Sciences*, *115*(39), 9761-9766.

Danecek, Petr, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A. DePristo, Robert E. Handsaker et al. "The variant call format and VCFtools." *Bioinformatics* 27, no. 15 (2011): 2156-2158.

De La Torre, A. R., Roberts, D. R., & Aitken, S. N. (2014). Genome-wide admixture and ecological niche modelling reveal the maintenance of species boundaries despite long history of interspecific gene flow. Molecular Ecology, 23(8), 2046-2059.

Donoghue, M. J. (2008). A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences*, *105*(supplement_1), 11549-11555.

Donoghue, M. J., Eaton, D. A., Maya-Lastra, C. A., Landis, M. J., Sweeney, P. W., Olson, M. E., ... & Edwards, E. J. (2022). Replicated radiation of a plant clade along a cloud forest archipelago. *Nature Ecology & Evolution*, *6*(9), 1318-1329.

Duque, A., Stevenson, P. R., & Feeley, K. J. (2015). Thermophilization of adult and juvenile tree communities in the northern tropical Andes. *Proceedings of the National Academy of Sciences*, *112*(34), 10744-10749.

Dudley, E. C. (1978). Adaptive radiation in the Melastomataceae along an altitudinal gradient in Peru. *Biotropica*, 134-143.

Eaton, D. A., & Overcast, I. (2020). ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics*, *36*(8), 2592-2594.

Ehrendorfer, F. (1970). Evolutionary patterns and strategies in seed plants. *Taxon*, *19*(2), 185-195.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular ecology*, *14*(8), 2611-2620.

Fahey, T. J., Sherman, R. E., & Tanner, E. V. (2016). Tropical montane cloud forest: environmental drivers of vegetation structure and ecosystem function. *Journal of Tropical Ecology*, *32*(5), 355-367.

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., ... & Feeley, K. J. (2018).
Widespread but heterogeneous responses of Andean forests to climate
change. *Nature*, *564*(7735), 207-212.

Farfan-Rios, W., Feeley, K. J., Myers, J. A., Tello, S., Sallo-Bravo, J., Malhi, Y., ... & Silman, M.
R. (2025). Amazonian and Andean tree communities are not tracking current climate warming.
Proceedings of the National Academy of Sciences, 122(34), e2425619122.

Feeley, K. J., Silman, M. R., Bush, M. B., Farfan, W., Cabrera, K. G., Malhi, Y., ... & Saatchi, S.
(2011). Upslope migration of Andean trees. *Journal of Biogeography*, *38*(4), 783-791.

Freeman, B. G., Strimas-Mackey, M., & Miller, E. T. (2022). Interspecific competition limits
bird species' ranges in tropical mountains. *Science*, *377*(6604), 416-420.

Funk, V. A., Gostel, M., Devine, A., Kelloff, C. L., Wurdack, K., Tuccinardi, C., ... &
Coddington, J. (2017). Guidelines for collecting vouchers and tissues intended for genomic work
(Smithsonian Institution): Botany Best Practices. *Biodiversity Data Journal*, (5), e11625.

Gailing, O., & Curtu, A. L. (2014). Interspecific gene flow and maintenance of species integrity
in oaks. *Annals of Forest Research*, *57*(1), 5-18.

Gentry, A. H. (1982). Neotropical floristic diversity: phytogeographical connections between
Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean
orogeny?. *Annals of the Missouri Botanical Garden*, *69*(3), 557-593.

Grant, V. (1971). Plant Speciation. New York: Columbia University Press.

Green, R. E., Krause, J., Briggs, A. W., Maricic, T., Stenzel, U., Kircher, M., ... & Pääbo, S.
(2010). A draft sequence of the Neandertal genome. science, 328(5979), 710-722.

Griffiths, A. R., Silman, M. R., Farfan-Rios, W., Feeley, K. J., Cabrera, K. G., Meir, P., ... &
Dexter, K. G. (2021). Evolutionary diversity peaks at mid-elevations along an Amazon-to-Andes
elevation gradient. *Frontiers in ecology and evolution*, *9*, 680041.

Grubb, P. J. (1977). Control of forest growth and distribution on wet tropical mountains: with
special reference to mineral nutrition. *Annual Review of Ecology and Systematics*, 83-107.

Guerrero, B. I., Guerra, M. E., & Rodrigo, J. (2022). Simple sequence repeat (SSR)-based
genetic diversity in interspecific Plumcot-type (Prunus salicina× Prunus armeniaca)
hybrids. *Plants*, *11*(9), 1241.

Hamrick, J. L., & Murawski, D. A. (1990). The breeding structure of tropical tree
populations. *Plant species biology*, *5*(1), 157-165.

Hauck, N. R., Yamane, H., Tao, R., & Iezzoni, A. F. (2006). Accumulation of nonfunctional S-
haplotypes results in the breakdown of gametophytic self-incompatibility in tetraploid
Prunus. *Genetics*, *172*(2), 1191-1198.

Hillyer, R., & Silman, M. R. (2010). Changes in species interactions across a 2.5 km elevation gradient: effects on plant migration in response to climate change. *Global Change Biology*, *16*(12), 3205-3214.

Hodel, R. G., Zimmer, E., & Wen, J. (2021). A phylogenomic approach resolves the backbone of Prunus (Rosaceae) and identifies signals of hybridization and allopolyploidy. *Molecular Phylogenetics and Evolution*, *160*, 107118.

Hodel, R. G., Winslow, S. K., Liu, B. B., Johnson, G., Trizna, M., White, A. E., ... & Wen, J. (2023). A phylogenomic approach, combined with morphological characters gleaned via machine learning, uncovers the hybrid origin and biogeographic diversification of the plum genus. *bioRxiv*, 2023-09.

Hubbell, S. P. (2005). Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional ecology*, *19*(1), 166-172.

Hughes, C., & Eastwood, R. (2006). Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the Andes. *Proceedings of the National Academy of Sciences*, *103*(27), 10334-10339.

International Peach Genome Initiative, Verde, I., Abbott, A. G., Scalabrin, S., Jung, S., Shu, S., ... & Rokhsar, D. S. (2013). The high-quality draft genome of peach (Prunus persica) identifies unique patterns of genetic diversity, domestication, and genome evolution. *Nature genetics*, *45*(5), 487-494.

Janzen, D. H. (1967). Why mountain passes are higher in the tropics. *The American Naturalist*, *101*(919), 233-249.

Jankowski, J. E., Ciecka, A. L., Meyer, N. Y., & Rabenold, K. N. (2009). Beta diversity along environmental gradients: implications of habitat specialization in tropical montane landscapes. *Journal of Animal ecology*, *78*(2), 315-327.

Jankowski, J. E., Graham, C. H., Parra, J. L., Robinson, S. K., Seddon, N., Touchton, J. M., & Tobias, J. A. (2012). The role of competition in structuring tropical bird communities. *Ornitologia Neotropical*, *23*, 115-124.

Judd, E. J., Tierney, J. E., Lunt, D. J., Montañez, I. P., Huber, B. T., Wing, S. L., & Valdes, P. J. (2024). A 485-million-year history of Earth's surface temperature. *science*, *385*(6715), eadk3705.

Kulmuni, J., Wiley, B., & Otto, S. P. (2024). On the fast track: hybrids adapt more rapidly than parental populations in a novel environment. *Evolution Letters*, *8*(1), 128-136.

Lagomarsino, L. P., Condamine, F. L., Antonelli, A., Mulch, A., & Davis, C. C. (2016). The abiotic and biotic drivers of rapid diversification in A ndean bellflowers (Campanulaceae). *New Phytologist*, *210*(4), 1430-1442.

Larson, D. A., Vargas, O. M., Vicentini, A., & Dick, C. W. (2021). Admixture may be extensive
among hyperdominant Amazon rainforest tree species. *New Phytologist*, *232*(6), 2520-2534.

Lee, S., & Wen, J. (2001). A phylogenetic analysis of Prunus and the Amygdaloideae (Rosaceae)
using ITS sequences of nuclear ribosomal DNA. *American Journal of Botany*, *88*(1), 150-160.

Lepais, O., Petit, R. J., Guichoux, E., Lavabre, J. E., Alberto, F., Kremer, A., & Gerber, S. (2009).
Species relative abundance and direction of introgression in oaks. *Molecular Ecology*, *18*(10),
2228-2242.

Levin, D. A. (2012). Mating system shifts on the trailing edge. *Annals of Botany*, *109*(3), 613-
620.

Linan, A. G., Myers, J. A., Edwards, C. E., Zanne, A. E., Smith, S. A., Arellano, G., ... & Tello, J.
S. (2021). The evolutionary assembly of forest communities along environmental gradients:
recent diversification or sorting of pre-adapted clades?. *New Phytologist*, *232*(6), 2506-2519.

Lotsy, J. P. (1925). *Evolution considered in the light of hybridization*. Canterbury College.

Luebert, F., & Weigend, M. (2014). Phylogenetic insights into Andean plant
diversification. *Frontiers in Ecology and Evolution*, *2*, 27.

McCartney-Melstad, E., Gidiş, M., & Shaffer, H. B. (2019). An empirical pipeline for choosing
the optimal clustering threshold in RADseq studies. *Molecular Ecology Resources*, *19*(5), 1195-
1204.

Meier, C. L., Rapp, J., Bowers, R. M., Silman, M., & Fierer, N. (2010). Fungal growth on a
common wood substrate across a tropical elevation gradient: temperature sensitivity, community
composition, and potential for above-ground decomposition. *Soil Biology and*
Biochemistry, *42*(7), 1083-1090.

Mendelson, T. C., Inouye, B. D., & Rausher, M. D. (2004). Quantifying patterns in the evolution
of reproductive isolation. *Evolution*, *58*(7), 1424-1433.

Menon, M., Bagley, J. C., Friedline, C. J., Whipple, A. V., Schoettle, A. W., Leal-Sàenz, A., ... &
Eckert, A. J. (2018). The role of hybridization during ecological divergence of southwestern
white pine (Pinus strobiformis) and limber pine (P. flexilis). *Molecular Ecology*, *27*(5), 1245-
1260.

Murawski, D. A., & Hamrick, J. L. (1991). The effect of the density of flowering individuals on
the mating systems of nine tropical tree species. *Heredity*, *67*(2), 167-174.

Nason, J. D., & Hamrick, J. L. (1997). Reproductive and genetic consequences of forest
fragmentation: Two case studies of neotropical... *Journal of Heredity*, *88*(4).

Nevado, B., Atchison, G. W., Hughes, C. E., & Filatov, D. A. (2016). Widespread adaptive evolution during repeated evolutionary radiations in New World lupins. *Nature communications*, *7*(1), 12384.

Nürk, N. M., Madriñán, S., Carine, M. A., Chase, M. W., & Blattner, F. R. (2013). Molecular phylogenetics and morphological evolution of St. John's wort (Hypericum; Hypericaceae). *Molecular Phylogenetics and Evolution*, *66*(1), 1-16.

Pease, J. B., & Hahn, M. W. (2015). Detection and polarization of introgression in a five-taxon phylogeny. Systematic biology, 64(4), 651-662.

Pease, J. B., Haak, D. C., Hahn, M. W., & Moyle, L. C. (2016). Phylogenomics reveals three sources of adaptive variation during a rapid radiation. PLoS biology, 14(2), e1002379.

Pease, J. B., Brown, J. W., Walker, J. F., Hinchliff, C. E., & Smith, S. A. (2018). Quartet Sampling distinguishes lack of support from conflicting support in the green plant tree of life. *American journal of botany*, *105*(3), 385-403.

Pérez-Escobar, O. A., Chomicki, G., Condamine, F. L., Karremans, A. P., Bogarín, D., Matzke, N. J., ... & Antonelli, A. (2017). Recent origin and rapid speciation of Neotropical orchids in the world's richest plant biodiversity hotspot. *New Phytologist*, *215*(2), 891-905.

Pérez-Escobar, O. A., Zizka, A., Bermúdez, M. A., Meseguer, A. S., Condamine, F. L., Hoorn, C., ... & Chomicki, G. (2022). The Andes through time: evolution and distribution of Andean floras. *Trends in Plant Science*, *27*(4), 364-378.

Pérez Zabala, J. A. (2022). *Taxonomic Diversity, Phylogeny, and Diversification of the Environmental Niche of the Genus Prunus L. with emphasis on the New World Tropics* (Doctoral dissertation, UC Davis).

Pires, J. M., Dobzhansky, T. H., & Black, G. A. (1953). An estimate of the number of species of trees in an Amazonian forest community. *Botanical Gazette*, *114*(4), 467-477.

Pitman, N. C., Terborgh, J., Silman, M. R., & Nuñez V, P. (1999). Tree species distributions in an upper Amazonian forest. *Ecology*, *80*(8), 2651-2661.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945-959.

Quinlan, E. J., Layman, C. A., & Silman, M. R. (2025). Climate-Mediated Hybridisation and the Future of Andean Forests. *Journal of Biogeography*, *52*(6), e15113.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic biology*, *67*(5), 901-904.

Rapp, J. M., & Silman, M. R. (2012). Diurnal, seasonal, and altitudinal trends in microclimate across a tropical montane cloud forest. *Climate Research*, *55*(1), 17-32.

Rehder A (1940) Manual of cultivated trees and shrubs hardy in North America exclusive of the
subtropical and warmer temperate regions. Dioscorides, Portland, OR.

Sanín, M. J., Kissling, W. D., Bacon, C. D., Borchsenius, F., Galeano, G., Svenning, J. C., ... &
Pintaud, J. C. (2016). The Neogene rise of the tropical Andes facilitated diversification of wax
palms (Ceroxylon: Arecaceae) through geographical colonization and climatic niche
separation. *Botanical Journal of the Linnean Society*, *182*(2), 303-317.

Schley, R. J., Pennington, R. T., Pérez-Escobar, O. A., Helmstetter, A. J., de la Estrella, M.,
Larridon, I., ... & Klitgård, B. (2020). Introgression across evolutionary scales suggests
reticulation contributes to Amazonian tree diversity. *Molecular Ecology*, *29*(21), 4170-4185.

Schley, R. J., Twyford, A. D., & Pennington, R. T. (2022). Hybridization: a 'double-edged
sword' for Neotropical plant diversity. *Botanical Journal of the Linnean Society*, *199*(1), 331-
356.

Schley, R. J., Piñeiro, R., Nicholls, J. A., Pezzini, F. F., Farbos, A., Lewis, G. P., ... & Pennington,
R. T. (2025). Rampant Reticulation in a Rapid Radiation of Tropical Trees-Insights from Inga
(Fabaceae). *Systematic biology*, syaf027.

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *the american
naturalist*, *156*(S4), S4-S16.

Segovia, R. A., Pennington, R. T., Baker, T. R., Coelho de Souza, F., Neves, D. M., Davis, C. C.,
... & Dexter, K. G. (2020). Freezing and water availability structure the evolutionary diversity of
trees across the Americas. *Science Advances*, *6*(19), eaaz5373.

Sempere, T., Picard, D., & Plantard, O. (2005). Assessing and dating Andean uplift by
phylogeography and phylochronology: early Miocene emergence of Andean cloud forests. In *6th
International Symposium on Andean Geodynamics (ISAG 2005, Barcelona), Extended
Abstracts* (pp. 663-665). IRD Éditions.

Simpson, G. G. (1953). *The major features of evolution*. Columbia University Press.

Sokoloff, D. D., Ignatov, M. S., Remizowa, M. V., Nuraliev, M. S., Blagoderov, V., Garbout, A.,
& Perkovsky, E. E. (2018). Staminate flower of Prunus s. l.(Rosaceae) from Eocene Rovno
amber (Ukraine). *Journal of Plant Research*, *131*, 925-943.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.

Stroud, J. T., & Losos, J. B. (2016). Ecological opportunity and adaptive radiation. *Annual
Review of Ecology, Evolution, and Systematics*, *47*(1), 507-532.

Stull, G. W., Pham, K. K., Soltis, P. S., & Soltis, D. E. (2023). Deep reticulation: the long legacy
of hybridization in vascular plant evolution. *The Plant Journal*, *114*(4), 743-766.

Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. (2018). Adaptive introgression: a plant
perspective. *Biology letters*, *14*(3), 20170688.

Su, N., Hodel, R. G., Wang, X., Wang, J. R., Xie, S. Y., Gui, C. X., ... & Wen, J. (2023).
Molecular phylogeny and inflorescence evolution of Prunus (Rosaceae) based on RAD-seq and
genome skimming analyses. *Plant diversity*, *45*(4), 397-408.

Tao, R., & Iezzoni, A. F. (2010). The S-RNase-based gametophytic self-incompatibility system
in Prunus exhibits distinct genetic and molecular features. *Scientia Horticulturae*, *124*(4), 423-
433.

Ter Steege, H., Pitman, N. C., Sabatier, D., Baraloto, C., Salomão, R. P., Guevara, J. E., ... &
Silman, M. R. (2013). Hyperdominance in the Amazonian tree flora. *Science*, *342*(6156),
1243092.

Terborgh, J. (1971). Distribution on environmental gradients: theory and a preliminary
interpretation of distributional patterns in the avifauna of the Cordillera Vilcabamba,
Peru. *Ecology*, *52*(1), 23-40.

Terborgh, J. (1977). Bird species diversity on an Andean elevational gradient. *Ecology*, *58*(5),
1007-1019.

Tsuda, Y., Semerikov, V., Sebastiani, F., Vendramin, G. G., & Lascoux, M. (2017). Multispecies
genetic structure and hybridization in the Betula genus across Eurasia. *Molecular Ecology*, *26*(2),
589-605.

Urrego, D. H., Silman, M. R., & Bush, M. B. (2005). The Last Glacial Maximum: stability and
change in a western Amazonian cloud forest. *Journal of Quaternary Science: Published for the*
Quaternary Research Association, *20*(7-8), 693-701.

Van Schaik, C. P., Terborgh, J. W., & Wright, S. J. (1993). The phenology of tropical forests:
adaptive significance and consequences for primary consumers. *Annual Review of ecology and*
Systematics, 353-377.

Verde, I., Jenkins, J., Dondini, L., Micali, S., Pagliarani, G., Vendramin, E., ... & Schmutz, J.
(2017). The Peach v2. 0 release: high-resolution linkage mapping and deep resequencing
improve chromosome-scale assembly and contiguity. *BMC genomics*, *18*, 1-18.

Ward, M., Dick, C. W., Gribel, R., & Lowe, A. J. (2005). To self, or not to self… A review of
outcrossing and pollen-mediated gene flow in neotropical trees. Heredity, 95(4), 246-254.

Wen, J., Berggren, S. T., Lee, C. H., Ickert-Bond, S., Yi, T. S., Yoo, K. O., ... & Potter, D. (2008).
Phylogenetic inferences in Prunus (Rosaceae) using chloroplast ndhF and nuclear ribosomal ITS
sequences.

Williams Jr, J. H., Friedman, W. E., & Arnold, M. L. (1999). Developmental selection within the angiosperm style: using gamete DNA to visualize interspecific pollen competition. *Proceedings of the National Academy of Sciences*, *96*(16), 9201-9206.

Wu, C. I. (2001). The genic view of the process of speciation. *Journal of evolutionary biology*, *14*(6), 851-865.

Xie, S. Y., Lin, X. H., Wang, J. R., Ma, D. K., Su, N., Xu, C., ... & Liu, B. B. (2025). Unraveling evolutionary pathways: allopolyploidization and introgression in polyploid Prunus (Rosaceae). The Plant Journal, 123(1), e70320.

Zhao, L., Jiang, X. W., Zuo, Y. J., Liu, X. L., Chin, S. W., Haberle, R., ... & Wen, J. (2016). Multiple events of allopolyploidy in the evolution of the racemose lineages in Prunus (Rosaceae) based on integrated evidence from nuclear and plastid data. *PLoS One*, *11*(6), e0157123.

891 Table 1. Summary of significant D-statistic tests, with *P. myrtifolia* set to P4.

P1	P2	P3	dstat	Z	ABBA	BABA	nloci
P. antioquensis	*P. stipulata*	*P. debilis*	-0.69	4.75	8.38	45.13	152
P. stipulata	*P. antioquensis*	*P. debilis*	0.69	4.60	45.13	8.38	152
P. amplifolia, P. stipulata	*P. antioquensis*	*P. debilis*	0.66	4.16	43.19	8.94	163
P. antioquensis	*P. amplifolia, P. stipulata*	*P. debilis*	-0.66	4.01	8.94	43.19	163
P. stipulata	*P. pleiantha*	*P. detrita, P. debilis*	0.13	2.94	793	605.44	3968
P. pleiantha	*P. stipulata*	*P. detrita, P. debilis*	-0.13	2.87	605.44	793	3968
P. herthae, P. huantensis	*P. pleiantha*	*P. detrita, P. debilis*	0.11	2.81	864.08	686.51	4485
P. stipulata	*P. huantensis, P. antioquensis*	*P. pleiantha*	0.09	2.71	1190.20	989.74	6218
P. huantensis, P. antioquensis	*P. stipulata*	*P. pleiantha*	-0.09	2.71	989.74	1190.2	6218
P. pleiantha	*P. herthae, P. huantensis*	*P. detrita, P. debilis*	-0.11	2.67	686.51	864.08	4485
P. stipulata	*P. pleiantha*	*P. detrita*	0.13	2.66	744.38	578.5	3725
P. pleiantha	*P. antioquensis*	*P. debilis*	0.51	2.63	20.25	6.5	142
P. pleiantha	*P. stipulata*	*P. detrita*	-0.13	2.62	578.5	744.38	3725
P. antioquensis	*P. pleiantha*	*P. debilis*	-0.51	2.58	6.5	20.25	142
P. pleiantha	*P. stipulata*	*P. debilis*	-0.27	2.53	68.5	119.25	576
P. herthae, P. huantensis, P. antioquensis, P. brittoniana	*P. pleiantha*	*P. detrita*	0.11	2.52	813.45	654.78	4203

893 Table 2. Summary of within species genomic diversity statistics.

Species	# of samples	Filter	# of loci	Average π	Average H_o	Average H_e
P. integrifolia	10	50	53574	0.42 ± 0.13	0.15 ± 0.064	0.42 ± 0.0025
		70	17592	0.41 ± 0.14	0.17 ± 0.073	0.40 ±0.0015
P. huantensis/ brittoniana	12	50	24457	0.41 ± 0.13	0.12 ± 0.078	0.41 ± 0.0012
		70	1960	0.38 ± 0.15	0.16 ± 0.10	0.37 ± 0.0031
P. stipulata	7	50	24457	0.45 ± 0.11	0.095 ± 0.096	0.45 ± 0.0024
				0.42 ± 0.13	0.11 ± 0.11	0.42 ± 0.0064
P. pleiantha	7	50	23299	0.45 ± 0.11	0.12 ± 0.081	0.44 ± 0.0015
		70	7295	0.43 ± 0.12	0.13 ± 0.091	0.42 ± 0.0013
P. herthae	3	50	6251	0.62 ± 0.077	0.085 ± 0.03	0.61 ± 0.0067
		70	699	0.51 ± 0.073	0.064 ± 0.0072	0.51 ± 0
P. debilis/detrita	3	50	1747	0.64 ± 0.061	0.064 ± 0.058	0.064 ± 0.0009

895   Table 3. Summary of within-individual heterozygosity estimates.

Species	Individual	H_o (50% filter)	H_e (50% filter)	H_o (70% filter)	H_e (70% filter)
P. integrifolia	139_integrifolia	0.11	0.42	0.12	0.41
	147_integrifolia	0.31	0.42	0.36	0.4
	148_integrifolia	0.14	0.42	0.16	0.4
	149_integrifolia	0.18	0.42	0.21	0.4
	151_integrifolia	0.10	0.42	0.12	0.4
	155_integrifolia	0.16	0.42	0.19	0.4
	156_integrifolia	0.12	0.42	0.14	0.4
	157_integrifolia	0.17	0.42	0.2	0.4
	212_integrifolia	0.09	0.41	0.1	0.4
	253_integrifolia	0.12	0.42	0.14	0.41
P. huantensis /brittoniana	108_huantensis	0.16	0.41	0.23	0.37
	115_huantensis	0.1	0.41	0.13	0.38
	116_huantensis	0.22	0.41	0.29	0.37
	117_huantensis	0.044	0.4	0.07	0.37
	119_huantensis	0.072	0.41	0.093	0.37
	123_huantensis	0.069	0.4	0.12	0.37
	124_huantensis	0.2	0.4	0.25	0.37
	130_huantensis	0.05	0.41	0.079	0.38
	131_huantensis	0.15	0.41	0.22	0.37
	132_huantensis	0.26	0.41	0.36	0.37
	179_brittoniana	0.064	0.41	0.08	0.37
	210_brittoniana	0.03	0.4	0.05	0.38
P. stipulata	171_stipulata	0.28	0.45	0.32	0.42
	173_stipulata	0.17	0.45	0.18	0.42
	175_stipulata	0.07	0.45	0.07	0.41
	285_stipulata	0.058	0.45	0.1	0.42
	286_stipulata	0.03	0.45	0.041	0.41
	312_stipulata	0.037	0.45	0.032	0.42
	332_stipulata	0.02	0.45	0.021	0.43
P. pleiantha	162_pleiantha	0.12	0.44	0.13	0.43
	163_pleiantha	0.22	0.44	0.25	0.43
	164_pleiantha	0.18	0.44	0.19	0.42
	165_pleiantha	0.2	0.44	0.22	0.42
	166_pleiantha	0.023	0.44	0.021	0.42
	167_pleiantha	0.05	0.44	0.052	0.42
	169_pleiantha	0.051	0.44	0.056	0.42
P. herthae	226_herthae	0.087	0.62	0.057	0.51
	236_herthae	0.054	0.61	0.063	0.51
	237_herthae	0.11	0.62	0.072	0.51
P. debilis/ detrita	180_debilis	0.13	0.64	NA	NA
	225_debilis	0.036	0.64	NA	NA
	107_detrita	0.026	0.64	NA	NA

Figure 1. (a) Map of samples included in this study. Black dot indicates the location of the
ABERG transect in Manu National Park, Peru where most samples (n=33) were collected, also
shown in picture to the right. Red points indicate locations of additional samples included in the
study. Further details on individual samples provided in Table S1. (b) Box plots showing
elevational distributions of the in-group Andean species across three different sources. "Aberg"
includes all known individuals along the ABERG transect >10 dbh over 20 years. "Acha"
includes all Bolivian material reviewed by Achá (2013) and "Zabala" includes all material

reviewed by Pérez-Zabala (2022) from Venezuela to Bolivia. Pérez-Zabala considered *P.*
huantensis to be *P. brittoniana* so those samples are grouped together on the figure under *P.*
brittoniana. Figure does not include new "varieties" of species proposed by either Achá or Pérez-
Zabala.

Figure 2. Species-level topology resulting from RAxML and ExaBayes phylogenetic analyses.
Nodes show RAxML bootstrap support values where black circles indicate values > 75 and
values < 75 are shown on white nodes. All Bayesian posterior probabilities were equal to 1. The
tree is rooted to the reference (*P. persica*) and all three non-Andean species are shown in grey.
Dashed arrows summarize patterns of historical admixture inferred from D-statistics.

Figure 3. Time-calibrated phylogeny generated from BEAST analysis. Posterior probabilities are shown for nodes when < 1. Node
bars represent the 95% highest probability density (HPD) interval for node ages. Red stars indicate the fossil calibration points.

Figure 4. Patterns of genetic structure among individuals of sympatric species. Each vertical bar
represents a single individual and the y-axis indicates the proportion of genetic ancestry of that
individual assigned to different inferred genetic clusters, designated with different colors.
Species are ordered on the x-axis according to their elevational range, from low to high.
STRUCTURE detected K=9 distinct genetic clusters as the most likely grouping (Figure S3).

927   Table S1. Summary of samples included in individual-level phylogeny and genetic structure
928   analyses. Samples in bold were used to construct the species-level phylogeny.
929

Species determination	Sample ID	Voucher (Herbarium)	Material	Sampling locality
P. detrita	**107_detrita**	**EJQ82 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. debilis	**180_debilis**	**Loza et al. 388 (LPB)**	**Herbarium**	**Madidi NP, Bolivia**
P. debilis	225_debilis	JSB6352A (CUZ)	Fresh	Manu NP, Peru
P. stipulata	**171_stipulata**	**EJQ114 (LPB)**	**Fresh**	**Cotapata, Bolivia**
P. stipulata	173_stipulata	EJQ116 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	175_stipulata	EJQ118 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	285_stipulata	JSB6320 (CUZ)	Fresh	Manu NP, Peru
P. stipulata	286_stipulata	JSB6321(CUZ)	Fresh	Manu, NP, Peru
P. stipulata	312_stipulata	Cayola 3259	Herbarium	La Paz, Bolivia
P. stipulata	332_stipulata	Villalobosos 81	Herbarium	
P. integrifolia	139_integrifolia	JSB6366 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	**147_integrifolia**	**JSB6402 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. integrifolia	148_integrifolia	JSB6396 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	149_integrifolia	JSB6403 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	151_integrifolia	JSB6400 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	155_integrifolia	JSB6397 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	156_integrifolia	JSB6391 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	157_integrifolia	JSB6392 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	212_integrifolia	Meneses 7535 (LPB)	Herbarium	Nor Yungas, Cotapata, Bolivia
P. integrifolia	253_integrifolia	JSB7255 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	162_pleiantha	JSB6382 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	**163_pleiantha**	**JSB6388 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. pleiantha	164_pleiantha	JSB6386 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	165_pleiantha	JSB6387 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	166_pleiantha	JSB6385 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	167_pleiantha	JSB6380 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	169_pleiantha	JSB6390 (CUZ)	Fresh	Manu NP, Peru
P. herthae	226_herthae	JSB6343 (CUZ)	Fresh	Manu NP, Peru
P. herthae	236_herthae	JSB6333 (CUZ)	Fresh	Manu NP, Peru
P. herthae	**237_herthae**	**JSB6334 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. amplifolia	**206_amplifolia**	**Beck 33550 (LPB)**	**Herbarium**	**La Paz, Bolivia**
P. antioquensis	**177_antioquensis**	**Araujo & Canqui 348 (LPB)**	**Herbarium**	**Madidi, Bolivia**
P. huantensis	108_huantensis	JSB6316 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	115_huantensis	JSB6310 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	**116_huantensis**	**JSB6358 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. huantensis	117_huantensis	JSB6360 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	119_huantensis	JSB6362 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	123_huantensis	JSB6353 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	124_huantensis	JSB6357 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	130_huantensis	JSB6375 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	131_huantensis	JSB6311 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	132_huantensis	JSB6312 (CUZ)	Fresh	Manu NP, Peru

P. brittoniana	**179_brittoniana**	**Paniagua et al. 5782A (LPB)**	**Herbarium**	**Pelechuco-Mojo, Bolivia**
P. brittoniana	210_brittoniana	Beck 35044 (LPB)	Herbarium	Sud Yungas, Cotapata, Bolivia
Non-Andean:				
P. myrtifolia	**1_myrtifolia**	**Silica sample only**	**Fresh**	**Fairchild Botanical Garden, Miami, Florida, USA**
P. caroliniana	**2_caroliniana**	**Silica sample only**	**Fresh**	**Weymouth Woods, North Carolina, USA**

Figure S1. Maximum likelihood individual-level phylogeny reconstructed in RAxML.
Individuals were selected to include those with at least 1000 loci per sample and up to 10
individuals per species. Sites were filtered to only include those with at least 25% coverage
across samples. Tree was rooted to the reference (*P. persica*) and nodes show bootstrap support
value.

Figure S2. Quartet sampling results with quartet concordance (QC)/quartet differential
(QD)/quartet informativeness (QI) indicated next to nodes. Nodes are colored according to QC
scores, indicating how often the concordant quartet is inferred over both discordant quartets.
Outgroup samples *P. persica* and *P. caroliniana* are removed for clarity. Figure created with
https://github.com/ShuiyinLIU/QS_visualization.

Figure S3. Plot of estimators used to select the optimal number of genetic clusters (K) in the
STRUCTURE analysis. Estimated log mean probability assesses how well the model with a
particular K explains the genetic data. As the log-likelihood values are negative, the best-fit K-
value is the least negative. Delta K estimates the rate of change in the log probability of data
between successive K values. The K value corresponding to the highest delta K and estimated log
mean probability (K=9) was considered the best estimate of the number of subpopulations in this
dataset.