1 TITLE: UPSLOPE DIVERSIFICATION AND GENE FLOW AMONG CONGENERIC
2 ANDEAN TREES
3

4 Ellen J. Quinlan[1][*], James B. Pease[2], Jhonatan Sallo Bravo[3], Alfredo F. Fuentes[4,5], William
5 Farfan-Rios[1,6], and Miles R. Silman[1,6]
6
[1]Department of Biology, Wake Forest University, Winston-Salem, NC, USA 27109
[2]Department of Evolution, Ecology, and Organismal Biology, The Ohio State University,
Columbus, OH, USA 43210
[3]Facultad de Ciencias Biológicas, Universidad Nacional de San Antonio Abad del Cusco, Cusco,
Perú
[4]Latin America Department, Missouri Botanical Garden, St. Louis, MO, USA 63110
[5]Herbario Nacional de Bolivia, Universidad Mayor de San Andrés, La Paz, Bolivia
[6]The Sabin Center for Environment and Sustainability, Wake Forest University, Winston-Salem,
NC, USA 27109
16
17 *Corresponding Author: Ellen J. Quinlan, quinej18@wfu.edu

18   ABSTRACT

The high biodiversity of tropical Andean forests is characterized by many closely related

species which turnover rapidly across elevation and environmental gradients. Yet, the eco-

evolutionary relationships among these congeneric species remain poorly understood –

specifically, how they diversify, sort into their elevational niches, and the extent to which species

boundaries are maintained through time. In this study, we use the genus *Prunus* (cherries) as a

model to investigate these processes by (1) reconstructing the phylogenetic relationships and

timing of divergence among sympatric species and (2) assessing inter-specific gene flow and its

correspondence to evolutionary and ecological distance. Extensive population-level sampling

was conducted for species which co-occur along an elevational gradient in Manu National Park,

Peru. Additional samples were included for species which co-occur across a gradient in Madidi

National Park, Bolivia. Phylogenetic relationships reconstructed a pattern of upslope adaptive

diversification concurrent with Andean orogeny during the Miocene. However, despite these

ancient diversification events, gene flow analyses found species are not reproductively isolated,

and gene flow has occurred through time. Additionally, gene flow is widespread and does not

appear to be constrained by evolutionary or ecological distance. Together, these findings provide

insight into the processes that have shaped tree diversity in the tropical Andes and suggest that

during adaptive radiations, observable ecological, and morphological differences may emerge

long before complete phylogenomic isolation, even amid ongoing gene flow.

Keywords: Andes, biodiversity, diversification, elevation gradient, gene flow, hybridization,
tropical trees

                                    INTRODUCTION

Most ecological theory surrounding diversity and coexistence assumes species are

discrete evolutionary units (e.g. Hubbell 2000, Chesson 2000), though interspecific gene flow is

known to have been common throughout the evolutionary history of plants (reviewed by Stull et

al. 2023). No system lends itself to tests of these theories better than hyper-diverse tropical

forests. Until recently, hybridization was considered an extremely rare occurrence in tropical

trees, as hybrids were thought to be poor competitors in high-diversity systems and few

individuals with intermediate morphologies have been identified (e.g. Ashton 1969; Ehrendorfer

1970; Gentry 1982). Mounting genomic evidence suggests hybridization has been a key driver

of diversification in the Neotropical flora (reviewed by Schley et al. 2022), and may be an

important evolutionary mechanism for maintaining high tree diversity with low population

densities (Cannon and Lerdau 2015; Quinlan et al. 2025). Clarifying how species diversify and

assemble into communities, and the degree to which species boundaries persist through time is

essential for understanding the origin and maintenance of tropical tree biodiversity and its

conservation.

The tendency for inter-specific gene flow and hybridization to occur amongst closely

related, sympatric trees has been widely documented in temperate zones (e.g. oaks – Cavender-

Bares et al. 2018, spruce – De La Torre et al. 2014, birch – Tsuda et al. 2017, poplar – Chhatre et

al. 2018, pine – Menon et al. 2018), yet species cohesion and morphological similarity is often

maintained (Grant 1971; Gailing and Curtu 2014; Cannon and Petit 2019). This may be partially

due to assortative mating and variable mating behaviors (selfing, conspecific outcrossing, and

inter-specific gene flow), determined by population densities and pollinator abundance (Hamrick

and Murawski 1990; Ward et al. 2005; Cannon and Lerdau 2015). When present, conspecific

pollen can outcompete heterospecific pollen, reinforcing assortative mating (Williams et al.

1999). However, incomplete reproductive isolation and the ability to accept heterospecific pollen

may become adaptively beneficial when conspecific pollen is rare, as the benefit of avoiding

local extinction overrides potential negative consequences of hybridization such as reduced

hybrid fitness (Barton 2001; Wu 2001; Cannon and Lerdau 2015). Hybridization may be

particularly beneficial in times of rapid environmental change, as it can produce novel gene

combinations within a single generation, increasing adaptive potential and population resilience

(Anderson and Stebbins 1954; Baskett and Gomulkiewicz 2011; Suarez-Gonzalez 2018; Brauer

et al. 2023; Kulmuni et al. 2024). Such density-dependent mating has been reported in oaks

(Lepais et al. 2009), and may be particularly relevant to tropical trees which maintain extremely

low local population densities (Pires et al. 1953; Pitman et al. 1999; ter Steege et al. 2013).

Most tropical forest trees are highly outcrossed (Bawa et al. 1985a, b; Hamrick and

Murawski 1990; Ward et al. 2005), though individuals retain variable capacities for different

mating behaviors, even within populations and across flowering events (Murawski and Hamrick

1991; Nason and Hamrick 1997). Self-fertilization has been shown to be inversely correlated

with population density (Hamrick and Murawski 1990; Ward et al. 2005), but few studies have

assessed the extent of inter-specific gene flow. Those that have (e.g. Schley et al. 2020 –

Brownea, Larson et al. 2021 – *Eschweilera,* Schley et al. 2025 – *Inga*), detected wide-spread

admixture. Evolutionary distance may be one important constraint on inter-specific gene flow

among congeners in sympatry, as hybridization is thought to decrease with increasing divergence

due to the accumulation of pre- and post-zygotic barriers through time (Mendelson et al. 2004;

Levin 2012; Comeault and Matute 2018). Linan et al. (2021) found evidence of this in the

Mascarene Island *Diospyros* clade, suggesting the processes by which species diversify and

assemble into communities may have lasting effects on their eco-evolutionary dynamics. For

example, adaptive radiations across gradients typically result in step-wise diversification patterns

with sister taxa occurring closest in niche proximity (Simpson 1953; Schluter 2000; Stroud and

Losos 2016), whereas communities assembled predominantly via the immigration and sorting of

pre-adapted clades demonstrate greater phylogenetic distance among co-occurring taxa

(Donoghue 2008; Linan et al. 2021).

In tropical montane forests, genera commonly comprise many rare species which turn

over rapidly with elevation (Terborgh 1971, Gentry 1982). Turnover in the abiotic environment

and biotic interactions leads to narrow, shoestring-like distributions and the packing of many

closely related species across elevation gradients (e.g. Janzen 1967, Jankowski et el. 2009,

Hillyer and Silman 2010, Freeman et al. 2022). In the tropical Andes, this is true across taxa (e.g.

birds – Terborgh 1977; fungi – Meier et al. 2010; microbes – Nottingham et al. 2018; mammals –

Patterson et al. 2003, Mena and Pacheco 2020), and is especially true for forest trees (e.g.

Jankowski et al. 2012). While many tree species have elevation ranges of <500 m, the

distributions of entire genera are often much broader (Griffiths et al. 2020), with congeneric

species occurring in partial sympatry, especially at range margins. At the genus level,

immigration and sorting of pre-adapted clades has been shown to be the primary mechanism

structuring tree diversity across elevation gradients in the Andes (Griffiths et al. 2020; Segovia et

al. 2020; Linan et al. 2021). However, many examples of adaptive diversification have been

found within individual plant clades (e.g. Melastomataceae – Dudley 1978, *Espletia* – Pouchon

et al. 2018, lupines - Hughes and Eastwood 2006, Nevado et al. 2016, bellflowers – Lagomarsino

et al. 2016, *Hypericums* - Nörk et al. 2013, orchids - Pérez-Escobar et al. 2017, and bromeliads –

Givnish et al. 2011, Silvestro et al. 2014). Still, few studies have explored how congeners within

individual tree clades have diversified and assembled into their elevation niches, and

characterizing these evolutionary relationships is essential for understanding potential gene flow

and its ecological implications.

Prunus L. (Rosaceae) is one such genus that exhibits high congeneric turnover with

elevation in the Andes (Figure 1). Globally, it has diversified into a range of environments, and

though typically considered a temperate group (Wen et al. 2008; Chin et al. 2014), a recent

revision found its highest diversity to occur in the tropics and primarily the Neotropics (Pérez-

Zabala 2022). *Prunus* is one of the only tree genera with species spanning the entire Andes-

Amazon gradient, from tree-line (~3800 m) into the lowlands (<500 m), though the ranges of

individual species are typically 500-1000 m or less (Achá 2013). It is a keystone clade, whose

fruits are important in the diets of frugivorous birds, primates, Andean bears, and humans (Achá

2013). The genus also contains many globally important crop species (peach, plum, almond,

cherries) and others used as ornamentals, timber, and medicine (Andro and Riffaud 1995, Bortiri

et al. 2001, Lee and Wen 2001, Wen et al. 2008).

The genus *Prunus* is divided into three major clades that correspond to inflorescence type

– solitary, corymbose, and racemose (Rehder 1940, Su et al. 2023). All species endemic to the

Americas and tropical regions are racemose (Perez-Zabala 2022). Though historically

understudied (Hodel et al. 2021), the exclusively South American species form a strongly

supported clade nested within the tropical Racemose group (Hodel et al. 2023). Studies have

placed the origin of the genus in either North America (Chin et al. 2014) or East Asia (Hodel et

al. 2023) during the Paleocene (ca. 68-55 Ma), with the three major clades diverging by the

early-mid Eocene as part of the Boreotropical flora (Chin et al. 2014, Su et al. 2023, Hodel et al.

2023). Divergence in the tropical Racemose group occurred in several regions including Asia,

Africa, North America, and Europe ca. 55-40 Ma (Hodel et al. 2023). Chin et al. (2014)

hypothesized migration into South America occurred via the dispersal of a North American

relative ca. 34 Ma through Central America and the Caribbean. Hodel et al. (2023) found

evidence of a long period of stasis in the Neotropical lineages followed by rapid diversification

during the Miocene.

While most species in the solitary-flower and corymbose groups are diploids, the

racemose lineage is known for higher ploidy levels across temperate and tropical species and the

lineage is thought to be the result of a single or multiple ancient hybridization events during the

Eocene (Chin et al. 2104, Zhao et al. 2016, Su et al. 2023, Hodel et al. 2023). Domesticated

Prunus (particularly plums) readily hybridize in agricultural settings (Cici and Acker 2010),

resulting in the creation of some crops (e.g. *P. armeniaca* L. x *P. cerasifera* Ehrh. – 'black

apricots,' *P. salicina* x *P. armeniaca* Lindl. – plumcots, pluots, and apriums; Guerrero et al.

2022), as well as among cultivated varieties and wild conspecifics (*P. yedoensis* - Baek et al.

2018). Species in this genus are highly outcrossing and typically self-incompatible with

generalist insect pollinators (Cici and Acker 2010), though we are unaware of any study that has

specifically looked at either breeding systems or pollination in Neotropical *Prunus*.

Here, we use *Prunus* as a model to investigate the patterns, timing, and extent of

divergence among partially sympatric tree species which turn over across elevation gradients in

the tropical Andes, and the degree to which species boundaries are maintained through time.

Specifically, we ask: *(1) how and when co-occurring species diversified into their elevational*

niches, (2) whether species are reproductively isolated, and, if not, (3) is inter-specific gene flow

related to evolutionary or ecological distance. To do this, we reconstructed the phylogenetic

relationships among sympatric species, dated the timing of divergence, and tested for historical

introgression and recent admixture. The results of these analyses expand current understanding

of the origin and maintenance of tropical Andean tree diversity, and provide important context

for predicting how these species may respond to rapid environmental change.

157   MATERIALS AND METHODS

Study site and sample collection

Sampling for this study primarily focused on species which co-occur across the Andes

Biodiversity and Ecosystem Research Group (ABERG; andesbiodiversity.org) plot network in

Manu National Park, Peru (Figure 2). This site is located on the eastern slope of the tropical

Andes in southeastern Peru and is a 3.5 km elevational gradient from the Andean highlands to

the Amazonian lowlands. With 23-1 ha and 33-0.1 ha forest plots along a series of descending

ridges, this is the most continuous transect in the tropical Andes. To date, ~19,000 individual

trees have been documented in these plots from 100 families, 292 genera, and 1255 species

(Farfan-Rios 2019).

Intensive field sampling was conducted for *Prunus* species along this transect, with

botanists searching in lateral transects centered on the 1-ha plots (approximately every 250 m)

for up to 6 hrs. Leaf tissue was field collected and stored in silica following the Smithsonian

Institution guidelines for collecting vouchers and tissues intended for genomic work (Funk et al.

2017). Samples were also included for species known to co-occur across an additional plot

network spanning 4 km of elevation in the Madidi region of Bolivia known as the Madidi

Project, and from at least one external population when possible. These samples were collected

from herbarium specimens previously not treated with alcohol or fresh tissue stored in silica

collected from individuals in the Cotapata region of Bolivia (Table 1, Figure 2). In total, we

sampled 158 individuals from 13 species including two outgroups (*P. myrtifolia* and *P.*

caroliniana) and a reference (Table 1). See Appendix for species descriptions adapted from

Pérez-Zabala 2022 and Achá 2013.

ddRAD-seq data

High molecular weight DNA was extracted from ~50 mg of dried leaf tissue using Aboul-

Maaty and Oraby's (2019) modified CTAB protocol for non-model plants with a few

modifications: samples were ground with a bead-beater, incubated at -20 ℃ overnight (and up to

24 hrs), and the DNA pellet was washed twice with 500 µl 70% ethanol. DNA was checked for

quality and quantity on a 1.5% agarose gel and Qubit before being sent to Floragenex

(Beaverton, Oregon, USA) for ddRAD library prep and sequencing. Samples were ligated with 6

bp barcodes and digested with the PstI/MseI+0 enzyme pair. Genomic data were demultiplexed

and assembled to sample using ipyrad v. 0.9.95 (Eaton and Overcast 2020) with an average of

$2,215,444 \pm 1,419,356$ million reads (mean ± s.d.) recovered per sample.

Reads were assembled into orthologous loci using reference-based assembly in ipyrad.

Default settings were used to trim and filter reads and the clustering threshold was set to 0.85 as

recommended by the ipyrad manual and McCartney-Melstad et al. (2019) for phylogenetic

analyses. For the reference-based assembly, reads were mapped to the *Prunus persica* (peach)

genome (GCF_000346465.2; The International Peach Genome Initiative et al. 2013, Verde et al.

2017). The proportion of mapped reads varied across samples (mean = 67%, s.d. = 13%), but

variation in the quality of sample preservation and DNA extractions likely explain much of the

variation. When calling consensus alleles, we allowed for up to 4 alleles per sample following

Donoghue et al. (2022), as all *Prunus* species in the tropical Racemose group studied to date are

tetraploids (2n=32; Hodel et al. 2023), but ploidy is not known for most of the species in our

study. The final reference-mapped dataset comprised 35,099 loci, representing non-overlapping

genomic regions where data from at least 4 samples passed filtering. Further downstream

analyses were performed using vcftools and the ipyrad-analysis toolkit, which allows for filtering

and subsampling loci, genomic windows, and/or SNPs for phylogenetic inference and D-

statistics.

Individual-level phylogeny

A maximum likelihood (ML) phylogeny was reconstructed from a subset of

representative samples with the most data present per species (up to 10 individuals per species, at

least 1000 loci per sample; n=47), including conspecifics from multiple localities when possible.

The ipa.window_extracter module in the ipyrad-analysis toolkit was used to extract data from the

8 largest scaffolds (the chromosomes) and filter sites to only maintain those shared by at least

25% of individuals. We selected the 25% threshold after testing various other filtering thresholds

(10%, 30%, 40%, 50%), as 25% provided for the best tree resolution while also including as

much information as possible. The filtered and concatenated sequence matrix is 2.8 Mb in length,

containing 107,708 SNPs and 65.8% missing data. This alignment was used in RAxML

(Stamatakis 2014) under the GTR+GAMMA evolutionary model to infer a best tree under the

rapid hill-climbing algorithm, with support values from 100 bootstrap replicates. The tree was

rooted to the reference genome *P. persica* (a native of China and member of the Solitary-flower

group and Amygdalus subgenus), and included two additional Racemose outgroup species – *P.*

caroliniana (native of the eastern United States and Carribbean) and *P. myrtifolia* (native to the

Caribbean, Central America, and lowland South America).

Species-level phylogeny

To construct the species-level phylogeny, we selected one representative specimen for

each species in the individual-level tree (n=13). The samples selected grouped with other

members of their species on the individual-level tree and had the most data present. Again, we

used the ipa.window_extracter module in the ipyrad-analysis toolkit to extract data from the

chromosomes and filter sites, selecting those shared by at least 25% of all species (after testing

multiple filtering thresholds). The resulting sequence matrix is 4.2 Mb with 62,262 SNPs and

63.6% missing data. To reconstruct the phylogenetic relationships, we used both maximum

likelihood (RAxML) and Bayesian (ExaBayes) analyses. We ran RAxML using GTR+GAMMA

to infer a best tree under the rapid hill-climbing algorithm with support values from 100

bootstrap replicates. For ExaBayes, we ran two Metropolis-coupling replicates with four

coupled-chains (each with three heated chains) for 1×10^6 MCMC generations, sampled every

500 generations. Estimated sample size (ESS) for all parameters and branch lengths were

summarized with the ProcParam tool and confirmed as sufficiently sampled. The majority-rule

consensus tree was generated with the ExaBayes 'consense' tool with a 25% burn-in.

To assess the robustness of the resolved relationships and distinguish between conflicted

and poorly informed branches, we used Quartet Sampling (Pease et al. 2018;

https://www.github.com/ fephyfofum/quartetsampling). This method randomly samples groups

of four taxa from the provided tree and then evaluates each of the three possible unrooted

topologies to determine which best fits the data. The frequency at which each of these possible

topologies appears is recorded and then aggregated across quartets to determine whether they are

concordant or discordant with the larger tree topology. Discordance is quantified with four

metrics – quartet concordance (QC), quartet differential (QD), quartet informativeness (QI), and

quartet frequency (QF). We performed Quartet Sampling on the resulting RAxML tree with the

full concatenated alignment, allowing 100 replicates per branch (Figure S2).

Divergence time dating

To assess the timing of divergence among species, we generated a time-calibrated

phylogeny using BEAST v. 2.7.7 (Bouckaert et al., 2019) with the same alignment used for the

species-level phylogenies. We used an optimized relaxed molecular clock and Yule speciation

process with the GTR+GAMMA substitution model. A nodal calibration was set for the crown

age of *Prunus* using a normal prior with an offset of 61.5 Mya and a standard deviation of 3.0

Mya (Chin et al. 2014; Su et al. 2023). The crown age of the racemose group was calibrated from

a staminate flower of *Prunus hirsutipetala* D.D. Sokoloff, Remizowa et Nuraliev preserved in

the late Eocene (Priabonian) Rovno amber from northwestern Ukraine (Sokoloff et al. 2018),

using a normal prior with an offset of 35.0 Mya and standard deviation of 2.0 Mya (following Su

et al. 2023). To ensure a more rapid convergence of MCMC runs, we used our RAxML topology

as our starting tree. The Markov chain Monte Carlo (MCMC) run was conducted for 20,000,000

generations and trees sampled every 2000 generations producing a total of 10,000 trees. The

resulting log files were checked in Tracer v.1.5 (Rambaut et al. 2018) and we used TreeAnnotator

v.1.10.4 to generate the maximum clade credibility tree discarding 25% of trees as burn-in before

visualizing the results in FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Tests of historical introgression

To test for historical introgression, we used the four-taxon D-statistic (ABBA-BABA),

which quantifies allele frequency patterns that are incongruent with the hypothesized species tree

topology (Durand et al. 2011). Sub-sampling alleles across sets of three species and an outgroup

[((P1, P2), P3) O], the ancestral allele is denoted as "A" and the derived allele as "B". With this

topology there are two parsimony-informative patterns of discordant sites, of which the pattern

"ABBA" represents sites where P2 and P3 share the derived allele and the pattern "BABA"

represents sites where P1 and P3 share the derived allele (Hibbins and Hahn 2022). The

frequencies of ABBA and BABA site patterns are then summed across many genomic loci.

Under incomplete lineage sorting (ILS) these discordant patterns are expected to be equal (D=0),

and if the number of ABBA and BABA sites differ significantly this asymmetry is inferred as

evidence of introgression between the species sharing the derived state ("B") more frequently

(Hibbins and Hahn 2022). Here, we ran the ABBA-BABA tests using the ipa.baba tool as

implemented in ipyrad, which is specifically designed for analysis of RAD-seq datasets. We used

the species-level RAxML tree as our starting tree hypothesis and constrained the tests setting P4

to *P. myrtifolia*, so all tests would focus on ingroup species. To assess significance, we performed

1000 bootstrap replicates for each test with loci resampled with replacement. Z-scores were then

calculated as the number of bootstrap standard deviations that the D-stat deviates from zero

(Table 2). Z-scores > 2.5 were considered significant.

Tests of recent admixture

To assess genetic structure and test for recent admixture, we used the Bayesian clustering

algorithm STRUCTURE v. 2.3.4 (Pritchard et al. 2000) as implemented in ipyrad with the set of

samples included in the individual-level phylogeny (minus the outgroups). SNPs were filtered

conservatively to only include those with at least 50% coverage across individuals and then

randomly sampled to one SNP per locus to avoid linkage. This resulted in a total of 1493 SNPs

included in the analysis. The model was run with a burn-in of 125,000 and run-length of 500,000

generations, testing K=1-10 with 10 replicates per K. The assignment of samples into K distinct

genetic groups was assessed using DeltaK, which has been shown to be more robust to

accurately estimating the correct number of clusters in STRUCTURE, particularly with high

amounts of missing data as is common in RADseq (Evanno et al. 2005; Figure S3).

Estimates of genomic diversity

Genetic diversity was estimated within individuals and species to compare to the

admixture results. In VCFtools v.1.16 (Danecek et al. 2011), sequences were filtered using two

different missing data thresholds, retaining loci present within at least 50% or 70% of the

samples for each species. Sites were also filtered if they had a minor allele frequency less than

0.05. We calculated nucleotide diversity (π) for each species – the average pairwise nucleotide

differences between sequences in a sample at a site as well as observed (H_o) and expected

heterozygosity (H_e) – the proportion of loci where an individual is heterozygous vs. the

proportion expected by Hardy-Weinberg equilibrium – within samples and averaged across the

species. For these calculations, *P. debilis* and *P. detrita* samples were grouped together as well as

P. huantensis and *P. brittoniana,* given that STRUCTURE (Figure 5) did not detect any distinct

genetic clusters among these species' pairs. Species with <3 samples were excluded from these

analyses given the lack of statistical power from such a small sample size.

304   RESULTS

Phylogenomic relationships and elevational niche diversification

Both the species-level maximum likelihood (RAxML) and Bayesian (ExaBayes)

phylogenetic analyses reconstructed the same evolutionary relationships among the focal Andean

Prunus species and their outgroups (Figure 3). The trees were rooted to the reference (*P.*

persica), with *P. caroliniana* diverging before the tropical species and *P. myrtifolia* sister to the

Andean species, supporting the results of prior studies (Chin et al. 2014; Hodel et al. 2023).

Among these partially sympatric Andean species, diversification generally occurred from the

lowlands to the highlands, with the lineage including *P. debilis* and *P. detrita* diverging first,

followed by the mid-elevation cloud-forest species (*P. herthae, P. pleiantha, P. stipulata, P.*

integrifolia) and finally the high-elevation species as the most recent diversification events (*P.*

huantensis, P. brittoniana, and P. antioquensis). Statistical support for the inferred relationships

was assessed using bootstrap support (RAxML), posterior probabilities (ExaBayes) and Quartet

Sampling. While the tree nodes are generally well-supported (all posterior probabilities = 1, most

bootstrap values >75; Figure 3), Quartet Sampling (Pease et al. 2018) indicated that low

bootstrap support values are not likely the result of insufficient phylogenetic signal, as QI values

ranged from 0.8 – 1.0. Moreover, the lack a strong skew in discordance at most nodes (QD > 0.5)

suggests that incomplete lineage sorting (ILS) may instead be responsible (Pease et al. 2018).

One node resulted in both weak support and a discordant skew (0.21/0.22/1, *P. huantensis*, *P.*

brittoniana, *P. antioquensis; Figure S2*), indicating a possible secondary evolutionary history

(Pease et al. 2018).

Divergence time estimation

The time-calibrated phylogeny generated in BEAST recovered the same tree topology as

RAxML and ExaBayes with similar branch support (Figure 4). The estimated divergence time

for P. myrtifolia and the ancestor of the Amazonian and Andean species is 27.3 Ma (95% highest

posterior density [HPD] 22.4-31.7). The Andean species began diverging at the start of the

Miocene, with the low-elevation *P. debilis/P. detrita* lineage branching off 23.5 Ma (95% HPD

19.4-28.1), and from each other ~17.6 Ma (95% HPD 12.8-22.6). The cloud forest species

followed, with *P. pleiantha* diverging ~20.4 Ma (95% HPD 16.3-24.4), and *P. integrifolia* ~18.0

Ma (95% HPD 14.2-21.7). The remaining lineages emerged in the mid to late-Miocene (15.9-

9.3) with the highest elevation clade diverging last, with *P. huantensis* ~11.6 Ma (95% HPD 8.6-

14.9) and *P. antioquensis/P. brittoniana* ~9.3 Ma (95% HPD 6.2-12.3).

Tests of historical introgression and recent admixture

D-statistics indicated significant historical introgression events between *P. pleiantha* x *P.*

debilis/P. detrita, *P. pleiantha* x *P. huantensis/P. antioquensis*, and *P. debilis* x *P. antioquensis*

(Figure 3, Table 2). There is also evidence of introgression between *P. pleiantha* x *P. debilis* and

P. pleiantha x *P. detrita*, although it is difficult to disentangle this from introgression between *P.*

pleiantha and the ancestor of *P. debilis/P. detrita*. The introgression event detected between *P.*

pleiantha x *P. huantensis/ P. antioquensis* is further supported by the results of Quartet Sampling

(Figure S2).

STRUCTURE analysis found K=9 to be the most likely number of distinct genetic

clusters within the dataset, determined by mean log probability and DeltaK values (Figures 5,

S3). Individuals largely clustered according to their morphologically assigned species; however,

the number of detected clusters was fewer than the expected number of named taxonomic species

in the analysis (n=10). *P. integrifolia* and *P. amplifolia* are largely reproductively isolated, though

low levels of admixture were detected in *P. integrifolia* and only one sample was included for *P.*

amplifolia. Admixture is widespread amongst all other species, particularly *P. pleiantha*, *P.*

stipulata, *P. herthae, P. debilis, P. huantensis, P. brittoniana and P. antioquensis,* and is unrelated

to evolutionary or ecological distance.

Estimates of genomic diversity

Within each species, the individuals with the highest observed heterozygosity (H_o) were

also the most admixed in the STRUCTURE results (Tables 3 and 4, Figure 5), and this was

consistent with different amounts of missing data. For example, in *P. pleiantha*, individuals 163,

164, and 165 showed high levels of admixture from multiple groups and had the highest H_o

estimates (50%: 0.18 – 0.22, 70%: 0.19 – 0.25; Table 4). Across species and filtering thresholds,

nucleotide diversity (π) was high (50%: 0.41 – 0.64, 70%: 0.37 – 0.51) compared to H_o, π

tracked with H_e as expected, and H_o (50%: 0.06 – 0.15, 70%: 0.06 – 0.17) was far below H_e

(50%: 0.41 – 0.64; 70%: 0.37 – 0.51; Table 3). However, the estimates for *P. herthae* and the *P.*

debilis/detrita group should be interpreted cautiously given the low sample sizes for each.

363   DISCUSSION

364   The phylogenomic data presented here suggest that congeneric *Prunus* species known to

co-occur as elevational replacements in the tropical Andes diversified upslope from the lowlands

into the highlands during the Miocene (lower montane lineage ~23.5 Ma, cloud forest species

~20.4 – 13.4, high-elevation clade ~11.6 – 9.3 Ma) in a pattern consistent with adaptive

diversification. However, despite the ancient diversification events, species are not

reproductively isolated, as gene flow has occurred among species through time and is not related

to evolutionary or ecological distance. Together, these findings provide insight into the processes

that have shaped tree diversity across elevation gradients in the tropical Andes, and suggest that

during adaptive radiations, observable ecological and morphological differences may emerge

long before complete genomic isolation, even in the presence of ongoing gene flow.

Upslope diversification and biogeographic implications

375   Phylogenomic reconstructions and divergence time dating support the hypothesis that *Prunus*

diversified upslope, likely in response to novel habitat opportunities as has been observed in

other plant clades (e.g. Melastomataceae – Dudley 1978, *Espletia* – Pouchon et al. 2018, lupines

- Hughes and Eastwood 2006, Nevado et al. 2016, bellflowers – Lagomarsino et al. 2016,

Hypericums - Nörk et al. 2013, orchids - Pérez-Escobar et al. 2017, and bromeliads – Givnish et

al. 2011, 2014, Silvestro et al. 2014, Males 2018). Though this does not exclude the possibility

that some species may have adapted elsewhere and immigrated in once their preferred habitat

became available, the predominant pattern reconstructed from the phylogenomic analyses is one

of step-wise, upslope adaptive diversification. In agreement with prior studies, we found *P.*

myrtifolia (endemic to the Caribbean, Central America, and lowland South America) to be

ancestral to the rest of the Andean species included in this analysis, supporting the idea that this

lineage or its ancestor colonized South America via migration through the Caribbean and Central

America (Chin et al. 2014; Hodel et al. 2023). While our divergence time estimate for *P.*

myrtifolia of ~27 Ma is slightly later than the Neotropical colonization of ~34 Ma predicted by

Chin et al. (2014), it is still within the climatic shift from warmhouse to transitional state that

coincided with an onset of glaciation in Antarctica (~33.9 – 27.29 Ma; Judd et al. 2024) and the

estimated end of the boreotropical forests in the Northern Hemisphere.

In our analyses, diversification of the Andean species occurred throughout the Miocene,

beginning with the lineage that includes two of the lowest elevation species (*P. debilis* and *P.*

detrita) ~23.5 Ma and ending with the high-elevation clade ~11.6 Ma (*P. huantensis*) - ~9.3 Ma

(*P. antioquensis* and *P. brittoniana*). This is consistent with the timing of major uplift events and

estimated paleo-elevations in the North-Central and Central Andes (Boschman 2021). It also

supports the findings of Hodel et al. (2023), which detected a long period of stasis followed by

rapid diversification of Neotropical *Prunus* during the Miocene. Although the exact timing, pace,

and sequence of Andean orogeny is a subject of ongoing study and is heterogeneous across

space, the late Oligocene and Miocene (~25-5 Ma) are known to have been a period of intense

and accelerated uplift in the Eastern Cordillera of the North-Central and Central Andes

(Boschman et al. 2021; Pérez-Escobar et al. 2022).

Andean cloud forests are distinct habitats that occur from ~1200 m to the upper forest limits

at ~3200-3800 m (depending on latitude), and are characterized by persistent cloud cover and

immersion. Although they are the most speciose habitats in the Andes (Pérez-Escobar 2022),

persistence in these environments often requires unique adaptations to cope with the reduced

light availability, high precipitation, and humidity such as thick, pachyphyllic leaves, low growth

rates, and ability to withstand heavy accumulations of epiphytes (Grubb 1977; Rapp and Silman

2012; Fahey et al. 2016). The lower montane cloud forest species in our analysis (median

elevation ranges 1200 - 2500 m; *P. pleiantha, P. integrifolia, P. herthae, P. stipulata*) diverged

during the early to mid-Miocene. This aligns with the predicted origination of cloud forests

(Sempere et al. 2005) and the estimated divergence times for other cloud forest lineages (Luebert

and Weigend 2014). The highest elevation species in our study (median elevation ranges ≥3000

m; *P. huantensis, P. antioquensis, P. brittoniana*) diverged during the late Miocene, as the final

uplift phase began in the North-Central and Central Andes (Boschman et al. 2021; Pérez-Escobar

2022). Species in this elevation zone are exposed to unique high-elevation stressors such as night

frost, with temperatures dipping below 0°C, requiring further physiological specialization (Pérez-

Escobar 2022).

Gene flow through time

Analyses also indicated that these partially sympatric, congeneric trees are not

reproductively isolated. Introgression and admixture have occurred throughout the evolutionary

history of the Andean *Prunus* clade. D-statistics detected at least five significant ancient

introgression events. Two of these events occurred among lineages separated by great

evolutionary and ecological distance (*P. pleiantha* x *P. huantensis/P. antioquensis*, and *P. debilis*

x *P. antioquensis*). Introgression was also detected between *P. pleiantha* and the ancestor of *P.*

debilis/P. detrita (*P. pleiantha* x *P. debilis/P. detrita*) as well as among *P. pleiantha* x *P. debilis*

and *P. pleiantha* x *P. detrita* individually. Although difficult to disentangle, especially with few

samples and loci, it suggests that interspecific hybridization occurred among the ancestors of

these lineages and has continued as they diverged.

Historical introgression among species separated in ecological space across the elevation

gradient is plausible for several reasons. While the medians of their elevation ranges may be

quite different, they are found in close geographic proximity, and their ranges do overlap (or are

much closer) at the range margins. *Prunus* species are thought to be obligate outcrossers with

generalist pollinators, as is common in tropical trees (Ashton 1969; Bawa 1974, 1979), though

the extent of self-incompatibility across the genus and its potential breakdown with

polyploidization is still unclear (Hauck et al. 2006, Tao et al. 2010, Aguiar et al. 2015). As some

generalist pollinators forage over kilometers (Bawa et al. 1985), opportunity for introgression is

possible, especially given that many Andean species flower semi-synchronously during the dry

season. Additionally, species ranges are not stagnant through time. Pollen records from Lago

Consuelo (1360 m a.s.l., located in southeastern Peru adjacent to the Bolivian border) show that

during the last full glacial period (43.5 – 22,000 yr BP), individual cloud forest species migrated

downslope at least 1000 – 1700 vertical m in response to lower cloud base and cooler

temperatures (-5°C; Urrego et al. 2005). Today, these species are as far apart as they have likely

been during the Quaternary, thus these historical introgression events may be signatures of past

ecological overlap.

Tests of recent admixture also found gene flow to be widespread among Andean *Prunus*

species. Though *P. integrifolia* and *P. amplifolia* are (to an extent) genomically isolated, gene

flow is rampant among all other species. The isolation of *P. integrifolia* is particularly interesting

as this species has the widest elevational range and is one of the most abundant species in the

tropical Andes. Gene flow was especially high among *P. pleiantha, P. stipulata*, and *P. herthae*,

all cloud forest species with largely overlapping elevation ranges. However, these species did

mostly form monophyletic groups in our individual-level phylogeny (Figure S1), indicating that

despite gene flow, these are distinct evolutionary lineages. Given that *P. pleiantha* was identified

in most of the historical introgression events and those events are further supported by the

genetic clusters detected by STRUCTURE, it seems this species readily hybridizes when in

contact with close relatives. Further investigation is needed to test this and identify the

directionality of such gene flow. The algorithm did not distinguish *P. debilis* and *P. detrita* as

distinct genetic clusters, possibly due to how closely related these species are, introgression, and

the low number of individuals for both species in the dataset. Similarly, *P. huantensis* and *P.*

brittoniana are indistinguishable, which is unsurprising given that *P. brittoniana* was nested

within *P. huantensis* in the individual-level phylogeny (Figure S1). This supports the most recent

taxonomic revision by Pérez-Zabala (2022) which considered *P. brittoniana* and *P. huantensis* to

be the same species given their morphological and ecological similarities. Finally, the genetic

cluster indicated in orange (Figure 5) is the only cluster that does not appear to be dominant in

any one species, possibly representative of a co-occurring species not captured in our sampling

or a ghost ancestral lineage (Ottenburghs 2020; Tricou et al. 2022).

Estimates of genomic diversity further support this evidence of recent admixture, as the

individuals in each species that appear the most admixed in STRUCTURE also have the highest

observed heterozygosity (H_o). *P. integrifolia* also shows moderate H_o but it is more consistent

across individuals, likely a result of its larger population sizes. Additionally, nucleotide diversity

(π) is moderate to high and expected heterozygosity exceeds observed ($H_e \gg H_o$) across species,

a signal of a possible historical bottleneck. This is because during bottlenecks, H_o declines much

faster than π and recovers much slower, as H_o is more sensitive to allele frequency changes (Nei

et al. 1975; Chakroborty and Nei 1977). Future work will aim to test this.

Conclusions

Overall, these results suggest that adaptive diversification following the emergence of

novel gradients has been important for within-genus diversification of Andean trees, particularly

in the genus *Prunus*, while immigration and sorting of pre-adapted clades may be the dominant

mechanism shaping community assembly at higher-level taxonomic levels (Segovia et al. 2020,

Griffiths et al. 2020, Linan et al. 2021). For most of the Neogene, *Prunus* diversified with the

rise of the Andes, developing sequential elevational replacements with narrow ranges, specially

adapted to live in radically different environmental conditions from the Amazonian lowlands to

Andean tree line. This has important implications in the context of climate change, as species are

now shifting their ranges to track their preferred temperatures (Feeley et al. 2011; Fadrique et al.

2018). Though most species will not be able to migrate fast enough to keep pace with climatic

shifts and are instead experiencing range retractions at the warmer-end (Duque et al. 2015;

Farfan-Rios et al. 2025 – *in review*), any upslope range shift may increasingly result in range

overlap among congeneric species and potential gene flow (e.g. Quinlan et al. 2025)

Here, we found evidence of both historical introgression (possibly in response to

Quaternary range shifts) and widespread admixture among co-occurring *Prunus* species. Yet

despite gene flow, these species have maintained distinct evolutionary lineages with identifiable

ecological and morphological differences since the Miocene. This aligns with the syngameon

model, wherein species maintain incomplete reproductive barriers but primarily experience

divergent selection (Lotsy 1925; Grant 1971). Though oaks are the classic example (e.g. Cannon

and Petit 2020), such genomic mutualism may be an important yet under-appreciated mechanism

in the maintenance of diversity in hyper-diverse tropical forests, where individuals

predominantly outcross but retain some capacity for self-fertilization and interspecific

hybridization at low population densities (Wu 2001; Cannon and Lerdau 2015). Further work is

required to understand the prevalence of syngameons in tropical forests and their role in species'

responses to rapid environmental change, namely resilience and avoidance of local extinction.

Incorporating these complex dynamics into demographic models may be important for

accurately predicting the futures of Andean forests and effective conservation and management

under current climate change.

505                REFERENCES

506 Aboul-Maaty, N. A. F., & Oraby, H. A. S. (2019). Extraction of high-quality genomic DNA from
507 different plant orders applying a modified CTAB-based method. *Bulletin of the National*
508 *Research Centre*, *43*(1), 1-10.

509 Achá, S. (2013). El género Prunus (Rosaceae) en Bolivia: Novedades en su sistemática.

510 Aguiar, B., Vieira, J., Cunha, A. E., Fonseca, N. A., Iezzoni, A., van Nocker, S., & Vieira, C. P.
511 (2015). Convergent evolution at the gametophytic self-incompatibility system in Malus and
512 Prunus. *PloS one*, *10*(5), e0126138.

513 Anderson, E., & Stebbins Jr, G. L. (1954). Hybridization as an evolutionary stimulus. *Evolution*,
514 378-388.

515 Andro, M. C., & Riffaud, J. P. (1995). Pygeum africanum extract for the treatment of patients
516 with benign prostatic hyperplasia: a review of 25 years of published experience. *Current*
517 *therapeutic research*, *56*(8), 796-817.

518 Ashton, P. S. (1969). Speciation among tropical forest trees: some deductions in the light of
519 recent evidence. *Biological journal of the Linnean Society*, *1*(1-2), 155-196.

520 Baek, S., Choi, K., Kim, G. B., Yu, H. J., Cho, A., Jang, H., ... & Mun, J. H. (2018). Draft
521 genome sequence of wild Prunus yedoensis reveals massive inter-specific hybridization between
522 sympatric flowering cherries. *Genome biology*, *19*, 1-17.

523 Barton, N. H. (2001). The role of hybridization in evolution. *Molecular ecology*, *10*(3), 551-568.

524 Baskett, M. L., & Gomulkiewicz, R. (2011). Introgressive hybridization as a mechanism for
525 species rescue. *Theoretical Ecology*, *4*, 223-239.

526 Bawa, K. S. (1974). Breeding systems of tree species of a lowland tropical community.
527 Evolution, 85-92.

528 Bawa, K. S. (1979). Breeding systems of trees in a tropical wet forest. *New Zealand Journal of*
529 *Botany*, *17*(4), 521-524.

530 Bawa, K. S., Perry, D. R., & Beach, J. H. (1985). Reproductive biology of tropical lowland rain
531 forest trees. I. Sexual systems and incompatibility mechanisms. *American journal of*
532 *botany*, *72*(3), 331-345.

533 Bawa, K. S., Bullock, S. H., Perry, D. R., Coville, R. E., & Grayum, M. H. (1985). Reproductive
534 biology of tropical lowland rain forest trees. II. Pollination systems. *American journal of*
535 *botany*, *72*(3), 346-356.

536 Boschman, L. M. (2021). Andean mountain building since the Late Cretaceous: A paleoelevation
537 reconstruction. *Earth-Science Reviews*, *220*, 103640.

Bortiri, E., Oh, S. H., Jiang, J., Baggett, S., Granger, A., Weeks, C., ... & Parfitt, D. E. (2001).
Phylogeny and systematics of Prunus (Rosaceae) as determined by sequence analysis of ITS and
the chloroplast trnL-trnF spacer DNA. *Systematic Botany*, *26*(4), 797-807.

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M., Gavryushkina, A.,
... & Drummond, A. J. (2019). BEAST 2.5: An advanced software platform for Bayesian
evolutionary analysis. *PLoS computational biology*, *15*(4), e1006650.

Brauer, C. J., Sandoval-Castillo, J., Gates, K., Hammer, M. P., Unmack, P. J., Bernatchez, L., &
Beheregaray, L. B. (2023). Natural hybridization reduces vulnerability to climate change. *Nature*
Climate Change, *13*(3), 282-289.

Cannon, C. H., & Lerdau, M. (2015). Variable mating behaviors and the maintenance of tropical
biodiversity. *Frontiers in Genetics*, *6*, 183.

Cannon, C. H., & Petit, R. J. (2020). The oak syngameon: more than the sum of its parts. *New*
Phytologist, *226*(4), 978-983.

Cavender-Bares, J., Kothari, S., Meireles, J. E., Kaproth, M. A., Manos, P. S., & Hipp, A. L.
(2018). The role of diversification in community assembly of the oaks (Quercus L.) across the
continental US. *American Journal of Botany*, *105*(3), 565-586.

Chakraborty, R., & Nei, M. (1977). Bottleneck effects on average heterozygosity and genetic
distance with the stepwise mutation model. *Evolution*, 347-356.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annual review of Ecology*
and Systematics, *31*(1), 343-366.

Chhatre, V. E., Evans, L. M., DiFazio, S. P., & Keller, S. R. (2018). Adaptive introgression and
maintenance of a trispecies hybrid complex in range-edge populations of Populus. *Molecular*
ecology, *27*(23), 4820-4838.

Chin, S. W., Shaw, J., Haberle, R., Wen, J., & Potter, D. (2014). Diversification of almonds,
peaches, plums and cherries–molecular systematics and biogeographic history of Prunus
(Rosaceae). Molecular phylogenetics and evolution, 76, 34-48.

Cici, S. Z. H., & Van Acker, R. C. (2010). Gene flow in Prunus species in the context of novel
trait risk assessment. *Environmental biosafety research*, *9*(2), 75-85.

Comeault, A. A., & Matute, D. R. (2018). Genetic divergence and the number of hybridizing
species affect the path to homoploid hybrid speciation. *Proceedings of the National Academy of*
Sciences, *115*(39), 9761-9766.

Danecek, Petr, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A.
DePristo, Robert E. Handsaker et al. "The variant call format and VCFtools." *Bioinformatics* 27,
no. 15 (2011): 2156-2158.

De La Torre, A. R., Roberts, D. R., & Aitken, S. N. (2014). Genome-wide admixture and ecological niche modelling reveal the maintenance of species boundaries despite long history of interspecific gene flow. Molecular Ecology, 23(8), 2046-2059.

Donoghue, M. J. (2008). A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences*, *105*(supplement_1), 11549-11555.

Donoghue, M. J., Eaton, D. A., Maya-Lastra, C. A., Landis, M. J., Sweeney, P. W., Olson, M. E., ... & Edwards, E. J. (2022). Replicated radiation of a plant clade along a cloud forest archipelago. *Nature Ecology & Evolution*, *6*(9), 1318-1329.

Duque, A., Stevenson, P. R., & Feeley, K. J. (2015). Thermophilization of adult and juvenile tree communities in the northern tropical Andes. *Proceedings of the National Academy of Sciences*, *112*(34), 10744-10749.

Dudley, E. C. (1978). Adaptive radiation in the Melastomataceae along an altitudinal gradient in Peru. *Biotropica*, 134-143.

Durand, E. Y., Patterson, N., Reich, D., & Slatkin, M. (2011). Testing for ancient admixture between closely related populations. *Molecular biology and evolution*, *28*(8), 2239-2252.

Eaton, D. A., & Overcast, I. (2020). ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics*, *36*(8), 2592-2594.

Ehrendorfer, F. (1970). Evolutionary patterns and strategies in seed plants. *Taxon*, *19*(2), 185-195.

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular ecology*, *14*(8), 2611-2620.

Fahey, T. J., Sherman, R. E., & Tanner, E. V. (2016). Tropical montane cloud forest: environmental drivers of vegetation structure and ecosystem function. *Journal of Tropical Ecology*, *32*(5), 355-367.

Fadrique, B., Báez, S., Duque, Á., Malizia, A., Blundo, C., Carilla, J., ... & Feeley, K. J. (2018). Widespread but heterogeneous responses of Andean forests to climate change. *Nature*, *564*(7735), 207-212.

Farfan-Rios, W. (2019). *Forest responses to climate change along an Andes-to-Amazon elevational gradient*. Wake Forest University.

Farfan-Rios et al. 2025 – in press

Feeley, K. J., Silman, M. R., Bush, M. B., Farfan, W., Cabrera, K. G., Malhi, Y., ... & Saatchi, S. (2011). Upslope migration of Andean trees. *Journal of Biogeography*, *38*(4), 783-791.

Freeman, B. G., Strimas-Mackey, M., & Miller, E. T. (2022). Interspecific competition limits bird species' ranges in tropical mountains. *Science*, *377*(6604), 416-420.

Funk, V. A., Gostel, M., Devine, A., Kelloff, C. L., Wurdack, K., Tuccinardi, C., ... & Coddington, J. (2017). Guidelines for collecting vouchers and tissues intended for genomic work (Smithsonian Institution): Botany Best Practices. *Biodiversity Data Journal*, (5), e11625.

Gailing, O., & Curtu, A. L. (2014). Interspecific gene flow and maintenance of species integrity in oaks. *Annals of Forest Research*, *57*(1), 5-18.

Gentry, A. H. (1982). Neotropical floristic diversity: phytogeographical connections between Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny?. *Annals of the Missouri Botanical Garden*, *69*(3), 557-593.

Givnish, T. J., Barfuss, M. H., Van Ee, B., Riina, R., Schulte, K., Horres, R., ... & Sytsma, K. J. (2011). Phylogeny, adaptive radiation, and historical biogeography in Bromeliaceae: insights from an eight-locus plastid phylogeny. *American journal of botany*, *98*(5), 872-895.

Grant, V. (1971). Plant Speciation. New York: Columbia University Press.

Griffiths, A. R., Silman, M. R., Farfan-Rios, W., Feeley, K. J., Cabrera, K. G., Meir, P., ... & Dexter, K. G. (2021). Evolutionary diversity peaks at mid-elevations along an Amazon-to-Andes elevation gradient. *Frontiers in ecology and evolution*, *9*, 680041.

Grubb, P. J. (1977). Control of forest growth and distribution on wet tropical mountains: with special reference to mineral nutrition. *Annual Review of Ecology and Systematics*, 83-107.

Guerrero, B. I., Guerra, M. E., & Rodrigo, J. (2022). Simple sequence repeat (SSR)-based genetic diversity in interspecific Plumcot-type (Prunus salicina× Prunus armeniaca) hybrids. *Plants*, *11*(9), 1241.

Hamrick, J. L., & Murawski, D. A. (1990). The breeding structure of tropical tree populations. *Plant species biology*, *5*(1), 157-165.

Hauck, N. R., Yamane, H., Tao, R., & Iezzoni, A. F. (2006). Accumulation of nonfunctional S-haplotypes results in the breakdown of gametophytic self-incompatibility in tetraploid Prunus. *Genetics*, *172*(2), 1191-1198.

Hibbins, M. S., & Hahn, M. W. (2022). Phylogenomic approaches to detecting and characterizing introgression. *Genetics*, *220*(2), iyab173.

Hillyer, R., & Silman, M. R. (2010). Changes in species interactions across a 2.5 km elevation gradient: effects on plant migration in response to climate change. *Global Change Biology*, *16*(12), 3205-3214.

Hodel, R. G., Zimmer, E., & Wen, J. (2021). A phylogenomic approach resolves the backbone of
Prunus (Rosaceae) and identifies signals of hybridization and allopolyploidy. *Molecular*
Phylogenetics and Evolution, *160*, 107118.

Hodel, R. G., Winslow, S. K., Liu, B. B., Johnson, G., Trizna, M., White, A. E., ... & Wen, J.
(2023). A phylogenomic approach, combined with morphological characters gleaned via machine
learning, uncovers the hybrid origin and biogeographic diversification of the plum
genus. *bioRxiv*, 2023-09.

Hubbell, S. P. (2005). Neutral theory in community ecology and the hypothesis of functional
equivalence. *Functional ecology*, *19*(1), 166-172.

Hughes, C., & Eastwood, R. (2006). Island radiation on a continental scale: exceptional rates of
plant diversification after uplift of the Andes. *Proceedings of the National Academy of*
Sciences, *103*(27), 10334-10339.

International Peach Genome Initiative, Verde, I., Abbott, A. G., Scalabrin, S., Jung, S., Shu, S., ...
& Rokhsar, D. S. (2013). The high-quality draft genome of peach (Prunus persica) identifies
unique patterns of genetic diversity, domestication, and genome evolution. *Nature*
genetics, *45*(5), 487-494.

Janzen, D. H. (1967). Why mountain passes are higher in the tropics. *The American*
Naturalist, *101*(919), 233-249.

Jankowski, J. E., Ciecka, A. L., Meyer, N. Y., & Rabenold, K. N. (2009). Beta diversity along
environmental gradients: implications of habitat specialization in tropical montane
landscapes. *Journal of Animal ecology*, *78*(2), 315-327.

Jankowski, J. E., Graham, C. H., Parra, J. L., Robinson, S. K., Seddon, N., Touchton, J. M., &
Tobias, J. A. (2012). The role of competition in structuring tropical bird
communities. *Ornitologia Neotropical*, *23*, 115-124.

Judd, E. J., Tierney, J. E., Lunt, D. J., Montañez, I. P., Huber, B. T., Wing, S. L., & Valdes, P. J.
(2024). A 485-million-year history of Earth's surface temperature. *science*, *385*(6715), eadk3705.

Kulmuni, J., Wiley, B., & Otto, S. P. (2024). On the fast track: hybrids adapt more rapidly than
parental populations in a novel environment. *Evolution Letters*, *8*(1), 128-136.

Lagomarsino, L. P., Condamine, F. L., Antonelli, A., Mulch, A., & Davis, C. C. (2016). The
abiotic and biotic drivers of rapid diversification in A ndean bellflowers (Campanulaceae). *New*
Phytologist, *210*(4), 1430-1442.

Larson, D. A., Vargas, O. M., Vicentini, A., & Dick, C. W. (2021). Admixture may be extensive
among hyperdominant Amazon rainforest tree species. *New Phytologist*, *232*(6), 2520-2534.

Lee, S., & Wen, J. (2001). A phylogenetic analysis of Prunus and the Amygdaloideae (Rosaceae)
using ITS sequences of nuclear ribosomal DNA. *American Journal of Botany*, *88*(1), 150-160.

Lepais, O., Petit, R. J., Guichoux, E., Lavabre, J. E., Alberto, F., Kremer, A., & Gerber, S. (2009).
Species relative abundance and direction of introgression in oaks. *Molecular Ecology*, *18*(10),
2228-2242.

Levin, D. A. (2012). Mating system shifts on the trailing edge. *Annals of Botany*, *109*(3), 613-
620.

Linan, A. G., Myers, J. A., Edwards, C. E., Zanne, A. E., Smith, S. A., Arellano, G., ... & Tello, J.
S. (2021). The evolutionary assembly of forest communities along environmental gradients:
recent diversification or sorting of pre-adapted clades?. *New Phytologist*, *232*(6), 2506-2519.

Lotsy, J. P. (1925). *Evolution considered in the light of hybridization*. Canterbury College.

Luebert, F., & Weigend, M. (2014). Phylogenetic insights into Andean plant
diversification. *Frontiers in Ecology and Evolution*, *2*, 27.

McCartney-Melstad, E., Gidiş, M., & Shaffer, H. B. (2019). An empirical pipeline for choosing
the optimal clustering threshold in RADseq studies. *Molecular Ecology Resources*, *19*(5), 1195-
1204.

Meier, C. L., Rapp, J., Bowers, R. M., Silman, M., & Fierer, N. (2010). Fungal growth on a
common wood substrate across a tropical elevation gradient: temperature sensitivity, community
composition, and potential for above-ground decomposition. *Soil Biology and
Biochemistry*, *42*(7), 1083-1090.

Mendelson, T. C., Inouye, B. D., & Rausher, M. D. (2004). Quantifying patterns in the evolution
of reproductive isolation. *Evolution*, *58*(7), 1424-1433.

Menon, M., Bagley, J. C., Friedline, C. J., Whipple, A. V., Schoettle, A. W., Leal-Sàenz, A., ... &
Eckert, A. J. (2018). The role of hybridization during ecological divergence of southwestern
white pine (Pinus strobiformis) and limber pine (P. flexilis). *Molecular Ecology*, *27*(5), 1245-
1260.

Mena, J. L., & Pacheco, V. (2022). Mountains and traits: environmental heterogeneity and
mammal assemblages along an elevational gradient in the Northern Andes. *Studies on
Neotropical Fauna and Environment*, *57*(3), 227-239.

Murawski, D. A., & Hamrick, J. L. (1991). The effect of the density of flowering individuals on
the mating systems of nine tropical tree species. *Heredity*, *67*(2), 167-174.

Nason, J. D., & Hamrick, J. L. (1997). Reproductive and genetic consequences of forest
fragmentation: Two case studies of neotropical... *Journal of Heredity*, *88*(4).

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The bottleneck effect and genetic variability
in populations. *Evolution*, 1-10.

Nevado, B., Atchison, G. W., Hughes, C. E., & Filatov, D. A. (2016). Widespread adaptive
evolution during repeated evolutionary radiations in New World lupins. *Nature*
communications, *7*(1), 12384.

Nottingham, A. T., Fierer, N., Turner, B. L., Whitaker, J., Ostle, N. J., McNamara, N. P., ... &
Meir, P. (2018). Microbes follow Humboldt: temperature drives plant and soil microbial diversity
patterns from the Amazon to the Andes. *Ecology*, *99*(11), 2455-2466.

Nürk, N. M., Madriñán, S., Carine, M. A., Chase, M. W., & Blattner, F. R. (2013). Molecular
phylogenetics and morphological evolution of St. John's wort (Hypericum;
Hypericaceae). *Molecular Phylogenetics and Evolution*, *66*(1), 1-16.

Ottenburghs, J. (2020). Ghost introgression: spooky gene flow in the distant
past. *Bioessays*, *42*(6), 2000012.

Patterson, B. D., Solari, S., & Velazco, P. M. (2012). The role of the Andes in the diversification
and biogeography of Neotropical mammals. *Bones, clones, and biomes: the history and*
geography of Recent Neotropical mammals, 351-378.

Pease, J. B., Brown, J. W., Walker, J. F., Hinchliff, C. E., & Smith, S. A. (2018). Quartet
Sampling distinguishes lack of support from conflicting support in the green plant tree of
life. *American journal of botany*, *105*(3), 385-403.

Pérez-Escobar, O. A., Chomicki, G., Condamine, F. L., Karremans, A. P., Bogarín, D., Matzke,
N. J., ... & Antonelli, A. (2017). Recent origin and rapid speciation of Neotropical orchids in the
world's richest plant biodiversity hotspot. *New Phytologist*, *215*(2), 891-905.

Pérez-Escobar, O. A., Zizka, A., Bermúdez, M. A., Meseguer, A. S., Condamine, F. L., Hoorn,
C., ... & Chomicki, G. (2022). The Andes through time: evolution and distribution of Andean
floras. *Trends in Plant Science*, *27*(4), 364-378.

Perez Zabala, J. A. (2022). *Taxonomic Diversity, Phylogeny, and Diversification of the*
Environmental Niche of the Genus Prunus L. with emphasis on the New World Tropics (Doctoral
dissertation, UC Davis).

Pires, J. M., Dobzhansky, T. H., & Black, G. A. (1953). An estimate of the number of species of
trees in an Amazonian forest community. *Botanical Gazette*, *114*(4), 467-477.

Pitman, N. C., Terborgh, J., Silman, M. R., & Nuñez V, P. (1999). Tree species distributions in an
upper Amazonian forest. *Ecology*, *80*(8), 2651-2661.

Pouchon, C., Fernández, A., Nassar, J. M., Boyer, F., Aubert, S., Lavergne, S., & Mavárez, J. (2018). Phylogenomic analysis of the explosive adaptive radiation of the Espeletia complex (Asteraceae) in the tropical Andes. *Systematic biology*, *67*(6), 1041-1060.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945-959.

Quinlan, E. J., Layman, C. A., & Silman, M. R. (2025). Climate-Mediated Hybridisation and the Future of Andean Forests. *Journal of Biogeography*, *52*(6), e15113.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic biology*, *67*(5), 901-904.

Rapp, J. M., & Silman, M. R. (2012). Diurnal, seasonal, and altitudinal trends in microclimate across a tropical montane cloud forest. *Climate Research*, *55*(1), 17-32.

Rehder A (1940) Manual of cultivated trees and shrubs hardy in North America exclusive of the subtropical and warmer temperate regions. Dioscorides, Portland, OR.

Schley, R. J., Pennington, R. T., Pérez-Escobar, O. A., Helmstetter, A. J., de la Estrella, M., Larridon, I., ... & Klitgård, B. (2020). Introgression across evolutionary scales suggests reticulation contributes to Amazonian tree diversity. *Molecular Ecology*, *29*(21), 4170-4185.

Schley, R. J., Twyford, A. D., & Pennington, R. T. (2022). Hybridization: a 'double-edged sword' for Neotropical plant diversity. *Botanical Journal of the Linnean Society*, *199*(1), 331-356.

Schley, R. J., Piñeiro, R., Nicholls, J. A., Pezzini, F. F., Farbos, A., Lewis, G. P., ... & Pennington, R. T. (2025). Rampant Reticulation in a Rapid Radiation of Tropical Trees-Insights from Inga (Fabaceae). *Systematic biology*, syaf027.

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *the american naturalist*, *156*(S4), S4-S16.

Segovia, R. A., Pennington, R. T., Baker, T. R., Coelho de Souza, F., Neves, D. M., Davis, C. C., ... & Dexter, K. G. (2020). Freezing and water availability structure the evolutionary diversity of trees across the Americas. *Science Advances*, *6*(19), eaaz5373.

Sempere, T., Picard, D., & Plantard, O. (2005). Assessing and dating Andean uplift by phylogeography and phylochronology: early Miocene emergence of Andean cloud forests. In *6th International Symposium on Andean Geodynamics (ISAG 2005, Barcelona), Extended Abstracts* (pp. 663-665). IRD Éditions.

Silvestro, D., Schnitzler, J., Liow, L. H., Antonelli, A., & Salamin, N. (2014). Bayesian estimation of speciation and extinction from incomplete fossil occurrence data. *Systematic biology*, *63*(3), 349-367.

Simpson, G. G. (1953). *The major features of evolution*. Columbia University Press.

Sokoloff, D. D., Ignatov, M. S., Remizowa, M. V., Nuraliev, M. S., Blagoderov, V., Garbout, A.,
& Perkovsky, E. E. (2018). Staminate flower of Prunus s. l.(Rosaceae) from Eocene Rovno
amber (Ukraine). *Journal of Plant Research*, *131*, 925-943.

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.

Stroud, J. T., & Losos, J. B. (2016). Ecological opportunity and adaptive radiation. *Annual
Review of Ecology, Evolution, and Systematics*, *47*(1), 507-532.

Stull, G. W., Pham, K. K., Soltis, P. S., & Soltis, D. E. (2023). Deep reticulation: the long legacy
of hybridization in vascular plant evolution. *The Plant Journal*, *114*(4), 743-766.

Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. (2018). Adaptive introgression: a plant
perspective. *Biology letters*, *14*(3), 20170688.

Su, N., Hodel, R. G., Wang, X., Wang, J. R., Xie, S. Y., Gui, C. X., ... & Wen, J. (2023).
Molecular phylogeny and inflorescence evolution of Prunus (Rosaceae) based on RAD-seq and
genome skimming analyses. *Plant diversity*, *45*(4), 397-408.

Tao, R., & Iezzoni, A. F. (2010). The S-RNase-based gametophytic self-incompatibility system
in Prunus exhibits distinct genetic and molecular features. *Scientia Horticulturae*, *124*(4), 423-
433.

Ter Steege, H., Pitman, N. C., Sabatier, D., Baraloto, C., Salomão, R. P., Guevara, J. E., ... &
Silman, M. R. (2013). Hyperdominance in the Amazonian tree flora. *Science*, *342*(6156),
1243092.

Terborgh, J. (1971). Distribution on environmental gradients: theory and a preliminary
interpretation of distributional patterns in the avifauna of the Cordillera Vilcabamba,
Peru. *Ecology*, *52*(1), 23-40.

Terborgh, J. (1977). Bird species diversity on an Andean elevational gradient. *Ecology*, *58*(5),
1007-1019.

Tricou, T., Tannier, E., & de Vienne, D. M. (2022). Ghost lineages highly influence the
interpretation of introgression tests. *Systematic Biology*, *71*(5), 1147-1158.

Tsuda, Y., Semerikov, V., Sebastiani, F., Vendramin, G. G., & Lascoux, M. (2017). Multispecies
genetic structure and hybridization in the Betula genus across Eurasia. *Molecular Ecology*, *26*(2),
589-605.

Urrego, D. H., Silman, M. R., & Bush, M. B. (2005). The Last Glacial Maximum: stability and
change in a western Amazonian cloud forest. *Journal of Quaternary Science: Published for the
Quaternary Research Association*, *20*(7-8), 693-701.

Verde, I., Jenkins, J., Dondini, L., Micali, S., Pagliarani, G., Vendramin, E., ... & Schmutz, J.
(2017). The Peach v2. 0 release: high-resolution linkage mapping and deep resequencing
improve chromosome-scale assembly and contiguity. *BMC genomics*, *18*, 1-18.

Ward, M., Dick, C. W., Gribel, R., & Lowe, A. J. (2005). To self, or not to self… A review of
outcrossing and pollen-mediated gene flow in neotropical trees. Heredity, 95(4), 246-254.

Wen, J., Berggren, S. T., Lee, C. H., Ickert-Bond, S., Yi, T. S., Yoo, K. O., ... & Potter, D. (2008).
Phylogenetic inferences in Prunus (Rosaceae) using chloroplast ndhF and nuclear ribosomal ITS
sequences.

Williams Jr, J. H., Friedman, W. E., & Arnold, M. L. (1999). Developmental selection within the
angiosperm style: using gamete DNA to visualize interspecific pollen competition. *Proceedings*
of the National Academy of Sciences, *96*(16), 9201-9206.

Wu, C. I. (2001). The genic view of the process of speciation. *Journal of evolutionary*
biology, *14*(6), 851-865.

Zhao, L., Jiang, X. W., Zuo, Y. J., Liu, X. L., Chin, S. W., Haberle, R., ... & Wen, J. (2016).
Multiple events of allopolyploidy in the evolution of the racemose lineages in Prunus (Rosaceae)
based on integrated evidence from nuclear and plastid data. *PLoS One*, *11*(6), e0157123.

Table 1. Summary of samples included in individual-level phylogeny and genetic structure
analyses. Samples in bold were used to construct the species-level phylogeny.
822

Species determination	Sample ID	Voucher (Herbarium)	Material	Sampling locality
P. detrita	**107_detrita**	**EJQ82 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. debilis	**180_debilis**	**Loza et al. 388 (LPB)**	**Herbarium**	**Madidi NP, Bolivia**
P. debilis	225_debilis	JSB6352A (CUZ)	Fresh	Manu NP, Peru
P. stipulata	**171_stipulata**	**EJQ114 (LPB)**	**Fresh**	**Cotapata, Bolivia**
P. stipulata	173_stipulata	EJQ116 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	175_stipulata	EJQ118 (LPB)	Fresh	Cotapata, Bolivia
P. stipulata	285_stipulata	JSB6320 (CUZ)	Fresh	Manu NP, Peru
P. stipulata	286_stipulata	JSB6321(CUZ)	Fresh	Manu, NP, Peru
P. stipulata	312_stipulata	Cayola 3259	Herbarium	La Paz, Bolivia
P. stipulata	332_stipulata	Villalobosos 81	Herbarium	
P. integrifolia	139_integrifolia	JSB6366 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	**147_integrifolia**	**JSB6402 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. integrifolia	148_integrifolia	JSB6396 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	149_integrifolia	JSB6403 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	151_integrifolia	JSB6400 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	155_integrifolia	JSB6397 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	156_integrifolia	JSB6391 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	157_integrifolia	JSB6392 (CUZ)	Fresh	Manu NP, Peru
P. integrifolia	212_integrifolia	Meneses 7535 (LPB)	Herbarium	Nor Yungas, Cotapata, Bolivia
P. integrifolia	253_integrifolia	JSB7255 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	162_pleiantha	JSB6382 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	**163_pleiantha**	**JSB6388 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. pleiantha	164_pleiantha	JSB6386 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	165_pleiantha	JSB6387 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	166_pleiantha	JSB6385 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	167_pleiantha	JSB6380 (CUZ)	Fresh	Manu NP, Peru
P. pleiantha	169_pleiantha	JSB6390 (CUZ)	Fresh	Manu NP, Peru
P. herthae	226_herthae	JSB6343 (CUZ)	Fresh	Manu NP, Peru
P. herthae	236_herthae	JSB6333 (CUZ)	Fresh	Manu NP, Peru
P. herthae	**237_herthae**	**JSB6334 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. amplifolia	**206_amplifolia**	**Beck 33550 (LPB)**	**Herbarium**	**La Paz, Bolivia**
P. antioquensis	**177_antioquensis**	**Araujo & Canqui 348 (LPB)**	**Herbarium**	**Madidi, Bolivia**
P. huantensis	108_huantensis	JSB6316 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	115_huantensis	JSB6310 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	**116_huantensis**	**JSB6358 (CUZ)**	**Fresh**	**Manu NP, Peru**
P. huantensis	117_huantensis	JSB6360 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	119_huantensis	JSB6362 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	123_huantensis	JSB6353 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	124_huantensis	JSB6357 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	130_huantensis	JSB6375 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	131_huantensis	JSB6311 (CUZ)	Fresh	Manu NP, Peru
P. huantensis	132_huantensis	JSB6312 (CUZ)	Fresh	Manu NP, Peru

P. brittoniana	**179_brittoniana**	**Paniagua et al. 5782A (LPB)**	**Herbarium**	**Pelechuco-Mojo, Bolivia**
P. brittoniana	210_brittoniana	Beck 35044 (LPB)	Herbarium	Sud Yungas, Cotapata, Bolivia
Outgroups:				
P. myrtifolia	**1_myrtifolia**	**Silica sample only**	**Fresh**	**Fairchild Botanical Garden, Miami, Florida, USA**
P. caroliniana	**2_caroliniana**	**Silica sample only**	**Fresh**	**Weymouth Woods, North Carolina, USA**

824   Table 2. Summary of significant D-statistic tests, with *P. myrtifolia* set to P4.

P1	P2	P3	dstat	Z	ABBA	BABA	nloci
P. antioquensis	*P. stipulata*	*P. debilis*	-0.69	4.75	8.38	45.13	152
P. stipulata	*P. antioquensis*	*P. debilis*	0.69	4.60	45.13	8.38	152
P. amplifolia, P. stipulata	*P. antioquensis*	*P. debilis*	0.66	4.16	43.19	8.94	163
P. antioquensis	*P. amplifolia, P. stipulata*	*P. debilis*	-0.66	4.01	8.94	43.19	163
P. stipulata	*P. pleiantha*	*P. detrita, P. debilis*	0.13	2.94	793	605.44	3968
P. pleiantha	*P. stipulata*	*P. detrita, P. debilis*	-0.13	2.87	605.44	793	3968
P. herthae, P. huantensis	*P. pleiantha*	*P. detrita, P. debilis*	0.11	2.81	864.08	686.51	4485
P. stipulata	*P. huantensis, P. antioquensis*	*P. pleiantha*	0.09	2.71	1190.20	989.74	6218
P. huantensis, P. antioquensis	*P. stipulata*	*P. pleiantha*	-0.09	2.71	989.74	1190.2	6218
P. pleiantha	*P. herthae, P. huantensis*	*P. detrita, P. debilis*	-0.11	2.67	686.51	864.08	4485
P. stipulata	*P. pleiantha*	*P. detrita*	0.13	2.66	744.38	578.5	3725
P. pleiantha	*P. antioquensis*	*P. debilis*	0.51	2.63	20.25	6.5	142
P. pleiantha	*P. stipulata*	*P. detrita*	-0.13	2.62	578.5	744.38	3725
P. antioquensis	*P. pleiantha*	*P. debilis*	-0.51	2.58	6.5	20.25	142
P. pleiantha	*P. stipulata*	*P. debilis*	-0.27	2.53	68.5	119.25	576
P. herthae, P. huantensis, P. antioquensis, P. brittoniana	*P. pleiantha*	*P. detrita*	0.11	2.52	813.45	654.78	4203

830   Table 3. Summary of within species genomic diversity statistics.

Species	# of samples	Filter	# of loci	Average π	Average H_o	Average H_e
P. integrifolia	10	50	53574	0.42 ± 0.13	0.15 ± 0.064	0.42 ± 0.0025
		70	17592	0.41 ± 0.14	0.17 ± 0.073	0.40 ±0.0015
P. huantensis/ brittoniana	12	50	24457	0.41 ± 0.13	0.12 ± 0.078	0.41 ± 0.0012
		70	1960	0.38 ± 0.15	0.16 ± 0.10	0.37 ± 0.0031
P. stipulata	7	50	24457	0.45 ± 0.11	0.095 ± 0.096	0.45 ± 0.0024
				0.42 ± 0.13	0.11 ± 0.11	0.42 ± 0.0064
P. pleiantha	7	50	23299	0.45 ± 0.11	0.12 ± 0.081	0.44 ± 0.0015
		70	7295	0.43 ± 0.12	0.13 ± 0.091	0.42 ± 0.0013
P. herthae	3	50	6251	0.62 ± 0.077	0.085 ± 0.03	0.61 ± 0.0067
		70	699	0.51 ± 0.073	0.064 ± 0.0072	0.51 ± 0
P. debilis/detrita	3	50	1747	0.64 ± 0.061	0.064 ± 0.058	0.064 ± 0.0009

832   Table 4. Summary of within-individual heterozygosity estimates.

Species	Individual	H_o (50% filter)	H_e (50% filter)	H_o (70% filter)	H_e (70% filter)
P. integrifolia	139_integrifolia	0.11	0.42	0.12	0.41
	147_integrifolia	0.31	0.42	0.36	0.4
	148_integrifolia	0.14	0.42	0.16	0.4
	149_integrifolia	0.18	0.42	0.21	0.4
	151_integrifolia	0.10	0.42	0.12	0.4
	155_integrifolia	0.16	0.42	0.19	0.4
	156_integrifolia	0.12	0.42	0.14	0.4
	157_integrifolia	0.17	0.42	0.2	0.4
	212_integrifolia	0.09	0.41	0.1	0.4
	253_integrifolia	0.12	0.42	0.14	0.41
P. huantensis /brittoniana	108_huantensis	0.16	0.41	0.23	0.37
	115_huantensis	0.1	0.41	0.13	0.38
	116_huantensis	0.22	0.41	0.29	0.37
	117_huantensis	0.044	0.4	0.07	0.37
	119_huantensis	0.072	0.41	0.093	0.37
	123_huantensis	0.069	0.4	0.12	0.37
	124_huantensis	0.2	0.4	0.25	0.37
	130_huantensis	0.05	0.41	0.079	0.38
	131_huantensis	0.15	0.41	0.22	0.37
	132_huantensis	0.26	0.41	0.36	0.37
	179_brittoniana	0.064	0.41	0.08	0.37
	210_brittoniana	0.03	0.4	0.05	0.38
P. stipulata	171_stipulata	0.28	0.45	0.32	0.42
	173_stipulata	0.17	0.45	0.18	0.42
	175_stipulata	0.07	0.45	0.07	0.41
	285_stipulata	0.058	0.45	0.1	0.42
	286_stipulata	0.03	0.45	0.041	0.41
	312_stipulata	0.037	0.45	0.032	0.42
	332_stipulata	0.02	0.45	0.021	0.43

P. pleiantha	162_pleiantha	0.12	0.44	0.13	0.43
	163_pleiantha	0.22	0.44	0.25	0.43
	164_pleiantha	0.18	0.44	0.19	0.42
	165_pleiantha	0.2	0.44	0.22	0.42
	166_pleiantha	0.023	0.44	0.021	0.42
	167_pleiantha	0.05	0.44	0.052	0.42
	169_pleiantha	0.051	0.44	0.056	0.42
P. herthae	226_herthae	0.087	0.62	0.057	0.51
	236_herthae	0.054	0.61	0.063	0.51
	237_herthae	0.11	0.62	0.072	0.51
P. debilis/ detrita	180_debilis	0.13	0.64	NA	NA
	225_debilis	0.036	0.64	NA	NA
	107_detrita	0.026	0.64	NA	NA

Figure 1. Box plots showing elevational distributions of the in-group Andean species across three
different sources. "Aberg" includes all known individuals along the ABERG transect >10 dbh
over 20 years. "Acha" includes all Bolivian material reviewed by Achá (2013) and "Zabala"
includes all material reviewed by Perez-Zabala (2022) from Venezuela to Bolivia. Perez-Zabala
considered *P. huantensis* to be *P. brittoniana* so those samples are grouped together on the figure
under *P. brittoniana*. Figure does not include new "varieties" of species proposed by either Achá
or Perez-Zabala.

Figure 2. (a) Map of samples included in this study. Black dot indicates the location of the
ABERG transect in Manu National Park, Peru where most samples were collected for this study.
Red points indicate locations of additional samples included in the study. (b) Image of the
ABERG transect highlighted by the box in (a), showing location of 1-ha plots. Further details on
individual samples provided in Table 1.

Figure 3. Species-level topology resulting from RAxML and ExaBayes phylogenetic analyses.
Nodes show RAxML bootstrap support values where black circles indicate values > 75 and
values < 75 are shown on white nodes. All Bayesian posterior probabilities were equal to 1. The
tree is rooted to the reference (*P. persica*) and all three outgroup species are shown in grey.
Dotted arrows summarize patterns of historical admixture inferred from D-statistics.

863   Figure 4. Time-calibrated phylogeny generated from BEAST analysis. Posterior probabilities are
864   shown for nodes when < 1. Node bars represent the 95% highest probability density (HPD)
865   interval for node ages. Red stars indicate the fossil calibration points.

871 Figure 5. Patterns of genetic structure among individuals of sympatric species. Each vertical bar
872 represents a single individual and the y-axis indicates the proportion of genetic ancestry of that
873 individual assigned to different inferred genetic clusters, designated with different colors.
874 STRUCTURE detected K=9 distinct genetic clusters as the most likely grouping (Figure S3).

Figure S1. Maximum likelihood individual-level phylogeny reconstructed in RAxML. Individuals were selected to include those with at least 1000 loci per sample and up to 10 individuals per species. Sites were filtered to only include those with at least 25% coverage across samples. Tree was rooted to the reference (*P. persica*) and nodes show bootstrap support value.

Figure S2. Quartet sampling results with quartet concordance (QC)/quartet differential
(QD)/quartet informativeness (QI) indicated next to nodes. Nodes are colored according to QC
scores, indicating how often the concordant quartet is inferred over both discordant quartets.
Outgroup samples *P. persica* and *P. caroliniana* are removed for clarity. Figure created with
https://github.com/ShuiyinLIU/QS_visualization.

Figure S3. Plot of estimators used to select the optimal number of genetic clusters (K) in the
STRUCTURE analysis. Estimated log mean probability assesses how well the model with a
particular K explains the genetic data. As the log-likelihood values are negative, the best-fit K-
value is the least negative. Delta K estimates the rate of change in the log probability of data
between successive K values. The K value corresponding to the highest delta K and estimated log
mean probability (K=9) was considered the best estimate of the number of subpopulations in this
dataset.