In Vitro Evidence to Support Amphotericin B and Flucytosine Combination Therapy for

Talaromycosis

Heera Natesan Sambath*, **[1], Shawin Vitsupakorn*[1,2], Kaushik Sreerama Reddy*[3], Lottie

Brown[4,5], Nguyen Thi Mai (Last name) Thu (First name)[1], Matthew Burke[1], Jialin Liu[1], Emily

Evans[6], Charles Giamberardino[1], John Perfect[1], Ngo Thi (Last name) Hoa (First name)[7,8,9],

Thuy Le**[1,7]

[1] Division of Infectious Diseases and International Health, Duke University School of Medicine,

Durham, North Carolina, USA

[2] The Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[3] Division of Rheumatology and Clinical Immunology, University of Pittsburgh School of

Medicine, Pittsburgh, Pennsylvania, USA

[4] Institute of Infection and Immunity, City St George's University of London, School of Health &

Medical Sciences, London, UK

[5] St George's University Hospitals NHS Foundation Trust, London, UK

[6] Emory University School of Medicine, Atlanta, Georgia, USA

[7] Tropical Medicine Research Center for Talaromycosis, Biomedical Research Center, Pham

Ngoc Thach University of Medicine, Ho Chi Minh City, Vietnam

[8] Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam

[9] Centre for Tropical Medicine and Global health, University of Oxford, UK

* Authors contributed equally

**Corresponding authors

Abstract: 249 words

Main text: 2984 words

Conflict of Interest

TL receives investigator-initiated research grant from Gilead Science to contribute to the

ongoing phase III Liposomal Amphotericin B and Flucytosine Antifungal Therapy (LAmB-FAST)

randomized clinical trial (clinicaltrials.gov NCT06525389). All other authors declare no conflict of

interest related to this work.

Funding

This work was supported by grants from the National Institutes of Health to TL [R01AI143409,

R21AI162367, R21AI159397, U01AI169358, P30AI064518]. LB is funded by the National

Institute for Health Research (NIHR) and received fellowship grants from the Federation of

European Microbiological Societies, Dowager Countess Eleanor Peel Trust, the British Society

for Medical Mycology and the Wingate Foundation. The funders had no role in the design and

conduct of the study.

Previous Presentation:

A part of this work was presented at the 11[th] Trends in Medical Mycology (TIMM-11)

conference, held in Athens, Greece, in April 2023.

Corresponding Authors:

1. Heera Natesan Sambath, MBBS. Division of Infectious Diseases and International
   Health, Duke University School of Medicine, Alex H. Sands Basic Sciences Research
   Building, 303 Research Drive, Room 346, Duke University, Durham, NC, USA. Phone:
   +1 (919)-684-0952. Email: Heera.natesan.sambath@duke.edu (ORCID ID: 0009-0007-
   8729-6121).

2. Thuy Le, MD, DPhil. Division of Infectious Diseases and International Health, Duke
   University School of Medicine, Hanes House, 315 Trent Drive, Room 257, Duke

51   University Medical Center, Box 102359, Durham, NC, USA. Phone: +1 (919)-668-5053.

52   Email: Thuy.le@duke.edu (ORCID ID: 0000-0002-3393-6580).

Abstract

Background: *Talaromyces marneffei* causes talaromycosis, a life-threatening fungal disease with limited treatment options. The standard treatment of amphotericin B (AmB) induction followed by itraconazole consolidation still results in 15% to 30% mortality. This study aimed to investigate the potential of AmB and flucytosine (5FC) combination therapy to enhance antifungal activity.

Methods: The in vitro antifungal activity of AmB and 5FC alone and in combination against 60 *T. marneffei* clinical isolates was evaluated using a validated colorimetric antifungal susceptibility assay and the checkerboard method. The minimum inhibitory concentration (MIC) was defined as the lowest drug concentration inhibiting ≥ 95% fungal growth (MIC_{95}) for both AmB and 5FC. The combination effect between AmB and 5FC against *T. marneffei* was determined using fractional inhibitory concentration index. Combination effects were further tested using a time-kill assay.

Results: The MIC_{95} was 0.25 – 2 µg/mL (geometric mean [GM] 0.68 µg/mL) for AmB, and 0.03 – 0.5 µg/mL (GM 0.28 µg/mL) for 5FC. Full synergy was observed in 4 isolates (7%), and indifference was observed in the remaining 56 isolates (93%). The time-kill experiments revealed a concentration-dependent fungicidal activity of AmB, and concentration-independent fungistatic effect of 5FC. Synergy between AmB and 5FC was confirmed, showing greater than 2-log_{10} reduction in colony forming units when used in combination. No antagonism was observed.

Conclusions: Our study demonstrated in vitro evidence of synergistic activity between AmB and 5FC against *T. marneffei*, providing the evidence to support in vivo and clinical trial testing of AmB and 5FC combination therapy, and dosing reduction strategies of 5FC.

Lay Summary: This study demonstrated in vitro synergy between amphotericin B and

flucytosine against *Talaromyces marneffei*, providing the proof of concept to test this antifungal

combination in a clinical trial of talaromycosis.

Keywords: Talaromycosis; *Talaromyces marneffei*; combination therapy; amphotericin B,

flucytosine.

Background

The need to improve the treatment of talaromycosis

The thermally dimorphic fungus *Talaromyces marneffei* causes talaromycosis, a life-threatening

invasive fungal disease endemic in Southeast Asia. Over just three decades, talaromycosis has

emerged from a rare disease to a leading opportunistic infection among people with advanced

HIV and other immunocompromised conditions, with an estimated 25,000 cases annually [1].

Current treatment options for talaromycosis are limited to just two drugs: amphotericin B (AmB)

for induction therapy for the first 2 weeks, followed by itraconazole consolidation and

maintenance therapy for at least 12 months [2]. Despite appropriate therapy, mortality remains

unacceptably high at 15% to 30% [3,4]. AmB is a large molecule with poor tissue penetration

[5], which reduces its efficacy in treating deep-seated multi-organ infections. Treatment effect is

further compromised by toxicity of AmB and poor pharmacokinetic profiles of itraconazole. Over

50% of patients on AmB experienced grade 3 or higher adverse events [2]. Itraconazole is

better tolerated but has poor oral bioavailability (only 20% to 40%), non-linear pharmacokinetics,

and a narrow therapeutic index [6]. The high mortality on treatment and limited treatment

options prompt the inclusion of *T. marneffei* on the WHO Fungal Priority Pathogen List [7],

calling for research on new therapeutics and treatment strategies for talaromycosis.

Rationale for antifungal combination therapy

Some novel antifungal drugs have promising potential for talaromycosis but take a long time to

be realized. A combination of existing therapy offers a promising strategy. Combination therapy

using drugs from different antifungal classes has been used to optimize the treatment of

invasive fungal diseases, including cryptococcosis, invasive aspergillosis and candidiasis [8–

10]. Combining drugs with distinct and complementary mechanisms of action can enhance

therapeutic efficacy by leveraging synergistic interactions, increasing drug penetration,

accelerating fungal clearance, and ultimately improving patient outcomes. Enhanced antifungal

activity could permit lower dosing, thereby reducing dose-related toxicity, and may help prevent

the emergence of resistance associated with monotherapy [8].

Rationale for amphotericin B and flucytosine (5FC) combination therapy for talaromycosis

AmB and 5FC act through different mechanisms which can potentiate one another to enhance

fungal killing*.* In vitro studies of AmB and 5FC in *Cryptococcus* and *Candida* showed varied

results ranging from synergy to indifference and, in the case of 5FC-resistant isolates,

antagonism. Nevertheless, these findings translated into improved fungal clearance in vivo

models and survival in clinical trials [11–15]. AmB and 5FC combination has been shown in

several clinical trials of cryptococcal meningitis to enhance fungal clearance in the cerebrospinal

fluid (CSF) and reduce mortality by 30% to 40% compared to AmB alone [16,17]. 5FC has

excellent oral bioavailability, reaching a maximum serum concentration of 50 µg/mL within 2

hours [18], and good penetration into the deep tissues and CSF [18,19]. 5FC given at 25 mg/kg

four times daily is generally safe and well-tolerated [16,17].

We and others have shown that 5FC has potent in vitro antifungal activity against *T. marneffei.*

The minimum inhibitory concentrations (MICs) range between 0.015 and 1.0 µg/mL, which is

75-folds lower than serum concentrations achieved with standard dosing [20–22]. Given potent

in vitro activity of 5FC against *T. marneffei,* robust mortality benefit of AmB and 5FC

combination therapy in cryptococcal disease, and the favorable safety profile, we hypothesize

that the AmB and 5FC combination may enhance the in vitro fungicidal activity against *T.*

marneffei and aim to test this hypothesis with the goal to expand treatment options and improve

outcomes of talaromycosis.

Materials and Methods

Ethics statement

This study utilized *T. marneffei* clinical isolates obtained from participants of the multi-center

Itraconazole *versus* Amphotericin for Penicilliosis (IVAP) randomized controlled trial, which was

approved by all five study sites in Vietnam, and the Oxford Tropical Research Ethical

Committee (OxTREC) in the UK, and the Vietnam Ministry of Health (MoH) clinical trial

registration: ISRCTN59144167, MoH approval: 3300/QD-BYT). All participants gave informed

consent for specimens to be stored in the IVAP biobank stored at Duke University and used in

this research (Duke IRB protocol: Pro00101915).

Talaromyces marneffei isolates

For the current study, 60 *T. marneffei* clinical isolates were randomly selected from 336 isolates

collected and whole-genome sequenced from the IVAP trial [2,3]. A sample size of 60 is

considered more than sufficient to obtain a normal sampling distribution [23]. We selected 30

isolates from northern Vietnam and 30 from southern Vietnam to represent the two

phylogenetically and geographically distinct clades [3]. The isolates were prepared as outlined

in **Supplementary File 1**.

Antifungal susceptibility testing

Antifungal drugs were prepared as outlined in **Supplementary File 2** and susceptibility testing

was performed using a validated Clinical and Laboratory Standards Institute (CLSI)-based

colorimetric assay [20], as outlined in **Supplementary File 3**. Although a 95% inhibition (MIC_{95})

threshold was used as the MIC endpoint for AmB, and MIC_{50} threshold was used for 5FC as per

CLSI, MIC_{95} was used for both AmB and 5FC to allow evaluation of combination effect by the

checkerboard assay.

AmB and 5FC combination effect in the checkerboard assay

Figure 1 illustrates the checkerboard assay setup, which enables simultaneous determination of

the MICs for each drug alone and in combination using a fungal inoculum of $1 - 5 \times 10^3$ colony

forming units per mL (CFUs/mL). The combination effect of AmB and 5FC were characterized

using the fractional inhibitory concentration index (FICI) [24], calculated according to **Equation**

A (Supplementary File 4). Classification of drug-drug interaction based on FICI values has

varied across studies. Older studies classify FICI values between 0.5 and 1 as 'additive or

partially synergy', and values between 1 and 4 as 'indifferent' [11,13]. Newer studies broadly

define FICI values between 0.5 and 4 as 'indifference' [8,9]. As prior classification system may

overestimate synergy, we used the more conservative and simple classification system to

assess interaction: (1) synergy (FICI ≤ 0.5), (2) indifference (0.5 < FICI ≤ 4), and (3) antagonism

(FICI > 4). Synergy was defined as a minimum four-fold decrease in the MICs with AmB and

5FC combination compared to each drug alone.

Validation of AmB and 5FC synergy by time kill assay

Time-kill assay was performed to evaluate the antifungal dynamics of AmB and 5FC

combination over time. A *T. marneffei* isolate which showed evidence of synergy of AmB and

5FC combination in the checkerboard experiments was tested. Time-kill assay procedures are

summarized in the **Supplementary File 5**. Fungal growth was measured by CFUs/mL and

converted to $\log_{10}$ CFU/mL to obtain normal distribution for analysis. A fungicidal effect of a drug

was defined as a minimum 3-$\log_{10}$ reduction in fungal burden from the starting inoculum,

whereas a reduction of less than 3-$\log_{10}$ is classified as fungistatic effect [25,26]. Synergy was

defined as a reduction of at least 2-$\log_{10}$ in fungal growth when the drugs are tested in

combination compared to the most active agent alone [25].

Quality control

Quality control procedures, including *Candida krusei* ATCC 6258 and the internal *T. marneffei*

strain 11CN-20-091, are detailed in **Supplementary File 6.**

Statistical analysis

The mode and geometric mean (GM) of MICs, and the mean and standard deviation of the FICs

and FICI for 60 isolates were calculated, along with 95% confidence interval (95% CI). The

mean differences in MICs between groups – AmB alone, 5FC alone, *versus* their combination –

were compared using paired student t-tests. MIC data follow a geometric sequence with a

common ratio of 2 (i.e., 2-fold serial dilutions), therefore the data was $\log_2$ transformed to

demonstrate fold-changes in **Figure 2.** For time-kill curves, we $\log_{10}$ transformed CFUs/mL and

generated trajectories for each testing conditions over 120 hours. All statistical analyses were

performed, and graphs were created using GraphPad Prism Version 8.4.0. (GraphPad software,

Boston, MA, USA). Illustrations were created using BioRender (Toronto, Ontario, Canada).

Results

MICs of AmB and 5FC alone and in combination

Table 1 summarizes the MICs for each drug, and the FICs and FICI of the 60 *T. marneffei*

isolates. Individual values for each isolate are shown in **Supplementary Table 1 (File 7)**. The

MIC_{95} range for AmB was 0.25 – 2 µg/mL (GM 0.68 µg/mL). The MIC_{95} and MIC_{50} ranges for

5FC were 0.06 – 2 µg/mL (GM 0.28 µg/mL) and 0.03 – 0.5 µg/mL (GM 0.13 µg/mL),

respectively. **Figure 2A and 2B** show the MIC_{95} of AmB and 5FC when tested alone *versus* in

combination. The MIC of AmB decreased three-fold on average when 5FC was added, and the

MIC of 5FC decreased four-fold on average when AmB was added ($P < 0.0001$ for both groups

by paired t-test).

AmB and 5FC combination effect by checkerboard assay

Figure 2C shows the distribution of the FICI values and classification of combination effect for

all isolates. Synergistic effects between AmB and 5FC were observed in four isolates (7%), all

from patients in northern Vietnam. Indifferent effects were observed in the remaining 56 isolates

(93%), and no antagonism was detected in any isolate.

Concentration-dependent antifungal dynamics of AmB and 5FC against T. marneffei by time-kill

assay

For the time-kill assay, we tested the *T. marneffei* isolate 11CN-21-012 which demonstrated

synergy between AmB and 5FC in the checkerboard assay. **Figure 3A** demonstrates a

concentration-dependent fungicidal effect of AmB against *T. marneffei* with a 3-$\log_{10}$ reduction in

CFUs/mL at concentrations of 0.5, 1, and 2 times the MIC compared to the starting inoculum.

5FC however demonstrated a concentration-independent antifungal effect*,* with less than 2-$\log_{10}$

reduction compared to the starting inoculum, even at concentrations as high as 20 times the

MIC, and was thus considered fungistatic against *T. marneffei* (**Figure 3B**).

Synergy between AmB and 5FC combination by time-kill assay

Because AmB exhibited such a robust fungicidal activity at its MIC (**Figure 3A**), and this

concentration would have masked any potential synergistic effects of 5FC, we tested sub-MIC

concentrations of AmB in combination with 5FC in order to allow a clear delineation of any

additional antifungal effect of 5FC on AmB. Sub-MIC concentrations of AmB also mimic true

drug concentrations achieved in the CSF due to poor tissue penetration of AmB [27].

The combinations of AmB (at 0.25 and 0.5 times the MIC) and a broad range of 5FC

concentrations (1 – 20 times the MIC) resulted in a 2- to 3-$\log_{10}$ reduction in CFUs by 96 hours

compared to AmB alone. This is indicative of robust synergy between AmB and 5FC.

Synergistic effect was evident even at the lowest concentrations of AmB and 5FC (0.25 times

and 1 time the MIC, respectively) **(Figure 4)**. Increasing the concentration of either drugs

beyond these levels did not lead to further enhancement of antifungal activity. No evidence of

antagonism was observed up to 120 hours.

Discussion

Summary of major findings

The current treatment of talaromycosis relies on only two drugs with suboptimal efficacy and safety profiles [2]. Antifungal combination therapy has shown to improve fungicidal activity and clinical outcomes for other serious fungal diseases, but had not been investigated for talaromycosis [9]. In this study, we performed a robust in vitro investigation of the antifungal effect of AmB and 5FC combination against *T. marneffei,* testing 60 *T. marneffei* clinical isolates representing two distinct phylogenetic and geographical clades in northern and southern Vietnam. We demonstrated full synergy (more than four-fold reduction of MICs bidirectionally) in 7% of isolates against *T. marneffei* using the checkerboard method. The remaining 93% of isolates showed a FICI between 0.52 and 1.13, which is classified by the more conservative classification system as indifference interaction. However, all 56 isolates demonstrated a three- to four-fold reduction in MICs with the AmB and 5FC combination, which would be classified as partial synergy by previous classification systems. We confirmed full synergy with the time-kill experiment. No antagonism was observed. Together, these findings provide promising in vitro evidence that AmB and 5FC combination therapy may be beneficial against talaromycosis.

Postulated mechanisms of synergy

We posit that the synergy between AmB and 5FC is due to their complementary mechanisms of action and permeability hypothesis. AmB extracts ergosterol from the fungal cell membrane, leading to leakage of intracellular ions and cell death [28]. This allows 5FC to permeate the cell wall and access the nucleus, where it converts to 5-fluorouracil and other metabolites that disrupt DNA and RNA synthesis [29]. AmB may also impair the fungal efflux mechanisms, thereby reducing the export of 5FC and resulting in higher intracellular concentration of 5FC [9]. Conversely, AmB is a large molecule that has variable penetration in the lungs, liver, kidneys, and a poor penetration in the brain and CSF [30,31]. The disruption of pyrimidine biosynthesis

by 5FC leads to structural defects in the fungal cell wall, facilitating membrane and tissue

penetration of AmB and potentiating the fungicidal effect of AmB [29]. We postulate that the

addition of 5FC to AmB may improve the treatment of deep-seated and multi-organ

dissemination nature of talaromycosis, and may reduce the incidence of disease relapse, and

incidence of immune reconstitution disease as the immune system of the host is restored [18].

Our findings in the context of existing literature

The MIC ranges observed for AmB and 5FC tested alone were consistent with the literature and

reflect the diversity of the clinical isolates. Interestingly, all four isolates showing full synergy

were from northern Vietnam, raising the question of whether treatment effect is clade-specific

[3]. The four isolates showing full synergy also had higher MICs for AmB (1–2 µg/mL);

however, not all isolates with higher AmB MICs exhibited full synergy. The remaining isolates

exhibited indifference, consistent with previous in vitro studies on *Cryptococcus* and *Candida*

spp.*,* which have reported a range of outcomes from synergy to indifference [26,32]. The

evidence of in vitro synergy in *Cryptococcus* has translated into robust antifungal effect and

mortality reduction in several randomized controlled trials, supporting AmB and 5FC

combination therapy as the first-line induction regimen for cryptococcosis [16,33,34]. While in

vitro synergy of AmB and 5FC combination against *Candida* has been inconsistent, excellent in

vivo activity has been observed. Mice treated with AmB and 5FC combination had substantially

higher survival rates than those treated with AmB or 5FC monotherapy (79.5% *vs* 15.9% and

13.7%, respectively) [35,36]. AmB and 5FC combination therapy is recommended for deep-

seated *Candida* infections, such as *Candida* endocarditis and endophthalmitis [37]. In vivo

studies and clinical trials are now needed to confirm the synergy of AmB and 5FC against *T.*

marneffei, with the ultimate goal to reduce mortality from talaromycosis.

Have we been overdosing 5FC in the setting of antifungal combination therapy?

5FC toxicity, including myelosuppression and hepatotoxicity, is dose dependent [19]. The

standard four times daily dosing of 5FC is impractical and can lead to adherence issues.

Emerging evidence suggests that lower doses of 5FC maintain efficacy while reducing toxicity.

The current recommended dose of 5FC (100mg/kg/day in 4 divided doses) rapidly achieves

peak serum concentrations of 60 – 80 µg/mL [19], which is more than 75-fold the MICs of 5FC

against *T. marneffei*. In the time-kill experiments, we showed that 5FC was fungistatic

regardless of concentration, and that the synergistic effect of adding 5FC was not dose-

dependent, suggesting that a higher 5FC dose or level is not more effective. In a murine model

of cryptococcal meningitis, the antifungal activity of 5FC at a 50mg/kg/day combined with AmB

caused near maximal fungal killing [33], similar to the findings from a murine model of invasive

candidiasis [38]. However, a recent non-comparative clinical trial in Uganda (n = 48) evaluated

5FC at a reduced dose (60mg/kg/day in 3 divided doses) for patients with cryptococcal

meningitis and observed slower CSF fungal clearance rate than historical data for the standard

5FC dosing [39]. This is potentially an issue of 5FC penetration into the CSF, which need to be

confirmed by pharmacokinetic data. Our data suggest that lower and/or less frequent dosing of

5FC may offer a more practical, safer, and equally effective alternative to the standard dosing of

5FC for talaromycosis.

Strengths and limitations

Our study has several limitations. Firstly, our evaluation of combination effect in the 60 clinical

isolates was based on the checkerboard method, which provides a "snapshot" of fungal growth

inhibition at a fixed point in time. Disagreement between different methods of measuring drug

activity (such as the checkerboard assays, E-test and time-kill assay) has been reported [26].

While we investigated the temporal interaction between AmB and 5FC in a time-kill assay, our

analysis was limited to a single isolate due to resource constraints, so the findings may not be

generalizable across genetically diverse isolates. For efficiency, we recommend the

checkerboard method as a screening tool for synergy and antagonism, followed by further

assessment with time-kill assay and in vivo studies. Secondly, no consensus exists for

calculating the FICI for agents with discordant endpoints (e.g., MIC_{50} for 5FC *versus* MIC_{95} for

AmB). We chose MIC_{95} threshold for 5FC in the evaluation of combination therapy to allow a

compatible calculation of FICI in the checkerboard assay. Thirdly, only isolates from Vietnam

were included in our study. Future studies should test isolates from a range of endemic regions

to determine whether the fungicidal activity of AmB and 5FC is consistent across different

geographical and genetic clades of *T. marneffei* beyond Vietnam. Finally, we recognize that in

vitro interactions may not reflect real-world conditions, where variations in pharmacokinetics,

pH, protein binding, host immune response, and the presence of other drugs and co-infections

can impact clinical efficacy. In vivo studies are essential to understand the pharmacokinetic and

pharmacodynamic properties of combination therapy. Despite these limitations, our study has

notable strengths. We tested a large number of representative clinical isolates, undertook

rigorous quality assurance measures, including the use of an internal control strain in all

experiments and CFU counts to ensure inoculum consistency.

Clinical Implications

Antifungal combination therapy may improve treatment efficacy in deep-seated and

disseminated talaromycosis, reducing the risk of relapse and mortality, and allowing for lower

and less toxic dosage. Our study identifies AmB and 5FC as a promising synergistic

combination and supports its testing in the ongoing Liposomal Amphotericin B and Flucytosine

Antifungal Therapy (LAmB-FAST) clinical trial (NCT06525389). Our findings also pave the way

for broader testing of antifungal combination therapies against talaromycosis and other invasive

fungal infections.

Acknowledgements

We thank patients of the IVAP trial for donating samples for this research and healthcare staff in

Vietnam who led recruitment.

Author Contributions

TL, HNS, SV, and NTMT conceived and designed the study. SV, HNS, NTMT, and KSR

generated the data. HNS, SV, NTMT, and JL performed the statistical analyses. SV, KSR, HNS,

LB, and NTMT drafted the manuscript. EE, CG, JP, LB and NTH provided clinical and technical

input for the research. All authors reviewed and approved the final manuscript.

References

1. Narayanasamy S, Dat VQ, Thanh NT, et al. A global call for talaromycosis to be recognised as a neglected tropical disease. Lancet Glob Health **2021**; 9:e1618–e1622.

2. Le T, Kinh NV, Cuc NTK, et al. A Trial of Itraconazole or Amphotericin B for HIV-Associated Talaromycosis. N Engl J Med **2017**; 376:2329–2340.

3. Sephton-Clark P, Nguyen T, Hoa NT, et al. Impact of pathogen genetics on clinical phenotypes in a population of Talaromyces marneffei from Vietnam. Genetics **2023**; 224:iyad100.

4. Le T, Wolbers M, Chi NH, et al. Epidemiology, Seasonality, and Predictors of Outcome of AIDS-Associated Penicillium marneffei Infection in Ho Chi Minh City, Viet Nam. Clin Infect Dis Off Publ Infect Dis Soc Am **2011**; 52:945–952.

5. Vogelsinger H, Weiler S, Djanani A, et al. Amphotericin B tissue distribution in autopsy material after treatment with liposomal amphotericin B and amphotericin B colloidal dispersion. J Antimicrob Chemother **2006**; 57:1153–1160.

6. Stott KE, Le T, Nguyen T, et al. Population Pharmacokinetics and Pharmacodynamics of Itraconazole for Disseminated Infection Caused by Talaromyces marneffei. Antimicrob Agents Chemother 65:e00636-21.

7. WHO fungal priority pathogens list to guide research, development and public health action. Geneva: World Health Organization; 2022. Licence: CC BY-NC-SA 3.0 IGO.

8. Wake RM, Allebone-Salt PE, John LLH, et al. Optimizing the Treatment of Invasive Candidiasis—A Case for Combination Therapy. Open Forum Infect Dis **2024**; 11:ofae072.

9.   Johnson MD, MacDougall C, Ostrosky-Zeichner L, Perfect JR, Rex JH. Combination

antifungal therapy. Antimicrob Agents Chemother **2004**; 48:693–715.

10. Marr KA, Schlamm HT, Herbrecht R, et al. Combination Antifungal Therapy for Invasive

Aspergillosis. Ann Intern Med **2015**; 162:81–89.

11. Shadomy S, Wagner G, Espinel-Ingroff E, Davis BA. In vitro studies with combinations of 5-

fluorocytosine and amphotericin B. Antimicrob Agents Chemother **1975**; 8:117–121.

12. Dupont B, Drouhet E. In vitro synergy and antagonism of antifungal agents against yeast-

like fungi. Postgrad Med J **1979**; 55:683–686.

13. Ghannoum MA, Fu Y, Ibrahim AS, et al. In vitro determination of optimal antifungal

combinations against Cryptococcus neoformans and Candida albicans. Antimicrob Agents

Chemother **1995**; 39:2459–2465.

14. Hamilton JD, Elliott DM. Combined activity of amphotericin B and 5-fluorocytosine against

Cryptococcus neoformans in vitro and in vivo in mice. J Infect Dis **1975**; 131:129–137.

15. Perfect JR, Durack DT. Treatment of Experimental Cryptococcal Meningitis with

Amphotericin B, 5-Fluorocytosine, and Ketoconazole. J Infect Dis **1982**; 146:429–435.

16. Day JN, Chau TTH, Wolbers M, et al. Combination Antifungal Therapy for Cryptococcal

Meningitis. N Engl J Med **2013**; 368:1291–1302.

17. Molloy SF, Kanyama C, Heyderman RS, et al. Antifungal Combinations for Treatment of

Cryptococcal Meningitis in Africa. N Engl J Med **2018**; 378:1004–1017.

18. Stott KE, Ahmadu A, Kajanga C, et al. Population pharmacokinetics and CSF penetration of

flucytosine in adults with HIV-associated cryptococcal meningoencephalitis. J Antimicrob

Chemother **2023**; 78:1015–1022.

19. Sigera LSM, Denning DW. Flucytosine and its clinical usage. Ther Adv Infect Dis **2023**;

10:20499361231161387.

20. Thu NTM, Borda H, Vitsupakorn S, et al. Development and validation of a colorimetric

antifungal susceptibility testing method for the dimorphic fungus Talaromyces marneffei.

Med Mycol **2023**; 61:myad111.

21. Fang L, Liu M, Huang C, et al. MALDI-TOF MS-Based Clustering and Antifungal

Susceptibility Tests of Talaromyces marneffei Isolates from Fujian and Guangxi (China).

Infect Drug Resist **2022**; 15:3449.

22. Zhang J, Liu H, Xi L, Chang YC, Kwon-Chung KJ, Seyedmousavi S. Antifungal

Susceptibility Profiles of Olorofim (Formerly F901318) and Currently Available Systemic

Antifungals against Mold and Yeast Phases of Talaromyces marneffei. Antimicrob Agents

Chemother **2021**; 65:e00256-21.

23. Chang H-J, Huang K-C, Wu C-H. Determination of Sample Size in Using Central Limit

Theorem for Weibull Distribution. Int J Inf Manag Sci **2006**; 17:31–46.

24. Synergism Testing: Broth Microdilution Checkerboard and Broth Macrodilution Methods. In:

Clinical Microbiology Procedures Handbook. John Wiley & Sons, Ltd, 2016: 5.16.1-

5.16.23. Available at:

https://onlinelibrary.wiley.com/doi/abs/10.1128/9781555818814.ch5.16. Accessed 20 April

2022.

25. Klepser ME, Ernst EJ, Lewis RE, Ernst ME, Pfaller MA. Influence of test conditions on

antifungal time-kill curve results: proposal for standardized methods. Antimicrob Agents

Chemother **1998**; 42:1207–1212.

26. Lewis RE, Diekema DJ, Messer SA, Pfaller MA, Klepser ME. Comparison of Etest,

chequerboard dilution and time–kill studies for the detection of synergy or antagonism

between antifungal agents tested against Candida species. J Antimicrob Chemother **2002**;

49:345–351.

27. Nau R, Sörgel F, Eiffert H. Penetration of Drugs through the Blood-Cerebrospinal

Fluid/Blood-Brain Barrier for Treatment of Central Nervous System Infections. Clin

Microbiol Rev **2010**; 23:858–883.

28. Maji A, Soutar CP, Zhang J, et al. Tuning sterol extraction kinetics yields a renal-sparing

polyene antifungal. Nature **2023**; 623:1079–1085.

29. Banerjee D, Umland TC, Panepinto JC. De Novo Pyrimidine Biosynthesis Connects Cell

Integrity to Amphotericin B Susceptibility in Cryptococcus neoformans. mSphere **2016**;

1:10.1128/msphere.00191-16.

30. Christiansen KJ, Bernard EM, Gold JW, Armstrong D. Distribution and activity of

amphotericin B in humans. J Infect Dis **1985**; 152:1037–1043.

31. Stott KE, Beardsley J, Whalley S, et al. Population Pharmacokinetic Model and Meta-

analysis of Outcomes of Amphotericin B Deoxycholate Use in Adults with Cryptococcal

Meningitis. Antimicrob Agents Chemother **2018**; 62:e02526-17.

32. Te Dorsthorst DTA, Verweij PE, Meletiadis J, et al. In Vitro Interaction of Flucytosine

Combined with Amphotericin B or Fluconazole against Thirty-Five Yeast Isolates

Determined by both the Fractional Inhibitory Concentration Index and the Response

Surface Approach. Antimicrob Agents Chemother **2002**; 46:2982–2989.

33. O'Connor L, Livermore J, Sharp AD, et al. Pharmacodynamics of Liposomal Amphotericin B

and Flucytosine for Cryptococcal Meningoencephalitis: Safe and Effective Regimens for

Immunocompromised Patients. J Infect Dis **2013**; 208:351–361.

34. Brouwer AE, Rajanuwong A, Chierakul W, et al. Combination antifungal therapies for HIV-

associated cryptococcal meningitis: a randomised trial. The Lancet **2004**; 363:1764–1767.

35. Thaler M, Bacher J, O'Leary T, Pizzo PA. Evaluation of Single-Drug and Combination

Antifungal Therapy in an Experimental Model of Candidiasis in Rabbits with Prolonged

Neutropenia. J Infect Dis **1988**; 158:80–88.

36. Rabinovich S, Shaw BD, Bryant T, Donta ST. Effect of 5-fluorocytosine and amphotericin B

on Candida albicans infection in mice. J Infect Dis **1974**; 130:28–31.

37. Pappas PG, Kauffman CA, Andes DR, et al. Clinical Practice Guideline for the Management

of Candidiasis: 2016 Update by the Infectious Diseases Society of America. Clin Infect Dis

Off Publ Infect Dis Soc Am **2016**; 62:e1-50.

38. Hope WW, Warn PA, Sharp A, et al. Optimization of the Dosage of Flucytosine in

Combination with Amphotericin B for Disseminated Candidiasis: a Pharmacodynamic

Rationale for Reduced Dosing. Antimicrob Agents Chemother **2007**; 51:3760–3762.

39. Rutakingirwa MK, Skipper CP, Dai B, et al. Evaluating the Use of Lower Dose Flucytosine

for the Treatment of Cryptococcal Meningitis: A Clinical Trial. Clin Infect Dis **2025**; :ciaf432.

 Figures and Tables

Figure 1. Checkerboard assay method demonstrating the in vitro interaction between

amphotericin B and flucytosine against *Talaromyces marneffei*.

The positive control well ("PC", position A1) contains no drug and allows for uninhibited growth.

The negative control well ("NC", position H10) contained no inoculum. A well showing synergy

(fractional inhibitory concentration index, FICI ≤ 0.5) is highlighted in blue with horizontal lines

(position F6). Wells showing indifference (0.5 < FICI ≤ 4) are in blue with purple outline

(positions D4 – D6, E4 – E6, F4, F5). A well showing antagonism (FICI > 4) is in blue with red

outline ("A", position B10).

Created in BioRender. Natesan Sambath, H. (2025) https://BioRender.com/h4r0z3t

<u>Abbreviations:</u> 5FC, flucytosine; AmB, amphotericin B; MIC, minimum inhibitory concentration.

Figure 2. Changes in the minimum inhibitory concentrations of amphotericin B and

flucytosine when tested in combination *versus* alone, and the classification of

combination effect observed for 60 *Talaromyces marneffei* clinical isolates.

(A) and (**B**) show the MICs of AmB and 5FC tested alone and in combination, with the dashed

lines showing the directionality of changes between the MIC alone and in combination for each

isolate. Absolute MIC data follows geometric sequence with a common ratio of 2, i.e., 2-fold

serial dilution, therefore $\log_2$ MIC values were plotted to explain the fold reduction in MICs. **(A)**

Minimum inhibitory concentrations (MIC) at 95% inhibition of amphotericin B (AmB) when tested

alone *versus* in combination with flucytosine (5FC) against 60 *T. marneffei* clinical isolates. The

MIC of AmB decreased three-fold on average when combined with 5FC (Paired t-test, $P <$

0.0001). **(B)** MIC of 5FC tested alone *versus* in combination with AmB against 60 *T. marneffei*

isolates. The MIC of 5FC decreased four-fold on average when combined with AmB (Paired t-

test, $P < 0.0001$). **(C)** Distribution of the fractional inhibitory concentration index (FICI) for AmB

and 5FC combination in 60 *T. marneffei* isolates. Data are presented as individual FICI data

points, with mean and standard deviation.

Abbreviations: 5FC, flucytosine; AmB, amphotericin B; FICI, fractional inhibitory concentration

index; MIC, minimum inhibitory concentration.

Figure 3. Time-kill assay demonstrating the concentration-dependent fungicidal effect of

amphotericin B and the concentration-independent fungistatic effect of flucytosine,

tested alone against *Talaromyces marneffei* isolate 11CN-21-012.

(A) The fungicidal effects of AmB testing at concentrations 0.25 to 2 times the minimum

inhibitory concentration (MIC) of AmB. **(B)** The fungistatic effects of 5FC testing at

concentrations 0.25 to 20 times the MIC of 5FC. AmB showed fungicidal activity, resulting in a

3-log$_{10}$ reduction in colony forming units per milliliter (CFUs/mL) at concentrations of 0.5 or

above the MIC by 96 hours. 5FC resulted in less than a 1 to 2-log$_{10}$ CFUs/mL reduction in the

fungal count over 120 hours and was thus considered fungistatic against *T. marneffei.*

Abbreviations: 5FC, flucytosine; AmB, amphotericin B; MIC, minimum inhibitory concentration;

LOD, limit of detection; CFU, colony forming unit.

Figure 4. Time-kill assay demonstrating the synergy of amphotericin B and flucytosine

combination, tested at sub-MIC concentrations of amphotericin B in combination with

various concentrations of flucytosine against *Talaromyces marneffei* isolate 11CN-21-

012.

Amphotericin B (AmB, red) at 0.25 times the MIC **(A)** and 0.5 times the MIC **(B)** concentrations

were tested in the time kill assay, as these sub-MIC concentrations were not rapidly fungicidal

(see the results presented in **Figure 3**), thus permitting the evaluation of the combination. Both

AmB concentrations were tested with flucytosine (5FC, blue) at 1, 5, 10, and 20 times the MIC.

T. marneffei growth dynamics were measured by colony forming units (CFUs) over time at 0-,

24-, 48-, 72-, 96-, and 120-hours post-inoculation. The no-drug control group (black) was used

to determine growth inhibition. At both AmB concentrations of 0.25 and 0.5 times the MIC, the

AmB and 5FC combination (green) showed a $\geq$ 2-$\log_{10}$ CFUs/mL reduction when compared to

AmB (the more active agent) or 5FC alone, demonstrating a synergistic effect between the

combination. Antagonism was not seen, even at the highest concentration of 5FC over 120

hours.

<u>Abbreviations:</u> 5FC, flucytosine; AmB, amphotericin B; MIC, minimum inhibitory concentration;

LOD, limit of detection; CFU, colony forming unit.

Table 1. Summary of the minimum inhibitory concentrations and the combination effect between amphotericin B and flucytosine against 60 *Talaromyces marneffei* clinical isolates.

	MIC_{95} (µg/mL)		FIC_{AmB}	MIC_{95} (µg/mL)		FIC_{5FC}	FICI $\text{FIC}_{\text{AmB}}+\text{FIC}_{\text{5FC}}$
	AmB	AmB+5FC		5FC	5FC+AmB		
Synergy *(FICI ≤ 0.5), n= 4 (7%)*							
Mean	1.68	0.42	0.25	0.35	0.04	0.14	0.39
	(95% CI: 1.20-2.36)	(95% CI: 0.30-0.59)	(0)	(95% CI: 0.15-0.85)	(95% CI: 0.02-0.09)	(0.08)	(0.08)
Indifference *(0.5 < FICI ≤ 4), n = 56 (93%)*							
Mean	0.64	0.23	0.46	0.27	0.06	0.34	0.80
	(95% CI: 0.55-0.75)	(95% CI: 0.18-0.31)	(0.23)	(95% CI: 0.22-0.34)	(95% CI: 0.04-0.08)	(0.25)	(0.20)
All isolates, *n = 60*							
Mean	0.68	0.24	0.45	0.28	0.06	0.33	0.77
	(95% CI: 0.58-0.80)	(95% CI: 0.19-0.32)	(0.23)	(95% CI: 0.22-0.34)	(95% CI: 0.04-0.08)	(0.25)	(0.22)
Mode	1	0.5	0.5	0.25	0.06	0.5	0.75

The MICs of AmB and 5FC were defined as the lowest drug concentration that resulted in at least 95% inhibition of fungal growth. The geometric means with the 95% confidence intervals are reported for the MICs of AmB and 5FC, alone, and in combination. The arithmetic means with the standard deviations are reported for FICs of AmB and 5FC, and the FICI for 60 isolates.

Abbreviations: 5FC, flucytosine; 95% CI, 95% confidence interval; AmB, amphotericin B; FIC, fractional inhibitory concentration; FICI, fractional inhibitory concentration index; MIC, minimum inhibitory concentration.