FDA-approved drug repurposing in zebrafish identifies thyroid hormone and other compounds
as potential antithrombotics

Murat Yaman[1†], Hongyu Su[2†], Jacqueline K. Lee[1], Allison C. Ferguson[1], Katherine M. Sowell[1],

Jenny Xun[3], David Wu[3], Clayton J. Habiger[1,4], David A. Hanauer[1,5], Martin C. Clasby[6,7], Jason C.

Rech[6,7], Jordan A. Shavit[1,8]*

1. Department of Pediatrics, University of Michigan, Ann Arbor, MI

2. Department of Pharmacology, University of Michigan, Ann Arbor, MI

3. Department of Computer Science, University of Michigan, Ann Arbor, MI

4. Children's Mercy Hospital, Kansas City, MO

5. Department of Learning Health Sciences, University of Michigan, Ann Arbor, MI

6. Vahlteich Medicinal Chemistry Core, University of Michigan, Ann Arbor, MI

7. College of Pharmacy, University of Michigan, Ann Arbor, MI

8. Department of Human Genetics, University of Michigan, Ann Arbor, MI

†Equal contribution

*Correspondence: Jordan A. Shavit, Department of Pediatrics, University of Michigan, Room

8301, Medical Science Research Building III, 1150 West Medical Center Dr, Ann Arbor, MI 48109;

email: jshavit@umich.edu

Abstract word count: 190

Number of figures and tables: 6 figures and 3 tables

Number of supplementary figures and tables: 5 figures and 7 tables

Number of references: 58

Scientific category: Thrombosis

ABSTRACT

Venous thromboembolism (VTE) is a highly prevalent medical condition with limited therapeutic options and an incomplete understanding of its acquired and inherited subtypes. The zebrafish is a model with the benefits of external development, fecundity, optical transparency, and hemostasis that demonstrates conservation with mammals. We utilized zebrafish as a phenotypic screening tool to identify novel therapeutic options for preventing VTE. A library of FDA-approved compounds was screened for suppression of acquired (elevated estrogen) and spontaneous (protein C deficiency) thrombosis. We found that thyroid hormone, receptor tyrosine kinase (RTK) inhibitors, and proton-pump inhibitors (PPIs) effectively modulated levels of thrombosis, particularly in the estrogen-induced model. These also showed a more favorable hemostatic profile than standard therapies, suggesting alternative mechanisms. Genome editing of thyroid hormone receptor proved that thyroid hormone action is on target. A retrospective electronic health record (EHR) analysis found that thyroid-hormone prescriptions in hormonal contraceptive users correlated with a higher VTE risk, potentially limiting direct repurposing but highlighting thyroid signaling as a pathway involved in estrogen-induced thrombosis. Together, these data identify several drug classes that can be tailored to specific subtypes of VTE and help elucidate distinct pathways driving thrombosis.

Key words

Venous thromboembolism, thrombosis, anticoagulant therapies, zebrafish, drug repurposing, FDA-approved compounds, estrogen-induced thrombosis, thyroid signaling, spontaneous thrombosis.

INTRODUCTION

Venous thromboembolism (VTE) includes deep vein thrombosis (DVT) and pulmonary embolism, characterized by the formation of pathologic intravascular blood clots in the venous system, leading to morbidity and mortality. VTE affects approximately 900,000 individuals/year in

the United States, and accounts for 60,000-100,000 deaths, imposing a heavy burden and costs

on healthcare systems.[1,2] The causes of VTE are varied, involving inherited factors such as

protein C (PC for protein, *proc* for gene) deficiency, and elevated estrogen levels from hormonal

treatments like combined hormonal contraceptives (CHCs) and pregnancy.[3] The PC pathway is

a crucial regulator of the coagulation cascade, playing a significant role in preventing intravascular

thrombosis and modulating inflammation.[4] The risk is heightened by factors such as hormonal

therapies, smoking, genetic thrombophilia, family history, and aging.[5-7] VTE is linked to numerous

complications, including clot progression, recurrence, post-thrombotic syndrome, and significant

bleeding due to anticoagulation.

Warfarin, direct oral anticoagulants (DOACs), and thrombolytic therapies are the primary

medications for the management of VTE, but they also pose a substantial risk of major bleeding,

and may carry up to a 9% fatality rate.[8,9] Despite recent advances in safer antithrombotic agents

(DOACs), these medications still parallel the risk profiles of warfarin, with a propensity for

bleeding.[10] DOACs accounted for nearly 40% of emergency department visits due to oral

anticoagulant bleeding, with hospitalization rates similar to those of warfarin.[11] Compounding this

issue is the limited variety of antithrombotic drugs compared to the breadth of pharmacological

classes available for other cardiovascular conditions such as hypertension, heart failure,

hyperlipidemia, and arrhythmia - each having at least 4-5 classes of medications, with between 2

to 10 agents per class. This disparity underscores the pressing medical need for innovative and

safer VTE treatments.[10,12]

To address this, we have leveraged the zebrafish model to investigate coagulation

disorders, which are characterized by rapid development from a single-cell to a fully formed larva

with a functional heart and circulatory system within 72 hours.[13,14] This rapid lifecycle, combined

with high fecundity and optical transparency of embryos, positions zebrafish as an efficient and

cost-effective platform for small molecule high-throughput screening.[13,15-17] This allows screening

of a living vascular system rather than isolated proteins or cells.[18,19] These "mini-organisms" have

been utilized in assays suited to various multiwell-plate formats, allowing for phenotypic-based

screenings and collection of *in vivo* efficacy data, rapidly propagating research from preliminary

screens to Phase I clinical trials.[20,21] The structural and functional homologies between zebrafish

and mammalian coagulation factors facilitate the modeling of diverse VTE pathologies,

encompassing venous thrombosis.[13,14,22,23] Furthermore, zebrafish larvae develop thrombosis in

response to estrogen.[24]

Utilizing models of estrogen-induced (acquired) and spontaneous (genetic) thrombosis,

we have identified 28 FDA-approved compounds from diverse drug families, with multiple agents

modulating thrombosis across both models. This overlap indicates potential shared

pharmacologic pathways across acquired and genetic VTE. Additionally, the identified candidate

compounds were found to have a relatively lower hemostatic risk compared to standard

anticoagulants. Comparative analysis revealed distinct clotting patterns between acquired and

spontaneous thrombosis, suggesting pathological differences. Notably, standard-of-care

therapies failed to achieve a complete anticoagulant effect in the estrogen-induced model. Thyroid

hormone was found to have efficacy via on-target thyroid receptor signaling in both acquired and

inherited thrombosis, which aligns with literature relating hyper/hypothyroidism to VTE risk.[25-29] A

retrospective EHR review revealed that thyroid hormone prescriptions in hormonal contraceptive

users correlates with increased VTE risk. In summary, we have used zebrafish to reveal

pathological differences in thrombosis and limitations of current therapies, which could aid in

developing safer and more effective therapies for VTE tailored for specific VTE subtypes, and

suggesting thyroid signaling as a valuable mechanistic target.

METHODS

Fish strains and husbandry

Zebrafish were raised in accordance with animal care guidelines as approved by the

University of Michigan Animal Care and Use Committee. Two transgenic zebrafish lines were

utilized, *Tg(fabp-fgb-egfp)* and *Tg(proc^{-/-}::fabp-fgb-egfp)*.[30] The former expresses enhanced green

fluorescent protein (eGFP) tagged fibrinogen, allowing for the visualization of thrombosis, while

the latter is also deficient in PC, serving as a model for spontaneous thrombosis.[31] All animal

procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Michigan (Protocol Number: PRO00010679). These experiments were conducted in

accordance with institutional and national ethical guidelines for animal research.

Drug screening

1,600 compounds from the FDA Approved & Pharmacopeia Drug Library

(MedChemExpress, Cat No: HY-L066) were screened for thrombotic effects in zebrafish models

of estrogen-induced (acquired) thrombosis. Embryos were treated at a concentration of 25 µM at

3 dpf. At 5 dpf, clotting in the caudal vein was assessed by an observer blinded to condition using

a semi-quantitative and reproducible grading system ranging from 0 to 3, indicating no thrombosis

through severe thrombosis.

For the acquired model, the *Tg(fabp-fgb-egfp)* line was additionally treated with mestranol

(prodrug of ethinylestradiol, 25 µM) at 4 dpf to induce thrombosis. Three embryos/compound were

tested in this model. For the initial screening, a Z' score was calculated according to the formula

$Z' = 1 - (\frac{3*SD \ of \ sample + 3*SD \ of \ control}{|Mean \ of \ sample - Mean \ of \ control|})$ where SD refers to the standard deviation of the

respective groups.[32] Molecules which passed a 50% reduction in thrombosis underwent

subsequent validation (**Supplemental Table 1**), including dose-response curves for the most

promising candidates (**Supplemental Table 2**) using 4-6 concentrations (max 100 µM) with at

least 12 embryos, and repeated at least twice. In the case of apixaban, we tested concentrations

up to 400 µM but did not explore further because this would have required raising the vehicle

control (DMSO) above 1%, which results in toxicity. To assess spontaneous (genetic) thrombosis,

18-24 embryos per compound were tested without mestranol, and with repetition at least twice.

Embryos were monitored daily for lethality and morphological abnormalities (heartbeat, blood

flow, pericardial edema, body curvature, and developmental delay). Any concentration that

induced toxicity was omitted from analysis.

Dose-response curves were generated by initially transforming the concentration values

to a log10 scale, with zero concentrations adjusted to 0.001 µM. Baseline correction was

performed using the response of DMSO-treated groups as controls. The regression lines for the

genetic thrombosis model were obtained using four-parameter log(agonist) versus response

models. In the acquired model, data normalization was conducted between zero and three,

referencing the average clotting scores of the DMSO and 25 µM mestranol-treated groups,

respectively, following baseline correction. From the resulting dose-response curves, the

inhibitory concentration values corresponding to 30% (IC_{30}) and 50% (IC_{50}) changes in

thrombosis grades were extrapolated. In a parallel study, corrected total cell fluorescence (CTCF)

values were obtained as described to compare the semi-quantitative grading system versus

quantification of the fluorescence signals.[33] All statistical analyses were performed using

GraphPad Prism software (version 9.1.0).

Laser-mediated endothelial injury

Laser-mediated endothelial injury was conducted using an Olympus IX73 inverted

microscope equipped with the Micropoint focusing system (Andor Technology, Belfast, UK) and

a pulsed dye laser, as previously described.[34] Embryos were anesthetized with 0.16 mg/mL

tricaine methanesulfonate at 5 dpf and then immobilized in 0.8% low-melting-point agarose on a

glass coverslip. Targeted ablation of the endothelium was carried out in the posterior cardinal vein

(PCV) at the fifth somite distal to the anal pore by delivering 99 laser pulses with a power level

set to 18 and observed for thrombotic occlusion up to two minutes. In parallel with the endothelial

injury procedure, the level of fluorescent thrombosis from each embryo was also recorded to

confirm that the applied concentrations of the respective drugs were within a range that mitigates

thrombosis.

Knockdown studies using CRISPR/Cas9 gene editing

CRISPR single-guide RNA (sgRNA) sequences targeting early exons were designed with

the Knockout Guide Design Tool (synthego.com). These sgRNA sequences, complexed with the

spCas9 enzyme, were injected into embryos at the 1-cell stage. The success of the injections was

verified by PCR and agarose gel electrophoresis. Sequences of the sgRNAs and PCR primers

are provided in **Supplemental Table 3**.

Electronic health record (EHR) data analyses

This study was approved by the University of Michigan Institutional Review Board

(Protocol Number: HUM00210605). Using the DataDirect EHR cohort discovery and data

extraction software at the University of Michigan, a search was conducted for female patients who

had an outpatient encounter (clinic appointments or emergency room visits) from January 1, 2000

to December 31st, 2019, and were between ages 12-50 at the time of the encounter. Patient data

including smoking history, encounter dates, body mass index (BMI), ICD9 (International

Classification of Diseases) and ICD10 diagnosis codes (billing and/or physician-entered data),

outpatient medications prescribed, and comorbidities which were represented via the Charlson

Comorbidity Index.

The patients were placed in three separate categories based on which medications they

were prescribed during the timeframe of interest. These categories included: (1) both hormonal

contraception (HC) and thyroid medication (desiccated thyroid extract, levothyroxine, or

liothyronine), (2) HC without thyroid medications, or (3) thyroid medications without HC

medications. This determination was based on prescription data and included HC written date

(StartHC), HC end date (EndHC), along with thyroid medication written date (StartThy) and thyroid

medication end date (EndThy). Because liothyronine and desiccated thyroid extract have different

doses than levothyroxine, they were converted to levothyroxine equivalent doses.[35] Intrauterine

contraceptive devices and progesterone only HC were excluded. Overlap between hormonal

contraception and thyroid medication prescriptions was determined by using SQL code (code

logic available upon request).

After patients were placed in these different categories, the rate of thrombosis was

determined using thrombosis-related ICD9 and 10 events. Any diagnosis that was from a

historical source was excluded. Because DataDirect software includes related diagnoses (e.g.

chronic and acute DVT), any non-acute thrombotic event was excluded. If there were multiple

encounters with varying dates, the initial encounter where a thrombotic event occurred was used.

If no BMI or comorbidities were noted during the encounter, the nearest record was used. If there

were no BMI or comorbidity data documented for that patient, the field was left blank. For

comparative purposes, diagnoses were simplified to either DVT, pulmonary embolism,

thrombophlebitis, stroke, abdominal thrombosis, or arterial thrombosis. For example, DVT of the

left lower extremity was simplified to DVT). Full diagnoses codes and equivalent simplified

diagnosis key available upon request.

Image analyses comparing differences in clotting patterns

To examine clotting pattern differences between thrombosis models, we developed

convolutional neural network (CNN) and support vector machine (SVM) algorithms using Python

in a Google Colab environment. In total, 2,072 images from six independent experiments were

analyzed, consisting of 1,066 images from the acquired thrombosis model and 1,006 images from

the spontaneous model (three experiments per model).

For the CNN analysis, built with PyTorch, images were resized to 224 × 224 pixels. Images

containing minimal green fluorescence were excluded by computing the percentage of bright

green pixels, which reduced the dataset to 810 images. These images were later processed

through three convolutional layers and two fully connected layers. The data were then split into

80% training and 20% testing sets. This experiment was run three times, and the performance

metrics were averaged across runs.

In the case of SVM algorithm, built with scikit-learn, all images (without the green-pixel

filtering step) were resized to 15 × 15 pixels, flattened into feature vectors, and split into training

and testing sets.

Both models were evaluated on accuracy, balanced accuracy, area under the curve

(AUC), sensitivity, and specificity. The analytic codes, along with an example use case

walkthrough for running the models on a subset of the dataset, are available in the GitHub

repository (https://github.com/ymnmurat/Zebrafish-Imaging-Classification).

Compound structure-based analysis

The lead compounds identified from the high throughput screen were sorted and binned

based on common structural features and functional groups. Additional compounds were

identified by substructure searching using the Chemical Abstracts Services (CAS) SciFinder

Discovery Platform™ based on structural and functional group similarity.

Statistical analyses

Ordinal scale statistics were employed to test the effects of the respective treatments

based on observed thrombosis grades. The *MASS* library in R (v4.3.3) was utilized to derive

proportional odds logistic regression (polr) models as previously described.[36,37] Time to occlusion

analyses were conducted using multivariate Cox proportional hazards regression models with the

survival library in R.[38,39] Following the development of the respective models, marginal means

were estimated, and Dunnett's multiple test corrections were applied for multiple comparisons

using the *emmeans* library.[40]

RESULTS

Distinct drug classes mitigate estrogen-induced thrombosis

We employed a comprehensive library of 1,600 FDA-approved compounds and pharmacopeia drugs to perform a screen for prevention of thrombosis induced by mestranol, a synthetic estrogen analog used for hormonal contraception (**Figure 1A-B**). The calculated Z' score was 0.4273 (**Figure 1C**), indicating reliable assay performance. In the first pass, a total of 117 compounds showed potency in reducing acquired thrombosis beyond an arbitrarily selected threshold of 50% reduction, according to our grading system. These candidates underwent further validation to confirm their therapeutic potential.

Subsequent analysis involved establishing dose-response relationships and determining potencies via IC_{30} and IC_{50} measurements, while concentrations demonstrating toxicity were omitted. As curves were generated using an ordinal scale (grades 0 to 3), we compared this method to a fully quantitative approach by calculating Corrected Total Cell Fluorescence (CTCF) values from the same images used for grading (**Figure 1D**). In a representative evaluation with elbasvir and L-thyroxine treatments, we observed a high correlation (Spearman's ρ=0.85) between grading and CTCF values, which supports the former system as a practical alternative. However, an earlier dose response shift by the CTCF curve, in comparison to the ordinal scale shift, indicated that the fully quantitative system was more sensitive to wide changes in fluorescence intensity than the grading system (particularly the slope). For instance, a shift in fluorescence levels caused by 5 µM elbasvir were graded as '2,' while CTCF measurements indicated that there was actually more than half reduction in thrombosis. Similarly, 0.06 µM L-thyroxine caused ~40% decrease in thrombosis which only became detectable with the grading system at 0.13 µM.

Among the validated compounds, we observed distinct therapeutic classes; steroid/hormone related, RTK/VEGFR inhibitors, proton-pump inhibitors, neuromodulators, and immune system modulators (**Table 1, Supplemental Table 4**), demonstrating multiple

mechanisms available for prevention of estrogen-induced thrombosis. Interestingly, some of the

drugs such as esomeprazole and the RTK/VEGFR inhibitors posed pro-coagulant effects in the

absence of mestranol, suggesting a competition between the molecular mechanisms is induced

by these compounds and estrogens. Among the candidates, only thyroid hormones were able to

reduce thrombosis at nanomolar concentrations.

Drug-response profiles between the acquired and genetic models overlap

Parallel screening of ~800 compounds conducted in the *proc* knockout line (genetic

model) identified some hits found in the acquired model. Several categories of compounds,

including hormone-related treatments, neuromodulators, proton-pump inhibitors, and RTK

inhibitors, were effective across both, while others were model-specific (**Table 1, Supplemental**

Table 4). Although the two models represent distinct thrombotic processes, this overlap indicates

shared pharmacologic pathways that can be targeted in both acquired and genetic VTE.

Hormone-related compounds, such as the aromatase inhibitor exemestane and thyroid

hormones, demonstrated anticoagulant effects in the *proc* mutant line, while the thyroid receptor

antagonist (compound 1-850) showed only mild activity in estrogen-induced thrombosis and no

clear effect for spontaneous thrombosis. The progestin inhibitor mifepristone was notable for its

procoagulant profile in the genetic model as well. Neuromodulators like meclofenoxate and

lofepramine had a noticeable impact on acquired thrombosis but were less active in the *proc*

mutant. However, a cannabinoid receptor agonist AM-251 and a GABA receptor modulator

etomidate were effective in both.[41,42] Loxapine (dopamine and serotonin receptor antagonist), in

contrast, led to procoagulant effects in the absence of estrogens.[43] Elbasvir, an antiviral drug,

also revealed a strong anticoagulant profile for estrogen-induced thrombosis with no equivalent

effect in the genetic model.

PPIs exhibited divergent effects, with dexlansoprazole demonstrating a mild anticoagulant

shift relative to control in the genetic model, unlike esomeprazole, which was procoagulant.

SCH28080 (a non-standard PPI that structurally deviates from the archetype) was shown to

reduce acquired thrombosis. Overall, these data suggest a thrombosis-modulating role for proton

pumps. Similar to what we observed with esomeprazole, the RTK inhibitors nintedanib and

tivozanib displayed promising profiles in acquired thrombosis, yet procoagulant activity in *proc*

mutant, without any clear changes in the wild-type controls.

Interestingly, the immunomodulatory drugs buparvaquone, edaravone, and malotilate,

along with the antibiotic agents thiabendazole and nalidixic acid, demonstrated antithrombotic

effects in the genetic model, but were ineffective in preventing acquired thrombosis.

Given that standard-of-care anticoagulants, including warfarin and the DOACs, are

commonly prescribed for a wide spectrum of VTE cases, we next evaluated them in both models

and compared their effects to the candidate hits (**Table 1**). Surprisingly, there were minimal effects

on acquired thrombosis, which is concordant with our previous studies.[24,44] In contrast, warfarin

(IC_{50}: 50 nM) and dabigatran etexilate (IC_{50}: 11.72 µM) exhibited robust anticoagulant effects in

the *proc* mutants, consistent with their known inhibitory effects on clotting factors. Apixaban

reduced thrombosis by ~20%, but only at concentrations exceeding 200 µM. These outcomes

present an intricate drug response landscape, with certain agents showing distinct effects that

may align with different pathological thrombosis subtypes.

Upon closer examination of the control embryo images from both models, distinct clotting

patterns were observed (**Figure 2A**). Estrogen exposure resulted in a speckled distribution of

clots. In contrast, images from *proc* mutants exhibited a sprouting pattern with occasional areas

showing dense accumulation. These model-specific drug responses may explain why standard

anticoagulants fail to prevent estrogen-induced thrombosis suggesting the presence of different

pathological VTE subtypes. Using CNN and SVM algorithms, we were also able to distinguish

between the two types of thrombi with high sensitivity and accuracy (**Figure 2B**). The relatively

low specificity of SVM (0.69) was expected, as this approach lacked a preliminary filter to exclude

images with minimal fluorescence hence its deployment on all 2,072 images, including 1,125

graded 0-1 (none to low thrombosis).

Candidate compounds exhibit minimal effects on hemostasis compared with standard

anticoagulants

To further assess hemostatic balance and potential therapeutic efficacy, we applied a

laser-mediated endothelial injury assay for the candidate drugs while simultaneously confirming

effectiveness on thrombosis (**Figure 3**, **Supplemental Figure 1**) in comparison to the standard

anticoagulants. Time to occlusion (TTO) values for the candidate compounds closely aligned with

those of the control. Notably, significant delays in occlusion were observed for the standard

anticoagulants relative to negative control (as previously shown), despite no clear mitigation of

the acquired thrombosis phenotype at the concentrations tested.[22,45] Interestingly, some

candidates, including liothyronine and RTK inhibitors, delayed the occlusion times when estrogen

levels are elevated, yet this effect remained less pronounced than that observed with standard

anticoagulants, both with and without estrogens. These findings suggest that the candidate drugs

may effectively modulate clot formation in the acquired model with a reduced risk of bleeding with

respect to the approved standard anticoagulants that carry this risk inherently.

Limited efficacy of the standard anticoagulants is likely not due to drug-drug interactions

The observations of limited efficacy profiles against estrogen-induced thrombosis

associated with the standard-of-care therapies (apixaban, dabigatran and warfarin) raised the

possibility of potential drug-drug interactions with estrogens or issues with absorption. To address

this, simultaneous assessments of thrombosis levels and time to occlusion were performed

(**Figure 3**, **Supplemental Figure 1)**. Notably, treatments involving the standard anticoagulants,

with and without estrogens, resulted in significant delays in occlusion, suggesting effective

absorption of the standard anticoagulants without reducing thrombosis (**Supplemental Table 5,**

Supplemental Table 6). More striking were the estrogen/standard anticoagulant combination

groups, which exhibited even more pronounced delays in occlusion time. This outcome hinted at

a possible dynamic interaction between the clot-promoting effects of estrogens and the clot-

preventing actions of standard anticoagulants. The absence of delays in occlusion times in

estrogen-only controls further indicated that the mechanisms at play may be more complex than

initially assumed and warrant additional investigation.

Candidate drugs form distinct molecular clusters based on their structures

Based on the obtained compound classes, we further investigated structure-based

similarities between the candidates. Close similarities can be indicative of overlapping

mechanisms of actions of the compounds, and the relevant structures can also serve as a scaffold

for novel drug development strategies in the future. Accordingly, we observed distinct clusters

formed by thyroid hormones, RTK/VEGFR inhibitors and PPIs. Interestingly, laquinimod, an

immunomodulatory drug, also fell into a cluster with the RTK/VEGFR inhibitors, and the

arylhydrocarbon receptor (AhR) inhibitor benvitimod had structural similarities with the thyroid

hormones. These results suggest overlapping mechanisms with the compounds and their

respective clusters.

Terminal phenyl ring iodination is required for antithrombotic activity

Thyroid medications demonstrated potent antithrombotic effects and relatively low

hemostatic risk profiles. We therefore tested additional compounds with core structures similar to

thyroid hormone in both the acquired and genetic thrombosis models (**Table 2**). L-tyrosine, as

well as iodine addition at the 3 and 5 positions of the phenol ring (3,5-diiodo-L-tyrosine dihydrate),

did not affect thrombosis. Interestingly, L-thyronine, a non-iodinated analog of liothyronine, did

not alter thrombosis in either model. Conversely, tiratricol, which retains iodine in the terminal

phenyl ring but lacks the amine on the carboxyl chain, effectively modulated estrogen-induced

and spontaneous thrombosis with nanomolar potency (<100 nM). These observations indicate

that iodine groups at the terminal phenyl rings, functionally important residues in thyroid hormone,

play a major role in the ability of thyroid hormone to modulate thrombotic events.

Antithrombotic effects of thyroid hormone are on-target through known receptors

The iodine requirement suggested that thyroid hormone was likely mediating

antithrombotic effects through a thyroid hormone receptor (THR). Since medications can have

off-target effects, we investigated whether the thyroid hormone antithrombotic actions were

mediated via canonical THRs. We used CRISPR/Cas9 to knock down all known thyroid receptor

orthologs (*thr*) (**Figure 4**). In the controls (DMSO) and thyroid hormone (L-thyroxine) alone, there

were no alterations in thrombosis. However, the ability of L-thyroxine to reverse estrogen-induced

thrombosis was lost in the knockdown. Similarly, tiratricol was not able to prevent the estrogen-

induced phenotype under *thr* knockdown, strongly implicating thyroid receptor signaling

(**Supplemental Figure 2**). This pattern was also observed in the genetic VTE model, where L-

thyroxine significantly lost its antithrombotic efficacy following receptor knockdown, reinforcing

that these effects are receptor-mediated (**Supplemental Figure 3**).

We next explored whether the increased TTO observed with thyroid hormones was

mediated through its receptor (**Figure 5**; **Supplemental Figure 4**). Upon receptor knockdown,

the TTO values closely approximated baseline, suggesting that the effect is on target.

Interestingly, a marginal delay in TTO levels was still observed in knockdown groups that were

treated with L-thyroxine in combination with elevated estrogen levels or protein C deficiency,

indicating residual receptor activity after the *thr* knockdown.

***thrab* is the major paralog mediating antithrombotic activity of thyroid hormone**

We next tested each *thr* gene individually to determine which one(s) mediate the

antithrombotic effect. We found that L-thyroxine treatment in *thraa* and *thrb* knockdown groups

could still rescue acquired thrombosis similarly to the wild-type controls (**Figure 6**). However, L-

thyroxine treatments failed to reduce the increased thrombosis levels in the *thrab* knockdown and

the triple thyroid receptor knockdown groups, indicating that *thrab* mediates the antithrombotic

effect.

Electronic health records indicate clinical relevance of thyroid levels in estrogen-induced

thrombosis

To determine whether our findings were clinically relevant, we undertook an analysis of

University of Michigan EHR data for 47,710 subjects on hormonal contraception (HC) without

concurrent thyroid medications (**Table 3**). 321 (0.67%) developed thrombosis in comparison to

27 of 2,101 (1.27%) who were concurrently using HC and thyroid medication. This comparison

yielded an odds ratio of 1.9 for thyroid hormone medication plus HC compared to HC alone, with

a number needed to harm (NNH) of 167. Additionally, 30,921 patients were taking thyroid

medication without HC, of which 260 (0.83%) developed thrombosis. Comparing this group to

those taking both HC and thyroid medication resulted in an odds ratio of 1.5 for thrombosis and

an NNH of 227.

Notably, there were some differences in the patient characteristics (**Supplemental Table**

7). For example, the patients in the thyroid medication only group were older and had higher BMIs.

Smoking status was similar between the HC alone group and the HC with thyroid medication

group, but there were more patients with unknown smoking status in the thyroid medication alone

group. Interestingly, the estrogen dose was higher in the HC with thyroid medication group

compared to the HC alone group, while the levothyroxine equivalent dose was lower in the HC

with thyroid medication group compared to the thyroid medication alone group.

Comorbidities were more prevalent in the HC with thyroid medication group compared to

either HC or thyroid medication alone. Chronic pulmonary disease, malignancy, and diabetes

were observed more in the patients who use HC along with thyroid medications. Rates of liver

diseases and metastatic solid tumors were also high across patients using thyroid medications

irrespective of HC status (**Supplemental Figure 5**). As malignancy is a well-known prothrombotic

state, a subanalysis was performed wherein patients with recorded cancer (labeled malignancy

or solid tumor) were excluded. Ultimately, this subanalysis did not reveal a significant difference

in odds ratio compared to the original (data not shown).

As observations further supported the clinical relevance of thyroid hormones in thrombosis

risk, it is also important to note some discrepancies. Our experimental data suggest that thyroid

medications may be preventive for thrombosis. However, EHR data from 27 individuals who were

recorded as using oral contraceptives during thyroid therapies showed higher thrombosis rates.

This was in comparison to those who reported using either oral contraceptives or thyroid therapies

alone within the same timeframe.

DISCUSSION

VTE is a common medical condition with limited treatment options, many of which come

with a high risk of significant bleeding. Moreover, the thrombotic effect of estrogen therapy has

been recognized for many years, yet we still know little about the underlying mechanisms and

how well its pathology aligns with conventional thrombosis. To address the need for novel

antithrombotics, we utilized a robust zebrafish thrombosis reporter line, enabling efficient

screening of 1,600 small molecules for prevention of estrogen-induced thrombosis. The semi-

quantitative grading system by blinded observers closely approximated quantitative fluorescence

readings, offering a practical solution for rapidly and reliably identifying promising drug candidates

and determining meaningful IC_{30} and IC_{50} values. While effective, automating the fluorescence

quantitation process from embryo images could significantly augment our capabilities. Such

automation strategies could include removing autofluorescence, accurately capturing signals from

each individual embryo, and enhancing signal precision. These advancements are warranted for

further studies to better detect subtle changes in thrombosis levels at higher grades and fully

exploit the throughput potential of zebrafish thrombosis models.

Using IC_{30} values, we captured drugs capable of marginally shifting thrombosis grades,

helping us identify a broad range of potential drug families for future antithrombotic development.

In addition, IC_{50} values helped prioritize the small molecules with the most significant effects on

thrombosis and identified distinct mitigating drug classes. Intriguingly, standard-of-care

anticoagulants failed to prevent estrogen-induced thrombosis in the regimen used in this study,

despite our previous findings that they completely block induced thrombosis secondary to

endothelial injury.[22,45] This suggests that the impact of estrogen-induced thrombosis may be

partially regulated by other mechanisms besides the traditional coagulation pathway. In contrast,

we have observed that dosing regimen and timing are important and warfarin can partially prevent

thrombosis when it is initiated at least two days prior to the estrogen.[24] This is likely due to the

known delayed onset of warfarin's effect given the differential half-lives of the vitamin K dependent

coagulation factors.[46] While differences in study design limit direct comparisons, these results

suggest that the duration of estrogen exposure and the timing of anticoagulant initiation are

important determinants of efficacy. Nevertheless, in both studies standard-of-care anticoagulant

efficacy was markedly reduced or incomplete against the extended estrogen exposure regimen,

whereas several of the candidate compounds we identified maintained strong antithrombotic

activity under the same conditions. This contrast suggests that the drug classes observed in this

study may offer more effective therapeutic strategies for estrogen-induced thrombosis. Notably,

among these drug classes, hormone-related compounds, especially thyroid hormones, showed

strong antithrombotic effects at very low concentrations.

There were a number of hits that had overlapping effects on inherited and acquired

thrombosis, while others affected only one system (**Table 1**, **Figure 2**). Hormone-related

compounds such as exemestane and thyroid hormones demonstrated antithrombotic effects in

both models, indicating potential mechanistic overlap. The candidate hits that were effective in

both models (e.g., hormone-related compounds, some neuromodulators, and PPIs) suggest that

their effects in estrogen-induced thrombosis are not strictly due to a potential drug-drug interaction

with estrogen; rather, they may impart thrombosis-modulating effects independently. However,

several antiviral, antibiotic, and immunomodulatory drugs, as well as some neuromodulators,

were effective only in one type of thrombosis. PPIs and RTK/VEGFR inhibitors showed

antithrombotic effects in preventing estrogen-induced thrombosis but exhibited prothrombotic

profiles in the absence of mestranol, suggesting a potential competition taking place between

these drugs and estrogens. The contrasting activities of esomeprazole and omeprazole might be

explained by the higher bioavailability of the former, as opposed to the latter which is a racemic

mixture that includes esomeprazole.[47]

475   Utilizing both estrogen-induced and genetic thrombosis in zebrafish, we identified

distinctive features between these models. This supports the notion that there may be further

distinctions both within and between inherited and some forms of acquired thrombosis, and how

they arise. Recent studies suggest the possibility that estrogen-induced thrombosis might consist

of transglutaminase cross-linked fibrin(ogen), instead of fibrin. Transglutaminases are membrane-

bound proteins involved in cross-linking extracellular matrix components. They are involved in

numerous physiological and pathological processes including wound healing and angiogenesis.[48-

50] Lisetto and colleagues reported a high degree of conservation between zebrafish and human

transglutaminases. They observed speckled localization of transglutaminase activity, particularly

around the vasculature by the caudal veins of the tail region. Interestingly, this pattern coincides

with the distribution of estrogen-induced thrombosis that we observe.[51] It has long been known

that non-factor XIII transglutaminases crosslink fibrinogen ex vivo,[52] and a recent study

demonstrated that tissue transglutaminase modifies FXIII-directed fibrin(ogen) cross-linking in

vivo.[53]

489   In parallel with measuring thrombosis level changes and evaluating potential therapeutic

efficacy, we tested the candidate drugs' effects on hemostasis and found that clotting times

closely aligned with controls. In contrast, standard anticoagulants resulted in significant delays in

occlusion, as we have previously shown, without mitigating estrogen-induced thrombosis.[22,45]

These findings suggest that the candidate drugs could potentially modulate clot formation in the

acquired model without the hemostatic risk inherent in approved anticoagulants.

Given the lack of efficacy in the estrogen-induced thrombosis model, we investigated the

possibility of a mestranol interaction that might reduce the anticoagulant effects of standard blood

thinners. Our simultaneous assessments of induced thrombosis and time to occlusion indicated

that the limited efficacy was likely not due to such interactions, as anticoagulant treatments still

resulted in significant delays in occlusion. It was surprising to see that mestranol combined with

the DOACs led to further increases in time to occlusion, suggesting a potentiation of the

anticoagulant effect. This phenomenon is reminiscent of our past work on antithrombin deficiency

in zebrafish.[22] We found that loss of antithrombin, which should result in a prothrombotic state,

led to a delayed hemostatic profile, which was rescued by infusion of fibrinogen.[22] This suggested

a disseminated intravascular coagulation-like state, where there is an overall consumption of

fibrinogen due to unchecked thrombin activity, although the effects we observe in this study are

milder.

Though it was not as severe, a degree of bleeding risk was observed across some of the

candidate drugs as evidenced by delayed occlusion in combination with estrogens, including

liothyronine, mifepristone, RTK inhibitors, neuromodulators like etomidate and AM-251, and the

immunomodulatory drug laquinimod. In contrast, exemestane, L-thyroxine, and PPIs like

esomeprazole and dexlansoprazole exhibited normal occlusion. These findings highlight potential

increased hemostatic risks for patients with elevated estrogen who are using the former group of

agents.

Thyroid hormone medications demonstrated promising nanomolar efficacy and limited

hemostatic risks. Interestingly, L-thyronine, which is structurally similar to thyroid hormone but

does not have iodine, or L-thyrosine that lacks the terminal phenol ring present in thyroid hormone,

did not exhibit the antithrombotic effect. These structural nuances suggest that the presence of

iodine on the terminal ring, rather than the inner one, is crucial in the observed antithrombotic

effects. Additionally, the carboxyl chain in tiratricol, as opposed to the amine moiety in thyroid

hormone, enhances the potency of the former. We used CRISPR-mediated knockdown to test if

the thyroid hormone effect is on-target through a known receptor. Zebrafish have two paralogs

for the human thyroid receptor alpha (*THRA*), *thraa* and *thrab*, and we found that *thrab* is the

major paralog required for antithrombotic activity. This suggests that THRA modulation could be

an important target in humans for improving therapeutic strategies in estrogen-induced

thrombosis. Both estrogen and thyroid signaling are known to influence thromboembolism.[25-28,54-

58] Thus, thyroid hormone signaling could offer an alternative route for thrombosis prevention.

Further studies are warranted to elucidate the exact mechanisms underlying its involvement in

mediating thrombosis. Knockouts could be also employed to investigate the marginal delays in

TTO values that persisted after *thr* knockdown, indicating residual receptor activity from existing

THR levels in the embryos.

While our experimental studies suggest that thyroid medications may prevent thrombosis,

EHR data indicate higher thrombosis rates among individuals using thyroid agonists with oral

contraceptives, compared to those not using them concurrently. This discrepancy might be

explained by the low baseline incidence of thrombosis observed in such individuals (27 cases

among 2,101 patients), non-reported comorbidities, and medications not included in our analysis

leading to alterations in hemostatic balance that affect their risk. Additionally, review of the

literature indicates a higher prevalence of thrombosis when there is an imbalance or pathology in

the thyroid system.[25-28] However, it is unclear whether thrombosis occurs prior to treatment or

persists even when on thyroid medications. Hormonal contraceptives can increase thyroid

stimulating hormone (TSH) and thyroxine binding globulin without changing free thyroxine

levels.[29] In contrast, excess thyroid hormone exposures in our studies mimic a hyperthyroid state

which could be a factor for preventing estrogen-induced thrombosis. The EHR data do not

explicitly indicate whether the patients on thyroid hormone therapies are in a hyper- or euthyroid

state by the time of a thrombotic event. Moreover, other health conditions (e.g., cancer),

medications, or genetic predisposition, could lead to a pro-thrombotic state and/or affect

interactions with thyroid medications and estrogens. This creates a complex interplay that is much

less controlled than our zebrafish models. To gain a better understanding, analysis using a larger

population size and higher-resolution data from multicenter datasets is necessary. Further

zebrafish mechanistic studies are also essential to clarify causal links, as would a prospective

human study controlling and accounting for health status, thyroid medication use, oral

contraceptive use, and hormone levels.

In summary, we observed distinct drug-response profiles and possible pathological

differences between genetic and acquired thrombosis. Notably, PPIs, RTK/VEGFR inhibitors, and

hormone-related compounds, emerged with efficacy against estrogen-induced thrombosis.

Among them, thyroid hormones showed on-target effects with nanomolar potency in vivo, but

observational EHR data linking hormonal contraceptive-related thrombosis and thyroid imbalance

are still preliminary. Nonetheless, THR signaling stands out as a promising target for future

mechanistic studies and drug development.

Acknowledgements

This work was supported by National Institutes of Health (NIH) grants R01 ES032255 and R35

HL150784, and by the University of Michigan Frankel Cardiovascular Center's Michigan Biological

Research Initiative on Sex Differences in Cardiovascular Disease (M-BRISC) program (J.A.S.).

J.A.S. is the Henry and Mala Dorfman Family Professor of Pediatric Hematology/Oncology. We

would like to thank Drs. Chris Andrews and Corey Powell of CSCAR (Consulting for Statistics,

Computing and Analytics Research) for valuable discussions on statistical analyses. Additionally,

we extend our gratitude to Dr. Jasmine Luzum for her expert insights on the precision medicine

landscape for VTE therapies and management, and Donald McDonnell for helpful discussions.

Authorship Contributions

M.Y. and H.S. designed and performed research, analyzed data, and wrote the manuscript. M.Y.

and H.S. are co-first authors. The order of co-first authorship was determined based on the overall

scope and breadth of their respective contributions to the study. M.Y. led the comprehensive drug

screening efforts and coordinated the integration of experimental work across study segments.

H.S. directed and conducted the thyroid hormone-related investigations and contributed

substantially to manuscript development. J.K.L. and C.J.H. designed and performed research,

and analyzed data, J.X. and D.W. performed research and analyzed data, A.C.F. and K.M.S.

performed research, D.A.H., M.C.C., and J.R. supervised research and edited the manuscript,

and J.A.S. designed and supervised research, and edited the manuscript.

Conflict of interest Disclosures

J.A.S. has been a consultant for Sanofi, NovoNordisk, Biomarin, Takeda, Pfizer, Genentech,

CSL Behring, and Medexus. The other authors declare no relevant conflicts of interest.

Figure 1: Drug library screening in zebrafish models of thrombosis identifies potential anti-thrombotic agents. (A) Transgenic zebrafish models expressing eGFP-tagged fibrinogen were employed to semi-quantitatively assess thrombosis levels in the caudal veins using an ordinal scale from grades 0 to 3 (arrows indicate areas of thrombosis). Imaging and drug exposure protocols were standardized using a 96-well plate format, with 3 embryos per well. (B) Both acquired (estrogen-induced) and genetic (*proc* mutant/spontaneous) thrombosis models were subjected to drug treatments at 3 dpf. Thrombosis levels were subsequently graded at 5 dpf. For the acquired model, thrombosis was triggered by exposure to 25 µM mestranol at 4 dpf. (C) Screening in the acquired model evaluated 1,600 compounds at a concentration of 25 µM. The mean thrombosis level for each treatment (according to the grading scale in (A)) is depicted as a percentage change. An arbitrary threshold was set at 50% reduction which identified 117 hits with a Z' score of 0.4273. (D) Representative dose-response curves for the acquired model. Thrombosis scores were obtained for each treatment quantitatively (CTCF) and semi-quantitatively (ordinal scale grades 0-3) from the same images. Four-parameter log(agonist) vs. response curves were modeled based on the concentrations of elbasvir and L-thyroxine on the x-axis and thrombosis percentages on the y-axis, demonstrating a close approximation between qualitative and quantitative measurements. Spearman's correlation coefficient (*ρ*) for each group was 0.85. Error bars represent 95% confidence intervals. All data were collected by an observer blinded to the condition.

A	Acquired model	Genetic model

B	Metrics	Machine Learning Models	
		CNN	SVM
	Sensitivity	0.90	0.97
	Specificity	0.89	0.69
	Accuracy	0.90	0.83
	Area Under Curve (AUC)	0.90	0.83

Figure 2: Image analyses of the acquired and genetic models of thrombosis identify distinct clotting patterns. (A) Speckled pattern of thrombus distribution was observed in the images obtained from the acquired model, while a sprouting pattern with denser fluorescence signals was noted in the genetic model. (B) To test these differences, we developed convolutional neural networks (CNN) and support vector machine (SVM) models that distinguished between the two based on fluorescence images, achieving a high degree of sensitivity and accuracy using images from six independent experiments. Scale bars, 100 µm. All data were analyzed by an observer blinded to the condition.

Figure 3: Candidate compounds exhibit reduced hemostatic risk compared to standard anticoagulants. Time to occlusion (TTO) values for candidate compounds were assessed by laser-mediated endothelial injury and consistent with DMSO control, indicating a lower hemostatic risk compared to standard-of-care therapies. Some candidates increased occlusion times in the presence of estrogens, but this effect was less pronounced than that observed with standard anticoagulants, both with and without estrogens. In the box plots, median values are indicated in blue, and mean TTO values for each treatment are marked with a diamond. Statistical analyses were performed using multivariate Cox proportional hazards regression models with the *survival* and *emmeans* libraries in R (v4.3.3). ****: p<0.0001, **: p<0.01, *: p<0.05, ns: not significant. All data were collected by an observer blinded to the condition.

Figure 4: The antithrombotic effect of thyroid hormone is mediated through known receptors. CRISPR sgRNA/Cas9 complexes were injected into single-cell embryos that were then treated with L-thyroxine (0.6 μM) at 3 dpf and mestranol (25 μM) at 4 dpf. Statistical analyses were performed using proportional odds logistic regression (polr) models with MASS and *emmeans* libraries in R (v4.3.3). ****: $p < 0.0001$, ns: not significant. wt: wild-type (uninjected *fabp-fgb-egfp* control group), *-thr*: combined knockdown of all thyroid hormone receptor paralogs *thraa*, *thrab*, and *thrb*. Magenta borders indicate mestranol treated groups. All data were collected by an observer blinded to the condition. Sample sizes for each group, corresponding to the thrombosis grades, are indicated within their respective stack.

Genotype	wt	wt	wt	wt	-thr	-thr	-thr	-thr
L-thyroxine	-	-	+	+	-	-	+	+
Mestranol	-	+	-	+	-	+	-	+
n =	41	41	41	34	57	58	54	55

Figure 5: Thyroid hormone modulates hemostasis through known receptors in the estrogen-induced thrombosis model. CRISPR sgRNA/Cas9 complexes were injected into single-cell embryos that were then treated with L-thyroxine (0.6 µM) at 3 dpf, and mestranol (25 µM) at 4 dpf, followed by assessment of laser-mediated endothelial injury at 5 dpf. Increased TTO observed in wild-type embryos treated with thyroid hormone was diminished upon *thr* knockdown, indicating involvement of thyroid receptor signaling. Increased TTO levels were still present in the L-thyroxine-mestranol combination groups. ****: $p < 0.0001$, ***: $p < 0.001$, **: $p < 0.01$, *: $p < 0.05$, ns: not significant. wt: wild-type (uninjected *fabp-fgb-egfp* control group), *-thr*: combined knockdown of all thyroid hormone receptor paralogs *thraa*, *thrab*, and *thrb*. All data were collected by an observer blinded to the condition.

Figure 6: *thrab* receptor modulates the antithrombotic effect of thyroid hormone. After
CRISPR-mediated individual knockdown of each canonical thyroid hormone receptor (*-thraa*, *-thrab*, and *-thrb*) or combination (*-thr*), embryos were co-treated with L-thyroxine (3 µM) and
mestranol (25 µM) at 4 dpf and evaluated 24 hours later. Magenta borders indicate mestranol
treated groups. ****: $p < 0.0001$, *: $p < 0.05$, ns: not significant. All data were collected by an
observer blinded to the condition. Sample sizes for each group, corresponding to the thrombosis
grades, are indicated within their respective stack.

658 **TABLES**

Compounds	IC$_{30}$		Compounds	IC$_{30}$	
	Acquired	Genetic		Acquired	Genetic
AM-251	6.72	12.87	Exemestane	2.95	13.72
Meclofenoxate	6.03	No effect	Mestranol*	8.47[#]	11.91[#]
Loxapine	19.4[#]	No effect[#]	1-850	6.07	No effect
Lofepramine	6.04	No effect	Pantoprazole	14.27	No effect
Etomidate	23.08	18.94	Dexlansoprazole	10.69	5.19
Laquinimod	1.43	10.01	Omeprazole	14.57	No effect
Bendazac	1.51	No effect	Esomeprazole*	8.38	5.16
Tiaprofenic acid	2.63	No effect	SCH28080	6.11	No effect
Buparvaquone	No effect	0.69	Ilaprazole	14.79	No effect
Edaravone	No effect	9.12	Elbasvir	1.35	No effect
Malotilate	No effect	24.05	Thiabendazole	No effect	14.34
Belinostat	47.45	No effect	Nalidixic acid	No effect	6.34
Diphenylcyclopropenone	7.89	No effect	Indole-3-carboxylic acid	7.89	No effect
Tivozanib*	0.67	1.68	Benvitimod	2.82	No effect
Nintedanib*	6.93	4.06	Phenindione	42.82	9.64
L-Thyroxine	0.2	0.15	Apixaban	No effect	No effect
Liothyronine	0.28	0.34	Warfarin	No effect	0.02
Mifepristone*	7.44	10.10	Dabigatran etexilate	No effect	4.58
			Rivaroxaban	No effect	No effect

Table 1: IC$_{30}$ values for the candidate compounds tested on acquired (estrogen-induced)
and genetic (spontaneous/PC deficient) models of thrombosis. The values (µM) were
obtained from dose response curves. No effect, no IC$_{30}$ detected from the dose response curves.
Hits that are not concordant between the models are highlighted in gray. #: IC$_{30}$ values are derived
from the procoagulant profiles of the respective drugs in the acquired model with no mestranol
treatment. *: compounds with procoagulant effects in the genetic model.

Compounds	Structures	IC$_{30}$ (µM)		IC$_{50}$ (µM)	
		Acquired	Genetic	Acquired	Genetic
L-thyroxine		0.2	0.15	0.83	0.32
Liothyronine		0.28	0.34	0.36	0.8
L-tyrosine		No effect	No effect	No effect	No effect
L-thyronine		No effect	No effect	No effect	No effect
3,5-diiodo-L-tyrosine		No effect	No effect	No effect	No effect
3,3',5-triiodothyroacetic acid (tiratricol)		0.09	<0.1	0.24	<0.1

Table 2: IC$_{30}$ and IC$_{50}$ values for thyroid hormone-related structures, highlighting the importance of iodine groups in antithrombotic effects. #, IC$_{30}$/IC$_{50}$ values are derived from the procoagulant profiles in the acquired model with no additional estrogen treatments by 4 dpf. *, compounds with procoagulant effects in the genetic model.

University of Michigan Data			
	HC alone	**HC with Thyroid Medication**	**Thyroid Medication Alone**
All Patients			
Thrombosis	321	27	260
All	47,710	2,101	30,921
Percentage	**0.67%**	**1.27%**	**0.83%**
Odds Ratio	**1.9 (1.29-2.82)**		
		1.5 (1.04-2.26)	
Duration			
Days of HC	674 (668-681)	557 (528-585)	
Days of Thyroid Medication		557 (528-585)	847 (833 - 860)

670 **Table 3: Significant increase in thrombosis rates for the combination HC and thyroid**
671 **medication group.** Patients with reported concordant use of HC and thyroid medications have
672 statistically significant increase in thrombosis rates.

Supplemental Figure 1: Thrombosis grades confirm effective drug concentrations in relation to TTO levels in Figure 3. The concentrations of candidate drugs used in the laser injury experiments in Figure 3 were confirmed to be within an effective range for modulating thrombosis. Thrombosis grades were collected simultaneously with the TTO evaluations from the same embryos. Magenta borders indicate treatments that included mestranol (25 µM) exposure. ****: p<0.0001, **: p<0.01, *: p<0.05, ns: not significant (p-values are calculated against DMSO and estrogen controls, black and magenta colored asterisks, respectively). Sample sizes for each group, corresponding to the thrombosis grades, are indicated within their respective stack.

Supplemental Figure 2: The antithrombotic effect of thyroid hormone analog (tiratricol) is mediated through known receptors. CRISPR sgRNA/Cas9 complexes were injected into single-cell embryos that were then treated with tiratricol (0.1 μM) at 3 dpf and mestranol (25 μM) at 4 dpf. Statistical analyses were performed using proportional odds logistic regression (polr) models with MASS and *emmeans* libraries in R (v4.3.3). To prevent infinitesimal values that would be generated by the regression models, each thrombosis grade group was incremented by one count across all treatment groups. ****: $p < 0.0001$, ***: $p < 0.001$. wt: wild-type (uninjected *fabp-fgb-egfp* control group), *-thr*: combined knockdown of all thyroid hormone receptor paralogs *thraa*, *thrab*, and *thrb*. Magenta borders indicate mestranol treated groups. All data were collected by an observer blinded to the condition. Sample sizes for each group, corresponding to the thrombosis grades, are indicated within their respective stack.

Supplemental Figure 3: The antithrombotic effect of thyroid hormone is mediated through known receptors in the genetic VTE model. CRISPR sgRNA/Cas9 complexes were injected into single-cell embryos that were then treated with L-thyroxine (0.6 µM) at 3 dpf and observed at 5 dpf. Thrombosis grades were collected simultaneously with the TTO evaluations, ensuring consistent sample use as presented in **Supplemental Figure 4**. Statistical analyses were performed using proportional odds logistic regression (*polr*) models with MASS and *emmeans* libraries in R (v4.3.3). ****: $p < 0.0001$, *: $p < 0.05$. *-proc*: genetic thrombosis line (*proc^{-/-}::fabp-fgb-egfp* control group), *-thr*: combined knockdown of all thyroid hormone receptor paralogs *thraa*, *thrab*, and *thrb*. All data were collected by an observer blinded to the condition. Sample sizes for each group, corresponding to the thrombosis grades, are indicated within their respective stack.

Supplemental Figure 4: Thyroid hormone modulates hemostasis through known receptors in the genetic model. Following thrombosis grading (**Supplemental Figure 3**), viable embryos underwent laser-mediated endothelial injury at 5 dpf. Increased TTO observed in genetic thrombosis treated with thyroid hormone was diminished upon *thr* knockdown, supporting involvement of thyroid receptor signaling. A marginal TTO delay in the *-thr* group with L-thyroxine treatment suggests residual receptor activity across *thr* knockdown samples. ***: p<0.001, *: p<0.05, ns: not significant. *-proc* (genetic thrombosis line: *proc⁻ᐟ⁻::fabp-fgb-egfp* control group), *-thr*, combined knockdown of all thyroid hormone receptor paralogs *thraa*, *thrab*, and *thrb*. All data were collected by an observer blinded to the condition.

Supplemental Figure 5: Increased incidence of comorbidities in the EHR dataset for the
HC with thyroid medication and thyroid medication alone groups. Notably, the HC with
thyroid medication group had higher rates of chronic pulmonary disease, malignancy, and
diabetes. The thyroid medication alone group showed increased incidence of mild liver disease,
hemiplegia, and metastatic solid tumors, otherwise there were no statistical differences between
the groups in specific comorbidities.

Plate No & Well Position	Drug Name	Plate No & Well Position	Drug Name	Plate No & Well Position	Drug Name
1_8A	Everolimus	7_3D	Methotrexate	13_4H	p-Toluic acid
1_8H	Belinostat	7_5D	Amenamevir	13_5B	Alosetron (Hydrochloride)
1_9A	Sorafenib Tosylate	7_6F	Thalidomide	13_7G	Crizotinib (hydrochloride)
1_10A	Rapamycin	7_8G	Entacapone	13_8D	Crizotinib
1_10D	Telaprevir	7_10C	Rolapitant	13_11G	Desfesoterodine
1_10F	Rilmenidine hemifumarate	8_2F	Opicapone	14_6D	Liothyronine (sodium)
1_11A	Ilaprazole	8_4H	Avermectin B1	14_7E	Malotilate
2_5B	Daclatasvir dihydrochloride	8_7B	Repaglinide	14_7G	Cefotaxime (sodium salt)
2_5E	Dapagliflozin	9_11D	Elbasvir	14_8A	Liothyronine
2_7A	Tiaprofenic acid	10_2D	Iguratimod	14_9D	Sapropterin (dihydrochloride)
2_8E	Daclatasvir	10_5D	Fluralaner	14_10C	Docetaxel
2_10E	Niraparib hydrochloride	10_6D	Idarubicin (hydrochloride)	15_5H	Bendamustine (hydrochloride)
2_11E	Neflamapimod	10_9H	Ziprasidone (hydrochloride monohydrate)	15_7D	Acarbose
3_2D	Promestriene	10_11D	Racecadotril	15_7F	Omeprazole (sodium)
3_2E	Nintedanib esylate	11_4G	Pantoprazole (sodium hydrate)	15_10C	Tropisetron
3_4D	Valemetostat (tosylate)	11_3A	Meclofenoxate (hydrochloride)	15_10D	Etomidate (hydrochloride)
3_6C	Lenvatinib	11_4G	Pantoprazole (sodium hydrate)	16_3D	Milnacipran ((1S-cis) hydrochloride)
3_8G	Delamanid	11_6A	Atosiban (acetate)	16_5B	Detomidine (hydrochloride)
3_8H	Tivozanib	11_8A	Ampiroxicam	16_6A	Lithocholic acid
3_9D	Ripretinib	11_10E	Parecoxib (Sodium)	16_10G	Chlorothiazide
3_11G	Enarodustat	11_11A	Boldenone Undecylenate	17_8D	Oxytetracycline
4_6A	Velpatasvir	11_11H	Bendazac	18_6D	Liranaftate
4_7C	Idelalisib	11_5D	Pantoprazole (sodium)	18_10B	Bupivacaine (hydrochloride)
4_8A	Bempedoic acid	11_4G	Pantoprazole (sodium hydrate)	18_10C	Nicorandil
4_9D	Dantrolene (sodium hemiheptahydrate)	12_2F	L-Thyroxine (sodium salt)	19_2A	Loperamide (hydrochloride)
4_11B	Rifamycin S	12_5A	Conivaptan (hydrochloride)	19_2H	Clomipramine (hydrochloride)
4_11G	Laquinimod	12_5E	Dalbavancin (hydrochloride)	19_3H	Phenoxybenzamine (hydrochloride)
5_2E	Fostamatinib (disodium hexahydrate)	12_6D	L-Thyroxine	19_5F	Ciclopirox (olamine)
5_5A	Lomustine	12_7A	Ornidazole (Levo-)	19_7C	Ornidazole
5_5C	Deflazacort	12_7G	Pleconaril	19_9B	Niclosamide
5_5F	Cladribine	12_9A	Glecaprevir	19_11B	Domperidone
5_6A	Cytarabine	12_9C	Filgotinib	20_3D	Sodium nitroprusside
5_6D	Cyproterone acetate	12_10E	L-Hydroxyproline	20_3H	Naftifine (hydrochloride)
5_7B	Rupatadine (Fumarate)	12_10G	trans-Tranilast	20_4H	Ampicillin (sodium)
5_8E	Exemestane	12_11A	Indole-3-carboxylic acid	20_5G	Tafluprost
5_10A	Anastrozole	12_11B	Quercetin	20_7D	Metformin
5_11B	Telotristat ethyl	12_11G	Hemin	20_7H	Atorvastatin
6_9H	Pirarubicin (Hydrochloride)	13_2F	Ammonium glycyrrhizinate	20_9B	Altrenogest
6_10F	Isocarboxazid	13_4D	Nintedanib	20_11G	Methylthiouracil

727 **Supplemental Table 1: MedChemExpress HY-L066 Library acquired thrombosis model hits**
728 **(n=117)**

Compounds	Company - Catalog No	Compounds	Company - Catalog No
AM-251	Cayman Chemical - 71670	1-850	Cayman Chemical - 17248
Meclofenoxate (hydrochloride)	Cayman Chemical - 28283	Pantoprazole sodium	Sigma - PHR1604
Loxapine succinate	Sigma - L106	Dexlansoprazole	Sigma - SML2624
Lofepramine hydrochloride	Sigma - BP697	Omeprazole	Sigma - PHR1059
Etomidate	Selleckchem - S1329	Esomeprazole magnesium	Sigma - PHR1585
Laquinimod	Sigma - SML2504	SCH28080	MedChemExpress - HY-103261
Bendazac	Selleckchem - S5361	Ilaprazole	Sigma - SML2322
Diphenylcyclopropenone	Sigma - 177377	Elbasvir	Selleckchem - S5652
Belinostat	MedChemExpress - HY-10225	Benvitimod	Sigma - SML3430
Tivozanib	Sigma - SML3132	Indole-3-carboxylic acid	Sigma - 284734
Nintedanib	Sigma - SML2848	Phenindione	Selleckchem - S1921
L-Thyroxine	Sigma - T1775	Apixaban	Sigma - SML3285
Liothyronine	MedChemExpress - HY-113082	Warfarin	Sigma – A2250
Mifepristone M8046	Sigma - M8046	Dabigatran etexilate	Sigma - SML2351
Exemestane	Sigma - PHR1634	Rivaroxaban	Sigma - SML2844
Mestranol*	Sigma - 855871		

729   **Supplemental Table 2: Validated hit compounds**

thr paralog	Exon	sgRNA sequence
thraa	4	AGAUAACUGGGAACGUACCC
thrab	5	CAUCCUUCUCCAGAUAGCUG
thrb	4	UCAUCCUUGUCCAGGUAACU

thr paralog	Forward primer	Reverse primer	Amplicon size (base pairs)
thraa Exon 4	CATGTGATGCAGCGGTAATGA	TGCAGTCGCCTATTCTCTCCT	140
thrab Exon 5	TGTGCACAAGCGATAGCAGA	GCATGGCAGGGTACATCC	140
thrb Exon 4	CTCACATGTAATGCAGCGATAGTG	ATGACGAGGTCTGTTGTCTTGG	359

730 **Supplemental Table 3: Sequences of sgRNAs and PCR primers**

Compounds	IC$_{30}$		IC$_{50}$	
	Acquired	Genetic	Acquired	Genetic
AM-251	6.72	12.87	No effect	14.02
Meclofenoxate	6.03	No effect	10.69	No effect
Loxapine	19.4[#]	No effect[#]	No effect	No effect
Lofepramine	6.04	No effect	No effect	No effect
Etomidate	23.08	18.94	26.91	No effect
Laquinimod	1.43	10.01	2.89	No effect
Bendazac	1.51	No effect	9.36	No effect
Tiaprofenic acid	2.63	No effect	5.13	No effect
Buparvaquone	No effect	0.69	No effect	1.08
Edaravone	No effect	9.12	No effect	13.1
Malotilate	No effect	24.05	No effect	59.99
Belinostat	47.45	No effect	No effect	No effect
Diphenylcyclopropenone	7.89	No effect	No effect	No effect
Tivozanib*	0.67	1.68	3.53	No effect
Nintedanib*	6.93	4.06	9.64	No effect
L-Thyroxine	0.2	0.15	0.83	0.32
Liothyronine	0.28	0.34	0.36	0.8
Mifepristone*	7.44	10.10	11.32	No effect
Exemestane	2.95	13.72	5.71	17.80
Mestranol*	8.47[#]	11.91[#]	10.19	15.47
1-850	6.07	No effect	No effect	No effect
Pantoprazole	14.27	No effect	27.73	No effect
Dexlansoprazole	10.69	5.19	13.77	18.80
Omeprazole	14.57	No effect	21.35	No effect
Esomeprazole*	8.38	5.16	10.62	No effect
SCH28080	6.11	No effect	6.62	No effect
Ilaprazole	14.79	No effect	73.45	No effect
Elbasvir	1.35	No effect	1.71	No effect
Thiabendazole	No effect	14.34	No effect	No effect
Nalidixic acid	No effect	6.34	No effect	No effect
Indole-3-carboxylic acid	7.89	No effect	10.48	No effect
Benvitimod	2.82	No effect	4.66	No effect
Phenindione	42.82	9.64	66.04	19.93
Apixaban	No effect	No effect	No effect	No effect
Warfarin	No effect	0.02	No effect	0.05
Dabigatran etexilate	No effect	4.58	No effect	11.72
Rivaroxaban	No effect	No effect	No effect	No effect

Supplemental Table 4: IC$_{30}$ and IC$_{50}$ values for the candidate compounds tested on acquired (estrogen-induced) and genetic (spontaneous/PC deficient) models of thrombosis. The values (μM) were obtained from dose response curves. No effect, no IC$_{30}$/IC$_{50}$ detected from the dose response curves. Hits that are not concordant between the models are highlighted in gray. #, IC$_{30}$/IC$_{50}$ values are derived from the procoagulant profiles of the respective drugs in the acquired model with no mestranol treatment. *, compounds with procoagulant effects in the genetic model.

Drug	Estimate	Std.error	Statistic	Adj.p.value
DMSO control	0.16	0.11	1.49	0.14
Apixaban (25 µM)	1.26	0.29	4.34	1.40E-05
Dabigatran etexilate (25 µM)	1.04	0.34	3.08	0.002
Warfarin (25 µM)	0.16	0.11	1.49	0.14
Exemestane (10 µM)	0.39	0.26	1.50	0.13
Liothyronine (0.4 µM)	0.68	0.30	2.29	0.02
L-thyroxine (0.6 µM)	0.19	0.20	0.94	0.34
Tivozanib (3 µM)	0.69	0.27	2.51	0.016
Esomeprazole (8 µM)	0.40	0.31	1.30	0.19
Nintedanib (6 µM)	0.66	0.21	3.12	0.0018
Dexlansoprazole (8 µM)	0.18	0.20	0.89	0.37
Mifepristone (6 µM)	0.49	0.24	2.02	0.04
Etomidate (10 µM)	0.81	0.24	3.32	0.0009
Laquinimod (5 µM)	0.54	0.19	2.83	0.0047
AM-251(5 µM)	0.72	0.27	2.72	0.006

738 **Supplemental Table 5: Time to occlusion batch corrected statistics comparing the effect**
739 **of candidate drugs versus their mestranol combinations in Figure 3.** These data show the
740 impact of combining the drugs with mestranol on time to occlusion.

Drug	Drug alone vs DMSO control			
	Estimate	Std.error	Statistic	Adj.p.value
Apixaban (25 µM)	-1.21	0.21	-5.79	9.14E-08
Dabigatran etexilate (25 µM)	-1.54	0.24	-6.52	9.24E-10
Warfarin (25 µM)	0	0	inf	inf
Exemestane (10 µM)	-0.33	0.22	-1.51	0.64
Liothyronine (0.4 µM)	-0.36	0.23	-1.54	0.62
L-thyroxine (0.6 µM)	-0.70	0.16	-4.39	0.00014
Tivozanib (3 µM)	0.14	0.22	0.61	0.99
Esomeprazole (8 µM)	0.57	0.24	2.27	0.19
Nintedanib (6 µM)	-0.03	0.19	-0.15	0.99
Dexlansoprazole (8 µM)	-0.09	0.17	-0.54	0.99
Mifepristone (6 µM)	-0.06	0.22	-0.29	0.99
Etomidate (10 µM)	0.50	0.21	2.42	0.14
Laquinimod (5 µM)	0.13	0.17	0.74	0.97
AM-251(5 µM)	1.10	0.21	5.15	3.33E-06

Drug	Drug + Mestranol vs Mestranol control			
Apixaban (25 µM)	-2.30	0.26	-8.80	4.03E-14
Dabigatran etexilate (25 µM)	-2.43	0.29	-8.49	4.95E-14
Warfarin (25 µM)	0	0	inf	inf
Exemestane (10 µM)	-0.54	0.22	-2.49	0.12
Liothyronine (0.4 µM)	-0.87	0.25	-3.49	0.006
L-thyroxine (0.6 µM)	-0.73	0.18	-3.96	0.0009
Tivozanib (3 µM)	-0.39	0.22	-1.74	0.49
Esomeprazole (8 µM)	0.33	0.25	1.32	0.75
Nintedanib (6 µM)	-0.52	0.17	-3.03	0.026
Dexlansoprazole (8 µM)	-0.10	0.18	-0.58	0.99
Mifepristone (6 µM)	-0.39	0.21	-1.90	0.39
Etomidate (10 µM)	-0.14	0.21	-0.71	0.98
Laquinimod (5 µM)	-0.25	0.17	-1.49	0.65
AM-251(5 µM)	0.55	0.22	2.46	0.13

741 *inf: infinitesimal*

742 **Supplemental Table 6: Time to occlusion batch corrected statistics comparing the effect**
743 **of drugs versus the DMSO and mestranol control groups, respectively, in Figure 3.**These
744 data show the impact of the drugs and mestranol combinations against the reference groups
745 including DMSO and mestranol control**.**

University of Michigan Data			
	HC alone	HC with Thyroid Medication	Thyroid Med Alone
Baseline Characteristics			
Total	47,710	2,125	30,731
Age	25.1 (25.0-25.2)	32.6 (32.2-33.0)	37.7 (37.6-37.8)
BMI	26.0 (25.9-26.1)	25.5 (25.0-25.9)	30.2 (30.1-30.3)
Smoking Status			
Never	79.2%	79.3%	52.0%
Former	8.2%	14.2%	15.0%
Current	5.2%	4.0%	9.1%
Unspecified	7.4%	2.5%	23.2%
Medication			
Thyroid			
Levothyroxine		1850 (87.1%)	27,981 (91.1%)
Reconstituted Thyroid		164 (7.7%)	1,720 (5.6%)
Liothyronine		111 (5.2%)	1,028 (3.4%)
Duration (days)			
		557 (529-586)	847 (834 - 860)
Levothyroxine-eq Dose (mg)			
		0.090 (0.088-0.092)	0.097 (0.096-0.98)
Hormonal contraception			
Estrogen			
Estrogen dose (mg)	0.030 (0.029-0.030)	0.037 (0.031-0.044)	
Estrogen only	18 (<1%)	19 (<1%)	
Progesterone (%)			
1st generation	13984 (29.3%)	762 (36.2%)	
2nd generation	8060 (16.9%)	417 (19.8%)	
3rd generation	21237 (44.5%)	752 (35.7%)	
4th generation	4,409 (9.2%)	173 (8.2%)	
Route (%)			
PO	43979 (92.2%)	1990 (93.7%)	
Vaginal Ring	2,834 (5.94%)	99 (4.6%)	
Patch	895 (1.9%)	34 (1.6%)	
Duration (days)			
	674 (668-681)	557 (528-585)	
Diagnosis			
DVT	131 (41.1%)	8 (32%)	149 (57.8%)
Pulmonary Embolism	79 (24.8%)	12 (48%)	59 (22.9%)
Thrombophlebitis	64 (20.1%)	3 (12%)	17 (6.6%)
Stroke	26 (8.2%)	2 (8%)	6 (2.3%)
Abdominal Thrombosis	12 (3.8%)	-	18 (7.0%)
Arterial Thrombosis	7 (2.2%)	-	9 (3.5%)

747   **Supplemental Table 7: Patient, medication, and diagnosis characteristics.**

REFERENCES

1. Grosse SD, Nelson RE, Nyarko KA, Richardson LC, Raskob GE. The economic burden of incident venous thromboembolism in the United States: a review of estimated attributable healthcare costs. *Thrombosis research* 2016:137:3-10.DOI: 10.1016/j.thromres.2015.11.033

2. Prevention USCfDCa. 2025 August 24. Data and Statistics on Venous Thromboembolism. <https://www.cdc.gov/blood-clots/data-research/facts-stats/index.html>. Accessed 2025 August 24.

3. Gupta A, Patibandla S. Protein C Deficiency. StatPearls. Treasure Island (FL): StatPearls Publishing Copyright © 2024, StatPearls Publishing LLC.; 2024.

4. Danese S, Vetrano S, Zhang L, Poplis VA, Castellino FJ. The protein C pathway in tissue inflammation and injury: pathogenic role and therapeutic implications. *Blood* 2010:115(6):1121-1130.DOI: 10.1182/blood-2009-09-201616

5. LaVasseur C, Neukam S, Kartika T, Samuelson Bannow B, Shatzel J, DeLoughery TG. Hormonal therapies and venous thrombosis: Considerations for prevention and management. *Res Pract Thromb Haemost* 2022:6(6):e12763.DOI: 10.1002/rth2.12763

6. Lutsey PL, Zakai NA. Epidemiology and prevention of venous thromboembolism. *Nat Rev Cardiol* 2023:20(4):248-262.DOI: 10.1038/s41569-022-00787-6

7. Bistervels IM, Scheres LJJ, Hamulyák EN, Middeldorp S. Sex matters: Practice 5P's when treating young women with venous thromboembolism. *J Thromb Haemost* 2019:17(9):1417-1429.DOI: 10.1111/jth.14549

8. Klok FA, Huisman MV. How I assess and manage the risk of bleeding in patients treated for venous thromboembolism. *Blood* 2020:135(10):724-734.DOI: 10.1182/blood.2019001605

9. Wallvik N, Renlund H, Själander A. Bleeding risk in patients with venous thromboembolic events treated with new oral anticoagulants. *J Thromb Thrombolysis* 2021:52(1):315-323.DOI: 10.1007/s11239-020-02319-w

10. Lucà F, Oliva F, Abrignani MG, Di Fusco SA, Parrini I, Canale ML, Giubilato S, Cornara S, Nesti M, Rao CM, Pozzi A, Binaghi G, Maloberti A, Ceravolo R, Bisceglia I, Rossini R, Temporelli PL, Amico AF, Calvanese R, Gelsomino S, Riccio C, Grimaldi M, Colivicchi F, Gulizia MM. Management of Patients Treated with Direct Oral Anticoagulants in Clinical Practice and Challenging Scenarios. *J Clin Med* 2023:12(18).DOI: 10.3390/jcm12185955

11. Geller AI, Shehab N, Lovegrove MC, Rose KO, Weidle NJ, Goring SK, Budnitz DS. Emergency visits for oral anticoagulant bleeding. *Journal of General Internal Medicine* 2020:35:371-373.DOI: 10.1007/s11606-019-05391-y

12. Zaidi SRH, Jenkins SM. Anticoagulant Therapy In Pregnancy. StatPearls. Treasure Island (FL): StatPearls Publishing Copyright © 2024, StatPearls Publishing LLC.; 2024.

13. Weyand AC, Shavit JA. Zebrafish as a model system for the study of hemostasis and thrombosis. *Curr Opin Hematol* 2014:21(5):418-422.DOI: 10.1097/moh.0000000000000075

14. Raghunath A, Ferguson AC, Shavit JA. Fishing for answers to hemostatic and thrombotic disease: Genome editing in zebrafish. *Res Pract Thromb Haemost* 2022:6(5):e12759.DOI: 10.1002/rth2.12759

15. Williams CH, Hong CC. Zebrafish small molecule screens: Taking the phenotypic plunge. *Comput Struct Biotechnol J* 2016:14:350-356.DOI: 10.1016/j.csbj.2016.09.001

16. Kretz CA, Weyand AC, Shavit JA. Modeling Disorders of Blood Coagulation in the Zebrafish. *Curr Pathobiol Rep* 2015:3(2):155-161.DOI: 10.1007/s40139-015-0081-3

17. Santoriello C, Zon LI. Hooked! Modeling human disease in zebrafish. *J Clin Invest* 2012:122(7):2337-2343.DOI: 10.1172/jci60434

18. Ayyoub S, Orriols R, Oliver E, Ceide OT. Thrombosis Models: An Overview of Common In Vivo and In Vitro Models of Thrombosis. *Int J Mol Sci* 2023:24(3).DOI: 10.3390/ijms24032569

19. Kithcart A, MacRae CA. Using Zebrafish for High-Throughput Screening of Novel Cardiovascular Drugs. *JACC Basic Transl Sci* 2017:2(1):1-12.DOI: 10.1016/j.jacbts.2017.01.004

20. North TE, Goessling W, Walkley CR, Lengerke C, Kopani KR, Lord AM, Weber GJ, Bowman TV, Jang IH, Grosser T, Fitzgerald GA, Daley GQ, Orkin SH, Zon LI. Prostaglandin E2 regulates vertebrate haematopoietic stem cell homeostasis. *Nature* 2007:447(7147):1007-1011.DOI: 10.1038/nature05883

21. Cutler C, Multani P, Robbins D, Kim HT, Le T, Hoggatt J, Pelus LM, Desponts C, Chen YB, Rezner B, Armand P, Koreth J, Glotzbecker B, Ho VT, Alyea E, Isom M, Kao G, Armant M, Silberstein L, Hu P, Soiffer RJ, Scadden DT, Ritz J, Goessling W, North TE, Mendlein J, Ballen K, Zon LI, Antin JH, Shoemaker DD. Prostaglandin-modulated umbilical cord blood hematopoietic stem cell transplantation. *Blood* 2013:122(17):3074-3081.DOI: 10.1182/blood-2013-05-503177

22. Liu Y, Kretz CA, Maeder ML, Richter CE, Tsao P, Vo AH, Huarng MC, Rode T, Hu Z, Mehra R, Olson ST, Joung JK, Shavit JA. Targeted mutagenesis of zebrafish antithrombin III triggers disseminated intravascular coagulation and thrombosis, revealing insight into function. *Blood* 2014:124(1):142-150.DOI: 10.1182/blood-2014-03-561027

23. Ku CJ, Yu X, Zhao QY, Grzegorski SJ, Daniel JG, Ferguson AC, Shavit JA. Loss of protein C vs protein S results in discrepant thrombotic phenotypes. *Blood Adv* 2025:9(3):545-557.DOI: 10.1182/bloodadvances.2024013237

24. Yu X, Zhao QY, Yaman M, Emly SM, Lee JK, Su H, Ferguson AC, Nagaswami C, Chaturantabut S, Goessling W, Weisel JW, Auchus RJ, Shavit JA. Hormone-induced thrombosis is mediated through non-canonical fibrin (ogen) aggregation and a novel estrogen target in zebrafish. *bioRxiv* 2024:10.1101/2024.11.13.623199.DOI: 10.1101/2024.11.13.623199

25. Antonijevic N, Matic D, Beleslin B, Mikovic D, Lekovic Z, Marjanovic M, Uscumlic A, Birovljev L, Jakovljevic B. The Influence of Hyperthyroidism on the Coagulation and on the Risk of Thrombosis. *J Clin Med* 2024:13(6).DOI: 10.3390/jcm13061756

26. Wang Y, Ding C, Guo C, Wang J, Liu S. Association between thyroid dysfunction and venous thromboembolism: A systematic review and meta-analysis. *Medicine (Baltimore)* 2023:102(11):e33301.DOI: 10.1097/md.0000000000033301

27. Zhang L, Li K, Yang Q, Lin Y, Geng C, Huang W, Zeng W. Associations between deep venous thrombosis and thyroid diseases: a two-sample bidirectional Mendelian randomization study. *Eur J Med Res* 2024:29(1):327.DOI: 10.1186/s40001-024-01933-1

28. Hieber M, von Kageneck C, Weiller C, Lambeck J. Thyroid Diseases Are an Underestimated Risk Factor for Cerebral Venous Sinus Thrombosis. *Front Neurol* 2020:11:561656.DOI: 10.3389/fneur.2020.561656

29. Raps M, Curvers J, Helmerhorst FM, Ballieux BE, Rosing J, Thomassen S, Rosendaal FR, van Vliet HA. Thyroid function, activated protein C resistance and the risk of venous thrombosis in users of hormonal contraceptives. *Thromb Res* 2014:133(4):640-644.DOI: 10.1016/j.thromres.2013.12.041

30. Vo AH, Swaroop A, Liu Y, Norris ZG, Shavit JA. Loss of fibrinogen in zebrafish results in symptoms consistent with human hypofibrinogenemia. *PLoS One* 2013:8(9):e74682.DOI: 10.1371/journal.pone.0074682

31. Ku C-J, Grzegorski S, Shavit JA. Loss of Protein C in Zebrafish Results in Impaired Neutrophil Migration after Tissue Injury. *Blood* 2020:136:22.DOI: 10.1182/blood-2020-142541

32. Zhang JH, Chung TD, Oldenburg KR. A Simple Statistical Parameter for Use in
Evaluation and Validation of High Throughput Screening Assays. *J Biomol Screen*
1999:4(2):67-73.DOI: 10.1177/108705719900400206
33. McCloy RA, Rogers S, Caldon CE, Lorca T, Castro A, Burgess A. Partial inhibition of
Cdk1 in G 2 phase overrides the SAC and decouples mitotic events. *Cell Cycle*
2014:13(9):1400-1412.DOI: 10.4161/cc.28401
34. Rost MS, Grzegorski SJ, Shavit JA. Quantitative methods for studying hemostasis in
zebrafish larvae. *Methods Cell Biol* 2016:134:377-389.DOI:
10.1016/bs.mcb.2016.01.003
35. Jonklaas J, Bianco AC, Bauer AJ, Burman KD, Cappola AR, Celi FS, Cooper DS, Kim
BW, Peeters RP, Rosenthal MS, Sawka AM. Guidelines for the treatment of
hypothyroidism: prepared by the american thyroid association task force on thyroid
hormone replacement. *Thyroid* 2014:24(12):1670-1751.DOI: 10.1089/thy.2014.0028
36. Richter CE, Raghunath A, Griffin MS, Yaman M, Arruda VR, Samelson-Jones BJ, Shavit
JA. Loss of factor VIII in zebrafish rebalances antithrombin deficiency but has a limited
bleeding diathesis. *bioRxiv* 2024:10.1101/2024.02.28.582609.DOI:
10.1101/2024.02.28.582609
37. Venables W, Ripley B. Modern Applied Statistics with S Fourth Edition, 2002.
38. Therneau T. A package for survival analysis in S. 2015.
39. Therneau T, Grambsch P. Modeling survival data: extending the Cox model Springer-
Verlag. 2000.
40. Lenth R. Emmeans: Estimated Marginal Means, Aka Least-Squares Means, R Package
Version 1.10. 4; 2024.
41. Fiori JL, Sanghvi M, O'Connell MP, Krzysik-Walker SM, Moaddel R, Bernier M. The
cannabinoid receptor inverse agonist AM251 regulates the expression of the EGF
receptor and its ligands via destabilization of oestrogen-related receptor α protein. *Br J*
Pharmacol 2011:164(3):1026-1040.DOI: 10.1111/j.1476-5381.2011.01384.x
42. Valk BI, Struys M. Etomidate and its Analogs: A Review of Pharmacokinetics and
Pharmacodynamics. *Clin Pharmacokinet* 2021:60(10):1253-1269.DOI: 10.1007/s40262-
021-01038-6
43. Popovic D, Nuss P, Vieta E. Revisiting loxapine: a systematic review. *Ann Gen*
Psychiatry 2015:14:15.DOI: 10.1186/s12991-015-0053-3
44. Yu X, Weyand AC, Shavit JA. Surprisingly Rapid Onset of Estrogen-Induced Venous
Thrombosis in an Animal Model. *Blood* 2018:132:418.DOI: 10.1182/blood-2018-99-
120325
45. Grzegorski SJ, Hu Z, Liu Y, Yu X, Ferguson AC, Madarati H, Friedmann AP, Reyon D,
Kim PY, Kretz CA, Joung JK, Shavit JA. Disruption of the kringle 1 domain of
prothrombin leads to late onset mortality in zebrafish. *Sci Rep* 2020:10(1):4049.DOI:
10.1038/s41598-020-60840-7
46. Vigano D'Angelo S, Comp PC, Esmon CT, D'Angelo A. Relationship between protein C
antigen and anticoagulant activity during oral anticoagulation and in selected disease
states. *J Clin Invest* 1986:77(2):416-425.DOI: 10.1172/jci112319
47. Kendall MJ. Review article: esomeprazole--the first proton pump inhibitor to be
developed as an isomer. *Aliment Pharmacol Ther* 2003:17 Suppl 1:1-4.DOI:
10.1046/j.1365-2036.17.s1.1.x
48. Upchurch HF, Conway E, Patterson JR. MK, Maxwell MD. Localization of cellular
transglutaminase on the extracellular matrix after wounding: Characteristics of the matrix
bound enzyme. *Journal of Cellular Physiology* 1991:149(3):375-382.DOI:
10.1002/jcp.1041490304
49. Haroon ZA, Hettasch JM, Lai T-S, Dewhirst MW, Greenberg CS. Tissue
transglutaminase is expressed, active, and directly involved in rat dermal wound healing

and angiogenesis. *The FASEB Journal* 1999:13(13):1787-1795.DOI:
10.1096/fasebj.13.13.1787

50. Wang Z, Perez M, Lee E-S, Kojima S, Griffin M. The functional relationship between
transglutaminase 2 and transforming growth factor β1 in the regulation of angiogenesis
and endothelial–mesenchymal transition. *Cell Death & Disease* 2017:8(9):e3032-
e3032.DOI: 10.1038/cddis.2017.399

51. Lisetto M, Fattorini M, Lanza A, Gerdol M, Griffin M, Wang Z, Ferrara F, Sblattero D.
Biochemical and Functional Characterization of the Three Zebrafish Transglutaminases
2. *Int J Mol Sci* 2023:24(15).DOI: 10.3390/ijms241512041

52. Murthy SN, Lorand L. Cross-linked A alpha.gamma chain hybrids serve as unique
markers for fibrinogen polymerized by tissue transglutaminase. *Proc Natl Acad Sci U S*
A 1990:87(24):9679-9682.DOI: 10.1073/pnas.87.24.9679

53. Wei Z, Boateng NKK, Schmitt LR, Cline H, Fernandes Fonseca MT, Newberry A, Taylor
A, Adelmeijer J, Poole LG, Stravitz RT, Lee WM, Lisman T, Hansen KC, Luyendyk JP.
Integrated cross-linking by TG2 and FXIII generates hepatoprotective fibrin(ogen)
deposits in injured liver. *Blood* 2025:145(21):2507-2517.DOI:
10.1182/blood.2024026938

54. Ren B, Zhu Y. A New Perspective on Thyroid Hormones: Crosstalk with Reproductive
Hormones in Females. *Int J Mol Sci* 2022:23(5).DOI: 10.3390/ijms23052708

55. Santin AP, Furlanetto TW. Role of estrogen in thyroid function and growth regulation. *J*
Thyroid Res 2011:2011:875125.DOI: 10.4061/2011/875125

56. Mulder JE. Thyroid disease in women. *Med Clin North Am* 1998:82(1):103-125.DOI:
10.1016/s0025-7125(05)70596-4

57. Vasudevan N, Ogawa S, Pfaff D. Estrogen and thyroid hormone receptor interactions:
physiological flexibility by molecular specificity. *Physiol Rev* 2002:82(4):923-944.DOI:
10.1152/physrev.00014.2002

58. Zsarnovszky A, Kiss D, Jocsak G, Nemeth G, Toth I, Horvath TL. Thyroid hormone- and
estrogen receptor interactions with natural ligands and endocrine disruptors in the
cerebellum. *Front Neuroendocrinol* 2018:48:23-36.DOI: 10.1016/j.yfrne.2017.10.001

929