Complex modulation of visual attention to 3[rd] person interactions in wild macaques

Sofia M. Pereira[1,2], Suthirote Meesawat[3,4], Suchinda Malaivijitnond[3,4], Julia Ostner[1,2,5] & Oliver Schülke[1,2,5]

[1]Department of Behavioral Ecology, University of Göttingen, Göttingen, Germany

[2]Social Evolution in Primates Group, German Primate Center Leibniz Institute for Primate Research, Göttingen, Germany

[3]Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

[4]National Primate Research Center of Thailand, Chulalongkorn University, Saraburi, Thailand

[5]Leibniz ScienceCampus Primate Cognition, German Primate Center, Leibniz Institute for Primate Research, Göttingen, Germany

ORCID IDs
Sofia M. Pereira: 0000-0002-8352-4664
Suchinda Malaivijitnond: 0000-0003-0897-2632
Suthirote Meesawat: 0000-0001-8612-2436
Oliver Schülke: 0000-0003-0028-9425
Julia Ostner: 0000-0001-6871-9976

Abstract

Social animals rely on socio-cognitive skills to monitor their social environment and make informed decisions. Yet, visual attention to others' interactions in real-life scenes remains understudied, despite evidence for dedicated neuronal networks for the processing of real-life social information and for social interactions in particular. Here, we ask how subject characteristics, interaction valence, and social relationships between subject and stimuli interact to guide overt attention of 56 wild male Assamese macaques to 962 3[rd] person social scenes that unfolded spontaneously in their vicinity. Social interactions drew more and longer attention than social control scenes lacking the interaction, highlighting the relevance of interaction. The effect of both social bond strength and dominance relations between subject and stimulus on the probability to attend was independently

modulated by scene valence. Moreover, beyond reacting to the affordances of the social scene and

independent of their social status, males differed in how long they observed the interactions of

others. These findings suggest that visual attention is guided by the value an individual associates

with different aspects of the social scene based on previous experience in both the agonistic and the

affiliative realms and by current threat indicative of complex information integration in the brain.

Keywords: social cognition, information integration, valence, social status, social bonds

INTRODUCTION

Social neuroscience research employs a variety of social stimuli that vary in their naturalism[1]. On one

end of the spectrum the subject is chair-seated and presented with static images of social agents

and non-social controls, while their eye gaze and neuronal activity is tracked[2-4]. From there,

naturalism is increased by using videos that are more salient, but also noisier, stimuli[5,6]. Virtual

reality scenes can even more closely mimic real-life interactions, as avatars can display contingent

behavior creating a live-stream of interactions[7,8]. However, real-life agents are the most naturalistic,

yet also the most challenging to control[9,10] particularly in natural environments[1]. This study leverages

the strengths of naturalistic animal observations, where large numbers of subjects can be studied

allowing us to weigh individual differences in social attention against the affordances of the stimulus

situation including subject's affiliative and dominance relations with the stimuli.

Social attention, i.e. monitoring conspecifics to extract information, is a crucial building block

of advanced social cognition[11], involving specialized brain networks[12,13]. Social primates are willing to

forgo an immediate food reward for the privilege to watch conspecifics in videos[14]. Yet, not all social

information is equally relevant. For example, rhesus monkeys (*Macaca mulatta*) pay with forgone

juice rewards for viewing a picture of the face of a dominant male, but have to be paid to look at a

subordinate[15]. Visual attention for faces provides information about others' overt spatial attention,

i.e. about gaze direction[16], and social signals encoded in ritualized facial expressions[2]. In fact,

macaques pay more attention to facial expressions associated with threat than to those associated

with benign intent and even less to neutral faces [17,18] , and more readily follow the gaze of others if

they show an emotional as opposed to a neutral expression [19]. Single neurons in the amygdala show

selective activity for threat, friendly and neutral facial expressions[2].

Thus, social attention targets faces, gazes, and facial expressions as well as relevancies

unrelated to perceptual aspects of the stimulus, e.g. stimulus dominance status, all of which can be

tested with static information. Testing with video recorded dynamic stimuli shows that sustained
attention depends on additional ethogram elements contained in the video sequence[20,21]: Rhesus
macaques looked longer at areas of a frame with a grooming solicitation, aggression or submission[20],
and chimpanzees (*Pan troglodytes*) pay more attention to interacting dyads than solitary controls[21].
During fMRI, macaques engage a prefrontal cortical area specifically when watching videos of two
others interacting compared to two others acting each on their own or simply resting; there is a
brain for interaction[6].

Presenting videos of longer natural scenes with multiple real-life agents hugely advances our
understanding of selective social attention in complex environments and the associated
neurophysiology[1,13]. However, dual eye-tracking and electro-physiology demonstrated that direct
interaction with real-life agents draws even more attention and involves different networks across
prefrontal cortical and amygdala areas than watching stills or even videos of the very same
agents[10,22]. Neurons in the anterior cingulate cortex predict the behavior of others during social
interaction in an iterated prisoner's dilemma task and disruption selectively hampers mutually
beneficial interactions in rhesus macaques[9]. A network in the dorsomedial prefrontal cortex codes
for social partner identity and tracks these partners' behavior as well as the distribution of food
rewards in a real-life social task[23]. Thus, studies employing real-life interactions promise to shed new
light on the processes involved in social information processing particularly if agents are freely
moving[24].

Social attention is dictated by the affordances of the stimulus situation, but can also vary
considerably between individuals. Social status within a group can be a strong predictor of how
much attention an individual pays their group mates[25–27]. In a population with clearly defined and
repeatable inter-individual differences in sociability, low-sociable rhesus macaques paid more
attention to any social information in videos than high-sociable monkeys independent of social
status[28]. Early life adversity, such as experimental maternal immune activation during gestation, can
also affect social attention in juvenile offspring possibly via epigenetic effects[29]. Both the genotype
upstream of the serotonin transporter gene[30,31] and administered serotonin precursor[32] affect social
attention, suggesting the involvement of a serotonergic pathway in generating the observed inter-
individual differences[33]. Additionally, an oxytocinergic pathway has been implicated in linking
individual differences in affiliation and dominance with visual attention[34].

This study focuses on how individuals selectively attend to 3rd person situations, i.e.
interactions among others[13], as they spontaneously unfold (or not) in the vicinity, excluding
interactions involving themselves. We propose that the relevance of agonistic and affiliative 3rd

person interactions differs considerably, and that relational aspects add an additional layer of

relevance[35] . For example, when conflicts arise, they can quickly escalate and involve the subject

directly, either through redirected aggression[36] or by soliciting their help[37]. As a result, social

attention for aggressive interactions may serve a risk avoidance function and is likely modulated by

the dominance relationship between the subject and the interacting parties. In contrast, the

relevance of affiliation between others is often less immediate and typically indirect. Rates of

grooming and agonistic support are often correlated[37] which means that the state of the grooming

network provides indirect information about the alliance network in the group[38]. Information about

3rd person grooming will affect the subject only indirectly and usually not immediately, but when it

needs to decide in a later situation whom to support or which affiliative relationships to invest in.

Yet sometimes, primates also directly and immediately interfere in the grooming interactions of

others to gain access to desirable partners or to manipulate others' relationships[39,40].

In a population of wild Assamese macaques in Thailand, we used behavioral event recording

to quantify the overt attention of 56 adult males to 962 agonistic and affiliative 3rd person

interaction unfolding in subject's vicinity (≤5m) or controls where two other adult males came into

close proximity of ≤1.5m without interacting by means of vocalization, facial expression, touch or

pose gesture. With statistical models we show that valence of the social scene (agonism, affiliation,

control) interacted with dominance relations and social bond strength between the subject and the

two stimulus monkeys to predict the occurrence of overt attention to the scene suggesting non-

additive integration of different relevancies during processing of 3rd person real-life interactions. We

further identify individual differences in sustained attention for agonistic or affiliative interactions

that are independent of subject rank. Note that we refer to the affiliative relationship between two

males as their social bond strength to enhance accessibility of the results; strictly speaking, social

bonds are only the strong, equitable, stable (and supportive) among all affiliative relationships[41].

METHODS

Study site and subjects

This study was conducted at the Phu Khieo Wildlife Sanctuary in North-Eastern Thailand (PKWS, 16°

05'-35' N, 101° 20'-55' E, 1573km^2) as part of a long-term research project on a wild population of

Assamese macaques ongoing since 2005[42]. PKWS is part of the well-preserved Western Isaan Forest

Complex covering an area of more than 6,500km^2 [43], which houses a diverse community of large

mammals and predators, an indicator of low disturbance levels[43]. We collected data on 56 adult

Assamese macaque males belonging to five different neighboring multimale–multifemale groups
with 39 to 75 individuals each and a minimum of eight adult males and ten adult females.

Social attention data collection

Social attention data were collected from June 2022 until April 2023 through event sampling of
behavior towards social scenes spontaneously unfolding in the group. A social scene was considered
valid for recording each time a stationary adult male subject in the vicinity (≤ 5 m) had an
unobstructed view and the human observer clearly saw the onset of the social scene. We recorded
head gaze (but also eye gaze if possible) towards three different social scene *conditions* that all
started with two adult males (the stimulus individuals) arriving within 1.5 m of proximity of one
another and continued either (1) as non-interacting *controls* if the males remained in close proximity
without explicitly interacting via touch, vocalization or facial expression or (2) as *affiliation* if the two
males only exchanged friendly behaviors, and (3) as *agonism* if the interaction included aggressive or
submissive behaviors. Given the specificity of parameters and viewing conditions necessary to
accurately record each attention event, these data were collected opportunistically outside the focal
animal follows. Upon the unambiguous onset of a social attention event, we recorded the following
variables: identities of the subject and two interacting stimulus individuals, behaviors displayed
during the stimulus interactions, start and end times of stimulus interactions, timing of looks relative
to the onset of the stimulus interaction and duration of each look. We additionally scored the
interaction for conspicuousness (0/1) as a means to control statistically for loud or tumultuous
interactions drawing attention[35]. We collected data from 962 social attention events, i.e. 17±9
events for each of the 56 subjects.

Social bond strength predictor

$$DDSI_{sub-max} = \max(DDSI_{sub-stim1}, \; DDSI_{sub-stim2})$$

Dominance relation predictor

$$\Delta R_{sub-max} = R_{sub} - \max(R_{stim1}, R_{stim2})$$

Control

Affiliation

Agonism

Subject

Figure 1: This field study integrates data on visual attention to social scenes spontaneously unfolding
in the vicinity of a subject (sub) with information about the affiliative and dominance relationships
among all males in five study groups. Social scenes with two adult males in close proximity were
classified as non-interacting controls or interactions reflecting affiliation or agonism. The relationship
of the subject to the two stimulus males (stim 1, stim 2) in the scene was quantified as the
dominance relation to the higher-ranking of the stimulus males (Rank difference sub-max) and the
maximum social bond strength to either of the two (DDSI$_{sub-max}$).

Collection of social relationship data

Concomitant with the social attention data collection, all adult males were systematically observed

as focal animals in 40 min protocols[44] from June 2022 through April 2023. We recorded continuously

[45] all affiliative and agonistic (both aggressive and submissive) interactions including the identities of

the actor and receiver as well as all approaches into and departures from a 1.5 m sphere around the

focal animal. To inform our dominance and social bond metrics (see below) about the state of

relationships at the onset of the study in June 2022, we included data from January-May 2022. A

total of 1,915.9h of continuous focal animal data were collected (mean ± standard deviation per

male 33.6h ± 12.1).

Dominance scoring – Elo-rating

For the assessment of dominance, we scored submissive behaviors (i.e. bare teeth, make room, give
ground) in decided dyadic conflicts (either unprovoked submission or submission following
aggression) registered during our continuous focal animal observations and ad libitum recordings
from January 2022 through April 2023. To avoid long burn-in periods when males immigrate or
mature, we employed Bayesian Elo-rating[46] that estimates start values from the data and captures
the state and dynamics of male dominance relationships, which can fluctuate remarkably in
Assamese macaques[47]. We used the elo_seq_bayes and extract_elo_b functions of the
EloRating.Bayes package[48] in R version 4.3.0[49]. A short burn-in period of one month was applied for
each male immigrating into a study group or maturing into an adult male based on his morphological
characteristics (fully developed testes and canines, full body length larger than that of an adult
female). For our statistical analysis, we extracted Elo-scores for the day that attention to a social
scene was observed, modeled Elo-scores standardized per group[50] and include group ID as a random
effect.

Social bond strength – DDSI

As a measure of the strength of dyadic social bond strength, we calculated Dynamic Dyadic Sociality
Indices (DDSI) and followed De Moor et al.[51] and Stranks et al.[47] in their application of this method
originally developed by Kulik and Mundry[52]. We focused strictly on adult male dyads, including only
interactions between adult males in our computations. We based our DDSI scores on the time each
dyad spent in body contact and allo-grooming, with each of these behaviors being weighed by the
inverse of their frequency. Scores for all dyads started at the standard value of 0.5 (DDSI scores
range from 0 to 1), indicative of a neutral relationship. Since affiliative interactions were rare, DDSI
values were not well differentiated during the first half of the study. Thus, in all models, social bond
strength was estimated as the DDSI value for that dyad at the end of the study period when
calculations included a maximum of information.

Statistical analysis
Looking probability model

We first tested the probability of a subject to look versus to not look at a scene as the response
variable. Looks were only considered when occurring in the first 5 seconds after the start of social
scene or else we scored a zero. To estimate how the probability of looking at a particular social
scene varies across different conditions, and combinations of dominance relation and social bond

strength, we fitted a Generalized Linear Mixed Model (GLMM)[53] with binomial error structure and

logit link function[54]. We included a three-way interaction term between condition (agonism,

affiliation, and non-interacting control), dominance relation between the subject and the higher

ranking of the two stimulus individuals ($\Delta R_{sub-max}$ = Elo subject minus Elo higher ranking stimulus),

and maximum DDSI value between the observer and each one of the interacting partners ($DDSI_{sub-max}$).

After we found that the three-way interaction term was not significant (see supplementary

materials) we excluded it and ran a reduced model including the two-way interactions between the

terms of the previous three-way interactions (see Table S2). In a last step, we excluded the non-

significant interaction term dominance relation x max. DDSI and report below the final model

including only two of the three 2-way interaction terms. In all models, we included rank of the

subject (R_{sub}) as well as conspicuity of the interaction as further fixed effects.

To control for their effects and avoid pseudo-replication, we included random intercept

effects for study group, identity of the subject, identity of the actor, and identity of the receiver in

the social scene. To avoid an overconfident model and secure a type I error rate at the nominal level

of 0.05 we included all theoretically identifiable random slopes[55], namely those of the three-way

interaction term and all fixed effects within each random intercept. Though we initially aimed at

including also all correlations among random intercepts and slopes[56] , we were forced to exclude

them to allow for successful model convergence.

Prior to fitting the model, we first re-leveled the different levels of our categorical predictor

for condition, defining controls (non-interacting) as the reference level for more intuitive result

interpretation. We then inspected the distributions of dominance relation ($\Delta R_{sub-max}$), maximum DDSI

($DDSI_{sub-max}$), and rank of the subject (R_{sub}), which overall appeared roughly symmetrical and free of

outliers. We z-transformed these predictors to a mean of zero and standard deviation of one to ease

model convergence. We manually dummy coded and centered the control predictor (conspicuity of

interaction) before including it as a random slope. The null model comprised only conspicuousness

and the random effects.

The sample analyzed with the looking probability model comprised a total of 962 social

attention situations, of which 337 were non-interacting controls, 354 affiliative interactions, and 271

agonistic events. These observations concerned a total of 56 subjects and 57 stimulus individuals.

Looks occurred in a total of 364 situations, out of the total 962.

Looking duration model

In order to quantify variation in the duration of sustained attention, we ran a looking duration model

using a subset of our full dataset corresponding only to situations where looks occurred. Here, we

used *looking time* as a response, which had been widely used in previous human and non-human

primate studies as an indicator of preferences and discriminative abilities[57], potentially indicating a

certain type of motivation to sustain attention.

To estimate the extent to which looking time varied across different conditions, and

combinations of dominance relations and social bond strengths, we fitted a Generalized Linear

Mixed Model (GLMM)[53] with gamma error structure and log link function[58]. This model was strictly

identical to our looking probability model in both fixed and random effects structure. Nonetheless,

to control for the effect that longer social scenes may have on looking duration, we here included

the duration of the stimulus interaction as a further fixed effect. We removed all correlations among

random intercepts and slopes[56] in an attempt to achieve successful model convergence. Since we

faced convergence issues from the start, we ultimately fitted a Bayesian model using the brms

package in R. The model was run using 4 MCMC chains of 4,000 iterations per chain with

adapt_delta set at 0.99 and a max_treedepth of 15, including 1,000 "warm up" iterations during

which no divergent transitions were reported. This resulted in 12,000 posterior samples per model.

We assessed the distributions of dominance relations ($\Delta R_{sub-max}$), maximum social bond

strength ($DDSI_{sub-max}$), and rank of the subject (R_{sub}), which overall appeared roughly symmetrical and

free of outliers. We z-transformed these predictors to a mean of zero and standard deviation of one

to ease model convergence. We further inspected the distribution of the response (looking duration)

and the added fixed effects of duration of the stimulus situation, both of which had heavily right-

skewed distributions. To deal with this, we log transformed interaction duration. Given that we were

fitting a model with a log link function, we kept the response as is. We manually dummy coded and

centered the control predictor (conspicuity of interaction) before including it as a random slope. We

defined weakly informative priors for all model parameters as to not strongly restrict posterior

distributions. For all fixed effects we used a Normal (0, 1) prior, Exponential (1) priors for the

standard deviations of random effects, and Exponential (1) for the shape parameter.

The sample analyzed with this model comprised a total of 441 social scenes that were

looked at including those where looks occurred later than 5 seconds after scene onset. Ninety-nine

scenes were non-interacting controls, 160 affiliative interactions, and 182 agonistic interactions.

These observations involved 56 different males as subjects. Looking duration ranged from 1 to 49

seconds.

Statistical model implementation

The looking probability model was fitted in R (version 4.4.3)[49], using the function glmer of the

package lme4 (version 1.1-35.1)[59]. We determined confidence limits of model estimates and fitted

values by means of a parametric bootstrap (N=1,000 bootstraps; function bootMer of the package

lme4). We determined model stability by dropping the levels of the grouping factors one by one,

fitting the full model to each of the subsets, and finally comparing the range of estimates with those

obtained for the full data set. This revealed moderate to good stability of the estimates (see Table 1

in the results section). We determined the significance of individual fixed effects by means of

likelihood ratio tests[60] comparing the full model with models lacking them one at a time (R function

drop1).

The looking duration model was fitted in R (version 4.4.3)[49], using the function brm of the

package brms (version 2.22.0), which builds on the computational framework Stan [https://mc-

stan.org] to fit Bayesian models[61]. The convergence and stability of the models were assessed with

R-hat values, which were below 1.01[62] and a visual inspection of the plots of the chains[63]. In

addition, all Bulk and Tail Effective Sample Sizes (ESS) for the estimated posteriors were >6000 [61,62].

RESULTS

Results are based on the behavior of 56 male subjects that had the opportunity to visually attend to

962 social scenes involving any combination of two males co-residing in their group. Social scenes

included 271 agonistic events, 354 affiliative interactions, and 337 non-interacting control situations

where two adult males were in close proximity of each other and did not interact via vocal exchange,

facial expression, or tactile behavior.

Figure 2: Variation in visual attention to 3rd person social scenes as predicted from the looking
probability model. Subjects generally attended more often to interactions than control scenes and to
agonistic than affiliative interactions. Both the 2a) negative effect of dominance relations (subject
minus dominant stimulus) and the 2b) positive effect of social bond strength (subject to stimuli) were
modulated independently by condition and were stronger for agonistic than for affiliative
interactions. Subject rank 2c) did not affect visual attention. Bubbles are average probabilities per
observed value of the predictor with size representing the number of observations, lines are model
fitted estimates and shades their confident intervals (Tab. 1). Data came from 56 subjects and 962
social scenes.

Selective attention (looking probability)

In the looking probability model, the three-way interaction term between condition, dominance relations, and maximum DDSI was not significant and subsequently was excluded from our first reduced model (Tab. S1). To arrive at our final model, we further excluded the non-significant two-way interaction maximum DDSI x dominance relations (Tab. S2). The final full model differed significantly from the null model including only the control variable conspicuity of the interaction and the random effects of subject ID and group (Chi^2 = 34.02, df = 11, P<0.001).

Based on the parameter estimates and conditional on the interaction effects, subjects were

more likely to attend to interactions than to non-interactive controls (Tab. 1, Fig. 2). The higher

above the subject the higher-ranking of the stimulus individual was, i.e. the more negative the

dominance relations, the more likely was the subject to pay attention to an agonistic interaction and

the lower below the subject the stimulus individuals were, the less likely the event was attended to.

This effect of dominance relations on overt attention differed between conditions. Specifically, the

effect of dominance relations was stronger (more negative estimate) for non-interacting controls

than for affiliative interaction stimuli, but not stronger than for agonistic scenes.

Furthermore, stimulus valence modulated the effect of social bond strength on visual

attention. The closer the subject was affiliated to the stimuli (the higher the maximum DDSI), the

more likely they paid attention which was an effect exacerbated for agonistic interactions, still

pronounced for affiliative 3rd person interactions and weakest for non-interacting control scenes.

Beyond the relational effects of dominance relations, the individual dominance rank of the

subject did not affect its probability to monitor stimuli across conditions (see Fig. 3). Situations that

were more conspicuous were more likely to be looked at.

Table 1. *Model estimates for the reduced looking probability model, including as predictors only the*
interaction term between condition and dominance relations between the subject and the dominant
of the stimulus individuals. The full model was significantly different from the null model including
only control factors (conspicuousness of the interaction), and random effects (Chi^2 = 34.02, df = 11,
P<0.001). Significant effects of test predictors are highlighted in bold. Significance of the different
levels of condition was assessed by releveling the intercept.

	Estimate	Std. Error	z value	Pr(>|z|)
Intercept	-1.629	0.185	-8.817	
Affiliation vs. Control	**0.887**	**0.226**	**3.925**	**<0.001**
Agonism vs. Control	**1.658**	**0.264**	**6.267**	**<0.001**
Agonism vs. Affiliation	**0.770**	**0.241**	**3.198**	**0.001**
Dominance relations	**-0.614**	**0.191**	**-3.208**	**0.001**
Social bond strength (max DDSI)	-0.043	0.221	-0.192	0.847
Subject rank	0.099	0.137	0.726	0.468
Conspicuousness	1.777	0.411	4.321	<0.001
Affiliation vs. Control x Dominance relations	**0.426**	**0.208**	**2.048**	**0.041**
Agonism vs. Control x Dominance relations	0.243	0.253	0.96	0.337
Agonism vs. Affiliation x Dominance relations	-0.175	0.230	-0.764	0.445
Affiliation vs. Control x Social bond strength	0.126	0.216	0.581	0.561
Agonism vs. Control x Social bond	**0.64**	**0.279**	**2.293**	**0.022**

strength				
Agonism vs. Affiliation x Social bond strength	0.515	0.269	1.914	0.056

Figure 3: Sustained attention for 3rd person interactions was longer than for non-interaction social control scenes and longer for agonistic than for affiliative interactions (Tab. 2) with distributions of posterior estimates for the looking duration model shown in 3a). Violin plots are drawn for raw data 3b) and whisker plot shows model estimates with 95% credible interval.

Sustained attention (looking duration)

We used a Bayesian model to estimate the effects of the same predictor variables on the time spent looking at the social scene. Posterior probabilities for the three-way-interaction between dominance relations, social bond strength and condition were centered around zero (Tab. S3). In a reduced model without the 3-way interaction term, we found no evidence supporting either of the three 2-way interaction effects (all posterior probabilities centered around zero). The final model (Tab. 2) including only main effects provided clear evidence for sustained attention to vary with condition. Males demonstrated longer attention for agonistic interactions between other males than for scenes with two non-interacting stimulus males (posterior probabilities >0.95 for difference between control and each of agonistic and affiliative scenes; Fig. 3). Furthermore, there was moderate

evidence for longer looking time towards agonistic compared to affiliative interactions (posterior

probability 0.84). Looking time during agonistic situations was on average 0.55s longer compared to

non-interacting controls, and 0.23s longer than affiliation. Dominance relations and social bonds to

the stimulus males and subject rank did not affect sustained attention.

Table 2. Model estimates for the reduced looking duration model including only main effects;
posterior means are reported with their standard errors and 95% credible intervals for the fixed
effects. Estimates are based on 4,000 posterior draws.

	Estimate	Est.Error	Q2.5	Q97.5
Intercept	0.171	0.206	-0.234	0.573
Affiliation vs. Control	**0.239**	**0.164**	**-0.086**	**0.565**
Agonism vs. Control	**0.380**	**0.215**	**-0.082**	**0.774**
Agonism vs. Affiliation	0.141	0.244	-0.362	0.592
Dominance relations	0.014	0.105	-0.198	0.218
Social bond strength (max DDSI subject – stimulus)	0.035	0.109	-0.184	0.251
Subject rank	-0.042	0.121	-0.263	0.194
Log interaction time	0.095	0.058	-0.016	0.210
Conspicuousness	0.359	0.178	-0.020	0.684

Inter-individual differences

Random effects from both models suggest that individuals do not systematically differ in attention

to just any type of social scene. Instead, significant individual differences were specific to and

differed between sustained attention to affiliative scenes (mean individual random effect for

affiliative scenes ± stddev. = 0.0013 ± 0. 174; Fig. 4a) and agonistic scenes (0.0012 ± 0.192; Fig.4b).

These random effects were extracted from a model with and are therefore controlled for subject

dominance rank.

Figure 4: Inter-individual variation in average looking duration. Posterior distributions are shown for a) affiliative scenes and b) agonistic scenes for all study subjects relative to the estimated population average. Black dots show the posterior mean for each subject. Positive estimates indicate above-average looking durations, whereas negative estimates indicate below-average looking durations.

DISCUSSION

Our study contributes to the development of real-life social neuroscience by elucidating the neural mechanisms underlying social information integration. Social interactions are inherently complex, requiring individuals to make predictions not only about their own actions but also about those of others, informed by observations of 3rd person interactions[13]. We found that individuals allocate increased visual attention to 3rd person interactions, particularly those involving agonistic versus affiliative behaviors, and that different dimensions of social relationships between the subject and third parties independently modulate this attentional bias. Furthermore, we observed significant individual differences in sustained attention to others' interactions, underscoring the importance of

considering personal factors in understanding social cognition. These results have important
implications for our understanding of how the brain integrates social information and how emotion
regulation and adaptive benefits influence sustained attention.

3[rd] person interactions draw more attention than non-interacting social controls

Using as a control a social scene with two individuals in close proximity without an interaction, we
clearly demonstrated that both the probability of overt visual attention and the duration of
sustained visual attention were increased for 3[rd] person interactions. The three social conditions
were matched for salience by requiring the scene to have a clear starting point where two males
arrived in close proximity, i.e. all scenes were initially dynamic. By requiring this conservative control
condition, we exclude the possibility that interaction scenes draw more attention, simply because
perceptually two individuals may be more salient[21]. Exceedingly dynamic scenes that involved quick
movements and/or vocalizations either in the agonistic or the affiliative context (e.g. ritualized male-
infant-male interactions)[64] were scored to be conspicuous which was a control variable in all models.
Thus, visual attention was not only guided by stimulus salience alone, but also by the mental
representation of its social relevance[6,65,66] and was sustained into a range where MEG studies
suggest different information integration events can be detected (1,100 and 2,000 ms)[67]. To our
knowledge, the only other study to demonstrate that 3[rd] person interactions attract more visual
attention is a three player experiment with rhesus macaques that identified neurons in the
prefrontal cortex to be differentially activated when two others interacted[23].

Attention guided by valence of others' interactions

Visual attention was further guided by interaction valence, adding ecological validity to previous
findings on selective attention for and differential processing of threat faces over photos of friendly
teeth-chattering[2,68]. Here the naturalism scale seems particularly relevant, because the two facial
expressions differ in important ways; teeth chattering is a repetitive and thus inherently more
dynamic and auditory display than a threat face and therefore may be more difficult to identify from
a still photograph. Even if videos of agonistic versus affiliative interactions of conspecifics are used,
chimpanzees attend equally long to both conditions[21]. Yet even in our real-life scenes in proximity of
the subject, monkeys attended more to agonistic than affiliative interactions of others.

Agonistic interactions may be more relevant, because the subject could be drawn into the
conflict by redirected aggression[69] or be solicited to help one of the opponents[70,71]. The fact that

watching others fight increases rates of self-directed behaviors indicative of increased emotional
arousal[72,73], suggests that subjects do not only gather information about the state of the dominance
relations of others, but also about imminent threat to themselves. Others' affiliative interactions
may also lead to direct involvement of the subject, if they decide to interfere by joining or stopping
the affiliation[39,40], but this decision lies with the subject and not with a 3rd party and may therefore
not require the same level of attention as 3rd person agonism.

Despite the primacy of agonism, subjects still paid more often and more sustained attention
to 3rd person friendly interactions than non-interacting controls. Primates look towards the mother
(the main affiliate), if they hear an infant in distress from the other direction[74] and look longer at a
staged conflict, if it is between close affiliation partners compared to other dyads[75]. Selective
attention for 3rd person friendly interactions must underlie such findings, but had not been
previously demonstrated.

Two dimension of social relationships independently guide visual attention

Variation in imminent threat may also explain how the social relationships between the subject and
the stimulus individuals have moderated valence effects on selective attention. Since aggression
flows down the hierarchy in macaques[76], imminent threat increases the higher above the subject the
dominant stimulus male. This effect appears most pronounced in non-interactive control scenes,
because in controls, attention drops to zero for scenes that only have males subordinate to the
subject; at the largest dominance differences (left of Fig. 2a) control scenes did not attract more
attention than agonistic or affiliative scenes. Attention to non-interacting control scenes with
individuals out-ranking the subject may serve in uncertainty-reduction[77]. Attention for conflicts was
much higher than for controls, even when both males were subordinate to the subject, so that the
rank-difference had a slightly weaker effect. Dominance difference did not moderate visual attention
to affiliative interaction scenes which suggests that it is not the mere presence of a higher ranking
individual that adds relevancy, but the action of a dominant.

We expected that monitoring close affiliates may also contribute towards risk-avoidance,
given that the conflicts of closely bonded partners are the most likely to trigger risky interactions for
the observing party[78,79] which requires behavioral coordination between close affiliates[36,80].
Matching these expectations, we found the attention-grabbing effect of social bond strength with
stimuli to vary with stimulus valence: the social bond effect was most pronounced for conflicts and
less so for affiliation and control scenes. This highlights again the increased relevance of potentially

risky situations. In other real-life observational studies on captive populations, female rhesus

macaques and mandrills of both sexes also paid more attention to their closest partner and their kin,

respectively[35,81]. This is in contrast to a recent study on male rhesus macaques[34]; the grooming

relationship did not affect visual attention for pictures of a group mate, whereas the aggressive

relationship with a stimulus clearly correlated with attention for this male[34]. By moving from the

presentation of still photographs to real-life stimuli and from individual stimuli[81] to interactions, we

demonstrate that two dimensions of a subject's relationships to others have independent effects on

the allocation of visual attention which is more pronounced for agonism than affiliation of others.

Individual differences in social attention

For good reasons, the most controlled studies employing brain-imaging or electrophysiology often

suffer from small sample sizes which unfortunately prohibits investigation of individual differences in

social attention and underlying processing[1]. Here, we find no evidence for individuals to generally

differ in their social attention or for social status. Instead, the 56 male Assamese macaques varied in

different ways in how they attended to agonistic interactions or to affiliation between others.

In a well-powered study on rhesus macaques, significant year-to-year repeatability was found in

attention to threat over neutral faces, but not in attention to friendly teeth-chattering over neutral

faces[82]. Differences between four rhesus macaques in the propensity to give aggression, grooming,

and approaches over receiving these behaviors were all associated to some degree with the social

attention they gained in a distractor task[34]. Neither of the rhesus macaque studies, however,

controlled for dominance rank prior to extracting information on individual differences. We can

clearly show here that despite dominance changing throughout the study period, individuals

maintain differences in how they attend to conflicts of others, suggesting stable differences in

underlying regulation via serotoninergic and oxytocinergic pathways[33,34].

Conclusion – information integration for visual attention

Together, these results suggest that different relevancies of social information are integrated in

complex ways. Interaction versus non-interacting control, valence of the interaction, and two

separate relational aspects do not have simple additive effects on social attention. The social value

of attention is evident from increased attention to others' affiliation over non-interacting controls

that is unaffected by dominance differences or social bonds towards the stimuli. All other results can

be explained by risk-avoidance alone where information is not gathered for its own sake, but

emotion regulation prepares and activates the subject in the face of imminent danger. Thus, social

attention serves at least two goals in tandem, which explains why it involves several brain areas in

the dorsomedial prefrontal cortex, dorsal anterior cingulate cortex, the ventral striatum and the

amygdala perhaps with pronounced mixed selectivity at the neuronal and network level[9,23,83,84].

ACKNOWLEDGEMENTS

We are grateful to the National Research Council of Thailand and the Department of National Parks,

Wildlife and Plant Conservation of Thailand for permission from to conduct research in a protected

area (permits: 0907.4/9313 April 2[nd] 2022, 0907.4/19458 September 3[rd] 2023). We thank the

superintendent of PKWS Wichanon Saenphala for his collaboration, Roger Mundry for statistical

advice, and the team of foreign and Thai field researchers that supported data collection. This study

was funded by the Deutsche Forschungsgesellschaft DFG- SFB 1528 Cognition of Interaction (project-

ID 454648639).

AUTHOR CONTRIBUTIONS

SP Investigation, Data curation, Formal analysis, Visualization, Writing - Original draft

SMa Resources, Project administration, Writing - Review & editing

SMe Resources, Project administration, Writing - Review & editing

JO Conceptualization, Funding acquisition, Data curation, Resources, Supervision, Writing - Review &

editing

OS Conceptualization, Funding acquisition, Data curation, Resources, Supervision, Writing - Original

draft

CONFLITCS OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY

Replication data are available at GRO.data https://doi.org/10.25625/XFN4B4

ETHICAL STATEMENT

The study was conducted with permission of relevant Thai authorities under a benefit sharing
agreement and was purely observational, completely non-invasive and did not involve food
provisioning or capture of the animals.

REFERENCES

1. Fan, S., Monte, O. D. & Chang, S. W. C. Levels of naturalism in social neuroscience research.

iScience **24**, (2021).

2. Gothard, K. M., Battaglia, F. P., Erickson, C. A., Spitler, K. M. & Amaral, D. G. Neural responses to

facial expression and face identity in the monkey amygdala. *J. Neurophysiol.* **97**, 1671–1683

(2007).

3. Kret, M. E., Muramatsu, A. & Matsuzawa, T. Emotion processing across and within species: A

comparison between humans (Homo sapiens) and chimpanzees (Pan troglodytes). *J. Comp.*

Psychol. **132**, 395–409 (2018).

4. Schacht, A. & Vrtička, P. Spatiotemporal pattern of appraising social and emotional relevance:

Evidence from event-related brain potentials. *Cogn. Affect. Behav. Neurosci.* **18**, 1172–1187

(2018).

5. Berg, D. J., Boehnke, S. E., Marino, R. A., Munoz, D. P. & Itti, L. Free viewing of dynamic stimuli by

humans and monkeys. *J. Vis.* **9**, 19 (2009).

6. Sliwa, J. & Freiwald, W. A. A dedicated network for social interaction processing in the primate

brain. *Science* **356**, 745–749 (2017).

7. Siebert, R. *et al.* A naturalistic dynamic monkey head avatar elicits species-typical reactions and

overcomes the uncanny valley. *eNeuro* **7**, (2020).

8. Wilson, V. A. D. *et al.* Macaque gaze responses to the primatar: a virtual macaque head for social

cognition research. *Front. Psychol.* **11**, (2020).

9. Haroush, K. & Williams, Z. M. Neuronal prediction of opponent's behavior during cooperative

social interchange in primates. *Cell* **160**, 1233–1245 (2015).

10. Dal Monte, O., Piva, M., Morris, J. A. & Chang, S. W. C. Live interaction distinctively shapes social

gaze dynamics in rhesus macaques. *J. Neurophysiol.* **116**, 1626–1643 (2016).

11. Seyfarth, R. M. & Cheney, D. L. Social cognition. *Anim. Behav.* **103**, 191–202 (2015).

12. Klein, J. T., Shepherd, S. V. & Platt, M. L. Social attention and the brain. *Curr. Biol. CB* **19**, R958-

962 (2009).

13. Freiwald, W. A. Social interaction networks in the primate brain. *Curr. Opin. Neurobiol.* **65**, 49–58

(2020).

14. Andrews, M. W., Bhat, M. C. & Rosenblum, L. A. Acquisition and long-term patterning of joystick

selection of food-pellet vs social-video reward by Bonnet Macaques. *Learn. Motiv.* **26**, 370–379

(1995).

15. Deaner, R. O., Khera, A. V. & Platt, M. L. Monkeys pay per view: adaptive valuation of social

images by rhesus macaques. *Curr. Biol. CB* **15**, 543–548 (2005).

16. Lindsay, G. W. Attention in psychology, neuroscience, and machine learning. *Front. Comput.*

Neurosci. **14**, 29 (2020).

17. Lacreuse, A. *et al.* Testosterone may increase selective attention to threat in young male

macaques. *Horm. Behav.* **58**, 854–863 (2010).

18. McFarland, R. *et al.* Social interactions through the eyes of macaques and humans. *PLoS ONE* **8**,

e56437 (2013).

19. Teufel, C., Alexis, D. M., Clayton, N. S. & Davis, G. Mental-state attribution drives rapid, reflexive

gaze following. *Atten. Percept. Psychophys.* **72**, 695–705 (2010).

20. Adams, G. K., Ong, W. S., Pearson, J. M., Watson, K. K. & Platt, M. L. Neurons in primate

prefrontal cortex signal valuable social information during natural viewing. *Philos. Trans. R. Soc. B*

Biol. Sci. **376**, 20190666 (2021).

21. Lewis, L. S. *et al.* Chimpanzees and children are curious about social interactions. *Proc. R. Soc. B*

Biol. Sci. **292**, 20242242 (2025).

22. Dal Monte, O. *et al.* Widespread implementations of interactive social gaze neurons in the

primate prefrontal-amygdala networks. *Neuron* **110**, 2183-2197.e7 (2022).

23. Báez-Mendoza, R., Mastrobattista, E. P., Wang, A. J. & Williams, Z. M. Social agent identity cells

in the prefrontal cortex of interacting groups of primates. *Science* **374**, eabb4149 (2021).

24. Xing, F., Sheffield, A. G., Jadi, M. P., Chang, S. W. & Nandy, A. S. Dynamic modulation of social

gaze by sex and familiarity in marmoset dyads. *eLife* **14**, (2025).

25. Chance, M. R. A. Attention structure as the basis of primate rank orders. *Man* **2**, 503–518 (1967).

26. Shepherd, S. V., Deaner, R. O. & Platt, M. L. Social status gates social attention in monkeys. *Curr.*

Biol. **16**, R119–R120 (2006).

27. Cheng, J. T. *et al.* Eye gaze and visual attention as a window into leadership and followership: A

review of empirical insights and future directions. *Leadersh. Q.* **34**, 101654 (2023).

28. Capitanio, J. P. Sociability and responses to video playbacks in adult male rhesus monkeys

(Macaca mulatta). *Primates* **43**, 169–177 (2002).

29. Machado, C. J., Whitaker, A. M., Smith, S. E. P., Patterson, P. H. & Bauman, M. D. Maternal

Immune Activation in Nonhuman Primates Alters Social Attention in Juvenile Offspring. *Biol.*

Psychiatry **77**, 823–832 (2015).

30. Mosher, C. P., Zimmerman, P. E. & Gothard, K. M. Videos of conspecifics elicit interactive looking

patterns and facial expressions in monkeys. *Behav. Neurosci.* **125**, 639–652 (2011).

31. Watson, K. K., Ghodasra, J. H. & Platt, M. L. Serotonin transporter genotype modulates social

reward and punishment in rhesus macaques. *PLOS ONE* **4**, e4156 (2009).

32. Weinberg-Wolf, H. *et al.* The effects of 5-hydroxytryptophan on attention and central serotonin

neurochemistry in the rhesus macaque. *Neuropsychopharmacology* **43**, 1589–1598 (2018).

33. Weinberg-Wolf, H. & Chang, S. W. C. Differences in how macaques monitor others: Does

serotonin play a central role? *WIREs Cogn. Sci.* **10**, e1494 (2019).

34. Liu, S., Huang, J., Chen, S., Platt, M. L. & Yang, Y. Multi-dimensional social relationships shape

social attention in monkeys. *eLife* **14**, RP104460 (2025).

35. Schülke, O., Dumdey, N. & Ostner, J. Selective attention for affiliative and agonistic interactions

of dominants and close affiliates in macaques. *Sci. Rep.* **10**, 5962 (2020).

36. Watts, D. P., Colmenares, F. & Arnold, K. Redirection, consolation, and male policing: hor targets

of aggression interact with Bystanders. in *Natural conflict resolution* (eds Aureli, F. & de Waal, F.

B. M.) 281–304 (University of California Press, Berkley, 2000).

37. Schino, G. Grooming and agonistic support: a meta-analysis of primate reciprocal altruism.

Behav. Ecol. **18**, 115–120 (2007).

38. Ostner, J. & Schülke, O. The evolution of social bonds in primate males. *Behaviour* **151**, 871–906

(2014).

39. Mielke, A. *et al.* Bystanders intervene to impede grooming in Western chimpanzees and sooty

mangabeys. *R. Soc. Open Sci.* **4**, (2017).

40. Mielke, A., Bruchmann, C., Schülke, O. & Ostner, J. Grooming interventions in female rhesus

macaques as social niche construction. *Anim. Behav.* **173**, 105–114 (2021).

41. Schülke, O. *et al.* Quantifying within-group variation in sociality—covariation among metrics and

patterns across primate groups and species. *Behav. Ecol. Sociobiol.* **76**, 50 (2022).

42. Ostner, J. & Schülke, O. Linking sociality to fitness in primates: a call for mechanisms. in *Advances*

in the Study of Behavior (eds Naguib, M. et al.) vol. 50 127–175 (Academic Press, 2018).

43. Borries, C., Larney, E., Kreetiyutanont, K. & Koenig, A. The diurnal primate community in a dry

evergreen forest in Phu Khieo Wildlife Sanctuary, Northeast Thailand. *Nat. Hist. Bull. Siam Soc.* **50**,

75–88 (2002).

44. Altmann, J. Observational study of behavior: sampling methods. *Behaviour* **49**, 227–266 (1974).

45. Martin, P. & Bateson, P. P. G. *Measuring Behaviour: An Introductory Guide.* (Cambridge Univ.

Press, Cambridge, 2006).

46. Neumann, C. & Fischer, J. Extending Bayesian Elo-rating to quantify the steepness of dominance

hierarchies. *Methods Ecol. Evol.* **14**, 669–682 (2023).

47. Stranks, J., Heistermann, M., Sangmaneedet, S., Schülke, O. & Ostner, J. The dynamics of sociality

and glucocorticoids in wild male Assamese macaques. *Horm. Behav.* **164**, 105604 (2024).

48. Neumann, C. EloRating.Bayes (GitHub Repository: https://github.com/gobbios/EloRating.Bayes).

(2024).

49. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing (2023).

50. Levy, E. *et al.* A comparison of dominance rank metrics reveals multiple competitive landscapes

in an animal society. *Proc. Biol. Sci.* **287**, 20201013 (2020).

51. De Moor, D., Roos, C., Ostner, J. & Schülke, O. Bonds of bros and brothers: Kinship and social

bonding in postdispersal male macaques. *Mol. Ecol.* **29**, 3346–3360 (2020).

52. Kulik, L. PhD Thesis: Development and consequences of social behavior in rhesus macaques

(Macaca mulatta). (Univ. Leipzig, Leipzig, 2015).

53. Baayen, R. H. *Analyzing Linguistic Data: A Practical Introduction to Statistics Using R*. vol. 49

(Cambridge Univ. Press, Cambridge, 2015).

54. McCullagh, P. *Generalized Linear Models*. (Routledge, London, 2019).

55. Schielzeth, H. & Forstmeier, W. Conclusions beyond support: overconfident estimates in mixed

models. *Behav. Ecol.* **20**, 416–420 (2009).

56. Barr, D. J. Random effects structure for testing interactions in linear mixed-effects models. *Front.*

Psychol. **4**, (2013).

57. Winters, S., Dubuc, C. & Higham, J. P. Perspectives: The looking time experimental paradigm in

studies of animal visual perception and cognition. *Ethology* **121**, 625–640 (2015).

58. Zuur, A. F. *A Beginner's Guide to Generalized Additive Models with R*. (Highland Statistics Limited,

Newburgh, NY, 2012).

59. Bates, D., Maechler, M. & Bolker, B. M. *Lme4: Linear Mixed-Effects Models Using S4 Classes. R*

Package (Version 0.999375-34). R-Project.Org/Package=lme4. (Institute for Statistics and

Mathematics, University of Economics and Business, Vienna, 2011).

60. Dobson, A. & Barnett, A. *An Introduction to Generalized Linear Models, Fourth Edition.* (Chapman

and Hall/CRC, 2018). doi:10.1201/9781315182780.

61. Bürkner, P.-C. Bayesian Item Response Modeling in *R* with **brms** and *Stan. J. Stat. Softw.* **100**, 1–

54 (2021).

62. Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. Rank-Normalization, Folding,

and Localization: An Improved Rˆ for Assessing Convergence of MCMC (with Discussion). *Bayesian*

Anal. **16**, 667–718 (2021).

63. Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. Visualization in Bayesian

Workflow. *J. R. Stat. Soc. Ser. A Stat. Soc.* **182**, 389–402 (2019).

64. Kalbitz, J., Schülke, O. & Ostner, J. Triadic male-infant-male interaction serves in bond

maintenance in male Assamese macaques. *PLoS ONE* **12**, e0183981 (2017).

65. Gobel, M. S., Tufft, M. R. A. & Richardson, D. C. Social beliefs and visual attention: how the social

relevance of a cue influences spatial orienting. *Cogn. Sci.* **42 Suppl 1**, 161–185 (2018).

66. Ninomiya, Taheri, Noritake, A., & Isoda, Masaki. Live agent preference and social action

monitoring in the macaque mid-superior temporal sulcus region. *Proc. Natl. Acad. Sci.* **118**,

e2109653118.

67. Ulloa, J. L., Puce, A., Hugueville, L. & George, N. Sustained neural activity to gaze and emotion

perception in dynamic social scenes. *Soc. Cogn. Affect. Neurosci.* **9**, 350–357 (2014).

68. Lacreuse, A., Schatz, K., Strazzullo, S., King, H. M. & Ready, R. Attentional biases and memory for

emotional stimuli in men and male rhesus monkeys. *Anim. Cogn.* **16**, 861–871 (2013).

69. Aureli, F. & Schaik, C. P. V. Post-conflict Behaviour in Long-tailed Macaques (*Macaca fascicularis*

): II. Coping with the Uncertainty. *Ethology* **89**, 101–114 (1991).

70. Young, C., Majolo, B., Schülke, O. & Ostner, J. Male social bonds predict partner recruitment in

cooperative aggression in wild Barbary macaques. *Anim. Behav.* **80**, 675–682 (2014).

71. Perry, S., Barrett, H. C. & Manson, J. H. White-faced capuchin monkeys show triadic awareness in

their choice of allies. *Anim. Behav.* **67**, 165–170 (2004).

72. Whitehouse, J., Micheletta, J., Kaminski, J. & Waller, B. M. Macaques attend to scratching in

others. *Anim. Behav.* **122**, 169–175 (2016).

73. Paschek, N., Müller, N., Heistermann, M., Ostner, J. & Schülke, O. Subtypes of aggression and

their relation to anxiety in Barbary macaques. *Aggress. Behav.* **45**, 120–128 (2019).

74. Cheney, D. L. & Seyfarth, R. M. Vocal recognition in free-ranging vervet monkeys. *Anim. Behav.*

28, 362–367 (1980).

75. Whitehouse, J. & Meunier, H. An understanding of third-party friendships in a tolerant macaque.

Sci. Rep. **10**, 9777 (2020).

76. Deag, J. M. Aggression and submission in monkey societies. *Anim. Behav.* **25**, 465–474 (1977).

77. Iki, S. & Adachi, I. Monkeys increase scratching when encountering unexpected good fortune. *Sci.*

Rep. **15**, 8386 (2025).

78. Kalbitz, J., Ostner, J. & Schülke, O. Strong, equitable and long-term social bonds in the dispersing

sex in Assamese macaques. *Anim. Behav.* **113**, 13–22 (2016).

79. Schülke, O., Bhagavatula, J., Vigilant, L. & Ostner, J. Social bonds enhance reproductive success in

male macaques. *Curr. Biol.* **20**, 2207–2210 (2010).

80. Dunbar, R. I. M. & Shultz, S. Bondedness and sociality. *Behaviour* **147**, 775–803 (2010).

81. Schino, G. & Sciarretta, M. Patterns of social attention in mandrills, *Mandrillus sphinx. Int. J.*

Primatol. **37**, 752–761 (2016).

82. Diaz, A. A., Hernández-Pacheco, R. & Rosati, A. G. Individual differences in sociocognitive traits in

semi-free-ranging rhesus monkeys (Macaca mulatta). *Am. J. Primatol.* **87**, e23660 (2025).

83. Leopold, D. A. The big mixup: Neural representation during natural modes of primate visual

behavior. *Curr. Opin. Neurobiol.* **88**, 102913 (2024).

84. Chang, S. W. C., Gariépy, J.-F. & Platt, M. L. Neuronal reference frames for social decisions in

primate frontal cortex. *Nat. Neurosci.* **16**, 243–250 (2013).