Complex modulation of visual attention to 3rd person interactions in wild macaques

Sofia M. Pereira[1,2], Suthirote Meesawat[3,4], Suchinda Malaivijitnond[3,4], Julia Ostner[1,2,5] & Oliver Schülke[1,2,5]

[1]Department of Behavioral Ecology, University of Göttingen, Göttingen, Germany

[2]Social Evolution in Primates Group, German Primate Center Leibniz Institute for Primate Research, Göttingen, Germany

[3]Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

[4]National Primate Research Center of Thailand, Chulalongkorn University, Saraburi, Thailand

[5]Leibniz ScienceCampus Primate Cognition, German Primate Center, Leibniz Institute for Primate Research, Göttingen, Germany

ORCID IDs

Sofia M. Pereira: 0000-0002-8352-4664

Suchinda Malaivijitnond: 0000-0003-0897-2632

Suthirote Meesawat: 0000-0001-8612-2436

Oliver Schülke: 0000-0003-0028-9425

Julia Ostner: 0000-0001-6871-9976

Abstract

Social animals rely on socio-cognitive skills to monitor their social environment and make informed decisions. Visual attention to others' interactions in real-life scenes, however, remains understudied although dedicated neuronal networks have been identified for the processing of real-life social information and for social interactions in particular. Here we ask how subject characteristics, interaction valence, and social relationships between subject and stimuli interact to guide overt attention of 57 wild Assamese macaques to 962 3rd person social scenes that spontaneously unfolded in their vicinity. Social interactions drew more and longer attention than control scenes lacking the interaction, highlighting the relevance of interaction. The effects of both affiliative relationship strength and dominance rank difference between subject and stimulus on the probability to attend

were independently modulated by valence of the scene. Beyond reacting to the affordances of the

social scene and independent of their social status, males differed in how long they attended to the

interactions of others. Thus, visual attention is guided by the value an individual associates with

different aspects of the social scene based on previous experience in both the agonistic and the

affiliative realm and by current threat indicative of complex information integration in the brain.

Keywords: social cognition, information integration, valence, social status, social bonds

INTRODUCTION

Social neuroscience research employs a variety of social stimuli that vary in their naturalism[1]. On one

end of the spectrum the subject is chair-seated and presented with static images of social agents and

non-social controls, while their eye gaze and neuronal activity is tracked[2–4]. From there, naturalism is

increased by using videos that are more salient, but also noisier, stimuli[5,6]. Virtual reality scenes can

even more closely mimic real-life interactions, as avatars can display contingent behavior creating a

live-stream of interactions[7,8]. However, real-life agents are the most naturalistic, yet also the most

challenging to control[9,10] particularly in natural environments[1]. This study leverages the strengths of

naturalistic animal observations, where large numbers of subjects can be studied allowing us to weigh

individual differences in social attention against the affordances of the stimulus situation including

subject's affiliative and dominance relations with the stimuli.

Social attention, i.e. monitoring conspecifics to extract information, is a crucial building block of

advanced social cognition[11], involving specialized brain networks[12,13]. Social primates are willing to

forgo an immediate food reward for the privilege to watch conspecifics in videos[14]. Yet, not all social

information is equally relevant. For example, rhesus monkeys (*Macaca mulatta*) pay with forgone

juice rewards for viewing a picture of the face of a dominant male, but have to be paid to look at a

subordinate[15]. Visual attention for faces provides information about others' overt spatial attention,

i.e. about gaze direction[16], and social signals encoded in ritualized facial expressions[2]. In fact,

macaques pay more attention to facial expressions associated with threat than to those associated

with benign intent and even less to neutral faces [17,18] and more readily follow the gaze of others, if

they show an emotional as opposed to a neutral expression [19]. Single neurons in the amygdala show

selective activity for threat, friendly and neutral facial expressions[2].

Thus, social attention targets faces, gazes, and facial expressions as well as relevancies unrelated to

perceptual aspects of the stimulus, e.g. stimulus dominance status, all of which can be tested with

static information. Testing with video recorded dynamic stimuli shows that sustained attention

depends on additional ethogram elements contained in the video sequence[20]; rhesus macaques

looked longer at areas of a frame with a grooming solicitation, aggression or submission[20]. During

fMRI, macaques engage a prefrontal cortical area specifically when watching videos of two others

interacting compared to two others acting each on their own or simply resting; there is a brain for

interaction[6].

Presenting videos of longer natural scenes with multiple real-life agents hugely advances our

understanding of selective social attention in complex environments and the associated

neurophysiology[1,13]. However, dual eye-tracking and electro-physiology demonstrated that direct

interaction with real-life agents draws even more attention and involves different networks across

prefrontal cortical and amygdala areas than watching stills or even videos of the very same agents[10,21].

Neurons in the anterior cingulate cortex predict the behavior of others during social interaction in an

iterated prisoner's dilemma task and disruption selectively hampers mutually beneficial interactions

in rhesus macaques[9]. A network in the dorsomedial prefrontal cortex codes for social partner identity

and tracks these partners' behavior as well as the distribution of food rewards in a real-life social

task[22]. Thus, studies employing real-life interactions promise to shed new light on the processes

involved in social information processing particularly if agents are freely moving[23].

Social attention is dictated by the affordances of the stimulus situation, but can also vary

considerable between individuals. Social status within a group can be a strong predictor of how much

attention an individual pays their group mates[24–26]. In a population with clearly defined and repeatable

inter-individual differences in sociability, low-sociable rhesus macaques paid more attention to any

social information in videos than high-sociable monkeys independent of social status[27]. Early life

adversity, such as experimental maternal immune activation during gestation, can also affect social

attention in juvenile offspring possibly via epigenetic effects[28]. Both, the genotype upstream of the

serotonin transporter gene[29,30] and administered serotonin precursor[31] affect social attention

suggesting the involvement of a serotonergic pathway in generating the observed inter-individual

differences[32]. Additionally, an oxytocinergic pathway has been implicated in linking individual

differences in affiliation and dominance with visual attention[33].

This study focusses on how individuals selectively attend to 3rd person situations, i.e.

interactions among others[13], as they spontaneously unfold (or not) in the vicinity, excluding

interactions involving themselves. We propose that the relevance of agonistic and affiliative 3rd person

interactions differs considerably, and that relational aspects add an additional layer of relevance[34] .

For example, when conflicts arise, they can quickly escalate and involve the subject directly, either

through redirected aggression[35] or by soliciting their help[36]. As a result, social attention for aggressive

interactions may serve a risk avoidance function, and is likely modulated by the dominance

relationship between the subject and the interacting parties. In contrast, the relevance of affiliation

between others is often less immediate and typically indirect. Rates of grooming and agonistic support

are often correlated[36] which means that the state of the grooming network provides indirect

information about the alliance network in the group[37]. Information about grooming will affect the

subject indirectly when it needs to decide whom to support or which affiliative relationships to invest

in. Sometimes, primates also directly and immediately interfere in the grooming interactions of others

to gain access to desirable partners or to manipulate others' relationships[38,39].

In a population of wild Assamese macaques in Thailand, we used behavioral event recording

to quantify the overt attention of 57 adult males to 962 agonistic and affiliative 3rd person interaction

unfolding in subject's proximity (≤5m) or controls where two other adult males came into close

proximity of ≤1.5m without interacting by means of vocalization, facial expression, touch or pose

gesture. With statistical models we show that valence of the social scene (agonism, affiliation, control)

interacted with rank difference and affiliation strength between the subject and the stimulus monkeys

to predict the occurrence of overt attention to the scene suggesting non-additive integration of

different relevancies during processing of 3rd person real-life interactions. We further identify

individual differences in social attention for agonistic interactions that are independent of subject

social status.

METHODS

Study site and subjects

This study was conducted at the Phu Khieo Wildlife Sanctuary in North-Eastern Thailand (PKWS, 16°

05'-35' N, 101° 20'-55' E, 1573km^2) as part of a long-term research project on a wild population of

Assamese macaques ongoing since 2005[40]. PKWS is part of the well-preserved Western Isaan Forest

Complex covering an area of more than 6,500km^2 [41], which houses a diverse community of large

mammals and predators, an indicator of low disturbance levels[41]. We collected data on 57 adult

Assamese macaque males belonging to five different neighboring multimale-multi-female groups with

39 to 75 individuals each and a minimum of eight adult males and ten adult females.

Social attention data collection

Social attention data were collected from June 2022 until April 2023 through event sampling of behavior towards social scenes spontaneously unfolding in the group. A social scene was considered valid for recording each time a stationary adult male subject in the vicinity had an unobstructed view and the human observer clearly saw the onset of the social scene. We recorded head gaze (but also eye gaze if possible) towards three different social scene *conditions* that all started with two adult males (the stimulus individuals) arriving within 1.5 m of proximity of one another and continued either (1) as non-interacting *controls* if the males remained in close proximity without explicitly interacting via touch, vocalization or facial expression or (2) as *affiliation* if the two males only exchanged friendly behaviors, and (3) as *agonism* if the interaction included aggressive or submissive behaviors. Given the specificity of parameters and viewing conditions necessary to accurately record each attention event, these data were collected opportunistically outside the focal animal follows. Upon the unambiguous onset of a social attention event, we recorded the following variables: identities of the subject and two interacting stimulus individuals, behaviors displayed during the stimulus interactions, start and end times of stimulus interactions, timing and duration of each look. We additionally scored the interaction for conspicuousness (0/1) as a means to control statistically for loud or tumultuous interactions drawing attention[34]. We collected data from 962 social attention events, i.e. 17±9 events for each of the 57 subjects.

Figure 1: This field study integrates data on visual attention to social scenes spontaneously unfolding in the vicinity of a subject with information about the affiliative and dominance relationships among all males in five study groups. Social scenes with two adult males in close proximity were classified as non-interacting controls or interactions reflecting affiliation or agonism. The relationship of the subject to the males in the scene was quantified as the dominance difference to the higher-ranking of the other males (Rank difference sub-max) and the maximum affiliative relationship strength to either of the two (DDSI$_{sub-max}$).

Collection of social relationship data

Concomitant with the social attention data collection, all adult males were systematically observed as focal animals in 40 min protocols[42] from June 2022 through April 2023. We recorded continuously [43] all affiliative and agonistic (both aggressive and submissive) interactions including the identities of the actor and receiver as well as all approaches into and departures from a 1.5 m sphere around the focal animal. To inform our dominance and affiliation metrics (see below) about the state of relationships at the onset of the study in June 2022, we included data from January-May 2022. A total of 1,915.9h of continuous focal animal data were collected (mean ± standard deviation per male 33.6h ± 12.1).

Dynamic dominance score – Elo-rating

For the assessment of dominance, we scored submissive behaviors (i.e. bare teeth, make room, give ground) in decided dyadic conflicts (either unprovoked submission or submission following aggression) registered during our continuous focal animal observations and ad libitum recordings from January 2022 through April 2023. To avoid long burn-in periods when males immigrate or mature, we employed Bayesian Elo-rating[44] that estimates start values from the data and captures the state and dynamics of male dominance relationships, which can fluctuate remarkably in Assamese macaques[45]. We used the elo_seq_bayes and extract_elo_b functions of the EloRating.Bayes package[46] in R version 4.3.0[47]. A short burn-in period of one month was applied for each male immigrating into a study group or maturing into an adult male based on his morphological characteristics (fully developed testes and canines, full body length). For our statistical analysis, we extracted Elo-scores for the day that attention to a social scene was observed, modeled Elo-scores standardized per group[48] and include group ID as a random effect.

Dynamic affiliative relationship score - DDSI

As a measure of the strength of dyadic affiliative relationships, we calculated Dynamic Dyadic Sociality

Indices (DDSI) and followed De Moor et al.[49] and Stranks et al.[45] in their application of this method

originally developed by Kulik and Mundry[50]. We focused strictly on adult male dyads, including only

interactions between adult males in our computations. We based our DDSI scores on the time each

dyad spent in body contact and allo-grooming, with each of these behaviors being weighed by the

inverse of their frequency. Scores for all dyads started at the standard value of 0.5 (DDSI scores range

from 0 to 1), indicative of a neutral relationship. Since affiliative interactions were rare, DDSI values

were not well differentiated during the first half of the study. Thus, in all models, the dyadic affiliative

relationships strength was the DDSI value for that dyad at the end of the study period when

calculations included a maximum of information.

Statistical analysis

Looking probability model

We first tested the probability of a subject to look versus to not look at a scene as the response

variable. Looks were only considered when occurring in the first 5 seconds after the start of social

scene or else we scored a zero. To estimate how the probability of looking at a particular social scene

varies across different conditions, and combinations of dominance relationships and affiliative

relationship, we fitted a Generalized Linear Mixed Model (GLMM)[51] with binomial error structure and

logit link function[52]. We included a three-way interaction term between condition (agonism,

affiliation, and non-interacting control), rank difference between the subject and the higher ranking

of the two stimulus individuals ($\Delta R_{sub\text{-}max}$ = Elo subject minus Elo higher ranking stimulus), and

maximum DDSI value between the observer and each one of the interacting partners ($DDSI_{sub\text{-}max}$).

After we found that the three-way interaction term was not significant (see supplementary materials)

we excluded it and ran a reduced model including the two-way interactions between the terms of the

previous three-way interactions (see Table S2). In a last step, we excluded the non-significant

interaction term rank difference x DDSI and report below the final model including only two of the

three 2-way interaction terms. In all models, we included rank of the bystander (R_{sub}) as well as

conspicuity of the interaction as further fixed effects.

To control for their effects and avoid pseudo-replication, we included random intercept

effects for study group, identity of the subject, identity of the actor, and identity of the receiver in the

social scene. To avoid an overconfident model and secure a type I error rate at the nominal level of

0.05 we included all theoretically identifiable random slopes[53], namely those of the three-way

interaction term and all fixed effects within each random intercept. Though we initially aimed at

including also all correlations among random intercepts and slopes[54] , we were forced to exclude them

to allow for successful model convergence.

Prior to fitting the model, we first re-leveled the different levels of our categorical predictor

for condition, defining controls (non-interacting) as the reference level for more intuitive result

interpretation. We then inspected the distributions of rank difference ($\Delta R_{sub-max}$), maximum DDSI

($DDSI_{sub-max}$), and rank of the subject (R_{sub}), which overall appeared roughly symmetrical and free of

outliers. We z-transformed these predictors to a mean of zero and standard deviation of one to ease

model convergence. We manually dummy coded and centered the control predictor (conspicuity of

interaction) before including it as a random slope. The null model comprised only conspicuousness

and the random effects.

The sample analyzed with the looking probability model comprised a total of 962 social

attention situations, of which 337 were non-interacting controls, 354 affiliative interactions, and 271

agonistic events. These observations were linked to a total of 57 subjects and 57 stimulus individuals.

Looks occurred in a total of 364 situations, out of the total 962.

Looking duration model

In order to quantify variation in the duration of sustained attention, we ran a looking duration model

using a subset of our full dataset corresponding only to situations where looks occurred. Here, we

used *looking time* as a response, which had been widely used in previous human and non-human

primate studies as an indicator of preferences and discriminative abilities[55], potentially indicating a

certain type of motivation to sustain observation.

To estimate the extent to which looking time varied across different conditions, and

combinations of dominance and affiliative relationships, we fitted a Generalized Linear Mixed Model

(GLMM)[51] with gamma error structure and log link function[56]. This model was strictly identical to our

looking probability model in both fixed and random effects structure. Nonetheless, to control for the

effect that longer social scenes may have on looking duration, we here included the duration of the

stimulus interaction as a further fixed effect. We removed all correlations among random intercepts

and slopes[54] in an attempt to achieve successful model convergence. Since we faced convergence

issues from the start, we ultimately fitted a Bayesian model using the brms package in R. The model

was run using 4 MCMC chains of 4,000 iterations per chain with adapt_delta set at 0.99 and a

max_treedepth of 15, including 1,000 "warm up" iterations during which no divergent transitions

were reported. This resulted in 12,000 posterior samples per model.

We assessed the distributions of rank difference ($\Delta R_{sub-max}$), maximum DDSI ($DDSI_{sub-max}$), and

rank of the bystander (R_{sub}), which overall appeared roughly symmetrical and free of outliers. We z-

transformed these predictors to a mean of zero and standard deviation of one to ease model

convergence. We further inspected the distribution of the response (looking duration) and the added

fixed effects of duration of the stimulus situation, both of which had heavily right-skewed

distributions. To deal with this, we log transformed interaction duration. Given that we were fitting a

model with a log link function, we kept the response as is. We manually dummy coded and centered

the control predictor (conspicuity of interaction) before including it as a random slope. We defined

weakly informative priors for all model parameters as to not strongly restrict posterior distributions.

For all fixed effects we used a Normal (0, 1) prior, Exponential (1) priors for the standard deviations of

random effects, and Exponential (1) for the shape parameter.

The sample analyzed with this model comprised a total of 441 social scenes, of which 99 were

non-interacting controls, 160 affiliative interactions, and 182 agonistic events. These observations

involved 48 different males as bystanders. Looking duration ranged from 1 to 49 seconds.

Statistical model implementation

The looking probability model was fitted in R (version 4.4.3)[47], using the function glmer of the package

lme4 (version 1.1-35.1)[57]. We determined confidence limits of model estimates and fitted values by

means of a parametric bootstrap (N=1,000 bootstraps; function bootMer of the package lme4). We

determined model stability by dropping the levels of the grouping factors one by one, fitting the full

model to each of the subsets, and finally comparing the range of estimates with those obtained for

the full data set. This revealed moderate to good stability of the estimates (see Table 1 in the results

section). We determined the significance of individual fixed effects by means of likelihood ratio tests[58]

comparing the full model with models lacking them one at a time (R function drop1).

The looking duration model was fitted in R (version 4.4.3)[47], using the function brm of the

package brms (version 2.22.0), which builds on the computational framework Stan [https://mc-

stan.org] to fit Bayesian models[59]. The convergence and stability of the models were assessed with R-

hat values, which were below 1.01[60] and a visual inspection of the plots of the chains[61]. In addition, all

Bulk and Tail Effective Sample Sizes (ESS) for the estimated posteriors were >6000 [59,60].

RESULTS

Results are based on the behavior of 57 male subjects that had the opportunity to visually attend to 962 social scenes involving any combination of two males co-residing in their group. Social scenes included 271 agonistic events, 354 affiliative interactions, and 337 non-interacting control situations where two adult males were in close proximity of each other and did not interact via vocal exchange, facial expression, or tactile behavior.

Figure 2: Variation in visual attention to 3rd person social scenes as predicted from the looking probability model. Subjects generally attended more often to interactions than control scenes and to agonistic than affiliative interactions. Both the 2a) negative effect of rank difference (subject minus dominant stimulus) and the 2b) positive effect of affiliative relationship strength (subject to stimuli) were modulated independently by condition and were stronger for agonistic than for affiliative interactions. Subject dominance score 2c) did not affect visual attention. Bubbles are average probabilities per observed value of the predictor with size representing the number of observations, lines are model fitted estimates and shades their confident intervals (Tab. 1). Data came from 57 subjects and 962 social scenes.

Selective attention (looking probability)

In the looking probability model, the three-way interaction term between condition, rank difference, and maximum DDSI was not significant and subsequently was excluded from our first reduced model

(Tab. S1). To arrive at our final model, we further excluded the non-significant two-way interaction
maximum DDSI x rank difference (Tab. S2). The final full model differed significantly from the null
model including only the control variable conspicuity of the interaction and the random effects of
subject ID and group (Chi^2 = 34.02, df = 11, P<0.001).

Based on the parameter estimates and conditional on the interaction effects, subjects were
more likely to attend to interactions than to non-interactive controls (Tab. 1, Fig. 2). The higher above
the subject the higher-ranking of the stimulus individual was, i.e. the more negative the rank
difference, the more likely was the subject to pay attention to an agonistic interaction and the lower
below the subject the stimulus individuals were, the less likely the event was attended to. This effect
of dominance rank difference on overt attention differed between conditions. Specifically, the effect
of rank difference was stronger (more negative estimate) for non-interacting controls than for
affiliative interaction stimuli, but not stronger than for agonistic scenes.

Furthermore, stimulus valence modulated the effect of DDSI on visual attention. The closer
the subject was affiliated to the stimuli (the higher the maximum DDSI), the more likely they paid
attention which was an effect exacerbated for agonistic interactions, still pronounced for affiliative 3rd
person interactions and weakest for non-interacting control scenes.

Beyond the relational effects of rank difference, the individual dominance rank of the subject
did not affect its probability to monitor stimuli across conditions (see Fig. 3). Situations that were more
conspicuous were more likely to be looked at.

Table 1. *Model estimates for the reduced looking probability model, including as predictors only the*
interaction term between condition and rank difference between the subject and the dominant of the
stimulus individuals. The full model was significantly different from the null model including only
control factors (conspicuousness of the interaction), and random effects (Chi^2 = 34.02, df = 11,
P<0.001). Significant effects of test predictors are highlighted in bold. Significance of the different levels
of condition was assessed by releveling the intercept.

	Estimate	Std. Error	z value	Pr(>|z|)
Intercept	-1.629	0.185	-8.817	
Affiliation vs. Control	**0.887**	**0.226**	**3.925**	**<0.001**
Agonism vs. Control	**1.658**	**0.264**	**6.267**	**<0.001**
Agonism vs. Affiliation	**0.770**	**0.241**	**3.198**	**0.001**
Rank difference	**-0.614**	**0.191**	**-3.208**	**0.001**
Max DDSI	-0.043	0.221	-0.192	0.847
Subject rank	0.099	0.137	0.726	0.468
Conspicuousness	1.777	0.411	4.321	<0.001
Affiliation vs. Control x Rank difference	**0.426**	**0.208**	**2.048**	**0.041**
Agonism vs. Control x Rank difference	0.243	0.253	0.96	0.337

Agonism vs. Affiliation x Rank difference	-0.175	0.230	-0.764	0.445
Affiliation vs. Control x Max DDSI	0.126	0.216	0.581	0.561
Agonism vs. Control x Max DDSI	**0.64**	**0.279**	**2.293**	**0.022**
Agonism vs. Affiliation x Max DDSI	**0.515**	**0.269**	**1.914**	**0.056**

Figure 3: Sustained attention for 3rd person interactions was longer than for non-interaction social control scenes and longer for agonistic than for affiliative interactions (Tab. 2) with distributions of posterior estimates for the looking duration model shown in 3a). Violin plots are drawn for raw data 3b) and whisker plot shows model estimates with 95% credible interval.

Sustained attention (looking duration)

We used a Bayesian model to estimate the effects of the same predictor variables on the time spent looking at the social scene. The full model was a better fit for to the data than the null model (ELPD difference 16.5, SE 9.6). Comparison of expected log pointwise predictive density between the full looking duration and the associated null model with the conspicuousness control variable and the random effects of subject ID and group revealed a loo difference of 9.6 with a standard-deviation of 8.6. In the looking duration model, posterior probabilities for the three-way-interaction between rank difference, affiliative relationship strength and condition were centered around zero (Tab. S3). In a reduced model without the 3-way interaction term, we found no evidence supporting either of the

three 2-way interaction effects (all posterior probabilities centered around zero). The final model (Tab.

2) including only main effects provided clear evidence for sustained attention to vary with condition.

Males demonstrated longer attention for agonistic interactions between other males than for scenes

with two non-interacting stimulus males (posterior probabilities >0.95 for difference between control

and each of agonistic and affiliative scenes). Furthermore, there was moderate evidence for longer

looking time towards agonistic compared to affiliative interactions (posterior probability 0.84).

Looking time during agonistic situations was on average 1.7s longer compared to non-interacting

controls, and 0.9s longer than affiliation. Social relationships to the stimulus males and subject

dominance did not affect sustained attention.

Table 2. *Model estimates for the reduced looking duration model including only main effects; posterior*
means are reported with their standard errors and 95% credible intervals for the fixed effects.
Estimates are based on 4,000 posterior draws.

	Estimate	Est.Error	Q2.5	Q97.5
Intercept	0.171	0.206	-0.234	0.573
Affiliation vs. Control	**0.239**	**0.164**	**-0.086**	**0.565**
Agonism vs. Control	**0.380**	**0.215**	**-0.082**	**0.774**
Agonism vs. Affiliation	0.141	0.244	-0.362	0.592
Rank difference subject – highest	0.014	0.105	-0.198	0.218
Max DDSI subject – stimulus	0.035	0.109	-0.184	0.251
Subject rank	-0.042	0.121	-0.263	0.194
Log interaction time	**0.095**	**0.058**	**-0.016**	**0.210**
Conspicuousness	**0.359**	**0.178**	**-0.020**	**0.684**

Inter-individual differences

Random effects from the both models suggest that individuals do not systematically differ in attention

to just any type of social scene. Instead, significant individual differences were specific to and differed

between sustained attention to agonistic scenes (mean individual random effect for agonistic scenes

± stddev. = 0.0012 ± 0.1921) and affiliative scenes (0.0013 ± 0. 1743). Random effects were extracted

from a model with and are therefore controlled for subject dominance rank.

DISCUSSION

Our study makes a significant contribution to the development of real-life social neuroscience by

elucidating the neural mechanisms underlying social information integration. Social interactions are

inherently complex, requiring individuals to make predictions not only about their own actions but

also about those of others, informed by observations of third-party interactions[13]. Our findings reveal

that individuals allocate increased visual attention to third-party interactions, particularly those

involving agonistic versus affiliative behaviors, and that different dimensions of social relationships

between the subject and third parties independently modulate this attentional bias. Furthermore, we

observed significant individual differences in sustained attention to third-party interactions,

underscoring the importance of considering personal factors in understanding social cognition. These

results have important implications for our understanding of how the brain integrates social

information and how emotion regulation and adaptive benefits influence sustained attention.

3rd person interactions draw more attention that non-interacting social controls

Using as a control a social scene with two individuals in close proximity without an interaction, we

clearly demonstrated that both the probability of overt visual attention and the duration of sustained

visual attention were increased for 3rd person interactions. The three social conditions were matched

for salience by requiring the scene to have a clear starting point where two males arrived in close

proximity, i.e. all scenes were initially dynamic. Exceedingly dynamic scenes that involved quick

movements and/or vocalizations either in the agonistic or the affiliative context (e.g. ritualized male-

infant-male interactions)[62] were scored to be conspicuous which was a control variable in all models.

Thus, visual attention was not only guided by stimulus salience alone, but also by the mental

representation of its social relevance[6,63,64] and was sustained into a range where MEG studies suggest

different information integration events can be detected (1,100 and 2,000 ms)[65]. To our knowledge,

the only other study to demonstrate that 3rd person interactions attract more visual attention is a

three player experiment with rhesus macaques that identified neurons in the prefrontal cortex to be

differentially activated when two others interacted[22].

Attention guided by valence of others' interactions

Visual attention was further guided by interaction valence, adding ecological validity to previous

findings on selective attention for and differential processing of threat faces over photos of friendly

teeth-chattering[2,66]. Here the naturalism scale seems particularly relevant, because the two facial

expressions differ in important ways; teeth chattering is a repetitive and thus an inherently more

dynamic and also a more auditory display than a threat face and therefore may be more difficult to

identify from a still photograph. Yet even in our real-life scenes in proximity of the subject, monkeys

were more likely to attend to agonistic than affiliative interactions of others. .

Agonistic interactions are more relevant, because the subject could be drawn into the conflict

by redirected aggression[67] or be solicited to help one of the opponents[68,69]. The fact that watching

others fight increases rates of self-directed behaviors indicative of increased emotional arousal[70,71],

suggests that subjects do not only gather information about the state of the dominance relations of

others, but also about imminent threat to themselves. Others' affiliative interactions may also lead to

direct involvement of the subject, if they decide to interfere by joining or stopping the affiliation, but

this decision lies with the subject and not with a 3rd party and may therefore not require the same

level of attention as 3rd person agonism.

Despite the primacy of agonism, subjects still paid more often and more sustained attention

to 3rd person friendly interactions than non-interacting controls. Primates look towards the mother

(the main affiliate), if they hear an infant in distress from the other direction[72] and look longer at a

staged conflict, if it is between close affiliation partners compared to other dyads[73]. Selective attention

for 3rd person friendly interactions must underlie such findings, but had not been previously

demonstrated.

Two dimension of social relationships independently guide visual attention

Variation in imminent threat may also explain how the social relationships between the subject and

the stimulus individuals have moderated valence effects on selective attention. Since aggression flows

down the hierarchy in macaques[74], imminent threat increases the higher above the subject the

dominant stimulus male. This effect appears most pronounced in non-interactive control scenes,

because in controls, attention drops to zero for scenes that only have males subordinate to the

subject; at the largest rank differences (left of Fig. 2a) control scenes did not attract more attention

than agonistic or affiliative scenes. Attention to non-interacting control scenes with individuals out-

ranking the subject may serve in uncertainty-reduction[75]. Attention for conflicts was much higher than

for controls, even when both males were subordinate to the subject, so that the rank-difference had

a slightly weaker effect. Rank difference did not moderate visual attention to affiliative interaction

scenes which suggests that it is not the mere presence of a higher ranking individual that adds

relevancy, but the action of a dominant.

We expected that monitoring close affiliates may also contribute towards risk-avoidance,

given that the conflicts of closely bonded partners are the most likely to trigger risky interactions for

the observing party[76,77] which requires behavioral coordination between close affiliates[35,78]. Matching

these expectations, we found the attention-grabbing effect of affiliative relationships with stimuli to

vary with stimulus valence: the affiliative relations effect was most pronounced for conflicts and less

so for affiliation and control scenes. This highlights again the increased relevance of potentially risky

situations. In other real-life observational studies on captive populations, female rhesus macaques

and mandrills of both sexes also paid more attention to their closest partner and their kin,

respectively[34,79]. This is in contrast to a recent study on male rhesus macaques[33]; the grooming

relationship did not affect visual attention for pictures of a group mate, whereas the aggressive

relationship with a stimulus clearly correlated with attention for this male[33]. By moving from the

presentation of still photographs to real-life stimuli and from individual stimuli[79] to interactions, we

demonstrate that two dimensions of a subject's relationships to others have independent effects on

the allocation of visual attention which is more pronounced for agonism than affiliation of others.

Individual differences in social attention

The most controlled studies employing brain-imaging or electrophysiology often suffer from small

sample sizes prohibiting investigation of individual differences in social attention and underlying

processing[1]. Here, we find no evidence for individuals to generally differ in their social attention or for

social status. Instead, the 57 male Assamese macaques varied only in how they attended to agonistic

interactions and not to affiliation between others. These results match with a well-powered study on

rhesus macaques that found year-to-year repeatability in attention to threat over neutral faces but

not in attention to friendly teeth-chattering over neutral faces[80]. In contrast, differences between four

monkeys in the propensity to give aggression, grooming, and approaches over receiving these

behaviors were all associated to some degree with social attention in a distractor task[33]. Neither of

the rhesus macaque studies, however, controlled for dominance rank prior to extracting information

on individual differences. We can clearly show here that despite dominance changing throughout the

study period (Fig. 1), individuals maintain differences in how they attend to conflicts of others,

suggesting stable differences in underlying regulation via serotoninergic and oxytocinergic

pathways[32,33].

Conclusion – information integration for visual attention

Together, these results suggest that different relevancies of social information are integrated in

complex ways. Interaction versus non-interacting control, valence of the interaction, and two separate

relational aspects do not have simple additive effects on social attention. The social value of attention

is evident from increased attention to others' affiliation over non-interacting controls that is

unaffected by status differences or social bonds towards the stimuli. All other results can be explained

by risk-avoidance alone where information is not gathered for its own sake, but for immediate

preparation and activation in the face of danger. Thus, social attention serves at least two goals in

tandem which explains why it involves several brain areas in the dorsomedial prefrontal cortex, dorsal
anterior cingulate cortex, the ventral striatum and the amygdala perhaps with pronounced mixed
selectivity at the neuronal and network level[9,22,81,82].

ACKNOWLEDGEMENTS

We are grateful to the National Research Council of Thailand and the Department of National Parks,
Plant and Wildlife Conservation, Thailand for permission from to conduct research in a protected area
(permits: 0907.4/9313 April 2rd 2022, 0907.4/19458 September 3nd 2023). We thank the
superintendent of PKWS Wichanon Saenphala for his collaboration, Roger Mundry for statistical
advice, and the team of foreign and Thai field researchers that supported data collection. This study
was funded by the Deutsche Forschungsgesellschaft DFG- SFB 1528 Cognition of Interaction (project-
ID 454648639).

AUTHOR CONTRIBUTIONS

SP Investigation, Data curation, Formal analysis, Visualization, Writing - Original draft
SMa Resources, Project administration, Writing - Review & editing
SMe Resources, Project administration, Writing - Review & editing
JO Conceptualization, Funding acquisition, Data curation, Resources, Supervision, Writing - Review &
editing
OS Conceptualization, Funding acquisition, Data curation, Resources, Supervision, Writing - Original
draft

CONFLITCS OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY

Replication data are available at GRO.data https://doi.org/10.25625/XFN4B4

ETHICAL STATEMENT

The study was conducted with permission of relevant Thai authorities under a benefit sharing

agreement and was purely observational, completely non-invasive and did not involve food

provisioning or capture of the animals.

REFERENCES

1. Fan, S., Monte, O. D. & Chang, S. W. C. Levels of naturalism in social neuroscience research.

iScience **24**, (2021).

2. Gothard, K. M., Battaglia, F. P., Erickson, C. A., Spitler, K. M. & Amaral, D. G. Neural responses to

facial expression and face identity in the monkey amygdala. *J. Neurophysiol.* **97**, 1671–1683

(2007).

3. Kret, M. E., Muramatsu, A. & Matsuzawa, T. Emotion processing across and within species: A

comparison between humans (Homo sapiens) and chimpanzees (*Pan troglodytes*). *J. Comp.*

Psychol. **132**, 395–409 (2018).

4. Schacht, A. & Vrtička, P. Spatiotemporal pattern of appraising social and emotional relevance:

Evidence from event-related brain potentials. *Cogn. Affect. Behav. Neurosci.* **18**, 1172–1187

(2018).

5. Berg, D. J., Boehnke, S. E., Marino, R. A., Munoz, D. P. & Itti, L. Free viewing of dynamic stimuli by

humans and monkeys. *J. Vis.* **9**, 19 (2009).

6. Sliwa, J. & Freiwald, W. A. A dedicated network for social interaction processing in the primate

brain. *Science* **356**, 745–749 (2017).

7. Siebert, R. *et al.* A naturalistic dynamic monkey head avatar elicits species-typical reactions and

overcomes the uncanny valley. *eNeuro* **7**, (2020).

8. Wilson, V. A. D. *et al.* Macaque gaze responses to the primatar: a virtual macaque head for social

cognition research. *Front. Psychol.* **11**, (2020).

9. Haroush, K. & Williams, Z. M. Neuronal prediction of opponent's behavior during cooperative

social interchange in primates. *Cell* **160**, 1233–1245 (2015).

10. Dal Monte, O., Piva, M., Morris, J. A. & Chang, S. W. C. Live interaction distinctively shapes social

gaze dynamics in rhesus macaques. *J. Neurophysiol.* **116**, 1626–1643 (2016).

11. Seyfarth, R. M. & Cheney, D. L. Social cognition. *Anim. Behav.* **103**, 191–202 (2015).

12. Klein, J. T., Shepherd, S. V. & Platt, M. L. Social attention and the brain. *Curr. Biol. CB* **19**, R958-

962 (2009).

13. Freiwald, W. A. Social interaction networks in the primate brain. *Curr. Opin. Neurobiol.* **65**, 49–

58 (2020).

14. Andrews, M. W., Bhat, M. C. & Rosenblum, L. A. Acquisition and long-term patterning of joystick

selection of food-pellet vs social-video reward by Bonnet Macaques. *Learn. Motiv.* **26**, 370–379

(1995).

15. Deaner, R. O., Khera, A. V. & Platt, M. L. Monkeys pay per view: adaptive valuation of social

images by rhesus macaques. *Curr. Biol. CB* **15**, 543–548 (2005).

16. Lindsay, G. W. Attention in psychology, neuroscience, and machine learning. *Front. Comput.*

Neurosci. **14**, 29 (2020).

17. Lacreuse, A. *et al.* Testosterone may increase selective attention to threat in young male

macaques. *Horm. Behav.* **58**, 854–863 (2010).

18. McFarland, R. *et al.* Social interactions through the eyes of macaques and humans. *PLoS ONE* **8**,

e56437 (2013).

19. Teufel, C., Alexis, D. M., Clayton, N. S. & Davis, G. Mental-state attribution drives rapid, reflexive

gaze following. *Atten. Percept. Psychophys.* **72**, 695–705 (2010).

20. Adams, G. K., Ong, W. S., Pearson, J. M., Watson, K. K. & Platt, M. L. Neurons in primate

prefrontal cortex signal valuable social information during natural viewing. *Philos. Trans. R. Soc. B*

Biol. Sci. **376**, 20190666 (2021).

21. Dal Monte, O. *et al.* Widespread implementations of interactive social gaze neurons in the

primate prefrontal-amygdala networks. *Neuron* **110**, 2183-2197.e7 (2022).

22. Báez-Mendoza, R., Mastrobattista, E. P., Wang, A. J. & Williams, Z. M. Social agent identity cells

in the prefrontal cortex of interacting groups of primates. *Science* **374**, eabb4149 (2021).

23. Xing, F., Sheffield, A. G., Jadi, M. P., Chang, S. W. & Nandy, A. S. Dynamic modulation of social

gaze by sex and familiarity in marmoset dyads. *eLife* **14**, (2025).

24. Chance, M. R. A. Attention structure as the basis of primate rank orders. *Man* **2**, 503–518 (1967).

25. Shepherd, S. V., Deaner, R. O. & Platt, M. L. Social status gates social attention in monkeys. *Curr.*

Biol. **16**, R119–R120 (2006).

26. Cheng, J. T. *et al.* Eye gaze and visual attention as a window into leadership and followership: A

review of empirical insights and future directions. *Leadersh. Q.* **34**, 101654 (2023).

27. Capitanio, J. P. Sociability and responses to video playbacks in adult male rhesus monkeys

(Macaca mulatta). *Primates* **43**, 169–177 (2002).

28. Machado, C. J., Whitaker, A. M., Smith, S. E. P., Patterson, P. H. & Bauman, M. D. Maternal

Immune Activation in Nonhuman Primates Alters Social Attention in Juvenile Offspring. *Biol.*

Psychiatry **77**, 823–832 (2015).

29. Mosher, C. P., Zimmerman, P. E. & Gothard, K. M. Videos of conspecifics elicit interactive looking

patterns and facial expressions in monkeys. *Behav. Neurosci.* **125**, 639–652 (2011).

30. Watson, K. K., Ghodasra, J. H. & Platt, M. L. Serotonin transporter genotype modulates social

reward and punishment in rhesus macaques. *PLOS ONE* **4**, e4156 (2009).

31. Weinberg-Wolf, H. *et al.* The effects of 5-hydroxytryptophan on attention and central serotonin

neurochemistry in the rhesus macaque. *Neuropsychopharmacology* **43**, 1589–1598 (2018).

32. Weinberg-Wolf, H. & Chang, S. W. C. Differences in how macaques monitor others: Does

serotonin play a central role? *WIREs Cogn. Sci.* **10**, e1494 (2019).

33. Liu, S., Huang, J., Chen, S., Platt, M. L. & Yang, Y. Multi-dimensional social relationships shape

social attention in monkeys. *eLife* **14**, RP104460 (2025).

34. Schülke, O., Dumdey, N. & Ostner, J. Selective attention for affiliative and agonistic interactions

of dominants and close affiliates in macaques. *Sci. Rep.* **10**, 5962 (2020).

35. Watts, D. P., Colmenares, F. & Arnold, K. Redirection, consolation, and male policing: hor targets

of aggression interact with Bystanders. in *Natural conflict resolution* (eds Aureli, F. & de Waal, F.

B. M.) 281–304 (University of California Press, Berkley, 2000).

36. Schino, G. Grooming and agonistic support: a meta-analysis of primate reciprocal altruism.

Behav. Ecol. **18**, 115–120 (2007).

37. Ostner, J. & Schülke, O. The evolution of social bonds in primate males. *Behaviour* **151**, 871–906

(2014).

38. Mielke, A. *et al.* Bystanders intervene to impede grooming in Western chimpanzees and sooty

mangabeys. *R. Soc. Open Sci.* **4**, (2017).

39. Mielke, A., Bruchmann, C., Schülke, O. & Ostner, J. Grooming interventions in female rhesus

macaques as social niche construction. *Anim. Behav.* **173**, 105–114 (2021).

40. Ostner, J. & Schülke, O. Linking sociality to fitness in primates: a call for mechanisms. in

Advances in the Study of Behavior (eds Naguib, M. et al.) vol. 50 127–175 (Academic Press, 2018).

41. Borries, C., Larney, E., Kreetiyutanont, K. & Koenig, A. The diurnal primate community in a dry

evergreen forest in Phu Khieo Wildlife Sanctuary, Northeast Thailand. *Nat. Hist. Bull. Siam Soc.* **50**,

75–88 (2002).

42. Altmann, J. Observational study of behavior: sampling methods. *Behaviour* **49**, 227–266 (1974).

43. Martin, P. & Bateson, P. P. G. *Measuring Behaviour: An Introductory Guide*. (Cambridge Univ.

Press, Cambridge, 2006).

44. Neumann, C. & Fischer, J. Extending Bayesian Elo-rating to quantify the steepness of dominance

hierarchies. *Methods Ecol. Evol.* **14**, 669–682 (2023).

45. Stranks, J., Heistermann, M., Sangmaneedet, S., Schülke, O. & Ostner, J. The dynamics of

sociality and glucocorticoids in wild male Assamese macaques. *Horm. Behav.* **164**, 105604 (2024).

46. Neumann, C. EloRating.Bayes (GitHub Repository: https://github.com/gobbios/EloRating.Bayes).

(2024).

47. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing (2023).

48. Levy, E. *et al.* A comparison of dominance rank metrics reveals multiple competitive landscapes

in an animal society. *Proc. Biol. Sci.* **287**, 20201013 (2020).

49. De Moor, D., Roos, C., Ostner, J. & Schülke, O. Bonds of bros and brothers: Kinship and social

bonding in postdispersal male macaques. *Mol. Ecol.* **29**, 3346–3360 (2020).

50. Kulik, L. PhD Thesis: Development and consequences of social behavior in rhesus macaques

(*Macaca mulatta*). (Univ. Leipzig, Leipzig, 2015).

51. Baayen, R. H. *Analyzing linguistic data: A practical introduction to statistics using R*. vol. 49

(Cambridge Univ. Press, Cambridge, 2015).

52. McCullagh, P. *Generalized linear models*. (Routledge, London, 2019).

53. Schielzeth, H. & Forstmeier, W. Conclusions beyond support: overconfident estimates in mixed

models. *Behav. Ecol.* **20**, 416–420 (2009).

54. Barr, D. J. Random effects structure for testing interactions in linear mixed-effects models. *Front.*

Psychol. **4**, (2013).

55. Winters, S., Dubuc, C. & Higham, J. P. Perspectives: The looking time experimental paradigm in

studies of animal visual perception and cognition. *Ethology* **121**, 625–640 (2015).

56. Zuur, A. F. *A beginner's guide to generalized additive models with R*. (Highland Statistics Limited,

Newburgh, NY, 2012).

57. Bates, D., Maechler, M. & Bolker, B. M. *Lme4: Linear mixed-effects models using S4 classes. R*

Package (Version 0.999375-34). R-Project.Org/Package=lme4. (Institute for Statistics and

Mathematics, University of Economics and Business, Vienna, 2011).

58. Dobson, A. & Barnett, A. *An introduction to generalized linear models, Fourth Edition*. (Chapman

and Hall/CRC, 2018). doi:10.1201/9781315182780.

59. Bürkner, P.-C. Bayesian item response modeling in *R* with **brms** and *Stan*. *J. Stat. Softw.* **100**, 1–

54 (2021).

60. Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. Rank-normalization, folding,

and localization: An improved R² for assessing convergence of MCMC . *Bayesian Anal.* **16**, 667–

718 (2021).

61. Gabry, J., Simpson, D., Vehtari, A., Betancourt, M. & Gelman, A. Visualization in Bayesian

workflow. *J. R. Stat. Soc. Ser. A Stat. Soc.* **182**, 389–402 (2019).

62. Kalbitz, J., Schülke, O. & Ostner, J. Triadic male-infant-male interaction serves in bond

maintenance in male Assamese macaques. *PLoS ONE* **12**, e0183981 (2017).

63. Gobel, M. S., Tufft, M. R. A. & Richardson, D. C. Social beliefs and visual attention: how the social

relevance of a cue influences spatial orienting. *Cogn. Sci.* **42 Suppl 1**, 161–185 (2018).

64. Ninomiya, Taheri, Noritake, A., & Isoda, Masaki. Live agent preference and social action

monitoring in the macaque mid-superior temporal sulcus region. *Proc. Natl. Acad. Sci.* **118**,

e2109653118.

65. Ulloa, J. L., Puce, A., Hugueville, L. & George, N. Sustained neural activity to gaze and emotion

perception in dynamic social scenes. *Soc. Cogn. Affect. Neurosci.* **9**, 350–357 (2014).

66. Lacreuse, A., Schatz, K., Strazzullo, S., King, H. M. & Ready, R. Attentional biases and memory for

emotional stimuli in men and male rhesus monkeys. *Anim. Cogn.* **16**, 861–871 (2013).

67. Aureli, F. & Schaik, C. P. V. Post-conflict behaviour in long-tailed macaques (*Macaca fascicularis*

): II. Coping with the Uncertainty. *Ethology* **89**, 101–114 (1991).

68. Young, C., Majolo, B., Schülke, O. & Ostner, J. Male social bonds predict partner recruitment in

cooperative aggression in wild Barbary macaques. *Anim. Behav.* **80**, 675–682 (2014).

69. Perry, S., Barrett, H. C. & Manson, J. H. White-faced capuchin monkeys show triadic awareness

in their choice of allies. *Anim. Behav.* **67**, 165–170 (2004).

70. Whitehouse, J., Micheletta, J., Kaminski, J. & Waller, B. M. Macaques attend to scratching in

others. *Anim. Behav.* **122**, 169–175 (2016).

71. Paschek, N., Müller, N., Heistermann, M., Ostner, J. & Schülke, O. Subtypes of aggression and

their relation to anxiety in Barbary macaques. *Aggress. Behav.* **45**, 120–128 (2019).

72. Cheney, D. L. & Seyfarth, R. M. Vocal recognition in free-ranging vervet monkeys. *Anim. Behav.*

28, 362–367 (1980).

73. Whitehouse, J. & Meunier, H. An understanding of third-party friendships in a tolerant macaque.

Sci. Rep. **10**, 9777 (2020).

74. Deag, J. M. Aggression and submission in monkey societies. *Anim. Behav.* **25**, 465–474 (1977).

75. Iki, S. & Adachi, I. Monkeys increase scratching when encountering unexpected good fortune.

Sci. Rep. **15**, 8386 (2025).

76. Kalbitz, J., Ostner, J. & Schülke, O. Strong, equitable and long-term social bonds in the dispersing

sex in Assamese macaques. *Anim. Behav.* **113**, 13–22 (2016).

77. Schülke, O., Bhagavatula, J., Vigilant, L. & Ostner, J. Social bonds enhance reproductive success

in male macaques. *Curr. Biol.* **20**, 2207–2210 (2010).

78. Dunbar, R. I. M. & Shultz, S. Bondedness and sociality. *Behaviour* **147**, 775–803 (2010).

79. Schino, G. & Sciarretta, M. Patterns of social attention in mandrills, *Mandrillus sphinx*. *Int. J.*

Primatol. **37**, 752–761 (2016).

80. Diaz, A. A., Hernández-Pacheco, R. & Rosati, A. G. Individual differences in sociocognitive traits in

semi-free-ranging rhesus monkeys (Macaca mulatta). *Am. J. Primatol.* **87**, e23660 (2025).

81. Leopold, D. A. The big mixup: Neural representation during natural modes of primate visual

behavior. *Curr. Opin. Neurobiol.* **88**, 102913 (2024).

82. Chang, S. W. C., Gariépy, J.-F. & Platt, M. L. Neuronal reference frames for social decisions in

primate frontal cortex. *Nat. Neurosci.* **16**, 243–250 (2013).