A female-specific CB$_1$R-gated subcortical circuit orchestrates defensive homeostasis in risk assessment

Xue Liu[1,2,3,7], Hongren Huang[1,2,7], Xiaolong Feng[1,2], Xulin Li[1,2], Jialin Ye[1,2], Huan Yang[4], Jingjing Liu[1,5], Zicong Liang[3], Zhihan Guo[1,2], Ruyi Cai[6], Shangxuan Cai[6], Yulong Li[6], Zhaofa Wu[4], Liping Wang[1,2,*], Feng Wang[1,2,3,8,*]

[1]Shenzhen Key Laboratory of Neuropsychiatric Modulation, Shenzhen-Hong Kong Institute of Brain Science, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China

[2]Guangdong Provincial Key Laboratory of Brain Connectome and Behavior, the Brain Cognition and Brain Disease Institute, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China

[3]Department of Neuroscience, Faculty of Life and Health Sciences, Shenzhen University of Advanced Technology, Shenzhen 518107, China

[4]State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China.

[5]Shenzhen Key Laboratory of Viral Vectors for Biomedicine, Translational Research Center for the Nervous System, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, 518055, China.

[6]State Key Laboratory of Membrane Biology, Peking University School of Life Sciences, Beijing 100871, China

[7]These authors contributed equally

[8]Lead contact

*Correspondence: lp.wang@siat.ac.cn (L.W.), feng.wang@siat.ac.cn (F.W.)

Highlights

SC CB_1R^+ neurons encode visual survival threats and initiate risk assessment in both sexes

CB_1R^+ SC-LHb GABAergic circuit maintains defensive homeostasis in risk assessment in females

eCBs gate this circuit by disinhibiting presynaptic GABA release exclusively in females

Disruption of this mechanism female-specifically impairs risk assessment and stress coping

In Brief

Liu et al. identify a female-specific subcortical circuit in which eCB signaling, via CB_1R on GABAergic terminals in the SC, suppresses transmission to the LHb to maintain precise defensive homeostasis in risk assessment. These findings uncover the circuit and neuronal mechanisms that contribute to the sexually dimorphic maintenance of defensive homeostasis.

SUMMARY

Risk assessment in defensive behavior is an adaptive mechanism shaped by natural selection, enabling individuals to evaluate potential threats and thereby maintain defensive homeostasis. However, it remains unknown whether specific neural circuits maintain their behavioral homeostasis in a sex-specific manner. To address this, we investigated its neural basis by hierarchical behavior analysis framework with neural circuit dissection. In mice of both sexes, visual survival threats activated cannabinoid 1 receptor (CB_1R)-expressing neurons in the superior colliculus (SC) to initiate consistent risk assessment. Deletion of CB_1R in SC GABAergic neurons female-specifically impairs risk assessment by disinhibiting GABA release in SC-lateral habenula (LHb) projections, resulting in shortened risk assessment. This disruption furthermore increases the occurrence of abnormal spontaneous behavior following chronic stress exclusively in females. We identified a female-specific SC-LHb GABAergic circuit gated by CB_1R maintains defensive homeostasis in risk assessment. Our findings reveal how a conserved neuromodulatory system sex-specifically gates a subcortical circuit to orchestrate distinct survival strategies.

Key Words: Sex differences; Risk assessment; Defensive homeostasis; Cannabinoid 1 receptor; Superior colliculus; Lateral habenula

INTRODUCTION

Risk assessment during defensive behavior represents an evolutionarily conserved survival mechanism[1-3], refined through natural selection to optimize trade-offs[3,4] between immediate survival threats, such as predation risk, and long-term fitness benefits, such as reproductive success. However, how the brain maintains defensive homeostasis in a sex-specific manner remains unknown. Females exhibit heightened risk-assessment capacities[5,6] to prioritize survival under gestational and lactational constraints[7-9], manifesting as increased risk avoidance[10] and perinatal anxiety[11,12]. Conversely, males display stable risk-taking preferences shaped by selective pressures for resource acquisition[13] and intrasexual competition[14], where consistent behavioral strategies enhance reproductive success. These sexually dimorphic behavioral strategies, driven by asymmetric reproductive investment[15] are evident not only in defensive behaviors[16,17] but also across a range of innate behaviors including mating[18-20] and aggression[21-23] to sex recognition[24-26].

Population viability often depends more critically on female survival in many species[27,28], Robert Trivers' theory of parental investment[29] suggests that females exhibit risk-averse behaviors to protect their greater reproductive investments, a notion supported by our recent findings of sex-specific defensive strategies in mice facing looming threats[30], suggesting the existence of sexually dimorphic neural mechanisms coordinate by evolutionary pressures. Neural circuits modulating sex-dimorphic innate behaviors frequently involve subcortical hubs that pair conserved anatomy with sex-specific neuromodulation[19,31]. However, it remains

unclear whether subcortical structures mediate sex-dimorphic innate defensive behavior, particularly those requiring rapid risk assessment under threat.

The superior colliculus (SC), a conserved hub for processing life-threatening visual cues across vertebrates[32-34] , and the cannabinoid 1 receptor (CB$_1$R), known for rapid modulation of survival behaviors[35-37], emerge as promising candidates. Optogenetic activation of SC parvalbumin$^+$ neurons prolongs escape time in female mice[38], highlighting an evolutionarily driven sexual dimorphism in defensive control. Cannabinoid 1 receptor (CB$_1$R), an ancient fear modulator across vertebrates[39,40], demonstrates sex-specific characteristics in modulating learned fear and fear-related disorders[41,42], as well as its distribution in the brain[43,44]. It also contributes to the regulation of behaviors essential for survivals, including feeding[35], sleep[36], metabolism[37] and social behavior[45,46] as well as reproductive strategies, such as maternal behavior[47,48] and sexual behavior[49]. It is noteworthy that endocannabinoid (eCB), as an endogenous ligand of CB$_1$R, achieves synthesis and functional action within milliseconds[50], matching the temporal demands of life-or-death defensive responses mediated by subcortical circuits such as the SC. We therefore hypothesize that the CB$_1$R in SC may gate defensive homeostasis through a sex-specific subcortical fast regulatory mechanism.

Here, we resolve this gap by integrating the hierarchical behavior analysis framework (HBAF)[51] with neural circuit dissection. We demonstrate that risk assessment in response to visual threats is encoded by SC CB$_1$R$^+$ neurons in both sexes, while CB$_1$R in GABAergic SC-lateral habenula (LHb) projections is necessary

for female-specific defensive homeostasis in risk assessment. eCB release from the LHb activates the CB_1R on the terminal projecting from the SC, while the deletion of CB_1R in SC GABAergic neurons disinhibits presynaptic GABA release, resulting in a female- specific impairment in risk assessment. Disruption of CB_1R in SC GABAergic neurons induces female-specific stress vulnerability. Importantly, we identify a neural substrate for evolutionarily optimized sex differences in survival strategies, with female-specific defensive homeostasis in risk assessment directly linked to vertebrate-conserved eCB system. We demonstrate how this circuit integrates evolutionary constraints with real-time threat evaluation and establish a general mechanistic paradigm in which conserved neuromodulatory systems maintain sex-specific defensive homeostasis through subcortical disinhibition.

RESULTS

SC CB_1R^+ neurons are necessary for processing visual threats in both sexes

To access CB_1R^+ neurons genetically, we generated a CB_1R^{iCre} knock-in mouse line (Figure S1). *In situ* hybridization confirmed $>$ 80% of *CB_1R* mRNA co-localization with Cre protein in both sexes (Figure S1K), exceeded 90% in the superior colliculus (SC) (Figure S1I, K). Quantitative analysis revealed that CB_1R^+ cells make up more than 50% of the SC neuronal populations, independent of sex (Figure S2A-B). To investigate activation due to threats, we profiled c-Fos mRNA expression patterns following looming stimuli (Figure S3A). Looming significantly increased c-Fos mRNA expression and its co-localization with Cre proteins in the SC of both sexes (Figure S3B-E). Remarkably, $>$ 80% of c-Fos$^+$ neurons expressed CB_1R (Figure

S3F), indicating predominant engagement of CB_1R^+ neurons in threat processing. Additionally, we monitored real-time SC CB_1R^+ neuronal activity via calcium imaging during looming exposure (Figure 1A-B). By injecting either unilateral AAV2/9-EF1α-DIO-GCaMP6s or a control virus into the SC of CB_1R^{iCre} mice, we enabled cell-specific recordings (Figure 1A). During looming presentation, we observed strong calcium transients in SC CB_1R^+ neurons in both male and female mice (Figure 1C-D), indicating sex-conserved recruitment.

Given the heterogeneity of SC neurons[52], we molecularly characterized the CB_1R^+ populations using multiplex fluorescent *in situ* hybridization (FISH) (Table S2). Spatial mapping showed that $CB_1R\square$ neurons predominantly co-expressed glutamatergic markers (VGLUT2 (*Slc17a6*)$\square$: 28.86±4.83%; CaMKIIα$\square$: 26.72±6.62%) but rarely GABAergic markers (VGAT (*Slc32a1*)$\square$: 21.19±3.89%; Dlx5/6$\square$ <0.01%) (Figure 1E-I, S2C-D) with no observed sex differences.

CB_1R in SC GABAergic neurons governs female-specific defensive homeostasis in risk assessment

To dissociate the role of CB_1R in SC GABAergic and glutamatergic neurons, CB_1R-flox mice were bilaterally injected with AAV2/9-Dlx5/6-Cre-GFP (CB_1R cKO) or AAV2/9-Dlx5/6-GFP (CB_1R Ctrl) (Figure 2A) and AAV-CaMK$\square$α-Cre-GFP (CaMK$\square$α-cKO) or AAV-CaMK$\square$α-GFP (CaMK$\square$α-Ctrl) (Figure S4A). This enabled selective deletion of CB_1R in GABAergic (Dlx5/6$^+$) or glutamatergic (CaMK$\square$α$^+$) neurons (Figure 2B, S4B). Dlx5/6-driven AAVs targeted 94.38±1.94% of GABAergic neurons and 26.52±6.80% of glutamatergic neurons (Figure S5), while

CaMK□α-driven AAVs transduced approximately 71±5.33% of glutamatergic neurons (Figure S5D, S5E). Using a hierarchical behavioral analysis framework (HBAF)[51], we quantified defensive responses across groups (Figure 2C, S4C). In Dlx5/6 CB$_1$R cKO mice, most escaped to the refuge after looming stimuli (Figure 2D-2E), while some exhibited alternative strategies like freezing or vigilance (Figure S7, Table S3). Female Dlx5/6 CB$_1$R cKO mice exhibited accelerated defensive behavior, with reduced response latency and return time to the refuge (Figure 2E), unlike males (Figure 2D). Refuge duration remained consistent across groups (Figure 2D-E). In contrast, CaMK□α CB$_1$R cKO mice exhibited reduced defensive behavior in both sexes, as evidenced by fewer flight-to-refuge events (Figure S4D, E). Males additionally showed shortened refuge time, a phenotype absent in females (Figure S4D, E).

HBAF-derived 16-movements classification (Table S4, validated in 3D feature space, Figure 2F, 2H, 2I, S4F) identified four behavioral phases: exploring, escaping, hiding, and threat-informed exploring (Figure 2G, S4G). Movement fractions during pre-looming and hiding in the refuge stage were comparable across groups (Figure S6A-D). Female CB$_1$R cKO mice showed impaired risk assessment during escaping, with decreased looking up (Figure 2K), faster flight initiation probability (Figure 2M), as well as reduced nose elevation and time-to-maximum speed (Figure 2O, Movie S1, S2), while males showed no comparable changes (Figure 2J, 2L, 2N). These alterations, including a leftward shift in flight probability distribution (Figure 2M), indicate compromised risk evaluation in female CB$_1$R cKO mice. Post-refuge

exploration revealed increased sniffing in females (Figure S6F) not in males (Figure S6E), suggesting compensatory environmental sampling. Principal component analysis (PCA) of the 16-dimensional movement space detected no behavioral differences in the four phases (Figure S6G-H). Unlike Dlx5/6 CB_1R cKO mice, CaMK□α CB_1R cKO showed no sex-dependent phenotypes during escaping (Figure S4H-4K and Movie S3, S4, Table S5) and hiding (Figure S4L, S4M), likely due to the low specificity (~70%) of CaMKIIα for glutamatergic neurons. Subsequent experiments focused on SC GABAergic neurons, where CB_1R exhibits sexually dimorphic regulatory functions.

CB_1R^+ SC-LHb GABAergic circuit governs female-specific defensive homeostasis in risk assessment

To identify the specific circuits through which downstream SC $CB_1R^+Dlx5/6^+$ neurons mediate female risk assessment during defensive behavior, we conducted anterograde neuronal tracing from $CB_1R^+Dlx5/6^+$ neurons in the SC. This revealed over 30 projection targets without sex differences in fiber density (Figure S8). Parallel cell-specific labeling showed comparable CB_1R^+ $Dlx5/6^+$ neurons density between sexes (Figure S9), highlighting conserved circuit architecture. c-Fos screening revealed that LHb neurons were selectively activated by looming stimuli in females following SC CB_1R ablation in Dlx5/6 neurons (Figure 3A-3D). Given the LHb's established role in aversive processing[53-60] and eCB-mediated coping[61], we hypothesized that the CB_1R^+ SC-LHb GABAergic circuit regulates female risk assessment. Retrograde tracing confirmed dense SC CB_1R^+ $Dlx5/6^+$ axons innervation

of the LHb (Figure S10). Fiber photometry recordings used to detect eCB dynamics via the eCB sensor[62-64] (Figure 3E) showed looming stimuli increased eCB and calcium signals in the unilateral LHb of flight but not non-flight-responding mice (Figure 3F-3I), with no sex differences in signal dynamics (Figure 3J). These results suggest that looming induces elevated post-synaptic neuronal activity and eCB release in downstream nuclei in both sexes, specifically in the active flight response. LHb-specific CB_1R antagonism female-specifically impaired risk assessment (Figure 3K-3N), unlike interventions in reuniens thalamic nucleus or zona incerta (ZI) (Figure S11).

Optogenetic activation of CB_1R^+ SC-LHb GABAergic projections (Figure 4A-4D) disrupted female risk assessment (Figure 4F) but not males (Figure 4E), while chemogenetic inhibition (Figure 4G-4I) prolonged female risk assessment (Figure 4K), evidenced by increased latency to initiate flight and return to the refuge. These effects were not observed in CB_1R^+ GABAergic SC-ZI (Figure S12A-S12D) or SC-lateral posterior thalamus (LP) (Figure S12E-S12H) protections, confirming the circuit's necessity. Male defensive behavior (Figure 4J) and spontaneous behavior in both sexes remained unaffected (Figure 4L-P, Table S6). In summary, CB_1R in SC GABAergic projections to the LHb are necessary for female to exhibit full, deliberative risk assessment comparable to males. Their genetic deletion compromised defensive homeostasis in risk assessment, manifesting shortened and inadequate threat evaluation prior to flight.

CB_1R-mediated suppression of GABA release from SC terminal onto the LHb

underlies female-specific phenotypes

Given that presynaptic CB_1R inhibits neurotransmitter release[65,66], we measured GABA dynamics in the LHb using a genetically encoded fluorescent sensor for *in vivo* imaging (Figure 5A-5C). Female Dlx5/6 CB_1R cKO mice exhibited elevated GABA release during looming (Figure 5E), unlike males (Figure 5D), indicating disinhibited transmission from CB_1R^+ SC GABAergic projections to the LHb. The LHb expressed high levels of diacylglycerol lipase alpha (Dagla) (Figure 5G), essential for synthesizing 2-arachidonoylglycerol (2-AG)[67,68]. Locally deletion of *Dagla* by bilaterally injecting AAV-EF1α-Cre-EGFP virus into the LHb of Daglα-floxed mice (Figure 5F-5I) induced female-specific risk assessment deficits (Figure 5K), similar to those seen in Dlx5/6 CB_1R cKO mice, with no equivalent effects in males (Figure 5J). These findings suggest that 2-AG retrograde signaling in the LHb gates presynaptic GABA release from CB_1R^+ SC GABAergic terminals, thereby ensuring that females maintain defensive homeostasis in risk assessment comparable to males. Disruption of this circuit through CB_1R knockout, circuit modulation, or 2-AG depletion consistently impairs female-specific defensive homeostasis in threat evaluation.

The loss of CB_1R in SC GABAergic neurons underlies a female-specific vulnerability to stress

Considering that anxiety can accelerate defensive responses to looming stimuli[69], we assessed rat stress susceptibility in SC GABAergic CB_1R cKO and Ctrl mice (Figure 6A). Chronic stress resulted in comparable weight stability across groups (Figure 6B),

but female Dlx5/6 CB_1R cKO mice showed increased anxiety-like behavior, evidenced by reduced open-arms time in elevated plus maze post-stress (Figure 6C). Stress-induced behavioral reorganization included reduced locomotion (Figure 6G, 6H, 6J, 6K, Table S6), increased vertical exploration, such as rearing, with decreased nose height (Figure 6I, 6L), and decreased horizontal exploration, such as sniffing, with increased nose height (Figure 6I, 6L) in male and female CB_1R Ctrl and cKO groups. The stressor elicits a strategic behavioral shift from active exploration to defensive vigilance, manifesting as reduced locomotion and heightened vertical investigation, which potentiates visual information acquisition for environmental surveillance. Female CB_1R cKO mice exhibited a pathological triad post-stress, characterized by anhedonia-like reduced sniffing (10–20 min) (Figure 6L), hypervigilance associated with increased corner occupancy (40–50 min) (Figure 6K), and compulsion-like elevated grooming (50–60 min) (Figure 6L). This progression resembles human anxiety disorders (DSM-5), confirming female-specific vulnerability. Machine learning analysis quantified significant behavioral dissimilarity induced by stress (Figure S13, S14). Collectively, CB_1R deletion in SC Dlx5/6[+] neurons triggers a female-specific pathogenic cascade characterized by elevated GABA release in the LHb, impaired risk assessment, and heightened vulnerability to stress-induced psychopathology. This defines an evolutionarily conserved circuit that governs sex-specific survival plasticity through a vertebrate-conserved neuromodulatory system.

DISCUSSION

This study identifies a female-specific CB_1R-gated SC-LHb GABAergic circuit that orchestrates defensive homeostasis in risk assessment. We demonstrate how evolutionarily conserved eCB signaling enables subsecond-scale precision in risk assessment through sexually dimorphic modulation of a subcortical pathway. Our findings establish a neural mechanism whereby life-history trade-offs are implemented at the circuit level, prioritizing female survival through specialized neuromodulatory control of threat processing.

CB_1R as a sexually dimorphic neuromodulator in the homeostatic maintenance of risk assessment

CB_1R, a key component of the evolutionarily conserved eCB system across vertebrates[40], modulates various innate behaviors[70,71] and exhibits sexual differences in neural function[72]. Although associated with locomotor activity[63,73,74], nociceptive processing[75] and affective states[45,76], here we reveal that CB_1R in the SC are necessary for visual threat-induced defensive behavior in both sexes, expanding previous studies upon its role in defensive behavior[77,78]. Notably, although visual threats activate SC CB_1R^+ neurons universally, CB_1R-mediated modulation of the SC-LHb GABAergic circuit specifically governs risk computation in females. This dissociation between conserved neural activation and sex-specific neuromodulation[79] reveals a previously unrecognized organizational principle for implementing sexually dimorphic behaviors.

The functional compartmentalization of eCB signaling observed here extends previous demonstrations of nucleus-[80-82] and cell type-specific[83-88]. Specifically, a recent study revealed that 2-AG acts on neurons to enhance synaptic inhibition, while

anandamide targets astrocytes to promote synaptic potentiation[89]. This cell type- and endocannabinoid-specific signaling paradigm provides critical support that eCB signaling achieves functional specialization, including the sex-specific modulation of the SC-LHb GABAergic circuit observed here, by leveraging distinct molecular and cellular targets. Despite conserved anatomical projections between the SC and LHb, CB_1R affect GABAergic transmission within this pathway exclusively in females, facilitating more deliberate threat evaluation, an adaptation likely driven by evolutionary pressures that prioritize female survival. This specialization contrasts with its non-dimorphic role in glutamatergic circuits regulating escape behavior[34,90-93], whereas GABAergic neurons provide inhibitory control over these responses[94]. Such functional segregation may reflect evolutionary optimization where conserved subcortical pathways are overlaid with sex-specific modulatory controls to accommodate different survival strategies.

GABAergic neurons orchestrate behaviors by regulating excitation-inhibition balance[95], gating signal throughput[96], and facilitating disinhibition[97], which allows for precise control over the intensity, direction, and timing. Different GABAergic subtypes across brain regions encode specific fear behavior[98], regulate defensive outputs[90], and coordinate vigilance[99,100]. Critically, these GABAergic functions themselves are often sexually dimorphic[101], contributing to sex differences in defensive behavior. Thus, these findings suggest that CB_1R modulates defensive behavior through complementary mechanisms, orchestrating behavioral initiation via glutamatergic regulation in both sexes, while maintaining defensive homeostasis

through sexually dimorphic modulation via GABAergic circuits. This dual arrangement may support immediate individual survival and long-term species-level reproductive strategies.

This functional dimorphism mediated via CB_1R parallels mechanisms such as tachykinin receptor 1 activation in preoptic area of the hypothalamus neurons, which drives male-specific mating behavior[19]. Both findings highlight how evolutionarily distinct neuromodulatory systems, the endocannabinoid and substance P systems, can be deployed within specialized circuits to promote sex-specific behaviors. We propose that such dimorphism may originate from early developmental programming, as eCB signaling has been shown to regulate microglia-dependent synaptic pruning and astrogenesis, thereby shaping sex differences in juvenile social behavior[102]. Further supporting a developmental basis, endocrine signals have been shown to influence the connectivity and gene expression profiles of hormone-responsive neurons[31], creating latent sex biases that are selectively activated by specific threat contexts in adulthood.

Subcortical GABAergic circuit architecture governs sex-specific threats evaluation

Subcortical circuit governing sex-specific survival strategies exhibit conserved anatomy alongside different neuromodulation. While sexually dimorphic neural architectures are well-documented in hypothalamic[20,103] and limbic hubs[104], manifesting as sex differences in gene expression[105], synaptic activity[106,107], and cell density[103], the SC-LHb circuit demonstrates functional sexual dimorphism without structural differences, as previously reported[52] and confirmed here. The SC, a conserved detector of survival-relevant visual threats across vertebrates, processes

threat information through glutamatergic neurons[34,90-93,108] and GABAergic neurons[94]. Our findings reveal a specific sexually dimorphic mechanism within the CB_1R^+ SC-LHb GABAergic circuit. We show that female-specific activation of the LHb depends on CB_1R-mediated retrograde 2-AG signaling that selectively enhances presynaptic inhibition of GABA release. This neuromodulatory specialization aligns with life-history trade-offs that prioritize female survival due to greater reproductive investment.

The LHb is broad involvement in encoding aversion, anxiety, and stress[54,109,110], positions it as a strategic hub for integrating female-specific survival behaviors. Our work identifies that the LHb integrates female-specific survival behaviors. Its broad involvement across these behaviors suggests that the LHb may function as a central integrator of survival-stress information, with sex differences in connectivity[53,111] potentially underlying differential susceptibility to stress-related disorders. This integrative function may also account for the heightened threat reactivity often seen in individuals with emotional disorders. Beyond risk assessment, the LHb also contributes to risk preference evaluation[112] and parental behavior[113,114] via distinct afferent pathways, highlighting its broader function in balancing survival-reproduction trade-offs. The engagement of these functions in the LHb highlights its central role in female-specific adaptive behaviors, with eCB signaling providing essential neuromodulatory flexibility to dynamically prioritize competing demands, based on internal and external contexts.

More broadly, our results align with growing evidence that subcortical structures,

including the BNST[19,24,115] and VMH[18,21,31], encode sexually dimorphic behaviors in a cell-type-specific mechanisms involving neuronal populations expressing Tac1[19], serotonin[115], aromatase[24], or progesterone-receptor[18,21]. This supports a unifying paradigm in which evolutionarily conserved subcortical circuits use genetically specified neuronal subpopulations to regulate innate behaviors in a sex-dependent manner. Such functional specialization, often mediated by inhibitory signaling, is conserved across species[116-119] while allowing for species-specific adaptations. Importantly, dysregulation of this circuits can disrupt defensive homeostasis, as evidenced by the LHb's involvement in female-biased disorders such as PTSD[120,121] and postpartum anxiety[122,123], highlighting the clinical relevance of these evolutionarily tuned circuits.

Defensive homeostasis, evolutionary trade-offs and therapeutic translation for sex-biased disorders

The behavioral homeostasis theory suggests that the evolutionary function of defensive behavior is to rapidly maximize an organism's overall readiness to cope with new survival threats and to minimize unnecessary energy expenditure[124-126]. This concept aligns with recently identified homeostatic systems regulating fundamental needs such as sleep[127] and social interaction[128]. Building on our previous findings that complete defensive behavior includes not only threat detection and escape to shelter, but also the critical process of re-emergence for risk reassessment and environmental exploration[30], we introduce the concept of defensive homeostasis. This homeostatic process involves the entire sequence of responses to life-threatening stimuli,

emphasizing the optimization of an organism's ability to rapidly evaluate whether external stimuli present imminent survival risks. Notably, such homeostatic systems often exhibit inherent sexual dimorphism, likely rooted in genetic and epigenetic mechanisms—as seen in the sex-specific imprinting of hypothalamic genes[129] which may predispose differential defensive strategies between males and females. Although both sexes showed consistent escape responses to immediate threats, females displayed quicker re-exploration and more thorough risk reassessment after reaching safety[30]. This enhanced defensive homeostasis in females can be understood through the life-history trade-off theory proposed by William Williams in 1957. Due to asymmetric parental investment, with females bearing the costs of gestation and lactation, female behavioral strategies prioritize survival through more cautious and comprehensive risk assessment. We propose that the comparable assessment duration, between sexes served in females as a proactive phase for gathering and processing more detailed information, thereby \ qualitatively enhancing their defensive homeostasis. The reduction of this duration following CB_1R deletion in SC GABAergic neurons highlights the role of CB_1R in facilitating the deliberative processing characteristic of female defensive strategy.

Chronic stress exposure further highlights these evolutionary trade-offs, where CB_1R dysfunction transforms adaptive vigilance into female-specific psychopathology. Consistent with the role of CB_1R in managing acute[130] and chronic stress[45,131], Dlx5/6 CB_1R-KO mice exhibit maladaptive behaviors exclusively in females, including reduced exploration, heightened thigmotaxis, and compulsive

grooming. These phenotypes closely align with the core symptoms of anxiety and depression as outlined in the DSM-5. This maladaptation may arise from disrupted threat calibration within the CB_1R^+ SC-LHb GABAergic circuit. From an evolutionary perspective, this vulnerability reflects the costs of optimizing female survival strategies. Females with CB_1R dysfunction may overestimate risks, a strategy that could theoretically enhance offspring protection during maternal investment[132] but imposing significant energetic costs. Conversely, males maintain more accurate defensive homeostasis in risk assessment, supporting competitive resource acquisition and reproductive success[13]. These sexually dimorphic behavioral strategies collectively enhance species survival and reproductive adaptability. Clinically, aberrant risk calibration is mechanistically linked to disorders predominantly affecting females, such as pathological hypervigilance in post-traumatic stress disorder (PTSD)[133], which correlated with habenular hyperactivity[134,135]. Thus, enhancing CB_1R signaling in this circuit presents a promising therapeutic approach for addressing sex-specific psychopathologies.

Limitations of the study

The Dlx5/6 promotor exhibits approximately 20% glutamatergic specificity, yet fails to fully resolve the heterogeneity among GABAergic subpopulations in the SC. Future studies should employ intersectional genetics, such as Cre/Flp-dependent transgenic mice with viral tools, for subpopulation-specific manipulation. Additionally, the continuous stress modeling protocol limited the dissection of stage-specific adaptations. Future designs should incorporate *in vivo* recordings during chronic

stress exposure to characterize temporal dynamics. Finally, therapeutic translation warrants testing CB_1R-positive allosteric modulators[136] for rescuing female-specific deficits.

FIGURES and FIGURE LEGENDS

Figure 1. SC CB$_1$R$^+$ neurons are activated during visually evoked defensive behavior

(A) Left: Schematic representing viral injection of AAV2/9-EF1α-DIO-GCaMP6s or AAV2/9-EF1α-DIO-EGFP into the unilateral SC of CB$_1$R-iCre-EGFP mice. Right: Representative image showing the recording site within the SC. Scale bar, 500 μm and 50 μm (enlarged view).

(B)   Experimental timeline and schematic of the looming stimuli paradigm used

to record calcium signals via fiber photometry in a refuge-containing open field apparatus.

(C and D) Left: Average calcium transients for the entire test group in males (C) and females (D). Shaded areas represent error bars, while the gray shadow indicates the looming stimuli event. Right: Plots showing calcium transients during looming stimuli in males (C) and females (D). Unpaired Student's t test with Mann-Whitney test was performed.

(E) Representative sections showing gene expression in the superior colliculus (SC). *Cnr1*, cannabinoid receptor 1; *Slc32a1*, solute carrier family 32 (GABA vesicular transporter), member 1; *Slc17a6*, solute carrier family 17 (sodium-dependent inorganic phosphate cotransporter), member 6; *CaMKIIα*, calcium/calmodulin-dependent protein kinase II alpha; *Dlx5*, distal-less homeobox 5; *Dlx6*, distal-less homeobox 6, displayed from left to right.

(F) Neural subtypes cell plots color-coded by gene expression. magenta, *Cnr1*; red, *Slc32a1*; green, *Slc17a6*; cyan, *CaMKIIα*; yellow, *Dlx5*; and gray, *Dlx6*.

(G) Multiplexed fluorescent *in situ* hybridization of *Cnr1* with *Slc32a1, Slc17a6, CaMKIIα, Dlx5, Dlx6* (from left to right) in the SC within a single representative section. Scale bar, 500 μm and 50 μm (enlarged view).

(H and I) Quantification of neural subtypes (I) and double-positive types, expressed as a percentage of DAPI. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed.

Data are presented as mean ± SEM. n=7 male Ctrl group (C), 6 male GCaMP6s group;

7 female Ctrl group, 6 female GCaMP6s group (D); 13 sections from 3 males and 3

females (H, I). ns (not significant), *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$

0.0001. For detailed p values, refer to Table S1.

Figure 2. Deletion of CB$_1$R in SC Dlx5/6$^+$ neurons female-specifically impairs defensive homeostasis in risk assessment

(A) Schematic representation of viral injection of AAV-Dlx5/6-GFP or AAV-Dlx5/6-Cre-GFP into the bilateral SC of CB$_1$R-floxed mice.

(B) Left: Representative images showing AAV-Dlx5/6-GFP (top) and

AAV-Dlx5/6-Cre-GFP (bottom) expression in the SC, alongside CB_1R mRNA (red) and Dlx5/6$^+$ cells bodies (green) in CB_1R Ctrl and CB_1R cKO group. Scale bar, 500 μm and 20 μm (enlarged view). White arrows indicate CB_1R mRNA fluorescent signals colocalized with GFP positive signals. Right: Quantification of CB_1R^+ cells in Dlx5/6$^+$ neurons in CB_1R Ctrl and CB_1R cKO groups for both sexes.

(C) Experimental timeline and schematic paradigm of virus injection and the looming test in a refuge-containing apparatus.

(D and E) Defensive strategy (left) and flight-to-refuge behaviors (from left to right: response latency, return time, time spent in the refuge) in male (D) and female mice (E) across CB_1R Ctrl and CB_1R cKO group.

(F) Sixteen movement types during defensive behavior, represented by corresponding colors.

(G) Schematic paradigm of the different stages of defensive behavior: exploring, escaping, hiding and threat-informed exploring.

(H and I) Ethograms showing the 16 movement types in males (H) and females (I) across groups (CB_1R Ctrl group on the left, CB_1R cKO group on the right). Scale bar, 1 s.

(J-K) Trajectories (left and middle) and quantification of fractions (right) of mice during the escaping stage (from the onset of looming to the end of flight) in males (J) and females (K) across CB_1R Ctrl and CB_1R cKO groups. Symbols indicate mouse locations: five-pointed star for looming onset, triangle for flight initiation, and circle for flight termination.

(L and M) Distribution and positions of mice during flight, looking up, and freezing from 5 s pre-looming to 15 s post-looming in males (L) and females (M) across groups (CB_1R Ctrl on the left, CB_1R cKO on the right).

(N and O) Left: Real-time speed and nose height from 5 s pre-looming to 15 s post-looming in males (N) and females (O) across groups. Right: Quantification of the time taken to reach maximum speed and mean nose height during looming stimuli in males (N) and females (O).

Data are presented as mean ± SEM (B, D, E, N, O) or mean (J-O). n=5 male CB_1R Ctrl, 4 male CB_1R cKO, 4 female CB_1R Ctrl, 5 female CB_1R cKO (B); 28 male CB_1R Ctrl, 24 male CB_1R cKO (D), 20 female CB_1R Ctrl, 21 female CB_1R cKO (E); 22 male CB_1R Ctrl and CB_1R cKO, 16 female CB_1R Ctrl and CB_1R cKO (H-O). Two-Way ANOVA with Sidak's multiple comparisons post hoc test (B, J-K) and Unpaired Student's t test with Mann-Whitney test (D, E, N, O) were performed. ns (not significant), *$p < 0.05$, ***$p < 0.001$. For detailed p values, refer to Table S1.

Figure 3. LHb serves as a key downstream target of the SC that mediates female-specific modulation of defensive homeostasis in risk assessment

(A) Left: Schematic representation of viral injections of AAV-Dlx5/6-GFP or AAV-Dlx5/6-Cre-GFP into the bilateral SC of CB$_1$R-floxed mice. Right: Representative images showing Dlx5/6$^+$ GFP signals expression in the SC. Scale bar, 500 μm.

(B) Experimental timeline and schematic paradigm of the looming test.

(C) Representative images of c-Fos expression in the LHb in response to looming stimuli in female CB_1R Ctrl and CB_1R cKO mice. Scale bar, 200 μm.

(D) Quantification of c-Fos expression in different brain regions across sexes.

Abbreviations: LHb, lateral habenula; LP, lateral posterior thalamus; ZI, zona incerta, PVT, paraventricular thalamic nucleus; Re, reuniens thalamic nucleus.

(E) Experimental timeline (top), schematic paradigm (middle) and representative image (bottom) of the viral infection in LHb neurons. Scale bar, 500 μm

(F, G) Heatmap visualization of endocannabinoid (eCB) and calcium (lower panel) dynamics (left), signals (middle), and peak signals (right) in male (F) and female (G) mice during flight response to looming stimuli. Shaded areas around the means indicate error bars; the gray shadow marks the looming stimuli event.

(H, I) eCB and calcium fluorescence changes in male (F) and female (G) mice during non-flight response to looming stimuli.

(J) Statistical graph showing the area under the curve (AUC) of eCB and calcium signaling in male and female mice with flight (left) or non-flight (right) response under looming stimulus.

(K) Timeline and schematic representation of pharmacological experiment during the looming test.

(L) Representative image showing the placement of track cannula in the bilateral LHb. Scale bar, 500 μm.

(M and N) Metrics of defensive behavior (from left to right: response latency, return time, time spend in the refuge and freezing time in the refuge after looming stimuli)

for male (M) and female mice (N) in vehicle (Veh) and NESS 0327 (NESS) groups.

Data are presented as mean $\pm$ SEM. n=5 per group (D), 4 per group (H), 5 per group

(F-J), 8 male Veh, 10 male NESS (K), 9 female Veh, 10 female NESS (L). Unpaired

Student's t test with Mann-Whitney test (D, K, L) and two-way ANOVA with Sidak's

multiple comparisons post hoc test (H) were performed. ns (not significant), *p < 0.05,

**p < 0.01, ***p < 0.001, ****p < 0.0001. For detailed p values, refer to Table S1.

Figure 4. Modulating SC-LHb $CB_1R^+Dlx5/6^+$ projections female-specifically affects defensive homeostasis in risk assessment

(A) Schematic representation of unilateral blue light activation targeting CB_1R^+ SC-LHb Dlx5/6$^+$ projections.

(B) Protocol for optogenetic stimulation.

(C) Representative image showing ChR2 virus expression in the SC of a CB$_1$R-iCre-EGFP mouse and the location of the fiber track (blue, DAPI; red, ChR2-mCherry; dotted border line, fiber track). Scale bars, 500 μm.

(D) Left: Schematics of patch-clamp slice recording during optogenetic stimulation of SC CB$_1$R$^+$Dlx5/6$^+$ neurons. Right: Light-pulse train triggered spikes (red) in ChR2-mCherry–positive neurons.

(E and F) Metrics of defensive behavior in males (E) and females (F) (from left to right: response latency, return time and time spend in the refuge after looming) under optogenetic activation of CB$_1$R$^+$ SC-LHb Dlx5/6$^+$ terminals.

(G) Schematic representation of bilateral chemogenetic inhibition targeting CB$_1$R$^+$ SC-LHb Dlx5/6$^+$ projections.

(H) Representative image showing CB$_1$R$^+$Dlx5/6$^+$ neurons in the SC and the fiber track location (blue, DAPI; red, ChR2-mCherry; dotted border line, fiber track). Scale bars, 500 μm.

(I) Defensive strategy in males (left) and females (right).

(J and K) Metrics of defensive behavior (from left to right: response latency, return time, time spent in the refuge, and freezing time in the refuge after looming stimuli) in male (J) and female mice (K) in mCherry and hM4Di group.

(L) Ethograms depicting 14 movements in males (top) and females (bottom) across different groups (mCherry and hM4Di group) in spontaneous behavior. Each color corresponds to a specific movement.

(M, N) A random forest classifier trained on 14-dimensional movement fractions to

predict group labels (left), and PCA visualization of behavioral features (right) for male (m) and female (n) in the mCherry and hM4Di groups in spontaneous behavior. (O, P) Comparison of the movement fraction, the sniffing to grooming ratio between male (L) and female (M) mice in the mCherry and hM4Di groups.

Data are presented as mean ± SEM. n=20 male mCherry, 18 male ChR2 (E), 18 female mCherry and ChR2 (F),11 male mCherry and hM4Di (J), 9 female mCherry, 12 female hM4Di (K). n=12 male mCherry group, 14 male hM4Di group (O); 11 female mCherry group, 13 female hM4Di group (P). Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed (O, P). Unpaired Student's t test with Mann-Whitney test was performed. ns (not significant), $*p < 0.05$, $**p < 0.01$. For detailed p values, refer to Table S1.

Figure 5. SC-LHb GABA transmission is required for risk assessment in a female-specific manner

(A) Schematic representation of viral injection: AAV-Dlx5/6-GFP or AAV-Dlx5/6-Cre-GFP into the bilateral SC (left) and AAV-EF1α-iGABASnFR into the unilateral LHb (right) of CB₁R-floxed mice.

(B) Representative image showing Dlx5/6+ GFP signals in the SC and iGABASnFR signals in the LHb, and the position of the fiber track (blue, DAPI; green, Dlx5/6+ GFP or iGABASnFR signals; dotted border line, fiber track). Scale bars, 500 μm.

(C) Apparatus for looming test and schematic of the recording system to measure GABA signals using fiber photometry during looming stimuli.

(D and E) Heatmap of iGABASnFR fluorescence changes (left), signals (middle), and

peak signals (right) in male (D) and female (E) mice. Shaded areas around the means indicate error bars; the gray shadow marks the looming stimuli event.

(F) Schematic representation of viral injection: AAV-EF1α-EGFP or AAV-EF1α-Cre-EGFP into the unilateral LHb (left) of Daglα-floxed mice.

(G) Representative images showing AAV-EF1α-EGFP (left) and AAV-EF1α-Cre-EGFP (right) expression in the LHb (red, Daglα mRNA; green, EGFP signals). Scale bar, 200 μm and 20 μm (enlarged view).

(H) Percentage of DAGLα^{+} cells in LHb neurons in DAGLα Ctrl and DAGLα cKO group.

(I) Schematic paradigm for the looming test.

(J and K) Metrics of defensive behavior (from left to right: response latency, return time, time spent in the refuge and freezing time in the refuge after looming stimuli) of male (J) and female mice (K) in DAGLα Ctrl and DAGLα cKO groups.

Data are presented as mean ± SEM. n = 8 male CB$_1$R Ctrl, 9 male CB$_1$R cKO (D), 14 female CB$_1$R Ctrl, 8 female CB$_1$R cKO (E), 4 per group (H), 16 male DAGLα Ctrl and DAGLα cKO (J), 18 female DAGLα Ctrl, 19 female DAGLα cKO (K). Unpaired Student's t test with Mann-Whitney test. ns (not significant), *p < 0.05. For detailed p-values, refer to Table S1.

Figure 6. CB₁R deletion in SC GABAergic neurons increases abnormal behavior patterns specifically in females after stress

(A) Schematic representation of viral injection of AAV-Dlx5/6-GFP or AAV-Dlx5/6-Cre-GFP into the bilateral SC of CB₁R-floxed mice and the timeline of behavioral teats conducted pre- and post- chronic stress.

(B, C) Changes in body weight and percentage of time spent in open arms during elevated plus maze test of mice pre- and post- chronic stress across different groups.

#*p* < 0.05.

(D) Trajectories showing speed during 60-min of open field activity.

(E) Representative trajectories illustrating different movements in the open field apparatus. Each of the 14 colors corresponds to a specific movement type.

(F) Density map showing the spatial distribution of dorsal points during 60-min of open field (left) and representative trajectories in the center, periphery, and corner (right) of the apparatus.

(G, J) Travel distance per minute throughout the experiment for males (G) and females (J).

(I, L) Comparison of the fraction (10-min epochs) between male (I) and female (L) mice in the CB_1R Ctrl and CB_1R cKO groups.

(H, K) Pie charts showing relative percentage of time spent in different areas for males (H) and females (K). Inner and outer rings represent pre- and post-stress percentages, respectively, while left and right halves correspond to CB_1R Ctrl (left) and CB_1R cKO (right) groups.

(M, N) Nose height during different movements in the 60-min open field apparatus (bottom) for males (M) and females (N).

Data are presented as mean ± SEM (B, C, G, J, M, N) and as mean (I, L, H, K). n=18 pre-stress, 17 post-stress male CB_1R Ctrl; 18 pre-stress, 16 post-stress male CB_1R cKO; 17 pre-stress, 20 post-stress female CB_1R Ctrl; 17 pre-stress, 18 post-stress female CB_1R cKO. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed. #*p* < 0.05; ##*p* < 0.01; ###*p* < 0.001; ####*p* < 0.001 pre-stress

versus post-stress; *p* < 0.05; **p* < 0.01 post-stress CB_1R Ctrl versus CB_1R cKO For

detailed *p* values, refer to Table S1.

STAR ★ METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**

- **RESOURCE AVAILABILITY**

  ○ Lead contact

  ○ Materials availability

  ○ Data and code availability

- **EXPERIMENTAL MODEL AND PARTICIPANT DETAILS**

  ○ Animals

  ○ Viruses

- **METHOD DETAILS**

  ○ Experimental design and randomization

  ○ Stereotaxic surgeries

  ○ Cannula implantation

  ○ Drugs delivery

  ○ Histology

  ○ Behavioral assays

  ○ 3D Behavioral data processing

  ○ Chronic rat stress model

  ○ Patch-clamp electrophysiology

  ○ Optical fiber implantation

- ○ Optogenetics

- ○ Fiber photometry

- **QUANTIFICATION AND STATISTICAL ANALYSIS**

  - ○ Statistics

ACKNOWLEDGMENTS

This work was supported by the Brain Science and Brain-like Intelligence Technology-National Science and Technology Major Project (2022ZD0208300 to F.W.), the National Natural Science Foundation of China (32371062 to F.W., 31630031 and 31930047 to L.W, Research Fund for International Senior Scientists T2250710685), the Guangdong Provincial Key Laboratory of Brain Connectome and Behavior (2023B1212060055 to L.W.), the Guangdong Basic and Applied Basic Research Foundation (2022A1515110120 to X.L.), the Shenzhen Key Basic Research Project (JCYJ20220818100805013 to F.W.), and the Financial Support for Outstanding Talents Training Fund in Shenzhen (to L.W.). Behavioral experiments focusing on chronic stress was conducted at the Shenzhen Brain Science Infrastructure. We are grateful to Dr. Xuemei Liu for conducting the preliminary experiments. We would like to express our sincere gratitude to the members of Feng Wang's lab for their helpful comments and kind suggestions regarding the manuscript.

AUTHOR CONTRIBUTIONS

F.W. and L.W. conceived the project, designed the experiments and proposed the concept. X.Liu and H.H. conducted the majority of experiments and analyzed the corresponding data. X.F. and X.Li conducted the optogenetic, morphological, and

pharmacological experiments and analyzed the corresponding data. X.Liu and Z.G. conducted the rat stress experiments. J.Y., H.H., X.Liu, and Z.L. contributed to 3D behavioral analysis. R.C., S.C., and Y.L. developed and provided eCB3.0 sensor. H.Y. conducted the RCamp3 and eCB3.0 sensor recordings under the supervision of Z.W.. F.W., X.Liu, H.H., and L.W. wrote the manuscript with input from all authors. L.W. and F.W. supported all aspects of this project.

DECLARATION OF INTERESTS

The authors declare no competing interests.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

During the preparation of this work, the AI tool DeepSeek and Monica were used to assist with language editing. Upon its use, the authors thoroughly reviewed and further edited the entire text. The authors take full responsibility for the content of the publication.

REFERENCES

1. Tovote, P., Fadok, J.P., and Lüthi, A. (2015). Neuronal circuits for fear and anxiety. Nature reviews. Neuroscience *16*, 317-331. 10.1038/nrn3945.

2. Motta, S.C., Goto, M., Gouveia, F.V., Baldo, M.V., Canteras, N.S., and Swanson, L.W. (2009). Dissecting the brain's fear system reveals the hypothalamus is critical for responding in subordinate conspecific intruders. Proceedings of the National Academy of Sciences of the United States of America *106*, 4870-4875. 10.1073/pnas.0900939106.

3. Blanchard, D.C., Griebel, G., Pobbe, R., and Blanchard, R.J. (2011). Risk assessment as an evolved threat detection and analysis process. Neuroscience and biobehavioral reviews *35*, 991-998. 10.1016/j.neubiorev.2010.10.016.

4. Tseng, Y.T., Schaefke, B., Wei, P., and Wang, L. (2023). Defensive responses: behaviour, the brain and the body. Nature reviews. Neuroscience *24*, 655-671. 10.1038/s41583-023-00736-3.

5. Blank, D.A. (2019). Diversity of risk assessment patterns in antipredator behavior of goitered gazelles. Behavioural processes *158*, 211-218. 10.1016/j.beproc.2018.12.008.

6. Jolles, J.W., Boogert, N.J., and van den Bos, R. (2015). Sex differences in risk-taking and associative learning in rats. Royal Society open science *2*, 150485. 10.1098/rsos.150485.

7. Liker, A., Freckleton, R.P., Remeš, V., and Székely, T. (2015). Sex differences in parental care: Gametic investment, sexual selection, and social environment. Evolution; international journal of organic evolution *69*, 2862-2875. 10.1111/evo.12786.

8. Janicke, T., Häderer, I.K., Lajeunesse, M.J., and Anthes, N. (2016). Darwinian sex roles confirmed across the animal kingdom. Science advances *2*, e1500983. 10.1126/sciadv.1500983.

9. Kokko, H., and Jennions, M.D. (2008). Parental investment, sexual selection and sex ratios. Journal of evolutionary biology *21*, 919-948. 10.1111/j.1420-9101.2008.01540.x.

10. Harris, C.R., Jenkins, M.J.J., and Making, D. (2006). Gender differences in risk assessment: Why do women take fewer risks than men? *1*, 48-63.

11. Brummelte, S., and Galea, L.A. (2016). Postpartum depression: Etiology, treatment and consequences for maternal care. Hormones and behavior *77*, 153-166. 10.1016/j.yhbeh.2015.08.008.

12. Lonstein, J.S., Lévy, F., and Fleming, A.S. (2015). Common and divergent psychobiological mechanisms underlying maternal behaviors in non-human and human mammals. Hormones and behavior *73*, 156-185. 10.1016/j.yhbeh.2015.06.011.

13. Zipple, M.N., Chang Kuo, D., Meng, X., Reichard, T.M., Guess, K., Vogt, C.C., Moeller, A.H., and Sheehan, M.J. (2025). Competitive social feedback amplifies the role of early life contingency in male mice. Science (New York, N.Y.) *387*, 81-85. 10.1126/science.adq0579.

14. Archer, J. (2019). The reality and evolutionary significance of human psychological sex differences. Biological reviews of the Cambridge Philosophical Society *94*, 1381-1415. 10.1111/brv.12507.

15. Trivers, R.L. (2017). Parental investment and sexual selection. In Sexual selection and the descent of man, (Routledge), pp. 136-179.

16. Arakawa, H. (2019). Age and sex differences in the innate defensive behaviors of C57BL/6 mice exhibited in a fear conditioning paradigm and upon exposure to a predatory odor. Physiology & behavior *204*, 264-274. 10.1016/j.physbeh.2019.02.030.

17. Blanchard, D.C., Shepherd, J.K., De Padua Carobrez, A., and Blanchard, R.J. (1991). Sex effects in defensive behavior: baseline differences and drug interactions. Neuroscience and biobehavioral reviews *15*, 461-468. 10.1016/s0149-7634(05)80132-0.

18. Inoue, S., Yang, R., Tantry, A., Davis, C.H., Yang, T., Knoedler, J.R., Wei, Y., Adams, E.L., Thombare, S., Golf, S.R., et al. (2019). Periodic Remodeling in a Neural Circuit Governs Timing of Female Sexual Behavior. Cell *179*, 1393-1408 e1316. 10.1016/j.cell.2019.10.025.

19. Bayless, D.W., Davis, C.O., Yang, R., Wei, Y., de Andrade Carvalho, V.M., Knoedler, J.R., Yang, T., Livingston, O., Lomvardas, A., Martins, G.J., et al. (2023). A neural circuit for male sexual behavior and reward. Cell *186*, 3862-3881 e3828. 10.1016/j.cell.2023.07.021.

20. Yang, C.F., Chiang, M.C., Gray, D.C., Prabhakaran, M., Alvarado, M., Juntti, S.A., Unger, E.K., Wells, J.A., and Shah, N.M. (2013). Sexually dimorphic neurons in the ventromedial hypothalamus govern mating in both sexes and aggression in males. Cell *153*, 896-909. 10.1016/j.cell.2013.04.017.

21. Yang, T., Bayless, D.W., Wei, Y., Landayan, D., Marcelo, I.M., Wang, Y., DeNardo, L.A., Luo, L., Druckmann, S., and Shah, N.M. (2023). Hypothalamic neurons that mirror aggression. Cell *186*, 1195-1211 e1119. 10.1016/j.cell.2023.01.022.

22. Nair, A., Karigo, T., Yang, B., Ganguli, S., Schnitzer, M.J., Linderman, S.W., Anderson, D.J., and Kennedy, A. (2023). An approximate line attractor in the hypothalamus encodes an aggressive state. Cell *186*, 178-193.e115. 10.1016/j.cell.2022.11.027.

23. Yan, R., Wei, D., Varshneya, A., Shan, L., Dai, B., Asencio, H.J., 3rd, Gollamudi, A., and Lin, D. (2024). The multi-stage plasticity in the aggression circuit underlying the winner effect. Cell *187*, 6785-6803.e6718. 10.1016/j.cell.2024.09.030.

24. Bayless, D.W., Yang, T., Mason, M.M., Susanto, A.A.T., Lobdell, A., and Shah, N.M. (2019). Limbic Neurons Shape Sex Recognition and Social Behavior in Sexually Naive Males. Cell *176*, 1190-1205 e1120. 10.1016/j.cell.2018.12.041.

25. Wang, Y., Song, X., Chen, X., Zhou, Y., Ma, J., Zhang, F., Wei, L., Qi, G., Yadav, N., Miao, B., et al. (2025). Integrating reproductive states and social cues in the control of sociosexual behaviors. Cell *188*, 3530-3549.e3524. 10.1016/j.cell.2025.04.035.

26. Wei, A., Zhao, A., Zheng, C., Dong, N., Cheng, X., Duan, X., Zhong, S., Liu, X., Jian, J., Qin, Y., et al. (2025). Sexually dimorphic dopaminergic circuits determine sex preference. Science (New York, N.Y.) *387*, eadq7001. 10.1126/science.adq7001.

27. Watts, J.C., and Fitzpatrick, C.L. (2024). The effects of intersexual interactions on survival can drive the evolution of female ornaments in the absence of mate limitation. Journal of evolutionary biology *37*, 1356-1367. 10.1093/jeb/voae121.

28. Kokko, H., and Jennions, M. (2003). It takes two to tango. Trends in Ecology & Evolution *18*, 103-104. https://doi.org/10.1016/S0169-5347(03)00009-0.

29. Trivers, R. (2002). Natural Selection and Social Theory : Selected Papers of Robert Trivers (Oxford University Press, Incorporated).

30. Liu, X., Feng, X., Huang, H., Huang, K., Xu, Y., Ye, S., Tseng, Y.T., Wei, P., Wang, L., and Wang, F. (2022). Male and female mice display consistent lifelong ability to address potential life-threatening cues using different post-threat coping strategies. BMC biology *20*, 281. 10.1186/s12915-022-01486-x.

31. Knoedler, J.R., Inoue, S., Bayless, D.W., Yang, T., Tantry, A., Davis, C.H., Leung, N.Y., Parthasarathy, S., Wang, G., Alvarado, M., et al. (2022). A functional cellular framework for

sex and estrous cycle-dependent gene expression and behavior. Cell *185*, 654-671 e622. 10.1016/j.cell.2021.12.031.

32.   Fratzl, A., Koltchev, A.M., Vissers, N., Tan, Y.L., Marques-Smith, A., Stempel, A.V., Branco, T., and Hofer, S.B. (2021). Flexible inhibitory control of visually evoked defensive behavior by the ventral lateral geniculate nucleus. Neuron *109*, 3810-3822.e3819. 10.1016/j.neuron.2021.09.003.

33.   Wei, P., Liu, N., Zhang, Z., Liu, X., Tang, Y., He, X., Wu, B., Zhou, Z., Liu, Y., Li, J., et al. (2015). Processing of visually evoked innate fear by a non-canonical thalamic pathway. Nat Commun *6*, 6756. 10.1038/ncomms7756.

34.   Shang, C., Chen, Z., Liu, A., Li, Y., Zhang, J., Qu, B., Yan, F., Zhang, Y., Liu, W., Liu, Z., et al. (2018). Divergent midbrain circuits orchestrate escape and freezing responses to looming stimuli in mice. Nature communications *9*, 1232. 10.1038/s41467-018-03580-7.

35.   Koch, M., Varela, L., Kim, J.G., Kim, J.D., Hernandez-Nuno, F., Simonds, S.E., Castorena, C.M., Vianna, C.R., Elmquist, J.K., Morozov, Y.M., et al. (2015). Hypothalamic POMC neurons promote cannabinoid-induced feeding. Nature *519*, 45-50. 10.1038/nature14260.

36.   Topchiy, I., and Kocsis, B. (2025). CB-1 receptor agonist drastically changes oscillatory activity, defining active sleep. Proceedings of the National Academy of Sciences of the United States of America *122*, e2411063122. 10.1073/pnas.2411063122.

37.   Busquets-García, A., Bolaños, J.P., and Marsicano, G. (2022). Metabolic Messengers: endocannabinoids. Nature metabolism *4*, 848-855. 10.1038/s42255-022-00600-1.

38.   Shang, C., Liu, Z., Chen, Z., Shi, Y., Wang, Q., Liu, S., Li, D., and Cao, P. (2015). BRAIN CIRCUITS. A parvalbumin-positive excitatory visual pathway to trigger fear responses in mice. Science (New York, N.Y.) *348*, 1472-1477. 10.1126/science.aaa8694.

39.   Rangari, V.A., O'Brien, E.S., Powers, A.S., Slivicki, R.A., Bertels, Z., Appourchaux, K., Aydin, D., Ramos-Gonzalez, N., Mwirigi, J., Lin, L., et al. (2025). A cryptic pocket in CB1 drives peripheral and functional selectivity. Nature *640*, 265-273. 10.1038/s41586-025-08618-7.

40.   McPartland, J.M., Matias, I., Di Marzo, V., and Glass, M. (2006). Evolutionary origins of the endocannabinoid system. Gene *370*, 64-74. 10.1016/j.gene.2005.11.004.

41.   Mizuno, I., Matsuda, S., Tohyama, S., and Mizutani, A. (2022). The role of the cannabinoid system in fear memory and extinction in male and female mice. Psychoneuroendocrinology *138*, 105688. 10.1016/j.psyneuen.2022.105688.

42.   Bowers, M.E., and Ressler, K.J. (2016). Sex-dependence of anxiety-like behavior in cannabinoid receptor 1 (Cnr1) knockout mice. Behavioural brain research *300*, 65-69. 10.1016/j.bbr.2015.12.005.

43.   Liu, X., Li, X., Zhao, G., Wang, F., and Wang, L. (2020). Sexual dimorphic distribution of cannabinoid 1 receptor mRNA in adult C57BL/6J mice. The Journal of comparative neurology *528*, 1986-1999. 10.1002/cne.24868.

44.   Castelli, M.P., Fadda, P., Casu, A., Spano, M.S., Casti, A., Fratta, W., and Fattore, L. (2014). Male and female rats differ in brain cannabinoid CB1 receptor density and function and in behavioural traits predisposing to drug addiction: effect of ovarian hormones. Current pharmaceutical design *20*, 2100-2113. 10.2174/13816128113199990430.

45.   Guo, B., Xi, K., Mao, H., Ren, K., Xiao, H., Hartley, N.D., Zhang, Y., Kang, J., Liu, Y., Xie, Y., et al. (2024). CB1R dysfunction of inhibitory synapses in the ACC drives chronic social isolation stress-induced social impairments in male mice. Neuron *112*, 441-457.e446.

10.1016/j.neuron.2023.10.027.

46. Wei, D., Allsop, S., Tye, K., and Piomelli, D. (2017). Endocannabinoid Signaling in the Control of Social Behavior. Trends Neurosci *40*, 385-396. 10.1016/j.tins.2017.04.005.

47. Schechter, M., Pinhasov, A., Weller, A., and Fride, E. (2012). Blocking the postpartum mouse dam's CB1 receptors impairs maternal behavior as well as offspring development and their adult social-emotional behavior. Behavioural brain research *226*, 481-492. 10.1016/j.bbr.2011.10.016.

48. Vargish, G.A., Pelkey, K.A., Yuan, X., Chittajallu, R., Collins, D., Fang, C., and McBain, C.J. (2017). Persistent inhibitory circuit defects and disrupted social behaviour following in utero exogenous cannabinoid exposure. Molecular psychiatry *22*, 56-67. 10.1038/mp.2016.17.

49. Rodríguez-Manzo, G., and Canseco-Alba, A. (2023). The endogenous cannabinoid system modulates male sexual behavior expression. Frontiers in behavioral neuroscience *17*, 1198077. 10.3389/fnbeh.2023.1198077.

50. Heinbockel, T., Brager, D.H., Reich, C.G., Zhao, J., Muralidharan, S., Alger, B.E., and Kao, J.P. (2005). Endocannabinoid signaling dynamics probed with optical tools. J Neurosci *25*, 9449-9459. 10.1523/JNEUROSCI.2078-05.2005.

51. Ye, J., Xu, Y., Huang, K., Wang, X., Wang, L., and Wang, F. (2025). Hierarchical behavioral analysis framework as a platform for standardized quantitative identification of behaviors. Cell Reports *44*. 10.1016/j.celrep.2025.115239.

52. Liu, Y., Savier, E.L., DePiero, V.J., Chen, C., Schwalbe, D.C., Abraham-Fan, R.J., Chen, H., Campbell, J.N., and Cang, J. (2023). Mapping visual functions onto molecular cell types in the mouse superior colliculus. Neuron *111*, 1876-1886.e1875. 10.1016/j.neuron.2023.03.036.

53. Liu, X., Huang, H., Zhang, Y., Wang, L., and Wang, F. (2022). Sexual Dimorphism of Inputs to the Lateral Habenula in Mice. Neuroscience bulletin *38*, 1439-1456. 10.1007/s12264-022-00885-y.

54. Hu, H., Cui, Y., and Yang, Y. (2020). Circuits and functions of the lateral habenula in health and in disease. Nature reviews. Neuroscience *21*, 277-295. 10.1038/s41583-020-0292-4.

55. Lazaridis, I., Tzortzi, O., Weglage, M., Märtin, A., Xuan, Y., Parent, M., Johansson, Y., Fuzik, J., Fürth, D., Fenno, L.E., et al. (2019). A hypothalamus-habenula circuit controls aversion. Molecular psychiatry *24*, 1351-1368. 10.1038/s41380-019-0369-5.

56. Manssuer, L., Ding, Q., Zhang, Y., Gong, H., Liu, W., Yang, R., Zhang, C., Zhao, Y., Pan, Y., Zhan, S., et al. (2023). Risk and aversion coding in human habenula high gamma activity. Brain : a journal of neurology *146*, 2642-2653. 10.1093/brain/awac456.

57. Lecca, S., Namboodiri, V.M.K., Restivo, L., Gervasi, N., Pillolla, G., Stuber, G.D., and Mameli, M. (2020). Heterogeneous Habenular Neuronal Ensembles during Selection of Defensive Behaviors. Cell Rep *31*, 107752. 10.1016/j.celrep.2020.107752.

58. Yang, H., Yang, J., Xi, W., Hao, S., Luo, B., He, X., Zhu, L., Lou, H., Yu, Y.Q., Xu, F., et al. (2016). Laterodorsal tegmentum interneuron subtypes oppositely regulate olfactory cue-induced innate fear. Nature neuroscience *19*, 283-289. 10.1038/nn.4208.

59. Lecca, S., Meye, F.J., Trusel, M., Tchenio, A., Harris, J., Schwarz, M.K., Burdakov, D., Georges, F., and Mameli, M. (2017). Aversive stimuli drive hypothalamus-to-habenula excitation to promote escape behavior. eLife *6*. 10.7554/eLife.30697.

60. Congiu, M., Mondoloni, S., Zouridis, I.S., Schmors, L., Lecca, S., Lalive, A.L., Ginggen, K., Deng, F., Berens, P., Paolicelli, R.C., et al. (2023). Plasticity of neuronal dynamics in the

lateral habenula for cue-punishment associative learning. Molecular psychiatry *28*, 5118-5127. 10.1038/s41380-023-02155-3.

61. Berger, A.L., Henricks, A.M., Lugo, J.M., Wright, H.R., Warrick, C.R., Sticht, M.A., Morena, M., Bonilla, I., Laredo, S.A., Craft, R.M., et al. (2018). The Lateral Habenula Directs Coping Styles Under Conditions of Stress via Recruitment of the Endocannabinoid System. Biological psychiatry *84*, 611-623. 10.1016/j.biopsych.2018.04.018.

62. Dong, A., He, K., Dudok, B., Farrell, J.S., Guan, W., Liput, D.J., Puhl, H.L., Cai, R., Wang, H., Duan, J., et al. (2022). A fluorescent sensor for spatiotemporally resolved imaging of endocannabinoid dynamics in vivo. Nature biotechnology *40*, 787-798. 10.1038/s41587-021-01074-4.

63. Dudok, B., Fan, L.Z., Farrell, J.S., Malhotra, S., Homidan, J., Kim, D.K., Wenardy, C., Ramakrishnan, C., Li, Y., Deisseroth, K., and Soltesz, I. (2024). Retrograde endocannabinoid signaling at inhibitory synapses in vivo. Science (New York, N.Y.) *383*, 967-970. 10.1126/science.adk3863.

64. Gunduz-Cinar, O., Castillo, L.I., Xia, M., Van Leer, E., Brockway, E.T., Pollack, G.A., Yasmin, F., Bukalo, O., Limoges, A., Oreizi-Esfahani, S., et al. (2023). A cortico-amygdala neural substrate for endocannabinoid modulation of fear extinction. Neuron *111*, 3053-3067.e3010. 10.1016/j.neuron.2023.06.023.

65. Oliet, S.H., Baimoukhametova, D.V., Piet, R., and Bains, J.S. (2007). Retrograde regulation of GABA transmission by the tonic release of oxytocin and endocannabinoids governs postsynaptic firing. The Journal of neuroscience : the official journal of the Society for Neuroscience *27*, 1325-1333. 10.1523/jneurosci.2676-06.2007.

66. Hentges, S.T., Low, M.J., and Williams, J.T. (2005). Differential regulation of synaptic inputs by constitutively released endocannabinoids and exogenous cannabinoids. The Journal of neuroscience : the official journal of the Society for Neuroscience *25*, 9746-9751. 10.1523/jneurosci.2769-05.2005.

67. Ogasawara, D., Deng, H., Viader, A., Baggelaar, M.P., Breman, A., den Dulk, H., van den Nieuwendijk, A.M., Soethoudt, M., van der Wel, T., Zhou, J., et al. (2016). Rapid and profound rewiring of brain lipid signaling networks by acute diacylglycerol lipase inhibition. Proceedings of the National Academy of Sciences of the United States of America *113*, 26-33. 10.1073/pnas.1522364112.

68. Bisogno, T., Howell, F., Williams, G., Minassi, A., Cascio, M.G., Ligresti, A., Matias, I., Schiano-Moriello, A., Paul, P., Williams, E.J., et al. (2003). Cloning of the first sn1-DAG lipases points to the spatial and temporal regulation of endocannabinoid signaling in the brain. The Journal of cell biology *163*, 463-468. 10.1083/jcb.200305129.

69. Li, L., Feng, X., Zhou, Z., Zhang, H., Shi, Q., Lei, Z., Shen, P., Yang, Q., Zhao, B., Chen, S., et al. (2018). Stress Accelerates Defensive Responses to Looming in Mice and Involves a Locus Coeruleus-Superior Colliculus Projection. Current biology : CB *28*, 859-871.e855. 10.1016/j.cub.2018.02.005.

70. Soria-Gomez, E., Bellocchio, L., Reguero, L., Lepousez, G., Martin, C., Bendahmane, M., Ruehle, S., Remmers, F., Desprez, T., Matias, I., et al. (2014). The endocannabinoid system controls food intake via olfactory processes. Nat Neurosci *17*, 407-415. 10.1038/nn.3647.

71. Zhao, Z., Soria-Gómez, E., Varilh, M., Covelo, A., Julio-Kalajzić, F., Cannich, A., Castiglione, A., Vanhoutte, L., Duveau, A., Zizzari, P., et al. (2020). A Novel Cortical Mechanism for

Top-Down Control of Water Intake. Current biology : CB *30*, 4789-4798.e4784. 10.1016/j.cub.2020.09.011.

72. Stapf, C.A., Keefer, S.E., McInerney, J.M., Cheer, J.F., and Calu, D.J. (2025). Dorsomedial Striatum (DMS) CB1R Signaling Promotes Pavlovian Devaluation Sensitivity in Male Long Evans Rats and Reduces DMS Inhibitory Synaptic Transmission in Both Sexes. eNeuro *12*. 10.1523/eneuro.0341-24.2024.

73. Soria-Gomez, E., Pagano Zottola, A.C., Mariani, Y., Desprez, T., Barresi, M., Bonilla-Del Río, I., Muguruza, C., Le Bon-Jego, M., Julio-Kalajzić, F., Flynn, R., et al. (2021). Subcellular specificity of cannabinoid effects in striatonigral circuits. Neuron *109*, 1513-1526.e1511. 10.1016/j.neuron.2021.03.007.

74. Zou, G., Xia, J., Luo, H., Xiao, D., Jin, J., Miao, C., Zuo, X., Gao, Q., Zhang, Z., Xue, T., et al. (2022). Combined alcohol and cannabinoid exposure leads to synergistic toxicity by affecting cerebellar Purkinje cells. Nature metabolism *4*, 1138-1149. 10.1038/s42255-022-00633-6.

75. Wang, H., Dong, P., He, C., Feng, X.Y., Huang, Y., Yang, W.W., Gao, H.J., Shen, X.F., Lin, S., Cao, S.X., et al. (2020). Incerta-thalamic Circuit Controls Nocifensive Behavior via Cannabinoid Type 1 Receptors. Neuron *107*, 538-551.e537. 10.1016/j.neuron.2020.04.027.

76. Shen, C.J., Zheng, D., Li, K.X., Yang, J.M., Pan, H.Q., Yu, X.D., Fu, J.Y., Zhu, Y., Sun, Q.X., Tang, M.Y., et al. (2019). Cannabinoid CB(1) receptors in the amygdalar cholecystokinin glutamatergic afferents to nucleus accumbens modulate depressive-like behavior. Nature medicine *25*, 337-349. 10.1038/s41591-018-0299-9.

77. Mederos, S., Blakely, P., Vissers, N., Clopath, C., and Hofer, S.B. (2025). Overwriting an instinct: Visual cortex instructs learning to suppress fear responses. Science (New York, N.Y.) *387*, 682-688. 10.1126/science.adr2247.

78. Chai, R., Wang, N., Nie, J., Xu, Z., Zhang, S., Deng, S., Wang, R., Li, M., Gao, X., Geng, R., et al. (2025). Endocannabinoids disinhibit the ventral tegmental nucleus of Gudden to dorsal premammillary nucleus pathway to enhance escape behavior following learned threat experience. Nature communications *16*, 4885. 10.1038/s41467-025-60080-1.

79. Bayless, D.W., Davis, C.O., Yang, R., Wei, Y., Miessler de Andrade Carvalho, V., Knoedler, J.R., Yang, T., Livingston, O., Lomvardas, A., Martins, G.J., et al. (2024). A neural circuit for male sexual behavior and reward. Cell *187*, 3783. 10.1016/j.cell.2024.05.047.

80. Shen, C.J., Zheng, D., Li, K.X., Yang, J.M., Pan, H.Q., Yu, X.D., Fu, J.Y., Zhu, Y., Sun, Q.X., Tang, M.Y., et al. (2019). Cannabinoid CB1 receptors in the amygdalar cholecystokinin glutamatergic afferents to nucleus accumbens modulate depressive-like behavior. Nat Med *25*, 337-349. 10.1038/s41591-018-0299-9.

81. Wang, H., Dong, P., He, C., Feng, X.Y., Huang, Y., Yang, W.W., Gao, H.J., Shen, X.F., Lin, S., Cao, S.X., et al. (2020). Incerta-thalamic Circuit Controls Nocifensive Behavior via Cannabinoid Type 1 Receptors. Neuron *107*, 538-551 e537. 10.1016/j.neuron.2020.04.027.

82. Xue, B., Zhang, M.X., Bi, X.C., Lai, S.P., Bie, X.T., Dong, Y., Li, J.F., Gao, F., Zhang, X., and Wang, Y. (2025). An Amygdala-hippocampus Circuit for Endocannabinoid Modulation of Anxiety Avoidance. Advanced science (Weinheim, Baden-Wurttemberg, Germany), e05121. 10.1002/advs.202505121.

83. Soria-Gomez, E., Busquets-Garcia, A., Hu, F., Mehidi, A., Cannich, A., Roux, L., Louit, I., Alonso, L., Wiesner, T., Georges, F., et al. (2015). Habenular CB1 Receptors Control the Expression of Aversive Memories. Neuron *88*, 306-313. 10.1016/j.neuron.2015.08.035.

84. Han, J., Kesner, P., Metna-Laurent, M., Duan, T., Xu, L., Georges, F., Koehl, M., Abrous, D.N., Mendizabal-Zubiaga, J., Grandes, P., et al. (2012). Acute cannabinoids impair working memory through astroglial CB1 receptor modulation of hippocampal LTD. Cell *148*, 1039-1050. 10.1016/j.cell.2012.01.037.

85. Skupio, U., Welte, J., Serrat, R., Eraso-Pichot, A., Julio-Kalajzic, F., Gisquet, D., Cannich, A., Delcasso, S., Matias, I., Fundazuri, U.B., et al. (2023). Mitochondrial cannabinoid receptors gate corticosterone impact on novel object recognition. Neuron *111*, 1887-1897 e1886. 10.1016/j.neuron.2023.04.001.

86. Soria-Gomez, E., Pagano Zottola, A.C., Mariani, Y., Desprez, T., Barresi, M., Bonilla-Del Rio, I., Muguruza, C., Le Bon-Jego, M., Julio-Kalajzic, F., Flynn, R., et al. (2021). Subcellular specificity of cannabinoid effects in striatonigral circuits. Neuron *109*, 1513-1526 e1511. 10.1016/j.neuron.2021.03.007.

87. Dudek, K.A., Paton, S.E.J., Binder, L.B., Collignon, A., Dion-Albert, L., Cadoret, A., Lebel, M., Lavoie, O., Bouchard, J., Kaufmann, F.N., et al. (2025). Astrocytic cannabinoid receptor 1 promotes resilience by dampening stress-induced blood-brain barrier alterations. Nature neuroscience *28*, 766-782. 10.1038/s41593-025-01891-9.

88. Bellocchio, L., Lafenêtre, P., Cannich, A., Cota, D., Puente, N., Grandes, P., Chaouloff, F., Piazza, P.V., and Marsicano, G. (2010). Bimodal control of stimulated food intake by the endocannabinoid system. Nature neuroscience *13*, 281-283. 10.1038/nn.2494.

89. Noriega-Prieto, J.A., Falcón-Moya, R., Noeker, J.A., Cai, R., Fundazuri, U.B., Eraso-Pichot, A., Cai, S., Guttipatti, P., Belisle, L., Rodríguez-Moreno, A., et al. (2025). Distinct endocannabinoids specifically signal to astrocytes or neurons in the adult mouse hippocampus. Nature neuroscience. 10.1038/s41593-025-02148-1.

90. Zhou, Z., Liu, X., Chen, S., Zhang, Z., Liu, Y., Montardy, Q., Tang, Y., Wei, P., Liu, N., Li, L., et al. (2019). A VTA GABAergic Neural Circuit Mediates Visually Evoked Innate Defensive Responses. Neuron *103*, 473-488.e476. 10.1016/j.neuron.2019.05.027.

91. Evans, D.A., Stempel, A.V., Vale, R., Ruehle, S., Lefler, Y., and Branco, T. (2018). A synaptic threshold mechanism for computing escape decisions. Nature *558*, 590-594. 10.1038/s41586-018-0244-6.

92. Xie, Z., Wang, M., Liu, Z., Shang, C., Zhang, C., Sun, L., Gu, H., Ran, G., Pei, Q., Ma, Q., et al. (2021). Transcriptomic encoding of sensorimotor transformation in the midbrain. eLife *10*. 10.7554/eLife.69825.

93. Zhao, Z., and Davis, M. (2004). Fear-potentiated startle in rats is mediated by neurons in the deep layers of the superior colliculus/deep mesencephalic nucleus of the rostral midbrain through the glutamate non-NMDA receptors. The Journal of neuroscience : the official journal of the Society for Neuroscience *24*, 10326-10334. 10.1523/jneurosci.2758-04.2004.

94. Li, C., Kühn, N.K., Alkislar, I., Sans-Dublanc, A., Zemmouri, F., Paesmans, S., Calzoni, A., Ooms, F., Reinhard, K., and Farrow, K. (2023). Pathway-specific inputs to the superior colliculus support flexible responses to visual threat. Science advances *9*, eade3874. 10.1126/sciadv.ade3874.

95. Yizhar, O., Fenno, L.E., Prigge, M., Schneider, F., Davidson, T.J., O'Shea, D.J., Sohal, V.S., Goshen, I., Finkelstein, J., Paz, J.T., et al. (2011). Neocortical excitation/inhibition balance in information processing and social dysfunction. Nature *477*, 171-178. 10.1038/nature10360.

96. Fagiolini, M., Fritschy, J.M., Low, K., Mohler, H., Rudolph, U., and Hensch, T.K. (2004).

Specific GABAA circuits for visual cortical plasticity. Science *303*, 1681-1683. 10.1126/science.1091032.

97. Pi, H.J., Hangya, B., Kvitsiani, D., Sanders, J.I., Huang, Z.J., and Kepecs, A. (2013). Cortical interneurons that specialize in disinhibitory control. Nature *503*, 521-524. 10.1038/nature12676.

98. Li, H.Q., Jiang, W., Ling, L., Pratelli, M., Chen, C., Gupta, V., Godavarthi, S.K., and Spitzer, N.C. (2024). Generalized fear after acute stress is caused by change in neuronal cotransmitter identity. Science *383*, 1252-1259. 10.1126/science.adj5996.

99. Munoz, W., Tremblay, R., Levenstein, D., and Rudy, B. (2017). Layer-specific modulation of neocortical dendritic inhibition during active wakefulness. Science *355*, 954-959. 10.1126/science.aag2599.

100. Lu, L., Ren, Y., Yu, T., Liu, Z., Wang, S., Tan, L., Zeng, J., Feng, Q., Lin, R., Liu, Y., et al. (2020). Control of locomotor speed, arousal, and hippocampal theta rhythms by the nucleus incertus. Nat Commun *11*, 262. 10.1038/s41467-019-14116-y.

101. Hart, M.P., and Hobert, O. (2018). Neurexin controls plasticity of a mature, sexually dimorphic neuron. Nature *553*, 165-170. 10.1038/nature25192.

102. VanRyzin, J.W., Marquardt, A.E., Argue, K.J., Vecchiarelli, H.A., Ashton, S.E., Arambula, S.E., Hill, M.N., and McCarthy, M.M. (2019). Microglial Phagocytosis of Newborn Cells Is Induced by Endocannabinoids and Sculpts Sex Differences in Juvenile Rat Social Play. Neuron *102*, 435-449 e436. 10.1016/j.neuron.2019.02.006.

103. Kim, Y., Yang, G.R., Pradhan, K., Venkataraju, K.U., Bota, M., García Del Molino, L.C., Fitzgerald, G., Ram, K., He, M., Levine, J.M., et al. (2017). Brain-wide Maps Reveal Stereotyped Cell-Type-Based Cortical Architecture and Subcortical Sexual Dimorphism. Cell *171*, 456-469.e422. 10.1016/j.cell.2017.09.020.

104. Gruene, T.M., Roberts, E., Thomas, V., Ronzio, A., and Shansky, R.M. (2015). Sex-specific neuroanatomical correlates of fear expression in prefrontal-amygdala circuits. Biol Psychiatry *78*, 186-193. 10.1016/j.biopsych.2014.11.014.

105. Rodríguez-Montes, L., Ovchinnikova, S., Yuan, X., Studer, T., Sarropoulos, I., Anders, S., Kaessmann, H., and Cardoso-Moreira, M. (2023). Sex-biased gene expression across mammalian organ development and evolution. Science (New York, N.Y.) *382*, eadf1046. 10.1126/science.adf1046.

106. Blume, S.R., Freedberg, M., Vantrease, J.E., Chan, R., Padival, M., Record, M.J., DeJoseph, M.R., Urban, J.H., and Rosenkranz, J.A. (2017). Sex- and Estrus-Dependent Differences in Rat Basolateral Amygdala. J Neurosci *37*, 10567-10586. 10.1523/JNEUROSCI.0758-17.2017.

107. Chen, P.B., Hu, R.K., Wu, Y.E., Pan, L., Huang, S., Micevych, P.E., and Hong, W. (2019). Sexually Dimorphic Control of Parenting Behavior by the Medial Amygdala. Cell *176*, 1206-1221.e1218. 10.1016/j.cell.2019.01.024.

108. Prévost, E.D., Phillips, A., Lauridsen, K., Enserro, G., Rubinstein, B., Alas, D., McGovern, D.J., Ly, A., Hotchkiss, H., Banks, M., et al. (2024). Monosynaptic Inputs to Ventral Tegmental Area Glutamate and GABA Co-transmitting Neurons. The Journal of neuroscience : the official journal of the Society for Neuroscience *44*. 10.1523/jneurosci.2184-23.2024.

109. Xin, Q., Wang, J., Zheng, J., Tan, Y., Jia, X., Ni, Z., Xu, Z., Feng, J., Wu, Z., Li, Y., et al. (2025). Neuron-astrocyte coupling in lateral habenula mediates depressive-like behaviors. Cell *188*, 3291-3309 e3224. 10.1016/j.cell.2025.04.010.

110. Fan, Z., Chang, J., Liang, Y., Zhu, H., Zhang, C., Zheng, D., Wang, J., Xu, Y., Li, Q.J., and Hu, H. (2023). Neural mechanism underlying depressive-like state associated with social status loss. Cell *186*, 560-576 e517. 10.1016/j.cell.2022.12.033.

111. Huang, H., Liu, X., Wang, L., and Wang, F. (2024). Whole-brain connections of glutamatergic neurons in the mouse lateral habenula in both sexes. Biol Sex Differ *15*, 37. 10.1186/s13293-024-00611-5.

112. Groos, D., Reuss, A.M., Rupprecht, P., Stachniak, T., Lewis, C., Han, S., Roggenbach, A., Sturman, O., Sych, Y., Wieckhorst, M., et al. (2025). A distinct hypothalamus-habenula circuit governs risk preference. Nat Neurosci *28*, 361-373. 10.1038/s41593-024-01856-4.

113. Lecca, S., Congiu, M., Royon, L., Restivo, L., Girard, B., Mazaré, N., Bellone, C., Telley, L., and Mameli, M. (2023). A neural substrate for negative affect dictates female parental behavior. Neuron *111*, 1094-1103.e1098. 10.1016/j.neuron.2023.01.003.

114. Benedict, J., Cudmore, R.H., Oden, D., Spruell, A., and Linden, D.J. (2025). The Lateral Habenula Is Necessary for Maternal Behavior in the Naturally Parturient Primiparous Mouse Dam. eNeuro *12*. 10.1523/eneuro.0092-24.2024.

115. Ravenelle, R., Lee, J., Fernandes-Henriques, C., Liu, J., Friedman, A.K., Likhtik, E., and Burghardt, N.S. (2025). Serotonergic modulation of the BNST-CeA pathway reveals sex differences in fear learning. Nature neuroscience *28*, 1897-1909. 10.1038/s41593-025-02025-x.

116. van Doeselaar, L., Stark, T., Mitra, S., Yang, H., Bordes, J., Stolwijk, L., Engelhardt, C., Kovarova, V., Narayan, S., Brix, L.M., et al. (2023). Sex-specific and opposed effects of FKBP51 in glutamatergic and GABAergic neurons: Implications for stress susceptibility and resilience. Proceedings of the National Academy of Sciences of the United States of America *120*, e2300722120. 10.1073/pnas.2300722120.

117. Jiang, T., Feng, M., Hutsell, A., and Lüscher, B. (2025). Sex-specific GABAergic microcircuits that switch vulnerability into resilience to stress and reverse the effects of chronic stress exposure. Molecular psychiatry *30*, 2297-2308. 10.1038/s41380-024-02835-8.

118. Zhang, M., Ni, Z., Ma, J., Liu, A., Liu, Y., Lou, Q., Dong, W.Y., Zhang, Z., Li, J., and Cao, P. (2025). A neural circuit for sex-dependent conditioned pain hypersensitivity in mice. Nature communications *16*, 3639. 10.1038/s41467-025-58851-x.

119. Tiihonen, J., Koskuvi, M., Storvik, M., Hyötyläinen, I., Gao, Y., Puttonen, K.A., Giniatullina, R., Poguzhelskaya, E., Ojansuu, I., Vaurio, O., et al. (2019). Sex-specific transcriptional and proteomic signatures in schizophrenia. Nature communications *10*, 3933. 10.1038/s41467-019-11797-3.

120. Yang, L., Guo, C., Zheng, Z., Dong, Y., Xie, Q., Lv, Z., Li, M., Lu, Y., Guo, X., Deng, R., et al. (2025). Stress dynamically modulates neuronal autophagy to gate depression onset. Nature *641*, 427-437. 10.1038/s41586-025-08807-4.

121. Xin, Q., Wang, J., Zheng, J., Tan, Y., Jia, X., Ni, Z., Xu, Z., Feng, J., Wu, Z., Li, Y., et al. (2025). Neuron-astrocyte coupling in lateral habenula mediates depressive-like behaviors. Cell *188*, 3291-3309.e3224. 10.1016/j.cell.2025.04.010.

122. Li, M. (2022). Lateral habenula neurocircuits mediate the maternal disruptive effect of maternal stress: A hypothesis. Zoological research *43*, 166-175. 10.24272/j.issn.2095-8137.2021.362.

123. Wang, B., and Li, M. (2025). Does lateral habenula mediate effects of gestational stress on rat

maternal behavior? Brain behavior and immunity integrative *9*. 10.1016/j.bbii.2024.100098.

124.   Eisenstein, E., and Eisenstein, D. (2006). A behavioral homeostasis theory of habituation and sensitization: II. Further developments and predictions. Reviews in the Neurosciences *17*, 533-558.

125.   Eisenstein, E., Eisenstein, D., and Smith, J.C. (2001). The evolutionary significance of habituation and sensitization across phylogeny: A behavioral homeostasis model. Integrative Physiological & Behavioral Science *36*, 251-265.

126.   Eisenstein, E.M., Eisenstein, D.L., Sarma, J.S., Knapp, H., and Smith, J.C. (2012). Some new speculative ideas about the "behavioral homeostasis theory" as to how the simple learned behaviors of habituation and sensitization improve organism survival throughout phylogeny. Communicative & Integrative Biology *5*, 233-239.

127.   Lee, S.S., Liu, Q., Cheng, A.H.R., Kim, D.W., Boudreau, D.M., Mehta, A., Keles, M.F., Fejfer, R., Palmer, I., Park, K.H., et al. (2025). Sleep need-dependent plasticity of a thalamic circuit promotes homeostatic recovery sleep. Science *388*, eadm8203. 10.1126/science.adm8203.

128.   Liu, D., Rahman, M., Johnson, A., Amo, R., Tsutsui-Kimura, I., Sullivan, Z.A., Pena, N., Talay, M., Logeman, B.L., Finkbeiner, S., et al. (2025). A hypothalamic circuit underlying the dynamic control of social homeostasis. Nature *640*, 1000-1010. 10.1038/s41586-025-08617-8.

129.   Gregg, C., Zhang, J., Butler, J.E., Haig, D., and Dulac, C. (2010). Sex-specific parent-of-origin allelic expression in the mouse brain. Science *329*, 682-685. 10.1126/science.1190831.

130.   Yasmin, F., Colangeli, R., Morena, M., Filipski, S., van der Stelt, M., Pittman, Q.J., Hillard, C.J., Teskey, G.C., McEwen, B.S., Hill, M.N., and Chattarji, S. (2020). Stress-induced modulation of endocannabinoid signaling leads to delayed strengthening of synaptic connectivity in the amygdala. Proceedings of the National Academy of Sciences of the United States of America *117*, 650-655. 10.1073/pnas.1910322116.

131.   Lesuis, S.L., Park, S., Hoorn, A., Rashid, A.J., Mocle, A.J., Salter, E.W., Vislavski, S., Gray, M.T., Torelli, A.M., DeCristofaro, A., et al. (2025). Stress disrupts engram ensembles in lateral amygdala to generalize threat memory in mice. Cell *188*, 121-140.e120. 10.1016/j.cell.2024.10.034.

132.   Mei, L., Yan, R., Yin, L., Sullivan, R.M., and Lin, D. (2023). Antagonistic circuits mediating infanticide and maternal care in female mice. Nature *618*, 1006-1016. 10.1038/s41586-023-06147-9.

133.   Ressler, K.J., Berretta, S., Bolshakov, V.Y., Rosso, I.M., Meloni, E.G., Rauch, S.L., and Carlezon, W.A., Jr. (2022). Post-traumatic stress disorder: clinical and translational neuroscience from cells to circuits. Nature reviews. Neurology *18*, 273-288. 10.1038/s41582-022-00635-8.

134.   Azevedo, H., Ferreira, M., Mascarello, A., Osten, P., and Guimarães, C.R.W. (2020). Brain-wide mapping of c-fos expression in the single prolonged stress model and the effects of pretreatment with ACH-000029 or prazosin. Neurobiology of stress *13*, 100226. 10.1016/j.ynstr.2020.100226.

135.   Zhukovskaya, A., Zimmerman, C.A., Willmore, L., Pan-Vazquez, A., Janarthanan, S.R., Lynch, L.A., Falkner, A.L., and Witten, I.B. (2024). Heightened lateral habenula activity during stress produces brainwide and behavioral substrates of susceptibility. Neuron *112*, 3940-3956.e3910. 10.1016/j.neuron.2024.09.009.

136. Liao, Y.Y., Zhang, H., Shen, Q., Cai, C., Ding, Y., Shen, D.D., Guo, J., Qin, J., Dong, Y., Zhang, Y., and Li, X.M. (2023). Snapshot of the cannabinoid receptor 1-arrestin complex unravels the biased signaling mechanism. Cell *186*, 5784-5797 e5717. 10.1016/j.cell.2023.11.017.

137. Allaire, J., Gandrud, C., Russell, K., and Yetman, C.J.R.p.v. (2017). networkD3: D3 JavaScript network graphs from R.

138. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D.A., François, R., Grolemund, G., Hayes, A., Henry, L., and Hester, J.J.J.o.o.s.s. (2019). Welcome to the Tidyverse. *4*, 1686.

139. Zhang, Y., Ke, J., Zhou, Y., Liu, X., Huang, T., and Wang, F. (2022). Sex-specific characteristics of cells expressing the cannabinoid 1 receptor in the dorsal horn of the lumbar spinal cord. J Comp Neurol *530*, 2451-2473. 10.1002/cne.25342.

140. Liu, X., Huang, H., Zhang, Y., Wang, L., and Wang, F. (2022). Sexual Dimorphism of Inputs to the Lateral Habenula in Mice. Neurosci Bull. 10.1007/s12264-022-00885-y.

141. Yang, X., Liu, Q., Zhong, J., Song, R., Zhang, L., and Wang, L. (2020). A simple threat-detection strategy in mice. BMC Biol *18*, 93. 10.1186/s12915-020-00825-0.

142. Tseng, Y.T., Zhao, B., Chen, S., Ye, J., Liu, J., Liang, L., Ding, H., Schaefke, B., Yang, Q., Wang, L., et al. (2022). The subthalamic corticotropin-releasing hormone neurons mediate adaptive REM-sleep responses to threat. Neuron *110*, 1223-1239.e1228. 10.1016/j.neuron.2021.12.033.

143. Silva, B.A., Mattucci, C., Krzywkowski, P., Murana, E., Illarionova, A., Grinevich, V., Canteras, N.S., Ragozzino, D., and Gross, C.T. (2013). Independent hypothalamic circuits for social and predator fear. Nature neuroscience *16*, 1731-1733. 10.1038/nn.3573.

STAR ⬤ METHODS

Detailed methods are provided in the online version of this paper and include the following:

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-Cre	Abcam	Cat #: ab190177, RRID: AB_2860024
Goat anti-GFP	Rockland	Cat #: 600-101-215, RRID: AB_218182
Rabbit anti-RFP	Rockland	Cat #: 600-401-379, RRID: AB_2209751
Sheep antibody to digoxigenin peroxidase	Roche	Cat #: 11207733910, RRID: AB_514500
Donkey anti-Goat, Alexa Fluor 488 conjugate	Jackson ImmunoResearch	Cat #: 705-545-003, RRID: AB_2340428
Donkey anti-Rabbit, Alexa Fluor 488 conjugate	Jackson ImmunoResearch	Cat #: 711-547-003, RRID: AB_2340620
Donkey anti-Rabbit, Alexa Fluor 594 conjugate	Jackson ImmunoResearch	Cat #: 705-585-003, RRID: AB_2340432
Bacterial and virus strains		
AAV-hEF1α-DIO-GCaMP6s-WPRE-pA	Taitool Bioscience Co., Ltd., Shanghai	Cat #: S0351-9
AAV-EF1α-DIO-EGFP	Fuqiang Xu at the SIAT	N/A
AAV2/9-Dlx5/6-Cre-GFP	Liping Wang's Lab at the SIAT	N/A
AAV2/9-Dlx5/6-GFP	Liping Wang's Lab at the SIAT	N/A
AAV2/9-EF1α-EGFP	Liping Wang's Lab at the SIAT	N/A
AAV2/9-hSyn -Cre-WPRE-pA	Taitool Bioscience Co., Ltd., Shanghai	S0278-9
rAAV-CaMKIIα-EGFP	Brain Case	Cat #: BC-0027
rAAV-CaMKIIα-Cre-EGFP	Brain Case	Cat #: BC-0165
AAV2/9-Dlx5/6-DIO-mCherry-WPRE-hGH pA	BrainVTA Co., Ltd., Wuhan	Cat #: PT-0850
AAV2/9-mDLX-DIO-hChR2(H134R)-mCherry-WPRE-hGH pA	BrainVTA Co., Ltd., Wuhan	Cat #: PT-0970
rAAV-mDlx5/6-DIO-hM4D(Gi)-	BrainVTA Co., Ltd.,	Cat #: PT-1191

mCherry-WPRE-hGH pA	Wuhan	
rAAV-mDlx-FLEX-tdTomato-T 2A-Synaptophysin-EGFP	Brain Case	Cat #: BC-0659
AAV2/2Rretro-hSyn-Cre-WPR E-pA	Taitool Bioscience Co., Ltd., Shanghai	Cat #: S0278-2R
AAV9-GfaABC1D-eCB3.0	Yulong Li's Lab at the Peking University	N/A
AAV9-hSyn-RCaMP3	Vigene Biosciences (this paper)	N/A
AAV2-EF1α-iGABASnFR	Fan Jia at the SIAT	N/A
AAV2/9-Dlx5/6-Cre on-ChR2-mCherry	Shanghai Sunbio Medical Biotechnology., Shanghai	N/A
AAV2/9-Dlx5/6-Cre out-ChR2-YFP	Shanghai Sunbio Medical Biotechnology., Shanghai	N/A
Chemicals, peptides, and Recombinant proteins		
NESS 0327	Cayman	Cat #: 10004148
DAPI	Sigma-Aldrich	Cat #: D9542
Clozapine N-oxide	Tocris	Cat #: 4936
Proteinase K	Roche	Cat #: 03654672103
Formamide	Sigma-Aldrich	Cat #: F9037
20×standard saline citrate (SSC)	Thermo Fisher Scientific	Cat #: AM9765
Ribonucleic acid from baker's yeast (S. cerevisiae)	Sigma-Aldrich	Cat #: R6750
Deoxyribonucleic acid from heparin sperm	Sigma-Aldrich	Cat #: D-7290
1× Amplification Diluent	APExBIO	CAT#: H1001
Protector RNase Inhibitor	Roche	Cat #: 3335399001
5×Denhardt's solution	Sigma-Aldrich	Cat #: D2532
DL-Dithiothreitol	Sigma-Aldrich	Cat #: 43816
diethyl pyrocarbonate (DEPC)	Sigma-Aldrich	Cat #: D5758
Fluoromount-G	Southern Biotech	Cat #: 0100-01
Fluorescein Tyramide TSA plus fluorescein kit	APExBIO	Cat#: K1084
Cy3 TSA Fluorescence System Kit	APExBIO	Cat#: K1051
Experimental models: Organisms/strains		
C57BL/6JNifdc mice	Zhejiang Vital River Laboratory Animal	RRID: IMSR_JAX:000664

	Technology Co., Ltd., Zhejiang	
CB₁R-floxed	Biocytogen Pharmaceuticals Co., Ltd., Beijing	N/A
DAGLα-floxed	Biocytogen Pharmaceuticals Co., Ltd., Beijing	N/A
CB₁R-iCre-EGFP	Biocytogen Pharmaceuticals Co., Ltd., Beijing	N/A
Sprague-Dawley	Zhejiang Vital River Laboratory Animal Technology Co., Ltd., Zhejiang	RRID: RRRC_00239
Oligonucleotides		
Primers for *Cnr1* FISH: GTTCAAGGAGAACGAGGACAAC CAGCAGGCAGAGCATACTACAG	This study	N/A
Primers for *VGLUT2* FISH: CCAAATCTTACGGTGCTACCTC TAGCCATCTTTCCTGTTCCACT	This study	N/A
Primers for *GAD2* FISH: TCTTTTCTCCTGGTGGCG TTGAGAGGCGGCTCATTC	This study	N/A
Primers for *c-Fos* FISH: GCCCACCGACCTGCCTGCAA CTTTCGGTGGGCAGCGGCAC	This study	N/A
Primers for Genotyping of CB₁R-iCre-EGFP: GCCTGCACAAGCACGCCAAT TGAAGGATCGCCGAGCAACTG GGCACAAGCTGGAGTACAAC	This study	N/A
Primers for Genotyping of CB₁R-floxed:	This study	N/A

AGGCAACACTCAGTGACA GC TAGGTGGATCTCAGGAATC TTGTAG		
Primers for Genotyping of DAGLα-floxed: GGTGCTGATTGAGGCCTAC CTGT AGCCCTGGCTTGTTTACCC AA	This study	N/A

Software and Algorithms		
Python v3.8	Python Software Foundation	https://www.python.org/, RRID: SCR_008394
R studio Desktop	Posit Software, PBC	https://posit.co/download/rstudio-desktop/
R v4.3.1	R Core Team	https://www.r-project.org/, RRID: SCR_001905
R package 'networkD3' v0.4 [137]	Allaire J et al., 2017	https://CRAN.R-project.org/package=networkD3
R package 'tidyverse' v2.0.0 [138]	Wickham H et al., 2019	https://doi.org/10.21105/joss.01686
TissueFAXS Viewer	TissueGnostics	https://www.tissuegnostics.com/cn
Adobe Photoshop	Adobe Systems Inc	https://www.adobe.com/, RRID:SCR_014199
ZEISS Zen Microscopy Software	Zeiss	https://www.micro-shop.zeiss.com/, RRID: SCR_013672
ANY-maze video tracking software	Stoelting Co., IL, USA	https://www.any-maze.com/, RRID: SCR_014289
Graphpad Prism v8.0	GraphPad Software Inc	https://www.graphpad.com/scientificsoftware/prism/, RRID: SCR_002798
Aegisub v3.2.2	Interscape	https://download.html.it/software/aegisub/
OlyVIA v4.1	Olympus	https://www.olympus-lifescience.com.cn/zh/, RRID:SCR_016167
BehaviorAtlas Analyzer v1.0	Bayone BioTech Co., Ltd. Guangdong, China	https://bayonesci.com/
BehaviorAtlas Capture v1.0	Bayone BioTech Co., Ltd. Guangdong, China	https://bayonesci.com/

Automatic behavior-monitor system	**infrared (AIBM)**	Liu Q et al., 2021	https://doi.org/10.1007/s12264-021-00661-4
Fiji		Schindelin J et al., 2012	https://imagej.net/
NetworkX		Python module	https://networkx.org/, RRID：SCR_016864

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Feng Wang (fengwang@siat.ac.cn).

Materials availability

This study did not generate new unique reagents.

Data and code availability

○ The data supporting the current study's findings are available at Zenodo at https://zenodo.org/uploads/16351971 and are publicly available as of the date of publication. Original videos will be shared by the lead contact upon request.

○ All original code has been deposited at GitHub at https://github.com/Feng-Wang-Research-Group/A-neural-circuit-for-female-specific-defensive-homeostasis-in-risk-assessment and is publicly available as of the date of publication.

○ Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

All experimental procedures adhered to the guidelines established by the Chinese Council on Animal Care and were approved by the Institutional Animal Care and Use Committee (IACUC) of the Shenzhen Institutes of Advanced Technology (SIAT), Chinese Academy of Sciences (No. SIAT-IACUC-200319-NS-WF-A1162 and No. SIAT-BSI-IRB-240613-NS-WF-A0048). The study utilized animals virgin male and female C57BL/6J and mutant mice aged 8~16 weeks and virgin male Sprague-Dawley (SD) rats at an age of 12~20 weeks. All C57BL/6J mice and SD rats were obtained from Zhejiang Vital River Laboratory Animal Technology Co., Ltd (Zhejiang, China). CB_1R-floxed mice were generous gifts from department of Anesthesiology and Perioperative Medicine, Xijing Hospital, Air Force Medical University, Xi'an, China. The CB_1R-iCre-EGFP mouse line was generated as described previously[139]. DAGLα-floxed mice were generated specifically for this study by Biocytogen Pharmaceuticals Co., Ltd., Beijing, China in this study. Genotyping was performed using primers listed in key resources table with correct insertion verified through *in situ* hybridization targeting for gene RNA. Mice were housed in a controlled environment at 22–25°C on a 12 h light/12 h dark cycle with *ad libitum* access to food and water. Behavioral tests and recordings were conducted during the 8:00~16:00.

Viruses

(1) *In vivo* **photometry recording experiments**, 300 nL/site AAV-EF1α-DIO-GCaMP6s (2×10^{12} genomic copies/mL; packaged by Taitool Bioscience Co., Ltd., Shanghai) or AAV-EF1α-DIO-EGFP (3×10^{12} genomic

copies/mL; packaged by Fuqiang Xu's lab at the SIAT) was injected into the unilateral SC (AP -3.8 mm, ML -0.8 mm, DV -1.8 mm from bregma) of CB_1R-iCre-EGFP mice at a rate of 100 nL/min. 200 nL/site AAV9-GfaABC1D-eCB3.0 (5.46×10^{12} genomic copies/mL; Yulong Li's Lab at the Peking University) and AAV9-hSyn-RCaMP3 (6×10^{14} genomic copies/mL; packaged by Vigene Biosciences Co., Ltd., Shangdong) was infused into the unilateral LHb (AP -1.85 mm, ML -0.55 mm, DV -2.55 mm from bregma) of adult C57BL/6J mice at a rate of 70 nL/min. 50 nL/site AAV-EF1α-iGABASnFR was infused into the unilateral LHb (AP -1.85 mm, ML -0.55 mm, DV -2.55 mm from bregma) of CB_1R-floxed mice at a rate of 70 nL/min.

(2) ***CB_1R conditional knockout in Dlx5/6+ neurons and control experiments***, 300 nL/site AAV-Dlx5/6-GFP or AAV-Dlx5/6-Cre-GFP (5×10^{12} genomic copies/mL; packaged by Liping Wang's Lab at SIAT) was injected into the bilateral SC (AP -3.8 mm, ML ±0.8 mm, DV -1.8 mm from bregma) at a rate of 100 nL/min. Cre recombination was successful in the target neurons, resulting Cre recombinase expression and the absence of CB_1R mRNA. (3) **Optogenetic manipulation of SC CB_1R^+ Dlx5/6+ projections to LHb, ZI, or LP**, 300 nL/site Cre dependent recombinant AAV constructs containing hChR2(H134R)-mCherry (AAV2/9-MDLX-DIO-hChR2(H134R)-mCherry-WPRE-hGH pA; 5×10^{12} genomic copies/ml; packaged by BrainVTA Co., Ltd., Wuhan) or control viruses (AAV2/9-Dlx5/6-DIO-mCherry-WPRE-hGH pA; 5×10^{12} genomic copies/ml; packaged by BrainVTA Co., Ltd., Wuhan) were injected into the unilateral SC (AP -3.8 mm, ML -0.8 mm, DV -1.8 mm from bregma) of CB_1R-iCre-EGFP mice at a rate

of 100 nL/min. (4) **Confirming SC CB$_1$R$^+$ terminals in the LHb**, 300 nL/site AAV-Dlx5/6-DIO-mCherry and AAV-mDlx5/6-FLEX-tdTomato-T2A-Synaptophysin-EGFP (packaged by Brain Case Co., Ltd., Shenzhen, title: 5×10^{12} genomic copies/mL) were injected into the unilateral SC of CB$_1$R-iCre-EGFP mice at a rate of 100 nL/min, respectively. (5) **Discriminating CB$_1$R$^+$ and CB$_1$R$^-$ projections in SC Dlx5/6$^+$ neurons**, a 1:1 mixture of viruses AAV2/9-Dlx5/6-Cre on-ChR2-mCherry and AAV2/9-Dlx5/6-Cre out-ChR2-YFP (packaged by Shanghai Sunbio Medical Biotechnology Co., Ltd., Shanghai, title: 5×10^{12} genomic copies/mL) was injected into the SC of CB$_1$R-iCre-EGFP mice at a rate of 100 nL/min. (6) **Chemogenetic manipulation of LHb CB$_1$R$^+$ GABAergic projections from the SC**, 300 nL/site Cre dependent recombinant AAV constructs containing hM4D(Gi)-mCherry (rAAV-mDlx5/6-DIO-hM4D(Gi)-mCherry-WPRE-hGH pA; 5×10^{12} genomic copies/ml; packaged by BrainVTA Co., Ltd., Wuhan) or control viruses (AAV2/9-Dlx5/6-DIO-mCherry-WPRE-hGH pA; 5×10^{12} genomic copies/ml; packaged by BrainVTA Co., Ltd., WuhanLtd., Wuhan) were injected into the bilateral SC of CB$_1$R-iCre-EGFP mice at a rate of 100 nL/min. (7) **DAGLα knock-out and control experiments**, 50 nL/site AAV-EF1α-GFP or AAV-EF1α-Cre-GFP (5×10^{12} genomic copies/mL; packaged by Liping Wang's Lab at SIAT) was injected into the bilateral LHb (AP -1.85 mm, ML ±0.50 mm, DV -2.75 mm from bregma) at a rate of 70 nL/min.

METHOD DETAILS

Experimental design and randomization

For all between-subject analysis, mice were randomly assigned to experimental conditions using cage-level randomization (2-3 littermates split between control and experimental groups) to account for litter effects while minimizing cage-related variance. Experimental replication was achieved through cohort stratification, with conditional CB_1R knock out models requiring ≥ 3 batches (n=3-6/ batches) and other behavioral and assays experiments requiring ≥ 2 batches. For 3D behavior analyses, conditions were randomized, and experimenters remained blinded to group assignments until statistical was completed. Spatial *FISH* experiments employed triple-masking protocols during both experimental execution and computational processing. In neurocircuitry mapping, fiber and cannula placements experiments, histological verification was performed to assess viral transduction efficiency and implant localization. Subjects with low transduction efficiency $<50\%$ or deviation from target coordinated were systematically excluded prior to data processing.

Stereotaxic surgeries

Stereotaxic injection was performed following the protocol outlined in our published work[140]. Mice were anesthetized with pentobarbital (i.p., 80 mg/kg) and positioned on a stereotaxic frame (RWD, Shenzhen, China). Erythromycin eye ointment was applied to protect the eyes. The scalp was shaved, and a ~1 cm incision was made to expose the cranium. Connective and muscle tissue were carefully removed, and the skull surface was leveled to ensure the height difference between bregma and lambda was

< 0.05 mm and aligned in the medial-lateral plane. The skull above the target brain region was thinned using a dental drill and carefully removed. Viral injections were performed using a 10-μL Hamilton syringe connected to a 33-gauge needle and a microsyringe pump (UMP3/Micro4, World Precision Instruments, Sarasota, USA). The syringe was filled with mineral oil prior to use. At the injection, the needle remained in place for approximately 5 min to prevent fluid from reflux. Following surgery and a recovery period of at least 7 days, behavioral tests were conducted.

Cannula implantation

For chemogenetics manipulation and pharmacological experiments, a stainless-steel bilateral cannula (O.D.0.41 mm×I.D.0.25 mm, c.c 1.0 mm, C 2.8 mm, G1 0.5 mm, G2 0.5 mm, RWD Life Science, Shenzhen, China) was implanted into the LHb. For pharmacological experiment, a bilateral cannula (O.D.0.48 mm×I.D.0.34 mm, c.c 2.4 mm, C 4.5 mm, G1 0.5 mm, G2 0.5 mm, RWD Life Science, Shenzhen, China) was implanted in the LHb (coordinates of AP: −1.85 mm, ML: ± 0.5 mm, DV: −2.55 mm), ZI (coordinates of AP: −2.03 mm, ML: ± 1.2 mm, DV: −4.4 mm) or a singular cannula (0.48 mm in diameter, 5 mm in length, RWD Life Science, Shenzhen, China) was implanted in the Re (coordinates of AP: −0.59 mm, ML: 0.1 mm, DV: −4.2 mm). Behavioral tests were performed one week later after implantation.

Drug delivery

For chemogenetic inhibition of SC-LHb $CB_1R^+Dlx5/6^+$ projections in freely behaving mice, Clozapine N-oxide (CNO, Tocris, Cat #:4936) was prepared by dissolved it in

vehicle solution (sterile saline with 0.2% DMSO). Aliquots were frozen and freshly diluted with saline to a final concentration of 3 μg/μL before bilaterally administrated into the LHb via a bilateral guide cannula. The cannula was connected to two Hamilton syringes via a polyethylene catheter. For each side of the LHb, 50 nL of CNO or vehicle solution was infused slowly at a rate of 70 nL/min using two syringe pumps. After infusion, mice were habituated for additional 5 min to allow CNO diffusion before the needle was withdrawn. Animals were observed for 30 min prior to behavioral tests.

For pharmacological experiments, the neutral CB_1R antagonist NESS0327 (Cayman Chemical, 10004184, Michigan, USA) was locally administered to the LHb via a guide cannula. NESS0327 was suspended in saline containing 0.2% Tween 80. On the day of the experiment, a stainless-steel injection cannula (o.d.:0.41; i.d.:0.25, RWD, Shenzhen, China) was inserted into the LHb. The injection cannula, connected to a micro-injection system housing a 10 μL Hamilton syringe via flexible polyethylene tubing, extended 0.4 mm beyond the guide cannula tip. NESS0327 (1 μM/L) was bilateral injected in the LHb at a volume of 50 nL/site. Animals were observed for 3 min before undergoing behavioral tests.

Histology

Fluorescence In Situ Hybridization (FISH)

FISH was carried out following a protocol similar to that previously described[140]. Complementary DNAs (cDNAs) of *CB_1R*, *GAD2*, *VGLUT2*, or *c-Fos* (sequences from Allen Brain Atlas: CB_1R, NCBI Accession: NM_007726.2; GAD2, NCBI

Accession: NM_008078.1; VGLUT2, NCBI Accession: NM_080853.2; c-Fos, NCBI

Accession: NM_010234.2) were cloned from a mouse cDNA library into a

pBluescript SK cloning vector. Sequencing was performed to verify the accuracy of

the cloned sequences. Antisense complementary RNA (cRNA) probes were

synthesized by *in vitro* transcription and labeled with digoxigenin. The following

primer sequences were used for cloning of CB_1R, GAD2, VGLUT2, and c-Fos:

CB_1R (forward/reverse):

GTTCAAGGAGAACGAGGACAAC/CAGCAGGCAGAGCATACTACAG

GAD2 (forward/reverse):

TCTTTTCTCCTGGTGGCG/TTGAGAGGCGGCTCATTC

VGLUT2 (forward/reverse):

CCAAATCTTACGGTGCTACCTC/TAGCCATCTTTCCTGTTCCACT

c-Fos (forward/reverse):

GCCCACCGACCTGCCTGCAA/CTTTCGGTGGGCAGCGGCAC

Spatial FISH

The specific probes for target RNA (Table S2) were designed by Spatial FISH Ltd

(Shenzhen, China). SC sections from adult male (n=3) and female (n=3) C57BL/6J

mice were fixed by 4% paraformaldehyde (PFA) and placed within a reaction

chamber for subsequent processing. After dehydration and denaturation with methanol,

hybridization buffer containing the specific probes were added to the chamber and

incubated overnight at 37°C. The samples were then washed three times with PBST,

followed by ligation of the target probes in a ligation mix at 25°C for 3 h. After

another three washes with PBST, rolling circle amplification was performed using Phi29 DNA polymerase at 30°C overnight. Fluorescent detection probes in hybridization buffer were then applied to the samples. Finally, the samples were dehydrated using an ethanol series and mounted with mounting medium. Images were captured using Leica THUNDER Imaging Systems, and signal dots were decoded to analyze RNA spatial position.

Immunohistochemistry

Mice were perfused with 20 ml of 1×PBS followed by 20 ml of 4% PFA. Brains were harvested, post-fixed in 4% PFA overnight at 4°C, and dehydrated in 30% sucrose for 48 hours. Using a cryostat (CM1950, Leica, Wetzlar, Germany), 40 μm sections were prepared. Free-floating brain sections were rinsed with PBS (3×5 min), PBST (0.3% Triton X-100 in PBS, 1×20 minutes), and then blocked with 3% normal donkey serum (Solarbio, Cat #:SL050) at room temperature. Sections were incubated overnight at 4°C with primary antibody, including rabbit anti-RFP (1:500 dilution, Rockland, Cat #:600-401-379), rabbit anti-Cre (1:500 dilution, Abcam, Cat #: ab190177), or goat anti-GFP (1:500 dilution, Rockland, Cat #: 600-101-215). After washing in PBST (3×5 min), sections were incubated at room temperature for 1 h with secondary antibodies: donkey anti-rabbit 594 antibody (1:500 dilution, Jackson ImmunoResearch, Cat #:711-585-152), donkey anti-rabbit 488 antibody (1:500 dilution, Jackson ImmunoResearch, Cat #:711-547-003) or donkey anti-goat 488 antibody (1:500 dilution, Jackson ImmunoResearch, Cat #:705-545-003). This was followed by three 10-min washes in 0.01 mol/L PBS. Finally, sections were mounted

and covered with the anti-fade reagent DAPI (1:5000 dilution, Sigma-Aldrich, Cat #: D9542).

Microscopy Imaging and Quantification of Input Neurons

Virus-labeled signals and immunofluorescence-positive cells were traced using an Olympus VS120 or VS200 microscope equipped with a 10× objective and analyzed with Adobe Photoshop software (Adobe, San Jose, USA). For classical image presentation, sections were photographed using a Carl Zeiss ApoTome.2 microscope (Zeiss, Oberkochen, Germany) and processed with Zen software (Zeiss) or TissueFAXS Imaging software (TissueGenostics, Austria) using a 20× objective with tiled imaging (stitched from multiple images). Cell counting was performed manually on every third 40 μm thick section, ensuring a 120 μm intervals between successive sections in the coronal plane. Brain regions were identified based on *The Mouse Brain* (Stereotaxic Coordinates, fourth edition, Franklin, 2013).

Axon projection mapping

To analyze the projection patterns of SC CB_1R^+ $Dlx5/6^+$ neurons, every third brain sections (40μm thick) spanning the whole brain was collected. Sections were fully washed with PBS, mounted, and cover-slipped using 50% glycerol. Images were captured using an Olympus microscopy with a 10× objective (Olympus VS200). Brain regions were identified using *The Mouse Brain* (Stereotaxic Coordinates, fourth edition, Franklin, 2013), and projection patterns were analyzed using ImageJ (ImageJ, RRID: SCR_003070). In detail, for each animal, all images were thresholded to remove background noise. Then, 2-5 regions of interest (ROIs) were selected for each

brain region, and the average pixel value of each ROI were calculated after background subtraction (fluorescence density excluding fiber-free regions). For each animal, the average fluorescence density of a brain region was determined by averaging the values across all ROIs. Relative density per region was expressed as a fraction of the total density (summed across all regions). Data processing was conducted using the tidyverse R package (v2.0.0), while interactive forced-directed graphs of relative density averages across neuroanatomical regions were generated using the networkD3 package (v0.4).

Behavioral assays

Looming behavior

To assess the defensive behavior of male and female mice in response to life-threatening stimuli, we performed 2D looming behavioral tests using an innately aversive overhead expanding spot (looming task, as previously described[141]). On Day 1, animals were acclimated to the test arena for 10 min. The arena consisted of an open-top acrylic cylinder ("open field", 50-cm diameter) connected to an adjacent alley ("refuge", 50 cm×10 cm), allowing free access between compartments. The entire setup was enclosed by a 30-cm high wall (Figure 1B). On Day 2, mice were allowed 5 minutes of free exploration in the open-field arena before being subjected to 1-2 looming trials, depending on the timing accuracy of the first-stimulus, with a 3-minute inter-stimulus interval. The looming stimuli was triggered automatically when mice entered a concentric circular zone (central area: 25 cm diameter) within the arena periphery. The looming stimuli consist of a dark disk expanding from 2° to

40° within 300 ms, maintaining this size for 50 ms before disappearing, and repeating 15 times at 30-ms intervals.

For the SC CB_1R knock-out experiments, a 3D looming behavioral tests was assessed using multi-view motion-capture device system, as previously described[30]. A transparent looming arena with the same parameter as the 2D behavioral tests was placed at the center of a supporting framework, equipped with four depth cameras (Inter, RealSense D435i) connected to the BehaviorAltas Capture system (version 1.01, Shenzhen Bayone Biotech Co.), ensuring full-view motion capture. Two monitors, mounted on a fixed shelf 45 cm above the bottom of the looming box, delivered uniform and stable white light for behavioral tests. Light intensity was set to 120 lux at the center of the circular trigger zone and 30 lux at the refuge midpoint, matching the brightness parameters of 2D looming behavior setup. Behavioral recordings lasted 15 min, capturing 30 frames per second. matching the brightness parameters of 2D looming behavior setup. The other experimental parameters were consistent with those used in the 2D tests.

Open field test (OFT) in the multi-view motion-capture device

The open field apparatus (50×50×50 cm, matte white polycarbonate) was calibrated to access anxiety-like behavior and locomotor activity. To collect unobstructed behavioral data for elaborate analysis, a multi-view video capture device was employed to record spontaneous behavior (Figure 6A)[51]. Briefly, a 50×50 cm transparent acrylic square enclosure with white ground was placed at the center of a support framework either 130×130×90 cm (Figure 4L) or a 90×90×75 cm (Figure 6).

Four cameras were rigidly mounted on structural support pillars, with pan-tilt heads finely adjusted to achieve full coverage of the arena. Behavioral recordings were synchronized (30 Hz, 640×480 or 960×540-pixel resolution) and processed using BehaviorAtlas Capture Software® v2.0.0.

3D Behavioral data processing

Data acquired from the BehaviorAtlas Capture System underwent three sequential stages as previously described[51]: unsupervised movement annotation, supervised label refinement with manual correction of misclassified movement labels, and ethological quantification.

Unsupervised movement annotation pipeline

Using pre-training body point data from images, the skeleton of each mouse included in the study was automatically reconstructed by BehaviorAltas Analyzer software, incorporating camera calibration information. Movements were classified by clustering skeleton kinematic features for each frame and assigning digital labels.

Supervised movement labeling

Movements with postural similarities, particularly those occurring in low-light refuge areas, were subjected to dual-criteria validation with skeleton kinematic parameters such as speed, head-to-limb angle, body length, and distances between the nose and limbs. Specific movements like grooming, pausing, and looking up were systematically reviewed and reclassified when discrepancies arose between 3D kinematic data and manual annotations (see below). Additionally, movements such as flight and freezing were manually annotated by experts due to significant

inter-individual variability in kinematic patterns.

Kinematic reclassification protocol

Movement	Threshold
Pausing	Average velocity (based on back coordinate) < 30 mm/s and the movement duration > 2 s
Hunching	Average nose height ranges from 30 to 45 mm and average back height > 25 mm
Looking up	Average nose height > 20mm and average front claw height < 15 mm
Right turning	Average velocity (based on back coordinate) > 50 mm/s and body angle from top-view (neck-back-root tail) >15°
Grooming	Average body angle in 3D space (neck-back-root tail) < 120°
Rotating	Average velocity (based on back coordinate) > 30 mm/s and body angle from top-view (neck-back-root tail) >20°

Ethograms

Four key defensive behavior metrics, flight probability, response latency, return time, and time in the refuge, were quantified through manual frame-by-frame analysis (Figure 2D-E). Hierarchical ethograms were constructed by segmenting movement fragments within specific time windows (Figure 2H-I). Trajectory analysis was performed to quantify mouse locomotion by tracking the coordinates of the mouse's back, integrating spatial and temporal movement parameters. Defensive behavior quantification curves were created by mapping three stereotypic responses, risk-assessment posturing, flight initiation, freezing duration, against threat-exposure timeline, with group-averaged movement probability on the y-axis and time on the x-axis. Peak response markers indicated behavior-specific latency (Figure 2L-M). Defensive behavior analysis also incorporated nasal elevation dynamics and locomotor velocity, derived from 3D pose estimation outputs (Figure 2N-O).

Sample classification and segregation

Movement classification was performed using a random forest classifier with PCA-reduced kinematic features, encompassing all movements and their frequencies. Random forest machine learning algorithms were applied to movement data for group classification, with accuracy evaluated through detailed confusion matrices.

Movement transition and central node definition

Movement transitions were analyzed using probability matrices that captured how often one movement followed another, disregarding the duration of each movement. Line thickness in the diagrams represented the frequency of transitions, while dot size indicated the likelihood of specific movement events (Figure S6I-L). Transitions with probabilities below 0.05, after statistical adjustments, were excluded. Network diagrams were generated using Python's NetworkX package (v2.8.4) with spring layout settings.

Elevated plus maze (EPM)

The EPM test was performed a using maze made of white acrylic glass, comprising of a central platform (5×5 cm), two opposing open arms (32.5×5 cm), and two opposing closed arms (32.5×5×32.5 cm). The maze was elevated 50 cm above the floor. Mice were placed at the junction of the open and closed arms, facing the closed arms. Behavioral tracking (5-minute sessions) was conducted using ANY-maze® v7.4 software, with body centroid tracking. Time spent in each area (center, open arms, closed arms) was quantified.

Between behavior trials, the arena was cleaned by scrubbing with a 25% ethanol

solution.

Chronic rat stress model

The experimental protocol was modified from previous studies[142,143]. In the chronic stress model, mice were exposed to a rat for 10 consecutive days. Each day, a mouse was placed in a cage containing a rat for 5 seconds and then co-housed with the rat using a ventilated plexiglass divider. This apparatus allowed odor transmission while preventing physical contact, ensuring continuous exposure to predator odors. Behavioral tests, including the open-field test and elevated-plus maze, were conducted at two time points: 3 days prior to the establishment of the model (Pre-stress) and on day 10 post-exposure to rat-induced stress (Post-stress).

Patch-clamp electrophysiology

Coronal brain slices (300 μm thick) containing the SC region (bregma 3.2–4.4 mm) were prepared from CB_1R-iCre-EGFP mice using standard procedures. Brains were quickly separated and chilled in ice-cold modified artificial cerebrospinal fluid (ACSF) containing (in mM): 110 Choline Chloride, 2.5 KCl, 1.3 NaH_2PO_4, 25 $NaHCO_3$, 1.3 Na-Ascorbate, 0.6 Na-Pyruvate, 10 Glucose, 2 $CaCl_2$, 1.3 $MgCl_2$. SC slices were sectioned in ice-cold modified ACSF using a Leica vibroslicer (VT-1200S). Following sectioning, slices were allowed to recover for 30 min in a storage chamber containing regular ACSF at 32~34 °C (in mM): 125 NaCl, 2.5 KCl, 1.3 NaH_2PO_4, 25 NaHCO3, 1.3 Na-Ascorbate, 0.6 Na-Pyruvate, 10 Glucose, 2 $CaCl_2$, 1.3 $MgCl_2$ (pH 7.3~7.4 when saturated with 95% O2/5% CO2). Slices were then maintained at ~25 °C until

transferred to the recording chamber. All solutions had an osmolarity of 300~320 mOsm/kg. For electrophysiological experiments, slices were visualized using infrared optics on an upright microscope (Eclipse FN1, Nikon Instruments) with a 40× water-immersion objective. The recording chamber was continuously perfused with oxygenated ACSF (2 ml/min) at 34 °C. Patch pipettes were pulled by a micropipette puller (Sutter P-2000 Micropipette Puller) and had a resistance of 3–5MΩ. Electrodes were filled with intracellular solution containing (in mM): 130 potassium gluconate, 1 EGTA, 10 NaCl, 10 HEPES, 2 $MgCl_2$, 0.133 CaCl2, 3.5Mg-ATP, 1 Na-GTP. In SC slices, mCherry-positive neurons were patched in current-clamp mode. Action potentials were elicited by stimulation SC neurons expressing ChR2 with blue light pulse (10 ms duration, 20 Hz).

Optical fiber implantation

Fifteen min after AAV injection, a ceramic ferrule with an optical fiber (200 μm in diameter, N.A. 0.37) was implanted, positioning the optical fiber tip above the LHb (AP: -1.90 mm, ML: ±0.50 mm, DV: -2.55 mm from bregma), ZI (AP: -1.79 mm, ML: +1.10 mm, DV: -4.3 mm from bregma), or LP (AP: -2.03 mm, ML: +1.20 mm, DV: -2.2 mm from bregma). Optogenetic stimulation was performed 4 weeks after fiber implantation.

Three weeks after AAV injection, a ceramic ferrule with an optical fiber (200 μm diameter, N.A. 0.37) was implanted, positioning the fiber tip above the SC (AP: -3.8 mm, ML: -0.8 mm, DV: -1.5 mm from bregma), or LHb (AP: -1.90 mm, ML: ±0.50 mm, DV: -2.45 mm from bregma). Fiber photometry recording was conducted one

week after fiber implantation.

Optogenetics

Optogenetic manipulations were performed using laser systems with wavelength-specific settings. Terminal activation in SC-LHb $CB_1R^+Dlx5/6^+$ experiments: a blue laser (473 nm; Aurora-300-470, NEWDOON, Hangzhou) was used to deliver 10 mW/mm² at the fiber tip, with 5-s illumination, 20 Hz frequency, and 10-ms pulse duration.

Fiber photometry

(1) Fluorescence signals of GCaMP6s and GABASnFR were recorded via fiber photometry as previously described[142], using 470-nm LEDs for target excitation and 410-nm LEDs for isosbestic control. The 410-nm light intensity was calibrated to 20 µW/mm² at the fiber tip to match the 470-nm intensity. (2) Fluorescence signals of eCB3.0 and RCaMP3 were recorded via fiber photometry (vista, Inper), using 470-nm, 560 -nm LEDs for target excitation and 410-nm LEDs for isosbestic control. The 410-nm light intensity (0.5 mW/mm²), 470-nm light intensity (0.8 mW/mm²), 561-nm light intensity (1.2 mW/mm²), was calibrated to at the fiber tip. Behavioral video synchronization was achieved through dedicated system software. Raw data were smoothed using a 5-25-point moving average variance, with variance analyzed across different experiments. ΔF/F, areas under the curve, and peak values were calculated using least-squares regression to scale the 410-nm to 470-nm or 560-nm traces.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistics

For all experiments, statistical analyses were performed using either unpaired two-tailed Student's t test or a two-way ANOVA followed by Holm-Sidak's multiple comparisons test. The statistical significance threshold was set at $\alpha=0.05$, with significance levels indicated as $*p < 0.05$, $**p < 0.01$, $*** p < 0.001$, $**** p < 0.0001$). Exact p-values are provided in the figure legends and Table S1. The "n" in these analyses represents the number of mice, cells, or brain sections. Statistical analyses were carried out using GraphPad Prism 10 (GraphPad Software Inc.). Additionally, a custom Python script was utilized to evaluate the differences in movement transition and postures through a permutation test (10,000 iterations), comparing mean disparities between the two groups. Comprehensive statistical information is available in the figure legends and Table S1.

Figure S1. CB₁R⁺ cells characterization in male and female brains, related

to Figure 1

(A)-(I) Diagram representation showing the colocalization of CB₁R mRNA (red) and Cre protein (green) positive cells in various brain regions of a female mouse, including the MOB (A), mPFC (B), MS (C), HIP (D), CPu (E), Amyg (F), VMH (G), MHb (H), PVT (H), SC (I), PAG (I), Cere (J). Scale bar, 200 μm and 50 μm (enlarged view).

(K) Statistical analysis illustrating the percentage of overlap between CB₁R

mRNA (red) and Cre protein (green) positive cells in the mouse brain. Data are presented as mean ± SEM (n = 6 per group). Two-Way ANOVA followed by Sidak's multiple comparisons post hoc test was performed. *$p < 0.05$. For detailed p values, refer to Table S1.

Abbreviations: MOB, main olfactory bulb; mPFC, medial prefrontal cortex; MS, medial septal nucleus; HIP, hippocampus; CPu, caudate putamen (striatum); Amyg, amygdala; VMH, ventromedial hypothalamus; MHb, medial habenular nucleus; PVT, paraventricular thalamic nucleus; SC, superior colliculus; PAG, periaqueductal gray; Cere, cerebellum.

Figure S2. SC neurons majority are CB₁R⁺ without sex differences, related to Figure 1

(A) Diagram showing the colocalization of CB₁R mRNA (red) and Cre (green) positive cell in the SC.

(B) Pie chart illustrating the proportions of *Cnr1⁺* and *Cnr1⁻* cells in the SC for male (top) and female (bottom) mice. Unpaired Student's t test with the Mann-Whitney test was performed.

(C) Representative fluorescence *in situ* hybridization images showing CB₁R⁺

Cre expression in GAD2 neurons in the SC (Green, Cre; Red, GAD2 mRNA). Scale bar, 500 μm, and 50 μm (enlarged view). White arrows indicate the GAD2 mRNA fluorescent signals co-labeled with CB_1R positive Cre signals in each panel.

(D) Percentage of $GAD2^+$ and CB_1R^+ GAD2 neurons in males and females. Two-Way ANOVA followed by Sidak's multiple comparisons post hoc test was performed.

Data are presented as means ± SEM, n = 4 per group (B), 5 per group (D). ns (not significant). For detailed p values, refer to Table S1.

Figure S3. Visual threat activates SC CB$_1$R$^+$ neurons, related to **Figure 1**

(A) Timeline of the experimental protocol for looming stimuli and a schematic representation of the paradigm using a refuge-containing open field apparatus.

(B-C) Representative images showing c-Fos mRNA expression in the SC under the control condition (B) and after the looming test (C) (red, c-Fos mRNA; green, Cre protein; scale bar, 500 μm and 50 μm (enlarged view). White arrows indicate c-Fos mRNA fluorescent signals co-labeled with CB$_1$R-positive Cre signals in each panel.

(D-E) Quantification of c-Fos mRNA-labeled neurons and their colocalization with CB$_1$R-positive Cre signals in the SC of males (D) and females (E).

Two-Way ANOVA followed by Sidak's multiple comparisons post hoc test was performed.

(F) Percentage of CB_1R^+ c-Fos mRNA labeled neurons across the total c-Fos mRNA-labeled neurons following the looming test in males and females. Unpaired Student's t test with Mann-Whitney test was performed.

Data are presented as means ± SEM, n=6 per group (D-F). *$p < 0.05$; **$p < 0.01$. For detailed p values, refer to Table S1.

Figure S4. Deletion of CB₁R in SC CaMKIIα⁺ neurons suppressed defensive behavior in both sexes, related to **Figure 2**

(A) Schematic representation of viral injection of AAV-CaMKIIα-GFP or AAV-CaMKIIα-Cre-GFP into the bilateral SC of CB₁R-floxed mice.

(B) Left: Representative images showing AAV-CaMKIIα-GFP (top) and AAV-CaMKIIα-Cre-GFP (bottom) expression in the SC, alongside CB_1R mRNA (red) and CaMKIIα$^+$ cells bodies (green) in CaMKIIα-Ctrl and CaMKIIα-cKO group. Scale bar, 500 μm and 20 μm (enlarged view). White arrows indicate CB_1R mRNA fluorescent signals colocalized with GFP positive signals. Right: Quantification of CB_1R^+ cells in CaMKIIα$^+$ neurons in CaMKIIα-Ctrl and CaMKIIα-cKO groups for both sexes.

(C) Experimental timeline and schematic paradigm of virus injection and the looming test in a refuge-containing apparatus.

(D and E) Defensive strategy (left) and flight-to-refuge behaviors (from left to right: response latency, return time, time spent in the refuge) in male (D) and female (E) mice across CaMKIIα-Ctrl and CaMKIIα-cKO group.

(F) Sixteen movement types during defensive behavior, represented by corresponding colors.

(G) Schematic paradigm of the different stages of defensive behavior: exploring, escaping, hiding and threat-informed exploring.

(H and I) Ethograms showing the 16 movement types in males (H) and females (I) across groups (CaMKIIα-Ctrl group on the left, CaMKIIα-cKO group on the right). Scale bar, 1 s.

(J-K) Trajectories (left and middle) and quantification of fractions (right) of mice during the escaping stage (from the onset of looming to the end of flight, J, K) and hiding in the refuge (L, M) in males (J, L) and females (K, M) across

CaMKIIα-Ctrl and CaMKIIα-cKO groups. Histograms show the percentage of mice occupying horizontal and vertical positions in the apparatus. Symbols indicate mouse locations: five-pointed star for looming onset, triangle for flight initiation, and circle for flight termination.

(N and O) Distribution and positions of mice during flight, looking up, and freezing from 5 s pre-looming to 15 s post-looming in males (N) and females (O) across groups (CaMKIIα-Ctrl on the left, CaMKIIα-cKO on the right).

(P and Q) Left: Real-time speed and nose height from 5 s pre-looming to 15 s post-looming in males (P) and females (Q) across groups. Right: Quantification of the time taken to reach maximum speed and mean nose height during looming stimuli in males (P) and females (Q).

Data are presented as mean ± SEM (B, D, E, P, Q) or mean (J-O). n=4 per group (B); 30 male CaMKIIα-Ctrl, 27 male CaMKIIα-cKO (D), 15 female CaMKIIα-Ctrl, 12 female CaMKIIα-cKO (E); 26 male CaMKIIα-Ctrl and 15 CaMKIIα-cKO, 15 female CaMKIIα-Ctrl and 8 CaMKIIα-cKO (K-R). Two-Way ANOVA with Sidak's multiple comparisons post hoc test (J-M) and Unpaired Student's t test with Mann-Whitney test (B, D, E, P, Q) were performed. ns (not significant), * $p < 0.05$. For detailed p values, refer to Table S1.

Figure S5. Selective GABAergic targeting with Dlx5/6-driven AAVs, related to Figure 2

(A and B) Representative images showing SC Dlx5/6[+] GFP signals in GAD2 (A) and VGLUT2 (B) SC neurons. Scale bar, 20 μm.

(C) Percentage of SC Dlx5/6[+] neurons that are positive for GAD2 (A) or VGLUT2 (B) mRNA as determined by *ISH* in males and females. Data are presented as mean ± SEM.

(D) Representative image showing CaMKIIα[+] GFP signals in the VGLUT2 SC neurons (green, CaMKIIα[+] GFP signals; red, VGLUT2 mRNA; blue, DAPI; scale bars, 20 μm).

(E) Percentage of SC CaMKIIα[+] neurons that are positive for VGLUT2 (E) mRNA as determined by *ISH* in males and females.

Data are presented as mean ± SEM. n = 4 per group (C), Two-Way ANOVA

with Sidak's multiple comparisons post hoc test was performed, ****$p < 0.0001$.

n = 3 for males and n=4 for females (E). Unpaired Student's t test with the

Mann-Whitney test. ns (not significant), ****$p < 0.0001$. For detailed p values,

refer to Table S1.

Table S3. Parameters in the Dlx5/6 experiments during looming stimuli, related to Figure 2

Type	Sex	Group	Speed (mm/s) Mean±SEM	Nose height (mm) Mean±SEM	Back height (mm) Mean±SEM
Flight	Male	Ctrl (n=22)	153.4±10.49	21.09±2.313	25.21±1.049
		cKO (n=22)	149±11.72	16.02±2.975	26.57±0.8789
	Female	Ctrl (n=16)	133.9±17.23	21.09±2.313	25.26±0.9304
		cKO (n=16)	166.2±9.934	16.02±2.975	23.68±1.177
Non-Flight	Male	Ctrl (n=6)	52.09±18.88	27.39±7.068	28.7±1.113
		cKO (n=2)	96±9.378	22.38±6.334	26.71±3.041
	Female	Ctrl (n=4)	62.47±10.22	27.42±4.827	27.07±1.087
		cKO (n=5)	81.16±23.89	27.41±7.222	28.71±2.119

Movement	Definition
Flight	Mose locomotes with relatively high speed following looming stimuli
Running	Mose locomotes with relatively high speed
Trotting	Mose locomotes with a slow and intermittent pace
Walking	Mose locomotes with relatively low speed
Left turning	Mouse bends the body to the left or turning body to the left while walking
Right turning	Mouse bends the body to the right or turning the body to the right while walking
Stepping	Mouse takes a step forward with a short distance
Rotating	Mouse turns around with low speed and moves a short distance
Sniffing	Mouse twitches its nose to explore the environment at a relatively low speed
Looking up	Mouse Looks up with four legs on the ground
Rising	Mouse lifts its front legs but doesn't stand up
Hunching	Mouse stands on its hind legs while the dorsal is bent
Rearing	Mouse stands on hind legs with straight back
Grooming	Mouse licks, grooms its fur with the forepaws, or scratches its fur using any limb
Pausing	Mouse is still, crouches, or lies down on the floor without obvious horizontal displacement
Freezing	Mouse is completely absence of movement

Table S4. Definition of movements, related to Figure 2

Table S5. Parameters in the CaMKIIα experiments during looming stimuli,

Type	Sex	Group	Speed (mm/s) Mean±SEM	Nose height (mm) Mean±SEM	Back height (mm) Mean±SEM
Flight	Male	Ctrl (n=26)	150.2±9.45	15.33±2.036	22.54±0.8555
		cKO (n=15)	151.5±12.17	15.4±2.073	23.23±1.099
	Female	Ctrl (n=15)	151.3±15.02	16.36±3.532	21.54±1.256
		cKO (n=8)	142.4±16.97	16.34±2.449	22.53±1.21
Non-Flight	Male	Ctrl (n=4)	35.65±7.68	35.55±9.42	26.66±2.281
		cKO (n=12)	48.26±4.861	35.15±4.261	29.94±1.479
	Female	Ctrl (n=0)			
		cKO (n=4)	45.1±4.248	42.15±6.576	28.46±2.759

related to Figure 2

Figure S6. Deletion CB₁R in SC Dlx5/6⁺ neurons increases female sniffing during threat-informed exploration, related to **Figure 2**

(A and B) Trajectories (left and middle) and quantification of fractions (right) of mice during the pre-looming stage in males (A) and females (B) across different groups.

(C and D) Trajectories (left and middle) and quantification of fractions (right) of mice during the hiding stage in males (C) and females (D) across different groups.

(E and F) Trajectories (left and middle) and quantification of fractions (right) of mice during the threat-informed exploring stage in males (E) and females (F) across different groups. Data are presented as mean ± SEM, n=22 (E) male CB_1R Ctrl, 22 (E) male CB_1R cKO, 16 (F) female CB_1R Ctrl, and 16 (F) female CB_1R cKO. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed. $*p < 0.05$. For detailed p values, refer to Table S1.

(G and H) Behavioral features across different stages (from left to right, pre-looming, escaping, hiding and threat-informed exploring) visualized by embedding 16-dimensional movement fractions into a low-dimensional space using principal component analysis (PCA) for males (G) and females (H).

(I-L) Movements transition probabilities across different stages (from left to right, pre-looming, escaping, hiding and threat-informed exploring) for the male CB_1R Ctrl (I), male CB_1R cKO (K), female CB_1R Ctrl (J), and female CB_1R cKO (L).

Each movement is represented as a state, with circle size indicating the number of movement bouts, where larger circles correspond to higher occurrence frequencies.

Figure S7. Flight-to-refuge deficits across groups following visual threats, related to Figure 2

(A and B) Ethograms depicting 16 movements from 60 s pre-looming to 60 s post-looming for males (A) and females (B) in the CB₁R Ctrl (top) and CB₁R cKO (bottom) group.

(C and D) Comparison of movement fractions during the pre-looming stage in males (C) and females (D) across different groups.

(E and F) Comparison of movement fractions during the looming stimuli stage in males (E) and females (F) across different groups.

(G and H) Comparison of movement fractions during the post-looming stage in males (C) and females (D) across different groups.

(I and J) Real-time speed and nose height data from 5 s pre-looming to 15 s post-looming for males (I) and females (J) across different groups.

Data are presented as mean ± SEM for C-H and as mean for I, J. n=6 male Ctrl group, 2 male CB_1R-KO group (C, E, G, I), 4 female CB_1R Ctrl group, 5 female CB_1R cKO group (D, F, H, J). For detailed data information, refer to Table S1.

Figure S8. Comparable downstream targeting of SC CB$_1$R$^+$Dlx5/6$^+$ neurons in males and females, related to Figure 3

(A) Schematic illustration of the surgical procedure for anterograde tracing of SC CB$_1$R$^+$ Dlx5/6$^+$ neurons.

(B) Representative image showing viral infection in SC CB$_1$R$^+$Dlx5/6$^+$ neurons

in a female mouse. Red, CB$_1$R$^+$Dlx5/6$^+$ starting cells; green, terminals of CB$_1$R$^+$Dlx5/6$^+$ cells. Scale bar, 500 µm.

(C-N) Representative image of various brain regions targeted by SC CB$_1$R$^+$Dlx5/6$^+$ neurons in a female mouse: Re (C), LH (C), LD (D), MD (D), PVT (E), ZI (F), PH (F), opt (F), LHb (G), PVG (H), PaF (H), APT (H), LP (H), LGN (H), PT (I), RPC (J), PP (J), CnF (K), MiTg (K), NLL (K), IC (L), GiA (L), CnF (M), mRt (M), and IOC (N).

Abbreviations: Re, reuniens thalamic nucleus; LH, lateral hypothalamus; LD, laterodorsal thalamus; MD, mediodorsal thalamus, PVT, paraventricular thalamus, LHb, lateral habenula; PVG, periventricular gray; PaF, parafascicular thalamic nucleus; APT, anterior pretectal nucleus, LP, lateral posterior thalamus; LGN, lateral geniculate nucleus; PH, posterior hypothalamus; ZI, zona incerta, opt, optic fiber; PT, pretectal nucleus (PT); RPC, red nucleus, parvicellular part; PP, peripeduncular nucleus; CnF, cuneiform nucleus; MiTg, microcellular tegmental nucleus; NLL, nucleus of the lateral lemniscus pontine reticular nucleus; IC, inferior colliculus, mRt, mesencephalic reticular format; GiA, gigantocellular reticular nucleus, alpha part; and IOC, inferior olive, subnucleus C of medial nucleus.

(O) Schematic representation showing outputs of SC CB$_1$R$^+$Dlx5/6$^+$ neurons.

(P) Sankey network showing projections of SC CB$_1$R$^+$Dlx5/6$^+$ neurons in male and female mice, with curve width representing connection strength.

(Q) Quantification of normalized fluorescence density in ipsilateral downstream

brain subregions targeted by SC $CB_1R^+Dlx5/6$ neurons in male and female mice.

Data are presented as mean ± SEM. n=3 each group. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed. For detailed *p* values, refer to Table S1.

Figure S9. SC CB$_1$R$^+$Dlx5/6$^+$ neurons dominate long-range projections, related to Figure 3

(A) Schematic representation of the virus tracing strategy.

(B) Representative fluorescent micrographs of a coronal brain slice showing the injection site in the SC (left) and an enlarged view of the dashed square area (right). Scale bars, 500 μm and 50 μm (enlarged view).

(C) Quantification of the percentage of CB_1R^+, CB_1R^-, and non-specific overlapped signals with mCherry and YFP fluorescence in Dlx5/6$^+$ starter neurons. Data are presented as mean ± SEM. n = 3 males, 4 females. Unpaired Student's t test with the Mann-Whitney test was performed. ****$p<$ 0.0001. For detailed p values, refer to Table S1.

(D-G) Representative images showing downstream projections of SC $CB_1R^+Dlx5/6^+$ neurons in various brain regions including the reuniens thalamic nucleus (Re), anterior hypothalamus (AH), lateral dorsal thalamus (LD), zona incerta (ZI), optic fiber (opt), lateral habenular (LHb), lateral posterior thalamus (LP), pregeniculate area (PrG) and the pontine reticular nucleus, oral part (PnO).

Figure S10. Retrograde tracing identifies CB$_1$R$^+$Dlx5/6$^+$ SC-LHb circuit, related to **Figure 3**

(A) Schematic representation of the AAV-Retro tracing strategy for SC-LHb projections using unilateral injection into the SC and LHb.

(B) Representative fluorescent micrographs showing Dlx5/6$^+$ mCherry expression in the LHb. Scale bar, 200 µm.

(C) Quantification of the CB$_1$R$^+$ SC-LHb population within Dlx5/6$^+$ neurons. Data are presented as mean ± SEM, n = 3 per group. Unpaired Student's t test with the Mann-Whitney test was performed. ns (not significant). For detailed *p* values, refer to Table S1.

(D and E) FISH analysis showing mCherry expression in the SC co-labeled with CB$_1$R mRNA signals in male (D) and female (E) mice. Scale bars, 500 µm and 50 µm (enlarged view).

Figure S11. CB₁R antagonism in Re or ZI lacks effect on defensive behavior in either sex, **related to Figure 3**

(A) Representative image showing the track cannula placement in the reuniens thalamic nucleus (Re).

(B) Schematic representation of the looming stimuli paradigm conducted in a refuge-containing open field apparatus.

(C and D) Metrics of defensive behavior (from left to right: response latency, return time, and time in the refuge after looming) for male (C) and female mice (D) in the vehicle (Veh) and NESS 0327 (NESS) groups.

(E) Representative image showing the track cannula in the zona incerta (ZI).

(F and G) Defensive behavior metrics (from left to right: response latency, return time, and time in the refuge after looming) for male (F) and female mice (G) in the Veh and NESS groups.

Data are presented as mean ± SEM. n=11 male Veh group, 12 male NESS group (C); 12 female Veh group, 11 female NESS group (D); 7 male Veh group,

8 male NESS group (F); 8 female Veh group, 7 female NESS group (G).

Unpaired Student's t test with the Mann-Whitney test was performed. For

detailed *p* values, refer to Table S1.

Figure S12. Activating CB$_1$R$^+$Dlx5/6$^+$ SC-ZI/LP terminals lacks effect on defensive behavior in either sex, related to **Figure 4**

(A and B) Viral strategy and histology of CB$_1$R-expressing SC-ZI Dlx5/6$^+$ terminals. The dotted box highlights the optic fiber track. Scale bar, 200 µm. SC, superior colliculus; ZI, zonal incerta.

(C and D) Statistic analysis of defensive behavior metrics, including time spent initiating a response, returning to the refuge, and remaining in the refuge after looming stimuli during opto-stimulation of SC-ZI$^{CB1R+ Dlx5/6+}$ terminals in males (C) and females (D).

(E and F) Viral strategy and histology analysis of the SC-LP$^{CB1R+Dlx5/6+}$ terminals. The dotted box highlights the optic fiber track. Scale bar, 200 µm. LP, lateral posterior thalamus.

(G and H) Statistic analysis of defensive behavior metrics, including the time

spent initiating a response, returning to the refuge, and remaining in the refuge after looming stimuli during opto-stimulation of SC $^{CB1R+Dlx5/6+}$ axons in the LP for males (G) and females (H).

Data are presented as mean ± SEM. n = 8 male mCherry group, 9 male ChR2 group (C); 7 female mCherry group, 10 female ChR2 group (D); 12 male mCherry group, 8 male ChR2 group (G), 10 female mCherry group, 11 female ChR2 group (H). Unpaired Student's t test with the Mann-Whitney test was performed. For detailed p values, refer to Table S1.

Table S6. Definition of movements and clusters, **related to Figure 5, Figure 7**

Clusters	Movement	Definition
Locomotion	Running	Mose locomotes with relatively high speed
	Trotting	Mose locomotes with a slow and intermittent pace
	Walking	Mose locomotes with relatively low speed
	Left turning	Mouse bends the body to the left or turning body to the left while walking
	Right turning	Mouse bends the body to the right or turning the body to the right while walking
	Stepping	Mouse takes a step forward with a short distance
Exploration	Sniffing	Mouse twitches its nose to explore the environment at a relatively low speed
	Looking up	Mouse Looks up with four legs on the ground
	Rising	Mouse lifts its front legs but doesn't stand up
	Hunching	Mouse stands on its hind legs while the dorsal is bent
	Rearing	Mouse stands on hind legs with straight back
	Jumping	Mouse jumps with its hind legs completely off the ground
Maintenance	Grooming	Mouse licks, grooms its fur with the forepaws, or scratches its fur using any limb
Nap	Pausing	Mouse is still, crouches, or lies down on the floor without obvious horizontal displacement

Figure S13. Chronic stress alerts spontaneous behavior dynamics, related to **Figure 6**

(A and B) Proportion of clusters during every 10-min epoch for male (A) and female (B) CB$_1$R Ctrl (top) and CB$_1$R cKO (bottom) groups. Data are presented

as mean ± SEM. Male CB_1R Ctrl (n=18 pre-stress, 17 post-stress); Male CB_1R cKO (n=18 pre-stress, 16 post-stress); Female CB_1R Ctrl (n=17 pre-stress, 20 post-stress); Female CB_1R cKO (n=17 pre-stress, 18 post-stress). #$p < 0.05$, ##$p < 0.01$, ###$p < 0.001$, ####$p < 0.001$ post-stress versus pre-stress in corresponding group. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed. For detailed p values, refer to Table S1.

(C and D) Comparison of movement transition probabilities for males (C) and females (D). From left to right, pre-stress (CB_1R cKO minus CB_1R Ctrl), post-stress (CB_1R cKO minus CB_1R Ctrl), CB_1R Ctrl group (post-stress minus pre-stress), CB_1R cKO group (post-stress minus pre-stress). Each movement bout is represented as a state.

(E and F) PCA visualization of behavioral features for males (G) and females (H) across different groups.

(G and H) Random forest classifier trained on 14-dimensional movement fractions and frequency to predict group labels for males (G) and females (H) across different groups.

Figure S14. Dynamics of spontaneous behavior over 60 min period, related to **Figure 6**

(A and B) Temporal fraction of each movement during pre- and post-stress stages for males (A) and females (B) in CB_1R Ctrl (top) and CB_1R cKO (bottom) groups. Data are presented as the mean normalized values over the time

course for each movement.

(C and D) Movement transition frequency during pre- and post-stress stages for males (C) and females (D) in CB_1R Ctrl and CB_1R cKO groups.

Data are presented as mean ± SEM. Sample sizes are consistent with those in Figure S13. #$p < 0.05$; ##$p < 0.01$; ###$p < 0.001$ post-stress versus pre-stress in corresponding group. Two-Way ANOVA with Sidak's multiple comparisons post hoc test was performed. For detailed p values, refer to Table S1.