Effects of partner workload and increasing environmental temperature on nestling

provisioning and body temperature in a declining aerial insectivore

Research Highlights

In adult male tree swallows, we found that experimental manipulations of heat dissipation

capacity had little effect on the provisioning rate, body temperature, nor growth of nestlings.

That males did not respond to enhanced heat dissipation suggests that they may provision below

the threshold of overheating. Rather, male provisioning was affected by the provisioning rate of

the male's partner and environmental temperature.

Megan C. Heft,[1]* Bronwen Hennigar[1], and Gary Burness[2]

Addresses:

[1]Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario,

Canada. Email: meganheft@trentu.ca

[2]Department of Biology, Trent University, Peterborough, Ontario, Canada. Email:

garyburness@trentu.ca; ORCID: 0000-0002-1695-7179

*Corresponding author: meganheft@trentu.ca

Keywords: heat dissipation limit theory, performance, bird**,** parental care, body temperature

Running title: Parental workload and environmental temperatures

Abstract:

With climate change, birds will face increasing thermoregulatory demands, which may alter

reproductive behaviors such as offspring provisioning. Experimental studies have shown that the

provisioning capacity of female tree swallows (*Tachycineta bicolor*) is limited by their risk of

overheating. Given that parental investment strategies may vary between sexes, the thermal

environment may have a different impact on males. We experimentally trimmed ventral feathers

from male tree swallows to create a "thermal window" through which they could dissipate heat.

We remotely monitored provisioning rate and core body temperature of males and their female

partners. At high temperatures, all males decreased their nestling provisioning rates irrespective

of trimming treatment. In addition, trimmed males maintained core body temperatures similar to

those of controls. This suggests that in contrast to previous work with females, males limit

provisioning rates to levels below the threshold at which they would overheat. Regardless of

male treatment, females adjusted their own activity to match that of their male partners; whether

there are costs to females is unknown. Combined, these studies highlight that sex-specific

differences in thermal physiology and behavior must be considered when predicting responses to

climatic warming.

Introduction

Climatic warming is predicted to result in an increase in extreme temperature and weather events (Sippel et al., 2020). In endotherms, prolonged exposure to high ambient temperatures has been associated with reduced reproductive success (e.g., Van de Ven et al., 2020; Pipoly et al., 2022, Levillan et al. 2025) and reports of mass mortality (Jones et al., 2018; Piatt et al., 2020; McKechnie et al., 2021). As thermal challenges are unlikely to subside given the current climatic estimates (IPCC, 2023), the extent to which individuals and populations will be able to effectively respond to thermal stressors is of growing concern (Altan et al., 2003; Heeter et al., 2023; Taff & Shipley, 2023).

Among endotherms, birds may be at particular risk of climatic warming given their relatively high body temperatures and elevated metabolic rates. When birds are faced with thermal stressors, they increase heat loss via evaporative or non-evaporative mechanisms (Tattersall et al., 2016) and/or allow body temperature (T_b) to increase which may result in hyperthermia (e.g., Tapper et al., 2020b; Freeman et al., 2022). In species that exhibit extensive parental care, high environmental temperatures may exacerbate the effects of heat generated by parents during flight and in other activities associated with provisioning offspring (Tapper et al., 2020a). In fact, an individual's rising body temperature and its inability to sufficiently dissipate metabolic heat generated during parental care may limit parental activity (heat-dissipation limit theory; Speakman & Król, 2010; Zagkle et al., 2022). The fitness benefits of increased provisioning rates must therefore be balanced against possible physiological damage induced by hyperthermia, including protein denaturation (Vazquez & Larson, 2013), depressed immune function (Alhenaky et al., 2017), and increased oxidative stress (Huang et al., 2015).

To date, a handful of avian field studies have explored experimentally how the risk of

overheating impacts the capacity of individuals to maintain performance during a breeding event.

For example, female tree swallows (*Tachycineta bicolor*) provided with an increased capacity to

dissipate heat via experimental trimming of feathers, provisioned their nestlings at higher rates

than their untrimmed counterparts, at least at highest temperatures (Tapper et al., 2020a). This

suggested that at high environmental temperatures, an individual's activity (and thereby capacity

to care for young) was determined by a thermal constraint, as predicted by the heat-dissipation

limit theory (Speakman & Król, 2010). In contrast, female blue tits (*Cyanistes caeruleus)* that

were trimmed of feathers did not increase provisioning, but instead displayed enhanced immune

function, suggesting a prioritization of self-maintenance (Nord & Nilsson, 2019; Andreasson et

al., 2020).

An individual's parental-investment strategy involves a trade-off between allocating

energy and resources between current and future reproductive attempts (Clutton-Brock, 1991;

Lemaître & Gaillard, 2017). In species that exhibit biparental care, thermal limits may play a role

in pair coordination, with possible effects on investment strategies. Recently, it was shown that

male tree swallows partnered with experimentally trimmed females increased their own

provisioning efforts at high environmental temperatures, despite not being trimmed themselves

(Tapper et al., 2020a). Male and female tree swallows are known the adjust their provisioning

efforts based on partner quality (Lendvai et al., 2018). However, if males were able to increase

their workload to better match that of their female partners, males were either working below

their physiological limit, or perhaps willing to endure physiological damage (e.g., Dick &

Guglielmo, 2019). In this sense, sex-specific investment strategies may emerge when males and

females are faced with the removal of a possible physiological constraint (Nilsson & Nord,

2018).

Considering that reproductive strategies can vary between sexes (Kappeler et al., 2023),

we investigated whether a male tree swallow's capacity to dissipate heat would alter its

reproductive investment, and we qualitatively compared our results with previous studies on

females (Tapper et al 2020a, 2020b). We experimentally trimmed ventral feathers from male tree

swallows to allow for enhanced heat loss and measured provisioning rates and body temperature

in trimmed individuals and compared them with untrimmed controls. Further, we explored the

response of their female partners (who were unmanipulated). We anticipated that the capacity to

dissipate metabolic heat would limit activity in adult male tree swallows (i.e., the heat dissipation

limit theory). We made the following predictions: (1) males with an augmented capacity to

dissipate heat (trimmed males) would alter their reproductive investment, and increase their

provisioning rate to nestlings across all environmental temperatures; (2) trimmed males would

maintain a similar body temperature as untrimmed controls, by adjusting their activity to

maintain T_b within a narrow range; (3) females paired with trimmed males would have higher

nestling provisioning rates than females paired with non-trimmed males, because the

provisioning rates of pairs are often coordinated (Lendvai et al., 2018); (4) nestlings from

trimmed males would reach a heavier mass than nestlings from control males.

Methods

Study species and general field methods

All research was approved by the Trent University Animal Care Committee, in accordance with the Canadian Council on Animal Care (Animal Use Protocol # 26534). We conducted this study from May-July 2021, on two nest-box breeding populations of tree swallows located at the Trent University Nature Areas, Peterborough, Ontario, Canada (44°21'N, 78°17'W) and at the Lakefield Sewage Lagoon, Lakefield, Ontario (44°24'58.3"N 78°15'26.8"W). The Nature Areas site contains ~40 boxes spread out in a grid across the site, while the Sewage Lagoon site consists of two large, central lagoons surrounded by a perimeter of ~80 nest boxes. Further details on the sites are found in Fischer et al. (2020). At our study sites, tree swallows begin constructing their nests in early May, typically rearing 1 brood per season. Clutch size varies between 3-7 eggs, laid approximately one day apart. Incubation typically lasts 14 days with nestlings hatching relatively synchronously. After hatch, the nestling period spans approximately 15-22 days with adults providing food and maintenance care until the nestlings fledge (Winkler et al., 2020).

We monitored nests boxes daily from May through July to determine the lay and hatch dates for each clutch. As each egg was laid, we labelled it with a marker to establish order of laying and incubation dates. Incubation day 0 was the date that the last egg was laid, and hatch day 0 was the date the majority of nestlings in a brood hatched.

Experimental manipulation of heat dissipation

When nestlings were between days 3-4 post-hatch, we captured adult male tree swallows and performed a feather-trimming manipulation. We assigned males randomly to either a Trimmed or Control treatment by flipping a coin (Trimmed: n=15, Control: n=20). Males in the Trimmed group had an approximately 3 x 2 cm area of ventral feathers trimmed with scissors, such that approximately 7% of body surface area was exposed, following Tapper et al. (2020a). To ensure the degree of trimming was consistent among individual males, we used a cardboard "stencil," placed over each individual's ventral surface. Males assigned to Control groups were handled for a similar length of time as the Trimmed individuals, but no feathers were removed.

Following trimming (or handling, in case of Controls), we implanted each male with either a temperature-sensitive passive integrated transponder (PIT) tag (accuracy ±0.5°C; Biotherm13; Biomark, Boise, Idaho, USA,) or attached a non-temperature-sensitive leg-mounted RFID tag (EM4100; #11001, GAO RFID, Ontario, 166 Canada). Tag type was randomized by coin flip. To implant a PIT tag, we first sterilized the incision site with ethanol, implanted the tag subcutaneously into the nape of the neck, and then sealed the incision site with tissue adhesive bond (Vetbond Tissue Adhesive, Fisher Scientific, Waltham, MA, USA). Prior to release we collected a 50-75 µL blood sample (as part of another study) and recorded body morphometrics (body mass, wing chord (flattened), and head-bill length). Handling time was < 10 minutes for each individual. Given limitations due to synchrony at hatch, loggers were rotated among nests and fewer birds were recorded than were initially trimmed and tagged. Sample size for temperature sensitive tagged males: Trimmed males: n= 4, Control males: n= 9. Sample size for RFID tagged males: Trimmed males: n= 9, Control males: n= 9.

To test female responsiveness to male trimming, we captured females at the same time as

males and equipped each female with the same tag type as their male mates received (i.e, females

paired with males that were equipped with temperature-sensitive tags were also assigned

temperature-sensitive tags). We collected a blood sample and recorded the same morphometrics

as for males; no females were trimmed (females paired with Trimmed males: n= 13; females

paired with Control males: n= 17). Individuals were then released. One female paired with a

Control male was tagged, but no nestbox visits were recorded while loggers were deployed. As a

result, the sample size for females is slightly lower than that of males.

Remote Monitoring

For all tagged adults we recorded nest-box visitation rates, while for individuals implanted with

temperature-sensitive tags we also recorded subcutaneous body temperature (T_b). We considered

T_b to be a proxy for core body temperature (McCafferty et al., 2015; Andreasson et al., 2023).

For individuals with temperature-sensitive tags, we recorded visitation rates and T_b using either a

Biomark HPR Plus or a Biomark Small Scale Monitoring System (AR650) reader. Because of a

limited number of Biomark readers, we cycled them among nests to capture adult visitation rates

during 2-3 day windows throughout nestling development (early, middle, and late from 2-5 days,

6-9 days, and 10-14 days post-hatch, respectively). Readers (connected to 12v batteries) were

placed at the base of each nest box and attached to a loop antenna mounted to the nest entrance.

Visits were recorded anytime an individual entered the nest or perched at the nest entrance. Both

types of Biomark readers were set to a 10 second 'delay' to limit tag detections when a bird spent

a prolonged period in the nest box (Tapper at al., 2020). Individual reads included the date, time,

and the bird identification code for a given tag (recorded when each tag was initially deployed).

To record visitation rates of individuals with leg mounted tags (rather than implanted

temperature-sensitive tags), we used Generation 2 RFID readers (Cellular Tracking Technology,

Rio Grande, NJ, USA) (Bonter and Bridge, 2011). Because we had a large number of Generation

2 RFID readers, we did not need to cycle them among nests. Nonetheless, to allow for

comparison with data from temperature sensitive tags, we blocked our non-temperature sensitive

detection data into 2-3 day windows to match the cycling of the Biomark readers.

Environmental Data

We collected hourly readings for environmental temperature (°C), relative humidity (%), and

windspeed (km ·hr-1), and total daily precipitation (mm) from the Trent University weather

station (1.5 km from the Trent University Nature Areas and 9.5 km from Lakefield Sewage

Lagoons). All environmental data were downloaded from Environment Canada (available at

https://climate.weather.gc.ca/index_e.html).

Nestling Morphometrics

Upon hatching of nestlings, we marked each nestling with a colored marker, allowing us to track

the growth rate of individuals (until they were large enough to band). We weighed nestlings

between the hours of ~1200 – 1800 on days 0 (hatch), 3, 6, 9 and 12 post-hatch on a digital

portable scale (Smart Weigh Digital Pro Pocket, ± 0.01 g). No measurements were recorded after

day 12 to prevent premature fledge. In our analysis, nestlings were assigned to either a Control (n

= 20) or Trimmed (n = 15) group based on the manipulation of the father, though no nestlings

were manipulated. Note that the sample sizes for nests of nestlings were greater than those for

adults used in provisioning rate analyses (above); due to logger limitations (cycling loggers

between boxes) four nests could not be included. However, males were still captured and

manipulated at these nests.

Data compilation

Nest box visitation rate

All data organization and analysis was performed in R (R version 4.2.2, R Core Team). To

quantify RFID reads (from both thermal and nonthermal tag types), we transformed raw RFID

reads into nest box visits using the 'visits' function from feedr (LaZerte et al., 2017; Tapper et

al., 2020a). We considered a nest box visit to be an index of parental provisioning. Subsequent

statistical analyses for parental provisioning visits were then completed using these transformed

data. Observations for adult provisioning visits were balanced within pairs (n=133 male

observations, n=144 female observations).

Because males were captured between days 3-4 of nestling development, we restricted

our analyses of provisioning visits to days 3-14 of nestling development. In addition, adults do

not typically feed nestlings between the hours of ~21:00 and 05:00 (Tapper et al., 2020a), so we

excluded this window from our provisioning rate data. In the case where successive RFID reads

occurred within 60 seconds (i.e., a bird was present in the box for multiple reads within the span

of 60 seconds), we defined this as a singular visit. In our analysis, provisioning rate was defined

as the average number of provisioning visits per hour of daytime activity (i.e., between 05:00 and

21:00 each day), or per hour that a logger was recording at a given nest box (if the logger was

removed partway through the day).

 Statistical analysis

 Provisioning rate

For our analyses of male and female provisioning rates, and male body temperature we evaluated

competing models with an Akaike Information Criterion corrected for small sample sizes

(AICc). In our analysis, we considered models with a ΔAICc of < 2 to be strongly supported

(Burnham & Anderson, 2004). For our provisioning rate and body temperature models, we

evaluated models based on model weights and likelihoods (Hurvich & Tsai, 1989).

For each of our feeding rate and body temperature analyses, we fit a series of candidate models

and report the number evaluated: male provisioning (n = 16), female provisioning (n = 16), and

male body temperature (n = 10). We also report a null model (intercept only) for reference. For

models with strong support, we report the 95% confidence intervals (CI) and associated effect

size for individual terms.

Statement on AI:

Artificial Intelligence (AI) was used during data analysis to refine visualizations and to suggest

corrections to code. Specifically, ChatGPT (OpenAI, 2025) was used to improve layouts, color

scheme, and labeling of figures, for clearer visualizations. All code was reviewed and checked

for accuracy.

Male provisioning rate

In our analysis of male provisioning rate, we were primarily interested in the response of males

to our experimental manipulation (trimming). To explore potential factors contributing to

variation in average number of visits per hour, we evaluated combinations of predictors

including manipulation (trimmed or control, i.e., untrimmed), tag type (thermal or nonthermal),

brood size (at hatch), clutch initiation date, average daily environmental temperature, relative

humidity and wind speed, and total daily precipitation. We evaluated environmental predictors

for collinearity by running a variance inflation factor test (all VIF <3; 'vif' function, Car

package, version 3.1-2; Fox and Weisberg, 2019). We also included the average number of visits

per hour for females (filtered from 05:00 to 21:00). Based on the nonlinear relation between male

provisioning rate and environmental temperature, we included a quadratic term for average daily

environmental temperature (T_a^2). Since previous work has shown that the effects of treatment

may vary based on environmental temperature (Tapper et al., 2020a), we also tested for an

interaction between average daily environmental temperature and manipulation (along with an

interaction between manipulation and the quadratic; T_a^2). To account for repeated measures from

the same individual over the brood rearing period, we included a random intercept for individual

male's identity.

Since thermal loggers were cycled among nests (sometimes being moved partway

through the morning) we favored an averaging approach for provisioning rates, where feeding

rate is defined as the total number of visits per day divided by the total number of hours logged

per day. As in previous work (Tapper et al., 2020a), the cycling of thermal loggers constained

our analysis of adult provisioning as rotations led to unequal observation effort at nest boxes of

birds equipped with thermal tags. While an hourly analysis was considered, we preferred the

average hourly provisioning rate, to account for unequal coverage and allow for comparisons

between tag types. However, we also conducted additional analysis of raw hourly provisioning

with focal hours captured using hourly provisioning rate and hourly environmental variables

(with the same models sets as described above for average hourly, to consider finer scale hourly

variation throughout the day). We provide the output of these analyses in supplementary table S1

and S2.

Male body temperature, T_b

We tested the impact of experimental trimming on male average hourly T_b (°C), with nestling

age, average number of visits per hour, manipulation (trimmed or untrimmed), and hour of the

day (filtered for diurnal readings; between 05:00 and 21:00), average daily environmental

temperature, relative humidity, and wind speed as main effects. As the relation between

environmental temperature (T_a) and body temperature (T_b) was not linear, we also tested

temperature as a linear (T_a), quadratic (T_a^2) and cubic (T_a^3) term. We detected five T_b

observations that had standardized residuals > 3.0, but since the values were within the range

reportedly previously (Tapper et al 2020b), we retained them in our analysis. To account for

repeated measures of T_b, we included a random intercept for individual male's identity.

Female provisioning rate

In addition to understanding factors that contribute to variation in male provisioning rate, we

were also interested in how a male's behavior, and particularly the experimental trimming

treatment, may impact that of the female partner. Due to heteroscedasticity and non-normality of

residuals in the female provisioning rate model, we applied a natural logarithmic transformation

(ln) to the response term (average number of visits per hour). Where relevant, we back-

transformed values and confidence intervals to the original scale. In our analyses of female

provisioning rate, we included the partner's manipulation (Trimmed or Control (ie.,

untrimmed)), brood size at hatch, clutch initiation date, and average daily environmental

temperature, relative humidity and wind speed, and total daily precipitation as main effects. We

also tested an interaction term of average daily environmental temperature and male

manipulation, since previous work suggested that untrimmed partners may respond to a partner's

trimming manipulation (Tapper et al., 2020a). To account for female identity, we included a

random intercept for individual bird identity. As with males, we report the average provisioning

rate for females (total visits per day divided by total hour recorded per day), but female

provisioning rate was also analyzed with hourly provisioning rate and environmental variables

(see supplementary materials). Note: We did not report T_b of adult females as female T_b has

previously been reported on in (Tapper et al., 2020b).

Nestling growth rate

We calculated nestling growth rate using a three-parameter logistic regression model, following

Tapper et al., (2020a). The logistic growth function models the sigmoidal increase in nestling

mass with age. For the nestling growth rate model, we calculated the asymptote (*A*; the upper

limit of the growth curve), the inflection point (*I*; the point at which the growth rate is greatest),

and growth rate constant (*K;* the steepness of the growth rate curve). For starting estimates in our

growth curves, we used the "SSlogis" function (Stats). We include the five measurements per

nestling (days 0, 3, 6, 9, and 12 post-hatch), which encompasses all major phases of the growth

curve (early, middle, and late; McCarty, 2001) and provides sufficient information for estimates.

Growth curves were fitted in a single hierarchical model for all nestlings, rather than

fitting individual treatment groups or individuals usings a nonlinear mixed-effect structure. We

account for the lack of independence between nestlings from the same brood with box identity as

a random effect. Similarly, to account for repeated measures from the same nestling, we also

included nestling identity as a random effect. Since individual variation can differ at different

stages of growth, we compared three growth models, each with different random effects

structures based on the three growth parameters (*A, I, K*). Each model included nestling and box

identity as random effects, but in each model we allowed for variance in one growth parameter

while holding the other two constant. All models included the same fixed effects and were

assessed for different random effect structures using maximum likelihood, to determine which

random effect structure best explained variance in the different growth curves. The model with

the highest log-likelihood was considered best supported. We included brood size at hatch as a

covariate to account for possible variation between large and small broods.

Results

Male provisioning rate was best predicted by environmental temperature and the provisioning

rate of female partners

Variation in average hourly provisioning rate by males was best predicted by environmental

temperature (T_a), its quadratic term (T_a^2), and average hourly female provisioning rate, each of

which appeared in our only model with strong support ($\Delta AIC_C < 2.0$; Table 1). Our top model

was $>10^5$ times more likely to be the top model than the null (Acc $W_i = 1.0$; ER = 1.0, Table 1),

we therefore report parameter estimates and effects from this model alone. As environmental

temperature (T_a) increased up to approximately 20°C, males increased their provisioning rate

(T_a; $\Box = 3.678$); although provisioning rates decreased at highest environmental temperatures

(T_a^2; $\Box = -0.099$; Fig 1; Table 2). Within a breeding pair, male provisioning rate increased with

increasing female provisioning rate ($\Box = 0.482$; Table 2). Contrary to our prediction - that

trimmed males would increase provisioning rates across all environmental temperatures, the

trimming manipulation was not a strong predictor of male provisioning rate. That is, while

manipulation appeared in our top model ($\Delta AIC_C < 2.0$; Table 1), the confidence intervals crossed

zero (Table 2). On average, male provisioning rate (visits·hr^{-1}) did not differ between treatments

(LSM ± SE: trimmed birds: 10.6 ± 0.50; control birds: 10.9 ± 0.40). Likewise, while we tested

whether the effects of male trimming were dependent on environmental temperature - as

previous work suggested that trimmed birds may be able to work harder at high temperatures

(Tapper et al., 2020a) - neither the interaction term of the first nor second order polynomial

(manipulation $\times T_a$; manipulation $\times T_a^2$) appeared in any of our top models. While hourly data

were also analyzed, the general patterns of male provisioning were consistent between both

temporal scales, with female provisioning rate appearing as a strong predictor at both the average

hourly (Table 1) and hourly scales (Table S1).

Male body temperature did not differ with experimental trimming, but increased with

environmental temperature and provisioning rate

We predicted that trimming manipulations would allow males to increase provisioning rates and

maintain a similar T_b as controls, because of their increased surface area to dissipate heat. The

top model explaining variation in male T_b was modeled with the linear, quadratic, and cubic term

for environmental temperature, along with manipulation and male provisioning rate ($\Delta AIC_C <$

2.0; Table 3). We found manipulation was not a strong predictor of body temperature (P = 0.23,

Table 4; Fig 2). Rather, male T_b increased with temperature (T_a) and its cubic term (T_a^3; both P

<0.001; Table 4, Fig 2). Similarly, male T_b increased with male provisioning rate (P<0.001;

Table 3). Male T_b was not significantly influenced by the quadratic term for temperature (T_a^2; P

= 0.36; Table 4), despite being retained in the top model. Note: Because sample size was limited

for trimmed males, these results must be interpreted with caution as the small sample size may

limit power to detect treatment effects.

Female provisioning rate was best predicted by environmental temperature and the provisioning

rate of male partners

The top model explaining variation in female provisioning rate included average daily

environmental temperature (T_a), male manipulation, and male provisioning rate ($\Delta AIC_C < 2.0$;

Table 5). There was no other competing model, and our top model was $>10^5$ times more likely to

be the top model than the null (Acc Wi = 1.0; ER = 1.0, Table 5), and we report parameter

estimates and effects from this model alone. Females slightly increased their provisioning rate as

environmental temperature increased ($\square$ = 0.02, Table 6; Fig 3). There was a small increase in

female provisioning rate as male provisioning rate increased ($\square$ = 0.05, Table 6). However,

while experimental manipulation of male partners appeared in our top model, the confidence

intervals crossed zero, indicating that females paired with trimmed males did not adjust their

provisioning rate based on male manipulation ($\square$ = -0.11, Table 6). Overall, female provisioning

rate (visits·hr^{-1}) did not differ with male manipulation (LSM ± SE: females partnered with

trimmed males: 12.5 ± 0.57; females partnered with control males 13.6 ± 0.46). As with male

provisioning rate, hourly data for females were also analyzed, with male provisioning rate and

environmental temperature in top models for both temporal scales (Table 5; Table S2).

Male trimming had a modest effect on nestling growth

Nestling growth rate was best modeled with a random effect structure for asymptotic mass (over

models with a random intercept for the inflection point or growth constant; logLik = -2002.56).

We found that male trimming (manipulation) did not affect the asymptote (A) of nestling mass (P

= 0.471, Table 7). On average, nestlings from trimmed males weighed 22.3 ± 1.43g (± SE) at

their asymptote (~ day 12 post-hatch) while nestlings from control males were similar at 21.8 ±

1.25g (± SE; Fig. 4; Fig. 5). However, we found that nestlings from trimmed males developed

slightly faster, reaching the inflection point of the growth curve (I) at 4.51 ± 0.25 days of age (±

SE) compared to controls at 4.29 ± 0.23 days of age (± SE, P = 0.014, Table 7). We also found

that nestlings from trimmed males had a slightly steeper curve for growth rate (i.e., growth rate

constant) than controls (K; P = 0.005, Table 7), growing faster after reaching the inflection point

of growth (nestlings from trimmed males: 2.10 ± 0.21; control nestlings: 1.88 ± 0.19; ± SE).

Nestling mass was similar between small and large broods at the asymptote (P = 0.225, Table 7),

inflection point (P = 0.975, Table 7), and for the growth rate (P = 0.599, Table 7).

Discussion

We found partial support for the hypothesis that nestling provisioning rates by male tree

swallows are limited by the male's risk of overheating (i.e., the heat dissipation limit theory,

Speakman & Król, 2010). Contrary to predictions, males with an enhanced capacity to dissipate

heat via feather trimming did not increase their provisioning rates above the rate of untrimmed

control males, nor did the two groups differ in average T_b. Rather, male provisioning rate was

largely predicted by environmental temperature and the provisioning rate of their female

partners. While we anticipated that nestlings of trimmed males would have increased growth

rates, our results were mixed. Taken together, our results did not provide strong support of the

heat dissipation limit theory.

Male tree swallows decreased nestling provisioning rates at high temperatures and adjusted

effort to match their female partner.

We anticipated that trimmed male tree swallows would maintain higher nestling provisioning

rates than untrimmed control males at highest environmental temperatures, as shown previously

in females (Tapper at al., 2020a). However, while we found that high environmental

temperatures reduced male provisioning, trimmed males were unable (or unwilling) to increase

provisioning above that of control males. A lack of response among males when provided with

an increased capacity to dissipate heat suggests males may be working below their upper thermal

limit. That birds may not be working maximally during parental care is an issue that has been

raised previously (Williams 2018). In breeding blue tits (*Cyanistes caeruleus*), feather trimming

also did not increase provisioning rates, however, trimmed males were better able to maintain

body condition at the same work rate as untrimmed controls (Nord & Nilsson, 2019).

Unfortunately, we could not measure change in body condition in our study due to difficulty in

recapturing male tree swallows. Although trimming did not increase male provisioning rate, the

increased ability to dissipate body heat may have buffered males from the negative consequences

of thermal stress. However, the effectiveness of this strategy is unclear. For example, high

ambient temperatures induce oxidative stress in zebra finches (*Taeniopygia guttata)*, however,

such damage was not ameliorated via trimming manipulations (Zagkle et al., 2022). Since both

trimmed and untrimmed tree swallow males decreased provisioning rates at high temperatures,

the physiological consequences of high environmental temperature may persist regardless of

manipulation.

In general, rates of male tree swallow provisioning appear to be closely tied to those of their

female partner. Previous work showed that experimental manipulation of female provisioning

rate (via increased nestling begging calls) increased the provisioning rate of males, suggesting

that males and females may coordinate care in a reciprocal manner (Lendvai et al., 2018). Males

may also adjust investment based on the relative attractiveness of their mate, with males

compensating for partners of perceived lower quality (i.e., reduced coloration; Limbourg et al.,

2013) or increasing efforts when paired with high quality partners (Mahr et al., 2012). If male

response is reflective of female investment or quality, then trimming manipulation may not fully

account for interactions between pairs in which female quality informs male efforts (Nolazco et

al., 2022).

Male body temperature increased with provisioning rate and environmental temperature but was

not affected by experimental trimming. We interpret these results with caution as our sample

sizes were limited, with only four trimmed males recorded with thermal tags.

Males with enhanced capacity to dissipate heat maintained a similar body temperature (T_b) as

control males, as we predicted. However, given that the experimental manipulation did not

increase male work rate, our results do not support (or conflict with) the heat dissipation limit

theory. However, we acknowledge that our sample sizes were limited for thermal tags (9 control

and 4 trimmed males) which increases uncertainty in our effect estimates. Previous

manipulations of an individual's insulation allowed adults to maintain a lower T_b for the same

work rate (Nord & Nilsson, 2019; Andreasson et al., 2020; Tapper et al., 2020b). Although we

assume that our manipulation increased dry heat loss in trimmed male tree swallows, both

trimmed and control males reduced activity at high temperatures, suggesting both groups were

avoiding hyperthermia. Perhaps reduced vascularization on the male's ventral surface (when

compared with females who have a brood patch) reduced the effectiveness of our feather

trimming in males and minimized the difference in heat loss between the trimmed and control

groups. However, in other species males were found to have increased heat loss as a result of

trimming, yet provisioning efforts were also not affected (Nord & Nilsson, 2019). An additional

explanation is that male tree swallows may simply be working well below their sustained

maximum (e.g., Williams, 2018) and are not at particular risk of overheating. If this were the

case, then our experimental manipulation of males may be ineffective at eliciting increased

parental effort.

An assumption of our test of the heat dissipation limit theory is that nest box visitation rate is an

accurate metric for nestling provisioning and parental activity, however, this may not always be

the case (e.g., Sarota and Williams 2019). Trimmed males may have increased investment in

other activities such as self-feeding (Kacelnik, 1984; Markman, 2014) or nest defense (Swihart

& Johnson, 1986; Patrick et al., 2022), which may have elevated T_b but would not have been

detected by our PIT tag readers. Differential investment strategies between males and females

may therefore complicate tests of the heat dissipation limit theory. That is, considering the

different activities associated with parental investment by males and females, and the cost-

benefits of each activity, heat limitations may impact aspects of male investment beyond

provisioning (Harrison et al., 2009).

Female provisioning rate was best predicted by environmental temperature and the provisioning

rate of male partners

In a companion study we experimentally trimmed feathers from female tree swallows and

observed an increased provisioning rate in both the trimmed females and their untrimmed male

partners (Tapper et al., 2020a). Coordination between males and females may vary depending on

the manipulated individual and the associated cost-benefit functions of care (Harrison et al.,

2009; Bründl et al., 2025). As a result, in the current study we tested the reverse, whether

females adjust their provisioning rate in response to experimental trimming of males. We found

that trimming of males did not affect either male or female provisioning rate; it may be that

female provisioning rate or quality drives male investment strategy more so than the reverse

(Pilastro, et al., 2003; Griggio, et al., 2005; Nolazco et al., 2022). Interestingly, while male tree

swallows reduced provisioning rates at the highest temperatures, females showed a slight

increase. This partial compensation by females may only be possible because females can

dissipate some heat via their vascularized brood patch.

Experimental trimming of males had a modest effect on nestling growth

We found slight differences in the growth trajectories of nestlings from trimmed and untrimmed

(control) males. Nestlings from both groups reached a similar asymptotic mass, however,

nestlings from trimmed males grew faster, despite reaching the midpoint of their growth later

than nestlings from untrimmed males. However, since males were trimmed between days 3-4 of

nestling development, the slight differences in timing (for the midpoint of growth) may not

reflect an actual treatment effect given the limited window of time for a true biological response.

In contrast, previous work with female tree swallows found that nestlings from trimmed mothers

reached a higher asymptotic mass. However, this was thought to be due to enhanced brooding

capabilities of trimmed mothers (i.e., increased heat transfer to nestlings via trimming) rather

than an increased provisioning rate (Tapper et al., 2020a). Since experimental trimming of males

did not increase provisioning rates, a lack of difference in asymptotic mass is perhaps not

surprising. Similarly, while nestlings of trimmed and untrimmed (control) fathers differed in

early growth trajectories (i.e., growth constant and midpoint of growth), the magnitude of these

effects was small and may not correspond to male trimming given the close overlap in timing

between male trimming and the midpoint of nestling growth. Nonetheless, the differences in

early growth rate suggests that male trimming, while not affecting provisioning rate, may have

altered other aspect of parental care. For example, the quality of prey items (Sinkovics et al.,

2021) and coordination of care between parents (Bebbington & Hatchwell, 2016) can each

impact nestling growth. Overall, it remains unknown whether the small differences we detected

in growth trajectories have ecological relevance.

Conclusion

We found limited evidence that provisioning rates by male tree swallows are constrained by the

male's risk of overheating, in contrast with the heat dissipation limit theory. Although males

decreased provisioning rates at high temperatures, they appeared to work routinely below a

thermal limit. Importantly within a pair, males and females coordinated provisioning with

environmental temperature acting as a driving factor. Our results contrast with previous

experimental work on female tree swallows, in which females with an enhanced capacity to

dissipate heat could maintain provisioning rates at high environmental temperatures (Tapper et al

2020a, 2020b). Overall, our study has relevance for aerial insectivores as individuals may face

fitness trade-offs at high temperatures (Van de Ven et al., 2019; Cunningham et al., 2021).

Different responses by males and females highlight the importance of considering sex-specific

differences in thermal physiology and behavior when predicting responses to climate change.

Acknowledgements

We thank Aleesa Schubert, Kaitlyn Baker and Sophia Luimes for assistance in the field, David

Riegert for advice with data analysis, and Simon Tapper for frequent advice and discussions.

Research funding was provided by a Natural Sciences and Engineering Research Council of

Canada (NSERC) - Discovery Grants (awarded to GB).

Data Accessibility:

All data will be archived on the Dryad Data repository

References

Alhenaky, A., Abdelqader, A., Abuajamieh, M., & Al-Fataftah, A. R. (2017). The effect of heat stress on intestinal integrity and Salmonella invasion in broiler birds. *Journal of Thermal Biology*, *70*, 9-14.

Andreasson, F., Hegemann, A., Nord, A., & Nilsson, J. Å. (2020). Experimental facilitation of heat loss affects work rate and innate immune function in a breeding passerine bird. *Journal of Experimental Biology*, *223*(8), jeb219790.

Andreasson, F., Rostedt, E., & Nord, A. (2023). Measuring body temperature in birds–the effects of sensor type and placement on estimated temperature and metabolic rate. *Journal of Experimental Biology*, *226*(24).

Ardia, D. R. (2007). Site-and sex-level differences in adult feeding behaviour and its consequences to offspring quality in tree swallows (*Tachycineta bicolor*) following brood-size manipulation. *Canadian Journal of Zoology*, *85*(8), 847-854.

Bebbington, K., & Hatchwell, B. J. (2016). Coordinated parental provisioning is related to feeding rate and reproductive success in a songbird. *Behavioral Ecology*, *27*(2), 652-659.

Burley, N. (1986). Sexual selection for aesthetic traits in species with biparental care. *The American Naturalist*, *127*(4), 415-445.

Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: Understanding AIC and BIC in model selection. *Sociological Methods and Research, 33*(2), 261–304.

Bründl, A. C., Lejeune, L. A., Deshpande, P., Mulvey, J. E., Thiney, A. C., Chaine, A. S., & Russell, A. F. (2025). Experimental support for partial compensation, not matched, partner response rules in blue tits. *Behavioral Ecology*, *36*(2), arae059.

Clutton-Brock, T. H. (1991). *The evolution of parental care* (Vol. 368). Princeton University Press.

Conradie, S. R., Woodborne, S. M., Cunningham, S. J., & McKechnie, A. E. (2019). Chronic, sublethal effects of high temperatures will cause severe declines in southern African arid-zone birds during the 21st century. *Proceedings of the National Academy of Sciences*, *116*(28), 14065-14070.

Cunningham, S. J., Gardner, J. L., & Martin, R. O. (2021). Opportunity costs and the response of birds and mammals to climate warming. *Frontiers in Ecology and the Environment*, *19*(5), 300-307.

Dakin, R., Lendvai, Á. Z., Ouyang, J. Q., Moore, I. T., & Bonier, F. (2016). Plumage colour is
associated with partner parental care in mutually ornamented tree swallows. *Animal*
Behavior, *111*, 111-118.

Dick, M. F., & Guglielmo, C. G. (2019). Flight muscle protein damage during endurance flight is
related to energy expenditure but not dietary polyunsaturated fatty acids in a migratory
bird. *Journal of Experimental Biology*, *222*(5), jeb187708.

Fox, J. and Weisberg, S. (2019). *An R Companion to Applied Regression*, 3rd edn. Thousand
Oaks, CA, USA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Freeman, M. C., Czenze, Z. J., Schoeman, K., & McKechnie, A. E. (2022). Adaptive varation in
the upper limits of avian body temperature. *Proceedings of the National Academy of*
Sciences 119 (26), e2116645119.

Guillemette, M., Woakes, A. J., Larochelle, J., Polymeropoulos, E. T., Granbois, J. M., Butler, P.
J., Pelletier, D., Frappell, P. B., & Portugal, S. J. (2016). Does hyperthermia constrain
flight duration in a short-distance migrant?. *Philosophical Transactions of the Royal*
Society B: Biological Sciences, *371*(1704), 20150386.

Griggio, M., Valera, F., Casas, A., & Pilastro, A. (2005). Males prefer ornamented females: a
field experiment of male choice in the rock sparrow. *Animal Behaviour*, *69*(6), 1243-
1250.

Hammond, K. A., & Diamond, J. (1997). Maximal sustained energy budgets in humans and
animals. *Nature*, *386*(6624), 457-462.

Harrison, F., Barta, Z., Cuthill, I., & Szekely, T. (2009). How is sexual conflict over parental
care resolved? A meta☐analysis. *Journal of Evolutionary Biology*, *22*(9), 1800-1812.

Hasegawa, M., Arai, E., Ito, S., & Wakamatsu, K. (2016). High brood patch temperature of less
colourful, less pheomelanic female Barn Swallows Hirundo rustica. *Ibis*, *158*(4), 808-
820.

Heeter, K. J., Harley, G. L., Abatzoglou, J. T., Anchukaitis, K. J., Cook, E. R., Coulthard, B. L.,
et al.. (2023). Unprecedented 21st century heat across the Pacific Northwest of North
America. *npj Climate and Atmospheric Science*, *6*(1), 5. https://doi.org/10.1038/s41612-
023-00340-3

Huang, C., Jiao, H., Song, Z., Zhao, J., Wang, X., & Lin, H. (2015). Heat stress impairs
mitochondria functions and induces oxidative injury in broiler chickens. *Journal of*
Animal Science, *93*(5), 2144-2153.

Hurvich, C. M., & Tsai, C. L. (1989). Regression and time series model selection in small
samples. *Biometrika*, *76*(2), 297-307.

IPCC. (2023). *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II*
and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate
Change (Core Writing Team, H. Lee & J. Romero, Eds.). Geneva, Switzerland:
Intergovernmental Panel on Climate Change. pp. 35–115.
https://doi.org/10.59327/IPCC/AR6-9789291691647

Johnson, M. S., & Speakman, J. R. (2001). Limits to sustained energy intake: V. Effect of cold-
exposure during lactation in Mus musculus. *Journal of Experimental Biology*, *204*(11),
1967-1977.

Jones, T., Parrish, J. K., Peterson, W. T., Bjorkstedt, E. P., Bond, N. A., Ballance, L. T., et al.
(2018). Massive mortality of a planktivorous seabird in response to a marine heatwave.
Geophysical Research Letters, 45, 3193–3202. https://doi.org/10.1002/2017GL076164

Kacelnik, A. (1984). Central place foraging in starlings (Sturnus vulgaris). I. Patch residence
time. *The Journal of Animal Ecology*, 283-299.

Kappeler, P. M., Benhaiem, S., Fichtel, C., Fromhage, L., Höner, O. P., Jennions, M. D., Kaiser,
S., Krüger, O., Schneider, J. M., Tuni, C., van Schaik, J., & Goymann, W. (2023). Sex
roles and sex ratios in animals. *Biological Reviews*, *98*(2), 462-480.

Kokko, H., & Jennions, M. D. (2008). Parental investment, sexual selection and sex ratios.
Journal of Evolutionary Biology, *21*(4), 919-948.

LaZerte, S. E., Reudink, M. W., Otter, K. A., Kusack, J., Bailey, J. M., Woolverton, A., Paetkau,
M., de Jong, A. and Hill, D. J. (2017). feedr and animalnexus.ca: a paired R package and
user-friendly Web application for transforming and visualizing animal movement data
from static stations. *Ecology and Evolution.* 7, 7884-7896.

Lemaître, J. F., & Gaillard, J. M. (2017). Reproductive senescence: new perspectives in the
wild. *Biological Reviews*, *92*(4), 2182-2199.

Lendvai, Á. Z., Akçay, Ç., Stanback, M., Haussmann, M. F., Moore, I. T., & Bonier, F. (2018).
Male parental investment reflects the level of partner contributions and brood value in
tree swallows. *Behavioral Ecology and Sociobiology*, *72*, 1-11.

Levillain, A., Reichert, S., & S. Massemin. (2025). The complex relationship between high
temperatues and avian breeding success: insights from a global review. *Ecology and*
Evolution, 15:e71771.

Liker, A., Freckleton, R. P., Remeš, V., & Székely, T. (2015). Sex differences in parental care:
gametic investment, sexual selection, and social environment. *Evolution*, *69*(11), 2862-
2875.

Limbourg, T., Mateman, A. C., & Lessells, C. M. (2013). Opposite differential allocation by
males and females of the same species. *Biology Letters*, *9*(1), 20120835.

Mahr, K., Griggio, M., Granatiero, M., & Hoi, H. (2012). Female attractiveness affects paternal
investment: experimental evidence for male differential allocation in blue tits. *Frontiers*
in Zoology, *9*, 1-8.

Markman, S. (2014). Parental self-feeding effects on parental care levels and time allocation in
Palestine sunbirds. *PLoS One*, *9*(12), e113890.

McCafferty, D. J., Gallon, S., & Nord, A. (2015). Challenges of measuring body temperatures of
free-ranging birds and mammals. *Animal Biotelemetry*, *3*, 1-10.

McCarty, J. P. (2001). Variation in growth of nestling tree swallows across multiple temporal
and spatial scales. *The Auk*, *118*(1), 176-190.

McKechnie, A. E., & Wolf, B. O. (2010). Climate change increases the likelihood of catastrophic
avian mortality events during extreme heat waves. *Biology Letters*, *6*(2), 253-256.

McKechnie, A. E., Hockey, P. A., & Wolf, B. O. (2012). Feeling the heat: Australian landbirds
and climate change. *Emu-Austral Ornithology*, *112*(2), i-vii.

McKechnie, A. E., Rushworth, I. A., Myburgh, F., & Cunningham, S.J. Mortality among birds
and bats during an extreme heat event in eastern South Africa. *Austral Ecology* 46, no. 4
(2021): 687-691.

Møller, A. P., & Birkhead, T. R. (1993). Certainty of paternity covaries with paternal care in
birds. *Behavioral Ecology and Sociobiology*, *33*, 261-268.

Nolazco, S., Delhey, K., Nakagawa, S., & Peters, A. (2022). Ornaments are equally informative
in male and female birds. *Nature Communications*, *13*(1), 5917.

Nilsson, J. Å., & Nord, A. (2018). Testing the heat dissipation limit theory in a breeding

passerine. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1878),

20180652.

Nord, A., & Nilsson, J. Å. (2019). Heat dissipation rate constrains reproductive investment in a

wild bird. *Functional Ecology*, *33*(2), 250-259.

OpenAI. (2025). ChatGPT (Mar 27 version) [Large language model]. https://chat.openai.com/

Patrick, S. C., Réale, D., Potts, J. R., Wilson, A. J., Doutrelant, C., Teplitsky, C., & Charmantier,

A. (2022). Differences in the temporal scale of reproductive investment across the

slow☐fast continuum in a passerine. *Ecology Letters*, *25*(5), 1139-1151.

Pérez, J. H., Krause, J. S., Chmura, H. E., Bowman, S., McGuigan, M., Asmus, A. L., Meddle, S.

L., Hunt, K. E., Gough, L., Boelman, N. T., & Wingfield, J. C. (2016). Nestling growth

rates in relation to food abundance and weather in the Arctic. *The Auk: Ornithological*

Advances, *133*(2), 261-272.

Piatt, J. F., Parrish, J. K., Renner, H. M., Schoen, S. K., Jones, T. T., et al.. (2020). Extreme

mortality and reproductive failure of common murres resulting from the northeast Pacific

marine heatwave of 2014-2016. *PloS one*, *15*(1), e0226087.

Pilastro, A., Griggio, M., & Matessi, G. (2003). Male rock sparrows adjust their breeding

strategy according to female ornamentation: parental or mating investment? *Animal*

Behaviour, *66*(2), 265-271.

Pipoly, I., Preiszner, B., Sándor, K., Sinkovics, C., Seress, G., Vincze, E., Bókony, V., & Liker,

A. (2022). Extreme hot weather has stronger impacts on avian reproduction in forests

than in cities. *Frontiers in Ecology and Evolution*, *10*, 825410.

Ratikainen, I. I., & Kokko, H. (2010). Differential allocation and compensation: who deserves

the silver spoon? *Behavioral Ecology*, *21*(1), 195-200.

Renner, S. S., & Zohner, C. M. (2018). Climate change and phenological mismatch in trophic

interactions among plants, insects, and vertebrates. *Annual Review of Ecology, Evolution,*

and Systematics, *49*(1), 165-182.

Robertson, J. K., Mastromonaco, G., & Burness, G. (2020). Evidence that stress-induced changes

in surface temperature serve a thermoregulatory function. *Journal of Experimental*

Biology, *223*(4), jeb213421.

Rodriguez, S., & Barba, E. (2016). Nestling growth is impaired by heat stress: an experimental
study in a Mediterranean great tit population. *Zoological Studies*, *55*.

Ruuskanen, S., Hsu, B. Y., & Nord, A. (2021). Endocrinology of thermoregulation in birds in a
changing climate. *Molecular and Cellular Endocrinology*, *519*, 111088.

Saeed, M., Abbas, G., Alagawany, M., Kamboh, A. A., Abd El-Hack, M. E., Khafaga, A. F., &
Chao, S. (2019). Heat stress management in poultry farms: A comprehensive overview.
Journal of thermal biology, *84*, 414-425.

Safari, I., Goymann, W., & Kokko, H. (2019). Male-only care and cuckoldry in black coucals:
does parenting hamper sex life?. *Proceedings of the Royal Society B*, *286*(1900),
20182789.

Santos, E. S. A., & Nakagawa, S. (2012). The costs of parental care: a meta‐analysis of the
trade‐off between parental effort and survival in birds. *Journal of Evolutionary Biology*,
25(9), 1911-1917.

Schroeder, J., Hsu, Y. H., Winney, I., Simons, M., Nakagawa, S., & Burke, T. (2016).
Predictably philandering females prompt poor paternal provisioning. *The American*
Naturalist, *188*(2), 219-230.

Senécal, S., Riva, J. C., O'Connor, R. S., Hallot, F., Nozais, C., & Vézina, F. (2021). Poor prey
quality is compensated by higher provisioning effort in passerine birds. *Scientific*
Reports, *11*(1), 11182.

Shuker, D. M., & Kvarnemo, C. (2021). The definition of sexual selection. *Behavioral Ecology*,
32(5), 781-794.

Siefferman, L., & Hill, G. E. (2005). Male eastern bluebirds trade future ornamentation for
current reproductive investment. *Biology Letters*, *1*(2), 208-211.

Sinkovics, C., Seress, G., Pipoly, I., Vincze, E., & Liker, A. (2021). Great tits feed their nestlings
with more but smaller prey items and fewer caterpillars in cities than in forests. *Scientific*
reports, *11*(1), 24161.

Sippel, S., Meinshausen, N., Fischer, E. M., Székely, E., & Knutti, R. (2020). Climate change
now detectable from any single day of weather at global scale. *Nature Climate Change*,
10(1), 35-41.

Solano-Brenes, D., Costa-Schmidt, L. E., Albo, M. J., & Machado, G. (2021). Differential

allocation in a gift-giving spider: males adjust their reproductive investment in response

to female condition. *BMC Ecology and Evolution*, *21*, 1-15.

Speakman, J. R., & Król, E. (2010). Maximal heat dissipation capacity and hyperthermia risk:

neglected key factors in the ecology of endotherms. *Journal of Animal Ecology*, *79*(4),

726-746.

Swihart, R. K., & Johnson, S. G. (1986). Foraging decisions of American robins: somatic and

reproductive tradeoffs. *Behavioral Ecology and Sociobiology*, *19*, 275-282.

Taff, C. C., & Shipley, J. R. (2023). Inconsistent shifts in warming and temperature variability

are linked to reduced avian fitness. *Nature Communications*, *14*(1), 7400.

Tapper, S., Nocera, J. J., & Burness, G. (2020a). Heat dissipation capacity influences

reproductive performance in an aerial insectivore. *Journal of Experimental Biology*,

223(10), jeb222232.

Tapper, S., Nocera, J. J., & Burness, G. (2020b). Experimental evidence that hyperthermia limits

offspring provisioning in a temperate-breeding bird. *Royal Society Open Science*, *7*(10),

201589.

Tattersall, G. J., Roussel, D., Voituron, Y., & Teulier, L. (2016). Novel energy-saving strategies

to multiple stressors in birds: the ultradian regulation of body temperature. *Proceedings*

of the Royal Society B: Biological Sciences, *283*(1839), 20161551.

Thackeray, S. J., Henrys, P. A., Hemming, D., Bell, J. R., Botham, M. S., Burthe, S., P.

Helaouet, et al. (2016). Phenological sensitivity to climate across taxa and trophic levels.

Nature, *535*(7611), 241-245.

Van de Ven, T. M., McKechnie, A. E., & Cunningham, S. (2019). The costs of keeping cool:

behavioural trade-offs between foraging and thermoregulation are associated with

significant mass losses in an arid-zone bird. *Oecologia*, *191*, 205-215.

Vazquez, R., & Larson, D. F. (2013). Plasma protein denaturation with graded heat

exposure. *Perfusion*, *28*(6), 557–559.

Velando, A., Drummond, H., & Torres, R. (2006). Senescent birds redouble reproductive effort

when ill: confirmation of the terminal investment hypothesis. *Proceedings of the Royal*

Society B: Biological Sciences, *273*(1593), 1443-1448.

Visser, M. E., te Marvelde, L., & Lof, M. E. (2012). Adaptive phenological mismatches of birds
and their food in a warming world. *Journal of Ornithology*, *153*, 75-84.

Williams, T. D. (2018). Physiology, activity and costs of parental care in birds. *Journal of
Experimental Biology*, *221*(17), jeb169433.

Winkler, D. W., Hallinger, K. K., Pegan, T. M., Taff, C. C., Verhoeven, M. A., Chang van
Oordt, D., Stager, M., Uehling, J. J., Vitousek, M. N., Andersen, M. J., Ardia, D. R.,
Belmaker, A., Ferretti, V., Forsman, A. M., Gaul, J. R., Llambias, P. E., Orzechowski, S.
C., Shipley, J. R., Wilson, M., & Yoon, H. S. (2020). Full lifetime perspectives on the
costs and benefits of lay□date variation in tree swallows. *Ecology*, *101*(9), e03109.

Zagkle, E., Martinez-Vidal, P. A., Bauchinger, U., & Sadowska, E. T. (2022). Manipulation of
heat dissipation capacity affects avian reproductive performance and output. *Frontiers in
Ecology and Evolution*, *10*, 86618

Table 1. Model selection results for factors that explain variation in average hourly provisioning rate by adult male tree swallows. Of the 16 models run, there was only a single model with strong support (<2 ΔAIC_C). Models were run with a random effect for individual bird identity (Trimmed: n=13, Control: n=18).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R^2m	ER
$T_a + T_a{}^2 +$ Female Provisioning Rate + Manipulation	6	629.81	0.00	1.0	1.0	0.735	1.00
Null Model	2	702.84	73.04	0.00	1.0	0.487	$>10^5$

Note: K = number of parameters, AIC_C= corrected Akaike Information Criterion, Δ AIC_C = difference of focal model AIC_C compared to the lowest model AIC_C; W_i = Akaike weight; Acc. W_i = Accumulated weight of models; R^2m = Marginal R^2; ER = Evidence ratio; T_a = Average daily temperature; $T_a{}^2$ = Quadratic term for average daily temperature; Female Provisioning Rate = Average hourly provisioning rate of females paired with males; Manipulation= Male manipulation (trimmed or untrimmed [Control]). The null model was included for reference but did not have strong support ($\Delta AIC_C > 2$).

779 **Table 2. Factors affecting the average hourly provisioning rate of adult male tree swallows.**

780 Estimates of fixed effects with 95% confidence intervals (CI) and *P*-values (bold signifies

781 statistical significance) are shown. Models were run with a random effect for individual bird

782 identity (Trimmed: n=13, Control: n=18).

Predictor	Estimate	95% CI	P-value
Intercept [Control]	-29.093	-46.59, -11.60	**0.001**
T_a	3.678	1.88, 5.47	**<0.001**
T_a^2	-0.099	-0.15, -0.05	**<0.001**
Female Provisioning Rate	0.482	0.36, 0.61	**<0.001**
Manipulation [Trimmed]	0.673	-1.84, 3.18	0.584

802 **Table 3. Model selection results for factors that explain variation in the body temperature**

803 **of adult male tree swallows.** Of the 10 models run, there was only a single model with strong

804 support (<2 ΔAIC_C). Models were run with a random effect for individual bird identity

805 (Trimmed, n= 4; Control, n= 9).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R^2m	ER
$T_a + T_a^2 + Ta^3 +$ Manipulation + Male Provisioning Rate	11	-21.41	**0.00**	1.00	1.00	0.091	1.00
Null Model	4	95.18	116.59	0.00	1.00	0.00	$>10^5$

808 Note: The null model was included for reference but did not have strong support ($\Delta AIC_C > 2$).

809 T_a^3 = Cubic term for average daily temperature, all other abbreviations are as in Table 1

Table 4. Factors affecting the body temperature of adult male tree swallows. Estimates of fixed effects with 95% confidence intervals (CI) and *P*-values are shown. Bold indicates statistical significance. Models were run with a random effect for individual bird identity (Trimmed, n= 4; Control, n= 9). Approximately 12% of variance in body temperature was accounted for by individual identity.

Predictor	Estimate	95% CI	P-value
Intercept [Control]	41.46	41.245, 41.674	**<0.001**
T_a	3.55	2.885, 4.222	**<0.001**
T_a^2	0.27	-0.314, 0.863	0.363
T_a^3	1.06	0.538, 1.598	**<0.001**
Manipulation [Trimmed]	0.24	-0.170, 0.657	0.226
Male Provisioning Rate	0.01	0.004, 0.012	**<0.001**

844 **Table 5. Model selection results for factors that explain variation in average hourly**

845 **provisioning rate of adult female tree swallows paired with trimmed or untrimmed males.**

846 There was only a single model with strong support (<2 ΔAIC_C). A total of 16 candidate models

847 were run with a random effect for individual bird identity (females partnered with Trimmed

848 males, n= 13; females partnered with Control males, n= 17).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R^2m	ER
T_a + Manipulation + Male Provisioning Rate	5	11.07	0.00	1.00	1.00	0.765	1.00
Null Model	2	85.71	74.64	0.00	1.00	0.399	$>10^5$

Note: The null model was included for reference but did not have strong support ($\Delta AIC_C > 2$).

Abbreviations are as in Table 1

Table 6. Factors affecting the average hourly provisioning rate of adult female tree

swallows. Estimates of fixed effects with 95% confidence intervals (CI) and *P*-values are shown.

Bold indicates statistical significance. Models were run with a random effect for individual bird

identity (females partnered with Trimmed males, n= 13; females partnered with Control males,

n= 17).

Predictor	Estimate	95% CI	P-value
Intercept [Control]	1.73	1.415, 2.041	**<0.001**
T_a	0.02	0.008, 0.0331	**0.002**
Male Provisioning Rate	0.05	0.039, 0.062	**<0.001**
Male Manipulation [Trimmed]	-0.11	-0.326, 0.102	0.291

861 **Table 7. Parameter estimates for growth trajectories in nestling tree swallows.** Estimates of

862 fixed effects with 95% confidence intervals (CI) and *P*-values are shown. Bold indicates

863 statistical significance. Models were run with a random effect for individual nestling identity as

864 well as box identity (Control fathers, n = 20 nests, n= 100 nestlings; Trimmed fathers, n = 15

Predictor	Predictor	Estimate	95% CI	P-value
Asymptote	Intercept	21.80	19.35, 24.26	**<0.001**
(*A*)	Manipulation	0.50	-0.847, 1.837	0.471
	Brood Size	-0.30	-0.774, 1.837	0.225
Inflection Point	Intercept	4.29	3.834, 4.739	**<0.001**
(*I*)	Manipulation	0.22	0.046, 0.397	**0.014**
	Brood Size	0.00	-0.088, 0.085	0.975
Growth Rate Constant	Intercept	1.88	1.502, 2.267	**<0.001**
(*K*)	Manipulation	0.21	0.063, 0.360	**0.005**
	Brood Size	-0.02	-0.093, 0.054	0.599

865 nests, n= 86 nestlings).

870 **Figure 1. Male average hourly provisioning rate decreased at high temperature (T_a), but**

871 **did not differ between trimmed (blue) and control (red) males.** Figure is drawn from the best-

872 supported model, with all other fixed effect values held at their means. Bands and trend lines

873 indicate the predicted means and 95% CI. Individual data points represent average provisioning

874 rate for each individual bird for a given temperature (Trimmed: n=13, Control: n=18).

Figure 2. Male hourly body temperature increased with ambient temperature but did not

differ between trimmed (blue) and control (red) males. Bands and trend lines indicate the

predicted means and 95% CI. Individual data points represent average hourly T_b for each

individual bird at a given temperature (Trimmed, n= 4; Control, n= 9). Note, this figure displays

only the subset of males with temperature-sensitive PIT tags.

Figure 3. Females increased provisioning rates with increasing temperature (T_a), but females partnered with trimmed males did not provision at a higher rate than those paired with control males. Sample sizes: females partnered with Trimmed males, n= 13; females partnered with Control males, n= 17.

Figure 4. Estimated growth trajectories for nestlings from control and trimmed males.
Nestlings from trimmed males reached a similar asymptotic mass as nestlings from control
males. Growth estimates were based on a logistic growth curve using scale, midpoint, and
asymptotes with random effects of nestling and box identity. Bands represent 95% CI. Each
point represents the mass of a nestling on a given developmental day. Sample sizes: Control, n =
20 nests, n= 100 nestlings; trimmed, n = 15 nests, n= 86 nestlings).

Figure 5. Estimates of each growth parameter (asymptote, inflection point, and growth constant) for nestlings from control and trimmed males. Each point represents the raw estimate for each nestling. Sample sizes: Control, n = 20 nests, n= 100 nestlings; trimmed, n = 15 nests, n= 86 nestlings).