Effects of partner workload and increasing environmental temperature on nestling provisioning and body temperature in a declining aerial insectivore

Research Highlights

In adult male tree swallows, we found that experimental manipulations of heat dissipation capacity had little effect on the provisioning rate, body temperature, nor growth of nestlings. That males did not respond to enhanced heat dissipation suggests that they may provision below the threshold of overheating. Rather, male provisioning was affected by the provisioning rate of the male's partner and environmental temperature.

Megan C. Heft,[1]* Bronwen Hennigar[1], and Gary Burness[2]

Addresses:

[1]Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario, Canada. Email: meganheft@trentu.ca

[2]Department of Biology, Trent University, Peterborough, Ontario, Canada. Email: garyburness@trentu.ca

*Corresponding author: meganheft@trentu.ca

Keywords: heat dissipation limit theory, performance, bird**,** parental care, body temperature

Running title: Parental workload and environmental temperatures

Abstract:

With climate change, birds will face increasing thermoregulatory demands, which may alter reproductive behaviors such as offspring provisioning. Experimental studies have shown that the provisioning capacity of female tree swallows (*Tachycineta bicolor*) is limited by their risk of overheating. Given that parental investment strategies may vary between sexes, the thermal environment may have a different impact on males. We experimentally trimmed ventral feathers from male tree swallows to create a "thermal window" through which they could dissipate heat. We remotely monitored provisioning rate and core body temperature of males and their female partners. At high temperatures, all males decreased their nestling provisioning rates irrespective of trimming treatment. In addition, trimmed males maintained core body temperatures similar to those of controls. This suggests that in contrast to previous work with females, males limit provisioning rates to levels below the threshold at which they would overheat. Females paired with trimmed males adjusted their own activity to match that of their male partners; whether there are costs to females is unknown. Combined, these studies highlight that sex-specific differences in thermal physiology and behavior must be considered when predicting responses to climatic warming.

Introduction

Climatic warming is predicted to result in an increase in extreme temperature and weather events (Sippel et al., 2020). In endotherms, prolonged exposure to high ambient temperatures has been associated with reduced reproductive success (e.g., Van de Ven et al., 2020; Pipoly et al., 2022, Levillan et al. 2025) and reports of mass mortality (Jones et al., 2018; Piatt et al., 2020; McKechnie et al., 2021). As thermal challenges are unlikely to subside given the current climatic estimates (IPCC, 2023), the extent to which individuals and populations will be able to effectively respond to thermal stressors is of growing concern (Altan et al., 2003; Heeter et al., 2023; Taff & Shipley, 2023).

Among endotherms, birds may be at particular risk of climatic warming given their relatively high body temperatures and elevated metabolic rates. When birds are faced with thermal stressors, they increase heat loss via evaporative or non-evaporative mechanisms (Tattersall et al., 2016) and/or allow body temperature (T_b) to increase which may result in hyperthermia (e.g., Tapper et al., 2020b; Freeman et al., 2022). In species that exhibit extensive parental care, high environmental temperatures may exacerbate the effects of heat generated by parents during flight and in other activities associated with provisioning offspring (Tapper et al., 2020a). In fact, an individual's rising body temperature and its inability to sufficiently dissipate metabolic heat generated during parental care may limit parental activity (heat-dissipation limit theory; Speakman & Król, 2010; Zagkle et al., 2022). The fitness benefits of increased provisioning rates must therefore be balanced against possible physiological damage induced by hyperthermia, including protein denaturation (Vazquez & Larson, 2013), depressed immune function (Alhenaky et al., 2017), and increased oxidative stress (Huang et al., 2015).

To date, a handful of avian field studies have explored experimentally how the risk of overheating impacts the capacity of individuals to maintain performance during a breeding event. For example, female tree swallows (*Tachycineta bicolor*) provided with an increased capacity to dissipate heat via experimental trimming of feathers, provisioned their nestlings at higher rates than their untrimmed counterparts, at least at highest temperatures (Tapper et al., 2020a). This suggested that at high environmental temperatures, an individual's activity (and thereby capacity to care for young) was determined by a thermal constraint, as predicted by the heat-dissipation limit theory (Speakman & Król, 2010). In contrast, female blue tits (*Cyanistes caeruleus)* that were trimmed of feathers did not increase provisioning, but instead displayed enhanced immune function, suggesting a prioritization of self-maintenance (Nord & Nilsson, 2019; Andreasson et al., 2020).

An individual's parental-investment strategy involves a trade-off between allocating energy and resources between current and future reproductive attempts (Clutton-Brock, 1991; Lemaître & Gaillard, 2017). In species that exhibit biparental care, thermal limits may play a role in pair coordination, with possible effects on investment strategies. Recently, it was shown that male tree swallows partnered with experimentally trimmed females increased their *own* provisioning efforts at high environmental temperatures, despite not being trimmed themselves (Tapper et al., 2020a). Male and female tree swallows are known the adjust their provisioning efforts based on partner quality (Lendvai et al., 2018). However, if males were able to increase their workload to better match that of their female partners, males were either working below their physiological limit, or perhaps willing to endure physiological damage (e.g., Dick & Guglielmo, 2019). In this sense, sex-specific investment strategies may emerge when males and females are faced with the removal of a possible physiological constraint (Nilsson & Nord, 2018).

Considering that reproductive strategies can vary between sexes (Kappeler et al., 2023),
we investigated whether a male tree swallow's capacity to dissipate heat would alter its
reproductive investment, and we qualitatively compared our results with previous studies on
females (Tapper et al 2020a, 2020b). We experimentally trimmed ventral feathers from male tree
swallows to allow for enhanced heat loss and measured provisioning rates and body temperature
in trimmed individuals and compared them with untrimmed controls. Further, we explored the
response of their female partners (who were unmanipulated). We anticipated that the capacity to
dissipate metabolic heat would limit activity in adult male tree swallows (i.e., the heat dissipation
limit theory). We made the following predictions: (1) males with an augmented capacity to
dissipate heat (trimmed males) would alter their reproductive investment, and increase their
provisioning rate to nestlings across all environmental temperatures; (2) trimmed males would
maintain a similar body temperature as untrimmed controls, by adjusting their activity to
maintain T_b within a narrow range; (3) females paired with trimmed males would have higher
nestling provisioning rates than females paired with non-trimmed males, because the
provisioning rates of pairs are often coordinated (Lendvai et al., 2018); (4) nestlings from
trimmed males would reach a heavier mass than nestlings from control males.
107
108
Methods
Study species and general field methods
All research was approved by the Trent University Animal Care Committee, in accordance with
the Canadian Council on Animal Care (Animal Use Protocol # 26534). We conducted this study
from May-July 2021, on two nest-box breeding populations of tree swallows located at the Trent
University Nature Areas, Peterborough, Ontario, Canada (44°21′N, 78°17′W) and at the
Lakefield Sewage Lagoon, Lakefield, Ontario (44°24'58.3"N 78°15'26.8"W). The Nature Areas
site contains ~40 boxes spread out in a grid across the site, while the Sewage Lagoon site
consists of two large, central lagoons surrounded by a perimeter of ~80 nest boxes. Further
details on the sites are found in Fischer et al. (2020). At our study sites, tree swallows begin
constructing their nests in early May, typically rearing 1 brood per season. Clutch size varies
between 3-7 eggs, laid approximately one day apart. Incubation typically lasts 14 days with
nestlings hatching relatively synchronously. After hatch, the nestling period spans approximately
15-22 days with adults providing food and maintenance care until the nestlings fledge (Winkler
et al., 2020).
124
125 We monitored nests boxes daily from May through July to determine the lay and hatch dates for
126 each clutch. As each egg was laid, we labelled it with a marker to establish order of laying and
127 incubation dates. Incubation day 0 was the date that the last egg was laid, and hatch day 0 was
128 the date the majority of nestlings in a brood hatched.
129
Experimental manipulation of heat dissipation
When nestlings were between days 3-4 post-hatch, we captured adult male tree swallows and
performed a feather-trimming manipulation. We assigned males randomly to either a Trimmed
or Control treatment by flipping a coin (Trimmed: n=15, Control: n=20). Males in the Trimmed
group had an approximately 3 x 2 cm area of ventral feathers trimmed with scissors, such that
approximately 7% of body surface area was exposed, following Tapper et al. (2020a). To ensure
the degree of trimming was consistent among individual males, we used a cardboard "stencil,"

placed over each individual's ventral surface. Males assigned to Control groups were handled for
a similar length of time as the Trimmed individuals, but no feathers were removed.
Following trimming (or handling, in case of Controls), we implanted each male with
either a temperature-sensitive passive integrated transponder (PIT) tag (accuracy ±0.5°C;
Biotherm13; Biomark, Boise, Idaho, USA,) or attached a non-temperature-sensitive leg-mounted
RFID tag (EM4100; #11001, GAO RFID, Ontario, 166 Canada). Tag type was randomized by
coin flip. To implant a PIT tag, we first sterilized the incision site with ethanol, implanted the tag
subcutaneously into the nape of the neck, and then sealed the incision site with tissue adhesive
bond (Vetbond Tissue Adhesive, Fisher Scientific, Waltham, MA, USA). Prior to release we
collected a 50-75 µL blood sample (as part of another study) and recorded body morphometrics
(body mass, wing chord (flattened), and head-bill length). Handling time was < 10 minutes for
each individual. Given limitations due to synchrony at hatch, loggers were rotated among nests
and fewer birds were recorded than were initially trimmed and tagged. Sample size for
temperature sensitive tagged males: Trimmed males: n= 4, Control males: n= 9. Sample size for
RFID tagged males: Trimmed males: n= 9, Control males: n= 9.
To test female responsiveness to male trimming, we captured females at the same time as
males and equipped each female with the same tag type as their male mates received (i.e, females
paired with males that were equipped with temperature-sensitive tags were also assigned
temperature-sensitive tags). We collected a blood sample and recorded the same morphometrics
as for males; no females were trimmed (females paired with Trimmed males: n= 13; females
paired with Control males: n= 17). Individuals were then released. One female paired with a
Control male was tagged, but no nestbox visits were recorded while loggers were deployed. As a
result, the sample size for females is slightly lower than that of males.
160
Remote Monitoring
For all tagged adults we recorded nest-box visitation rates, while for individuals implanted with
temperature-sensitive tags we also recorded subcutaneous body temperature (T_b). We considered
T_b to be a proxy for core body temperature (McCafferty et al., 2015; Andreasson et al., 2023).
For individuals with temperature-sensitive tags, we recorded visitation rates and T_b using either a
Biomark HPR Plus or a Biomark Small Scale Monitoring System (AR650) reader. Because of a
limited number of Biomark readers, we cycled them among nests to capture adult visitation rates
during 2-3 day windows throughout nestling development (early, middle, and late from 2-5 days,
6-9 days, and 10-14 days post-hatch, respectively). Readers (connected to 12v batteries) were
placed at the base of each nest box and attached to a loop antenna mounted to the nest entrance.
Visits were recorded anytime an individual entered the nest or perched at the nest entrance. Both
types of Biomark readers were set to a 10 second 'delay' to limit tag detections when a bird spent
a prolonged period in the nest box (Tapper at al., 2020). Individual reads included the date, time,
and the bird identification code for a given tag (recorded when each tag was initially deployed).
175
To record visitation rates of individuals with leg mounted tags (rather than implanted
temperature-sensitive tags), we used Generation 2 RFID readers (Cellular Tracking Technology,
Rio Grande, NJ, USA) (Bonter and Bridge, 2011). Because we had a large number of Generation
2 RFID readers, we did not need to cycle them among nests. Nonetheless, to allow for
comparison with data from temperature sensitive tags, we blocked our non-temperature sensitive
detection data into 2-3 day windows to match the cycling of the Biomark readers.
182

Environmental Data
We collected hourly readings for environmental temperature (°C), relative humidity (%), and
windspeed (km ·hr-1), and total daily precipitation (mm) from the Trent University weather
station (1.5 km from the Trent University Nature Areas and 9.5 km from Lakefield Sewage
Lagoons). All environmental data were downloaded from Environment Canada (available at
https://climate.weather.gc.ca/index_e.html).
189
Nestling Morphometrics
Upon hatching of nestlings, we marked each nestling with a colored marker, allowing us to track
the growth rate of individuals (until they were large enough to band). We weighed nestlings
between the hours of ~1200 – 1800 on days 0 (hatch), 3, 6, 9 and 12 post-hatch on a digital
portable scale (Smart Weigh Digital Pro Pocket, ± 0.01 g). No measurements were recorded after
day 12 to prevent premature fledge. In our analysis, nestlings were assigned to either a Control (n
= 20) or Trimmed (n = 15) group based on the manipulation of the father, though no nestlings
were manipulated. Note that the sample sizes for nests of nestlings were greater than those for
adults used in provisioning rate analyses (above); due to logger limitations (cycling loggers
between boxes) four nests could not be included. However, males were still captured and
manipulated at these nests.
201
Data compilation
<u>*Nest box visitation rate*</u>
All data organization and analysis was performed in R (R version 4.2.2, R Core Team). To
quantify RFID reads (from both thermal and nonthermal tag types), we transformed raw RFID
reads into nest box visits using the 'visits' function from feedr (LaZerte et al., 2017; Tapper et
al., 2020a). We considered a nest box visit to be an index of parental provisioning. Subsequent
statistical analyses for parental provisioning visits were then completed using these transformed
data. Observations for adult provisioning visits were balanced within pairs (n=133 male
observations, n=144 female observations).
211
Because males were captured between days 3-4 of nestling development, we restricted our
analyses of provisioning visits to days 3-14 of nestling development. In addition, adults do not
typically feed nestlings between the hours of ~21:00 and 05:00 (Tapper et al., 2020a), so we
excluded this window from our provisioning rate data. In the case where successive RFID reads
occurred within 60 seconds (i.e., a bird was present in the box for multiple reads within the span
of 60 seconds), we defined this as a singular visit. In our analysis, provisioning rate was defined
as the average number of provisioning visits per hour of daytime activity (i.e., between 05:00 and
21:00 each day), or per hour that a logger was recording at a given nest box (if the logger was
removed partway through the day).
221
Statistical analysis
<u>*Provisioning rate*</u>
For our analyses of male and female provisioning rates, we evaluated competing models with an
Akaike Information Criterion corrected for small sample sizes (AICc). In our analysis, we
considered models with a ΔAICc of < 2 to be strongly supported (Burnham & Anderson, 2004).
For our provisioning rate models, we evaluated models based on model weights and likelihoods

(Hurvich & Tsai, 1989). For models with strong support, we report the 95% confidence intervals
(CI) and associated effect size for individual terms.
230
231   *Statement on AI:*
232   Artificial Intelligence (AI) was used during data analysis to refine visualizations and to suggest
233   corrections to code. Specifically, ChatGPT (OpenAI, 2025) was used to improve layouts, color
234   scheme, and labeling of figures, for clearer visualizations. All code was reviewed and checked
235   for accuracy.
236
Male provisioning rate
In our analysis of male provisioning rate, we were primarily interested in the response of males
to our experimental manipulation (trimming). To explore potential factors contributing to
variation in average number of visits per hour, we evaluated combinations of predictors
including manipulation (trimmed or control, i.e., untrimmed), brood size (at hatch), clutch
initiation date, average daily environmental temperature, relative humidity and wind speed, and
total daily precipitation. We evaluated environmental predictors for collinearity by running a
variance inflation factor test (all VIF <3; 'vif' function, Car package, version 3.1-2; Fox and
Weisberg, 2019). We also included the average number of visits per hour for females (filtered
from 05:00 to 21:00). Based on the nonlinear relation between male provisioning rate and
environmental temperature, we included a quadratic term for average daily environmental
temperature (T_a^2). Since previous work has shown that the effects of treatment may vary based
on environmental temperature (Tapper et al., 2020a), we also tested for an interaction between
average daily environmental temperature and manipulation (along with an interaction between
manipulation and the quadratic; T_a^2). To account for repeated measures from the same individual
over the brood rearing period, we included a random intercept for individual male's identity.
253
Male body temperature, T_b
We tested the impact of experimental trimming on male average hourly T_b (°C), with nestling
age, average number of visits per hour, manipulation (trimmed or untrimmed), and hour of the
day (filtered for diurnal readings; between 05:00 and 21:00), average daily environmental
temperature, relative humidity, and wind speed as main effects. As the relation between
environmental temperature (T_a) and body temperature (T_b) was not linear, we also tested
temperature as a linear (T_a), quadratic (T_a^2) and cubic (T_a^3) term. We detected five T_b
observations that had standardized residuals > 3.0, but since the values were within the range
reportedly previously (Tapper et al 2020b), we retained them in our analysis.
263
Female provisioning rate
In addition to understanding factors that contribute to variation in male provisioning rate, we
were also interested in how a male's behavior, and particularly the experimental trimming
treatment, may impact that of the female partner. Due to heteroscedasticity and non-normality of
residuals in the female provisioning rate model, we applied a natural logarithmic transformation
(ln) to the response term (average number of visits per hour). Where relevant, we back-
transformed values and confidence intervals to the original scale. In our analyses of female
provisioning rate, we included the partner's manipulation (Trimmed or Control (ie.,
untrimmed)), brood size at hatch, clutch initiation date, and average daily environmental
temperature, relative humidity and wind speed, and total daily precipitation as main effects. We

also tested an interaction term of average daily environmental temperature and male
manipulation, since previous work suggested that untrimmed partners may respond to a partner's
trimming manipulation (Tapper et al., 2020a). To account for female identity, we included a
random intercept for individual bird identity. Note: We did not report T_b of adult females as
female T_b has previously been reported on in (Tapper et al., 2020b).
279
280

Nestling growth rate
We calculated nestling growth rate using a three-parameter logistic regression model,
following Tapper et al., (2020a). For the nestling growth rate model, we calculated the asymptote
(*A*; the upper limit of the growth curve), the inflection point (*I*; the point at which the growth rate
is greatest), and growth rate constant (*K;* the steepness of the growth rate curve). For starting
estimates in our growth curves, we used the "SSlogis" function (Stats).
To account for the lack of independence between nestlings from the same brood, we
included nest box identity as a random effect. Similarly, to account for repeated measures from
the same nestling, we also included nestling identity as a random effect. We evaluated random
effect structures for each of the three growth parameters (*A, I, K*). All models included the same
fixed effects and were assessed using AICc, the model with the lowest AICc was considered best
supported. Evidence ratios and Akaike weights were also calculated to assess relative model
support. We included brood size at hatch as a covariate to account for possible variation between
large and small broods.
295
296

Results
Male provisioning rate was best predicted by environmental temperature and the provisioning
rate of female partners
Variation in average hourly provisioning rate by males was best predicted by
environmental temperature (T_a), its quadratic term (T_a^2), and average hourly female provisioning
rate, each of which appeared in our only model with strong support ($\Delta AIC_C < 2.0$; Table 1). Our
top model was $>10^5$ times more likely to be the top model than the null (Acc $W_i = 1.0$; ER = 1.0,
Table 1), we therefore report parameter estimates and effects from this model alone. As
environmental temperature (T_a) increased up to approximately 20°C, males increased their
provisioning rate ($\beta = 3.678$); although provisioning rates decreased at highest environmental
temperatures ($\beta = -0.099$; Fig 1; Table 2). Within a breeding pair, male provisioning rate
increased with increasing female provisioning rate ($\beta = 0.482$; Table 2). Contrary to our
prediction - that trimmed males would increase provisioning rates across all environmental
temperatures, the trimming manipulation was not a strong predictor of male provisioning rate.
That is, while manipulation appeared in our top model ($\Delta AIC_C < 2.0$; Table 1), the confidence
intervals crossed zero (Table 2). On average, male provisioning rate (visits·hr^{-1}) did not differ
between treatments (LSM ± SE: trimmed birds: 10.6 ± 0.50; control birds: 10.9 ± 0.40).
Likewise, while we tested whether the effects of male trimming were dependent on
environmental temperature - as previous work suggested that trimmed birds may be able to work
harder at high temperatures (Tapper et al., 2020a) - neither the interaction term of the first nor
second order polynomial (manipulation × T_a; manipulation × T_a^2) appeared in any of our top
models.
319

Male body temperature did not differ with experimental trimming, but increased with
environmental temperature and provisioning rate
We predicted that trimming manipulations would allow males to increase provisioning
rates and maintain a similar T_b as controls, as a result of their increased surface area to dissipate
heat. The top model explaining variation in male T_b was modeled with the linear, quadratic, and
cubic term for environmental temperature, along with manipulation and male provisioning rate
($\Delta AIC_C < 2.0$; Table 3). We found manipulation was not a strong predictor of body temperature
(P = 0.23, Table 4; Fig 2). Rather, male T_b increased with temperature (T_a) and its cubic term
(T_a^3; both P <0.001; Table 4, Fig 2). Similarly, male T_b increased with male provisioning rate
(P<0.001; Table 3). Male T_b was not significantly influenced by the quadratic term for
temperature (T_a^2; P = 0.36; Table 4), despite being retained in the top model.
331
Female provisioning rate was best predicted by environmental temperature and the provisioning
rate of male partners
The top model explaining variation in female provisioning rate included average daily
environmental temperature (T_a), male manipulation, and male provisioning rate ($\Delta AIC_C < 2.0$;
Table 5). There was no other competing model, and our top model was $>10^5$ times more likely to
be the top model than the null (Acc Wi = 1.0; ER = 1.0, Table 5), and we report parameter
estimates and effects from this model alone. Females slightly increased their provisioning rate as
environmental temperature increased ($\beta = 0.02$, Table 6; Fig 3). There was a small increase in
female provisioning rate as male provisioning rate increased ($\beta = 0.05$, Table 6). However, while
experimental manipulation of male partners appeared in our top model, the confidence intervals
crossed zero, indicating that females paired with trimmed males did not adjust their provisioning
rate based on male manipulation ($\beta = -0.11$, Table 6). Overall, female provisioning rate
(visits·hr^{-1}) did not differ with male manipulation (LSM ± SE: females partnered with trimmed
males: 12.5 ± 0.57; females partnered with control males 13.6 ± 0.46).
346
Male trimming had a modest effect on nestling growth
Nestling growth rate was best modeled with a random effect structure for asymptotic mass (over
models with a random intercept for the inflection point or growth constant; $\Delta AIC_C < 2.0$, Table
7). We found that male trimming (manipulation) did not affect the asymptote (*A*) of nestling
mass (P = 0.471, Table 8). On average, nestlings from trimmed males weighed 22.3 ± 1.43g (±
SE) at their asymptote (~ day 12 post-hatch) while nestlings from control males were similar at
21.8 ± 1.25g (± SE; Fig. 4). However, we found that nestlings from trimmed males reaching the
inflection point of the growth curve (*I*) at 4.51 ± 0.25 days of age (± SE) compared to controls at
4.29 ± 0.23 days of age (± SE, P = 0.014, Table 8). We also found that nestlings from trimmed
males had a steeper curve for growth rate (i.e., growth rate constant) than controls (*K;* P = 0.005,
Table 8), growing faster after reaching the inflection point of growth (nestlings from trimmed
males: 2.10 ± 0.21; control nestlings: 1.88 ± 0.19; ± SE).
359
Discussion
We found partial support for the hypothesis that nestling provisioning rates by male tree
swallows are limited by the male's risk of overheating (i.e., the heat dissipation limit theory,
Speakman & Król, 2010). Contrary to predictions, males with an enhanced capacity to dissipate
heat via feather trimming did not increase their provisioning rates above the rate of untrimmed
control males, nor did the two groups differ in average T_b. Rather, male provisioning rate was

largely predicted by environmental temperature and the provisioning rate of their female partners. While we anticipated that nestlings of trimmed males would have increased growth rates, our results were mixed. Taken together, our results did not provide strong support of the heat dissipation limit theory.

Male tree swallows decreased nestling provisioning rates at high temperatures and adjusted effort to match their female partner.
We anticipated that trimmed male tree swallows would maintain higher nestling provisioning rates than untrimmed control males at highest environmental temperatures, as shown previously in females (Tapper at al., 2020a). However, while we found that high environmental temperatures reduced male provisioning, trimmed males were unable (or unwilling) to increase provisioning above that of control males. A lack of response among males when provided with an increased capacity to dissipate heat suggests males may be working below their upper thermal limit. That birds may not be working maximally during parental care is an issue that has been raised previously (Williams 2018). In breeding blue tits (*Cyanistes caeruleus*), feather trimming also did not increase provisioning rates, however, trimmed males were better able to maintain body condition at the same work rate as untrimmed controls (Nord & Nilsson, 2019). Unfortunately, we could not measure change in body condition in our study due to difficulty in recapturing male tree swallows. Although trimming did not increase male provisioning rate, the increased ability to dissipate body heat may have buffered males from the negative consequences of thermal stress. However, the effectiveness of this strategy is unclear. For example, high ambient temperatures induce oxidative stress in zebra finches (*Taeniopygia guttata)*, however, such damage was not ameliorated via trimming manipulations (Zagkle et al., 2022). Since both trimmed and untrimmed tree swallow males decreased provisioning rates at high temperatures, the physiological consequences of high environmental temperature may persist regardless of manipulation.

In general, rates of male tree swallow provisioning appear to be closely tied to those of their female partner. Previous work showed that experimental manipulation of female provisioning rate (via increased nestling begging calls) increased the provisioning rate of males, suggesting that males and females may coordinate care in a reciprocal manner (Lendvai et al., 2018). Males may also adjust investment based on the relative attractiveness of their mate, with males compensating for partners of perceived lower quality (i.e., reduced coloration; Limbourg et al., 2013) or increasing efforts when paired with high quality partners (Mahr et al., 2012). If male response is reflective of female investment or quality, then trimming manipulation may not fully account for interactions between pairs in which female quality informs male efforts (Nolazco et al., 2022).

Male body temperature increased with provisioning rate and environmental temperature, but was not affected by experimental trimming
Males with enhanced capacity to dissipate heat maintained a similar body temperature (T_b) as control males, as we predicted. However, given that the experimental manipulation did not increase male work rate, our results do not support (or conflict with) the heat dissipation limit theory. Previous manipulations of an individual's insulation allowed adults to maintain a lower T_b for the same work rate (Nord & Nilsson, 2019; Andreasson et al., 2020; Tapper et al., 2020b). Although we assume that our manipulation increased dry heat loss in trimmed male tree

swallows, both trimmed and control males reduced activity at high temperatures, suggesting both groups were avoiding hyperthermia. Perhaps reduced vascularization on the male's ventral surface (when compared with females who have a brood patch) reduced the effectiveness of our feather trimming in males and minimized the difference in heat loss between the trimmed and control groups. However, in other species males were found to have increased heat loss as a result of trimming, yet provisioning efforts were also not affected (Nord & Nilsson, 2019). An additional explanation is that male tree swallows may simply be working well below their sustained maximum (e.g., Williams, 2018) and are not at particular risk of overheating. If this were the case, then our experimental manipulation of males may be ineffective at eliciting increased parental effort.

An assumption of our test of the heat dissipation limit theory is that nest box visitation rate is an accurate metric for nestling provisioning and parental activity, however, this may not always be the case (e.g., Sarota and Williams 2019). Trimmed males may have increased investment in other activities such as self-feeding (Kacelnik, 1984; Markman, 2014) or nest defense (Swihart & Johnson, 1986; Patrick et al., 2022), which may have elevated T_b but would not have been detected by our PIT tag readers. Differential investment strategies between males and females may therefore complicate tests of the heat dissipation limit theory. That is, considering the different activities associated with parental investment by males and females, and the cost-benefits of each activity, heat limitations may impact aspects of male investment beyond provisioning (Harrison et al., 2009).

Female provisioning rate was best predicted by environmental temperature and the provisioning rate of male partners
In a companion study we experimentally trimmed feathers from female tree swallows and observed an increased provisioning rate in both the trimmed females and their untrimmed male partners (Tapper et al., 2020a). Coordination between males and females may vary depending on the manipulated individual and the associated cost-benefit functions of care (Harrison et al., 2009; Bründl et al., 2025). As a result, in the current study we tested the reverse, whether females adjust their provisioning rate in response to experimental trimming of males. We found that trimming of males did not affect either male or female provisioning rate; it may be that female provisioning rate or quality drives male investment strategy more so than the reverse (Pilastro, et al., 2003; Griggio, et al., 2005; Nolazco et al., 2022). Interestingly, while male tree swallows reduced provisioning rates at the highest temperatures, females showed a slight increase. This partial compensation by females may only be possible because females can dissipate some heat via their vascularized brood patch.

Experimental trimming of males had a modest effect on nestling growth
We found differences in the growth trajectories of nestlings from trimmed and untrimmed (control) males. Nestlings from both groups reached a similar asymptotic mass, however, nestlings from trimmed males grew faster, despite reaching the midpoint of their growth later than nestlings from untrimmed males. In contrast, previous work with female tree swallows found that nestlings from trimmed mothers reached a higher asymptotic mass. However, this was thought to be due to enhanced brooding capabilities of trimmed mothers (i.e., increased heat transfer to nestlings via trimming) rather than an increased provisioning rate (Tapper et al., 2020a). Since experimental trimming of males did not increase provisioning rates, a lack of

difference in asymptotic mass is perhaps not surprising. Similarly, while nestlings of trimmed
and untrimmed (control) fathers differed in early growth trajectories (i.e., growth constant and
midpoint of growth), the magnitude of these effects was small. Nonetheless, differences in early
growth trajectories suggests that male trimming, while not affecting provisioning rate, may have
altered other aspect of parental care. For example, the quality of prey items (Sinkovics et al.,
2021) and coordination of care between parents (Bebbington & Hatchwell, 2016) can each
impact nestling growth. However, it remains unknown whether the small differences we detected
in growth trajectories have ecological relevance.
466
467

Conclusion
We found limited evidence that provisioning rates by male tree swallows are constrained by the
male's risk of overheating, in contrast with the heat dissipation limit theory. Although males
decreased provisioning rates at high temperatures, they appeared to work routinely below a
thermal limit. Importantly within a pair, males and females coordinated provisioning with
environmental temperature acting as a driving factor. Our results contrast with previous
experimental work on female tree swallows, in which females with an enhanced capacity to
dissipate heat could maintain provisioning rates at high environmental temperatures (Tapper et al
2020a, 2020b). Overall, our study has relevance for aerial insectivores as individuals may face
fitness trade-offs at high temperatures (Van de Ven et al., 2019; Cunningham et al., 2021).
Different responses by males and females highlight the importance of considering sex-specific
differences in thermal physiology and behavior when predicting responses to climate change.
480
481

Acknowledgements
We thank Aleesa Schubert and Kaitlyn Baker for assistance in the field, David Riegert for advice
with data analysis, and Simon Tapper for frequent advice and discussions. Research funding was
provided by a Natural Sciences and Engineering Research Council of Canada (NSERC) -
Discovery Grants (awarded to GB).

Data Accessibility:

All data will be archived on the Dryad Data repository

490
491
492

References

Alhenaky, A., Abdelqader, A., Abuajamieh, M., & Al-Fataftah, A. R. (2017). The effect of heat

stress on intestinal integrity and Salmonella invasion in broiler birds. *Journal of Thermal*

Biology, *70*, 9-14.

Andreasson, F., Hegemann, A., Nord, A., & Nilsson, J. Å. (2020). Experimental facilitation of

heat loss affects work rate and innate immune function in a breeding passerine bird.

Journal of Experimental Biology, *223*(8), jeb219790.

Andreasson, F., Rostedt, E., & Nord, A. (2023). Measuring body temperature in birds–the effects

of sensor type and placement on estimated temperature and metabolic rate. *Journal of*

Experimental Biology, *226*(24).

Ardia, D. R. (2007). Site-and sex-level differences in adult feeding behaviour and its

consequences to offspring quality in tree swallows (*Tachycineta bicolor*) following

brood-size manipulation. *Canadian Journal of Zoology*, *85*(8), 847-854.

Bebbington, K., & Hatchwell, B. J. (2016). Coordinated parental provisioning is related to

feeding rate and reproductive success in a songbird. *Behavioral Ecology*, *27*(2), 652-659.

Burley, N. (1986). Sexual selection for aesthetic traits in species with biparental care. *The*

American Naturalist, *127*(4), 415-445.

Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: Understanding AIC and BIC

in model selection. *Sociological Methods and Research, 33*(2), 261–304.

https://doi.org/10.1177/0049124104268644

Bründl, A. C., Lejeune, L. A., Deshpande, P., Mulvey, J. E., Thiney, A. C., Chaine, A. S., &

Russell, A. F. (2025). Experimental support for partial compensation, not matched,

partner response rules in blue tits. *Behavioral Ecology*, *36*(2), arae059.

Clutton-Brock, T. H. (1991). *The evolution of parental care* (Vol. 368). Princeton University

Press.

Conradie, S. R., Woodborne, S. M., Cunningham, S. J., & McKechnie, A. E. (2019). Chronic,

sublethal effects of high temperatures will cause severe declines in southern African arid-

zone birds during the 21st century. *Proceedings of the National Academy of Sciences*,

116(28), 14065-14070.

Cunningham, S. J., Gardner, J. L., & Martin, R. O. (2021). Opportunity costs and the response of

birds and mammals to climate warming. *Frontiers in Ecology and the Environment*,

19(5), 300-307.

Dakin, R., Lendvai, Á. Z., Ouyang, J. Q., Moore, I. T., & Bonier, F. (2016). Plumage colour is

associated with partner parental care in mutually ornamented tree swallows. *Animal*

Behavior, *111*, 111-118.

Dick, M. F., & Guglielmo, C. G. (2019). Flight muscle protein damage during endurance flight is
related to energy expenditure but not dietary polyunsaturated fatty acids in a migratory
bird. *Journal of Experimental Biology*, *222*(5), jeb187708.

Fox, J. and Weisberg, S. (2019). *An R Companion to Applied Regression*, 3rd edn. Thousand
Oaks, CA, USA: Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Freeman, M. C., Czenze, Z. J., Schoeman, K., & McKechnie, A. E. (2022). Adaptive varation in
the upper limits of avian body temperature. *Proceedings of the National Academy of*
Sciences 119 (26), e2116645119. Guillemette, M., Woakes, A. J., Larochelle, J.,
Polymeropoulos, E. T., Granbois, J. M., Butler, P. J., Pelletier, D., Frappell, P. B., &
Portugal, S. J. (2016). Does hyperthermia constrain flight duration in a short-distance
migrant?. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
371(1704), 20150386.

Griggio, M., Valera, F., Casas, A., & Pilastro, A. (2005). Males prefer ornamented females: a
field experiment of male choice in the rock sparrow. *Animal Behaviour*, *69*(6), 1243-
1250.

Hammond, K. A., & Diamond, J. (1997). Maximal sustained energy budgets in humans and
animals. *Nature*, *386*(6624), 457-462.

Harrison, F., Barta, Z., Cuthill, I., & Szekely, T. (2009). How is sexual conflict over parental
care resolved? A meta-analysis. *Journal of Evolutionary Biology*, *22*(9), 1800-1812.

Hasegawa, M., Arai, E., Ito, S., & Wakamatsu, K. (2016). High brood patch temperature of less
colourful, less pheomelanic female Barn Swallows Hirundo rustica. *Ibis*, *158*(4), 808-
820.

Heeter, K. J., Harley, G. L., Abatzoglou, J. T., Anchukaitis, K. J., Cook, E. R., Coulthard, B. L.,
et al.. (2023). Unprecedented 21st century heat across the Pacific Northwest of North
America. *npj Climate and Atmospheric Science*, *6*(1), 5. https://doi.org/10.1038/s41612-
023-00340-3

Huang, C., Jiao, H., Song, Z., Zhao, J., Wang, X., & Lin, H. (2015). Heat stress impairs
mitochondria functions and induces oxidative injury in broiler chickens. *Journal of*
Animal Science, *93*(5), 2144-2153.

Hurvich, C. M., & Tsai, C. L. (1989). Regression and time series model selection in small
samples. *Biometrika*, *76*(2), 297-307.

IPCC. (2023). *Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (Core Writing Team, H. Lee & J. Romero, Eds.). Geneva, Switzerland: Intergovernmental Panel on Climate Change. pp. 35–115. https://doi.org/10.59327/IPCC/AR6-9789291691647

Johnson, M. S., & Speakman, J. R. (2001). Limits to sustained energy intake: V. Effect of cold-exposure during lactation in Mus musculus. *Journal of Experimental Biology*, *204*(11), 1967-1977.

Jones, T., Parrish, J. K., Peterson, W. T., Bjorkstedt, E. P., Bond, N. A., Ballance, L. T., et al. (2018). Massive mortality of a planktivorous seabird in response to a marine heatwave. *Geophysical Research Letters*, 45, 3193–3202. https://doi.org/10.1002/2017GL076164

Kacelnik, A. (1984). Central place foraging in starlings (Sturnus vulgaris). I. Patch residence time. *The Journal of Animal Ecology*, 283-299.

Kappeler, P. M., Benhaiem, S., Fichtel, C., Fromhage, L., Höner, O. P., Jennions, M. D., Kaiser, S., Krüger, O., Schneider, J. M., Tuni, C., van Schaik, J., & Goymann, W. (2023). Sex roles and sex ratios in animals. *Biological Reviews*, *98*(2), 462-480.

Kokko, H., & Jennions, M. D. (2008). Parental investment, sexual selection and sex ratios. *Journal of Evolutionary Biology*, *21*(4), 919-948.

LaZerte, S. E., Reudink, M. W., Otter, K. A., Kusack, J., Bailey, J. M., Woolverton, A., Paetkau, M., de Jong, A. and Hill, D. J. (2017). feedr and animalnexus.ca: a paired R package and user-friendly Web application for transforming and visualizing animal movement data from static stations. *Ecology and Evolution.* 7, 7884-7896. https://doi.org/10.1002/ece3.3240

Lemaître, J. F., & Gaillard, J. M. (2017). Reproductive senescence: new perspectives in the wild. *Biological Reviews*, *92*(4), 2182-2199.

Lendvai, Á. Z., Akçay, Ç., Stanback, M., Haussmann, M. F., Moore, I. T., & Bonier, F. (2018). Male parental investment reflects the level of partner contributions and brood value in tree swallows. *Behavioral Ecology and Sociobiology*, *72*, 1-11.

Levillain, A., Reichert, S., & S. Massemin. (2025). The complex relationship between high temperatues and avian breeding success: insights from a global review. *Ecology and Evolution,* 15:e71771.

Liker, A., Freckleton, R. P., Remeš, V., & Székely, T. (2015). Sex differences in parental care:
gametic investment, sexual selection, and social environment. *Evolution*, *69*(11), 2862-
2875.

Limbourg, T., Mateman, A. C., & Lessells, C. M. (2013). Opposite differential allocation by
males and females of the same species. *Biology Letters*, *9*(1), 20120835.

Mahr, K., Griggio, M., Granatiero, M., & Hoi, H. (2012). Female attractiveness affects paternal
investment: experimental evidence for male differential allocation in blue tits. *Frontiers*
in Zoology, *9*, 1-8.

Markman, S. (2014). Parental self-feeding effects on parental care levels and time allocation in
Palestine sunbirds. *PLoS One*, *9*(12), e113890.

McCafferty, D. J., Gallon, S., & Nord, A. (2015). Challenges of measuring body temperatures of
free-ranging birds and mammals. *Animal Biotelemetry*, *3*, 1-10.

McKechnie, A. E., & Wolf, B. O. (2010). Climate change increases the likelihood of catastrophic
avian mortality events during extreme heat waves. *Biology Letters*, *6*(2), 253-256.

McKechnie, A. E., Hockey, P. A., & Wolf, B. O. (2012). Feeling the heat: Australian landbirds
and climate change. *Emu-Austral Ornithology*, *112*(2), i-vii.

McKechnie, A. E., Rushworth, I. A., Myburgh, F., & Cunningham, S.J. Mortality among birds
and bats during an extreme heat event in eastern South Africa. *Austral Ecology* 46, no. 4
(2021): 687-691.

Møller, A. P., & Birkhead, T. R. (1993). Certainty of paternity covaries with paternal care in
birds. *Behavioral Ecology and Sociobiology*, *33*, 261-268.

Nolazco, S., Delhey, K., Nakagawa, S., & Peters, A. (2022). Ornaments are equally informative
in male and female birds. *Nature Communications*, *13*(1), 5917.

Nilsson, J. Å., & Nord, A. (2018). Testing the heat dissipation limit theory in a breeding
passerine. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1878),
20180652.

Nord, A., & Nilsson, J. Å. (2019). Heat dissipation rate constrains reproductive investment in a
wild bird. *Functional Ecology*, *33*(2), 250-259.

OpenAI. (2025). ChatGPT (Mar 27 version) [Large language model]. https://chat.openai.com/

Patrick, S. C., Réale, D., Potts, J. R., Wilson, A. J., Doutrelant, C., Teplitsky, C., & Charmantier,
A. (2022). Differences in the temporal scale of reproductive investment across the slow-
fast continuum in a passerine. *Ecology Letters*, *25*(5), 1139-1151.

Pérez, J. H., Krause, J. S., Chmura, H. E., Bowman, S., McGuigan, M., Asmus, A. L., Meddle, S.
L., Hunt, K. E., Gough, L., Boelman, N. T., & Wingfield, J. C. (2016). Nestling growth
rates in relation to food abundance and weather in the Arctic. *The Auk: Ornithological
Advances*, *133*(2), 261-272.

Piatt, J. F., Parrish, J. K., Renner, H. M., Schoen, S. K., Jones, T. T., et al.. (2020). Extreme
mortality and reproductive failure of common murres resulting from the northeast Pacific
marine heatwave of 2014-2016. *PloS one*, *15*(1), e0226087.

Pilastro, A., Griggio, M., & Matessi, G. (2003). Male rock sparrows adjust their breeding
strategy according to female ornamentation: parental or mating investment? *Animal
Behaviour*, *66*(2), 265-271.

Pipoly, I., Preiszner, B., Sándor, K., Sinkovics, C., Seress, G., Vincze, E., Bókony, V., & Liker,
A. (2022). Extreme hot weather has stronger impacts on avian reproduction in forests
than in cities. *Frontiers in Ecology and Evolution*, *10*, 825410.

Ratikainen, I. I., & Kokko, H. (2010). Differential allocation and compensation: who deserves
the silver spoon? *Behavioral Ecology*, *21*(1), 195-200.

Renner, S. S., & Zohner, C. M. (2018). Climate change and phenological mismatch in trophic
interactions among plants, insects, and vertebrates. *Annual Review of Ecology, Evolution,
and Systematics*, *49*(1), 165-182.

Robertson, J. K., Mastromonaco, G., & Burness, G. (2020). Evidence that stress-induced changes
in surface temperature serve a thermoregulatory function. *Journal of Experimental
Biology*, *223*(4), jeb213421.

Rodriguez, S., & Barba, E. (2016). Nestling growth is impaired by heat stress: an experimental
study in a Mediterranean great tit population. *Zoological Studies*, *55*.

Ruuskanen, S., Hsu, B. Y., & Nord, A. (2021). Endocrinology of thermoregulation in birds in a
changing climate. *Molecular and Cellular Endocrinology*, *519*, 111088.

Saeed, M., Abbas, G., Alagawany, M., Kamboh, A. A., Abd El-Hack, M. E., Khafaga, A. F., &
Chao, S. (2019). Heat stress management in poultry farms: A comprehensive overview.
Journal of thermal biology, *84*, 414-425.

Safari, I., Goymann, W., & Kokko, H. (2019). Male-only care and cuckoldry in black coucals:

does parenting hamper sex life?. *Proceedings of the Royal Society B*, *286*(1900),

20182789.

Santos, E. S. A., & Nakagawa, S. (2012). The costs of parental care: a meta-analysis of the trade-

off between parental effort and survival in birds. *Journal of Evolutionary Biology*, *25*(9),

1911-1917.

Schroeder, J., Hsu, Y. H., Winney, I., Simons, M., Nakagawa, S., & Burke, T. (2016).

Predictably philandering females prompt poor paternal provisioning. *The American*

Naturalist, *188*(2), 219-230.

Senécal, S., Riva, J. C., O'Connor, R. S., Hallot, F., Nozais, C., & Vézina, F. (2021). Poor prey

quality is compensated by higher provisioning effort in passerine birds. *Scientific*

Reports, *11*(1), 11182.

Shuker, D. M., & Kvarnemo, C. (2021). The definition of sexual selection. *Behavioral Ecology*,

32(5), 781-794.

Siefferman, L., & Hill, G. E. (2005). Male eastern bluebirds trade future ornamentation for

current reproductive investment. *Biology Letters*, *1*(2), 208-211.

Sinkovics, C., Seress, G., Pipoly, I., Vincze, E., & Liker, A. (2021). Great tits feed their nestlings

with more but smaller prey items and fewer caterpillars in cities than in forests. *Scientific*

reports, *11*(1), 24161.

Sippel, S., Meinshausen, N., Fischer, E. M., Székely, E., & Knutti, R. (2020). Climate change

now detectable from any single day of weather at global scale. *Nature Climate Change*,

10(1), 35-41.

Solano-Brenes, D., Costa-Schmidt, L. E., Albo, M. J., & Machado, G. (2021). Differential

allocation in a gift-giving spider: males adjust their reproductive investment in response

to female condition. *BMC Ecology and Evolution*, *21*, 1-15.

Speakman, J. R., & Król, E. (2010). Maximal heat dissipation capacity and hyperthermia risk:

neglected key factors in the ecology of endotherms. *Journal of Animal Ecology*, *79*(4),

726-746.

Swihart, R. K., & Johnson, S. G. (1986). Foraging decisions of American robins: somatic and

reproductive tradeoffs. *Behavioral Ecology and Sociobiology*, *19*, 275-282.

Taff, C. C., & Shipley, J. R. (2023). Inconsistent shifts in warming and temperature variability
are linked to reduced avian fitness. *Nature Communications*, *14*(1), 7400.

Tapper, S., Nocera, J. J., & Burness, G. (2020a). Heat dissipation capacity influences
reproductive performance in an aerial insectivore. *Journal of Experimental Biology*,
223(10), jeb222232.

Tapper, S., Nocera, J. J., & Burness, G. (2020b). Experimental evidence that hyperthermia limits
offspring provisioning in a temperate-breeding bird. *Royal Society Open Science*, *7*(10),
201589.

Tattersall, G. J., Roussel, D., Voituron, Y., & Teulier, L. (2016). Novel energy-saving strategies
to multiple stressors in birds: the ultradian regulation of body temperature. *Proceedings*
of the Royal Society B: Biological Sciences, *283*(1839), 20161551.

Thackeray, S. J., Henrys, P. A., Hemming, D., Bell, J. R., Botham, M. S., Burthe, S., P.
Helaouet, et al. (2016). Phenological sensitivity to climate across taxa and trophic levels.
Nature, *535*(7611), 241-245.

Van de Ven, T. M., McKechnie, A. E., & Cunningham, S. (2019). The costs of keeping cool:
behavioural trade-offs between foraging and thermoregulation are associated with
significant mass losses in an arid-zone bird. *Oecologia*, *191*, 205-215.

Vazquez, R., & Larson, D. F. (2013). Plasma protein denaturation with graded heat
exposure. *Perfusion*, *28*(6), 557–559.

Velando, A., Drummond, H., & Torres, R. (2006). Senescent birds redouble reproductive effort
when ill: confirmation of the terminal investment hypothesis. *Proceedings of the Royal*
Society B: Biological Sciences, *273*(1593), 1443-1448.

Visser, M. E., te Marvelde, L., & Lof, M. E. (2012). Adaptive phenological mismatches of birds
and their food in a warming world. *Journal of Ornithology*, *153*, 75-84.

Williams, T. D. (2018). Physiology, activity and costs of parental care in birds. *Journal of*
Experimental Biology, *221*(17), jeb169433.

Winkler, D. W., Hallinger, K. K., Pegan, T. M., Taff, C. C., Verhoeven, M. A., Chang van
Oordt, D., Stager, M., Uehling, J. J., Vitousek, M. N., Andersen, M. J., Ardia, D. R.,
Belmaker, A., Ferretti, V., Forsman, A. M., Gaul, J. R., Llambias, P. E., Orzechowski, S.
C., Shipley, J. R., Wilson, M., & Yoon, H. S. (2020). Full lifetime perspectives on the
costs and benefits of lay-date variation in tree swallows. *Ecology*, *101*(9), e03109.

Zagkle, E., Martinez-Vidal, P. A., Bauchinger, U., & Sadowska, E. T. (2022). Manipulation of

heat dissipation capacity affects avian reproductive performance and output. *Frontiers in*

Ecology and Evolution, *10*, 866182

Table 1. Model selection results for factors that explain variation in average hourly provisioning rate by adult male tree

swallows. There was only a single model with strong support (<2 ΔAIC_C).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R^2adj	ER
T_a + T_a^2 + Female Provisioning Rate + Manipulation	6	629.81	0.00	1.0	1.0	0.735	1.00
Null Model	2	702.84	73.04	0.00	1.0	0.487	$>10^5$

724   Note: K = number of parameters, AIC_C= corrected Akaike Information Criterion, ΔAIC_C = difference of focal model AIC_C compared

725   to the lowest model AIC_C; W_i = Akaike weight; Acc. W_i = Accumulated weight of models; R^2adj = Adjusted R^2; ER = Evidence ratio;

726   T_a = Average daily temperature; T_a^2 = Quadratic term for average daily temperature; Female Provisioning Rate = Average hourly

727   provisioning rate of females paired with males; Manipulation= Male manipulation (trimmed or untrimmed [Control]). The null model

728   was included for reference but did not have strong support ($\Delta AIC_C > 2$).

Table 2. Factors affecting the provisioning rate of adult male tree swallows. Estimates of fixed effects with 95% confidence intervals (CI) and *P*-values (bold signifies statistical significance) are shown.

Predictor	Estimate	95% CI	P-value
Intercept [Control]	-29.093	-46.59, -11.60	**0.001**
T_a	3.678	1.88, 5.47	**<0.001**
T_a^2	-0.099	-0.15, -0.05	**<0.001**
Female Provisioning Rate	0.482	0.36, 0.61	**<0.001**
Manipulation [Trimmed]	0.673	-1.84, 3.18	0.584

753 **Table 3. Model selection results for factors that explain variation in the body temperature of**

754 **adult male tree swallows.** There was only a single model with strong support (<2 ΔAIC$_C$).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R^2adj	ER
$T_a + T_a^2 + Ta^3 +$ Manipulation + Male Provisioning Rate	11	-21.41	**0.00**	1.00	1.00	0.091	1.00
Null Model	4	95.18	116.59	0.00	1.00	0.00	$>10^5$

757 Note: The null model was included for reference but did not have strong support (ΔAIC$_C > 2$). T_a^3

758 = Cubic term for average daily temperature, all other abbreviations are as in Table 1

770   **Table 4. Factors affecting the body temperature of adult male tree swallows.** Estimates of

771   fixed effects with 95% confidence intervals (CI) and *P*-values are shown. Bold indicates

772   statistical significance.

Predictor	Estimate	95% CI	P-value
Intercept [Control]	41.46	41.245, 41.674	**<0.001**
T_a	3.55	2.885, 4.222	**<0.001**
T_a^2	0.27	-0.314, 0.863	0.363
T_a^3	1.06	0.538, 1.598	**<0.001**
Manipulation [Trimmed]	0.24	-0.170, 0.657	0.226
Male Provisioning Rate	0.01	0.004, 0.012	**<0.001**

795   **Table 5. Model selection results for factors that explain variation in average hourly**

796   **provisioning rate of adult female tree swallows paired with trimmed or untrimmed males.**

797   There was only a single model with strong support (<2 ΔAIC_C).

Models	K	AICc	ΔAICc	Wi	Acc Wi	R²adj	ER
T_a + Manipulation + Male Provisioning Rate	5	11.07	0.00	1.00	1.00	0.765	1.00
Null Model	2	85.71	74.64	0.00	1.00	0.399	$>10^5$

800   Note: The null model was included for reference but did not have strong support ($\Delta AIC_C > 2$).

801   Abbreviations are as in Table 1

803 **Table 6. Factors affecting the provisioning rate of adult female tree swallows.** Estimates of

804 fixed effects with 95% confidence intervals (CI) and *P*-values are shown. Bold indicates

805 statistical significance.

Predictor	Estimate	95% CI	P-value
Intercept [Control]	1.73	1.415, 2.041	**<0.001**
T_a	0.02	0.008, 0.0331	**0.002**
Male Provisioning Rate	0.05	0.039, 0.062	**<0.001**
Male Manipulation [Trimmed]	-0.11	-0.326, 0.102	0.291

808   **Table 7. Model selection results for random effect structure that best explained variation in**
809   **growth trajectories for nestling tree swallows.**

Random Effect Structure	K	AICc	ΔAICc	Wi	Acc Wi	LogLik	ER
Asymptote (*A*)	9	4029.44	0.00	1.00	1.00	-2002.56	1.00
Inflection Point (*I*)	9	4366.51	337.07	0.00	1.00	-2171.10	$>10^5$
Growth Rate Constant (*K*)	9	4560.68	531.24	0.00	1.00	-2268.18	$>10^5$

810
811   Note: LogLik = Log-likelihood. Abbreviations are as in Table 1

825 **Table 8. Parameter estimates for growth trajectories in nestling tree swallows.** Estimates of

826 fixed effects with 95% confidence intervals (CI) and *P*-values are shown. Bold indicates

827 statistical significance. Manipulation refers to the manipulation of the male fathers (Trimmed or

828 Control).

Predictor	Predictor	Estimate	95% CI	P-value
Asymptote	Intercept	21.80	19.35, 24.26	**<0.001**
(*A*)	Manipulation	0.50	-0.847, 1.837	0.471
Inflection Point	Intercept	4.29	3.834, 4.739	**<0.001**
(*I*)	Manipulation	0.22	0.046, 0.397	**0.014**
Growth Rate Constant	Intercept	1.88	1.502, 2.267	**<0.001**
(*K*)	Manipulation	0.21	0.063, 0.360	**0.005**

833 **Figure 1. Male provisioning rate decreased at high temperature but did not differ between**

834 **trimmed (blue) and control (red) males.** Bands and trend lines indicate the predicted means

835 and 95% CI. Individual data points represent average provisioning rate for each individual bird

836 for a given temperature (Trimmed: n=13, Control: n=18).

840
841   **Figure 2. Male body temperature increased with ambient temperature but did not differ**

842   **between trimmed (blue) and control (red) males.** Bands and trend lines indicate the predicted

843   means and 95% CI. Individual data points represent averaged hourly T_b for each individual bird

844   at a given temperature (Trimmed, n= 4; Control, n= 9). Note, this figure displays only the subset

845   of males with temperature-sensitive PIT tags.

851   **Figure 3. Females increased provisioning rates with increasing environmental temperature**

852   **(T_a), but females partnered with trimmed males did not provision at a higher rate than**

853   **those paired with control males.** Sample sizes: females partnered with Trimmed males, n= 13;

854   females partnered with Control males, n= 17.

Figure 4. Estimated growth trajectories for nestlings from control and trimmed males.

Nestlings from trimmed males reached a similar asymptotic mass as nestlings from control

males. Growth estimates were based on a logistic growth curve using scale, midpoint, and

asymptotes. Bands represent 95% CI. Each point represents the mass of a nestling on a given

developmental day. Sample sizes: Control, n = 20 nests, n= 100 nestlings; trimmed, n = 15 nests,

n= 86 nestlings).