Comparative genomic insights into the action of suicide Thi4 thiazole synthases

Edmar R. Oliveira-Filho[1]*, Kristen Van Gelder[1], David Obe[2], Cătălin Voiniciuc[1], Mark A. Wilson[2], and Andrew D. Hanson[1,†]

[1] Horticultural Sciences Department, University of Florida, Gainesville, FL 32611, USA

[2] Department of Biochemistry & Redox Biology Center, University of Nebraska, Lincoln, NE 68588, USA

[†] Deceased

*Correspondence: ramosdeoli.edmar@ufl.edu

ABSTRACT

Suicide Thi4 thiazole synthases are mononuclear metal enzymes that form the thiazole moiety of thiamin from NAD^+, glycine, and a sulfur atom that is stripped from an active-site cysteine residue, causing enzyme inactivation. However, many prokaryotic Thi4 genes cluster on the chromosomal regions with genes encoding ThiS, ThiF, and other proteins that produce, relay, or use persulfide or thiocarboxylate sulfur. Such clusters include DUF6775 proteins, with domains of unknown function that feature a metal-binding motif. This comparative genomic evidence suggests that certain suicide Thi4s might use a persulfide or thiocarboxylate as sulfur donor instead of the active-site cysteine – i.e., that they can operate in a non-suicide mode – and that a metal cofactor reservoir supports Thi4 function. We explored both possibilities using *Escherichia coli* as a heterologous platform. A representative bacterial Thi4 (clustered with *thiS* and *thiF*) complemented an *E. coli* Δ*thiG* (thiazole auxotroph) single mutant better than a Δ*thiG* Δ*thiF* Δ*thiS* triple mutant, indicating that this Thi4 interacts with the *E. coli* sulfide transfer chain. A representative archaeal DUF6775 protein was expressed in *E. coli;* however, the bound metal was not detected in protein prepared from inclusion bodies. Collectively, this evidence indicates that suicide Thi4s do not necessarily always operate suicidally with respect to the sulfur donor or autonomously with respect to the metal cofactor.

Keywords: DUF6775, gene cluster, sulfur transfer chain, thiamin biosynthesis

1. Introduction

Plants, fungi, and certain prokaryotes produce the thiazole moiety of thiamin via a suicide Thi4 thiazole synthase (EC 2.4.2.60).[1-4] Yeast and plant suicide Thi4s use an active-site cysteine residue as sulfur donor for the reaction; whereby the cysteine is converted to dehydroalanine, inactivating the protein after a single reaction (**Figure 1a**).[2,5] The presence of this active-site cysteine distinguishes suicide Thi4s from non-suicide (catalytic) Thi4s (EC 2.4.2.59), which have a histidine or other residue in place of cysteine and use free sulfide as a sulfur donor.[6,7] Yeast and plant suicide Thi4s are mononuclear metal enzymes, with Fe(II) as the most likely cofactor. Even though metalation of purified Thi4s with Fe(II) is required to detect biochemical activity, [2,8] the metal may be loosely bound because it could not be detected in two of the four reported suicide Thi4 crystal structures.

Figure 1. The complex reaction mediated by suicide Thi4s and hypothetical alternative sulfur donors for this reaction. (a) The known reaction. NAD+ and glycine are converted to the adenylated thiazole product (ADT) in an Fe(II) (or other metal)-dependent reaction in which the sulfur donor is an active-site cysteine residue. This residue is converted to dehydroalanine (DHA). Red and blue colors show origins of the thiazole ring. The metal is weakly bound and

may dissociate from the protein during the reaction. (b) Hypothetical alternative sulfur donors, based on comparative biochemistry.

Comparative biochemistry suggests that Thi4s with an active-site cysteine could, in some cases, operate in a non-suicide mode, either by using an alternative sulfur donor that leaves the cysteine intact or by repairing the dehydroalanine residue left after the reaction (**Figure 1b**). Precedents for these hypothetical options are as follows. The bacterial thiazole synthase ThiG (EC 2.8.1.10), which is unrelated to Thi4 and forms the thiazole product via a different reaction, provides a direct evidence for an alternative sulfur donor.[1] The proximal sulfur donor for ThiG is a C-terminal thiocarboxylate group on the ThiS sulfur carrier protein; this group is formed from a persulfide borne on the sulfur carrier protein ThiI in a reaction mediated by the adenylyltransferase ThiF. The ultimate source of the persulfide sulfur, as in other sulfur-relay systems, is the action of a cysteine desulfurase on free cysteine.[1,9,10] Another precedent is 2-thiouridine biosynthesis by MnmA, for which the proximal sulfur donor is a persulfide group on the sulfur carrier protein TusE.[9] In these cases, genes encoding sulfur relay components often cluster on the chromosome with the gene for ThiG or MnmA, i.e. the enzyme that synthesizes the sulfur-containing product. A precedent for the repair of a dehydroalanine residue by a persulfide comes from the sacrificial sulfur insertase LarE from *Lactobacillus plantarum*; in this case the persulfide group is on a small-molecule thiol (Coenzyme-A), not on a protein carrier.[11]

Comparative biochemistry also suggests that suicide Thi4s, like other metalloenzymes, could be supported by metallochaperone proteins that deliver metal ions to their specific target and maintain metal homeostasis. Examples include the iron-trafficking chaperone HcdG, which delivers iron to the [Fe]-hydrogenase Hmd,[12] and the nickel-binding protein HypA, which transfers a nickel ion to the catalytic core of the [NiFe]-hydrogenase.[13] Like the sulfur-relay genes above, the genes encoding these chaperones are often clustered in operonic arrangements with the genes for the enzymes they service.[12,13]

These comparative biochemical and genomic considerations led us to briefly survey the genomic context of representative prokaryote suicide Thi4 genes. We saw obvious clustering with genes coding for various sulfur-relay proteins and their client biosynthetic enzymes, as well as with genes encoding for DUF6775, a domain of unknown function with predicted metal-binding residues. Applying the maxim[9] that: "A genetic, genomic or proteomic…indication that…a protein homologous to another sulfur carrier protein has a role in a particular biological

process should trigger a careful scrutiny for the involvement of a persulfide group in the bio-chemistry of the process", and extending this maxim to metallochaperones, we therefore made a fuller comparative genomic analysis and pilot genetic and biochemical tests to explore possible functional roles of sulfur-relay proteins and DUF6775 in the biochemistry of prokaryote suicide Thi4s.

2. Materials and methods

2.1. Bioinformatics tools and databases

Prokaryote metagenomes and genomes were searched primarily in the Integrated Microbial Genomes & Microbiomes system (IMG/M) at the DOE Joint Genome Institute (https://img.jgi.doe.gov/m/)[14] and secondarily in GenBank. The query was the suicide Thi4 protein sequence from Deltaproteobacteria bacterium RBG_16_54_11 (GenBank OGP80537); gene neighborhoods around similar suicide Thi4 genes were displayed in IGM/M via the 'Show neighborhood regions with the same top COG hit (via top homolog)' tool. Functional gene annotations were made or confirmed using the NCBI Conserved Domain Database (CDD).[15]

2.2. Plasmid constructs

All molecular manipulations followed standard protocols[16] or reagent/kit manufacturers' instructions. Codon-optimized nucleotide sequences with added compatible restriction sites were ordered from GenScript (Piscataway, NJ). Restriction cloning was used to clone genes using the following restriction enzymes (New England Biolabs, Ipswich, MA): *Eco*RI/*Xba*I for pBAD24; *Nde*I/*Hin*dIII for pET28b(+). Digestion products were gel-purified using GeneJET Gel Extraction Kits (Thermo Fisher Scientific, Waltham, MA), and assembled by T4 DNA Ligase (Thermo Fisher Scientific, Waltham, MA). Ligation products were transferred to *Escherichia coli* Top10 via electroporation using an *E. coli* Pulser apparatus (Bio-Rad Laboratories, Hercules, CA). Candidate clones (antibiotic-resistant) were then PCR-screened to select those with successful assembly. Oligonucleotide sequences are listed in **Table S1**. Recombinant plasmids were purified using GeneJET Plasmid Miniprep Kits (Thermo Fisher Scientific, Waltham, MA) and sequence-verified by Sanger or whole plasmid sequencing (Genewiz, South Plainfield, NJ).

2.3. Escherichia coli gene knockouts

E. coli MG1655 Δ*thiG* and Δ*thiF* Δ*thiS* Δ*thiG* mutants were generated through recombineering.[17] For the Δ*thiG* single knockout, the deletion cassette was amplified by PCR from *E. coli* Keio

JW5549-1 genomic DNA. To generate the ΔthiF ΔthiS ΔthiG mutant, a DNA fragment of the region encoding for the C-terminus of ThiE through to ThiF, ThiS, ThiG, and for the N-terminus of ThiH was amplified from *E. coli* MG1655 and cloned into pGE using NEBuilder HiFi DNA Assembly Cloning Kit (New England Biolabs, Ipswich, MA), generating pGE_*thiFthiSthiG*. A spectinomycin resistance cassette (Sm^R) flanked by FRT sites was amplified from pTs and inserted into pGE_*thiFthiSthiG* using the same kit. The Sm^R cassette replaced *thiF*, *thiS*, and *thiG,* leaving 200-bp flanking sequences (homology arms) for *thiE* and *thiH* upstream and downstream of Sm^R, respectively. Each deletion cassette was transferred via electroporation to *E. coli* MG1655 harboring pKD46. Recombinants were selected on LB (tryptone, 10 g/L; yeast extract, 5 g/L; and NaCl, 5 g/L) with kanamycin (50 μg/mL) or spectinomycin (100 μg/mL). Candidate clones were screened by colony PCR followed by agarose gel electrophoresis. pKD46 was purged from PCR-verified clones by overnight incubation at 42°C.

2.4 Functional complementation assays

MB1, and cOb THI4 encoding sequences were recoded for expression in *E. coli* (**Table S2**) and synthesized by GenScript (Piscataway, NJ). pBAD24 constructs harboring *thi4* were transformed into *E. coli* MG1655 ΔthiG and ΔthiF ΔthiS ΔthiG. pBAD24::Ta_*thi4* was used as a benchmark.[7]

To verify the production of soluble THI4 protein, recombinant MG1655 was cultured at 37°C and 250 rpm in 25 mL MOPS minimal medium[18] supplemented with 0.2% glycerol, micronutrients,[19] 100 μg/mL carbenicillin, and 100 nM thiamin. When OD_{600} reached ~0.5, cultures were cold-shocked in an ice bath for 30 min, L-Arabinose was added to a final concentration of 0.02% (w/v), before an overnight incubation at 18°C, 180 rpm. Cells were harvested by centrifugation (5000*g*, 15 min, 4°C) and stored at -80°C. Cell pellets were sonicated in 100 mM potassium phosphate (pH 7.2), 2 mM β-mercaptoethanol buffer. Protein concentration was determined by Bradford assay, and protein profiles (10 μg per sample) were visualized by SDS-PAGE on 12.5% gels stained with GelCode Blue Safe Protein Stain (Thermo Fisher Scientific, Waltham, MA).

Functional complementation was evaluated on the MOPS minimal medium with the supplements described above. Single colonies were grown in 3 mL of medium containing 100 nM thiamin for 24 hours. These were sub-cultured twice on thiamin-free MOPS supplemented with 1 mM cysteine, and 0.02 % (w/v) L-arabinose. Cells from the third culture were washed three times with basal MOPS (i.e., MOPS medium minus $MgSO_4$, $FeSO_4$, micronutrients, or thiamin) and used to prepare serial 10-fold dilutions starting with an OD_{600} of 1.5. These were

spot-plated on MOPS agar plates with 1 mM cysteine, 0.02% (w/v) L-arabinose, with or without 100 nM thiamin. Plates were incubated at 37°C for 3-4 days.

2.5. Recombinant DUF6775 production and analysis

Selected DUF6775-containing sequences were codon-optimized for *E. coli* (**Table S2**) and synthesized by GenScript (Piscataway, NJ). pET28b(+) constructs were transferred to *E. coli* BL21 (DE3) by electroporation. Recombinant BL21 was cultured on 50 mL LB plus kanamycin at 37°C, 250 rpm. When OD_{600} reached ~0.5, cultures were cold-shocked in an ice bath for 30 min, induced with 100 µM IPTG, and incubated overnight at 18°C, 180 rpm. Cells were harvested by centrifugation (5000*g*, 15 min, 4°C) and stored at -80°C. Cell pellets were sonicated in 25 mM HEPES-NaOH (pH 7.5), 100 mM KCl, 0.5% Triton X-100, and 5 mM dithiothreitol. Non-soluble fractions were collected by centrifugation (25000g, 30 min, 4°C), washed three times with the same buffer with 1% Triton X-100, with a short sonication cycle (3x 10s on, 50s off) between washes (25000*g*, 15 min, 4°C); and one time with 25 mM HEPES-NaOH (pH 7.5), 1 M KCl, 5 mM dithiothreitol. The obtained pellet was then resuspended in 25 mM HEPES-NaOH (pH 7.5), 100 mM KCl, and 5 mM dithiothreitol. Protein profiles were visualized by SDS-PAGE on 12.5% gels stained with GelCode Blue Safe Protein Stain (Thermo Fisher Scientific). Protein concentration was adjusted to 1 mg/mL by measuring absorbance at 280 nM and calculating concentrations based on the peptide's extinction coefficient (ProtParam).

Metals associated with recombinant DUF6775 recovered from inclusion bodies were measured using inductively coupled plasma-mass spectrometry (ICP-MS) using an Agilent 7500 cx ICP-MS equipped with a 96-well plate autosampler.[20] Samples were lyophilized and then incubated overnight with 100 µL of 70% trace metal-grade nitric acid at 65°C. After acid digestion for 14 hours, the samples were cooled to room temperature, centrifuged, and diluted 10-fold with 2% nitric acid spiked with 50 µg/L gallium (Ga) as the internal ICP-MS standard. The final diluted sample was injected into the ICP-MS autosampler for analysis, following the method described by Malinouski et al.[21] ICP-MS measurements were performed in mix reaction mode (3.5 mL H_2 and 1.5 mL He per minute) to eliminate polyatomic interferences in the collision cell. The samples were loaded using an autosampler (ESI, Omaha, NE) and injected through a 6-port injection valve with a 125 µL sample loop at a flow rate of 55 µL/min. The following ICP-MS operating conditions were used: Ar carrier flow: 1.0 L/min; Ar make-up flow: 0.1–0.2 L/min; forward power: 1,500 W; Ar plasma gas: 15 L/min; Ar auxiliary gas: 1 L/min. Metal concentrations were determined using an external calibration curve, with Ga (50 ppb) as the

internal standard.[22] Element ratios were calculated as the ratio of the element concentration to the protein concentration. All solutions for metal analysis were prepared using metal-grade water and nitric acid.

2.6 Structural model prediction

The structure of the conserved hypothetical archaeal DUF6775 protein was predicted using the AlphaFold 3 server.[23] The resulting model exhibited a high overall confidence, with a mean predicted Local Distance Difference Test (pLDDT) score of 94.6. Regions of lower confidence (70 > pLDDT > 50) were localized to helices α7 and α8, as well as loop regions corresponding to loops 8, 10, 15, and 17. Proteins with experimentally determined structures similar to the predicted structure of DUF6775 were identified using the DALI server.[24] Two structural homologs were identified: the zinc-dependent peptidase from *Methanocorpusculum labreanum* (PDB: 3LMC, UniProt: A2SQK8) and Archaemetzincin from *Methanopyrus kandleri* (PDB: 2X7M),[25] with DALI Z-scores of 13.9 and 14.3, respectively. These homologs were used as structural templates to infer the likely metal-binding site in DUF6775. The identity of the bound metal in 2X7M has been experimentally verified via anomalous difference Fourier maps,[25] although the methods used to assign the metals in 3LMC are unknown, as there is no supporting publication.

All predicted models and structural comparisons were visualized and analyzed using UCSF ChimeraX version 1.9.[26]

3. Results and discussion

3.1. Neighboring-gene analyses of suicide Thi4s

The suicide Thi4 gene from Deltaproteobacteria bacterium RBG_16_54_11 is typical of those seen in our initial survey to cluster with sulfur-relay and DUF6775 genes (**Figure 2a**). This Thi4 gene was used as query to extract gene neighborhood data from the IMG/M system[14] and GenBank for >400 suicide Thi4 genes from bacteria and archaea belonging to at least 26 different phyla; **Figure 2a** shows representative cases. From these data, we identified genes that cluster consistently in the ten-gene window around the Thi4 gene (**Figure 2b** and **Table S3**).

Figure 2. Gene neighborhoods around suicide Thi4s. (a) Representative examples from bacteria and archaea. (b) Frequencies (percent of metagenomes and genomes surveyed) with which genes for DUF6775, sulfur-generating enzymes, sulfur-relay proteins, sulfur-utilizing enzymes, and thiamin synthesis and salvage enzymes (ThiC, ThiD, ThiE, ThiL, ThiM), the carrier ThiW, and the regulator ThiR recur in the 10-gene window around Thi4. Abbreviations: ThiF, adenylyltransferase catalyzing adenylylation of the ThiS C terminus; CysK, cysteine synthase; ThiS, sulfur carrier; NifS, cysteine desulfurase; ThiI, sulfurtransferase; CysI, sulfite reductase; NifU, FeS cluster assembly scaffold protein; YiiM, MOSC (MOCO sulfurase C-terminal) domain protein; CysE, serine O-acetyltransfer-ase; LarE, sacrificial sulfur insertase; CysM, S-sulfocysteine synthase; FdhD, sulfurtransferase; MnmA, tRNA U34 2-thiouridine synthase; Fe-S, iron-sulfur cluster protein; BioB, biotin synthase; HcyBio, L-aspartate semialdehyde sulfurtransferase; SseA, 3-mercaptopyruvate sulfurtransferase.

Genes encoding DUF6775 were the most frequently clustered with Thi4, followed by a full range of genes for sulfur generation and relay proteins including ThiF and ThiS, and by genes for proteins that use the sulfur delivered by relays (MnmA, BioB, and various Fe-S proteins) (**Figure 2b**). These genes all clustered with Thi4 at similar or higher frequencies than thiamin synthesis, salvage, transport, and regulation genes (ThiC, ThiD, ThiE, ThiL, ThiM, ThiW, and ThiR) (**Figure 2b**). None of the genomes from **Figure 2a** encoded ThiG (**Table S4**), indicating that the Thi4-clustered ThiS and ThiF proteins in these genomes cannot be servicing ThiG. These patterns imply functional relationships between suicide Thi4s and sulfur transfer components at least as strong as those with other thiamin-related enzymes, carriers, and regulators.[27] Given the *a priori* plausibility of non-suicidal Thi4 action and of the dependence of Thi4s on a metal chaperone (see Introduction), the above comparative genomic evidence led us to run pilot experiments to test both possibilities.

3.2. Complementation tests for interaction of Thi4s with E. coli sulfur carrier protein ThiS

A straightforward possibility suggested by the clustering of Thi4 genes with ThiS and ThiF genes (**Figure 2**) is that, like ThiG, Thi4 can accept sulfur from the ThiS thiocarboxylate group formed by ThiF.[1] We therefore sought Thi4s that are (i) encoded by genes clustered with *thiS* and *thiF* genes, (ii) from mesophiles (and hence likely to function in *E. coli*), and (iii) that express well as soluble proteins in *E. coli*. Thi4s from *Methanobacterium* sp. MB1 (MB1) and *Candidatus* Omnitrophica bacterium (cOb) isolate bin.255 met these criteria (**Figure 2a** and **Figure S1**). These two Thi4s were then tested for their ability to complement an *E. coli* single Δ*thiG* mutant or a triple Δ*thiG* Δ*thiF* Δ*thiS* mutant. The logic was that, to the extent that the Thi4 can accept sulfur from the alien *E. coli* ThiS, complementation would be stronger in the single mutant than the triple mutant. The non-suicide Thi4 from *Thermovibrio ammonificans* (Ta) was included as a benchmark because it uses free sulfide as sulfur donor[7] and is consequently predicted to perform equally well in the single and triple mutants.

There was evidence for stronger complementation in the single mutant for cObThi4 but not for MB1Thi4, or – as expected – for TaThi4. Thus, as cells were serially passaged without thiamin, the growth of cells expressing cOb4Thi4 became significantly weaker in the triple mutant relative to the single mutant, which was not the case for MB1Thi4 or TaThi4 (**Figure 3a**). Weaker complementation by cObThi4 was also visible on agar plates using thiamin-depleted cells used as inoculum (**Figure 3b**). Because the native ThiS proteins of MB1 and cOb are similar to *E. coli* ThiS only in the C-terminal quarter (**Figure S2**), and ThiS family sulfur carriers

can be specific to particular sulfur acceptor enzymes,[28] lack of stronger complementation in the single mutant might arise from failure of MB1Thi4 to interact functionally with *E. coli* ThiS. Therefore, the negative result with MB1Thi4 does not necessarily exclude the ThiS of MB1 as a physiological sulfur donor for MB1Thi4 in the native host.

Figure 3. Complementation tests of interaction between two suicide Thi4s and the heterologous ThiF-ThiS system of *E. coli*. The *E. coli* host strains were deleted for *thiG* alone or for *thiF* and *thiS* as well as *thiG*. (a) Growth at 24 h of the liquid cultures minus thiamin used to inoculate the agar plate complementation tests. Error bars are SEM of three biological replicates. (b) Growth at 4 days on the complementation plates. Abbreviations: Vector, vector alone; TaThi4, non-suicide Thi4 from *Thermovibrio ammonificans* (benchmark); MB1Thi4, suicide Thi4 from *Methanobacterium* sp. MB1; cObThi4, suicide Thi4 from *Candidatus* Omnitrophica bacterium isolate bin.255. Note the poorer growth of cells expressing cObThi4 in the triple mutant host compared to the single mutant host in liquid medium minus thiamin (**, P<0.01) and on plates minus or plus thiamin, and the lack of such a difference for single and triple mutant cells expressing TaThi4 or MB1Thi4. Similar results were obtained in three independent experiments.

3.3. Predicted structure and metal-binding tests of a DUF6775 protein

DUF6775 genes cluster with Thi4s in two-thirds of the genomes analyzed (**Figure 2b**) and are annotated as putative metallopeptidases. Representative DUF6675 proteins have conserved cysteine and histidine residues for potential metal binding (**Figure S3**), so we analyzed their structure with AlphaFold 3. The proteins chosen for analysis came from a mesophile archaeon (metagenome SAMEA7390951, SAMEA). The AlphaFold DUF6775 model has a three-layer αβα sandwich fold (**Figure 4**) characteristic of metallopeptidases within the collagenase catalytic domain superfamily (CATH 3.40.390.10). Related proteins with known structures include PDB 3LMC (Northeast Structural Genomics Consortium Target MuR16) and 2X7M,[25] which contain bound metals (**Figure 4**). We note that the assignment of one of the metal sites in 3LMC as Fe is surprising given its 3xCys-1xHis tetrahedral coordination geometry and the assignment of Zn to the corresponding site of 2X7M. Unlike canonical archaemetzincins, which contain a HEXXHXXGX$_3$CX$_4$CXMX$_{17}$CXXC zinc-binding motif comprising four strictly conserved cysteine residues,[25] DUF6775 has a distinct predicted metal binding site with three conserved cysteines and one histidine (**Figure 4**) and it lacks the classical HEXXHXXGXXH/D motif that coordinates the second Zn^{2+} ion. These differences suggest a metal coordination environment in DUF6775 that is different from known zinc metallopeptidases with related structures.

Figure 4. Predicted structure of DUF6775 and comparison to experimentally determined metallopeptidases. (a) Superposition of the AlphaFold3-predicted structure of DUF6775 (pink) and PDB 3LMC (green). 3LMC contains two metals: Fe (orange) and Zn (grey), although the Fe site is unusual and requires validation. (b) The coordination environment in a close-up inset. Note that the His residues that coordinate the second Zn in 3LMC are absent in DUF6775. (c) Superposition of DUF6775 (pink) and PDB 2X7M, the structure of archaemetzincin (AmzA) from Methanopyrus kandleri.[26] The predicted metal in DUF6775 is orange, and the experimentally determined Zn in 2X7M are grey.

To experimentally identify any bound metals, a DUF6775 protein from the archaeal metagenome SAMEA7390951 was recovered from the insoluble fraction of *E. coli* lysates. An inclusion body preparation protocol yielded samples considerably enriched in DUF6775 protein with ~33 kDa (**Figure S4**). However, analysis of the inclusion body samples via inductively coupled plasma mass spectrometry (ICP-MS) did not identify any bound metals relative to the basal buffer control (**Table S5**).

4. Conclusion

This work presents comparative genomic evidence consistent with the possibility that, like ThiG thiazole synthase, at least some suicide Thi4s can use a sulfur relay system to provide the sulfur atom needed for the thiazole ring, and thus are not necessarily doomed to sudden death. Pilot experimental data consistent with this possibility are also presented. While not definitive, these data warrant future, deeper investigation of the role of ThiS family proteins as potential sulfur donors for Thi4s, e.g. by constructing and expressing in *E. coli* synthetic operons containing a suicide Thi4 and the ThiS, ThiF, and other sulfur relay genes with which the Thi4 gene clusters in its native host (see **Figure 2a**).

Here, we also unveiled a close genomic association between suicide Thi4s and DUF6775 family proteins, which remain uncharacterized but feature similar motifs to metal-dependent peptidases. Although we did not detect any metals bound to an archaeal DUF6775 tested in our pilot analysis, the protein was only found in inclusion bodies rather than in a soluble form that could be purified. Recombinant proteins in inclusion bodies may be poorly folded or otherwise incapable of binding metal, so the lack of ICP-MS-detectable metal in our samples does exclude the possibility that DUF6775 acts as a physiological cofactor for Thi4. It is worth noting that non-suicide Thi4s cluster strongly with genes encoding TRASH proteins known to bind various metals.[7] Therefore, these exploratory data warrant further biochemical and genetic investigation to elucidate the functions of prokaryotic Thi4s and of their neighboring genes that encode uncharacterized domains.

Funding

This work was supported primarily by the U.S. Department of Energy, Office of Science, Basic Energy Sciences under Award DE-SC0020153 awarded to A.D.H. and M.A.W.; by USDA NIFA Hatch project FLA-HOS-005796, and an Endowment from the C.V. Griffin, Sr. Foundation.

Author contributions

Edmar R. Oliveira-Filho: Conceptualization, Methodology, Investigation, Visualization, Writing – original draft. **Kristen Van Gelder:** Methodology, Investigation, Writing – review & editing. **David Obe:** Methodology, Investigation, Writing – review & editing. **Cătălin Voiniciuc**: Writing – review & editing. **Mark A. Wilson:** Conceptualization, Methodology, Investigation, Visualization, Writing – review & editing, Funding acquisition. **Andrew D. Hanson:** Conceptualization, Writing – original draft, Funding acquisition, Project administration.

Conflict of Interest Statement

The authors declare no competing financial interest.

Supporting Information

Figure S1. SDS-PAGE analysis of the soluble fraction of Δ*thiG* and Δ*thiF* Δ*thiG* Δ*thiS E. coli* strains expressing selected *thi4.*

Figure S2. Alignment of *E. coli* ThiS with representative ThiS family or ThiS-like proteins from Figure 2a whose genes cluster with suicide Thi4 genes.

Figure S3. Alignment of representative DUF6775 proteins from **Figure 2a** whose genes cluster with suicide *thi4* genes.

Figure S4. SDS-PAGE analysis of recombinant DUF6775 protein from archaeal metagenome SAMEA7390951.

Table S1. Primer sequences used in this work.

Table S2. Recoded nucleotide sequences of Thi4 and DUF6775 proteins tested in this study.

Table S3. Sulfide-related, thiamin-related, and DUF6775 genes in the 10-gene window around the suicide Thi4 gene in each of 40 genomes representative of the >400 genomes analyzed.

Table S4. Occurrence of Genes Coding for ThiG, ThiS, and ThiF Family Proteins in the Seven Representative Genomes from Figure 2a.

Table S5. Inductively Coupled Plasma–Mass Spectrometry (ICP-MS) Analysis of DUF6775 Protein Samples.

Acknowledgments

We dedicate this work to the memory of Andrew D. Hanson, whose leadership, wisdom, and generosity enriched the lives of all who worked with him.

We thank Dr. Javier Seravalli of the University of Nebraska Spectroscopy and Biophysics Core Facility for performing the ICP-MS measurements.

References

(1) Jurgenson, C. T.; Begley, T. P.; Ealick, S. E. The structural and biochemical foundations of thiamin biosynthesis. *Annu. Rev. Biochem.* **2009**, *78*, 569–603, DOI: 10.1146/annurev.biochem.78.072407.102340.

(2) Chatterjee, A.; Abeydeera, N. D.; Bale, S.; Pai, P. J.; Dorrestein, P. C.; Russell, D. H.; Ealick, S. E.; Begley, T. P. *Saccharomyces cerevisiae* THI4p is a suicide thiamine thiazole synthase. *Nature* **2011**, *478*, 542-546, DOI: 10.1038/nature10503.

(3) Woodward, J. B.; Abeydeera, N. D.; Paul, D.; Phillips, K.; Rapala-Kozik, M.; Freeling, M.; Begley, T. P.; Ealick, S. E.; McSteen, P.; Scanlon, M. J. A maize thiamine auxotroph is defective in shoot meristem maintenance. *Plant Cell* **2010**, *22*, 3305-3317, DOI: 10.1105/tpc.110.077776.

(4) Hwang, S.; Cordova, B.; Chavarria, N.; Elbanna, D.; McHugh, S.; Rojas, J.; Pfeiffer, F.; Maupin-Furlow, J. A. Conserved active site cysteine residue of archaeal THI4 homolog is essential for thiamine biosynthesis in *Haloferax volcanii*. *BMC Microbiol.* **2014**, *14*, 1-14, DOI: 10.1186/s12866-014-0260-0.

(5) Joshi, J.; Beaudoin, G. A.; Patterson, J. A.; García-García, J. D.; Belisle, C. E.; Chang, L. Y.; Li, L.; Duncan, O.; Millar, A. H.; Hanson, A. D. Bioinformatic and experimental evidence for suicidal and catalytic plant THI4s. *Biochem. J.* **2020**, *477*, 2055-2069, DOI: 10.1042/BCJ20200297.

(6) Eser, B. E.; Zhang, X.; Chanani, P. K.; Begley, T. P.; Ealick, S. E. From suicide enzyme to catalyst: the iron-dependent sulfide transfer in *Methanococcus jannaschii* thiamin thiazole biosynthesis. *J. Am. Chem. Soc.* **2016**, *138*, 3639-3642. DOI: 10.1021/jacs.6b00445.

(7) Joshi, J.; Li, Q.; García-García, J. D.; Leong, B. J.; Hu, Y.; Bruner, S. D.; Hanson, A. D. Structure and function of aerotolerant, multiple-turnover THI4 thiazole synthases. *Biochem. J.* **2021**, *478*, 3265-3279, DOI: 10.1042/BCJ20210565.

(8) Paul, D. Structural characterization of enzymes involved in biosynthesis of thiamin and salvage of uridine. Ph.D. Thesis, Cornell University, **2010**, https://ecommons.cornell.edu/bitstream/1813/17205/1/Paul,%20Debamita.pdf.

(9) Mueller, E. G. Trafficking in persulfides: delivering sulfur in biosynthetic pathways. *Nat. Chem. Biol.* **2006**, *2*, 185-194, DOI: 10.1038/nchembio779.

(10) Chatterjee, S.; Hausinger, R. P. Sulfur incorporation into biomolecules: Recent advances. *Crit. Rev. Biochem. Mol. Biol.* **2022**, *57*, 461-476, DOI: 10.1080/10409238.2022.2141678.

(11) Fellner, M.; Rankin, J. A.; Desguin, B.; Hu, J.; Hausinger, R. P. Analysis of the active site cysteine residue of the sacrificial sulfur insertase LarE from *Lactobacillus plantarum*. *Biochemistry* **2018**, *57*, 5513-5523, DOI: 10.1021/acs.biochem.8b00601.

(12) Arriaza-Gallardo, F. J.; Zheng, Y.C.; Gehl, M.; Nomura, S.; Fernandes-Queiroz, J. P.; Shima, S. [Fe]-Hydrogenase, cofactor biosynthesis and engineering. *ChemBioChem* **2023**, *24*, e202300330, DOI: 10.1002/cbic.202300330.

(13) Chan, K. H.; Lee, K. M.; Wong, K. B. Interaction between hydrogenase maturation factors HypA and HypB is required for [NiFe]-hydrogenase maturation. *PloS One* **2012**, *7*, e32592, DOI: 10.1371/journal.pone.0032592.

(14) Chen, I. M .A.; Chu, K.; Palaniappan, K.; Ratner, A.; Huang, J.; Huntemann, M.; Hajek, P.; Ritter, S. J.; Webb, C.; Wu, D.; Varghese, N. J. The IMG/M data management and analysis system v. 7: content updates and new features. *Nucleic Acids Res.* **2023**, *51(D1),* D723-D732, DOI: 10.1093/nar/gkac976.

(15) Wang, J.; Chitsaz, F.; Derbyshire, M.K.; Gonzales, N.R.; Gwadz, M.; Lu, S.; Marchler, G. H.; Song, J. S.; Thanki, N.; Yamashita, R. A.; Yang, M.; Zhang, D.; Zheng, C.; Lanczycki, C. J.; Marchler-Bauer, A. The conserved domain database in 2023. *Nucleic Acids Res.* **2022**, 51(D1):D384-D388. DOI: 10.1093/nar/gkac1096.

(16) Sambrook, J.; Russell, D. W. Molecular Cloning: A Laboratory Manual, 3rd edition; Cold Spring Harbor Laboratory Press: New York, 2001.

(17) Datsenko K. A.; Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A.* **2000**, 97(12):6640-5. DOI: 10.1073/pnas.120163297.

(18) Stewart, V.; Parales, J. Identification and expression of genes *narL* and *narX* of the *nar* (nitrate reductase) locus in *Escherichia coli* K-12. *J. Bacteriol.* **1988**, 170, 1589–1597, DOI: 10.1128/jb.170.4.1589-1597.1988.

(19) Neidhardt, F. C.; Bloch, P. L.; Smith, D. F. Culture medium for enterobacteria. *J. Bacteriol.* **1974**, 119 (3), 736– 747, DOI: 10.1128/jb.119.3.736-747.1974.

(20) Thomas, R. *Practical Guide to ICP-MS: A Tutorial for Beginners, Third Edition.* CRC Press, 2013, DOI: 10.1201/b14923.

(21) Malinouski, M.; Hasan, N.M.; Zhang, Y.; Seravalli, J.; Lin, J.; Avanesov, A.; Lutsenko, S.; Gladyshev, V. N. Genome-wide RNAi ionomics screen reveals new genes and regulation of human trace element metabolism. *Nat Commun.* **2014**, 5 (3301), DOI: 10.1038/ncomms4301.

(22) Nývltová, E.; Dietz, J. V.; Seravalli, J.; Khalimonchuk, O.; Barrientos, A. Coordination of metal center biogenesis in human cytochrome c oxidase. *Nat Commun.* **2022**, 13(1):3615. DOI: 10.1038/s41467-022-31413-1.

(23) Abramson, J.; Adler, J.; Dunger, J.; Evans, R.; Green, T.; Pritzel, A.; Ronneberger, O.; Willmore, L.; Ballard, A. J.; Bambrick, J.; Bodenstein, S. W.; Evans, D. A.; Hung, C. C.; O'Neill, M.; Reiman, D.; Tunyasuvunakool, K.; Wu, Z.; Žemgulytė, A.; Arvaniti, E.; Beattie, C.; Bertolli, O.; Bridgland, A.; Cherepanov, A.; Congreve, M.; Cowen-Rivers, A. I.; Cowie, A.; Figurnov, M.; Fuchs, F. B.; Gladman, H.; Jain, R.; Khan, Y. A.; Low, C. M. R.; Perlin, K.; Potapenko, A.; Savy, P.; Singh, S.; Stecula, A.; Thillaisundaram, A.; Tong, C.; Yakneen, S.; Zhong, E, D.; Zielinski, M.; Žídek, A.; Bapst, V.; Kohli, P.; Jaderberg, M.; Hassabis, D.; Jumper, J. M. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **2024**, 630(8016):493-500. DOI: 10.1038/s41586-024-07487-w.

(24) Holm, L.; Laiho, A.; Törönen, P.; Salgado, M. DALI shines a light on remote homologs: One hundred discoveries. *Protein Sci.* **2023**, 32(1):e4519. DOI: 10.1002/pro.4519.

(25) Waltersperger, S.; Widmer, C.; Wang, M.; Baumann, U. Crystal structure of archaemetzincin AmzA from *Methanopyrus kandleri* at 1.5 A resolution. *Proteins* **2010**, 78(12):2720-3. DOI: 10.1002/prot.22777.

(26) Meng, E. C.; Goddard, T. D.; Pettersen, E. F.; Couch, G. S.; Pearson, Z. J.; Morris, J. H.; Ferrin, T. E. UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci.* **2023**, 32(11):e4792. DOI: 10.1002/pro.4792.

(27) Fischbach, M.; Voigt, C. A. Prokaryotic gene clusters: a rich toolbox for synthetic biology. *Biotechnol. J. 2010*, *5*, 1277-1296, DOI: 10.1002/biot.201000181.

(28) Shigi, N. Biosynthesis and functions of sulfur modifications in tRNA. *Front. Genet. 2014*, *5*, 67, DOI: 10.3389/fgene.2014.00067.

Figure 1. The complex reaction mediated by suicide Thi4s and hypothetical alternative sulfur donors for this reaction. (a) The known reaction. NAD^+ and glycine are converted to the adenylated thiazole product (ADT) in an Fe(II) (or other metal)-dependent reaction in which the sulfur donor is an active-site cysteine residue. This residue is converted to dehydroalanine (DHA). Red and blue colors show origins of the thiazole ring. The metal is weakly bound and may dissociate from the protein during the reaction. (b) Hypothetical alternative sulfur donors, based on comparative biochemistry.

Figure 2. Gene neighborhoods around suicide Thi4s. (a) Representative examples from bacteria and archaea. (b) Frequencies (percent of metagenomes and genomes surveyed) with which genes for DUF6775, sulfur-generating enzymes, sulfur-relay proteins, sulfur-utilizing enzymes, and thiamin synthesis and salvage enzymes (ThiC, ThiD, ThiE, ThiL, ThiM), the carrier ThiW, and the regulator ThiR recur in the 10-gene window around Thi4. Abbreviations: ThiF, adenylyltransferase catalyzing adenylylation of the ThiS C terminus; CysK, cysteine synthase; ThiS, sulfur carrier; NifS, cysteine desulfurase; ThiI, sulfurtransferase; CysI, sulfite reductase; NifU, FeS cluster assembly scaffold protein; YiiM, MOSC (MOCO sulfurase C-terminal) domain protein; CysE, serine *O*-acetyltransferase; LarE, sacrificial sulfur insertase; CysM, *S*-sulfocysteine synthase; FdhD, sulfurtransferase; MnmA, tRNA U34 2-thiouridine synthase; Fe-S, iron-sulfur cluster protein; BioB, biotin synthase; HcyBio, L-aspartate semialdehyde sulfurtransferase; SseA, 3-mercaptopyruvate sulfurtransferase.

Figure 3. Complementation tests of interaction between two suicide Thi4s and the heterologous ThiF-ThiS system of *E. coli*. The *E. coli* host strains were deleted for *thiG* alone or for *thiF* and *thiS* as well as *thiG*. (a) Growth at 24 h of the liquid cultures minus thiamin used to inoculate the agar plate complementation tests. Error bars are SEM of three biological replicates. (b) Growth at 4 days on the complementation plates. Abbreviations: Vector, vector alone; TaThi4, non-suicide Thi4 from *Thermovibrio ammonificans* (benchmark); MB1Thi4, suicide Thi4 from *Methanobacterium* sp. MB1; cObThi4, suicide Thi4 from Candidatus Omnitrophica bacterium isolate bin.255. Note the poorer growth of cells expressing cObThi4 in the triple mutant host compared to the single mutant host in liquid medium minus thiamin (**, P<0.01) and on plates minus or plus thiamin, and the lack of such a difference for single and triple mutant cells expressing TaThi4 or MB1Thi4. Similar results were obtained in three independent experiments.

Figure 4. Predicted structure of DUF6775 and comparison to experimentally determined metallopeptidases. (a) Superposition of the AlphaFold3-predicted structure of DUF6775 (pink) and PDB 3LMC (green). 3LMC contains two metals: Fe (orange) and Zn (grey), although the Fe site is unusual and requires validation. (b) The coordination environment in a close-up inset. Note that the His residues that coordinate the second Zn in 3LMC are absent in DUF6775. (c) Superposition of DUF6775 (pink) and PDB 2X7M, the structure of archaemetzincin (AmzA) from *Methanopyrus kandleri*.[26] The predicted metal in DUF6775 is orange, and the experimentally determined Zn in 2X7M are grey.