Reduced legacy precipitation decreases microbial community growth efficiency and alters soil organic carbon in a California grassland

Linnea K. Hernandez[1], Nicole Didonato[2], Ljiljana Pasa-Tolic[2], Pete F. Chuckran[3], Mary K. Firestone[3,4], Ella T. Sieradzki[3,5], Mengting Maggie Yuan[3,6], Katerina Estera-Molina[1], Jeffrey Kimbrel[1], Paul Dijkstra[7], Jillian F. Banfield[3], Jennifer Pett-Ridge[1,8,9], Steven J. Blazewicz[1]

Physical and Life Science Directorate, Lawrence Livermore National Laboratory, Livermore, CA, USA.

Environmental Molecular Science Laboratory (EMSL), Earth and Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA

Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA, USA

Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

Current location: Department of Agroecology, Aarhus University, Aarhus, Denmark

Current location: Pacific Biosciences Research Center, University of Hawaii, Manoa, HI, USA

Center for Ecosystem Science and Society (ECOSS) and Department of Biological Sciences, Northern Arizona, University, Flagstaff, AZ, USA

Life & Environmental Science Department, University of California Merced, Merced, CA, USA

Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA, USA

Corresponding authors: Linnea K. Hernandez (honeker1@llnl.gov) and Steve Blazewicz

(blazewicz1@llnl.gov)

ABSTRACT

Changes in global patterns can leave a lasting legacy in semi-arid grasslands by reshaping

microbial growth dynamics and carbon cycling during the first wet-up in the autumn—a period

known for intense microbial activity and significant carbon emissions. To study the lasting impacts

of decreased winter rain, we implemented two precipitation regimes (100% vs. 50% mean annual

precipitation) in California Mediterranean-climate grassland field plots. After the dry season, soils

were rewetted in the laboratory with $H_2^{18}O$, and sampled at 0 h, 3 h, 24 h, 48 h, 72 h, and 168 h

post rewet. We quantified CO_2 efflux; measured microbial growth and mortality via quantitative

stable isotope (^{18}O) probing and 16S rRNA gene amplicon sequencing; and characterized the soil

organic carbon chemical composition, metagenomes, and metatranscriptomes. We found that

reduced winter precipitation imposed a strong legacy effect on microbial turnover; despite

maintaining similar respiration rates, microbial growth declined by ~1 order of magnitude,

yielding decreased community growth efficiency (CGE = gross community growth/net

respiration), and microbial mortality declined by ~ 2 orders of magnitude. Soil organic carbon also

shifted from lipid-like, amino-sugar-like, and protein-like compounds (indicative of microbial

necromass) to more oxidized lignin-like and tannin-like compounds (indicative of decomposing

plant-derived compounds). Meta-omics revealed distinct metabolic strategies linked to CGE. At

high-CGE, microbes appeared to consume more energetically favorable N-rich necromass

(released via high microbial turnover). This supported increased amino acid and peptidoglycan

biosynthesis and enhanced aromatic compound degradation, generating acetyl-CoA for the TCA

cycle that sustains higher energy yield and growth efficiency. At low-CGE, communities had

elevated carbohydrate metabolism and lipid turnover, consistent with increased investment in plant

detritus degradation and membrane repair and maintenance rather than growth. Together, our

findings demonstrate that reduced winter rainfall decreases microbial turnover. The lowered soil

organic carbon in combination with consistent C loss as CO_2 could eventually lead to a net C loss

from the soil.

1.   Introduction

Climate change is projected to alter global precipitation patterns, leading to more frequent

droughts and more intense rewetting events [1]. These changes have profound implications for soil

microbial communities, which are central to belowground biogeochemical cycling [2, 3].

Microbial diversity and activity are highly sensitive to moisture conditions [4, 5], and deviations

from typical moisture regimes are likely to influence carbon cycling and soil organic carbon

(SOC) storage [6].

Microbes control soil carbon fate through partitioning of carbon between biosynthesis of biomass

for growth and secondary metabolites, and production of CO_2 that is associated with energy

generating pathways. Microbial growth and biomass accumulation in the soil, followed by

microbial death, lead to the formation of microbial necromass, which can account for up to 50% of

the total SOC [3, 7]. This highlights the importance of microbial turnover in SOC accumulation

and persistence. One quantitative measure of microbial C allocation to growth vs. respiration is

carbon use efficiency (CUE; C-biomass $\div$ [C-biomass + C-CO_2]). Higher CUE is strongly

positively correlated with SOC accumulation [8], indicating that higher C allocation to biomass,

rather than to CO_2, enhances soil C sequestration.

Soil moisture is a primary regulator of microbial activity, growth, and respiration because water is

essential for chemical reactions, nutrient exchange, and cellular integrity. Under drought

conditions, microbial activity often shifts to adapt with changing microbial community

composition [9, 10] and gene expression [6, 11], typically resulting in decreased growth and

respiration [12] and ecosystem-dependent effects on CUE. For example, in grassland ecosystems,

drought can decrease both microbial growth and respiration without altering CUE [13, 14]. In

contrast, studies in forest, agricultural, and grassland ecosystems have reported increased CUE

under drought, which is attributed to reduced respiration, while growth rates remain stable [14,

15]. These findings indicate that the effects of drought on microbial activity and CUE may be

ecosystem dependent and can differ across studies.

In addition to drought, fluctuations in soil moisture driven by seasonal cycles further complicate

the dynamics of microbial growth and respiration. Seasonal changes in soil moisture, with

alternating periods of dry and wet conditions, exert strong controls on microbial processes. For

example, in grassland ecosystems, drier summer conditions often lead to reduced microbial growth

compared with wetter winter conditions [4, 15]. In addition to gradual changes in soil moisture

across seasons, abrupt shifts, such as rewetting after a dry period, can have dramatic effects on

microbial activity and carbon dynamics. Rewetting events often trigger a burst of microbial

respiration, frequently referred to as the "Birch effect" [16], which can result in substantial CO_2

efflux from soils, accounting for up to 10% of annual net ecosystem production [17, 18]. In a

California grassland, rewetting after the summer dry season caused a simultaneous burst in

microbial growth and mortality, indicating high microbial turnover during these events [19, 20].

These findings highlight the potential for rewetting to significantly influence both C release to the

atmosphere and the generation of microbial necromass for SOC sequestration in the soil.

Given the complexity of these ecosystem-specific responses, it is critical to employ measurement

approaches that capture the full scope of microbial carbon processing under natural soil conditions.

To address this, our study adopts the term community growth efficiency (CGE) to describe the

ratio of newly synthesized microbial biomass to total CO_2 released. Here, new microbial growth is

determined by $H_2{}^{18}O$ DNA quantitative stable isotope probing (qSIP), and the total CO released

represents carbon mineralized for energy to support growth, cell maintenance, and carbon

overflow [21]. Unlike traditional CUE measurements, which typically rely on single substrate

incubations (i.e. ^{13}C-glucose) and aim to quantify the absolute efficiency of carbon conversion

[22], CGE integrates carbon uptake from all available sources in situ and reflects the community-

level allocation of carbon under natural soil conditions. While the absolute values of CGE are not

directly comparable to those reported for CUE in the literature, the directional trends in CGE

provide meaningful insights into how microbial communities respond to environmental changes,

such as drought and rewetting, in terms of carbon allocation. By comparing CGE trends with

established CUE patterns, we aim to contextualize our findings within the broader framework of

microbial carbon cycling in soils, while maintaining transparency about the methodological

distinctions inherent to our approach.

With CGE as an integrative community-scale metric, we next consider how seasonal drying-

rewetting cycles and their legacies shift carbon allocation. While the immediate effects of soil

moisture on microbial growth and respiration have been well studied, the legacy effects of

moisture regimes—how past moisture conditions influence current microbial activity—remain less

understood. The concept of "ecological memory" describes how past events influence current

ecosystem responses; this concept is prevalent in research on aboveground ecosystems such as

plants [23] and is increasingly applied to soil microbiomes [24–27]. Canarini et al. [13] reported

that the microbial community composition differed between soils subjected to the 1st drought and

those subjected to the 10th drought, leading to altered predicted functional capacity and enzyme

production. However, other studies, such as Liu et al. (2022), reported no differences in microbial

composition or respiration between legacy drought and control soils after one month, indicating

variability in legacy effects [28]. More studies are needed to better characterize how drought

severity and ecosystem type influence these legacy responses. Understanding these legacy effects

is critical for predicting how changing precipitation patterns influence microbial ecophysiology

and carbon allocation, as well as soil organic carbon storage. Additionally, integrating microbial

growth dynamics and CGE with analyses of SOC composition and gene expression via

metatranscriptomics could provide deeper insights into how moisture regimes shape microbial

activity and carbon allocation.

In this study, we investigated the legacy effects of normal and reduced precipitation regimes

during the winter growing season on microbial growth dynamics following rewetting after the

summer dry period in a California annual grassland with a Mediterranean-type climate.

Specifically, we examined how these past precipitation regimes influenced (1) taxon-specific

microbial growth and mortality via 16S rRNA gene quantitative stable isotope probing (qSIP), (2)

CGE (microbial net growth divided by total CO_2 efflux), (3) SOC composition, and (4) microbial

gene expression via metatranscriptomics. We hypothesized that reduced precipitation during the

winter growing season would have a legacy effect, decreasing microbial growth and mortality

during rewetting six months later. Additionally, we hypothesized that these changes would

influence SOC profiles and microbial activity; soils exposed to normal precipitation regimes

would have higher CGE and exhibit increased expression of genes associated with biosynthesis,

whereas soils exposed to reduced precipitation regimes would have lower CGE and increased

expression of genes associated with energy acquisition. By improving our understanding of how

reduced rainfall impacts microbial growth, carbon allocation, and SOC composition dynamics, this

research provides critical insights into how climate change may affect SOC persistence. These

findings can inform landscape management strategies aimed at increasing SOC storage in soils.

2. Methods

2.1 Field site description

This field site is at the Hopland Research and Extension Center (HREC) located in northern California (39° 00' 14.6" N, 123° 05' 09.1" W) on territory originally occupied by the indigenous Pomo Nation. This site experiences a Mediterranean climate with cool, wet winters and warm, dry summers where most of annual precipitation occurs between October and March and most plant growth occurs from late February to early April [29]. Experimental plots were located on soils of the Squawrock–Witherell complex, have a pH of 5.9, total organic C content of 15.1 mg g^{-1}, and total N content of 1.4 mg g^{-1} [30].

In October 2017, sixteen 3.24 m^2 plots dominated by *Avena barbata* (wild oat grass) were randomly divided into two treatments that received either full (100% mean annual precipitation [MAP]; 930 mm) or reduced (50% MAP; 465 mm) precipitation. Precipitation was controlled controlled with rainout shelters constructed over each plot and a plaster pond liner placed belowground to 1 m depth to isolate each plot and reduce soil moisture flow between plots. The final rainfall occurred in May 2018, followed by the summer dry-down. In August 2018, at the end of the summer dry down when the gravimetric soil moisture was 3%, topsoil samples (0–15 cm, approximately 4700 cm^3) were collected from eight plots (n = 4 each for 50 and 100% MAP; total of 48 samples) with cleaned shovels (rinsed with Eliminase [DeconLabs, King of Prussia, PA] and water between samples), immediately placed into jumbo-sized Ziplock bags, and transferred to Lawrence Livermore National Laboratory where roots were removed and soils were homogenized by sieving (2 mm) to remove rocks and large plant debris in a greenhouse with a dry and hot atmosphere.

2.2 Laboratory ^{18}O-H$_2$O wet-up and sample collection

The samples in the laboratory were subjected to $H_2^{18}O$ labeling for quantitative growth analysis

(for details, see [30–33]). Briefly, 5 g of sieved soil from each of the eight plots was transferred

each to 11 Nalgene flat bottom vials (15 mL), for a total of 88 microcosms. One mL of isotopically

enriched water (98-atom $H_2^{18}O$) or natural abundance water (control) was slowly and evenly

pipetted onto the soil and gently mixed with the pipette tip, resulting in a final average gravimetric

soil moisture of 22%. Two vials each were then immediately sealed inside a 500 mL mason jar

with lids fitted with septa (to facilitate headspace sampling) and incubated at room temperature in

the dark. Parallel jars (total of four vials) were destructively harvested at multiple timepoints

following rewetting (0 h [before labeling/rewetting], 3 h, 24 h, 48 h, 72 h, and 168 h). For each

harvest, vials containing soil were flash frozen in liquid nitrogen and stored at -80 °C.

Headspace gas samples (5 mL) were collected at each soil harvesting time point with a gas-tight

syringe and transferred to 20 mL Wheaton serum bottles (DWK Life Sciences, Wertheim,

Germany) that had been purged and filled with N_2 (1 atm). Total headspace CO_2 was quantified

via gas chromatography equipped with a methanizer paired with a flame ionization detector

(GC2015, Shimadzu).

2.3   16S-qSIP and metagenomics

2.3.1   DNA extraction and fractionation

DNA was extracted from the soil samples via a modified phenol–chloroform protocol adapted

from [34]. Each sample was extracted in triplicate, and the final DNA extracts were combined. For

each extraction, soil (0.4 g) was added to a 2 mL Lysing Matrix E tube (MP Biomedicals, Irvine,

CA) and extracted twice as follows: each tube received 500 µL extraction buffer (5%

cetyltrimethylammonium bromide [CTAB], 0.5 M NaCl, 240 mM K_2HPO_4, pH 8.0) and 500 µL of

25:24:1 phenol:chloroform:isoamyl alcohol, shaken (30 s, 5.5 ms^{-1}) (FastPrep24, MP Biomedicals,

Santa Clara, CA), and then centrifuged for 5 min (16,100 × g). To remove residual phenol, the

supernatant was added to a pre-spun 2 mL phase lock gel tube (5 prime, Gaithersburg, MD) with

an equal volume of 24:1 chloroform:isoamyl alcohol, mixed by inverting, and centrifuged for 2

min (16,100 × g). The aqueous phases from both extractions were pooled, mixed with 7 µL of

RNAse (10 mg/mL) by inverting, incubated for 1 h (4 °C), and centrifuged for 15 min (16,100 ×

g). The supernatant was added to a new 1.7 mL microcentrifuge tube, and 1 µL of glycoblue (15

mg/mL) and 1 mL of 40% polyethylene glycol (PEG) 6000 in 1.6 M NaCl were added, mixed by

vortexing, and incubated at room temperature in the dark for 2 h. After centrifugation for 20 min

(16,100 × g), the pellet was rinsed with 1 mL of ice-cold 70% ethanol, air-dried, resuspended in 30

µL of 1x Tris-EDTA (TE), and stored at -80 °C.

To measure the degree of ^{18}O incorporation in the DNA, each sample was separated in a cesium

chloride density gradient via ultracentrifugation, as previously described [19]. For each sample, 5

µg of DNA in 150 µL of 1xTE was mixed with 1 mL gradient buffer (0.1 mol L^{-1} Tris, 0.1 mol L^{-1}

KCl, and 1 mmol L^{-1} EDTA) and 4.6 mL cesium chloride (CsCl) stock (1.885 g mL^{-1}), with a final

average density of 1.73 g mL^{-1}. The samples (~5.2 mL) were loaded into an ultracentrifuge tube

(13 mm × 51 mm Quick-Seal, Beckman Coulter Brea, CA) and spun at 20 °C for 108 h at 176,284

RCF$_{avg}$ in an Optima XE-90 ultracentrifuge (Beckman Coulter, Brea, CA) using a VTi65.2 rotor.

Immediately following centrifugation, the contents of each tube were separated into 36 fractions

(~200 µL each) via a high-throughput, automated SIP pipeline (HT-SIP) [35]. Each tube was

mounted in a fraction recovery system (Beckman Coulter, Brea, CA) where a 1260 isocratic pump

(Agilent Technologies, Santa Clara, CA) delivered water at 0.25 mL min^{-1} through a 25G needle

inserted through the top of the ultracentrifuge tube, and a side port needle routed to a 1260 Infinity

(Agilent Technologies, Santa Clara, CA) fraction collector was inserted into the bottom of the tube

to collect fractions in 96-well deep well plates. The density of each fraction was measured using an

AR200 digital refractometer (Reichart, Depew, NY) fitted with a prism covering to facilitate

measurement from 5 µL, as previously described [36]. The DNA in each fraction was purified and

concentrated using a Hamilton Microlab Star liquid handling system (Hamilton Technology, Reno,

NV) programmed to automate glycogen/PEG precipitation [35]. The washed DNA pellets were

suspended in 40 µL of 1xTE, and the DNA concentration of each fraction was quantified via a

PicoGreen fluorescence assay (Invitrogen, Waltham, MA). The fractions for each sample were

pooled into five groups on the basis of density (1.6400-1.7039 g mL^{-1}, 1.7040-1.7169 g mL^{-1},

1.7170-1.7299 g mL^{-1}, 1.7300-1.7449 g mL^{-1}, and 1.7450-1.7800 g mL^{-1}), and the fractions within

a pooled group were combined and sequenced.

2.3.2   Quantitative PCR

Bacterial 16S rRNA gene copy abundances in the SIP fractions were quantified by qPCR in

triplicate using the primers EUB 338F/EUB 518R [37] and averaged. Each 10 µL reaction

contained 1X Forget-Me-Not EvaGreen qPCR Master Mix (Biotium, Fremont, CA), 1.5 mM

MgCl$_2$, 0.2 µM of each primer, and 1 µL of template. Reactions were performed on a CFX384

thermal cycler (Bio-Rad, Hercules, CA) under the following cycling procedure: 95 °C for 2 min

followed by 35 cycles of 95 °C for 10 s, 60 °C for 10 s, and 72 °C for 10 s. Standard curves were

generated via 10-fold serial dilutions of genomic *Escherichia coli* DNA (ATTC, MG1655).

2.3.3.   16S sequencing and analysis

For 16S rRNA gene amplicon sequencing, DNA from pooled density fractions from samples

labeled with H$_2$^{18}O and the control (5 per sample-label combination for a total of 330) was

amplified in triplicate in 10-µL reactions using primers 515f and 806r [38]. Each reaction

contained 1 µL of sample and 9 µL of Phusion Hot Start II High Fidelity master mix (Thermo

Fisher Scientific, Waltham, MA) containing 1.5 mM MgCl2. The PCR conditions were as follows:

95 °C for 2 min followed by 20 cycles of 95 °C for 30 s, 64.5 °C for 30 s, and 72 °C for 15 s. The

triplicate PCR products were then pooled, diluted 10x, and used as a template in a subsequent dual

indexing reaction that used the same primers as above, which also included the Illumina flow cell

adaptor sequences and 8-nucleotide Golay barcodes (15 cycles identical to the initial amplification

conditions). The PCR conditions were identical to those described above but included 15 cycles.

Amplicons were purified with AMPure XP magnetic beads (Beckman Coulter, Brea, CA) and

quantified with a PicoGreen assay (Invitrogen, Waltham, MA) on a BioTek Synergy HT plate

reader (BioTek Technologies, Vinooski, VT). The samples were pooled at equimolar

concentrations, purified with AMPure XP beads (Beckman Coulter, Brea, CA), and quantified

using the KAPA Sybr Fast qPCR kit (Kapa Biosciences, Willmington, MA). Libraries were

sequenced on an Illumina MiSeq (Illumina, San Diego, CA) at Northern Arizona University's

Genetics Core Facility using a 300-cycle v2 reagent kit.

Demultiplexed reads were right trimmed to 140 nucleotides and quality filtered with DADA2

v1.14.1 (parameters maxN = 0, maxEE = c(2, 2), truncQ = 2) [39]. Chimeric sequences were

predicted *de novo* and removed with the removeBimeraDenovo() function in DADA2 using the

"consensus" method. The resulting amplicon sequence variant (ASV) table was filtered to keep

ASVs with lengths 252-254. Taxonomy was assigned with the RDP classifier v2.11 [40] against

training set 16.

2.3.4.   Taxon-specific growth and mortality rate calculations

Atom fraction excess (AFE) and growth, mortality, and net rates were calculated using quantitative

isotope probing (qSIP) in R (v.4.4.0) [41], as previously described [19]. Specifically, we estimated

the number of 16S rRNA gene copies per taxon in each pooled density fraction by multiplying the

sequencing-derived taxon-specific relative abundances by the total number of 16S gene copies as

measured by qPCR. To maximize the available information on the densities of taxa in the natural

abundance isotope treatment, we combined taxon-specific density estimates across the 3 h, 24 h,

48 h, 72 h, and 168 h time points for a total of 15 replicates. We limited our analyses to only those

taxa that occurred in at least 3 (of 15) replicates of the natural abundance treatment and in all 3

replicates of the ^{18}O treatment to reduce the likelihood of spurious density shifts for rare taxa.

Density shifts and atom fraction excess estimates, along with 95% confidence intervals, were

computed by bootstrapping (1000 iterations; [42]).

We estimated absolute population growth and mortality rates (# [g-soil]$^{-1}$ d^{-1}) as the changes in

16S gene copies per gram soil per time, over each of the time intervals of the incubation: 0-3 h, 0-

24 h, 0-48 h, 0-72 h, and 0-168 h. We assumed linear population growth over the course of the

incubations, and we estimated taxon-specific net absolute population growth rates and taxon-

specific absolute mortality rates for the five time intervals using the equations previously

described [19]. The fraction of oxygen in DNA derived from water (versus other organic sources)

was assumed to be 0.60 [43]. Uncertainty of growth and mortality rates (95% confidence interval)

was estimated using a bootstrapping procedure with 1000 iterations [43]. As highlighted in

previous work [43], calculated absolute abundances should be interpreted as approximations,

given the various challenges and biases inherent to the entire work flow. These include incomplete

extraction of DNA from soil microorganisms [44], differences in 16S rRNA gene copy numbers

among taxa, and biases introduced during amplification and sequencing [45, 46]. Consequently,

we emphasize that our calculations are best viewed as rough approximations for assessing taxon-

specific growth and mortality rates.

2.3.5.   Bulk 16S rRNA gene amplicon analysis

We used the sequence data per fraction to approximate the bulk microbial communities at each

timepoint. To this end, we first determined how much of the total DNA was represented in each

fraction by dividing the 16S RNA gene copy number from each fraction by the total across all

fractions. Next, we multiplied the total counts across all fractions by the proportion of DNA

represented in each fraction to obtain an adjusted new total count. Finally, the relative abundance

of each ASV was multiplied by the total count for each fraction to obtain an adjusted new count

for each ASV. This allowed us to sum the counts of each ASV across all fractions to obtain a

single value for each sample to obtain a count table. Downstream taxonomic analysis and bar

chart creation were performed via the phyloseq R package (v. 1.48.0) [47]. Taxa with relative

abundances less than 0.01% were filtered out.

2.3.6.   Proxy or Metric for Community growth efficiency (CGE)

To provide a quantitative assessment of carbon allocated to new growth relative to CO_2

production, the community growth efficiency (CGE) for each timepoint (t) was calculated from

net growth (*Gn*) and respiration (R; CO_2 efflux) as follows, where CGE_{max} was the highest CGE

across all samples:

$G_n = G_r \times t$

$CGE_{raw} = G_n/R$

$CGE = CGE_{raw}/CGE_{raw\text{-}max}$ (so that $0 \leq CGE \leq 1$)

2.3.4.   Metagenomic sequencing and analysis

Metagenomic analysis was performed on 9 samples from the 50% MAP treatment (3 plots at 0 h,

48 h, and 168 h; no samples from 24 h or 72 h were sequenced) and 15 samples from the 100%

MAP treatment (3 plots at 0 h, 24 h, 48 h, 72 h, and 168 h). For each sample, unfractionated DNA

and DNA from 5 density fractions per sample (total sample-fractions = 234) were sequenced

following a 2 × 151 indexed run recipe on an Illumina NovaSeq S4 platform (Illumina, San

Diego, CA) by Novogene Co. (novogene.com). For metagenomics analysis, Illumina adapters and

phiX sequences were removed from raw reads using BBmap (v. 38.79), which is included in the

BBtools package [48], and the remaining reads were quality trimmed using Sickle (v. 1.33) [49]

for paired-end reads. This process yielded $1.55 \times 10^8 \pm 1.5 \times 10^7$ (mean $\pm$ standard deviation) reads

per sample. For unfractionated samples, reads from triplicate samples (plots) from each time point

were coassembled, and for fractionated samples, reads from all fractions within a sample were

coassembled using MEGAHIT (v. 1.2.9) [50] with presets of "large-meta" and a minimum contig

length of 1000 bp. DRAM (v. 1.5.0) [51] was used for gene prediction and annotation using

KEGG orthology (KO) using Hidden Markov Models (HMM) for all KO entries within the

KOfam database [52], and with the carbohydrate active enzyme (CAZy) entries within dbcan [53].

A reference gene catalog was created by clustering the amino acid sequences of all the predicted

genes across all the unfractionated and fractionated coassemblies at the 95% confidence level

using Mmseq2 (v. 13.45111) [54] to create a dereplicated list of genes to map metatranscriptome

reads.

2.4.   RNA extraction and metatranscriptome sequencing and analysis

Metatranscriptomics was performed on all 48 samples (2 legacy precipitation treatments $\times$ 3 plots

$\times$ 6 timepoints), and a full description of RNA extraction and sequencing is provided in [32, 55].

Briefly, RNA was extracted from soils via the RNeasy PowerSoil Total RNA Kit (Qiagen, Hilden,

Germany) according to the manufacturer's instructions. Extracted RNA was treated with RNase-

free DNase (Qiagen, Hilden, Germany) and stored at -80 °C. The RNA concentration was

measured with a Qubit fluorometer (Invitrogen, Waltham, MA), and the quality was assessed with

a Nanodrop One Spectrophotometer (Thermo Fisher Scientific, Waltham, MA). The samples were

sent to the Joint Genome Institute (JGI; Berkeley, California) for rRNA depletion, library

preparation, and sequencing. Ribosomal (r) RNA was removed from 100 ng of total RNA via

Qiagen FastSelect 5S/16S/23S for bacterial rRNA depletion (and additional FastSelect plant

and/or yeast rRNA depletion) (Qiagen, Hilden, Germany) with RNA blocking oligo technology.

One heavily degraded sample was discarded (P04_x50_t0; t0 sample from Plot 4 of 50% legacy

moisture treatment), and the remaining 47 samples were reverse transcribed to create first-strand

complementary (c) DNA using the Illumina TruSeq Stranded mRNA Library prep kit (Illumina,

San Diego, CA), followed by second-strand cDNA synthesis, which incorporated dUTP to quench

the second strand during amplification. The double-stranded cDNA fragments were then A-tailed

and ligated to JGI dual-indexed Y-adapters, followed by enrichment of the library by 10 cycles of

PCR. The prepared libraries were quantified via a KAPA Biosystems next-generation sequencing

library qPCR kit (Kapa Biosciences, Willmington, MA) and run on a Roche LightCycler 480 real-

time PCR instrument. Sequencing of the flow cell was performed on the Illumina NovaSeq

sequencer (Illumina, San Diego, CA) using NovaSeq XP V1.5 reagent kits and an S4 flow cell,

following a 2×151 indexed run recipe.

The raw reads were downloaded from the JGI genome portal and quality control filtered via bbduk

(from the BBtools [v. 38.79] package [48]) and Sickle (v. 1.33) [49]. rRNA was then filtered from

the paired-end reads via SortMeRNA (v. 4.3.6) [56], followed by repair.sh (BBtools [v. 39.06]

package) to correct the read orientation. After all the filtering steps, $31.6 \times 10^6 \pm 1.5 \times 10^6$ reads per

sample remained. Reads were mapped via BBmap (BBtools [v. 39.06]) to the metagenome

reference gene catalog, and counts and transcripts per million (tpm) were calculated via coverM

(v. 0.70) [57]. The mapping rate was $23.2 \pm 8.4\%$ (except for P08_x100_t3 which had no reads

mapped and was removed from downstream analyses). The taxonomies of the reads were

determined via Kraken 2 (v. 2.1.3) [58]. Normalization of counts was performed to obtain

variance stabilized transformations (VSTs) using DESeq2 (v. 1.44.0) [59].

2.5   Fourier transform ion cyclotron resonance mass spectrometry (FTICR-MS)

For each sample, 1 g of lyophilized soil was defrosted, added to a clean tube, and extracted with 2

mL of Milli-Q water (>18.2 MΩ·cm resistivity). The samples were shaken at 1000 rpm at room

temperature for 2 hours on a vortex shaker and then centrifuged at 6000 rpm for 5 minutes, after

which the supernatant was removed. This process was repeated with another 2 mL of Milli-Q

water, and the supernatants were combined. Solid phase extraction (SPE) was performed on water

extracts using Bond Elut Priority Pollutant (PPL) cartridges (Agilent Technologies, Santa Clara,

CA) to remove salts and impurities that could interfere with mass spectrometry (MS) analysis

[60]. Water extracts were diluted to 5 mL with Milli-Q water and adjusted to pH 2 with ~2 µL of

85% H_3PO_4 prior to addition to methanol-activated PPL cartridges. The organic matter bound to

the cartridges was rinsed with 50 mL of 10 mM HCl, dried with nitrogen and eluted in 1.5 mL of

methanol. Following the water extraction, a modified metabolite, protein, lipid extraction

(MPLEx) [61, 62] was performed to yield polar and nonpolar liquid fractions. Ice-cold chloroform

and methanol (2:1, total ×6 mL) were added to the soil, which was vortexed after each addition.

Water was added (0.25 mL), and the samples were shaken for 1 hr at 1000 RPM. Another 1.25

mL of water was added at a final solvent ratio of 8:4:3 (chloroform:methanol:water), and the

samples were gently shaken before being incubated at 4 °C overnight. The samples were then

centrifuged at 6000 RPM for 5 minutes, yielding two layers. The layers were removed separately

and frozen at -80 °C until individual analysis.

Fourier transform ion cyclotron resonance (FTICR)-MS data were acquired using a 7 T Bruker

ScimaX FTICR-MS (Bruker, Billerica, MA) operated with quadripolar (2×) detection and located

in the Environmental Molecular Science Laboratory at Pacific Northwest National Laboratory

(Richland, WA, USA). External calibration was performed with sodium trifluoracetic acid

followed by shimming of the magnet to minimize 1× harmonic resonance peaks and tuning to

optimize the spectra for peak intensity, shape and resolution over the m/z range of 200–1000. The

extracts were directly infused into the electrospray source in negative ion mode at a voltage of +4

kV, a temperature of 200 °C, and a dry gas flow of 4 L/min in randomized order via a custom

automated cart. The ion accumulation time was set to 10 ms, and 2.1 s transients were added over

300 acquisitions in an 8 MW time domain for an estimated average resolution of ~680 K at m/z

400. Internal recalibration of each spectrum, post-acquisition, was performed with calibration lists

of standard organic matter (OM) components. Peak picking was performed during data analysis

using a signal-to-noise ratio (S/N) threshold of 7, a relative intensity threshold of 0.01, and an

absolute intensity threshold of 100,000.

The peak lists were exported, and Formularity software [63] was used to align the spectra within a

0.5 ppm threshold and assign formulas of $N \leq 2$, $S = 0$ and $P = 0$, and <0.5 ppm error for high

confidence assignments (average error <0.2 ppm). The peak lists were then corrected by

subtracting the peaks detected in the extraction blanks. Elemental ratios were calculated from the

molecular formulas assigned for each peak and averaged for each sample.

Analysis of the FTICR data was performed using MetaboDirect (v.0.2.7) [64] to 1) create profiles

of biochemical compound classes for each of the thousands of peaks in each sample's electrospray

ionization (ESI) FTICR-MS spectrum using van Krevelen diagrams and 2) calculate

thermodynamic parameters (nominal oxidation state of carbon [NOSC] and Gibb's free energy

[GFE]). Each peak was assigned a biochemical compound class on the basis of its predicted

formula and plotting of C, H, and O on a van Krevelen diagram as follows: s ($0<C:C\leq0.3$ and

$1.5 \leq H:C \leq 2.5$), unsaturated hydrocarbons ($0 \leq O:C \leq 0.125$ and $0.8 \leq H:C \leq 2.5$), proteins

($0.3 < O:C < 0.55$ and $1.5 \leq H:C \leq 2.3$), amino sugars ($0.55 < O:C \leq 0.7$ and $1.5 \leq H:C \leq 2.2$), lignin

($0.125 < O:C \leq 0.65$ and $0.8 \leq H:C < 1.5$), tannins ($0.6 < O:C \leq 1.1$ and $0.8 \leq H:C < 1.5$) and condensed

hydrocarbons (aromatics; $O \leq 200$ $O:C \leq 0.95$ and $0.2 \leq H:C < 0.8$) [65]. This method is

semiquantitative, and each compound class is then reported as relative abundance values on the

basis of counts (presence/absence). For the methanol extraction, two outliers were removed:

P10_x50_t72 and P13_x50_t71.

2.6   Statistical analysis

All the statistical tests were performed in R v. 4.4 [41].

2.6.1   Linear mixed effects models

Significant differences in the NOSC and GFE of unique compounds as well as the
growth/mortality rate, net growth, 16S rRNA gene abundance (biomass), bulk 16SrRNA phyla,
and CO_2 efflux between the legacy precipitation treatments over time post rewet were assessed by
fitting a linear mixed-effects (LME) model. LME models were created using the lmer() function in
the lmerTest package [66] with the following formula:

  LME model = *Value ~ Precipitation * Timepoint + (1|Plot)*

where *Value* is the measurement, *Precipitation* (50% vs 100%) and *Timepoint* (0 h, 3 h, 24 h, 48
h, 72 h, and 168 h) are the fixed effects, and *Plot* (location sample was collected from) is the
random intercept to account for repeated measures. To determine the significance of each fixed
effects term in the LME model, an analysis of variance (ANOVA) was performed using the
anova() function included in base R. To test for significant differences at each timepoint, we
computed estimated marginal means (least-squares means) and Tukey's honestly significant
difference (HSD)-adjusted pairwise comparisons with the emmeans() and pairs() functions in the
emmeans package (v. 1.11.1) [67].

2.6.2   G test for unique compounds

To test for the uniqueness of the FTICR compounds between the legacy precipitation treatments,
we used a G test of independence. First, we extracted all formulas that were detected in only the

50% or 100% legacy precipitation treatments, where a compound was considered detected if it

was present in at least two of the three plots and determined the counts for each compound class

(amino sugar-, carbohydrate-, condensed hydrocarbon-, lignin-, protein-, tannin-, and unsaturated

hydrocarbon-like compound). We tested for a shift in unique compound class composition

between treatments with a log-likelihood ratio (G-) test using the GTest() function in the

DescTools package (v. 0.99.60) [68]. We then performed a G test on each compound class

individually (with all other compounds in the "other" category) to determine a G value and

associated p value per class. False discovery rate (FDR)-corrected p values are reported.

2.7.3   Nonmetric multidimensional scaling

Nonmetric multidimensional scaling (NMDS) was performed to identify relationships between

samples and identify major drivers for each meta-omics dataset (FTICR compounds [WEOC and

MEOC] and transcript abundances [metatranscriptomics] at the KO functional level). Analysis

was performed on Bray☐Curtis dissimilarity matrices for transcript abundances and Jaccard

distance matrices for FTICR using the metaMDS function in the vegan package (v. 2.7-1) [69].

PERMANOVA was used to determine significant differences between treatment groups (i.e.,

timepoint relative to wet-up and legacy precipitation) using the adonis2 function in the vegan

package.

2.7.4   Partial least square regression

Partial least squares regression (PLSR) was used to identify genes whose related metabolic

pathways were most strongly associated with CGE. As predictor variables, the VST counts of all

KOs and the CAZy enzyme class were used across samples from all timepoints, except for 0 h.

The response variable was the log-transformed CGE metric. We selected the genes with variable

importance in projection (VIP) scores in the top 10% as those that contributed most to CGE. The

direction of correlation was determined by whether the coefficient was positive or negative. KO

genes were mapped onto KEGG metabolic pathways using the online tool Interactive Pathway

Explorer v3 (https://pathways.embl.de/).

2.7.5   Correlations

Pearson correlations were calculated between the log-transformed CGE metric and the transcript

abundances of CGE-associated genes (VST counts) within specific biosynthesis and degradation

pathways for key biomolecules and energy-related pathways from curated lists (**Table S1**). The

specific pathways within each of these categories are as follows: 1) the biosynthesis list included

amino acid, lipid, nucleotide, peptidoglycan, teichoic acid, and extracellular polymeric substance

[EPS] biosynthesis pathways; 2) the degradation lists included amino acid, lipid, nucleotide,

aromatic, and C-ring (subset of aromatic) degradation pathways; and 3) the energy-related lists

included central C metabolism, pyruvate oxidation (subset of central metabolism), electron

transport chain, adenosine triphosphate production (ATP-ase), fermentation, and the reactions

between phosphoenolpyruvate-oxalacetate-malate-pyruvate (POMP). Correlations were

performed between the sum of transcript abundances (VST counts) of all KOs within each

pathway and the log-transformed CGE metric. Linear regressions were performed to visualize

relationship between VST counts and CGE metric on plots and Pearson correlations to determine

degree and significance of correlation with r and p-values (FDR-corrected).

3.   3.   Results

3.1   Growth, microbial turnover and community growth efficiency (CGE)

Large differences in taxon-specific growth rates were evident between legacy precipitation

regimes (**Fig. 1a-d**). At 3 h post rewatering, the normal legacy precipitation (100% MAP)

treatment was dominated by the growth of *Proteobacteria*, *Firmicutes*, and *Actinobacteria* and the

mortality of *Proteobacteria* and *Actinobacteria*, in decreasing order of rates (**Fig. 1a, b**). In

contrast, growth under reduced legacy precipitation (50% MAP) was dominated by *Firmicutes*,

Proteobacteria, and *Actinobacteria* in decreasing order of rates, and mortality was dominated by

Actinobacteria (**Fig. 1c, d**). All taxa at the phylum level presented the highest growth rates at 3 h

across both legacy precipitation regimes, except for *Proteobacteria* at reduced precipitation,

which presented the highest growth rate at 24 h post rewatering. This finding reflects the

differences in the bulk communities representative of all active and non-active microbes, where

reduced legacy precipitation had a greater overall relative abundance of *Actinobacteria* across all

timepoints (p=3e-9; df=1; LME) and specifically at 0h before rewetting (p=4e-4, Tukey HSD;

Fig. 1e). Normal legacy precipitation had a greater overall relative abundance of *Proteobacteria*

across all timepoints (p=0.01, df = 1, LME; **Fig. 1e**) and specifically at 0 h (p=0.002; Tukey

HSD). Despite a greater proportion of *Firmicutes* growing after rewetting in the reduced legacy

precipitation treatment, the relative abundance of *Firmicutes* at the bulk level across all

timepoints, including before rewetting at 0 h, did not differ between the legacy precipitation

treatments (p>0.05, LME and Tukey HSD; **Fig. 1e**). These shifts in microbial communities at the

bulk level were also evident in beta diversity, where legacy precipitation treatment was a

significant driver of community composition (p=0.001, df = 1, PERMANOVA).

These overall patterns in growth dynamics were reflected at the community level, where growth

and mortality rates across timepoints were significantly higher for the normal than for the reduced

legacy precipitation treatments (p=4e-14 and p=8e-15, respectively, df=4, LME; **Fig. 2b, c**). The

highest growth and mortality rates under normal legacy precipitation occurred at 3 h and were

2.1×10^9 and 9.5×10^9 16S copies g-soil^{-1} d^{-1}, respectively, and the highest growth and mortality

rates under reduced legacy precipitation occurred at 24 h and 3 h and were 2.3×10^8 and 1.2×10^8

16S copies g-soil^{-1} d^{-1}, respectively.

The total biomass, measured as the total 16S rRNA gene abundance as determined by qPCR and

representing both active and inactive bacteria and archaea, was overall greater in the normal than

in the reduced legacy precipitation treatments (p=2e-11; df=4, LME; **Fig. 2a**). While the

cumulative new biomass, which was calculated from growth rates, was greater under normal

compared to reduced legacy precipitation (p=3e-9, df=4, LME; **Fig. 2d**), the CO_2 efflux was

similar in both treatments (p=0.4, df=4, LME; **Fig. 2e**). This resulted in a divergence in CGE,

which was greater in the normal compared to the reduced legacy precipitation treatments (p=8e-5,

df=4, LME; **Fig. 2f**), indicating a decline in CGE metric at reduced legacy precipitation associated

with an increase in soil CO_2 loss per unit of new growth.

3.2   Dynamics of meta-omics driven by rewatering and legacy precipitation

After observing the drastic impact of legacy precipitation on microbial communities, community

growth rates, and CGE, we sought to determine the impact of legacy precipitation on changes in

SOC, measured as water-extractable organic carbon (WEOC) and methanol-extractable organic

carbon (MEOC) from FTICR-MS, and gene transcript abundances (metatranscriptomics). For

these datasets, timepoint was the primary driver of changing SOC composition and gene

expression (WEOC [p<0.001], MEOC [p=0.01], gene expression [p<0.001; df=5;

PERMANOVA), whereas legacy precipitation was a secondary driver for only WEOC (p=0.006)

and MEOC (p=0.01, df=1, PERMANOVA) (**Fig. S1**).

3.2.1   Water-extractable organic carbon (WEOC)

The composition and thermodynamics of soil WEOC, which is indicative of dissolved organic

matter readily accessible to microbes, differed between timepoints and legacy precipitation. In

total, 119,705 and 122,285 compounds were detected in the reduced and normal legacy

precipitation treatments, respectively, and of these, 8,660 and 8,307 from the reduced and normal

legacy precipitation treatment, respectively, were assigned formulas and classified into compound

classes on the basis of their O:C and H:C ratios. Class-assigned compounds, averaged across

legacy precipitation treatments, were characterized as follows: lignin-like compounds dominated

with an average proportion of 47.9%, followed by condensed hydrocarbon-like compounds

(32.4%), tannin-like compounds (14.5%), protein-like compounds (2.0%), carbohydrate-like

compounds (1.0%), lipid-like compounds (1.0%), amino sugar-like compounds (0.8%), and

unsaturated hydrocarbon-like compounds (0.2%) (**Fig. S2c**).

A comparison of the reduced and normal legacy precipitation treatments revealed that there were

643 compounds were only present in the reduced legacy precipitation treatment ("unique" to

reduced precipitation), and 280 compounds were only present in the normal legacy precipitation

treatment ("unique" to normal precipitation). Within these unique compounds, there was a

significant shift in class-level categorization (p<2e□¹□, G=229, df=8, G test). In the reduced

legacy precipitation treatment, tannin- and lignin-like compounds, which are characteristic of

decomposing plant detritus, were overrepresented (p=4e-10, G=41, and p=6e-4, G=12,

respectively; G test), as were condensed hydrocarbons (p=4e-6, G=22.4, G test). In the normal

legacy precipitation treatment, protein-like (p=9e-11, G=45, G-test), lipid-like (p=1e-7, G=30, G-

test), carbohydrate-like (p=1e-3, G=11, G-test), and amino sugar-like (p=0, G=71, G-test)

compounds, characteristic of necromass, were overrepresented, as were unsaturated hydrocarbon-

like compounds (p=3e-5, G=18, G-test) (**Fig 3a, b**; see **Fig S3d** for the van Krevelen diagram with

all the compounds). This shift in the class composition of compounds unique to normal and

reduced legacy precipitation was associated with decreased mean Gibb's free energy (GFE) and

increased nominal oxidation state of carbon (NOSC) (p<2e-16, df=1, LME**; Fig. 3c, d**).

At the class level, there were trends across time after rewetting (**Fig. S3a**). Carbohydrate-like and

lipid-like compounds decreased in proportion in both the reduced and normal legacy precipitation

treatments over time between 0 h and 168 h (p<0.05 and p<0.001, respectively; df=5; Tukey

HSD). Amino sugar-like compounds decreased in proportion in only the normal legacy

precipitation treatment between 0 h and 168 h (p<0.05, df=5, Tukey HSD). Condensed

hydrocarbon-like compounds decreased in proportion in both the reduced and normal legacy

precipitation treatments between 0 h and 168 h (p<0.05, df=5, Tukey HSD) and unsaturated

hydrocarbons decreased in only the normal legacy precipitation treatment between 0 h and 168 h

(p<0.001, df=5, Tukey HSD). Lignins and tannins did not change significantly over time.

3.3.2   Methanol extractable organic carbon (MEOC)

The composition and thermodynamics of soil MEOC, which is indicative of more polar organic

matter that is not as readily accessible to microbes as WEOC is, also differed across timepoints

post rewetting and legacy precipitation. In total, 51,829 and 55,963 compounds were detected in

the reduced and normal legacy precipitation treatments, respectively, and of these, 5,358 and 5340

from the reduced and normal legacy precipitation treatment, respectively, were assigned formulas

and classified into compound classes. Class-assigned compounds across legacy precipitation

treatments were characterized as follows: lignin-like compounds dominated with an average

proportion of 38.9%, followed by condensed hydrocarbon-like compounds (17.1%), protein-like

compounds (15.2%), carbohydrate-like compounds (11.9%), amino sugar-like compounds (8.7%),

lipid-like compounds (5.7%), tannin-like compounds (2.1%), and unsaturated hydrocarbon-like

compounds (0.3%) (**Fig. S3e**). Compared with the WEOC, there were more protein-like,

carbohydrate-like, amino sugar-like, and lipid-like MEOCs and fewer tannin-like and condensed

hydrocarbon-like MEOC compounds. Compounds such as amino sugars and proteins are often

considered polar; however, they may contain hydrophobic regions that can sorb to soil minerals,

making them less bioavailable [70, 71].

There were 725 compounds only present in the reduced legacy precipitation treatment ("unique"

to reduced precipitation), and 293 compounds were only present in the normal legacy precipitation

treatment ("unique" to normal precipitation). Among these unique compounds, there was a

smaller shift in class-level categorization between legacy precipitation treatments than in the

WEOC, but this shift was still significant (p<2e^{-1}^{6}, G=87, df=8, G test) with carbohydrate-like

compounds being enriched in the normal legacy precipitation treatment (p=5e-4, G=16) (**Fig. S2a,**

b; see **Fig. S3f** for the van Krevelen diagram with all the compounds). Similar to WEOC, this shift

in the class composition of unique compounds from normal to reduced legacy precipitation was

associated with a lower mean GFE and higher NOSC (p=3e-7, df=1, LME**; Fig. S2c, d**).

At the class level, there were fewer trends across time post laboratory rewet than WEOC, with the

largest shift occurring between pre- (0 h) and post rewet (all other timepoints), particularly with a

decrease in amino sugars and carbohydrates at normal precipitation between 0 and 3 h (p=0.02,

df=18, Tukey HSD) (**Fig. S3b**). Furthermore, at 0 h, there was a greater abundance of amino

sugars and carbohydrates in the normal-precipitation treatment than in the reduced-precipitation

treatment (p=0.02 and 0.05, respectively; df=18; Tukey HSD) and a greater abundance of

condensed hydrocarbons in the reduced-precipitation than in the normal precipitation (p=0.04,

df=18, Tukey HSD; **Fig S3b**). Compared with those at 0 h, the abundance of condensed

hydrocarbons tended to increase with time at 48, 72, and 168 h post rewetting (p=0.02, p=0.03,

and p=0.04, respectively; df=18; Tukey HSD).

3.3.   Gene expression and community growth efficiency metric

Next, we identified genes and their metabolic pathways that were most associated with CGE

metric using PLSR. We screened all the KO-annotated genes and identified 644 that were

associated with CGE (top 10% of the VIP scores ranging from 1.6–3.5) (**Table S4**). Among these

genes, 389 were positively associated with CGE and 254 were negatively associated with CGE.

Transcripts for genes associated positively with CGE metric were involved in amino acid

metabolism (biosynthesis of threonine, methionine, lysine, and tryptophan, serine), nucleotide

metabolism (biosynthesis and degradation of purines and pyrimidines), xenobiotics metabolism

(degradation of benzoate and catechol) pathways, and energy (TCA cycle, oxidative

phosphorylation) (**Fig. 4a**), whereas transcription associated negatively with CGE metric were

associated with amino acid metabolism (biosynthesis of lysine, serine, and proline; degradation of

lysine), lipid metabolism (biosynthesis of fatty acid, glycerolipids, and glycerophospholipids),

carbohydrate metabolism (breakdown of trehalose, glycogen, and cellulose and pentose and

glucoronate metabolism), nucleotide metabolism (biosynthesis and degradation of purines and

pyrimidines), and energy (glycolysis, oxidative phosphorylation, and pentose phosphate pathway)

metabolism pathways (**Fig. 4b**).

Next, we focused on whether the expression of CGE-associated genes within specific biosynthesis,

degradation, and energy pathways from our curated lists was correlated with the CGE metric

(**Table 1; Table S2**). Within our curated lists we found 74 genes that were associated positively

with CGE and of these, 26% were involved in biosynthesis, 64% were involved in degradation,

Table 1. Summary of CGE-associated genes that belong to each pathway and sub-pathway within
the curated lists of biosynthesis, degradation, and energy genes

Category	Pathway	Sub-pathway (total number of genes)	Genes associated with CGE		Sum of genes within each pathway	
			Negative	Positive	Negative	Positive
Biosynthesis	Amino acid	Alanine (8)	0	2	5	11
		Chorismate (15)	0	1		
		Cysteine (15)	1	1		
		Lysine/methionine (5)	0	1		
		Serine (18)	2	2		
		Tryptophan (14)	0	1		
		Shared/ Other (154)	2	3		
	Lipid	Fatty acid (38)	1	0	3	0
		Glycerolipid (42)	1	0		
		Glycerophospholipid (42)	1	0		
	Nucleotide	Purine (32)	1	0	4	3
		Pyrimidine (15)	2	2		
		Ribonucleotide (7)	1	1		
	Peptidoglycan	Aminosugar (22)	0	1	1	2

		Peptidoglycan unit (22)	1	0		
		Remodeling (21)	0	1		
	EPS	Acetan/xanthan (12)	1	0	**1**	**1**
		Alginate (7)	0	1		
	Teichoic acid	from UND-PP to wall teichoic acid (42)	0	2	**0**	**2**
	Amino acid	Alanine/aspartate/asparagine/glutamate/glutamine (23)	2	1		
		Arginine/Proline (38)	3	3		
		Cysteine/methionine (45)	1	4		
		Lysine (22)	1	0	**13**	**12**
		Phenylalanine/tyrosine/tryptophan (101)	2	4		
		Serine/glycine/threonine (29)	2	0		
		Shared (52)	2	0		
	Lipid	Other	1	0	**1**	**0**
	Nucleotide	Purine	2	1	**4**	**3**
		Pyrimidine	2	2		
Degradation	Xenobiotics	Aminobenzoate (62)	1	2		
		Atrazine (16)	0	1		
		Benzoate (52)	1	8		
		Caprolactam (16)	0	1		
		Chlorocyclohexane and chlorobenzene (16)	0	1		
		Furfural (8)	1	0	**6**	**22**
		Naphthalene (18)	0	2		
		Nitrotoluene (18)	1	1		
		Shared (87)	2	4		
		Steroid (15)	0	1		
		Styrene (17)	0	1		
	Aromatic ring (AR)	Dihydroxylation of AR (6)	0	1		
		Dihydroxylation and meta-cleavage of AR (25)	1	1		
		Ortho-cleavage of dihydroxylated AR (9)	1	2	**2**	**10**
		Ortho-cleavage of halogenated AR (2)	0	1		
		Other pathway (36)	0	4		
		Ring removal from polycyclic AR (12)	0	1		
Energy	ATPase	ATP-binding (75)	2	3		
		F-type (29)	1	0	**4**	**3**
		P-type (14)	1	0		
	Central C metabolism	Glycolysis/Gluconeogenesis (46)	2	0		
		PPP (16)	1	0	**4**	**1**
		Pyruvate oxidation (11)	1	0		
		TCA cycle (49)	0	1		
	ETC	Cytochrome C oxidase (41)	3	0	**4**	**3**
		NADH reductase (80)	1	3		
	Fermentation	to acetate (10)	0	1	**0**	**1**
	Pyruvate ox.	Pyruvate => acetyl-CoA (29)	1	0	**1**	**0**
Total sum			**53**	**74**		

CGE-associated genes were determined by partial least squares regression (PLSR) analysis and selecting genes within the top 10 % of variable importance projection (VIP) scores (VIP > 1.6). Direction of association with CGE was determined as positive or negative depending on sign of coefficient. "Shared" sub-pathways signifies that gene is part of multiple sub-pathways. CGE, community growth efficiency; AR, aromatic ring; ox., oxidation; ETC, electron transport chain; EPS, extracellular polymeric substance; C, carbon

and 11% involved in energy metabolism. We found 53 genes that were associated negatively with

CGE and of these, 26% were involved in biosynthesis, 49% were involved in degradation, and

25% were involved in energy metabolism.

Within the biosynthesis category, the CGE metric was positively correlated with gene transcript

abundance of amino acid biosynthesis (16 genes; p=0.002, r=0.55, Pearson; **Fig. 5a**) and

peptidoglycan biosynthesis (3 genes; p=0.03, 0.35, Pearson; **Fig. 5d**), negatively correlated with

lipid biosynthesis (3 genes; p=0.002, r=-0.51, Pearson; **Fig. 5b**), and not correlated with nucleotide

biosynthesis (7 genes, p=0.3, r=0.17, Pearson; **Fig. 5c**), teichoic acid biosynthesis (2 genes; p=0.2,

r=0.29, Pearson; **Fig. S4a**), or EPS biosynthesis (2 genes; p=0.6, r=0.09, Pearson; **Fig. S4b**).

Overall, the CGE metric was positively correlated with the sum of gene transcript abundances

across all these biosynthesis pathways (total: 33 genes; p=0.004, r=0.50, Pearson; **Fig. 5e**).

The CGE metric was not correlated individually with different energy pathways (central C

metabolism, fermentation, electron transport chain, or ATPase; p>0.05, **Fig. S4c-f**) nor the sum of

transcript abundances across all energy metabolism pathways (total: 19 genes; p=0.6, r=-0.11,

Pearson; **Fig. S4h**). There were four genes within the central metabolism pathway associated

negatively with CGE (two glycolysis/gluconeogenesis, one pentose phosphate pathway, and one

pyruvate oxidation [pyruvate dehydrogenase E1 subunit gene; *PDH*]) compared to one gene

associated positively with CGE within the central metabolism pathway involved in the TCA cycle.

This suggests that although there were more genes involved in energy metabolism that associated

negatively with CGE, the transcript abundances of those positively associated with CGE were

greater.

The ratio of gene transcript abundance of biosynthesis to energy production was positively

correlated with CGE (**Fig. 5f**), an approximation of the definition of CGE (biosynthesis: energy).

A more accurate description would be the sum of the CGE-associated biosynthesis genes to CO_2-

producing genes. The pyruvate dehydrogenase E1 subunit gene (*PDH*; K00163) is the only CO_2-

producing energy gene associated with CGE (negatively) and within the pyruvate oxidation

pathway; the ratio of transcript abundance of CGE-associated biosynthesis genes to *PDH* is also

positively correlated with CGE (p=0.009, r=0.48, Pearson; **Fig S4i**).

Finally, within the degradation category, the CGE metric was positively correlated with aromatic

degradation (28 genes; p = 0.005, r-0.51, Pearson; **Fig. S5d**) and hydrocarbon ring degradation (a

subset of aromatic degradation genes that specifically act on rings; 12 genes; p=0.005, r=0.49,

Pearson; **Fig. S5e**), negatively correlated with lipid degradation (1 gene [K05939, *aas*], p=0.04,

r=-0.35; **Fig. S5b**), and not correlated with amino acid degradation (25 genes; p=0.4, r=0.15,

Pearson; **Fig. S5a**) or nucleotide degradation (7 genes; p=0.5, r=-0.08, Pearson; **Fig. S5b**). Overall,

the CGE metric was positively correlated with the sum of transcript abundance across all these

degradation genes (total: 59 unique genes; p = 0.04, r=0.38, Pearson; **Fig. S5f).**

We screened all CAZy-annotated genes and identified 26 that were strongly associated with CGE,

based on VIP scores in the top 10% (ranging from 1.6–3.5; **Fig. S6, Table S3**). Of these, 12 genes

were positively associated with CGE, while 14 were negatively associated with CGE. The gene

most positively associated with CGE (VIP = 3.5), was a glycosyltransferase (GT12) involved in

biosynthesis. Pfam annotation of this gene (PF13641.10; glycosyltransferase like family 2),

indicates a role in bacterial capsule formation. Three additional biosynthesis genes (GT76, GT33,

and GT22) were also positively associated with CGE and play a role in glycan biosynthesis, a

component of cell walls. While glycans are found in microbes and plants, these genes are likely

indicative of microbial biosynthesis because plant genes represented < 12.5 % of the sequences

(**Fig. S7)** and plants were not growing during the lab wet-up experiment. There was one glycan

biosynthesis gene (GT10) that was associated negatively with CGE. The CAZy gene that was

most negatively associated with CGE was PL42 (VIP = 2.7), a polysaccharide lyase (rhamnose-a-

1,4-D-glucuronate lyase) involved in pectin degradation, a structural component of plant tissues.

Most of the remaining CGE-associated genes encoded glycoside hydrolase (GH) enzymes that

degrade glycosidic bonds in different substrates to release glucose. Of the GH enzymes associated

negatively with CGE, four were specific to plant material substrates (arabinans, xylans,

glucuronoxylans), seven to substrates that are found in both plants and microbes (glucans,

trehalose, glycans, and galactans), and one that is specific to microbial substrates (chitosan). In

contrast, of the GH (and one PL) enzyme associated positively with CGE, three were specific to

plant material substrates (arabinans and pectin), two to substrates that are found in both plants and

microbes (glycans and galactans), and two that are specific to microbial substrates (mannans and

alginate). In addition to the greater number of CAZy genes that were negatively associated with

CGE, the total gene transcript abundance (normalized as VST) was also greater (7.3 ± 2.3 vs. 5.3

± 1.2 [VST-counts]; $p = 2.2e\text{-}16$, LME).

4.   Discussion

In this study, we found that reduced precipitation during the winter growing season in a

Mediterranean grassland ecosystem had lasting effects on SOC composition and microbial growth

and mortality dynamics during the subsequent rewetting and Birch effect. These legacy effects

persisted even six months later, after the summer dry period, when the soil moisture no longer

differed between the treatments. Specifically, reduced legacy precipitation resulted in: 1)

Microbial turnover (growth and mortality rates) that was up to two orders of magnitude lower,

despite unchanged respiration rates, leading to a decreased community growth efficiency (CGE),

and 2) compounds with decreased GFE and increased NOSC. We also identified microbial

functions associated with differences in CGEs between legacy precipitation treatments. A lower

CGE was linked to decreased biosynthesis, yet no changes in overall energy metabolism, leading

to a decreased ratio of transcripts involved in biosynthesis to energy metabolism. These results

suggest that reduced precipitation during the growing season—a scenario expected to become

more frequent under climate change—not only alters SOC composition but also impacts microbial

growth and metabolism during rewetting after the summer dry season. Under reduced legacy

precipitation, microbes are less efficient at incorporating carbon into biomass while maintaining

similar respiration rates, leading to similar C loss from the soil yet reduced SOC replenishment

through microbial turnover.

Reduced legacy precipitation decreases microbial turnover

Despite similar soil moisture conditions following the summer dry down, we observed that both

microbial growth rates and mortality rates were substantially lower—by one and two orders of

magnitude, respectively—during rewetting after a single growing season with 50% reduced

precipitation. This decrease in microbial growth in the California grassland soil is consistent with

previous findings, where growth rates at the drier Sedgwick site (383 mm y^{-1} MAP) were lower

than those at the wetter Hopland site (956 mm yr^{-1} MAP), corresponding to a ~60% decrease in

precipitation and shorter C residence times [4]. These results suggest that, over multiple growing

seasons with reduced rainfall, the observed decreases in microbial growth and mortality at

Hopland would likely lead to reduced soil C accumulation. Given that microbial necromass can

contribute up to 50% of SOC in grassland ecosystems [72], a consistent two-order magnitude

decrease in mortality each year would significantly reduce necromass inputs to SOC.

The observed changes in microbial growth may be linked to shifts in the composition and

physiological state of the microbial community induced by legacy precipitation conditions. Under

reduced legacy precipitation, the relative abundance of *Actinobacteria* was greater, that of

Proteobacteria was lower, and that of *Firmicutes* remained constant. However, *Firmicutes*, which

made up only ~5% of the total community before rewetting, accounted for ~20% and 40% of the

growing community at 3 h post rewetting in the normal and reduced legacy precipitation

treatments, respectively. This pattern is consistent with previous findings at this site, where

Firmicutes were also a major component of the growing community after being rewet [19]. In

contrast, *Actinobacteria* made up a greater proportion (~40%) of the total community under

reduced legacy precipitation but only ~20% of the growing community at 3 h post rewetting. Both

Firmicutes and *Actinobacteria* are recognized as drought-tolerant taxa [6, 73, 74]. However, their

survival strategies likely differ: *Actinobacteria* may persist with some active growth [73], whereas

Firmicutes typically survive by forming dormant spores [5]. We propose that rewetting triggers

the exit from dormancy in *Firmicutes*, leading to rapid growth, whereas *Actinobacteria* may

experience mortality due to rewetting stress after a prolonged period of low activity during the dry

season. Overall, our findings highlight that legacy precipitation conditions distinctly shape the

post-rewetting growth dynamics of *Firmicutes* and *Actinobacteria*, with *Firmicutes* exhibiting

enhanced recovery and *Actinobacteria* displaying reduced growth under reduced precipitation

regimes.

These changes in taxonomic turnover have implications not only for growth dynamics but also for

the biochemical nature of microbial necromass. Under normal legacy precipitation, dying taxa

were dominated by gram-negative *Proteobacteria*, whereas under reduced legacy precipitation,

gram-positive Actinobacteria dominated. This suggests a shift in necromass composition from

gram-negative to gram-positive origins, which differ in their macromolecule compositions (e.g.,

peptidoglycans, glycoproteins, lipoproteins, and phospholipids). Because cell wall components

can adsorb onto mineral surfaces [75] and the strength of this binding varies among different

macromolecules, these changes may influence both the type of compounds [76] entering the soil

carbon pool and their potential for stabilization.

Importantly, fungi likely contributed to microbial biomass turnover and respiration in this study.

While we did not measure fungal growth here, previous work in Hopland soils have shown that

fungal and bacterial/archaeal growth respond similarly to rewetting events [20]. Furthermore,

transcriptomic analyses indicated comparable ratios of bacteria + archaea to fungi across legacy

precipitation treatments and timepoints (**Fig. S7**), suggesting that fungal growth and mortality

likely followed similar patterns across legacy precipitation treatments as those of bacteria and

archaea. Future studies that directly quantify fungal growth and turnover will help refine microbial

CGE estimates and provide a more complete understanding of how precipitation variability

influences oil carbon cycling.

Influence of bioavailable substrate composition on CGE

Legacy precipitation influenced the composition of the organic carbon pool, altering the types of

WEOC substrates available to microbes. Consistent with previous studies, we found that reduced

precipitation led to a WEOC pool with fewer unique compounds typical of microbial biomass and

necromass, such as lipids, proteins, and amino sugars [75, 77, 78], and more unique compounds

derived from plant detritus, including lignin and tannins [79]. This shift to residual plant detritus

contributed to an increase in NOSC, indicating that a more oxidized SOC pool was consistent with

equal microbial mineralization but reduced new microbial biomass production in the reduced

precipitation treatments, resulting in greater C loss as CO_2.

In addition to substrate GFE, elemental stoichiometry (C:N:P) strongly influences microbial CUE

[80, 81]. The stoichiometry of these substrates and their hierarchy of energetic favorability can be

approximated as follows: microbial biomass/necromass (global average ~60:7:1) [82] > plant

detritus (grassland belowground ~200:6.7:1) [83]. Microbial biomass is most favorable because its

elemental ratios closely match those of the consuming microbes, minimizing the energetic cost of

assimilation. Under normal legacy precipitation conditions, rewetting induces rapid microbial

mortality, resulting in the release of a fresh pool of nutrient-rich biomass. This energetically

favorable substrate likely contributed to the observed increase in CGE at 3 h post rewetting. In

support of this finding, a high CGE coincided with increased expression of genes involved in

amino acid, peptidoglycan, and glycolipid biosynthesis, as well as several necromass

(bacterial/fungal cell wall)-degrading enzymes (for substrates including mannans and alginate), all

of which are indicative of active biomass degradation combined with growth. Previous work at

Hopland further supports this mechanism, suggesting that the nutrient-rich products and

biomolecules from microbial mortality generated following wet-up can fuel a substantial portion

of the microbial response to sudden increases in soil moisture [20]. Furthermore, necromass

accrual during the summer dry down could be made bioavailable to the microbes with the sudden

increase in moisture; amino sugars, indicative of necromass, were higher before rewetting in the

normal compared to the reduced legacy precipitation treatment and decreased significantly by 3 h

post rewet indicating rapid microbial consumption.

In contrast, plant detritus imposes a stoichiometric mismatch, requiring microbes to both expend

energy-producing extracellular enzymes and respire excess carbon to acquire sufficient nitrogen

[80]. In our study, we indeed found that low CGE was associated with greater diversity and

expression of genes encoding plant material-degrading enzymes (for substrates including pectin,

arabinans, glucuronoxylans, and xylans). This observation aligns with previous studies reporting a

negative correlation between enzyme production and CUE [15, 84], suggesting that increased

enzyme demand diverts carbon from biomass synthesis to energy generation. Additionally, gene

expression data suggests decreased P and N availability under reduced legacy precipitation—

conditions that are also associated with lower CUE [80, 85]. In this study, low CGE was

specifically associated with the expression of *senX3* [K07768] and *regX3* [K07776], genes

activated during phosphate limitation, and *ntrX* [K13599], a gene activated during nitrogen

limitation. Together, these findings highlight that legacy precipitation shapes both substrate

quality and microbial nutrient stress, ultimately influencing microbial CGE.

Microbial energy, degradation, and biosynthesis metabolisms associated with CGE

Consistent with our hypothesis that higher CGE would be associated with increased expression of

biosynthesis genes, CGE metric increased as transcript abundances for CGE-associated

biosynthesis genes in the metatranscriptome increased. However, contrary to our hypothesis that

higher CGE would be associated with decreased expression of energy genes, CGE metric did not

show an overall correlation with the combined transcript levels of CGE-associated energy

metabolism genes. However, as CGE metric increased, the ratio of biosynthesis:energy transcript

abundance also increased, indicating decreased transcriptional investment in energy-supplying

reactions per unit of anabolic transcription as growth yield increased. At the gene level, more

energy genes were negatively associated with CGE metric than positively (13 vs. 8; **Table 1**) and

were concentrated in central C metabolism pathways (glycolysis/gluconeogenesis, pentose-

phosphate pathway [PPP], and pyruvate oxidation). In contrast, there was one gene in the central

C metabolism pathway (TCA cycle) that was positively associated with CGE metric.

Although total biosynthetic activity was higher under normal legacy precipitation, lipid

biosynthesis was enriched at low CGE metric, especially for fatty acids, glycerolipids, and

glycerophospholipids. Flux through central carbon and PPP pathways can supply both the

precursor acetyl-CoA as well as NADPH to power fatty-acid elongation and desaturation [80].

The coordinated increase in lipid biosynthesis and lipid degradation at low CGE metric is most

consistent with a maintenance program rather than net biomass production characterized by

membrane repair and remodeling and futile lipid cycling under nutrient and legacy-moisture

stress. In this context, *fabB* (K00647), a core synthase of the fatty-acid pathway, can support

triacylglyceride formation for temporary carbon and energy storage [81] and cell-envelope

remodeling to maintain osmotic homeostasis [82]. Moreover, lipid catabolism can return P and C

to central metabolism during stress [83] and can be tightly coupled to lipid biosynthesis [84],

creating microbial turnover that depresses growth yield. This is characteristic of a futile cycle,

where opposing biochemical reactions occur simultaneously resulting in net energy loss without

productivity [86].

In contrast, peptidoglycan and amino acid biosynthesis were associated with high CGE metric,

consistent with rapid microbial biomass production. A second major characteristic of high CGE

was aromatic degradation that funnels carbon directly to the TCA cycle via succinyl-CoA. Entry

at succinyl-CoA bypasses glycolysis and pyruvate oxidation, avoiding upstream CO_2-releasing

reactions, and yielding substrate level ATP production. This reaction also yields succinate which

can continue through the TCA cycle supplying intermediates for amino acid biosynthesis,

particularly oxaloacetate which is upstream from the aspartate-family of amino acids (i.e. alanine,

lysine, threonine), the biosynthesis of which was associated with high CGE metric. Amino acids

are the building blocks of proteins, which make up 52–68% of the microbial cell composition

[87], including peptidoglycan components of the cell wall (specifically alanine and lysine) [88].

Aromatic degradation can also generate intermediates that can act as quorum sensing and

interspecies signal molecules that trigger cell-to-cell communication and instigate complex

microbial interactions [89]. Here, we found 11 genes that were positively associated with CGE

within the Quorum Sensing KEGG pathway (map 2024) [90] as opposed to just two that were

negatively associated with CGE metric. We also found genes associated with tryptophan

metabolism, another downstream pathway from benzoate degradation, that could produce indole,

a microbial signal molecule that can impact both microbial community composition and carbon

metabolism. Therefore, in addition to energy and biosynthesis precursors, aromatic degradation

could promote microbial communication and aggregate formations.

Conclusions

Climate change-driven increases in drought and rewetting events have the potential to strongly

affect microbial activity and SOC accrual in CA grassland ecosystems. Our study demonstrated

that while both normal and reduced winter precipitation conditions produced a pronounced Birch

effect—characterized by a CO_2 pulse following rewetting after the summer dry-down period—

reduced precipitation led to markedly lower microbial growth and mortality, resulting in

diminished necromass formation. Under typical precipitation conditions, high microbial mortality

following a rewet event generates an energetically favorable pool of necromass. This necromass

supports subsequent microbial growth which we saw in increased transcript abundances of

biosynthesis genes, such as amino acids and peptidoglycan. In contrast, reduced legacy

precipitation limits necromass availability and shifts the bioavailable SOC pool toward less

energetically favorable substrates, such as decomposed plant detritus. These metabolic

adjustments are associated with lower microbial community growth efficiency (**Fig. 6**) and

increased investment in stress response, such as lipid degradation coupled with biosynthesis for

membrane remodeling. Taken together, these findings highlight that legacy precipitation

conditions fundamentally alter microbial turnover, activity, and the composition of SOC. Over

time, such changes have the potential to reshape SOC stocks and the carbon balance of grassland

soils under future climate scenarios.

References

1. Sherwood S, Fu Q. A drier future? Science. 2014;343:737–9.

https://doi.org/10.1126/science.1247620.

2. Crowther TW, van den Hoogen J, Wan J, Mayes MA, Keiser AD, Mo L, et al. The global soil

community and its influence on biogeochemistry. Science. 2019;365:eaav0550.

https://doi.org/10.1126/science.aav0550.

3. Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie EL, et al. Life and

death in the soil microbiome: how ecological processes influence biogeochemistry. Nat Rev

Microbiol. 2022. https://doi.org/10.1038/s41579-022-00695-z.

4. Foley MM, Blazewicz SJ, McFarlane KJ, Greenlon A, Hayer M, Kimbrel JA, et al. Active

populations and growth of soil microorganisms are framed by mean annual precipitation in three

California annual grasslands. Soil Biol Biochem. 2023;177:108886.

https://doi.org/10.1016/j.soilbio.2022.108886.

5. Schimel JP. Life in Dry Soils: Effects of Drought on Soil Microbial Communities and

Processes. Annu Rev Ecol Evol Syst. 2018;49:409–32. https://doi.org/10.1146/annurev-ecolsys-

110617-062614.

6. Honeker LK, Pugliese G, Ingrisch J, Fudyma J, Gil-Loaiza J, Carpenter E, et al. Drought re-

routes soil microbial carbon metabolism towards emission of volatile metabolites in an artificial

tropical rainforest. Nat Microbiol. 2023;8:1480–94. https://doi.org/10.1038/s41564-023-01432-9.

7. Liang C, Kästner M, Joergensen RG. Microbial necromass on the rise: the growing focus on its

role in soil organic matter development. Soil Biol Biochem. 2020;:108000.

https://doi.org/10.1016/j.soilbio.2020.108000.

8. Tao F, Huang Y, Hungate BA, Manzoni S, Frey SD, Schmidt MWI, et al. Microbial carbon use

efficiency promotes global soil carbon storage. Nature. 2023;618:981–5.

https://doi.org/10.1038/s41586-023-06042-3.

9. Bouskill NJ, Wood TE, Baran R, Ye Z, Bowen BP, Lim H, et al. Belowground Response to

Drought in a Tropical Forest Soil. I. Changes in Microbial Functional Potential and Metabolism.

Front Microbiol. 2016;7:525. https://doi.org/10.3389/fmicb.2016.00525.

10. Bouskill NJ, Wood TE, Baran R, Hao Z, Ye Z, Bowen BP, et al. Belowground Response to

Drought in a Tropical Forest Soil. II. Change in Microbial Function Impacts Carbon Composition.

Front Microbiol. 2016;7:323. https://doi.org/10.3389/fmicb.2016.00323.

11. Malik AA, Swenson T, Weihe C, Morrison EW, Martiny JBH, Brodie EL, et al. Drought and

plant litter chemistry alter microbial gene expression and metabolite production. ISME J.

2020;14:2236–47. https://doi.org/10.1038/s41396-020-0683-6.

12. Wu Z, Dijkstra P, Koch GW, Peñuelas J, Hungate BA. Responses of terrestrial ecosystems to

temperature and precipitation change: a meta☐analysis of experimental manipulation. Glob Chang

Biol. 2011;17:927–42. https://doi.org/10.1111/j.1365-2486.2010.02302.x.

13. Canarini A, Wanek W, Watzka M, Sandén T, Spiegel H, Šantrůček J, et al. Quantifying

microbial growth and carbon use efficiency in dry soil environments via 18 O water vapor

equilibration. Glob Chang Biol. 2020;26:5333–41. https://doi.org/10.1111/gcb.15168.

14. Simon E, Canarini A, Martin V, Séneca J, Böckle T, Reinthaler D, et al. Microbial growth and

carbon use efficiency show seasonal responses in a multifactorial climate change experiment.

Commun Biol. 2020;3:584. https://doi.org/10.1038/s42003-020-01317-1.

15. Ullah MR, Carrillo Y, Dijkstra FA. Drought-induced and seasonal variation in carbon use

efficiency is associated with fungi:bacteria ratio and enzyme production in a grassland ecosystem.

Soil Biol Biochem. 2021;155:108159. https://doi.org/10.1016/j.soilbio.2021.108159.

16. Birch HF. The effect of soil drying on humus decomposition and nitrogen availability. Plant

Soil. 1958;10:9–31. https://doi.org/10.1007/bf01343734.

17. Lee X, Wu H-J, Sigler J, Oishi C, Siccama T. Rapid and transient response of soil respiration

to rain. Glob Chang Biol. 2004;10:1017–26. https://doi.org/10.1111/j.1529-8817.2003.00787.x.

18. Nguyen NB, Migliavacca M, Bassiouni M, Baldocchi DD, Gherardi LA, Green JK, et al.

Widespread underestimation of rain-induced soil carbon emissions from global drylands. Nat

Geosci. 2025. https://doi.org/10.1038/s41561-025-01754-9.

19. Blazewicz SJ, Hungate BA, Koch BJ, Nuccio EE, Morrissey E, Brodie EL, et al. Taxon-

specific microbial growth and mortality patterns reveal distinct temporal population responses to

rewetting in a California grassland soil. ISME J. 2020;14:1520–32.

https://doi.org/10.1038/s41396-020-0617-3.

20. Blazewicz SJ, Schwartz E, Firestone MK. Growth and death of bacteria and fungi underlie

rainfall-induced carbon dioxide pulses from seasonally dried soil. Ecology. 2014;95:1162–72.

https://doi.org/10.1890/13-1031.1.

21. Geyer KM, Kyker-Snowman E, Grandy AS, Frey SD. Microbial carbon use efficiency:

accounting for population, community, and ecosystem-scale controls over the fate of metabolized

organic matter. Biogeochemistry. 2016;127:173–88. https://doi.org/10.1007/s10533-016-0191-y.

22. He X, Abs E, Allison SD, Tao F, Huang Y, Manzoni S, et al. Emerging multiscale insights on

microbial carbon use efficiency in the land carbon cycle. Nat Commun. 2024;15:8010.

https://doi.org/10.1038/s41467-024-52160-5.

23. Peterson GD. Contagious disturbance, ecological memory, and the emergence of landscape

pattern. Ecosystems. 2002;5:329–38. https://doi.org/10.1007/s10021-001-0077-1.

24. Canarini A, Schmidt H, Fuchslueger L, Martin V, Herbold CW, Zezula D, et al. Ecological

memory of recurrent drought modifies soil processes via changes in soil microbial community. Nat

Commun. 2021;12:5308. https://doi.org/10.1038/s41467-021-25675-4.

25. Wang B, Allison SD. Drought legacies mediated by trait trade□offs in soil microbiomes.

Ecosphere. 2021;12. https://doi.org/10.1002/ecs2.3562.

26. Wang S, Thi Thu Hoang D, Anh The Luu, Mostafa T, Razavi BS. Environmental memory of

microbes regulates the response of soil enzyme kinetics to extreme water events: Drought-

rewetting-flooding. Geoderma. 2023;437:116593.

https://doi.org/10.1016/j.geoderma.2023.116593.

27. E. Evans S, D. Allison S, V. Hawkes C. Microbes, memory and moisture: Predicting microbial

moisture responses and their impact on carbon cycling. Funct Ecol. 2022;36:1430–41.

https://doi.org/10.1111/1365-2435.14034.

28. Liu L, Estiarte M, Bengtson P, Li J, Asensio D, Wallander H, et al. Drought legacies on soil

respiration and microbial community in a Mediterranean forest soil under different soil moisture

and carbon inputs. Geoderma. 2022;405:115425. https://doi.org/10.1016/j.geoderma.2021.115425.

29. Becchetti T, George M, McDougald N, Flavel D, Vaughn C, Forero L, et al. Rangeland

management series: Annual range forage production. University of California, Agriculture and

Natural Resources. 2016. https://doi.org/10.3733/ucanr.8018.

30. Fossum C, Estera-Molina KY, Yuan M, Herman DJ, Chu-Jacoby I, Nico PS, et al.

Belowground allocation and dynamics of recently fixed plant carbon in a California annual

grassland. Soil Biol Biochem. 2022;165:108519. https://doi.org/10.1016/j.soilbio.2021.108519.

31. Sieradzki ET, Greenlon A, Nicolas AM, Firestone MK, Pett-Ridge J, Blazewicz SJ, et al.

Functional succesision of actively growing soil microorganisms during rewetting is shaped by

precipitation history. Bioarxiv. 2022.

32. Chuckran PF, Estera-Molina K, Nicolas AM, Sieradzki ET, Dijkstra P, Firestone MK, et al.

Codon bias, nucleotide selection, and genome size predict in situ bacterial growth rate and

transcription in rewetted soil. Proc Natl Acad Sci U S A. 2025;122:e2413032122.

https://doi.org/10.1073/pnas.2413032122.

33. Nicolas AM, Sieradzki ET, Pett-Ridge J, Banfield JF, Taga ME, Firestone MK, et al. A subset

of viruses thrives following microbial resuscitation during rewetting of a seasonally dry California

grassland soil. Nat Commun. 2023;14:5835. https://doi.org/10.1038/s41467-023-40835-4.

34. Barnard RL, Osborne CA, Firestone MK. Changing precipitation pattern alters soil microbial

community response to wet-up under a Mediterranean-type climate. ISME J. 2015;9:946–57.

https://doi.org/10.1038/ismej.2014.192.

35. Nuccio EE, Blazewicz SJ, Lafler M, Campbell AN, Kakouridis A, Kimbrel JA, et al. HT-SIP:

a semi-automated stable isotope probing pipeline identifies cross-kingdom interactions in the

hyphosphere of arbuscular mycorrhizal fungi. Microbiome. 2022;10:199.

https://doi.org/10.1186/s40168-022-01391-z.

36. Buckley DH, Huangyutitham V, Hsu S-F, Nelson TA. Stable isotope probing with 15N

achieved by disentangling the effects of genome G+C content and isotope enrichment on DNA

density. Appl Environ Microbiol. 2007;73:3189–95. https://doi.org/10.1128/AEM.02609-06.

37. Fierer N, Jackson JA, Vilgalys R, Jackson RB. Assessment of soil microbial community

structure by use of taxon-specific quantitative PCR assays. Appl Environ Microbiol.

2005;71:4117–20. https://doi.org/10.1128/AEM.71.7.4117-4120.2005.

38. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, et al.

Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. Proc Natl

Acad Sci U S A. 2011;108 Suppl 1 supplement_1:4516–22.

https://doi.org/10.1073/pnas.1000080107.

39. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-

resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13:581–3.

https://doi.org/10.1038/nmeth.3869.

40. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naïve Bayesian classifier for rapid assignment of

rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol. 2007;73:5261–7.

https://doi.org/10.1128/aem.00062-07.

41. R core team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria; 2024.

42. Hungate BA, Mau RL, Schwartz E, Caporaso JG, Dijkstra P, van Gestel N, et al. Quantitative

microbial ecology through stable isotope probing. PeerJ. 2015.

https://doi.org/10.7287/peerj.preprints.1282.

43. Koch BJ, McHugh TA, Hayer M, Schwartz E, Blazewicz SJ, Dijkstra P, et al. Estimating

taxon☐specific population dynamics in diverse microbial communities. Ecosphere.

2018;9:e02090. https://doi.org/10.1002/ecs2.2090.

44. Martin-Laurent F, Philippot L, Hallet S, Chaussod R, Germon JC, Soulas G, et al. DNA

extraction from soils: Old bias for new microbial diversity analysis methods. Appl Environ

Microbiol. 2001;67:4397–4397. https://doi.org/10.1128/aem.67.9.4397-4397.2001.

45. Kanagawa T. Bias and artifacts in multitemplate polymerase chain reactions (PCR). J Biosci

Bioeng. 2003;96:317–23. https://doi.org/10.1016/S1389-1723(03)90130-7.

46. Kozarewa I, Ning Z, Quail MA, Sanders MJ, Berriman M, Turner DJ. Amplification-free

Illumina sequencing-library preparation facilitates improved mapping and assembly of (G+C)-

biased genomes. Nat Methods. 2009;6:291–5. https://doi.org/10.1038/nmeth.1311.

47. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and

graphics of microbiome census data. PLoS One. 2013;8:e61217.

https://doi.org/10.1371/journal.pone.0061217.

48. Bushnell B, Rood J, Singer E. BBMerge – Accurate paired shotgun read merging via overlap.

PLOS ONE. 2017;12:e0185056. https://doi.org/10.1371/journal.pone.0185056.

49. Joshi NA, Fass JN. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ

files. 2011.

50. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. MEGAHIT: an ultra-fast single-node solution

for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics.

2015;31:1674–6. https://doi.org/10.1093/bioinformatics/btv033.

51. Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, et al. DRAM for

distilling microbial metabolism to automate the curation of microbiome function. Nucleic Acids

Res. 2020;48:8883–900. https://doi.org/10.1093/nar/gkaa621.

52. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA:

KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics.

2020;36:2251–2. https://doi.org/10.1093/bioinformatics/btz859.

53. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server for

automated carbohydrate-active enzyme annotation. Nucleic Acids Res. 2018;46:W95–101.

https://doi.org/10.1093/nar/gky418.

54. Steinegger M, Söding J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. Nat Biotechnol. 2017;35:1026–8. https://doi.org/10.1038/nbt.3988.

55. Chuckran PF, Estera-Molina K, Huntemann M, Foster B, Roux S, Mukherjee S, et al.

Metatranscriptomes of California grassland soil microbial communities in response to rewetting.

Microbiol Resour Announc. 2024;13:e0032224. https://doi.org/10.1128/mra.00322-24.

56. Kopylova E, Noé L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs in

metatranscriptomic data. Bioinformatics. 2012;28:3211–7.

https://doi.org/10.1093/bioinformatics/bts611.

57. Aroney STN, Newell RJP, Nissen JN, Camargo AP, Tyson GW, Woodcroft BJ. CoverM: Read

alignment statistics for metagenomics. arXiv [q-bio.GN]. 2025.

58. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol.

2019;20:257. https://doi.org/10.1186/s13059-019-1891-0.

59. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol. 2014;15:550. https://doi.org/10.1186/s13059-014-0550-8.

60. Dittmar T, Koch B, Hertkorn N, Kattner G. A simple and efficient method for the solid-phase

extraction of dissolved organic matter (SPE-DOM) from seawater. Limnology and Oceanography:

Methods. 2008;6:230–5. https://doi.org/10.4319/lom.2008.6.230.

61. Nicora CD, Burnum-Johnson KE, Nakayasu ES, Casey CP, White RA III, Roy Chowdhury T,

et al. The MPLEx protocol for multi-omic analyses of soil samples. J Vis Exp. 2018.

https://doi.org/10.3791/57343-v.

62. Folch J, Lees M, Stanley GHS. A simple method for the isolation and purification of total

lipides from animal tissues. J Biol Chem. 1957.

63. Tolić N, Liu Y, Liyu A, Shen Y, Tfaily MM, Kujawinski EB, et al. Formularity: software for

automated formula assignment of natural and other organic matter from ultrahigh-resolution mass

spectra. Anal Chem. 2017;89:12659–65.

64. Ayala-Ortiz C, Graf-Grachet N, Freire-Zapata V, Fudyma J, Hildebrand G, AminiTabrizi R, et

al. MetaboDirect: an analytical pipeline for the processing of FT-ICR MS-based metabolomic

data. Microbiome. 2023;11:28. https://doi.org/10.1186/s40168-023-01476-3.

65. Tfaily MM, Chu RK, Toyoda J, Tolić N, Robinson EW, Paša-Tolić L, et al. Sequential

extraction protocol for organic matter from soils and sediments using high resolution mass

spectrometry. Anal Chim Acta. 2017;972:54–61. https://doi.org/10.1016/j.aca.2017.03.031.

66. Kuznetsova A, Brockhoff PB, Christensen RHB. LmerTest package: Tests in linear mixed

effects models. J Stat Softw. 2017;82. https://doi.org/10.18637/jss.v082.i13.

67. Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package. 2025.

68. Signorell A. DescTools: Tools for Descriptive Statistics. 2025.

69. Dixon P. VEGAN, a package of R functions for community ecology. J Veg Sci. 2003;14:927–

30. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x.

70. Young R, Avneri-Katz S, McKenna A, Chen H, Bahureksa W, Polubesova T, et al.

Composition-dependent sorptive fractionation of anthropogenic dissolved organic matter by

Fe(III)-montmorillonite. Soil Syst. 2018;2:14. https://doi.org/10.3390/soilsystems2010014.

71. Bahureksa W, Tfaily MM, Boiteau RM, Young RB, Logan MN, McKenna AM, et al. Soil

organic matter characterization by Fourier transform ion cyclotron resonance mass spectrometry

(FTICR MS): A critical review of sample preparation, analysis, and data interpretation. Environ

Sci Technol. 2021;55:9637–56. https://doi.org/10.1021/acs.est.1c01135.

72. Wang B, An S, Liang C, Liu Y, Kuzyakov Y. Microbial necromass as the source of soil

organic carbon in global ecosystems. Soil Biol Biochem. 2021;162:108422.

https://doi.org/10.1016/j.soilbio.2021.108422.

73. Metze D, Schnecker J, Canarini A, Fuchslueger L, Koch BJ, Stone BW, et al. Microbial

growth under drought is confined to distinct taxa and modified by potential future climate

conditions. Nat Commun. 2023;14:5895. https://doi.org/10.1038/s41467-023-41524-y.

74. Veach AM, Zeglin LH. Historical drought affects microbial population dynamics and activity

during soil drying and re-wet. Microb Ecol. 2020;79:662–74. https://doi.org/10.1007/s00248-019-

01432-5.

75. Miltner A, Bombach P, Schmidt-Brücken B, Kästner M. SOM genesis: microbial biomass as a

significant source. Biogeochemistry. 2012;111:41–55. https://doi.org/10.1007/s10533-011-9658-z.

76. Pouran HM. Bacterial cell-mineral interface, its impacts on biofilm formation and

bioremediation. In: Handbook of Environmental Materials Management. Cham: Springer

International Publishing; 2019. p. 535–56. https://doi.org/10.1007/978-3-319-73645-7_80.

77. Wang Q, Ding J, Zhang Z, Liang C, Lambers H, Zhu B, et al. Rhizosphere as a hotspot for

microbial necromass deposition into the soil carbon pool. Journal of Ecology. 2025;113:168–79.

https://doi.org/10.1111/1365-2745.14448.

78. Glaser B, Turrión M-B, Alef K. Amino sugars and muramic acid—biomarkers for soil

microbial community structure analysis. Soil Biol Biochem. 2004;36:399–407.

https://doi.org/10.1016/j.soilbio.2003.10.013.

79. Kögel-Knabner I. The macromolecular organic composition of plant and microbial residues as

inputs to soil organic matter. Soil Biol Biochem. 2002;34:139–62. https://doi.org/10.1016/s0038-

0717(01)00158-4.

80. Manzoni S, Taylor P, Richter A, Porporato A, Ågren GI. Environmental and stoichiometric

controls on microbial carbon-use efficiency in soils. New Phytol. 2012;196:79–91.

https://doi.org/10.1111/j.1469-8137.2012.04225.x.

81. Mendonca CM, Zhang L, Waldbauer JR, Aristilde L. Disproportionate carbon dioxide efflux in

bacterial metabolic pathways for different organic substrates leads to variable contribution to

carbon-use efficiency. Environ Sci Technol. 2024;58:11041–52.

https://doi.org/10.1021/acs.est.4c01328.

82. Cleveland CC, Liptzin D. C:N:P stoichiometry in soil: is there a "Redfield ratio" for the

microbial biomass? Biogeochemistry. 2007;85:235–52. https://doi.org/10.1007/s10533-007-9132-

0.

83. Heyburn J, McKenzie P, Crawley MJ, Fornara DA. Effects of grassland management on plant

C:N:P stoichiometry: implications for soil element cycling and storage. Ecosphere.

2017;8:e01963. https://doi.org/10.1002/ecs2.1963.

84. Mganga KZ, Sietiö O-M, Meyer N, Poeplau C, Adamczyk S, Biasi C, et al. Microbial carbon

use efficiency along an altitudinal gradient. Soil Biol Biochem. 2022;173:108799.

https://doi.org/10.1016/j.soilbio.2022.108799.

85. Xu Q, Li L, Guo J, Guo H, Liu M, Guo S, et al. Active microbial population dynamics and life

strategies drive the enhanced carbon use efficiency in high-organic matter soils. MBio.

2024;15:e0017724. https://doi.org/10.1128/mbio.00177-24.

86. Sharma AK, Khandelwal R, Wolfrum C. Futile cycles: Emerging utility from apparent futility.

Cell Metab. 2024;36:1184–203. https://doi.org/10.1016/j.cmet.2024.03.008.

87. Bremer H, Dennis PP. Modulation of chemical composition and other parameters of the cell at

different exponential growth rates. EcoSal Plus. 2008;3. https://doi.org/10.1128/ecosal.5.2.3.

88. Pellon G. Biosynthesis of peptidoglycans. Biochimie. 1978;60:XI–XII.

89. Díaz E, Jiménez JI, Nogales J. Aerobic degradation of aromatic compounds. Curr Opin

Biotechnol. 2013;24:431–42. https://doi.org/10.1016/j.copbio.2012.10.010.

90. Kanehisa Laboratories. KEGG pathway database. KEGG pathway databse.

www.genome.jp/kegg/pathway.html. Accessed 1 Jan 2023.

1151

Data and code availability statement

The datasets supporting the conclusions of this article are available as follows:

Metatranscriptomics data are available at NCBI Bioproject accession no. PRJNA1088039-

PRJNA1088070 and PRJNA1087216- PRJNA1087833 (doi:10.1128/mra.00322-24 [55]).

Metagenomics data are available at NCBI Bioproject accession no. PRJNA856348. 16S rRNA

gene sequencing data are available at PRJNA1311596. FTICR and CO_2 data are available at

doi:10.6084/m9.figshare.29825273. Code is available at

https://github.com/linneakh/CA_grassland_omics.

1160

Acknowledgements

1162   We thank Jeanette Yuko for creating the conceptual diagram in Figure 6.

1163

Funding

This research was supported by the LLNL 'Microbes Persist' Soil Microbiome SFA, funded by the

U.S. DOE Office of Biological and Environmental Research Genomic Science program (award

SCW1632). The original HREC moisture manipulation experiment and initial soil metagenomic

characterizations were supported by DOE BER awards DE-SC0020163 (MKF), DE-SC0016247

(MKF), SCW1589 (JPR), and SC1421 (JPR). Work at LLNL was conducted under the auspices of

the Department of Energy Contract DE-AC52-07NA27344.

1171

1172   **Author contributions:** JFB, JPR, MKF, and SJB conceived of and designed the experiment.

1173   MMY and KEM helped to run the field study and collect samples. ND and LPT processed samples

1174   and collected data for metabolomics. LKH, PFC, JK, and ETS analyzed data. LKH and SJB wrote

1175   the manuscript.

1176

1177

1178   **Figures**

1179

1180   Figure 1. Taxon-specific growth and mortality rates and bulk community composition. a) Growth
1181   rates for normal (100% MAP) and b) reduced (50% MAP) legacy precipitation. c) Mortality rates
1182   for normal and d) reduced legacy precipitation. e) Bulk changes in community composition as
1183   determined with 16S rRNA gene amplicon sequencing. Colors across panels indicate phylum-level
1184   classifications of taxa.

1185

1186

Figure 2. Microbial growth dynamics upon rewatering post summer dry down for normal (100% MAP) and reduced (50% MAP) legacy precipitation. a) Total biomass calculated from qPCR of 16S rRNA gene and representing both active and nonactive bacteria and archaea. b) Growth rates and c) morality rates. d) Cumulative biomass and e) cumulative CO_2 efflux at each time point post rewatering. f) Community growth efficiency (CGE) metric measured as ratio of new biomass to cumulative CO_2 efflux, normalized to a value between 0 and 1. Differences between legacy precipitation treatments were assessed with Linear Mixed-Effects models (panels a-f) followed by Tukey HSD to compare treatment at each time point (panel a, f). Linear regressions for each legacy moisture treatment were performed to capture linear trend over time (panels b-e) with best fit shown and R^2. For panel f, linear trend shown but R^2 were not significant for 50% legacy precipitation. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$

1198

1199

Figure 3. Soil water extractable organic carbon (WEOC) differs between normal (100% MAP) and reduced (50% MAP) legacy precipitation. A) Van Krevelen Diagram showing unique compounds (only compounds in either 50 or 100% MAP). Ratios of O:C vs H:C allow for predictions of compound classes (showed in doted square boxes). b) Relative abundances of compound classes of unique compounds found in 100%-only and 50%-only as well as for compounds found in both 50 and 100% legacy precipitation treatments. Compounds with a star are significantly overrepresented in respective treatment (G-test). (c-d) Violin plots and embedded boxplots (represent Q1-Q3, centerlines indicate the median and whiskers extend to the minimum and maximum values, excluding outliers [black points]) showing mean and distribution of c) nominal oxidation state of carbon (NOSC) and d) Gibbs free energy (GFE) for unique compounds within each precipitation treatment. For c-d: ***, $p < 0.001$ (unique compounds in normal vs reduced precipitation, LME, df = 1)

1212

1213

1214 Figure 4. KEGG global metabolic pathways of all genes associated a) positively with community
1215 growth efficiency (CGE) and b) negatively associated with CGE. Pathways are highlighted with
1216 color based on category and labeled with description. Categories are colored as: Red- nucleotide,
1217 tan- xenobiotic, purple- energy, blue- carbohydrate, green- lipid, orange-amino acid, and magenta-
1218 vitamin and cofactor metabolism. Pent., pentose; TCA, tricarboxylic acid

1219

1220

1221

Figure 5. Correlations between community growth efficiency (CGE) metric and transcript
abundances (normalized as variance stabilization transformation [VST]) of CGE-associated genes
within specific biosynthesis and energy pathways. Biosynthesis pathways include: a) amino acid,
b) lipid, c) nucleotide and d) peptidoglycan. e) The sum of transcript abundance across all
biosynthesis pathways (amino acid, peptidoglycan, lipids, nucleotides, extracellular polymeric
substances (EPS), and teichoic acids. f) Correlation between CGE metric and ratio of transcript
abundances of biosynthesis: energy genes. Red line indicates linear regression line for significant
correlations. Pearson correlation coefficient (r) and fdr-corrected p-values labeled on each plot. *,
$p < 0.05$; **, $p < 0.01$.

1231

1232

Figure. 6. Conceptual diagram showing flow of carbon between soil organic carbon (SOC),
biomass, and necromass carbon pools under normal and reduced legacy precipitation. Red lines
indicate allocation of carbon to CO_2 for energy, green lines represents allocation to growth, and
blue lines represent general direction and flow of carbon between pools. Thicker lines indicate
greater allocation and flow. Molecules drawn within SOC pool represent biomolecules (i.e.,
DNA), proteins, carbohydrates, lignins, and tannins. Microbial cells are depitcted inside the
biomass pool, and cell death is represented in necromass pool. Number of icons in each pool is
representative of the quanitity of each in each legacy precipitaiton regime.

1241

 Supplementary Figure Captions:

 Figure S1. Nonmetric multidimensional scaling (NMDS) plots of a) water extracted organic

 carbon (WEOC) and b) methanol extraction organic carbon (MEOC) measured using FTICR-MS

 with color indicating timepoint post rewet and shape indicating legacy precipitation, b) WEOC and

 d) MEOC measured using FTICR-MS with color indicating legacy precipitation, e) transcript

 abundances from metatranscriptomics normalized as variance stabilized transformations (VST) of

 read counts with color indication timepoint post rewet and shape indicating legacy precipitation.

 Jaccard distance was performed for MEOC and WEOC (a-d) and Bray☐Curtis dissimilarity was

 performed for gene transcripts (e-g). Statistical grouping of treatments was performed using

 PERMANOVA. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$

 Figure S2. Soil methanol extractable organic carbon (MEOC) differs between legacy precipitation

 regimes. A) Van Krevelen Diagram showing unique compounds (only compounds in either 50 or

 100% MAP). Ratios of O:C vs H:C allow for predictions of compound classes (showed in doted

 square boxes). b) Relative abundances of compound classes in 50%-only, 100%-only, and both

 precipitation treatments. Compounds with a star are significantly overrepresented in respective

 treatment (G-test). (c-d) Violin plots and embedded boxplots (represent Q1-Q3, centerlines

 indicate the median and whiskers extend to the minimum and maximum values, excluding outliers

 [black points]) showing mean and distribution of c) nominal oxidation state of carbon (NOSC) and

 d) Gibbs free energy (GFE) for unique compounds within each precipitation treatment. For c-d:

 ***, $p<0.001$ (unique compounds in normal vs reduced precipitation, Linear Mixed Effects, df =

 1)

Figure S3. Overview of FTICR-MS composition of WEOC and MEOC. Relative abundances of

compound classes across time and for a) WEOC and b) MEOC of each legacy precipitation

regime. Boxplots represent Q1-Q3, centerlines indicate the median and whiskers extend to the

minimum and maximum values, excluding outliers (black points). Linear Mixed Effect model p

values for precipitation (df=1) and timepoint (df=5) are results are labeled in each plot. Significant

pairwise Tukey HSD comparisons between precipitation at each timepoint and timepoints within

each precipitation are shown as lines above the boxes. c) Pie chart of compound classes in WEOC

averaged across all samples. d) Van Krevelen Diagram of each compound in WEOC across all

samples. e) Pie chart of compound classes in MEOC averaged across all samples. d) Van Krevelen

Diagram of each compound in MEOC across all samples. Colors of dotted outlines in Van

Krevelen Diagrams indicate compound classes. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$

1276

Figure S4. Correlations between community growth efficiency (CGE) metric and transcript

abundances (normalized as variance stabilization transformation [VST]) of CGE-associated genes

within specific biosynthesis and energy pathways. Biosynthesis pathways include: a) teichoic acid

and b) extracellular polymeric substances (EPS), and the energy pathways include c) central C

metabolism, d) fermentation, e) electron transport chain (ETC), f) ATPase, and g) pyruvate

oxidation. h) Correlation between CGE metric and all energy metabolism genes. d) Correlation

between CGE metric and ratio of transcript abundances of biosynthesis: PDH gene, the only CO_2-

producing energy gene associated with CGE. Red line indicates linear regression line. Pearson

correlation coefficient (r) and fdr-correct p-values are labeled in each plot. *, $p < 0.05$; **, $p <$

0.01.

1287

Figure S5. Correlations between community growth efficiency (CGE) metric and transcript

abundances (normalized as variance stabilization transformation [VST]) of CGE-associated genes

within degradation pathways. Degradation pathways include: a) amino acid, b) nucleotide, c)

aromatic, d) aromatic ring, and e) lipid degradation, and f) the sum of all these degradation

pathways. Red line indicates linear regression. Pearson correlation coefficient (r) and fdr-corrected

p-values are labeled on each plot. *, $p < 0.05$; **, $p < 0.01$.

1294

1295   Figure S6. CGE-associated CAZy genes determined using PLSR. VIP scores were adjusted based

1296   on whether the sign of the coefficient, indicating direction of correlation of each CAZy gene with

1297   CGE metric. Bars are colored based on the substrate each enzyme degrades, with the exception of

1298   the enzymes involved in biosynthesis, as denoted by "B". Substrate colors indicate sources of

1299   substrate (purple, animals; blue, microbes; brown, plants/microbes; green, plants).

1300

1301   Figure S7. Taxonomy of metatranscriptomics reads. a) Taxonomic profiles of metatranscriptomics

1302   reads to show abundances of prokaryotes (archaea and bacteria), eukaryotes (fungi, plants, and

1303   other), and viruses. b) Ratio of bacteria to fungal read counts across timepoints.

1304
1305