JOINT ANALYSIS OF HUMAN RETROELEMENTS-LINKED HISTONE MODIFICATION PROFILES REVEALS QUICKLY EVOLVING MOLECULAR PROCESSES CONNECTED WITH CANCER

Daniil Nikitin

"BostonGene Technologies" LTD
daniil.nikitin@bostongene.com
ORCID: 0000-0003-1029-1174

ABSTRACT

Human retroelements (REs), which comprise approximately 40% of the genome, have played a pivotal role in the evolution of key molecular processes, such as placental development, by introducing novel regulatory elements near host gene promoters and enhancers. Despite their genomic abundance and regulatory influence, the functional trajectories of REs remain poorly understood. Here, leveraging ChIP-seq profiles of histone modifications (H3K4me1, H3K4me3, H3K9ac, H3K27ac, H3K27me3, and H3K9me3) from five human cell lines deposited in the ENCODE database, we systematically ranked the regulatory impact of REs across 25,075 human genes. Gene sets enriched for promoter- and enhancer-associated RE-linked regulatory sites were identified. Consensus gene sets across cell lines were found to be associated with pathways involved in cancer progression, specifically chronic myeloid leukemia and small cell lung cancer, as well with host defense responses to infection with human T-cell lymphotropic virus type 1. These findings provide new insights into recent human evolution and highlight the ongoing influence of selfish genetic elements on genome regulation and disease susceptibility.

Keywords: retroelements, retrotransposons, genome evolution, chromatin, histone modifications

Introduction.

Human retroelements (REs, also called retrotransposons), namely LTRs, SINEs and LINEs, comprise 40% of the genome [1], [2]. They impact the host genome by providing regulatory sites [3] and novel protein coding genes, manifesting themselves in various human diseases [4]. For example, human endogenous retroviruses (LTR) derived protein coding genes suppress maternal anti fetal immune response in placenta (like syncytin-1 [5]). All REs transpose via an RNA intermediate and then a DNA copy is synthesised and inserted into other genomic locus. Therefore, REs bear transcription factor binding sites enriched with active chromatin marks (H3K4me1, H3K4me3, H3K29ac etc) that can rewire the host gene regulatory networks [6]. On the other hand, REs can be repressed by the host defense mechanisms, such as CpG methylation, which allows to mitigate deleterious effects of new genetic elements insertion near essential genes [7]. In a greater perspective, there is an evolutionary arms race between mobile selfish genetic elements and host defensive systems [8], which leads to such complex innovations as adaptive immunity, including both the microbial (CRISPR-Cas) and the vertebrates one (VDJ-recombination) [8].

In order to decipher the evolutionary impact of REs on human gene regulation, in 2018-2019 we analyzed genome profiles of transcription factors and chromatin modifications in connection with RE loci and their impact onto the adjacent gene expression. In 2018 we studied 225 transcription factors binding profiles in the K562 cell line and measured

regulatory impact of human REs, including the evolutionary young ones (inserted after the radiation of Old World monkeys) [1]. We found that such molecular processes as PDGF, TGF beta, EGFR, and p38 signaling were positively regulated by evolutionary young REs [1]. In 2019 we expanded the analysis, profiling 563 transcription factors binding in 13 cell lines [9], as well as the enhancer-specific chromatin modification H3K4me1 [10] and simultaneously five chromatin marks (H3K4me3, H3K9ac, H3K27ac, H3K27me3 and H3K9me3 [11]). We developed the original method called RetroSpect and showed that such human molecular processes as gene regulation by microRNAs, olfaction, color vision, fertilization, cellular immune response, and amino acids and fatty acids metabolism and detoxification were enriched with RE-specific regulation and hence were quickly evolving under the RE pressure [9].

Despite the significant progress being made, the field remains generally unexplored. Chromatin marks form a complex interactive system with readers, writers and erasers, collectively regulating gene expression [12]. In the previous articles we gathered gene-level RE regulatory impact scores for the following chromatin marks:
- H3K4me1 - active enhancer specific [13]
- H3K27ac - active enhancer specific [14]
- H3K4me3 - promoter specific [15]
- H3K9ac - promoter specific [16]
- H3K27me3 - polycomb repression specific [17]
- H3K9me3 - heterochromatin specific [18]

In the present work we jointly analyse active promoters, enhancers and heterochromatin RE-linked chromatin marks at the level of genes, based on the fact that REs can both activate and inactivate adjacent human genes because the defensive host heterochromatinisation which is relevant in pathology, for example in X-linked dystonia parkinsonism [19]. The proposed comprehensive intersection analysis based on the original RetroSpect method allowed to show that RE-linked promoter histone marks activate chronic myeloid leukemia and small cell lung cancer connected genes in all 5 cell lines under study. Moreover, the RE-linked promoter regulation impacts such general cancer associated processes as cell cycle and cellular senescence. Surprisingly, we observed RE-linked upregulation of human T-cell leukemia virus type 1 infection related genes and the level of promoter specific histone modifications. In contrast, active RE-linked enhancers are more cell type specific and do not form cell line consensus groups of enriched processes. Taken together, these findings push forward our understanding of human RE-impacted evolution of gene regulatory networks in health and disease.

Materials and methods.

Gene-level scores. Scores of RE-linked epigenetic marks regulatory impact were downloaded for 6 chromatin modifications (H3K4me1, H3K27ac, H3K4me3, H3K9ac, H3K27me3 and H3K9me3) and for 5 cell lines (K562, HepG2, GM12878, MCF-7, HeLa-S3) from the previously published papers [10], [11]. The scores were calculated based on more than 1.5 million histone tags for 25075 genes. The following scores were used in the current research:
- GRE, gene RE-linked enrichment score.
- NGRE, normalized gene RE-linked enrichment score.

Formulas and biological meaning of these scores are described in Nikitin et al., 2019a [10].

Selection of RE-linked regulatory impact enriched and deficient genes. According to the original RetroSpect approach [9], for each chromatin modification in each cell line we

plotted NGRE vs GRE in a scatter plot, build a trend line by the method of least squares [20], and top-10% and bottom-10% genes by their distance to the trend line were selected. Genes from the top-10% group (with NGRE score high compared to the GRE one) were considered RE-enriched, the latter group was deemed RE-deficient.

Plots and visualizations. Plots were drawn using the python matplotlib [21] and seaborn [22] packages, Venn diagrams were drawn using the python supervenn [23] package. Principal component analysis (PCA) with centroid approximation was done using the python scikit-learn package [24].

Gene Ontology analysis. Gene Ontology analysis was done using the ShinyGo service from South Dakota State University [25].

Results and discussion

Dimensional reduction and major data properties.
In order to study the degree of cell-specific and chromatin modification-specific differences for the dataset of NGRE and GRE scores in 6 chromatin modifications and 5 cell lines, we performed principal component analysis with centroid approximation (Fig. 1).

Figure 1. Principal component analysis of GRE and NGRE scores. A - PCA of GRE colored by cell line, B - PCA of GRE colored by histone modification, C - PCA of NGRE colored by cell line, D - PCA of NGRE colored by histone modification. Centroid approximations are

drawn with 95% confidence intervals. The palettes for cell lines in A and C are the same, as well as the palettes for chromatin modifications in B and D.

Mapping of different cell lines with the same histone modifications intersects to a high degree (Fig. 1B, 1D) with heterochromatin marks clustering differently compared to the active chromatin ones both for GRE (Fig. 1B) and NGRE (Fig. 1D). Notably, H3K4me1, the active enhancer mark, is mapped separately in both cases, and the second active enhancer one, H3K27ac, is located the most highly differing between the cell lines. Contrastingly, different cell lines do not form isolated groups, albeit they are from different organs and tissues (Fig. 1A, 1C). This corresponds with fundamental differences between chromatin marks in terms of their functioning, whereas cell line specific patterns are less pronounced.

Selection of RE-enriched and deficient genes for individual cell lines and chromatin modifications.

According to the original RetroSpect procedure [10], for all 5 cell lines and 6 chromatin modifications we correlated NGRE and selected RE-enriched and deficient genes (Fig. 2).

Figure 2. Correlations of NGRE (y-axis) and GRE scores for 5 cell lines (shown by rows, names written on the leftmost part of the plot grid) and 6 chromatin modifications (shown by rows, names written on the bottom part of the plot grid). For each plot the linear trend line was built using the least squares method, and top 10% genes with the highest distance to the curve from above are shown in green (RE-enriched genes), top 10% genes with the highest distance to the curve from below are shown in red (RE-deficient genes) and the rest 80% genes are blue.

Out of 30 all combinations, 5 ones showed negative correlations because of two highly divergent linear trends: H3K27ac, H3K4me3, H3K9ac in the cell line HeLa-S3, and H3K4me3, H3K9ac in the cell line MCF-7. The two heterochromatin histone modifications (H3K27me3 and H3K9me3) and the enhancer specific one H3K4me1 showed no negative linear trends, mimicking the patterns observed earlier by us [9]. The major difference of the current approach applied here is the fact that RE-enriched and deficient genes are extracted individually by cell lines and chromatin modification, whereas earlier the GRE scores of different cell lines and epigenomic features (such as transcription factors) were averaged [10]. At this stage of the analysis the functional significance of the selected RE-enriched and deficient genes is unclear, so further intersections and Gene Ontology approaches were applied as described below.

Intersection of the RE-enriched and deficient genes based on different cell line and histone modifications.

In order to understand how REs impact human gene regulation via different histone modifications and in different cell lines, we intersected RE-enriched and deficient genes by different modifications with each other, repeating the procedure in all 5 cell lines (Fig. 3).

*Figure 3. Supervenn plot showing intersections between RE-enriched (abbreviated as 'top')
and RE-deficient (abbreviated as 'bottom') for each histone modification (shown on each
row) for each cell line. A - GM12878, B - HeLa-S3, C - HepG2, D - K562, E - MCF-7. Each*

gene set is shown using a different color. Colored grey rectangles denote the successful intersection (common genes), white and light grey ones indicate no intersection.Intersection groups that have more than 500 genes have gene numbers written in the bottom of each panel.

The intersection analysis shows that different cell lines have different interplay of RE-linked active and repressive chromatin marks. For example, the K562 cell line (Fig. 3D) has a large cluster of genes (1161) that share promoter and enhancer active marks (H3K27ac, H3K4me3, H3K9ac) without any heterochromatin marks - this can be interpreted as genes that are actively regulated by REs. In contrast, in MCF-7 (Fig. 3E) such a cluster is less than 500 genes, and 692 genes are RE-enriched by active marks (H3K27ac, H3K4me3, H3K9ac) and both repressive marks simultaneously, which pinpoints ambiguity of RE-linked epigenetic regulation of the host genes. The same situation is observed in HepG2 (716 genes, Fig. 3C), whereas in GM12878 (Fig. 3A) and HeLa-S3 (Fig. 3B) both types of clusters are present: the one with the active marks only and the one with active and repressive marks (either H3K9me3 or H3K27me3, respectively). This can be connected with the fact the RE evolutionary pressure is exerted at different intensities between tissues, with placenta [26] and neocortex [27] being one of the most invaded by REs (transcriptionally active) and hence quickly evolving in the human lineage.

Promoter and enhancer-level RE-enriched genes selection among the five cell lines. Having the dataset of GRE and NGRE for 5 cell lines and 6 histone modifications, we got 12 sets of RE-enriched and deficient genes in each cell line. In order to get cell line consensus of promoters and enhancers specific RE-enriched genes, we defined enhancers and promoters RE-enriched genes in the following way for each cell line:
-   Enhancer specific genes that are intersection of RE-enriched genes by H3K4me1 and H3K27ac, and RE-enriched genes by heterochromatin marks (H3K9me13 and H3K27me3) are subtracted from this intersection.
-   Promoter specific genes that are intersection of RE-enriched genes by H3K4me3 and H3K9ac, and RE-enriched genes by heterochromatin marks (H3K9me13 and H3K27me3) are subtracted from this intersection in the same way.
For each cell line we obtained a variable number of promoter and enhancer RE-enriched genes, and we intersected them between all the cell lines to get the degree of cell-specificity in RE-linked genes regulation and regulatory evolution (Fig. 4).

Figure 4. Supervenn plot showing intersections of enhancer (enhancers_top) and promoter (promoter_top) specific RE-enriched genes in the 5 cell lines investigated in the current

study. The numbers on the right part of the plot are indicating gene numbers in the sets shown. Each gene set is shown using a different color. Colored grey rectangles denote the successful intersection (common genes), white and light grey ones indicate no intersection. Intersection groups that have more than 100 genes have gene numbers written in the bottom of each panel. Numbers on the top part of the plot are showing the number of gene sets that share the intersection (for cases with more than 100 genes).

The intersection patterns shown in Fig. 5 illustrate the degree of concordance between RE-enriched promoter- and enhancer-associated regulation across different cell lines. Notably, 92 promoter-associated RE-enriched genes were shared across all five cell lines (ranging from 548 to 1,745 genes per cell line), whereas only three enhancer-associated RE-enriched genes were common (from 194 to 566 genes per cell line). This disparity likely reflects the greater variability of enhancer-associated chromatin modifications (H3K4me1 and H3K27ac; Fig. 3), as well as the well-established observation that transcriptional regulation at promoters is more conserved across tissues than at enhancers [28]. Furthermore, promoter- and enhancer-specific RE-enriched gene sets exhibited minimal overlap within each cell line, suggesting that distinct subsets of REs - potentially corresponding to different RE classes - modulate gene expression at the promoter and enhancer levels, respectively. This hypothesis warrants further investigation through integrative computational analyses and experimental validation.

Functional characterisation of the cell line consensus promoter and enhancer-level RE-enriched genes.
The consensus RE-enriched promoter and enhancer genes are shown in the Table 1 below.

Table 1. The list of consensus RE-enriched promoter and enhancer genes.

Promoter RE-enriched genes									
MRPL23	ASMT	MYC	ELMO3	ZBTB46	EPC2	SMIM15	SNORD91A	EIF1B	MRPL43
TMEM50A	TMEM60	PLD6	SMIM15-AS1	KLLN	GADD45A	RNF214	MIR4795	RNF5P1	TRAPPC1
CRYZL1	ACD	MTMR11	RPL7	IP6K1	POLB	DNASE1L1	MYNN	CHD7	
ATP1B1	HMGB1	LAMB2	RAP2A	PSMA7	TTC30B	C5orf34	LZTS2	CNPPD1	
CTRL	CHMP2B	VAPA	MIR5587	AKAP17A	WASHC2C	USF2	KDM6A	RDH10	
WASHC2A	ZBTB7B	PTEN	ATP11B	ANKIB1	CHUK	PKP4	SLC25A5-AS1	OGT	
JADE1	RNF5	PET100	MIR6833	ARF1	TAPT1-AS1	MIR548H4	TXNDC17	ATR	
NAA38	ETV3	C3orf58	CGGBP1	HYAL3	SLC25A5	ATP13A1	ERBIN	CNIH3	
HECTD2	CTC1	CDK5	ST3GAL3	TTC21A	EIF4A1	CDKN2C	MIR1282	HNRNPDL	
CAPN15	MLF2	CPEB2	HNRNPA0	PTMS	LENG9	MCUB	CASC11	SNORD131	

Enhancer RE-enriched genes									
MIR4674	NALT1	LINC011 32							

The genes found were tested for functional enrichment using the Gene Ontology approach [29] via the ShinyGO portal [25] without the background genes, and the enrichment processes were filtered by False Discovery Rate [30] threshold 0.1. The resulting processes for the RE-enriched promoter genes are shown in Table 2.

Table 2. The list of enrichment results for RE-enriched promoter genes.

Enrichment FDR-corrected p-value	Number of genes from RE-enriched promoters in the process	Number of genes in the process	Fold Enrichment	The process with the GO knowledgebase hyperlink
0.02	4	92	12.6	Small cell lung cancer
0.064	3	76	11.4	Chronic myeloid leukemia
0.044	4	126	9.2	Cell cycle
0.016	6	222	7.8	Human T-cell leukemia virus 1 infection
0.064	4	156	7.4	Cellular senescence

Gene Ontology analysis (Table 2) revealed that human REs transcriptionally activate genes associated with two major cancer types - small cell lung cancer and chronic myeloid leukemia - as well as genes upregulated during infection with human T-cell leukemia virus type 1 (HTLV-1), a retrovirus that, similar to HIV, targets CD4□ T cells [31]. These findings raise the possibility that RE-mediated regulatory activity and evolutionary selection may operate, at least in part, through the transcriptional repurposing of pre-existing host defense mechanisms, a phenomenon exemplified by placental evolution [5]. Additional enriched gene sets, including cell cycle regulation and cellular senescence - both canonical hallmarks of cancer [32] - further support the notion that promoter-associated, RE-enriched regulatory programs are linked to oncogenesis. By contrast, no statistically significant functional enrichments were observed for the three enhancer-associated, RE-linked genes identified (*MIR4674*, *NALT1*, and *LINC01132*), two of which are non-coding.

Conclusion.
We investigated the regulatory impact of retroelements (REs) on human gene regulation by analyzing promoter-associated, enhancer-associated, and heterochromatin-associated histone modifications, using ChIP-seq whole-genome profiles from five human cell lines. Clustering and intersection analyses revealed that RE-associated regulatory activity at enhancers exhibits a high degree of cell type specificity, providing a foundation for further studies on the mechanisms underlying this specificity. In contrast, RE-associated regulation at active promoters appeared to be more conserved across different cell types. Notably, REs were found to activate pathways related to cancer-associated processes, including cellular

senescence and cell cycle regulation. These pathways are linked to cancer diseases such as small cell lung cancer and chronic myeloid leukemia, and RE-associated regulatory activity also impacts host cell defense mechanisms against human T-cell lymphotropic virus type 1. Collectively, these findings advance our understanding of the role of REs in human evolution and oncogenesis.

REFERENCES

1. *Nikitin D. et al.* Profiling of Human Molecular Pathways Affected by Retrotransposons at the Level of Regulation by Transcription Factor Proteins // Front. Immunol., 2018 Jan 30;9:30. doi: 10.3389/fimmu.2018.00030.
2. *Lander E.S. et al.* Initial sequencing and analysis of the human genome // Nature, 2001 Feb 15;409(6822):860-921. doi: 10.1038/35057062.
3. *Cardelli M.* The epigenetic alterations of endogenous retroelements in aging // Mech Ageing Dev. 2018 Sep;174:30-46. doi: 10.1016/j.mad.2018.02.002.
4. *Fan T.-J. and Cui J.* Human Endogenous Retroviruses in Diseases // Subcell Biochem. 2023;106:403-439. doi: 10.1007/978-3-031-40086-5_15.
5. *Wang Q. et al.* Molecular mechanisms of syncytin-1 in tumors and placental development related diseases // Discov Oncol. 2023 Jun 16;14(1):104. doi: 10.1007/s12672-023-00702-6.
6. *Göke J. and Ng H.H.* CTRL+INSERT: retrotransposons and their contribution to regulation and innovation of the transcriptome // EMBO Rep. 2016 Aug;17(8):1131-44. doi: 10.15252/embr.201642743.
7. *Chuong E.B. et al.* Regulatory activities of transposable elements: from conflicts to benefits // Nat Rev Genet. 2017 Feb;18(2):71-86. doi: 10.1038/nrg.2016.139.
8. *Koonin E.V.* Viruses and mobile elements as drivers of evolutionary transitions // Philos Trans R Soc Lond B Biol Sci. 2016 Aug 19;371(1701):20150442. doi: 10.1098/rstb.2015.0442.
9. *Nikitin D. et al.* Retroelement-Linked Transcription Factor Binding Patterns Point to Quickly Developing Molecular Pathways in Human Evolution // Cells. 2019a Feb 6;8(2):130. doi: 10.3390/cells8020130.
10. *Nikitin D. et al.* Retroelement-Linked H3K4me1 Histone Tags Uncover Regulatory Evolution Trends of Gene Enhancers and Feature Quickly Evolving Molecular Processes in Human Physiology // Cells. 2019b Oct 8;8(10):1219. doi: 10.3390/cells8101219.
11. *Igolkina A.A. et al.* H3K4me3, H3K9ac, H3K27ac, H3K27me3 and H3K9me3 Histone Tags Suggest Distinct Regulatory Evolution of Open and Condensed Chromatin Landmarks // Cells. 2019 Sep 5;8(9):1034. doi: 10.3390/cells8091034.
12. *Ueberheide B.M. et al.* On the Hunt for the Histone Code // Mol Cell Proteomics. 2024 Dec;23(12):100873. doi: 10.1016/j.mcpro.2024.100873.
13. *Rada-Iglesias A.* Is H3K4me1 at enhancers correlative or causative? // Nat Genet. 2018 Jan;50(1):4-5. doi: 10.1038/s41588-017-0018-3.
14. *Creyghton M.P. et al.* Histone H3K27ac separates active from poised enhancers and predicts developmental state // Proc Natl Acad Sci U S A. 2010 Dec 14;107(50):21931-6. doi: 10.1073/pnas.1016071107.
15. *Liang G. et al.* Distinct localization of histone H3 acetylation and H3-K4 methylation to the transcription start sites in the human genome // Proc Natl Acad Sci U S A. 2004 May 11;101(19):7357-62. doi: 10.1073/pnas.0401866101.
16. *Karmodiya K. et al.* H3K9 and H3K14 acetylation co-occur at many gene regulatory elements, while H3K14ac marks a subset of inactive inducible promoters in mouse embryonic stem cells // BMC Genomics. 2012 Aug 24;13:424. doi: 10.1186/1471-2164-13-424.
17. *Ku M. et al.* Genomewide Analysis of PRC1 and PRC2 Occupancy Identifies Two Classes of Bivalent Domains // PLoS Genet. 2008 Oct;4(10):e1000242. doi: 10.1371/journal.pgen.1000242.

18. *Nicetto D. et al.* H3K9me3-heterochromatin loss at protein-coding genes enables developmental lineage specification // Science. 2019 Jan 18;363(6424):294-297. doi: 10.1126/science.aau0583.

19. *Horváth V. et al.* Mini-heterochromatin domains constrain the cis-regulatory impact of SVA transposons in human brain development and disease // Nat Struct Mol Biol. 2024 Oct;31(10):1543-1556. doi: 10.1038/s41594-024-01320-8.

20. *Charnes A. et al.* The Equivalence of Generalized Least Squares and Maximum Likelihood Estimates in the Exponential Family // J Am Stat Assoc. 1976 Mar; 71(353):169-171. doi: 10.1080/01621459.1976.10481508.

21. "Matplotlib — Visualization with Python." Accessed: Apr. 22, 2025. [Online]. Available: https://matplotlib.org/

22. "An introduction to seaborn — seaborn 0.13.2 documentation." Accessed: Apr. 22, 2025. [Online]. Available: https://seaborn.pydata.org/tutorial/introduction.html

23. Fedor, gecko984/supervenn. (Apr. 15, 2025). Python. Accessed: Apr. 22, 2025. [Online]. Available: https://github.com/gecko984/supervenn

24. "PCA," scikit-learn. Accessed: Apr. 22, 2025. [Online]. Available: https://scikit-learn/stable/modules/generated/sklearn.decomposition.PCA.html

25. *Ge S.X. et al.* ShinyGO: a graphical gene-set enrichment tool for animals and plants // Bioinformatics. 2020 Apr 15;36(8):2628-2629. doi: 10.1093/bioinformatics/btz931.

26. *Yu M. et al.* Endogenous retrovirus-derived enhancers confer the transcriptional regulation of human trophoblast syncytialization // Nucleic Acids Res. 2023 Jun 9;51(10):4745-4759. doi: 10.1093/nar/gkad109.

27. *Garza R. et al.* Single-cell transcriptomics of human traumatic brain injury reveals activation of endogenous retroviruses in oligodendroglia // Cell Rep. 2023 Nov 28;42(11):113395. doi: 10.1016/j.celrep.2023.113395.

28. *Razin S.V. and Ulianov S.V.* Divide and Rule: Phase Separation in Eukaryotic Genome Functioning // Cells. 2020 Nov 15;9(11):2480. doi: 10.3390/cells9112480.

29. *Gene Ontology Consortium et al.* The Gene Ontology knowledgebase in 2023 // Genetics. 2023 May 4;224(1):iyad031. doi: 10.1093/genetics/iyad031.

30. *Benjamini Y. and Hochberg Y.* Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing // J R Statist Soc Ser B Methodol. 1995 Jan;57(1):289–300. doi: 10.1111/j.2517-6161.1995.tb02031.x.

31. *Hirons A. et al.* Human T-cell lymphotropic virus type-1: a lifelong persistent infection, yet never truly silent // Lancet Infect Dis. 2021 Jan;21(1):e2-e10. doi: 10.1016/S1473-3099(20)30328-5.

32. *Hanahan D.*, Hallmarks of Cancer: New Dimensions // Cancer Discov.2022 Jan; 12(1):e31-e46. doi: 10.1158/2159-8290.CD-21-1059.

Author information

Name	Daniil Nikitin
Working place	*"BostonGene Technologies" LTD*
Position	Senior bioinformatician

Academic degree	
Academic title	--
Phone	055 55775459
e-mail	daniil.nikitin@bostongene.com