1 **A tumour-host feed-forward loop contributes to malignant growth in chromosomal**
2 **instability-induced epithelial tumours**

4 Kaustuv Ghosh [1,3], Aishwarya Kunchur[1,3] and Marco Milán[1,2] *

6
[1]Institute for Research in Biomedicine (IRB Barcelona), The Barcelona Institute of Science and

Technology, Baldiri Reixac, 10, 08028 Barcelona, Spain.

[2]Institució Catalana de Recerca i Estudis Avançats (ICREA)

Pg. Lluís Companys 23, 08010 Barcelona, Spain

[3]These two authors contributed equally to this work

[4]Lead Contact

* Correspondence: marco.milan@irbbarcelona.org

Phone/Fax: 34 93 4034902/34 93 4037109

Running title: SASP-driven tumour host interactions

Keywords: aneuploidy, senescence, SASP, super-competition, hippo signalling

Number of figures: 7 (+ 7 supplementary)

Number of tables: 1 (+ 4 supplementary)

Number of characters: 63000 (with spaces)

Author ORCiD/ email id:

Kaustuv Ghosh: 0000-0001-8657-4889 / kaustuv.ghosh@irbbarcelona.org

Aishwarya Kunchur: 0000-0002-0653-6848 / aishwarya.kunchur@irbbarcelona.org

Abstract

Chromosomal instability (CIN), characterized by frequent changes in chromosome number and structure, is common in human carcinomas and often leads to aneuploidy. Drosophila has been instrumental in demonstrating that CIN can promote tumour growth and malignancy through aneuploidy-induced senescence, a state marked by cell-cycle arrest and high secretory activity. Despite extensive chromosomal heterogeneity, we show that these cells share a distinct transcriptional program, with most responses to aneuploidy and senescence regulated at the transcriptional level. Nearly 10% of the most upregulated genes encode secreted proteins of the senescence-associated secretory phenotype. Five of these proteins act additively, locally or systemically, to block proliferation and induce cell death in neighbouring tissues. This non-autonomous cell death feeds back to the tumour to enhance its growth, resembling super-competition and providing insight into tumour–host interactions relevant to human cancer.

Teaser: Aneuploidy-induced senescent cells activate the pro-survival Hippo-Yorkie pathway and the SASP drives tumour host interactions

Introduction

Chromosomal instability (CIN), an increased rate of errors in chromosome segregation during cell division, is one of the most common forms of genomic instability in cancer. It is prevalent in most solid tumours and is closely associated with chemotherapy resistance, reduced patient survival, and metastasis (*1–6*). Traditionally, CIN has been understood to be a driver of genetic diversity within tumours, mainly through the gradual gain of chromosomes that carry oncogenes and the loss of those containing tumour suppressor genes. However, beyond these slow, progressive changes, CIN can also cause sudden and dramatic alterations in the genome. For example, mis-segregated chromosomes can end up in micronuclei, where they often experience extensive DNA damage. This process, known as chromothripsis, leads to the fragmentation and chaotic rearrangement of chromosomes (*7*). In addition to fuelling cancer progression through both gradual and abrupt mechanisms, CIN also causes cytosolic double-stranded DNA, likely originating from ruptured micronuclei. This DNA is detected by cellular sensors, such as cGAS, which normally activate immune responses. However, cancer cells can exploit this pathway to reshape the tumour microenvironment in a way that suppresses immune activity and promotes metastasis (*8–10*).

Although low levels of CIN are widely recognized as a driver of cancer, excessive levels can be harmful, leading to cell death and the suppression of tumour growth (*11–14*). In normal cells, CIN often results in aneuploidy—an abnormal number of chromosomes—which can disrupt cell function (*15*). Studies in yeast and human cells have shown that this imbalance in the number of chromosomes alters protein ratios and interferes with key cellular processes like DNA replication, mitosis, and metabolism (*16–18*). One outcome of such metabolic disruption is the production of reactive oxygen species (ROS). Studies in *Drosophila* and human epithelial cells subjected to CIN have shown that cells with aneuploid karyotypes that affect a large proportion of the genome typically enter a state of senescence, where they permanently stop dividing and begin releasing inflammatory molecules (*19–22*). These molecules help the mammalian immune system identify and eliminate abnormal cells (*23*). Research in *Drosophila* has further revealed that aneuploidy-induced senescence promotes tumour development. The fly genome is organized into four chromosomes [two autosomes, a sex chromosome pair (X and Y), and a tiny dot-like autosome]. In epithelial tissues subjected to CIN, random chromosome mis-segregation events generate cells with whole-chromosome aneuploidies

(thus impacting a large proportion of the genome) that are expelled from tissues as a result of gene

dosage and proteome imbalances, proteostasis failure, and mitochondrial dysfunction (*19, 24, 25*).

Once extruded, these cells undergo apoptosis as a result of ROS-driven activation of JNK and

JAK/STAT signalling pathways (*19, 24, 26, 27*). When fly aneuploid cells are maintained in the tissue

by apoptosis inhibition, they often undergo a JNK-driven transition to a mesenchymal-like state and

enter a form of senescence marked by G2 cell cycle arrest (*19, 28*). Key molecules expressed by

senescent cells include the *Drosophila* orthologs of Wnt and IL-6 [Wingless (Wg) and Upd3,

respectively], which act on the epithelial tissue subjected to CIN to drive tumour growth, to enhance

cell motility of aneuploidy-induced delaminated cells, and even affect systemic hormonal signals that

prevent developmental maturation and cause organismal death (*24, 29–31*).

In this study, we used the *Drosophila* epithelial model of CIN-induced tumorigenesis to characterize

the transcriptional profile of aneuploidy-induced senescent cells. Despite the highly heterogeneous

chromosome content of these cells, we unravel a distinct transcriptional program. This unique

transcriptional program presents evidence that most cellular responses to aneuploidy and

senescence, including cell cycle arrest, activation of autophagy, reorganization of the actin

cytoskeleton, and a highly secretory phenotype, are regulated at the transcriptional level. In addition

to the signalling pathways involved in driving all these cellular responses, we unravel a pro-survival

function of the Hippo-Yorkie signalling pathway in aneuploid senescent cells. Besides the secreted

proteins well known to promote tumour growth, metastasis, and malignancy (*24, 29–31*), we

uncovered a role of dIlp8 and ImpL2 (fly orthologs of Relaxin and IGBP7, respectively (*32–34*)) in

blocking proliferative growth of both wildtype and tumour cells, and identified a role for cytokines Upd1

and Upd3 and the Drosophila Tumour Necrosis Factor (TNF-α) homologue Eiger in non-

autonomously driving cell death in neighbouring wild type cells. Most importantly, we present

evidence that this non-autonomous cell death feeds back to the tumour to enhance its growth, thus

paralleling the phenomenon of super-competition in cancer, an aggressive form of cell competition

where tumour cells actively eliminate neighbouring, healthy wild-type cells to expand its territory (*35–*

37). Overall, our data highlight potential therapeutic strategies to prevent CIN-driven tumour

development by targeting aneuploidy-induced senescent cells and disrupting the relevant interactions

with the tumour host.

Results

Cellular responses to aneuploidy and senescence are reflected at the transcriptional level

We used the wing primordium of *Drosophila*—a highly proliferative epithelial monolayer that forms a two-sided epithelial sac (Figure 1A)—as a model system to characterize the transcriptional profile of cells subjected to CIN. CIN, an increased rate of changes in chromosome structure and number, was induced by depletion of the spindle assembly checkpoint (SAC) gene *rod* (*rough deal*) using an RNAi form expressed with the GAL4/UAS system (*24*, *26*). The *engrailed-gal4* driver, which is restricted to the posterior (P) compartment of the wing primordium (Figure 1A), was used to limit the induction of CIN to a defined cell population. One of the most attractive features of this model of CIN is that it facilitates the identification of three well-defined cell populations with distinct behaviours (Figure 1B). On one hand, if apoptosis is blocked by the expression of p35 (a baculovirus protein that binds and represses effector caspases Dcp-1 and Drice (*38*)), those cells that become aneuploid - as a consequence of random chromosome missegregation events in mitosis caused by *rod* depletion - delaminate and become cell cycle-arrested, motile, and highly secretory. These cells can then be tracked with MMP1-GFP, which is a GFP-based transcriptional reporter of JNK (*39*) (Figure 1B, green cells). On the other hand, the epithelial tissue subjected to CIN (the P compartment), which consists of cells that remain largely euploid, enters a hyperplastic program that relies on high mitotic activity and unlimited growth potential (Figure 1B, cells located apically to the green cells). Finally, the anterior (A) compartment of these wing primordia (Figure 1B, cells labelled by the Ci marker in red), which are not subject to CIN and whose cells are euploid, will be treated as the neighbouring cell population. The transcriptional profile of these three cell populations (CIN, aneuploid cells, and neighbouring cell population) was compared to the transcriptional profile of control wing primordia expressing p35 under the control of the *engrailed-gal4* driver. Four distinct biological replicates were analysed. Principal Component Analysis (PCA) revealed significant transcriptomic differences between the five populations, showing strict clustering based on their respective traits (Figure 1C). Interestingly, the aneuploid cell population cluster presented the highest level of difference when compared to the compartment of origin [P(CIN)] or the control tissue [P(Control)]. Consistent with this observation, the aneuploid population showed a higher number of upregulated and downregulated genes than the CIN epithelium (Figure 1D, F, see also Figure S2A). However, there was a certain degree of overlap in

the type of genes that changed in these two populations when compared to control tissues. We thus

analysed the transcriptional profile that was unique to aneuploid cells. Remarkably, many of the

previously observed cellular responses to aneuploidy were reflected at the transcriptional level, as

shown by gene set enrichment analysis (GSEA, Figure 1E, Figure S1A, and Table S1). First, high

levels of proteotoxic stress caused by unbalanced karyotypes were reflected by strong upregulation

of genes involved in autophagy (e.g., *atg4a*, *atg8a*, *atg9*, *atg17*, and *atg18b*) and lysosome function

(e.g., *Lamp1*, *Vps13D*, *Vps16A, Vamp7,* and *Iqf*). Second, consistent with the activation of the JNK

stress signalling pathway as a consequence of mitochondrial dysfunction and ROS (*19*, *27*), many of

its target genes (e.g., *mmp1*, *ets21C*, and *scaf*) were strongly upregulated in aneuploid cells (Figure

1E, 2A, and Figure S1A). Third, the ability of aneuploid cells to become motile through modulation of

the actin cytoskeleton and a non-apoptotic role of caspases (*28*, *31*) was reflected by strong

upregulation of genes involved in these two processes, as well as in cell motility (e.g., *sqh*, *cortactin*,

rok, *fak*, *Arpc2-5, hid*, *rpr*, *Dronc*, *Drice*, and *Dcp-1;* Figures 1E, 2A, and Figure S1A). Finally, entry

into a senescence-like state [1] was also reflected at the transcriptional level, as shown by strong

downregulation of genes involved in cell cycle progression (e.g., *stg*, *cdk1*, *cycA*, and *cycB*) and DNA

replication (e.g., *DNA polymerase*, *Orc* and *Mcm* complexes, Figure 1E and 2B), and the clear

upregulation of genes involved in secretion (e.g., *Tango 5, 11, Sec 5, 6, 33, 24CD, and Rab 1, 2, 6,*

7, 10, and *35*) and encoding for secreted proteins (Figure 1E, 2A, and S1A). Indeed, 10% of the most

upregulated genes encoded for secreted proteins (Table 1 and S2). These included the systemic

signals Upd3 cytokine (involved in CIN-induced invasion and developmental delay (*30*, *31*)), Dilp8

relaxin (mediating developmental delay (*30*, *31*)), Pvf1 (PDGF- and VEGF-related factor 1 involved

in driving organ wasting (*40*)) and the mitogenic signals Wg and Wnt6 (*29*), many of them are known

targets of JNK. We confirmed the highly secretory phenotype of senescent cells by using three

fluorescent reporters to mark the enlargement of the Golgi and the endoplasmic reticulum (ER)

compartment, and an increase in the number of exosomes (Figure 2C). The transcriptional profile of

aneuploid cells that was shared with CIN comprised genes involved in the activation of ERK and

JAK/STAT signalling, previously shown to contribute to various aspects of CIN-induced

tumorigenesis, such as growth and invasiveness (*28*, *31*) (Figure 1G and S2B, C). In CIN cells, we

observed unique upregulation of the genes involved in DNA damage, consistent with the high

propensity of dividing cells subjected to SAC depletion to accumulate lagging or broken chromosomes

(Figure S2B, C). Overall, these results indicate that the cellular responses to aneuploidy and

senescence are reflected at the transcriptional level, and they reinforce the highly secretory

phenotype of aneuploidy-induced senescent cells.

A survival role of hippo/Yorkie in CIN-induced senescent cells

In CIN-induced tumours, the ability of aneuploid cells to become motile relies on modulation of the

actin cytoskeleton and a non-apoptotic role of caspases (*28*, *31*). Consistent with this, activation of

the apoptotic pathway in CIN tissues (monitored with cDcp1 antibodies, Figure 3D, E) was not

followed by apoptosis execution (monitored by TUNEL staining, Figure 3C). Activation of the apoptotic

pathway in aneuploidy-induced senescent cells was reflected by strong upregulation of genes

involved in apoptosis, including the pro-apoptotic genes *hid* and *reaper*, the initiator caspase *Dronc,*

and the effector caspases *Drice* and *Dcp-1 (*Figure 3A). The increase in *hid* expression was confirmed

by a *hid-lacZ* reporter (Figure 3B). Interestingly, the activity levels of effector Caspases, monitored by

the GC3Ai activity reporter (*41*), were relatively high in aneuploidy-induced senescent cells when

compared to control tissues (Figure 3B) despite the expression of the p35 baculovirus protein, a

potent inhibitor of these Caspases (*38*). However, there were no pyknotic nuclei in the tissue

suggesting that apoptosis execution is indeed actively blocked in these cells. In our search for

candidate pathways with well-known pro-survival activity, we observed that many elements of the

Hippo/Yorkie signalling cascade, including *crumbs* (*crb*), *fat* (*ft*), *Merlin* (*Mer*), *Ajuba* (*jub*), *kibra*, *hippo*

(*hpo*), and *yorkie* (*yki*), were transcriptionally activated in aneuploidy-induced senescent cells (Figure

3A). The activation of Yorkie in these cells was confirmed with the use of GFP-protein traps of Ajuba

(Jub-GFP, an upstream negative regulator of Hippo (*42*)) and Yorkie (Figure 3C, F), transcriptional

reporter and GFP-protein trap of *Diap1* (an E3 ubiquitin ligase regulated by Yorkie and involved in

inhibiting Dronc, (*43*), Figure 3E and S3), and transcriptional reporters of *bantam* (a miRNA regulated

by Yorkie and involved in inhibiting *hid* (*44*, *45*)), and *expanded* [an upstream regulator of the Hippo

pathway and Yorkie target (*46*), Figure 3D, S3]. Yorkie has been previously shown to be activated

by JNK (*47*), which is highly active in CIN-induced aneuploid cells and plays a pivotal role in defining

their senescent state (*19*). Consistently, the expression levels of Yorkie targets *Diap1* and *bantam*

were significantly reduced upon blockage of the JNK pathway by expression of a dominant negative

version of JNK (Bsk in *Drosophila* (*48*), Figure 3D, E, quantified in Figure 3D', E'). Given the well-

known anti-apoptotic role of the Yorkie pathway, we addressed whether Yorkie depletion had any

impact on the survival of CIN tissues. To this end, and based on the developmental role of Yorkie in

regulating growth and survival of epithelial tissues (*49*), we searched for a mild *UAS-yorkie-RNAi*

transgene. Interestingly, depletion of Yorkie (with the use of this mild transgene) did not have a major

effect on the size or survival of p35-overexpressing tissues but did have a mild effect on the size and

survival of otherwise wild-type tissues (Figure 4A, C, left panels, and quantified in Figure 4B, D). In

contrast, the use of this transgene induced a strong negative impact on the size of CIN tissue, and

this reduction was accompanied by a significant increase in the number of apoptotic cells (Figure 4A,

C, right panels, and quantified in Figure 4B, D). In experimental conditions where apoptosis was not

blocked, the CIN tissue collapsed and was transformed into isolated islands of dying cell populations

(Figure 4C, right panel). Taken together, these results indicate that the Hippo/Yorkie pathway acts as

an intrinsic anti-apoptotic pathway that contributes to dampening the deleterious effects of CIN and

maintaining senescent cells alive despite the activation of the apoptotic pathway. Consistent with the

proposal of this intrinsic anti-apoptotic pathway, we observed that by halving the doses of the pro-

apoptotic genes *reaper*, *hid* and *grim* (in *Df(H99)/+* larvae), the levels of cell death observed in CIN

tissues were reduced (Figure 4E, and quantified in Figure 4I), and aneuploid cells, labelled by the

expression of MMP1, were maintained in the tissue even in the absence of p35 (Figure 4E and F).

These cells were negative for apoptotic markers (Figure 4G) and were not the result of loss of Df(H99)

heterozygosity caused by CIN, as they expressed the *bantam-lacZ* reporter located in the

homologous chromosome (Figure 4H). Thus, aneuploid cells were not maintained in the tissue in this

experimental setting as a trivial consequence of inheriting the two chromosomes bearing the *Df(H99)*

deficiency. Most interestingly, under these circumstances, the tissue also overgrew (Figure 4E, F,

and quantified in Figure 4I).

A negative effect of CIN tumours on growth and survival of neighbouring cells

We noticed that the transcriptional profile of the neighbouring cell population (cells located in the

nearby A compartment and not subject to CIN, Figure 1B) showed upregulation of genes involved in

cell death and a reduction in translation activity (Figure S4A-C). We next examined the response of

this cell population (A compartment) to the nearby hyperplastic tissue (P compartment) subject to CIN

and additional blockage of the apoptotic pathway at different time points of development. As

anticipated, the P compartment showed a steady increase in size, whereas the size of the A

compartment was significantly reduced compared with controls (Figure 5A, quantified in Figure 5B).

After eight days of development, the A compartment reached the expected size of a control A

compartment, while the P compartment continued to grow. The differential growth dynamics of CIN

and neighbouring cell populations were also observed when analysing the ratio of their corresponding

sizes [P/A ratio; Figure 5B]. Interestingly, the negative impact of the CIN tissue on the growth

dynamics of the A compartment was accompanied by an increase in the number of dying cells

labelled by TUNEL staining (Figure 5A, quantified in Figure 5C), a reduction in mitotic activity labelled

with pH3 antibody (Figure 5D, quantified in Figure 5C), and a decrease in EdU incorporation (Figure

5E, quantified in Figure 5C). Non-autonomous cell death was also observed upon induction of CIN in

other regions of the wing primordium (Figure S4D).

We next addressed whether the observed response of the A compartment to the nearby CIN tissue

was driven by any of the signalling molecules secreted by aneuploidy-induced senescent cells (Table

1). Interestingly, Dilp8, a molecule secreted by damaged tissues and well known to block the

production of Ecdysone (*50, 51*), a systemic hormone driving the larval to pupal transition and

promoting proliferative growth, was among the most upregulated secreted proteins in aneuploidy-

induced senescent cells (Figure 2A and Table 1). Using a GFP protein trap reporter, we confirmed

the expression of Dilp8 in aneuploidy-induced senescent cells (Figure 6A). Depletion of *dilp8* in the

CIN tissue (with a *UAS-RNAi*) restored mitotic activity and EdU incorporation levels, but not the

number of apoptotic cells, in the adjacent A compartment (Figure 6B and C, quantified in Figure 6B'

and C', Figure S5A, B). Interestingly, Dilp8 overexpression reduced mitotic activity and the size of the

A compartment even further (Figure 6C, quantified in Figure 6C'). Since experimental manipulation

of Dilp8 levels in the P compartment was unable to have a significant effect on the size of the nearby

A compartment (Figure 6D), we searched for additional genes expressed in aneuploidy-induced

senescent cells with a known role in inhibiting proliferative growth and found ImpL2, an orthologue of

IGFBP7 that binds to and inhibits the function of *Drosophila* insulin-like peptides (*52, 53*) and

consequently the growth-promoting PI3K signalling pathway (Table 1), an upregulation that was

confirmed with a GFP protein trap reporter (Figure 6A). Depletion of *ImpL2* in the CIN tissue (with a

UAS-RNAi) restored mitotic activity and EdU incorporation levels (Figure 6B and C, quantified in

Figure 6B' and C') in the adjacent A compartment but not number of apoptotic cells (Figure S5A, B),

and ImpL2 overexpression reduced mitotic activity even further (Figure 6C, quantified in Figure 6C').

As observed in the case of Dilp8, experimental manipulation of ImpL2 levels had no significant impact

on the size of the A compartment either (Figure 6D). Interestingly, co-depletion of both *dilp8* and

impL2 was able to dampen the reduction in the size of the A compartment caused by the nearby CIN

tissue (Figure 6D). However, we noticed that the size of the P compartment was also dramatically

increased upon *dilp8* and *impL2* co-depletion (Figure 6E), thus resulting in visibly bigger tumours

(Figure 6F). These results are consistent with the previously reported systemic effects of these two

secreted factors (*51, 53, 54*), and point to an additive tumour-suppressive role of these two systemic

molecules, most probably by interfering with Ecdysone and dILPs, two growth-promoting systemic

hormones. The reduction in the proliferative activity of the A compartment was not observed in those

cells abutting the CIN tissue (Figure 5E and Figure S5D), thus pointing to a signal originating in the

P compartment and counteracting the negative effects of Dilp8 and ImpL2 on proliferation. Indeed,

the proliferative activity of these cells relies on the activity of the mitogenic molecule Wg produced by

senescent cells (Table 1, (*29*), see also Figure S5C), as Wg depletion reduced the levels of EdU

incorporation in these cells (Figure S5D, E).

A feed-forward loop driving tumour growth

The non-autonomous induction of cell death and reduction in tissue size caused by CIN tumours was,

at least in part, a consequence of activation of the apoptotic pathway as halving the doses of the pro-

apoptotic genes *reaper*, *hid* and *grim* (in *Df(H99)/+* larvae) caused a reduction in the levels of non-

autonomous cell death (Figure 7D, D') and restored the size of the nearby A compartment (Figure

7E). We noticed that targeted depletion of these three genes with the use of a synthetic micro-RNA

[*UAS-miRHG*, (*55*)] only in the P compartment did not have any effect on the levels of cell death and

on the size of the A compartment (Figure 7D, D', E). In search of secreted molecules involved in the

non-autonomous induction of the apoptotic pathway (Table 1), we noticed that aneuploidy-induced

senescent cells showed upregulation, on the one hand, of elements of the JAK/STAT pathway,

including ligands (cytokines Upd1, Upd2, and Upd3, Table 1) and the receptor Domeless (Dome,

Figure 7A), and, on the other hand, the TNF-α ligand Eiger (Figure 2A and Table 1), which activates

the JNK signaling pathway through its receptor Grindelwald (Grnd, (*56*)). Both JNK and JAK/STAT

signalling pathways are known in CIN tissues to induce apoptosis through transcriptional induction of

pro-apoptotic genes (*31*). We confirmed activation of the JAK/STAT pathway with a GFP

transcriptional reporter (STAT-GFP (*57*)), and, most interestingly, we observed that it was also

activated in the nearby A compartment (Figure 7B). We confirmed induction of *eiger* expression with

a transcriptional reporter (Figure 7C, see also (*27*)). We next addressed the potential contribution of

Upd ligands, the JAK/STAT pathway and Eiger to the effects of CIN tissues on the nearby wild type

cell population.

Expression of a truncated form of receptor Domeless lacking the intracellular domain in the CIN tissue

(Dome-ΔCYT) led to a marked reduction in the levels of STAT-GFP signalling, not only cell

autonomously but also in the nearby A compartment (Figure 7B). This finding is consistent with the

proposed role of Dome-ΔCYT in blocking JAK/STAT signalling by trapping the Upd ligands and

competing with the endogenous receptor (*30*). Interestingly, the non-autonomous reduction in STAT-

GFP levels caused by Dome-ΔCYT was accompanied by a decrease in the number of apoptotic cells

(Figure 7D, quantified in Figure 7D'). In contrast, depletion of Dome by RNAi, which is expected to

block ligand-dependent JAK/STAT signalling in a cell autonomous manner without trapping the Upd

ligands, did not affect the neighbouring A compartment (Figure 7D, quantified in Figure 7D'). We

found that both transgenes (*UAS-dome-ΔCYT* and *UAS-dome-RNAi*) caused a similar reduction in

the size of the P compartment, most probably caused by the developmental role of the JAK/STAT

pathway in regulating the growth of this compartment (*58*). The non-autonomous reduction in the

number of apoptotic cells caused by Dome-ΔCYT was also observed in tissues mutant for *hop* (the

Drosophila JAK orthologue) and upon depletion of the cytokines *upd1 or upd3* by RNAi in CIN and

aneuploid cells (Figure 7D, quantified in Figure 7D'). We observed that neither *upd1* depletion nor

Dome-ΔCYT expression in CIN tissue ameliorated the non-autonomous reduction in proliferation

dynamics (Figure S6A, quantified in Figure S6B).

In a similar manner, depletion of *eiger* or expression of a dominant negative version of its receptor

Grnd (Grnd-extra) in CIN tissues, known to trap Eiger ligand thus blocking its function (*56*), reduced

the number of apoptotic cells in the nearby A compartment (Figure 7D, quantified in Figure 7D').

Depletion of *grnd* by RNAi, which is expected to block Eiger-dependent JNK activation in a cell

autonomous manner without trapping the Eiger ligand, did not affect the apoptotic levels in the

neighbouring A compartment (Figure 7D, quantified in Figure 7D'). Most interestingly, depletion of

Upd1, Upd3 or Eiger rescued the reduction in the size of the A compartment (Figure 7E). Similar

results were observed with Grnd-extra (Figure 7E). All these results point to an additive role of multiple

molecules secreted from aneuploidy-induced senescent cells and acting in nearby wild type tissues

to induce cell death and compromise growth.

We next analysed whether the non-autonomous cell death had any impact on the growth of the

tumour itself. Interestingly, the reduction in the levels of non-autonomous cell death and the

restoration of the size of the A compartment caused by depletion of Eiger, the Upd1 and Upd3

cytokines or expression of their truncated receptors was in all cases accompanied by a reduction in

the size of the hyperplastic tissue (P compartment) subject to CIN and additional blockage of the

apoptotic pathway (Figure 7F, G). We noticed that depletion of *grindelwald* by RNAi did not have any

impact on the size of the P compartment, thus ruling out an autocrine role of Eiger in CIN tumors

[Figure 7F, see also (*27*)]. Depletion of several genes of the JAK/STAT signalling pathway by RNAi

was previously shown to have no impact on the size of CIN tumours either, thus ruling out also an

autocrine role of Upd cytokines (*31*). These results indicate that several SASP members act in an

additive manner on neighbouring tissues to reduce growth through cell death and that these non-

autonomous effects feed back to the tumour to reinforce its growth.

Although the classical apoptotic pathway appears to play a major role in the non-autonomous

induction of cell death and reduction in tissue size exerted by Upd cytokines and Eiger, as cell death

and tissue size were both largely rescued by halving the doses of the pro-apoptotic genes (Figure

7D, D', E), non-apoptotic cell death might also contribute to the observed effects on tissue size. On

the one hand, Eiger and the JNK pathway have been reported to induce also non-apoptotic cell death

through the production of ROS (*59*). On the other hand, Upd cytokines and the JAK/STAT pathway,

well known to induce autophagy in CIN tissues (*19*), might also cause cell death through autophagy.

Consistent with this proposal, we noticed that in those regions within the nearby A compartment with

the highest levels of apoptotic activity (monitored with an antibody to detect the cleaved version of

the effector caspase Dcp1, cDcp1), autophagy was also induced, as reflected by the accumulation in

puncta of a mCherry-tagged Atg8a under the control of the endogenous Atg8a promoter (Atg8a;

Figure S7A). This induction was largely dependent on Upd1 derived from CIN tissue, as *upd1*

depletion led to a significant reduction in the amount of Atg8a signal in the A compartment (Figure

S7A, A'). Compromising autophagic flux, by supplementing the larval food with chloroquine, led to a

significant reduction in the extent of cell death observed in the A compartment (Figure S7B, B'). We

noticed that caspase activation in CIN cells, which play a non-apoptotic role by inducing cell motility

in aneuploid cells, was not reduced with this treatment, which is consistent with their reported

activation through transcriptional induction of pro-apoptotic genes by JNK and JAK/STAT signalling

pathways (*31*). Overall, these observations point to the potential contribution of different forms of cell

death in the observed feed-forward loop involved in the growth of CIN tumours.

Discussion

The transcriptome of aneuploidy-induced senescence

From flies to mammals, CIN is widely recognized as a driver of cellular senescence, primarily through

the generation of highly aneuploid karyotypes (*19–22*), and cellular senescence and its associated

senescence-associated secretory phenotype (SASP) are major triggers of inflammation and

tumorigenesis (*60*). Here we have characterized the transcriptional profile of aneuploidy-induced

senescence in an epithelial model of CIN in *Drosophila* that recapitulates most emerging cellular

behaviours observed in mammalian epithelial tissues upon CIN (*10*, *22*, *24–28*, *61–63*). Thus, in fly

epithelial tissues subjected to CIN and upon additional blockade of the apoptotic pathway, aneuploid

cells are extruded from the epithelium and enter a state of senescence characterized by cell cycle

arrest in G2, increased motility, and a highly secretory phenotype that drives the hyperplastic growth

of the epithelium in a non-autonomous manner (*64*). Consequently, CIN tissues acquire a tumour-

like, unlimited growth potential and highly invasive behaviour. All these processes rely on the activity

of the JNK signalling pathway. As it occurs from yeast to mammals, aneuploidy activates autophagy

as the primary quality control mechanism for maintaining proteostasis in response to gene dosage

imbalance and proteotoxic stress (*65, 66*). Saturated use of the autophagy machinery in these cells

leads to the accumulation of dysfunctional mitochondria as a result of compromised mitophagy, and

these mitochondria become active sources of ROS (*19*), which ultimately activate JNK via Ask1 (*27*).

The use of the GAL4/UAS system to trigger CIN in only one half of the growing wing epithelium and

a GFP transcriptional reporter of JNK (MMP1-GFP, (*39*)) to track aneuploid cells has allowed us to

characterize the transcriptional profile of aneuploidy-induced senescent cells, the epithelial layer

subjected to CIN that enters a hyperplastic mode of growth, and the nearby wild-type population.

Remarkably, most cellular responses to aneuploidy and senescence are robustly reflected at the

transcriptional level, as shown by the up- or down-regulation of a large number of genes involved in

any of these responses (Figures 1 and 2). Thus, cell cycle progression is locked in aneuploid cells

due to the downregulation of most genes involved in DNA replication (including all subunits of the

DNA polymerase, and Orc and Mcm complexes) and G2/M transition (including *Dcdc25*/*stg*, *cdk1*,

cycA, and *cycB*). The induction of autophagy, a process that delivers its cargo to lysosomes for

degradation, is accompanied by the upregulation of many autophagy (*atg4a*, *atg8a*, *atg9*, *atg17,* and

atg18b) and lysosomal genes (e.g., *Lamp1*, *Vps13D*, *Vps16A, Vamp7,* and *lqf*). Similarly, the

reported ability of aneuploid cells to become motile through modulation of the actin cytoskeleton is

accompanied by the upregulation of many actin cytoskeleton proteins, including Sqh, Cortactin, Rok,

Fak, and Arpc2-5*.* One of the most remarkable properties of senescent cells is the SASP (*67*). Our

transcriptomic analysis has unravelled a significant upregulation of genes involved in secretion (e.g.,

Tango 5, 11, Sec 5, 6, 33, 24CD, and *Rab 1, 2, 6, 7, 10,* and *35*), reinforced with the use of cellular

reporters to unveil an enlarged secretory apparatus, and has allowed us to identify all the elements

of the SASP, where 10% of the most upregulated genes in these cells encode for secreted proteins

(Table S2). Remarkably, many of the cellular behaviors observed in our fly model of CIN have been

observed in recent years in mammalian tissues, thus reinforcing its translational impact. Thus,

mammalian cells with complex karyotypes caused by chromosome segregation defects are also

extruded from the epithelium (*61*), and they are cell cycle-arrested, exhibit features of senescence,

and produce pro-inflammatory signals (*22*). Most interestingly, CIN also drives tumorigenesis in

mammals [e.g. (*62*)] and promotes metastasis (*10*).

An anti-apoptotic role of Yorkie in senescent cells

Drosophila CIN tissues are exposed to the activation of pro-apoptotic pathways JNK and JAK/STAT

that lead to the elimination of aneuploid cells in the tissue (*24*, *31*). It is the blockage of the apoptotic

pathway by the expression of the baculovirus protein p35 what unravels a subversive role of JNK in

driving cellular senescence, and JAK/STAT in promoting cell motility. However, the apoptotic pathway is not completely blocked under these conditions as shown by the high activity levels of effector caspases in the tissue, which have been shown to contribute in a non-apoptotic manner to the invasive behaviour of aneuploid cells. These results point to the presence of an endogenous pathway that counteracts the pro-apoptotic activities of JNK and JAK/STAT. Similar to how senescent cells in mammals resist normal apoptotic pathways through expression of the Bcl2 anti-apoptotic protein (*68*), our results indicate that the Hippo/Yorkie pathway plays an analogous role in senescent fly cells. Activation of this pathway is most probably the result of JNK activation and cellular tension positively regulating the cortical localization of Ajuba expression to sequester Warts, thus allowing Yorkie to translocate into the nucleus and drive gene expression (*47, 69*). Although DIAP1 (a negative regulator of the initiator Caspase Dronc (*43*)) or bantam miRNA (which acts negatively on the hid 3'UTR (*70*)) could be two potential target genes of Yorkie that block the death of senescent cells, the finding that effector caspases show high activity in these cells suggests that Yorkie negatively impinges on the apoptotic pathway even at the level of cell death execution. These experimental data contribute to the identification of Yorkie as a potential therapeutic strategy to block CIN-induced tumorigenesis by targeting senescent cells.

The SASP mediates a tumour-host feed-forward loop: a case of super-competition

In our fly epithelial model of CIN, senescent cells show a remarkable ability to act at a paracrine, autocrine, and systemic level to drive tumour growth, invasiveness, and malignancy, respectively [2]. They exert their action through elements of the SASP, including mitogenic molecules Wg and Wnt6, Matrix metalloproteinase MMP1, and cytokines such as Upd3. Here we identify a new axis of interaction between senescent cells with the nearby wild type cell population that contribute to tumour growth (Figure 7H). While the CIN tissue enters a hyperplastic mode of growth in response to the mitogenic molecules Wg and Wnt6 secreted by senescent cells, the nearby wild type cell population shows a reduction in growth rates, which is accompanied by a reduction in proliferative activity and an increase in the number of apoptotic cells. We identified several components of the SASP driving the observed non-autonomous effects: the Relaxin-like molecule Dilp8, the Insulin-like growth factor antagonist ImpL2 (the fly ortholog of IGFBP-7), the Upd1 and Upd3 cytokine and the TNF-α-orthologue Egr. While Dilp8 and ImpL2 were shown to be required and sufficient to drive the non-

autonomous reduction in proliferative activity, Upd1, Upd3 and Eiger were found to be responsible

for the reduction in growth and increase in apoptotic levels of the neighbouring cell population. We

present evidence that Upd1 and Upd3 exerts its action by triggering the JAK/STAT signalling pathway

and by inducing cell death. Consistent with the reported activities of Dilp8 and ImpL2 as systemic

signals blocking the production or activity of those systemic hormones required for organismal growth

[eg. ecdysone and dILPs, (*34*, *50*, *51*, *71*)], their negative impact on the proliferative activity was

found to be extended to the tumour itself and not just to neighbouring tissues. These data expand

the repertoire of signalling activities exerted by senescent cells and provide another example of non-

cell-autonomous induction of cell death in the tumour host (*72*). Most interestingly, we present

evidence that the non-cell-autonomous induction of cell death induced by Upd cytokines and Eiger

is required to feed back to the tumour to enhance its growth, either as a consequence of the liberation

of nutrients and metabolites, or the expression of signalling molecules by dying cells before being

captured by circulating macrophages (Figure 7H). These tumour-host interactions parallel interface

interactions observed during super-competition, an aggressive form of cell competition where a cell

acquiring a specific mutation (often oncogenic) becomes "super-fit" and actively eliminates

neighbouring, healthy wild-type cells to expand its territory (*35–37*).

Materials and Methods

Fly maintenance, husbandry and transgene expression

Strains of *Drosophila melanogaster* were maintained on standard medium (4% glucose, 55 g/L yeast,

0.65% agar, 28 g/L wheat flour, 4 ml/L propionic acid, and 1.1 g/L nipagin) at 25°C in light/dark cycles

of 12 h. The resulting larvae after the egg laying (AEL) at 25°C were either transferred to 29°C, or

maintained at 25°C for specific experiments as mentioned in the figures, and were dissected on the

day AEL as mentioned in the figures. The sex of experimental larvae was not considered relevant to

this study and was not determined. The strains used are summarized in Table S4.

Standard induction of CIN, and endogenous labelling of different tissue subpopulations for

transcriptomic analysis

Virgin female flies carrying *UAS-myrT, MMP1-GFP* transgenes were crossed with males carrying

either *en-GAL4, UAS-p35* (Control) or *en-GAL4, UAS-p35, UAS-rod^RNAi* (CIN) transgenes. Fertilized

females were allowed to lay eggs at 25°C, and the resulting larvae were switched to 29°C 24 h AEL.

Wing imaginal discs were then dissected after either 5 days (in case of control) or 7 days (in case of

CIN) of induction at 29°C. Wing imaginal discs (40–50 in case of controls, and 30–40 in case of CIN)

were cleaned in ice-cold 1X PBS and were incubated in 10X trypsin-EDTA (Sigma-Aldrich) at room

temperature (RT) for 1 h, with regular nutation, and gentle pipetting. The single cell suspension was

then subjected to centrifugation at 2500*g* for 2.5 min. The supernatant was gently removed, and the

cells were stained with 1 ml of 1ug/ml of DAPI solution for 10 min at 4°C in low-light conditions, after

resuspending the cell pellet. The cells were immediately transferred to flow-tubes, and FACS sorting

was carried out on the different tissue subpopulations.

FACS sorting of different tissue subpopulations

Flow cytometry experiments were carried out using a FACS Aria Fusion sorter (Beckton Dickinson,

San Jose, California) equipped with a 4-laser configuration (405-488-561-640nm). Single cells were

gated according to scatter parameters, using FSC width to avoid doublets. Dead cells were excluded

using DAPI as a viability marker; only live single cells were analysed, and these were sorted on the

basis of their fluorescence pattern. For sorting different CIN subpopulations from CIN tissues,

P(Aneuploid) cells were distinguished as GFP+ RFP+ cells and P(CIN) cells as GFP- RFP+ cells,

and the remaining GFP- RFP- cells were labelled as A(Aneuploid) cells. Similarly, for sorting the two

subpopulations from the control tissues, RFP+ cells were attributed as P(control), while the RFP-

cells were attributed as A(control).   Fluorescence of GFP was detected using blue (488 nm) laser

excitation and collecting green fluorescence using a 530/30 BP filter. RFP fluorescence was obtained

using green-yellow laser excitation (561 nm) and collecting fluorescence using a 610/20 BP filter. A

dot plot combining GFP and RFP emission on single live cells was used to select them for sorting

into 1.5 ml Eppendorf tubes. A 70 µm nozzle was used to minimize the sorted volume into collection

vials. During the entire experiment, the cells were under 300 RPM agitation and were maintained at

4°C. For each subpopulation, 30000 cells were collected in RNase-free 1.5 ml microcentrifuge tubes

containing 10 µl of freshly prepared 2X Lysis Buffer following the methods from (*73*).

RNA purification, library preparation, and bulk RNA-sequencing

Total RNA from a total of 4 replicates of 5 distinct subpopulations was then purified at the IRB Barcelona Functional Genomics Core Facility, as described in (*73*). mRNA was isolated from total RNA using the NEBNext Poly(A) mRNA Magnetic Isolation Module Kit (New England Biolabs). NGS libraries were prepared from the purified mRNA using the NEBNext Ultra II RNA Library Prep Kit for Illumina (New England Biolabs). A total of 18 cycles of PCR amplification were applied to all libraries. The final libraries were quantified using the Qubit dsDNA HS assay (Invitrogen) and quality controlled with the Bioanalyzer 2100 DNA HS assay (Agilent). An equimolar pool was prepared with the 20 libraries and submitted for sequencing at the *Centro Nacional de Análisis Genómico* (CNAG). A final quality control by qPCR was performed by the sequencing provider before paired-end 150 nt sequencing on a NovaSeq6000 S4 (Illumina). More than 275 Gbp of reads were produced, with a minimum of 39 million paired-end reads per sample.

Data processing

Adapters from reads were removed using trim galore (v.0.6.7) (https://github.com/FelixKrueger/TrimGalore) with default parameters. Resulting reads were aligned to the dm6 *D. melanogaster* genome version using STAR (v.2.7.10a) (*74*). The count matrix was generated in R (https://www.R-project.org/) through the function "featureCounts" of the Rsubread package (*75*).

Differential expression

Differential expression was performed using the DESeq2 package (*76*). For plotting purposes and generating principal components, the count matrix was normalized using the "vst" function.

Gene set enrichment analysis

Gene set enrichment analysis was performed using the roastgsa package (*77*). GO and KEGG pathways were downloaded using the org.Dm.eg.db package.

Immunohistochemistry

Larvae were selected according to the day AEL mentioned in the individual figures, and wing imaginal discs were dissected in phosphate-buffered saline (PBS), fixed for 20 min in 4% formaldehyde in PBS, and stained with antibodies in PBS with 0.3% BSA, 0.2% Triton X-100. The antibodies and reagents used in this study are summarized in Table S4.

DNA synthesis

The Click-iT™ Plus EdU Alexa Fluor™ 647 Imaging Kit from Invitrogen (C10640) was used to measure DNA synthesis (S phase) in proliferating cells, following the manufacturer's instructions. EdU (5-ethynyl-2´-deoxyuridine) provided in the kit is a nucleoside analogue of thymidine and is incorporated into DNA during active DNA synthesis. Time of incubation with EdU: 15 min

TUNEL

Larvae were dissected in cold 1X PBS and fixed in 4% FA for 20 min. After fixation, three fast washes with PBT 0.3% Triton (PBS 1%, Triton X-100 0.3%) followed by four 15-min washes with PBT 0.3% Triton were performed. The samples were blocked with BBT (0.3% BSA, 0.3% Triton X-100) for 1 h rotating at RT and then incubated overnight with the primary antibody at 4℃. The next day, samples were subjected to three fast washes with PBT 0.3% Triton, and then they were washed four more times with PBT 0.3% Triton for 20 min each, followed by incubation with secondary antibodies diluted in BBT for 90 min rotating in the dark at RT. After incubation, the samples were subjected to three rounds of fast washes with PBT 0.3% Triton and four 20-min washes with PBT 0.3%. Samples were kept in BBT blocking overnight at 4ºC. The following day, samples were permeabilized with 495 μl of 100 mM Na-Citrate and 2.5 μl of 20% TritonX for 30 min at 65ºC and then subjected to three fast washes with PBT 0.3% Triton. They were then incubated twice for 5 min at RT with 50 μl of Equilibration buffer. Then 51 μl of TUNEL reaction mix (50 μl of TUNEL reaction buffer Component A, 1 μl TdT Enzyme, Biotium) was added and the samples were incubated for 2 h at RT. After incubation, the samples were subjected to three fast washes with PBT 0.3% Triton, followed by a 30-min incubation with DAPI at RT. Finally, three fast washes and three 10-min washes with PBT 0.3% Triton were performed. Stained tissues were kept in mounting media.

Confocal microscopy.

Samples were analysed using the following confocal microscopes: STELLARIS Confocal Platform with White Light Laser (Leica Microsystems), LSM 780 Zeiss, and LSM880 Zeiss fitted with a Fast Airyscan module (Carl Zeiss). Z-stacks were acquired using either the laser scanning confocal mode or the High-Resolution mode (Airyscan). Image acquisition was done with 20x, 25x, 40x, 63x, and 100x lenses. The most representative image is shown in all experiments.

Chloroquine treatment

Flies of desired genotypes were crossed and set up at 25°C and shifted to 29°C 8 h AEL. After 48 h at 29°C, a working concentration of 3 mg/ml of chloroquine (with water as controls) was added to the food. Larvae were processed 72 h thereafter. Experimental flies and control individuals were grown in parallel.

Tissue size, cell death and mitotic activity

The sizes of the whole wing disc (based on DAPI staining), the anterior (A) and posterior (P) compartments (based on Ci expression, in Figures 3-5) and the anterior (A) and posterior (P) areas of the wing pouch (labelled by Nubbin expression, in Figure 6 and 7) were measured manually using Fiji Software (NIH, USA) on apical sections (epithelium) of the wing disc obtained using a Zeiss LSM780 confocal microscope. A and P areas were normalized to the size of the respective control compartments. The left column of each graph represents the genotype which was considered as the control. The area of the signal to be quantified (TUNEL, cDcp-1, EdU, pH3, and mCh-Atg8) was set by manually adjusting the threshold and normalized to the area of the indicated larval territory (A or P compartments within the wing pouch) after making a maximum intensity Z-Projection.

Statistical analysis

Statistical analysis was performed by either One-way ANOVA when comparing more than two experimental groups or an unpaired equal-variance two-tailed Student's t-test when comparing the difference in means of a single experimental group with a control. Differences were considered significant when p values were less than 0.0001 (****), 0.001 (***), 0.01 (**), or 0.05 (*). All genotypes included in each histogram or scatter plot were subjected to the same experimental conditions (temperature and time of transgene induction) and analysed in parallel. Mean values and standard

549　deviations were calculated, and the corresponding statistical analysis and graphical representations

550　were carried out with GraphPad Prism 10.0 statistical software.

Data availability

552　Materials generated for the study are available from the corresponding author on request. Source

553　data are provided with this paper. The raw and processed RNAseq data have been deposited in the

554　GEO database under accession number GSE306996.

References

1.　B. Nguyen, C. Fong, A. Luthra, S. A. Smith, R. G. DiNatale, S. Nandakumar, H. Walch, W.
K. Chatila, R. Madupuri, R. Kundra, C. M. Bielski, B. Mastrogiacomo, M. T. A. Donoghue, A.
Boire, S. Chandarlapaty, K. Ganesh, J. J. Harding, C. A. Iacobuzio-Donahue, P. Razavi, E.
Reznik, C. M. Rudin, D. Zamarin, W. Abida, G. K. Abou-Alfa, C. Aghajanian, A. Cercek, P.
Chi, D. Feldman, A. L. Ho, G. Iyer, Y. Y. Janjigian, M. Morris, R. J. Motzer, E. M. O'Reilly, M.
A. Postow, N. P. Raj, G. J. Riely, M. E. Robson, J. E. Rosenberg, A. Safonov, A. N.
Shoushtari, W. Tap, M. Y. Teo, A. M. Varghese, M. Voss, R. Yaeger, M. G. Zauderer, N.
Abu-Rustum, J. Garcia-Aguilar, B. Bochner, A. Hakimi, W. R. Jarnagin, D. R. Jones, D.
Molena, L. Morris, E. Rios-Doria, P. Russo, S. Singer, V. E. Strong, D. Chakravarty, L. H.
Ellenson, A. Gopalan, J. S. Reis-Filho, B. Weigelt, M. Ladanyi, M. Gonen, S. P. Shah, J.
Massague, J. Gao, A. Zehir, M. F. Berger, D. B. Solit, S. F. Bakhoum, F. Sanchez-Vega, N.
Schultz, Genomic characterization of metastatic patterns from prospective clinical
sequencing of 25,000 patients. *Cell.* **185**, 563-575.e11 (2022).

2.　A. J. X. Lee, D. Endesfelder, A. J. Rowan, A. Walther, N. J. Birkbak, P. A. Futreal, J.
Downward, Z. Szallasi, I. P. M. Tomlinson, M. Howell, M. Kschischo, C. Swanton,
Chromosomal Instability Confers Intrinsic Multidrug Resistance. *Cancer Res.* **71**, 1858–1870
(2011).

3.　E. van Dijk, T. van den Bosch, K. J. Lenos, K. El Makrini, L. E. Nijman, H. F. B. van Essen,
N. Lansu, M. Boekhout, J. H. Hageman, R. C. Fitzgerald, C. J. A. Punt, J. B. Tuynman, H. J.
G. Snippert, G. J. P. L. Kops, J. P. Medema, B. Ylstra, L. Vermeulen, D. M. Miedema,
Chromosomal copy number heterogeneity predicts survival rates across cancers. *Nat.*
Commun. **12**, 3188 (2021).

4.　D. A. Lukow, E. L. Sausville, P. Suri, N. K. Chunduri, A. Wieland, J. Leu, J. C. Smith, V.
Girish, A. A. Kumar, J. Kendall, Z. Wang, Z. Storchova, J. M. Sheltzer, Chromosomal
instability accelerates the evolution of resistance to anti-cancer therapies. *Dev. Cell.* **56**,
2427-2439.e4 (2021).

5.　M. R. Ippolito, V. Martis, S. Martin, A. E. Tijhuis, C. Hong, R. Wardenaar, M. Dumont, J.
Zerbib, D. C. J. J. Spierings, D. Fachinetti, U. Ben-David, F. Foijer, S. Santaguida, Gene
copy-number changes and chromosomal instability induced by aneuploidy confer resistance
to chemotherapy. *Dev. Cell.* **56**, 2440-2454.e6 (2021).

6.　T. Davoli, A. W. Xu, K. E. Mengwasser, L. M. Sack, J. C. Yoon, P. J. Park, S. J. Elledge,
Cumulative Haploinsufficiency and Triplosensitivity Drive Aneuploidy Patterns and Shape
the Cancer Genome. *Cell.* **155**, 948–62 (2013).

7.　P. J. Stephens, C. D. Greenman, B. Fu, F. Yang, G. R. Bignell, L. J. Mudie, E. D. Pleasance,
K. W. Lau, D. Beare, L. A. Stebbings, S. McLaren, M.-L. Lin, D. J. McBride, I. Varela, S. Nik-
Zainal, C. Leroy, M. Jia, A. Menzies, A. P. Butler, J. W. Teague, M. A. Quail, J. Burton, H.
Swerdlow, N. P. Carter, L. A. Morsberger, C. Iacobuzio-Donahue, G. A. Follows, A. R.
Green, A. M. Flanagan, M. R. Stratton, P. A. Futreal, P. J. Campbell, Massive Genomic

Rearrangement Acquired in a Single Catastrophic Event during Cancer Development. *Cell*.
144, 27–40 (2011).

8. J. Li, M. J. Hubisz, E. M. Earlie, M. A. Duran, C. Hong, A. A. Varela, E. Lettera, M. Deyell, B.
Tavora, J. J. Havel, S. M. Phyu, A. D. Amin, K. Budre, E. Kamiya, J.-A. Cavallo, C. Garris, S.
Powell, J. S. Reis-Filho, H. Wen, S. Bettigole, A. J. Khan, B. Izar, E. E. Parkes, A. M.
Laughney, S. F. Bakhoum, Non-cell-autonomous cancer progression from chromosomal
instability. *Nature*. **620**, 1080–1088 (2023).

9. J. Li, M. A. Duran, N. Dhanota, W. K. Chatila, S. E. Bettigole, J. Kwon, R. K. Sriram, M. P.
Humphries, M. Salto-Tellez, J. A. James, M. G. Hanna, J. C. Melms, S. Vallabhaneni, K.
Litchfield, I. Usaite, D. Biswas, R. Bareja, H. W. Li, M. L. Martin, P. Dorsaint, J.-A. Cavallo,
P. Li, C. Pauli, L. Gottesdiener, B. J. DiPardo, T. J. Hollmann, T. Merghoub, H. Y. Wen, J. S.
Reis-Filho, N. Riaz, S.-S. M. Su, A. Kalbasi, N. Vasan, S. N. Powell, J. D. Wolchok, O.
Elemento, C. Swanton, A. N. Shoushtari, E. E. Parkes, B. Izar, S. F. Bakhoum, Metastasis
and Immune Evasion from Extracellular cGAMP Hydrolysis. *Cancer Discov.* **11**, 1212–1227
(2021).

10. S. F. Bakhoum, B. Ngo, A. M. Laughney, J.-A. Cavallo, C. J. Murphy, P. Ly, P. Shah, R. K.
Sriram, T. B. K. Watkins, N. K. Taunk, M. Duran, C. Pauli, C. Shaw, K. Chadalavada, V. K.
Rajasekhar, G. Genovese, S. Venkatesan, N. J. Birkbak, N. McGranahan, M. Lundquist, Q.
LaPlant, J. H. Healey, O. Elemento, C. H. Chung, N. Y. Lee, M. Imielenski, G. Nanjangud, D.
Pe'er, D. W. Cleveland, S. N. Powell, J. Lammerding, C. Swanton, L. C. Cantley,
Chromosomal instability drives metastasis through a cytosolic DNA response. *Nature*. **553**,
467–472 (2018).

11. A. D. Silk, L. M. Zasadil, A. J. Holland, B. Vitre, D. W. Cleveland, B. A. Weaver,
Chromosome missegregation rate predicts whether aneuploidy will promote or suppress
tumors. *Proc Natl Acad Sci U S A*. **110**, E4134-41 (2013).

12. K. M. Godek, M. Venere, Q. Wu, K. D. Mills, W. F. Hickey, J. N. Rich, D. A. Compton,
Chromosomal Instability Affects the Tumorigenicity of Glioblastoma Tumor-Initiating Cells.
Cancer Discov. **6**, 532–545 (2016).

13. B. a a Weaver, A. D. Silk, C. Montagna, P. Verdier-Pinard, D. W. Cleveland, Aneuploidy acts
both oncogenically and as a tumor suppressor. *Cancer Cell*. **11**, 25–36 (2007).

14. J. Monserrat, C. Morales Torres, L. Richardson, T. S. Wilson, H. Patel, M.-C. Domart, S.
Horswell, O.-R. Song, M. Jiang, M. Crawford, M. Bui, Y. Dalal, P. Scaffidi, Disruption of the
MSL complex inhibits tumour maintenance by exacerbating chromosomal instability. *Nat.*
Cell Biol. **23**, 401–412 (2021).

15. S. L. Thompson, D. A. Compton, Proliferation of aneuploid human cells is limited by a p53-
dependent mechanism. *J Cell Biol*. **188**, 369–381 (2010).

16. S. Santaguida, A. Amon, Short- and long-term effects of chromosome mis-segregation and
aneuploidy. *Nat Rev Mol Cell Biol*. **16**, 473–485 (2015).

17. J. Zhu, H.-J. Tsai, M. R. Gordon, R. Li, Cellular Stress Associated with Aneuploidy. *Dev.*
Cell. **44**, 420–431 (2018).

18. R. Li, J. Zhu, Effects of aneuploidy on cell behaviour and function. *Nat. Rev. Mol. Cell Biol.*
(2022), doi:10.1038/s41580-021-00436-9.

19. J. Joy, L. Barrio, C. Santos-Tapia, D. Romão, N. N. Giakoumakis, M. Clemente-Ruiz, M.
Milán, Proteostasis failure and mitochondrial dysfunction leads to aneuploidy-induced
senescence. *Dev. Cell*. **56**, 2043-2058.e7 (2021).

20. M. Barroso-Vilares, J. C. Macedo, M. Reis, J. D. Warren, D. Compton, E. Logarinho, Small-
molecule inhibition of aging-associated chromosomal instability delays cellular senescence.
EMBO Rep. **21**, e49248 (2020).

21. J. C. Macedo, S. Vaz, B. Bakker, R. Ribeiro, P. L. Bakker, J. M. Escandell, M. G. Ferreira, R.
Medema, F. Foijer, E. Logarinho, FoxM1 repression during human aging leads to mitotic
decline and aneuploidy-driven full senescence. *Nat. Commun.* **9**, 2834 (2018).

22. S. Santaguida, A. Richardson, D. R. Iyer, O. M'Saad, L. Zasadil, K. A. Knouse, Y. L. Wong,
N. Rhind, A. Desai, A. Amon, Chromosome Mis-segregation Generates Cell-Cycle-Arrested
Cells with Complex Karyotypes that Are Eliminated by the Immune System. *Dev Cell.* **41**,
638-651 e5 (2017).

23. R. W. Wang, S. Viganò, U. Ben-David, A. Amon, S. Santaguida, Aneuploid senescent cells
activate NF-κB to promote their immune clearance by NK cells. *EMBO Rep.* **22** (2021),
doi:10.15252/embr.202052032.

24. A. Dekanty, L. Barrio, M. Muzzopappa, H. Auer, M. Milan, Aneuploidy-induced delaminating
cells drive tumorigenesis in Drosophila epithelia. *Proc. Natl. Acad. Sci.* **109**, 20549–20554
(2012).

25. M. Clemente-Ruiz, J. M. Murillo-Maldonado, N. Benhra, L. Barrio, L. Perez, G. Quiroga, A.
R. Nebreda, M. Milan, L. Pérez, G. Quiroga, A. R. Nebreda, M. Milán, Gene Dosage
Imbalance Contributes to Chromosomal Instability-Induced Tumorigenesis. *Dev Cell.* **36**,
290–302 (2016).

26. S. Morais da Silva, T. Moutinho-Santos, C. E. Sunkel, A tumor suppressor role of the Bub3
spindle checkpoint protein after apoptosis inhibition. *J Cell Biol.* **201**, 385–393 (2013).

27. M. Muzzopappa, L. Murcia, M. Milan, M. Milán, M. Milan, Feedback amplification loop drives
malignant growth in epithelial tissues. *Proc Natl Acad Sci U S A.* **114**, E7291–E7300 (2017).

28. N. Benhra, L. Barrio, M. Muzzopappa, M. Milán, Chromosomal Instability Induces Cellular
Invasion in Epithelial Tissues. *Dev. Cell.* **47**, 161-174.e4 (2018).

29. E. Gracia-Latorre, L. Pérez, M. Muzzopappa, M. Milán, A single WNT enhancer drives
specification and regeneration of the Drosophila wing. *Nat. Commun.* **13**, 4794 (2022).

30. D. Romão, M. Muzzopappa, L. Barrio, M. Milán, The Upd3 cytokine couples inflammation to
maturation defects in Drosophila. *Curr. Biol.* **31**, 1780-1787.e6 (2021).

31. L. Barrio, A.-E. Gaspar, M. Muzzopappa, K. Ghosh, D. Romao, M. Clemente-Ruiz, M. Milán,
Chromosomal instability-induced cell invasion through caspase-driven DNA damage. *Curr.*
Biol. **33**, 4446-4457.e5 (2023).

32. A. Garelli, A. M. Gontijo, V. Miguela, E. Caparros, M. Dominguez, Imaginal Discs Secrete
Insulin-Like Peptide 8 to Mediate Plasticity of Growth and Maturation. *Science (80-.).* **336**,
579–582 (2012).

33. J. Colombani, D. S. D. S. Andersen, P. Leopold, P. Léopold, Secreted peptide Dilp8
coordinates Drosophila tissue growth with developmental timing. *Science.* **336**, 582–5
(2012).

34. Y. Kwon, W. Song, I. A. Droujinine, Y. Hu, J. M. Asara, N. Perrimon, Systemic Organ
Wasting Induced by Localized Expression of the Secreted Insulin/IGF Antagonist ImpL2.
Dev Cell. **33**, 36–46 (2015).

35. R. Nagata, N. Akai, S. Kondo, K. Saito, S. Ohsawa, T. Igaki, Yorkie drives supercompetition
by non-autonomous induction of autophagy via bantam microRNA in Drosophila. *Curr. Biol.*
32, 1064-1076.e4 (2022).

36. E. Moreno, K. Basler, dMyc transforms cells into super-competitors. *Cell.* **117**, 117–129
(2004).

37. E. Moreno, L. Valon, F. Levillayer, R. Levayer, Competition for Space Induces Cell
Elimination through Compaction-Driven ERK Downregulation. *Curr. Biol.* **29**, 23-34.e8
(2019).

38.   B. A. Hay, T. Wolff, G. M. Rubin, Expression of baculovirus P35 prevents cell death in
Drosophila. *Development.* **120**, 2121–2129 (1994).

39.   M. Uhlirova, D. Bohmann, JNK- and Fos-regulated Mmp1 expression cooperates with Ras to
induce invasive tumors in Drosophila. *EMBO J.* **25**, 5294–304 (2006).

40.   W. Song, S. Kir, S. Hong, Y. Hu, X. Wang, R. Binari, H.-W. Tang, V. Chung, A. S. Banks, B.
Spiegelman, N. Perrimon, Tumor-Derived Ligands Trigger Tumor Growth and Host Wasting
via Differential MEK Activation. *Dev. Cell.* **48**, 277-286.e6 (2019).

41.   S. Schott, A. Ambrosini, A. Barbaste, C. Benassayag, M. Gracia, A. Proag, M. Rayer, B.
Monier, M. Suzanne, A fluorescent toolkit for spatiotemporal tracking of apoptotic cells in
living Drosophila tissues. *Development.* **144**, 3840–3846 (2017).

42.   M. Das Thakur, Y. Feng, R. Jagannathan, M. J. Seppa, J. B. Skeath, G. D. Longmore, Ajuba
LIM Proteins Are Negative Regulators of the Hippo Signaling Pathway. *Curr. Biol.* **20**, 657–
662 (2010).

43.   P. Meier, J. Silke, S. J. Leevers, G. I. Evan, The Drosophila caspase DRONC is regulated by
DIAP1. *EMBO J.* **19**, 598–611 (2000).

44.   B. J. Thompson, S. M. Cohen, The Hippo pathway regulates the bantam microRNA to
control cell proliferation and apoptosis in Drosophila. *Cell.* **126**, 767–774 (2006).

45.   R. Nolo, C. M. Morrison, C. Tao, X. Zhang, G. Halder, The bantam microRNA is a target of
the hippo tumor-suppressor pathway. *Curr Biol.* **16**, 1895–1904 (2006).

46.   F. Hamaratoglu, M. Willecke, M. Kango-Singh, R. Nolo, E. Hyun, C. Tao, H. Jafar-Nejad, G.
Halder, The tumour-suppressor genes NF2/Merlin and Expanded act through Hippo
signalling to regulate cell proliferation and apoptosis. *Nat Cell Biol.* **8**, 27–36 (2006).

47.   G. Sun, K. D. Irvine, Ajuba family proteins link JNK to Hippo signaling. *Sci Signal.* **6**, ra81
(2013).

48.   H. K. Sluss, Z. Han, T. Barrett, D. C. Goberdhan, C. Wilson, R. J. Davis, Y. T. Ip, A JNK
signal transduction pathway that mediates morphogenesis and an immune response in
Drosophila. *Genes Dev.* **10**, 2745–2758 (1996).

49.   W. Kowalczyk, L. Romanelli, M. Atkins, H. Hillen, C. Bravo González-Blas, J. Jacobs, J. Xie,
S. Soheily, E. Verboven, I. M. Moya, S. Verhulst, M. de Waegeneer, L. Sansores-Garcia, L.
van Huffel, R. L. Johnson, L. A. van Grunsven, S. Aerts, G. Halder, Hippo signaling instructs
ectopic but not normal organ growth. *Science (80-.).* **378** (2022),
doi:10.1126/science.abg3679.

50.   J. A. Sanchez, D. Mesquita, M. C. Ingaramo, F. Ariel, M. Milán, A. Dekanty, Eiger/TNFα-
mediated Dilp8 and ROS production coordinate intra-organ growth in Drosophila. *PLOS
Genet.* **15**, e1008133 (2019).

51.   L. Boulan, D. Andersen, J. Colombani, E. Boone, P. Léopold, Inter-Organ Growth
Coordination Is Mediated by the Xrp1-Dilp8 Axis in Drosophila. *Dev. Cell.* **49**, 811-818.e4
(2019).

52.   N. Arquier, C. Géminard, M. Bourouis, G. Jarretou, B. Honegger, A. Paix, P. Léopold,
Drosophila ALS regulates growth and metabolism through functional interaction with insulin-
like peptides. *Cell Metab.* **7**, 333–8 (2008).

53.   B. Honegger, M. Galic, K. Köhler, F. Wittwer, W. Brogiolo, E. Hafen, H. Stocker, Imp-L2, a
putative homolog of vertebrate IGF-binding protein 7, counteracts insulin signaling in
Drosophila and is essential for starvation resistance. *J. Biol.* **7**, 10 (2008).

54.   J. A. Sanchez, D. Mesquita, M. C. Ingaramo, F. Ariel, M. Milán, A. Dekanty, Eiger/TNFα-
mediated Dilp8 and ROS production coordinate intra-organ growth in Drosophila. *bioRxiv*
(2019), , doi:10.1101/606707.

55. S. E. Siegrist, N. S. Haque, C. H. Chen, B. A. Hay, I. K. Hariharan, Inactivation of both Foxo
and reaper promotes long-term adult neurogenesis in Drosophila. *Curr Biol*. **20**, 643–648
(2010).

56. D. S. Andersen, J. Colombani, V. Palmerini, K. Chakrabandhu, E. Boone, M. Rothlisberger,
J. Toggweiler, K. Basler, M. Mapelli, A. O. Hueber, P. Leopold, The Drosophila TNF receptor
Grindelwald couples loss of cell polarity and neoplastic growth. *Nature*. **522**, 482–486
(2015).

57. E. A. Bach, L. A. Ekas, A. Ayala-Camargo, M. S. Flaherty, H. Lee, N. Perrimon, G. H. Baeg,
GFP reporters detect the activation of the Drosophila JAK/STAT pathway in vivo. *Gene Expr*
Patterns. **7**, 323–331 (2007).

58. C. Recasens-Alvarez, A. Ferreira, M. Milán, M. Milan, JAK/STAT controls organ size and
fate specification by regulating morphogen production and signalling. *Nat Commun*. **8**,
13815 (2017).

59. H. Kanda, T. Igaki, H. Okano, M. Miura, Conserved metabolic energy production pathways
govern Eiger/TNF-induced nonapoptotic cell death. *Proc Natl Acad Sci U S A*. **108**, 18977–
18982 (2011).

60. V. Gorgoulis, P. D. Adams, A. Alimonti, D. C. Bennett, O. Bischof, C. Bishop, J. Campisi, M.
Collado, K. Evangelou, G. Ferbeyre, J. Gil, E. Hara, V. Krizhanovsky, D. Jurk, A. B. Maier,
M. Narita, L. Niedernhofer, J. F. Passos, P. D. Robbins, C. A. Schmitt, J. Sedivy, K. Vougas,
T. von Zglinicki, D. Zhou, M. Serrano, M. Demaria, Cellular Senescence: Defining a Path
Forward. *Cell*. **179**, 813–827 (2019).

61. K. Rowald, M. Mantovan, J. Passos, C. Buccitelli, B. R. Mardin, J. O. Korbel, M. Jechlinger,
R. Sotillo, Negative Selection and Chromosome Instability Induced by Mad2 Overexpression
Delay Breast Cancer but Facilitate Oncogene-Independent Outgrowth. *Cell Rep.* **15**, 2679–
2691 (2016).

62. M. S. Levine, B. Bakker, B. Boeckx, J. Moyett, J. Lu, B. Vitre, D. C. Spierings, P. M.
Lansdorp, D. W. Cleveland, D. Lambrechts, F. Foijer, A. J. Holland, Centrosome
Amplification Is Sufficient to Promote Spontaneous Tumorigenesis in Mammals. *Dev Cell*.
40, 313-322 e5 (2017).

63. M. Clemente-Ruiz, M. Muzzopappa, M. Milán, M. Milan, Tumor suppressor roles of CENP-E
and Nsl1 in Drosophila epithelial tissues. *Cell Cycle*. **13**, 1450–1455 (2014).

64. E. Fusari, M. Muzzopappa, J. Gracia, M. Milán, Depletion of aneuploid cells is shaped by
cell-to-cell interactions. *Cell Genomics*. **5**, 100894 (2025).

65. S. Stingele, G. Stoehr, Z. Storchova, Activation of autophagy in cells with abnormal
karyotype. *Autophagy*. **9**, 246–248 (2013).

66. S. Santaguida, E. Vasile, E. White, A. Amon, Aneuploidy-induced cellular stresses limit
autophagic degradation. *Genes Dev*. **29**, 2010–2021 (2015).

67. B. Wang, J. Han, J. H. Elisseeff, M. Demaria, The senescence-associated secretory
phenotype and its physiological and pathological implications. *Nat. Rev. Mol. Cell Biol.* **25**,
958–978 (2024).

68. N. Martin, N. Popgeorgiev, G. Ichim, D. Bernard, BCL-2 proteins in senescence: beyond a
simple target for senolysis? *Nat. Rev. Mol. Cell Biol.* **24**, 517–518 (2023).

69. C. Rauskolb, S. Sun, G. Sun, Y. Pan, K. D. Irvine, Cytoskeletal Tension Inhibits Hippo
Signaling through an Ajuba-Warts Complex. *Cell*. **158**, 143–156 (2014).

70. J. Brennecke, D. R. Hipfner, A. Stark, R. B. Russell, S. M. Cohen, bantam encodes a
developmentally regulated microRNA that controls cell proliferation and regulates the
proapoptotic gene hid in Drosophila. *Cell*. **113**, 25–36 (2003).

71. A. Figueroa-Clarevega, D. Bilder, Malignant Drosophila Tumors Interrupt Insulin Signaling to
Induce Cachexia-like Wasting. *Dev Cell*. **33**, 47–55 (2015).

72. N. S. Katheder, R. Khezri, F. O'Farrell, S. W. Schultz, A. Jain, M. M. Rahman, K. O. Schink,
T. A. Theodossiou, T. Johansen, G. Juhasz, D. Bilder, A. Brech, H. Stenmark, T. E. Rusten,
Microenvironmental autophagy promotes tumour growth. *Nature*. **541**, 417–420 (2017).

73. E. Gonzalez-Roca, X. Garcia-Albéniz, S. Rodriguez-Mulero, R. R. Gomis, K. Kornacker, H.
Auer, Accurate Expression Profiling of Very Small Cell Populations. *PLoS One*. **5**, e14418
(2010).

74. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson,
T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. **29**, 15–21
(2013).

75. Y. Liao, G. K. Smyth, W. Shi, The R package Rsubread is easier, faster, cheaper and better
for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **47**, e47–e47
(2019).

76. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

77. A. Caballé-Mestres, A. Berenguer-Llergo, C. Stephan-Otto Attolini, Roastgsa: a comparison
of rotation-based scores for gene set enrichment analysis. *BMC Bioinformatics*. **24**, 408
(2023).

Acknowledgments

We thank E. Bach, J. Castelli-Gair, N. Tapon, M. Suzanne, G. Juhasz, the Bloomington *Drosophila*

Stock Center (USA), the Vienna *Drosophila* Resource Center (Austria), and the Developmental Studies

Hybridoma Bank (USA) for flies and antibodies, and IRB Barcelona's Advanced Digital Microscopy,

Functional Genomics, and Biostatistics and Bioinformatics Facilities for help. This work was funded by

the PID2019-110082GB-I00 and PID2022-137673NB-I00 grants from the Spanish Ministry of Science,

Innovation and Universities (MICIU) and the ERDF "Una manera de hacer Europa", and the European

Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant

agreement number 945352. We gratefully acknowledge institutional funding from the MICIU through the

Centres of Excellence Severo Ochoa Award, and from the CERCA Programme of the Catalan

Government.

Author Contributions

All the authors conceived and designed the experiments and analysed the data. K.G. and A.K.

performed the experiments, M.M. and K.G. prepared the final figures, and M. M. supervised the whole

project and wrote the paper.

Declaration of interests

The authors declare no competing interests.

(A, B) Control wing discs (**A**) and wing discs subjected to CIN (**B**), stained for DAPI (blue or black),

Ci (red, to label the A compartment), and MMP1 (green, to label the aneuploid cells) and expressing

the indicated transgenes in the P compartment under the control of the *en-gal4* driver. Scale bars,

50 μm**.** Days of dissection after egg laying are as indicated. **(C)** Principal Component (PC) analysis

showing the transcriptomic differences between the 5 different populations: A (Red) and P (Blue)

cells of control wing discs, and aneuploid (green), A (orange) and P (gray) cells of wing discs

subjected to CIN. **(D)** Venn diagram representing the number of genes up- or down-regulated

uniquely in aneuploid cells (green), and proliferating cells subjected to CIN cells or in both types of

cells (gray). **(E, G)** Bubble plots representing Gene Ontology (GO) enrichment analyses of aneuploid

(**E**) and aneuploid and proliferating cells (**G**) when compared to P cells of control discs. Size of the

bubble represents the number of measured genes, and the colour scale represents the associated

p-value. **(F)** Volcano plot representing all the genes that are differentially up- (red dots, FC>2 and p-

value<0.05) or down-regulated (blue dots FC<-2 and p-value<0.05), or not differentially expressed

(gray dots, p-value>0.05 or 2>FC>-2) in aneuploid cells when compared to P cells of control discs.

See also Figure S1, S2, and Table S1.

Transcriptomic profile of ANEUPLOID cells (unique features)

Figure 2. The transcriptome of aneuploidy-induced senescent cells points to a heightened secretory potential

(A, B) Volcano plot representing GO class-specific genes that are differentially upregulated (red area, FC>2 and p-value<0.05), downregulated (blue area, FC<-2 and p-value<0.05), or not differentially

expressed (white area, p-value>0.05 or 2>FC>-2) in aneuploid cells when compared to P cells of

control discs. GO classes (A): Autophagy - GO0006914, Lysosome - GO0005764, Cytoskeleton -

GO0005856, Secretion - GO0046903, Secretome - Custom Gene Set. GO classes (B): G1/S Transition

- GO0000082, G2/M Transition - GO0000086, Regulation of cell cycle - GO0051726, and DNA

Replication - GO0006260. Genes corresponding to each of the GO classes have a distinct colour code.

(C) High magnification images of wing discs expressing the indicated transgenes under the control of

the *en-gal4* driver and stained to visualize differential expression of fluorescent reporters (green, red,

or white) and DAPI (blue and white). Apical and basal sections are shown. Aneuploid and CIN cells are

marked by a dashed line. Scale bars, 10 µm. See also Tables S1 and S2.

Figure 3. The transcriptome of aneuploidy-induced senescent cells points to activation of the

apoptotic and Yorkie pathways

(A) Volcano plot representing selected genes in 'Cell death - GO0008219' (left), and 'hpo-yki signaling

- GO0035329' (right) classes that are differentially upregulated (red area, FC>1.3 and p-value<0.05),

downregulated (blue area, FC<-1.3 and p-value<0.05), or not differentially expressed (white area, p-

value>0.05 or 1.3>FC>-1.3) in aneuploid cells when compared to P cells of control discs. (**B-F**)

Control wing discs (first column) and wing discs subjected to CIN (remaining columns), stained to

visualize GC3Ai or *hid-lacZ* (green, **B**), Jub-GFP (green, **C**), *bantam-lacZ* (green, **D**), DIAP1-GFP

(green, **E**), Yorkie-GFP (green, **F**), DAPI (white, **B-F**), MMP1 (red, **F**), and expressing the indicated

transgenes in the P compartment under the control of the *en-gal4* driver. In **C**, aneuploidy-induced

senescent cells are marked by a yellow line. Scale bars, 50 µm. Discs were dissected 5 days AEL.

The boundary between A and P cells is labelled by a yellow line. Note in **D** and **E,** a reduction in the

expression levels of *bantam-lacZ* and *DIAP1-GFP* upon expression of a dominant negative form of

JNK (Bsk-DN). (**D'**, **E'**) Histograms plotting the P/A signal intensity ratio of *bantam-lacZ* (**D'**) and

DIAP1-GFP (**E'**) of wing discs expressing the indicated transgenes in the P compartment under the

control of the *en-gal4* driver. Ordinary one-way ANOVA was performed to analyse the data. $*p<0.05$,

$**p<0.01$, $***p<0.001$, $****$ $p<0.0001$, ns, not significant. See also Table S3. See also Figure S3, and

Tables S1 and S3.

Figure 4. Yorkie plays a pro-survival role in CIN tissues

(**A, C, E, F, G, H**) Control wing discs (first columns in **A**, **C**) and wing discs subjected to CIN (remaining columns), stained to visualize TUNEL (**A**, white), cDcp-1 (**C**, **E**, white, **G**, red), *bantam-lacZ* (red, **H**), Ci (green, **A**, **C**, **E**, **F**), Nubbin (red, **A**, **C**), MMP1 (red, **E**, green, **G**, **H**), and DAPI (white, **A, C, E, F, G**) expressing the indicated transgenes in the P compartment under the control of the *en-gal4* driver.

Scale bars, 50 µm, unless indicated otherwise**.** Discs were dissected at the indicated days of

development. The boundary between A and P cells is labelled by a green line, and the wing pouch

by a red dotted line. In **G**, **H**, high magnification images of the basal sections of CIN tissues, where

aneuploidy-induced senescent cells are found and marked by a yellow line. Red arrows in **G** point to

dying cells. (**B**, **D**, **I**) Histograms plotting the amount of TUNEL (**B**, left), MMP1 (**I,** right) and cDcp-1

signal (**D, and I,** left) in the P wing pouch compartment (normalized to the size of the corresponding

compartment), the normalized size of the P wing pouch compartment (**B**, **D**, right), and size ratio

between the P and A wing pouch compartments (**B, D and I,** centre) of wing discs expressing the

indicated transgenes in the P compartment under the control of the *en-gal4* driver. Ordinary one-way

ANOVA was performed to analyse the data plotted in **B, D, I;** * $p<0.05$, ** $p<0.01$, *** $p<0.001$, ****

$p<0.0001$, ns, not significant. See also Table S3.

Figure 5. A non-autonomous effect of CIN tumours on the proliferation, growth, and survival of the neighbouring tissue

(**A**, **D**, **E**) Control wing discs (first column) and wing discs subjected to CIN (remaining columns), and expressing p35 in the P compartment under the control of the *en-gal4* driver, stained for Ci (green, to label the A compartment), DAPI (white), Nubbin (red, to label the wing pouch cells, **A**, **D**), TUNEL (white, **A**), pH3 (white, **D**), and EdU (white, **E**). Scale bars, 50 μm. Days of dissection after egg laying

 are as indicated. The boundary between A and P cells is labelled by a green line. The tissue subjected

to CIN is marked in **A**. High magnification of the squared regions is shown in the lower panels in **D**, **E**.

Note the non-autonomous induction of cell death (**A**) and block of cell proliferation (**D**, **E**) in A cells. (**B**,

C) Histograms plotting in **B** the normalized size of the A and P compartments and the size ratio between

these compartments, and in **C** the amount of TUNEL, pH3, and EdU incorporation in the A wing pouch

compartment (normalized to the size of the corresponding compartment) of wing discs expressing the

indicated transgenes in the P compartment under the control of the *en-gal4* driver. Ordinary one-way

ANOVA was performed to analyse the data plotted in **B** and **C**. *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****

$p<0.0001$; ns, not significant. See also Figure S4 and Table S3.

Figure 6. A growth suppressor role of Dilp8/Relaxin and ImpL2/IGFBP7 in CIN tumours.

(**A-C**, **F**) Control wing discs (first column) and wing discs subjected to CIN (remaining columns),

stained to visualize Dilp8-GFP (green, **A**), ImpL2-GFP (green, **A**), EdU (white, **B**), pH3 (white, **C**),

DAPI (white in **A-C**, blue in **F**), Nubbin (red, to label the wing pouch cells, **C**, **F**), and Ci (green, to

label the A compartment, **B**, **C**, **F**), and expressing the indicated transgenes in the P compartment

under the control of the *en-gal4* driver. Scale bars, 50 µm**.** Discs were dissected 5 days after egg

laying. The boundary between A and P cells is labelled by either a green line or a yellow dashed line.

High magnifications of the squared regions are shown in the lower panels in **B**, **C**. (**B', C', D, E**)

Histograms plotting the amount of EdU incorporation and pH3 in the A wing pouch compartment

(normalized to the size of the corresponding compartment, **B'**, **C'**), and the normalized size of the A

(**D**) and P (**E**) wing pouch compartments of wing discs expressing the indicated transgenes in the P

compartment under the control of the *en-gal4* driver. Ordinary one-way ANOVA was performed to

analyse the data plotted in **B', C', D, E.** * $p < 0.05$; **$p < 0.01$; ***$p < 0.001$; ****$p < 0.0001$; ns, not

significant. See also Figure S5 and Table S3.

Figure 7. A feed-forward loop involved in the growth of CIN tumours

(A) Volcano plot representing selected genes in JAK-STAT Signaling - GO0097696 class that are

differentially upregulated (red area, FC>1.3 and p-value<0.05), downregulated (blue area, FC<-1.3

and p-value<0.05), or not differentially expressed (white area, p-value>0.05 or 1.3>FC>-1.3) in

aneuploid cells when compared to P cells of control discs. (**B, C, D, G**) Control wing discs (first

column) and wing discs subjected to CIN (remaining columns) stained to visualize STAT-GFP (green,

B), *eiger-lacZ* (red, **C**), TUNEL (white, **D**), Nubbin (red, to label the wing pouch cells, **D, G**), and Ci

(green, to label the A compartment, **C, D, G**), and DAPI (white in **B**, **D**, blue in **C**, **G**) and expressing

the indicated transgenes in the P compartment under the control of the *en-gal4* driver. Scale bars, 50

μm**.** Discs were dissected 5 days AEL. The boundary between A and P cells is labelled by a green or

a yellow dashed line, or by Ci staining. High magnification of the squared regions are shown in the

lower panels in **B**. (**D', E, F**) Histograms plotting the amount of TUNEL (**D'**), and the size of the A (**E**)

and P (**F**) wing pouch compartments of wing discs expressing the indicated transgenes in the P

compartment under the control of the *en-gal4* driver. Ordinary one-way ANOVA was performed.

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, **** $p < 0.0001$, ns, not significant. (H) Model: in epithelial tissues

subjected to restricted induction of CIN in the P cell population (yellow), aneuploid cells (green)

delaminate basally and enter a senescent state. Members of the SASP impact on the neighbouring

wild type cell population (the A cell population depicted in purple) locally of systemically to induce cell

death and regulate cell proliferation and growth. Non-autonomous cell death feeds back to the tumour

to enhance its growth. Cartoon was generated with Biorender.com. See also Tables S1 and S3.

Symbol	Gene name	Vertebrate homolog	Significance (p.Value)	Fold Change
RYa	Ryamide	NPY family	3,25E-08	21721233,75
NLaz	Neural Lazarillo	ApoD and RBP4	1,07E-35	681,212
aspr	asperous	DNER and EDIL3	1,32E-55	491,381
Ilp8	Insulin-like peptide 8	Relaxin	5,46E-81	198,57
upd2	unpaired 2	Leptin	3,26E-49	110,948
nec	necrotic	SERPIN	1,02E-75	89,581
PGRP-SA	Peptidoglycan recognition protein SA	PGLYRP3	2,88E-59	88,703
Mmp1	Matrix metalloproteinase 1	MMP	2,30E-215	83,782
Pvf1	PDGF- and VEGF-related factor 1	PDGF and VEGF	4,28E-31	75,409
Tig	Tiggrin	RGD-containing integrin ligand	1,96E-26	73,827
Sp7	Serine protease 7	Serine Proteases	5,87E-85	65,623
upd3	unpaired 3	Il-6	3,28E-34	56,97
CG30280	Unannotated	MFAP4	8,16E-05	43,886
sona	sol narae	ADAMTS	2,08E-120	41,327
tnc	tenectin	Mucin-type protein with RGD motif	4,07E-17	37,48
scaf	scarface	Serine Protease	2,42E-124	34,526
Mmp2	Matrix metalloproteinase 2	MMP	3,28E-39	31,565
frm	farmer	FLG	1,72E-144	30,547
Wnt4	Wnt oncogene analog 4	WNT	1,01E-14	28,179
Idgf2	Imaginal disc growth factor 2	CHI3L1, CHI3L2, and OVGP1	5,07E-07	26,81
daw	dawdle	NODAL	1,84E-14	26,357
Spn47C	Serpin 47C	SERPIN	4,32E-71	23,661
Pvf2	PDGF- and VEGF-related factor 2	PDGF and VEGF	5,62E-99	21,107
ths	thisbe	FGF8	1,46E-32	18,374
CG31869	Unannotated	KCP	2,31E-86	18,285
upd1	unpaired 1	Il-6	1,01E-26	18,278
mtg	mind the gap	Lectin	3,46E-90	13,673
ImpL2	Ecdysone-inducible gene L2	IGFBP7	1,43E-76	12,527
Pvf3	PDGF- and VEGF-related factor 3	PDGF and VEGF	1,21E-92	12,395
CecB	Cecropin B	Similar to Antimicrobial peptides	1,15E-03	12,284
spz	spatzle	Neurotrophin	7,22E-09	11,326
CG9095	Unannotated	Secretory phospholipase A2	3,11E-19	10,982
egr	eiger	TNF	2,52E-22	10,773
fon	fondue	NCOA5 and NUP98	3,01E-08	9,755
l(1)G0193	orion	Similar to Chemokines	2,25E-56	8,84
Dro	Drosocin	Similar to Antimicrobial peptides	1,43E-03	8,007
Spn55B	Serpin 55B	SERPIN	1,94E-64	7,591

Wnt6	Wnt oncogene analog 6	WNT	1,32E-29	6,846
wg	wingless	WNT	5,12E-62	6,798
Cp15	Chorion protein 15	No homolog identified	4,48E-04	6,417

Table 1. The secretome of aneuploidy-induced senescent cells.

Table representing the 40 most differentially upregulated genes (comparing the transcriptome of the

aneuploidy-induced senescent cells with control cells) which are annotated as genes encoding for

secreted proteins, along with their vertebrate homolog, adjusted p. values and shrunken fold-

changes. The genes labelled in red are those which have been mentioned in the current work. A

manually curated custom gene set, consisting of 261 known or predicted secreted proteins as inferred

from FlyBase is provided in Table S2. The transcriptome of the aneuploidy-induced senescent cells

was compared to posterior wild-type cells.