1   ***Mycobacterium tuberculosis* growth arrest on propionate at acidic pH is suppressed by**

2   **mutations in *phoPR* and pyrazinamide treatment**

4   Heather M. Murdoch[1,2], Shelby J. Dechow[1,2], Bassel J. Abdalla[1] and Robert B. Abramovitch[1*.]

5
6   [1]Department of Microbiology, Genetics & Immunology, Michigan State University, East Lansing,
7   MI,48824
8
9   [2]Co-first authors contributed equally to this study.

Disclosures

*Corresponding Author:

Robert B. Abramovitch

E-mail: abramov5@msu.edu

Phone: (517) 884-5416

Fax: (517) 353-8957

Abstract

Mycobacterium tuberculosis (Mtb) arrests its growth at acidic pH, when grown on specific single carbon sources, including propionate. However, Mtb grows well on propionate at pH 7.0, supporting that propionate can support growth as a sole carbon source. To understand the basis of the propionate-driven growth arrest at acidic pH, we performed a forward genetic selection for mutants that enable growth on propionate at pH 5.7. All the selected mutants had insertions in the two-component regulatory genes *phoR* or *phoP*. We hypothesized that growth arrest at acidic pH is caused by PhoPR diverting carbon from central carbon metabolism towards lipid anabolism and that when PhoPR is inactivated, growth is promoted through metabolizing propionate by the methyl citrate cycle (MCC) into pyruvate, a permissive carbon source for growth at acidic pH. Using chemical inhibition and mutants of the MCC pathway, we demonstrate that the enhanced growth is dependent on the MCC. Furthermore, stimulating lipid synthesis via the methylmalonyl-CoA pathway by adding vitamin B12 restricts growth in the *ΔphoPR* mutant and, conversely, restricting lipid anabolism by inhibiting the triacylglycerol (TAG) synthase *tgs1* enhances growth of the *ΔphoPR* mutant. Notably, CoA pools increased in the *ΔphoPR* mutant grown on propionate, directly supporting our model. Given the role of CoA metabolism in pyrazinamide sensitivity, we examined Mtb sensitivity to pyrazinamide on propionate at acidic pH and, surprisingly, observed that pyrazinamide treatment of WT Mtb suppresses growth arrest on propionate at acidic pH. In contrast, the *phoPR* mutant has enhanced sensitivity to pyrazinamide. Together, these findings support that propionate-driven growth arrest at acidic pH is caused by metabolic remodeling that is regulated by PhoPR and is associated with pyrazinamide sensitivity.

52
53 **Importance**
54

When grown on certain single carbon sources, such as propionate, Mtb arrests its growth at acidic

pH and establishes a state of non-replicating persistence (NRP). To understand the genetic basis

of this growth restriction, a genetic selection was performed to identify mutants unable to arrest

growth at acidic pH with propionate as a sole carbon source. The selection exclusively identified

mutants in the PhoPR two-component regulatory system, which functions to modulate cell

envelope lipids and redox homeostasis through the upregulation of lipid synthesis at acidic pH.

Using genetic and chemical inhibition studies, we demonstrate that PhoPR arrests growth at

acidic pH by diverting carbon away from the methyl citrate cycle towards lipid anabolism.

Surprisingly, treatment of Mtb with pyrazinamide at acidic pH on propionate, also enabled growth.

Therefore, this study defines new mechanisms by which Mtb integrates environmental signaling

to regulate growth, metabolism, and drug susceptibility. These findings are relevant to

pathogenesis, as PhoPR is essential for growth in macrophages and animals, environments with

varying pH and carbon source availability, depending on immune pressures. These data suggest

that drug susceptibility may be impacted by enhanced growth and metabolic capacity of Mtb in

acidic and propionate-rich environments, such as the within the macrophage or the granuloma.

Introduction

Mycobacterium tuberculosis (Mtb) is a slow-growing pathogen that replicates within macrophages or extracellularly. Depending on the conditions, the bacterium may be colonizing environments with specific carbon sources and varying pH. The combinations of pH, carbon source availability, and other environmental cues can impact Mtb growth rate and virulence. During growth in vitro, or in vivo, Mtb has a slow doubling time of 20 hours to 70 days[1-3]. Under low oxygen, starvation, or mildly acidic pH on specific carbon sources, Mtb enters a non-replicating persistent (NRP) state that allows the pathogen to arrest its growth while maintaining its viability[4]. Slow or arrested growth in mycobacteria is thought to play a role in establishing tuberculosis-like diseases and can promote phenotypic drug tolerance and persistence[4-7]. Therefore, a mechanistic understanding of how Mtb regulates its growth rate, in response to specific environmental cues, is necessary for understanding pathogenesis and characterizing antibiotic susceptibility.

One of the key regulators of pH-driven adaptations in Mtb is the PhoPR two-component regulatory system (TCS). PhoPR induces the expression of virulence-related genes in response to acidic pH, magnesium, chloride and CO_2[8-11]. These virulence-related genes include the ESX-1 secretion system and cell envelope lipid synthesis genes such as pks2, pks3 and pks4 which generate sulfolipid (SL) and other acylated trehalose lipids [12, 13]. Notably, phoPR mutants are highly attenuated in mice and guinea pigs [14, 15], making PhoPR a target for vaccine development[16], supporting that PhoPR and adaptation to acidic pH or related cues are critical for Mtb pathogenesis.

Our lab previously discovered that Mtb has restricted growth on specific single carbon sources at acidic pH. Mtb grows well on carbon sources that function at the intersection of glycolysis and the TCA cycle (e.g. pyruvate, acetate, oleic acid and cholesterol)[17]. In contrast, on other tested carbon sources, such as glycerol or propionate, Mtb fully arrests its growth at acidic pH and establishes a state of NRP. We refer to this phenotype as acid growth arrest (17,

18). We hypothesized that Mtb has evolved means to restrict growth at acidic pH on specific

carbon sources to regulate growth and support pathogenesis. Indeed, we previously selected for

suppressor mutants of acid growth arrest in *ppe51* that enabled Mtb growth on glycerol at acidic

pH, increased its replication in infected activated macrophages, and resulted in decreased

fitness(18, 19). The goal of this study is to define the mechanisms of acid growth arrest on

propionate as a sole carbon source.

Significant research has been conducted examining propionyl-CoA metabolism in Mtb,

given it is a catabolic product of cholesterol, a carbon source consumed by Mtb in the host(20).

Propionyl-CoA is toxic to Mtb and is detoxified by the methyl citrate cycle (MCC) or by

incorporation into methyl branched lipids (such as phthiocerol dimycocerosates (PDIM), SL and

acylated trehaloses) (21, 22).   The MCC is composed of the methyl-isocitrate lyases *icl1/icl2*

along with *prpC* and *prpD* (22). *icl1/icl2, prpC* and *prpD* are essential for growth in minimal media

supplemented with propionate (22-25). In the presence of vitamin B12, Mtb is able to survive in

the absence of the MCC, by promoting metabolism via methylmalonyl-CoA pathway, enabling the

sequestration of propionyl-CoA into methyl branched lipids such as PDIM (21, 22).

To define the mechanism of propionate-driven acid growth arrest, we conducted a forward

genetic selection for transposon (Tn) mutants that suppress acid growth arrest on propionate at

acidic pH. The selection exclusively identified Tn mutants in the PhoPR TCS. Based on these

findings, we propose a model where PhoPR arrests growth by diverting propionate towards the

synthesis of cell envelope lipids and, when mutated, growth is promoted by metabolism of

propionate by the MCC into pyruvate, a permissive carbon source for growth at acidic pH(17).

Additionally, we found that pyrazinamide (PZA) treatment also suppresses the acid growth arrest

phenotype in WT Mtb and on propionate PZA selectively sensitizes the *phoPR* mutant to killing

PZA. Together, these findings support a model linking acidic pH, carbon source, PhoPR signaling

and growth state for PZA activity.

Results

PhoPR is required for acid growth arrest on propionate.

We previously reported that Mtb grows on propionate at pH 7.0, but arrests growth on at pH 5.7, based on changes in optical density (17). To determine if Mtb is viable and non-replicating, we examined growth by optical density and CFUs over the course of 12 days of Mtb incubated on propionate or pyruvate at pH 5.7. Mtb arrested growth on propionate (Figure 1A) and was viable (Figure 1B), whereas pyruvate was permissive for growth. Therefore, like acid growth arrest on glycerol, Mtb is non-replicating and viable during propionate-induced acid growth arrest.

We hypothesized that growth arrest on propionate is a regulated process and not intrinsic to the carbon source itself, given that the carbon source can be metabolized at neutral pH and Mtb is viable. To test this hypothesis, a forward genetic selection was performed to identify suppressor mutants of acid growth arrest on propionate at acidic pH. An Mtb Erdman transposon (Tn) mutant library containing ~100,000 mutants was plated on agar plates containing minimal media buffered to pH 5.7 and supplemented with 2mM propionate as a sole carbon source. Mutants with enhanced acidic growth (EAG) phenotype formed colonies on the plates following 8 weeks of incubation. In total, 20 colonies were isolated, and 16 were confirmed as mutants that could grow on propionate (Figure 1C and Table S1). Using whole genome sequencing or inverse PCR, we identified Tn insertion sites in 12 of the 16 confirmed mutants and all sequenced Tn insertions were in *phoR or phoP,* including 6 independent insertions in *phoR and* one insertion in *phoP.* (Supplemental Table 1). Notably, several of the isolated *phoR* mutants also had mutations in the PDIM gene *ppsE*. We confirmed the phenotype of these mutants and observed the *phoR*::Tn and *phoP*::Tn mutants grew on propionate at acidic pH, with higher growth in the *phoR*::Tn mutant (Figure 1D). Using a *ΔphoPR* deletion mutant and complemented strain previously generated in the CDC1551 background (10), we confirmed that at pH 5.7 on propionate the *ΔphoPR* mutant demonstrated robust growth compared to the wildtype and the phenotype was complemented (Figure 1E). There was no significant difference between strains at neutral

pH (Figure S1A). Therefore, we conclude that PhoPR is required for acidic growth arrest on

propionate at acidic pH.

The *phoPR* mutant has enhanced growth on multiple carbon sources at acidic pH

Previously, we established that *phoPR* is required to slow the growth of Mtb on the

permissive carbon source pyruvate at acidic pH(17). Therefore, we sought to determine if *phoPR*

is required for growth arrest on other carbon sources. We observed that the *ΔphoPR* mutant

displays enhanced growth on specific carbon sources that are permissive for growth at acidic pH

(pyruvate and acetate) and normally non-permissive for growth at acidic pH (propionate and

succinate), with the strongest enhanced growth phenotype observed in propionate (Figure 2).

We previously observed that acid growth arrest on glycerol was dependent on both pH

and carbon concentration(19). Therefore, we defined the interactions of pH and propionate

concentration on Mtb growth by testing growth at varying pH levels (pH 5.0-7.0), and propionate

concentrations (0 mM and 20 mM) (Figure 3 A-F). This assay was conducted in 96 well plates

where Mtb exhibits less robust growth, as compared to standing flask growth conditions used in

prior experiments. At 8 mM and 20 mM propionate, the *phoPR* mutant exhibited enhanced growth

at pHs 5.0-5.7, (Figure 3 A-C), whereas at pHs 6.0-7.0, the mutant had similar growth to the wild-

type. The *ΔphoPR* mutant exhibited a higher OD at 20 mM propionate, suggesting that growth is

proportionate to carbon source availability. Therefore, acid growth arrest is induced at pH 5.7 or

below. Notably, the PhoPR regulon is strongly induced pH below 6.0(9), consistent with the

induction of PhoPR signaling driving the growth arrest phenotype that we see. *pks2* is strongly

induced at acidic pH in a *phoPR*-dependent manner and is required for the synthesis of sulfolipids.

At acidic pH, we have previously shown that Mtb promotes SL synthesis and accumulation in a

phoPR-dependent manner(17). We previously isolated a *pks2::*Tn mutant and hypothesized that

if enhanced growth is a consequence of diversion of carbon from the TCA cycle, then the *pks2::Tn*

mutant should have enhanced growth on propionate. We examined the growth of the WT,

phoR::Tn and *pks2::Tn* mutants on propionate at acidic pH and observed that the *pks2::*Tn mutant

displayed an intermediate impact, with higher growth than the WT, but less than that of the *phoR*

mutant (Figure 1D). Therefore, the enhanced growth is also dependent on *pks2*, implicating

changes in lipid synthesis or the cell envelope composition in growth arrest.

PhoPR is required for acidic pH-dependent cell death

We were surprised to observe the *phoPR* mutant grow on propionate at pH 5.0 (Figure 3A).

This highly acidic pH can lead to cell death. Therefore, we examined WT, Δ*phoPR* mutant and

complemented strains for growth and viability over a 12-day time course in 2 mM propionate as

a sole carbon source, incubated in standing flasks. As anticipated, we observed a significant

reduction of WT OD and CFUs at acidic pH, indicating cell death, and an increase of CFUs for

the Δ*phoPR* mutant (Figure 4)*.* Therefore, we conclude that cell death at acidic pH, is not an

intrinsic stress associated at acidic pH and instead dependent on PhoPR-dependent activities,

possibly related to remodeling of metabolism. Notably, the differences in growth between the pH

5.0 conditions in the dose response study (Figure 3A) and this experiment, are likely driven by

the more hypoxic conditions in the 96-well plates as compared to the standing flasks due to

differences in head space ratio, which is presumed to result in the more robust growth observed

in standing flasks.

PhoPR restricts growth at acidic pH by diverting carbon from the methyl citrate cycle

One of the key functions of PhoPR at acidic pH is envelope composition remodeling by

promoting lipid synthesis, including sulfolipids (SL), diacyltrehaloses (DAT) and

polyacyltrehaloses (PAT)(13). Deletion of *phoPR* results in a near complete loss of SL at acidic

pH, which is compensated for by the induction of triacylglycerol (TAG) (17). Propionyl-CoA is an

important substrate for the PhoPR-dependent synthesis of methyl-branched lipids such as SL,

DAT and PAT, as well as methyl-branched lipids synthesized independently of PhoPR, such as

PDIM and TAG. Given the partially enhanced growth of the *pks2* mutant (Figure 1D), we

hypothesized that growth arrest at acidic pH is due to PhoPR diverting carbon away from central

metabolism towards lipid metabolism, thus starving Mtb of carbon needed for growth. Indeed, a

similar phenomenon has been observed for hypoxia-driven growth arrest and the induction of *tgs1*

to produce TAG (26). We further hypothesize that in the PhoPR mutant, the propionate can be

metabolized into pyruvate via the methyl citrate cycle (MCC). Pyruvate is permissive for growth

at acidic pH, thus the redirection of propionate from lipid anabolism (in the WT) to pyruvate via

the MCC (in the *phoPR* mutant) could explain the enhanced growth of the Δ*phoPR* mutant on

propionate at acidic pH.

To first test this model, we examined if the MCC was required for growth on propionate at

acidic pH, using mutants in the methylisocitrate lyases *icl1/icl2* and complemented strains(24).

Lacking a *phoPR* mutant in the *icl1/icl2* mutant backgrounds, we used a chemical inhibitor of

phoPR signaling, ethoxzolamide (ETZ) (27), to downregulate PhoPR. The strains were cultured

in minimal media in propionate with or without ETZ, at acidic pH, and growth and viability were

monitored over 12 days(Figure 5A and S2A). In the DMSO control, the WT and *icl1* or *icl2* mutants

all had arrested growth at acidic pH, and the *icl1/icl2* exhibited a reduction of CFUs and OD, a

phenotype consistent with propionate toxicity in the absence of the MCC (Figure 5A and S2A). In

the ETZ-treated cells, we observed enhanced growth of the WT, consistent with PhoPR inhibition

driving growth at acidic pH (Figure 5A and S2A). The *icl1* and *icl2* mutants both had reduced

growth and viability (Figure 5A and S2A), with a the *icl2* mutant showing a greater reduction in

viability which could be driven by the direct regulation of Icl2 by propionate(28). Notably, at pH

7.0, the single *icl1* or *icl2* mutants grew well on propionate in both the DMSO and ETZ treated

cells, while the *icl1/icl2* mutant could not grow, due to propionate toxicity (Figure S3). To further

confirm the role of the MCC in *phoPR*-dependent enhanced growth, we examined if chemical

inhibition of Icl1/2 by itaconic acid (ITA) could inhibit the enhanced growth. Indeed, ITA completely

inhibited Mtb *phoPR* mutant growth on propionate at acidic pH (Figure S4). These data are

consistent with the MCC being required for enhanced growth of the Δ*phoPR* mutant at acidic pH.

To further validate the model that PhoPR-dependent diversion of carbon to lipid synthesis

promotes the growth arrest phenotype, we hypothesized that stimulating lipid synthesis in the

PhoPR mutant would suppress growth on propionate. To test this hypothesis, we examined

whether the induction of the methylmalonyl-CoA pathway, by the supplementation with vitamin

B12, reduces the growth of the PhoPR mutant. The WT, Δ*phoPR*, and complement strains were

grown in minimal media at acidic pH supplemented with propionate, with or without vitamin B12,

and growth and viability were monitored over 12 days. As hypothesized, vitamin B12 suppresses

the growth of the Δ*phoPR* mutant at acidic pH (Figure 5B and S2B), supporting that stimulating

lipid anabolism suppresses growth at acidic pH on propionate. Addition of vitamin B12 had no

impact on Mtb growth at pH 7.0 on propionate , in the WT or Δ*phoPR* mutant (Figure S2D),

supporting the phenotypes are acidic pH dependent.

Mtb increases the production of the TAG when *phoPR* is deleted, possibly as a

mechanism to balance redox homeostasis(17). We hypothesized that TAG synthesis may function

similarly as a sink for carbon and slow the growth of WT or the Δ*phoPR* mutant. To test this

hypothesis, we used a DosRST inhibitor, MSU-43996, that potently inhibits *tgs1* expression and

TAG synthesis (29). We examined the growth of WT, and the Δ*phoPR* mutant in the presence of

propionate, with or without 40 µM MSU-43996 at acidic pH and neutral pH (Figure 5C and S1B) .

When *tgs1* is inhibited, the WT exhibits enhanced growth, and the Δ*phoPR* mutant exhibits further

enhanced growth at acidic pH (Figure 5C and S2C); while no major differences are observed at

neutral pH (Figure S1B). These data further reinforce the hypothesis that lipid synthesis is slowing

or arresting Mtb growth at acidic pH.

PhoPR restricts CoA at acidic pH

We hypothesize that if PhoPR is diverting carbon away from central metabolism to slow growth, this adaptation may be associated with less available CoA in the WT at acidic pH. In the Δ*phoPR* mutant, metabolism of propionate into pyruvate would fuel the TCA cycle and promote increases in CoA pools. To test this hypothesis, we examined total CoA in the WT, Δ*phoPR* mutant, and complemented strains in 2mM propionate, 10 mM pyruvate or 10 mM glycerol at pH 5.7. During growth arrest in propionate and glycerol, WT CoA pools remained relatively stable over the course of six days (Figures 6A and S5). In contrast, on propionate CoA pools in the *phoPR* mutant increased threefold at day 3 and then lowered again at day 6 (Figure 6A). There is no change between day 0 and 6 in CoA in the growth arrested *phoPR* mutant in the presence of glycerol (Figure S5). In pyruvate, which is growth permissive, CoA pools increased slightly in the WT and were enhanced in the *phoPR* mutant, which also has enhanced growth in pyruvate (Figure 6C). Consistent with our hypothesis, diversion of carbon from the TCA cycle to lipid synthesis by the addition of vitamin B12, decreases the accumulation of CoA in the *phoPR* mutant (Figure 6B), which is also associated with decreased growth (Figure 5B).

In our findings, total CoA in the *phoPR* mutant on propionate at acidic pH is negatively associated with stimulation of lipid synthesis pathways and positively associated with growth. The decrease in CoA pools at Day 6 in the *phoPR* mutant suggested that additional lipid synthesis pathways may be engaged over time. The cultures are grown in standing flasks and we anticipate that growth over 6 days by the *phoPR* mutant will consume oxygen and induce the *dosRST* pathway. This would cause the strong induction of the TAG-synthase *tgs1* and provide an induced, second mechanism to restrict growth under the hypoxic conditions caused by the growth of the mutant. To test this hypothesis, we examined the impact of the *dosRST* inhibitor MSU-39446, on CoA pools. In the MSU-39446 treated *phoPR* mutant the CoA pools did not decrease at day 6 (Figure 6D), a phenotype consistent with the enhanced growth caused by MSU-39446 treatment (Figure 5C). Notably, CoA pools increased in the WT, a response consistent with the enhanced

growth of the WT (Figure 5C). Together, these data further support the hypothesis that growth

arrest on propionate at acidic pH is driven by restriction of central metabolism via synthesis of

PhoPR-dependent lipids.

Pyrazinamide suppresses growth arrest on propionate at acidic pH.

Pyrazinamide (PZA) activity is enhanced at acidic pH and its mechanism of action is

uncertain, but recent data suggests it may be associated with CoA metabolism and cell envelope

lipid homeostasis (17, 30-32). Therefore, we hypothesized that the WT strain and *phoPR* mutant

may exhibit differential sensitivity to PZA. To test this hypothesis, we examined the sensitivity of

WT, Δ*phoPR* mutant, and complemented strains to PZA across a dose range from 0-100 μM in

minimal medium supplemented with 2mM propionate at pH 5.7. Surprisingly, following 21 days of

incubation, PZA at concentrations of 2.5 and 10 μM suppressed acid growth arrest in the WT and

complemented strains (Figure 7A). However, the *phoPR* mutant was growth arrested under these

conditions, showing that it has enhanced sensitivity to PZA, as compared to the DMSO control,

where the *phoPR* mutant grew robustly. At a concentration of 1.25 μM, the growth of the WT and

phoPR mutant are about equal (Figure 7A) ,suggesting a complex interplay between PZA

concentration and modulation of growth at acidic pH on propionate.

The above assay was conducted in 96-well plates, and we sought to better characterize

the suppression of acid growth arrest by PZA in a standing flask-based acid growth arrest assay,

which exhibits more robust growth. WT, Δ*phoPR* mutant and complemented strains were

incubated at pH 5.7 with propionate as a sole carbon source, treated with 3.8 μM PZA, 20 μM

isoniazid (INH), or a DMSO control, and growth was monitored over 12 days. The WT and

complemented strains exhibited enhanced growth on PZA as compared to DMSO (Figure 7A and

7B), whereas the *phoPR* mutant was killed by PZA and showed robust growth in the DMSO control

(Figure 7D and 7E). We hypothesized that like enhanced growth of the Δ*phoPR* mutant, PZA may

also be causing enhanced growth by inhibiting lipid synthesis. To test this hypothesis, we

examined if additional of vitamin B12, limited the enhanced growth of PZA treated WT Mtb. As

hypothesized, vitamin B12 blocked the enhanced growth in the PZA treated Mtb (Figures S6 and

S7). INH inhibited the growth of all of the strains, showing the differential growth impact is PZA-

specific (Figure 7F and 7G). We also examined the impact of PZA sensitivity on the strains grown

on pyruvate and observed that that WT Mtb did not show enhanced growth on PZA, showing the

enhanced growth is propionate specific. (Figure S8). Notably, the Δ*phoPR* mutant was also more

sensitive to PZA on pyruvate (Figure S8) and INH on propionate (Fig. 7G), supporting that the

enhanced PZA sensitivity is not carbon source specific and the *phoPR* mutant is more sensitive

to distinct classes of antibiotics. To determine if the enhanced growth on WT Mtb on PZA is shared

between different strains, we also conducted this experiment in Mtb Erdman. We observed that

3.8 µM PZA caused enhanced growth of WT Erdman (Figure S9). No significant differences were

observed, between strains or treatments, when the experiment was replicated at neutral pH

(Figure S10), an expected result given the acidic pH-dependent activity of PZA. Together, these

findings show that PZA can result in enhanced growth or killing of Mtb depending on the pH,

concentration of PZA and PhoPR-signaling of the bacterium. This discovery generates further

complexity for understanding the mechanism of PZA function, as pH and carbon sources can vary

in different environments during infection, and differences in *phoPR* or *prpR* genotype and

function exist in clinical strains (33-35).

Discussion

Mtb remodels its physiology at acidic pH below ~6.5, demonstrating several pH-dependent

adaptations associated with pathogenesis and drug susceptibility(5, 36). Changes in carbon

metabolism are a substantial component of acidic pH-dependent adaptations. We and others

have observed that Mtb restricts its ability to grow on specific carbon sources at acidic pH, a

phenomenon we refer to as acid growth arrest. Previously, we found that Mtb arrests its growth

on glycerol and that this growth arrest is overcome by mutations *in ppe51* that promote enhanced

glycerol uptake (36). We concluded that growth arrest is overcome by increasing glycerol uptake

(36), possibly to overcome reduced glyceraldehyde-3-phosphate dehydrogenase activity at acidic

pH(37). *ppe51* mutants did not promote growth on other non-permissive carbon sources, including

propionate, leading us to hypothesize that acid growth arrest has different mechanisms for

different carbon sources.

The isolation of *phoPR* mutants that can grow on propionate strongly support our

hypothesis of carbon source specific metabolic restraints at acidic pH (Figure 8). For example,

phoPR mutants cannot grow on glycerol and *ppe51* EAG variants cannot grow on propionate.

Together, multiple independent observations support that the growth arrest on propionate is due

to the diversion of carbon towards the synthesis of PhoPR-dependent lipids (Figure 8). For

example, we observe growth arrest at acidic pH when methyl-branched lipid synthesis pathways

are activated on propionate, and when PhoPR-dependent pathways (SL, DAT and PAT), DosRST-

dependent pathways (TAG), or vitamin B12-dependent pathways (PDIM) are available to the

pathogen. Disruption of these pathways by deletion of *phoPR* (Fig. 8), interruption of the *phoPR*-

regulated gene *pks2* (Fig. 1*)*, inhibition of PhoPR-by treatment with ETZ (Fig. 5), inhibition of *tgs1*

expression by treatment with MSU-39446 (Fig. 8), or the absence of vitamin B12, results in the

enhanced acid growth phenotype. Notably, treatment with PZA also leads to enhanced acid

growth, suggesting part of PZA function may be related to inhibiting lipid synthesis. When lipid

synthesis is blocked, growth on propionate at acidic pH is dependent on the MCC, which is one

of primary mechanisms by which Mtb detoxifies propionate (38). Pyruvate is a permissive carbon

source at acidic pH, therefore, it is expected that if propionate is metabolized to pyruvate via the

MCC that it would promote growth (Figure 8). Given the many regulatory functions of PhoPR(39),

it remains possible that PhoPR is also regulating lipid-synthesis independent mechanisms to

arrest growth. Given the link between growth arrest and decreased CoA-pools, this could possibly

be a regulatory activity inhibiting central metabolism. We have also not explored the potential role

of cAMP signaling in the observed phenotypes, where PhoPR modulation of cAMP could be

impacting propionate detoxification mechanisms and cell envelope permeability(40, 41).

The presented data show that the PhoPR plays a role in the detoxification of propionate.

We observed that the DMSO treated WT strain and the single Δ*icl1* or Δ*icl2* mutants do not have

a survival defect (Figure. 5 and S2) and the double Δ*icl1/2* mutant has its viability reduced by ~1

log relatively to the starting CFUs. In contrast, when PhoPR signaling is inhibited by ETZ, viability

is reduced in Δ*icl2* mutants and in the Δ*icl1/2* double mutant by ~100 fold (Figure. 5 and S2). This

demonstrates that inhibition of PhoPR sensitizes Mtb to propionate toxicity. Therefore, PhoPR is

exerting dual functions under the tested conditions, to both arrest growth and detoxify propionate.

Mtb induces PhoPR at acidic pH on various carbon sources (glycerol, pyruvate,

propionate), and WT Mtb has a functional MCC, therefore, we do not believe that PhoPR is solely

induced specifically to detoxify propionate. Rather, we observed that the PhoPR mutant

experiences reductive stress at acidic pH, and we hypothesize that lipid synthesis serves the

purpose of oxidizing cofactors (such as NADPH to NADP+) to maintain redox homeostasis(17,

18). Consistent with this hypothesis, we observe that downregulation of SL production in the

phoPR mutant is associated with induction of other lipids such as TAG (17). Blocking both PhoPR

and DosRST pathways removes four of the major lipid synthesis pathways (SL, DAT, PAT and

TAG) and results in the most robust growth of propionate (Figure 5C, S2C). Pyruvate is known to

promote growth at acidic pH, and we hypothesize that by entering the TCA cycle near the

anaplerotic node provides Mtb the metabolic flexibility to maintain redox homeostasis by

balancing anabolism and catabolism, possibly via the PEP glyoxylate cycle(42, 43). In this

manner, if propionate is not diverted to lipid synthesis, it can be turned into a permissive carbon

source that balances redox homeostasis at the anaplerotic node, even in the absence of *phoPR*.

Similar to the mutations in *phoPR*, PZA at low concentrations can also suppress Mtb

growth arrest on propionate at acidic pH. Based on our model for propionate-dependent growth

arrest, we reason that PZA treatment may be acting to inhibit lipid synthesis, thus enabling more

propionate to be metabolized by the MCC into central metabolism. The mechanism of action of

PZA remains controversial, but recent evidence supports a role for CoA metabolism and changes

in cell envelope lipid synthesis playing a role in its activity (30, 44) . Resistance mutations to PZA

arise in PanD, which is involved in the synthesis of CoA. The PZA metabolite POA can bind to

PanD to inhibit CoA synthesis (32). Reduced CoA levels are predicted to result in lower levels of

lipid synthesis, as it is an essential cofactor in the initial steps of fatty acid activation. Notably, it

was recently reported that PDIM genes are strongly downregulated by treatment with PZA (45)

and PDIM mutants are associated with resistance to PZA (44). Therefore, we speculate enhanced

growth of the WT is due to reduced PDIM synthesis or other long-chain fatty acids that require

CoA for their synthesis. The small range of concentrations where this enhanced growth in the WT

is observed (~3 µM) may reflect a hypomorphic phenotype, where CoA metabolism is sufficiently

repressed by PZA to impact lipid metabolism, but not so much so to fully deplete CoA and kill the

cell. It is surprising that the $\Delta phoPR$ mutant is susceptible to PZA and does not have further

enhanced growth. Furthermore, the $\Delta phoPR$ mutant is more susceptible to PZA on both

propionate and pyruvate (Fig. 7 and Fig. S8), and INH on propionate (Fig. 7G), but rather

associated with enhanced growth at acidic pH. Because propionate is metabolized to pyruvate by

the MCC it is consistent with our model that PZA sensitivity would be similar on either propionate

and pyruvate. The $\Delta phoPR$ is also more sensitive to INH on propionate (Fig. 7G), suggesting

more general mechanisms be driving antibiotic susceptibility in the mutant. We speculate that an

alternative mechanism is driving PZA or INH sensitivity in the *phoPR* mutant, possibly differential

expression of a *phoPR* pathway independent on lipid anabolism. For example, a *ppe51* loss of

function mutant is reported to have enhanced susceptibility to PZA(46). *ppe51* is downregulated

in the *phoPR* mutant, suggesting it, or some other unknown PhoPR-regulated genes may play a

role in susceptibility that overcomes the expected further enhanced growth. Alternatively, the

phoPR mutant may have has increased permeability due to changes in the cell envelope or cAMP

signaling (13, 40, 47). It is possible that this increased permeability is increasing the intracellular

concentration of PZA or INH, accumulating to sufficient levels to inhibit growth.

Concluding remarks

The observation that PhoPR restricts growth on propionate at acidic pH is analogous to

the findings of Baek *et al*, who showed that the DosRST and hypoxia regulated gene *tgs1* inhibits

growth during hypoxia by diverting carbon from the TCA cycle(26). In both cases, growth

restriction is an environmental adaption and not an inherent metabolic limitation. TB disease is

caused by slow growing mycobacteria and persistence and growth arrest are associated with

drug tolerance. By understanding the complex mechanisms by which Mtb integrates

environmental cues, metabolism and growth arrest, we may define new drug targets that can

disrupt pathogenesis in a way the increases bacterial growth. Enhanced growth is associated with

reduced fitness in macrophages (36) and enhanced drug susceptibility *in vitro* (18). Therefore,

these non-traditional drug targets may have unique impacts during treatment, including treatment

shortening. Indeed, PZA was essential to shortening TB treatment times to six months and it is

tempting to speculate that some of this activity may be related to PZA enhancing Mtb growth in

mildly acidic and cholesterol rich environments, such as the macrophage or the granuloma. It is

also important to consider that *phoPR* and *prpR* have variants in clinical isolates (33-35),

supporting that the interplay of PhoPR and propionate observed in these studies, maybe

associated with differential pathogenesis and drug susceptibility and in human infections and drug

treatment.

Methods

Bacterial strains and growth

Mtb experiments were conducted with WT ERDMAN or WT CDC1551 as indicated. The Δ*phoPR*
deletion mutant, and complemented strains are in the CDC1551 background (10) and the *icl1/2*
mutants and complemented strains are in the Erdman background(24). Cultures were maintained
in 7H9 media supplemented with 10% OADC and .05% Tween-80. All single carbon source
experiments were performed in MMAT minimal media buffered to acidic pH 5.7 or pH 7.0(21),
using (1 g l−1 KH2PO4, 2.5 g l−1 Na2PO4, 0.5 g l−1 (NH4)2SO4, 0.15 g l−1 asparagine, 10 mg l−1
MgSO4, 50 mg l−1 ferric ammonium citrate, 0.1 mg l−1 ZnSO4, 0.5 mg l−1 CaCl2, and 0.05%
Tyloxapol) . For 12-day growth experiments, time points were taken every three days. For each
of our growth curve experiments, 2mM propionate was used unless stated otherwise. Chemical
inhibitors and supplements were used at the following concentrations: Ethoxzolamide at 40 μM,
MSU-39446 at 40 μM, Itaconic acid at 2 mM and vitamin B12 at 10ug/ml. Mtb was seeded in T-
25 standing tissue culture flasks in 8ml of minimal media at an initial cell density of 0.05 OD and
incubated at 37 °C and 5% CO_2. 500 μL of each culture was taken each day for optical density
measurements. Bacterial viability was assessed enumerating colony-forming units (CFUs) on
7H10+OADC plates.

Genetic selection

Transposon mutagenesis was performed in Mtb Erdman using the ϕMycoMarT7 Tn system as
previously described(48), generating a library with ~100,000 Tn mutants. The Tn mutant library
was plated on MMAT pH 5.7 agar plates supplemented with 2mM propionate as the sole carbon
source. Plates were incubated at 37°C, with mutants appearing around week 8 and isolated for
growth. Single-colony isolates were confirmed as EAG (enhanced acidic growth) mutants under
acidic conditions in liquid media MMAT pH 5.7 supplemented with 2mM propionate. The

transposon insertion sequences were confirmed using inverse PCR (49) or whole genome

sequencing.

<u>Whole Genome Sequencing</u>

Genomic DNA of selected mutants as well as the WT Erdman control was isolated, DNA libraries

were constructed, and sequenced using the Illumina MiSeqm in-paired end, 250 bp read format

(PE250). After the sequencing run, reads were demultiplexed and converted to FASTQ format

using the illumina bcl2 fastq (v1.8.4) script. The reads in the raw data files were then subjected

to trimming of low-quality bases and removal of adapter sequences using Trimmomatic v0.36,

with a 4 bp sliding window, and a read quality cutoff of below 15 or read length cutoff less than 36

bp. The trimmed reads were then aligned to the Erdman reference genome using the Burrow-

Wheeler alignments. Genome analysis tool kit base quality score recalibration, indel alignment,

and duplicate removal were applied, and SNP and INDEL discovery were performed.

<u>pH and propionate dose response combination growth assays</u>

Mtb cultures were incubated in a range of pH-buffered MMAT media ranging from pH 5.0-7.0 at a

starting OD of 0.2 in 96 well plates. Cultures were treated with a 2.5-fold serial dilution (0mM-

20mM) of propionate and incubated over a course of 21 days, with growth assessed by optical

density.

<u>Coenzyme A assay</u>

Coenzyme A was measured using a commercially available assay (Sigma-Aldrich). Cultures were

maintained in 7H9, and then seeded to an OD of 0.2 and resuspended in minimal media

supplemented with 2mM propionate in 8 ml of media. The assay measures both free CoA and

acyl-CoA. Total CoA was measured at Day 0, 3, and 6 and concentration determined using a

standard curve.

Pyrazinamide Sensitivity Assays

Dose response assays were conducted in a 96 well plate cultures seeded at OD 0.2 and in minimal media supplemented with 2mM propionate across a dose response from 100 μM-0 μM of PZA. Plates were incubated for 6 days in a sealed Ziploc bag at 37C and OD read on a plate reader. Standing flask assays were conducted using conditions as described earlier in the methods. WTCDC, $\Delta phoPR$ mutant, and complement were grown in minimal media at acidic pH supplemented with propionate and treated with DMSO, 3.84 uM Pyrazinamide, and 20uM Isoniazid using conditions described above. Growth was measured Day 0, 3, 6, 9, and 12. Viability was measured Day 0 and Day 12 by enumerating CFUs.

Acknowledgements

We thank Prof. John McKinney for sharing the *icl1/icl2* mutant strains. Research on this project was supported by grants from the NIAID (R01 AI116605 and R01 AI150855) and AgBioResearch.

Figure Legends

Figure 1. PhoPR arrests growth on propionate at acidic pH. A) Growth curves of WT Mtb CDC strain grown in acidic pH for 12 days, in media supplemented with propionate or pyruvate. B) Bacterial viability evaluated using CFUs plated at days 0 and 12, demonstrating the viability of the growth-arrested bacteria under propionate. C) Relative growth of 16 different transposon mutants with enhanced growth on propionate relative to the WT (dotted line). All sequenced mutants were in *phoR or phoP* (Supplemental Table 1). D) Growth curves of transposon mutants in *phoR*, *phoP* and *pks2*, grown at acidic pH in minimal media supplemented with 2mM propionate. E) Growth curves of $\Delta phoPR$ mutant and complemented strain compared to WT CDC strain showing growth of the *phoPR* mutant. Multiple comparison unpaired t-test was used for the

growth curves analysis while one-way ANOVA was used for the viability analysis,   * <0.05, **

<0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure 2. The ΔPhoPR has enhanced growth on specific carbon sources at acidic pH. Mtb

growth was evaluated on various carbon sources after 21 days incubation. The Δ*phoPR* mutant

exhibits enhanced growth at acidic pH on selected permissive (acetate and pyruvate) and non-

permissive carbon sources (propionate and succinate). Two-way ANOVA was used for statistical

analysis, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice

with similar results.

Figure 3. Enhanced growth of the Δ*phoPR* mutant is dependent on level of pH and

propionate concentration. Growth of the Δ*phoPR* mutant, the WT and complemented strains

was tested in minimal media supplemented with a range of concentrations of propionate (20 mM

to 0.16 mM) at different pHs (5.0-7-.0). The Δ*phoPR* mutant has enhanced growth in dependent

on propionate concentration at pHs of 5.0, 5.5 and 5.7 (**A-C**), but not 6-7 (**D-F**). A multiple

comparison unpaired t-test was used for this analysis, * <0.05, ** <0.01, *** <0.001, **** <0.0001.

Experiments were replicated at least twice with similar results.

Figure 4. PhoPR is required for cell death at pH 5.0. A) The WT, Δ*phoPR* mutant, and

complemented strain were incubated at acidic pH for 12 days, in minimal media supplemented

with propionate. The Δ*phoPR* mutant has enhanced growth compared to the WT, a phenotype

that is complemented. **B)** Viability assays, show that the WT and complemented strains exhibit

cell death at pH 5.0, while the *phoPR* mutant grows. A multiple comparison unpaired t-test was

used for this analysis, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at

least twice with similar results.

Figure 5. PhoPR restricts growth at acidic pH by diverting carbon from the methyl citrate cycle. A) Growth curves of the WT, knockout strain of isocitrate lyases 1 and/or 2, and the complemented strains, treated with DMSO or ethoxzolamide (ETZ), an indirect inhibitor of PhoPR, showing a significant reduction in growth in the *icl2* and double knockout strains. **B)** Growth curves conducted in wildtype, *phoPR* mutant, and complemented strains in minimal media buffered to pH 5.7 supplemented with 2mM propionate and vitamin B12 over a course of 12 days, showing vitamin B12 suppresses enhanced growth in the Δ*phoPR* mutant strain.**C)** Growth of the WT, Δ*phoPR* mutant, and complement strains treated with MSU-43996 or the DMSO control, showing an enhanced growth in the presence of MSU-43996 treatment. An unpaired t-test was used between individual groups and the WT for the growth curves, while a one-way ANOVA was used for the viability experiments, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure 6: Enhanced acid growth is associated with increased CoA pools. A) CoA pools in the Δ*phoPR* mutant, WT CDC, and complemented strain grown in minimal media supplemented with 2 mM propionate at acidic pH, showing enhanced metabolic capacity (i.e., higher CoA pools) in the Δ*phoPR* mutant during the first three days of growth ($p = 0.006$), which significantly decreases around day 6 ($p = 0.009$). **B)** CoA pools in the three strains, grown in minimal media supplemented with vitamin B12 at acidic pH, showing no significant differences between the three strains throughout the 6-day growth period. **C)** CoA pools in the three strains, grown in minimal media supplemented with 10 mM of pyruvate at acidic pH, showing enhanced metabolic capacity of the Δ*phoPR* mutant and WT through day 6. **D)** CoA pools in the three strains, grown in minimal media supplemented with 40 µM of MSU-43996 at acidic pH, showing enhanced metabolic capacity of the Δ*phoPR* mutant when *tgs1*, a is inhibited. Two-way ANOVA was used for this

analysis, *** <0.05, ** <0.01, *** <0.001, **** <0.0001.** Experiments were replicated at least twice

with similar results.

Figure 7: PZA suppresses acid growth arrest on propionate. A) Dose-response relationship

of pyrazinamide (100-0 µM) in the Δ*phoPR* null mutant, WT, and complement strains, conducted

in minimal media supplemented with 2mM propionate at acidic pH, demonstrating enhanced

growth of the WT and complemented strains on PZA at concentrations between 5.0-2.5 µM. **B**

and C) Assessment of the growth (B) and viability (C) of the three strains in minimal media

supplemented with 2mM propionate at acidic pH, treated with the vehicle as a control, showing

enhanced growth of the Δ*phoPR* mutant. **D and E)** Assessment of the growth (D) and viability (E)

of the three strains in minimal media supplemented with 2mM propionate at acidic pH treated with

PZA (3.84 µM), demonstrating enhanced growth of the WT and higher sensitivity of the Δ*phoPR*

mutant. **F and G)** Assessment of the growth (F) and viability (G) of the three strains in minimal

media supplemented with 2mM propionate at acidic pH treated with isoniazid (INH, 20 µM),

demonstrating similar levels of sensitivity in the three strains. Multiple comparison T-test was used

for the growth curves analysis, while Two-way ANOVA was used for the viability, *** <0.05, ** <0.01,**

*** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure 8: Model for Mtb growth arrest on propionate at acidic pH and mechanisms of

suppression by mutations in *phoPR* or pyrazinamide treatment. Propionate-dependent acid

growth arrest is a PhoPR-regulated response driven by diversion of carbon away from central

metabolism and into the synthesis lipids. At acidic pH, propionyl-CoA is used as a substrate to

synthesize methyl-branched lipids, either through the polyketide synthases Pks2, Pks3 and Pks4,

or through the methyl-malonyl pathway. Additionally, the enzyme MutAB, a vitamin B12-

dependent enzyme, catalyzes the reversible interconversion between malonyl-CoA and succinyl-

CoA, contributing to the synthesis of branched lipids. In either cases, low pH triggers remodeling

of central metabolism towards complex lipids synthesis to alter the mycomembrane structure as

a pH-dependent adaptation, consuming the CoA pools and slowing, or arresting growth in the

process. This pH-dependent adaptation in lost in the *phoPR* null mutant or under indirect PhoPR

chemical inhibition using ethoxzolamide (ETZ). Instead, the propionate is metabolized into

pyruvate by the methylcitrate cycle, a permissive carbon source, increasing CoA pools and

permitting fast growth. Additionally, at acidic pH, the *phoPR* null mutant upregulates triacylglycerol

synthesis using tgs1. Chemical inhibition of *tgs1* using MSU-39446 prevents diversion of CoA

pools into TAG synthesis and away from central metabolism, increasing CoA pools, metabolism

and growth in both the WT and *phoPR* null mutant. PZA treatment causes enhanced growth in

the WT, which may be driven by inhibiting lipid synthesis via depletion of CoA pools.

References

1. Gill WP, Harik NS, Whiddon MR, Liao RP, Mittler JE, Sherman DR. 2009. A replication clock for Mycobacterium tuberculosis. Nat Med 15:211-4.

2. Rohde KH, Veiga DF, Caldwell S, Balazsi G, Russell DG. 2012. Linking the transcriptional profiles and the physiological states of Mycobacterium tuberculosis during an extended intracellular infection. PLoS Pathog 8:e1002769.

3. Munoz-Elias EJ, Timm J, Botha T, Chan WT, Gomez JE, McKinney JD. 2005. Replication dynamics of Mycobacterium tuberculosis in chronically infected mice. Infect Immun 73:546-51.

4. Zheng H, Abramovitch RB. 2019. Host-pathogen interactions influencing Mycobacterium tuberculosis persistence and drug tolerance *In* Lewis K (ed), Persister Cells and Infectious Disease. Springer.

5. Baker JJ, Dechow SJ, Abramovitch RB. 2019. Acid Fasting: Modulation of Mycobacterium tuberculosis Metabolism at Acidic pH. Trends Microbiol 27:942-953.

6. Adams KN, Takaki K, Connolly LE, Wiedenhoft H, Winglee K, Humbert O, Edelstein PH, Cosma CL, Ramakrishnan L. 2011. Drug tolerance in replicating mycobacteria mediated by a macrophage-induced efflux mechanism. Cell 145:39-53.

7. Liu Y, Tan S, Huang L, Abramovitch RB, Rohde KH, Zimmerman MD, Chen C, Dartois V, VanderVen BC, Russell DG. 2016. Immune activation of the host cell induces drug tolerance in Mycobacterium tuberculosis both in vitro and in vivo. J Exp Med 213:809-25.

8. Walters SB, Dubnau E, Kolesnikova I, Laval F, Daffe M, Smith I. 2006. The *Mycobacterium tuberculosis* PhoPR two-component system regulates genes essential for virulence and complex lipid biosynthesis. Molecular Microbiology 60:312-330.

9. Abramovitch RB, Rohde KH, Hsu FF, Russell DG. 2011. *aprABC*: a *Mycobacterium tuberculosis* complex-specific locus that modulates pH-driven adaptation to the macrophage phagosome. Mol Microbiol 80:678-94.

10. Tan S, Sukumar N, Abramovitch RB, Parish T, Russell DG. 2013. Mycobacterium tuberculosis Responds to Chloride and pH as Synergistic Cues to the Immune Status of its Host Cell. PLoS Pathog 9:e1003282.

11. Dechow SJ, Goyal R, Johnson BK, Haiderer ER, Abramovitch RB. 2025. Carbon dioxide regulates Mycobacterium tuberculosis PhoPR signaling and virulence. Infect Immun 93:e0056824.

12. Feng L, Chen S, Hu Y. 2018. PhoPR Positively Regulates whiB3 Expression in Response to Low pH in Pathogenic Mycobacteria. J Bacteriol 200.

13. Asensio JG, Maia C, Ferrer NL, Barilone N, Laval F, Soto CY, Winter N, Daffe M, Gicquel B, Martin C, Jackson M. 2006. The virulence-associated two component PhoP-PhoR

system controls the biosynthesis of polyketide-derived lipids in *Mycobacterium*
tuberculosis. Journal of Biological Chemistry 281:1313-1316.

14. Perez E, Samper S, Bordas Y, Guilhot C, Gicquel B, Martin C. 2001. An essential role for
phoP in *Mycobacterium tuberculosis* virulence. Molecular Microbiology 41:179-187.

15. Martin C, Williams A, Hernandez-Pando R, Cardona PJ, Gormley E, Bordat Y, Soto CY,
Clark SO, Hatch GJ, Aguilar D, Ausina V, Gicquel B. 2006. The live *Mycobacterium*
tuberculosis phoP mutant strain is more attenuated than BCG and confers protective
immunity against tuberculosis in mice and guinea pigs. Vaccine 24:3408-3419.

16. Luabeya AKK, Rozot V, Imbratta C, Ratangee F, Shenje J, Tameris M, Mendelsohn SC,
Geldenhuys H, Fisher M, Musvosvi M, Young C, Mulenga H, Bilek N, Mabwe S, Jelsbak
IM, Rodriguez E, Puentes E, Doce J, Aguilo N, Martin C, Pillay C, Tait D, Russell M, Van
Der Merve A, Rutkowski K, Hunt D, Ginsberg A, Scriba TJ, Hatherill M, team A. 2025. Live-
attenuated Mycobacterium tuberculosis vaccine, MTBVAC, in adults with or without M
tuberculosis sensitisation: a single-centre, phase 1b-2a, double-blind, dose-escalation,
randomised controlled trial. Lancet Glob Health 13:e1030-e1042.

17. Baker JJ, Johnson BK, Abramovitch RB. 2014. Slow growth of Mycobacterium
tuberculosis at acidic pH is regulated by phoPR and host-associated carbon sources. Mol
Microbiol 94:56-69.

18. Baker JJ, Abramovitch RB. 2018. Genetic and metabolic regulation of Mycobacterium
tuberculosis acid growth arrest. Sci Rep 8:4168.

19. Dechow SJ, Baker JJ, Murto M, Abramovitch RB. 2022. ppe51 Variants Enable Growth of
Mycobacterium tuberculosis at Acidic pH by Selectively Promoting Glycerol Uptake. J
Bacteriol 204:e0021222.

20. VanderVen BC, Fahey RJ, Lee W, Liu Y, Abramovitch RB, Memmott C, Crowe AM, Eltis
LD, Perola E, Deininger DD, Wang T, Locher CP, Russell DG. 2015. Novel inhibitors of
cholesterol degradation in Mycobacterium tuberculosis reveal how the bacterium's
metabolism is constrained by the intracellular environment. PLoS Pathog 11:e1004679.

21. Lee W, VanderVen BC, Fahey RJ, Russell DG. 2013. Intracellular Mycobacterium
tuberculosis exploits host-derived fatty acids to limit metabolic stress. J Biol Chem
288:6788-800.

22. Savvi S, Warner DF, Kana BD, McKinney JD, Mizrahi V, Dawes SS. 2008. Functional
characterization of a vitamin B-12-dependent methylmalonyl pathway in *Mycobacterium*
tuberculosis: Implications for propionate metabolism during growth on fatty acids. Journal
of Bacteriology 190:3886-3895.

23. Gould TA, van de Langemheen H, Munoz-Elias EJ, McKinney JD, Sacchettini JC. 2006.
Dual role of isocitrate lyase 1 in the glyoxylate and methylcitrate cycles in Mycobacterium
tuberculosis. Mol Microbiol 61:940-7.

24. Munoz-Elias EJ, McKinney JD. 2005. Mycobacterium tuberculosis isocitrate lyases 1 and
2 are jointly required for in vivo growth and virulence. Nat Med 11:638-44.

25.   Upton AM, McKinney JD. 2007. Role of the methylcitrate cycle in propionate metabolism
685   and detoxification in Mycobacterium smegmatis. Microbiology 153:3973-82.

686   26.   Baek SH, Li AH, Sassetti CM. 2011. Metabolic regulation of mycobacterial growth and
687   antibiotic sensitivity. PLoS Biol 9:e1001065.

688   27.   Johnson BK, Colvin CJ, Needle DB, Mba Medie F, Champion PA, Abramovitch RB. 2015.
689   The Carbonic Anhydrase Inhibitor Ethoxzolamide Inhibits the Mycobacterium tuberculosis
690   PhoPR Regulon and Esx-1 Secretion and Attenuates Virulence. Antimicrob Agents
691   Chemother 59:4436-45.

692   28.   Bhusal RP, Jiao W, Kwai BXC, Reynisson J, Collins AJ, Sperry J, Bashiri G, Leung IKH.
693   2019. Acetyl-CoA-mediated activation of Mycobacterium tuberculosis isocitrate lyase 2.
694   Nat Commun 10:4639.

695   29.   Zheng H, Williams JT, Aleiwi B, Ellsworth E, Abramovitch RB. 2020. Inhibiting
696   Mycobacterium tuberculosis DosRST Signaling by Targeting Response Regulator DNA
697   Binding and Sensor Kinase Heme. ACS Chem Biol 15:52-62.

698   30.   Thiede JM, Dillon NA, Howe MD, Aflakpui R, Modlin SJ, Hoffner SE, Valafar F, Minato Y,
699   Baughn AD. 2021. Pyrazinamide Susceptibility Is Driven by Activation of the SigE-
700   Dependent Cell Envelope Stress Response in Mycobacterium tuberculosis. mBio
701   13:e0043921.

702   31.   Gopal P, Gruber G, Dartois V, Dick T. 2019. Pharmacological and Molecular Mechanisms
703   Behind the Sterilizing Activity of Pyrazinamide. Trends Pharmacol Sci 40:930-940.

704   32.   Gopal P, Yee M, Sarathy J, Low JL, Sarathy JP, Kaya F, Dartois V, Gengenbacher M, Dick
705   T. 2016. Pyrazinamide Resistance Is Caused by Two Distinct Mechanisms: Prevention of
706   Coenzyme A Depletion and Loss of Virulence Factor Synthesis. ACS Infect Dis 2:616-626.

707   33.   Liu Q, Zhu J, Dulberger CL, Stanley S, Wilson S, Chung ES, Wang X, Culviner P, Liu YJ,
708   Hicks ND, Babunovic GH, Giffen SR, Aldridge BB, Garner EC, Rubin EJ, Chao MC,
709   Fortune SM. 2022. Tuberculosis treatment failure associated with evolution of antibiotic
710   resilience. Science 378:1111-1118.

711   34.   Malaga W, Payros D, Meunier E, Frigui W, Sayes F, Pawlik A, Orgeur M, Berrone C,
712   Moreau F, Mazeres S, Gonzalo-Asensio J, Rengel D, Martin C, Astarie-Dequeker C,
713   Mourey L, Brosch R, Guilhot C. 2023. Natural mutations in the sensor kinase of the PhoPR
714   two-component regulatory system modulate virulence of ancestor-like tuberculosis bacilli.
715   PLoS Pathog 19:e1011437.

716   35.   Hicks ND, Yang J, Zhang X, Zhao B, Grad YH, Liu L, Ou X, Chang Z, Xia H, Zhou Y, Wang
717   S, Dong J, Sun L, Zhu Y, Zhao Y, Jin Q, Fortune SM. 2018. Clinically prevalent mutations
718   in Mycobacterium tuberculosis alter propionate metabolism and mediate multidrug
719   tolerance. Nat Microbiol 3:1032-1042.

720   36.   Dechow SJ, Abramovitch RB. 2024. Targeting Mycobacterium tuberculosis pH-driven
721   adaptation. Microbiology (Reading) 170.

37.   Gouzy A, Healy C, Black KA, Rhee KY, Ehrt S. 2021. Growth of Mycobacterium
tuberculosis at acidic pH depends on lipid assimilation and is accompanied by reduced
GAPDH activity. Proc Natl Acad Sci U S A 118.

38.   Savvi S, Warner DF, Kana BD, McKinney JD, Mizrahi V, Dawes SS. 2008. Functional
characterization of a vitamin B12-dependent methylmalonyl pathway in Mycobacterium
tuberculosis: implications for propionate metabolism during growth on fatty acids. J
Bacteriol 190:3886-95.

39.   Broset E, Martin C, Gonzalo-Asensio J. 2015. Evolutionary landscape of the
Mycobacterium tuberculosis complex from the viewpoint of PhoPR: implications for
virulence regulation and application to vaccine development. MBio 6:e01289-15.

40.   Khan H, Paul P, Goar H, Bamniya B, Baid N, Sarkar D. 2024. Mycobacterium tuberculosis
PhoP integrates stress response to intracellular survival by regulating cAMP level. Elife
13.

41.   McDowell JR, Bai G, Lasek-Nesselquist E, Eisele LE, Wu Y, Hurteau G, Johnson R, Bai
Y, Chen Y, Chan J, McDonough KA. 2023. Mycobacterial phosphodiesterase Rv0805 is a
virulence determinant and its cyclic nucleotide hydrolytic activity is required for propionate
detoxification. Mol Microbiol 119:401-422.

42.   Munoz-Elias EJ, McKinney JD. 2006. Carbon metabolism of intracellular bacteria. Cell
Microbiol 8:10-22.

43.   Sauer U, Eikmanns BJ. 2005. The PEP-pyruvate-oxaloacetate node as the switch point
for carbon flux distribution in bacteria. FEMS Microbiol Rev 29:765-94.

44.   Gopal P, Sarathy JP, Yee M, Ragunathan P, Shin J, Bhushan S, Zhu J, Akopian T, Kandror
O, Lim TK, Gengenbacher M, Lin Q, Rubin EJ, Gruber G, Dick T. 2020. Pyrazinamide
triggers degradation of its target aspartate decarboxylase. Nat Commun 11:1661.

45.   Ostrer L, Crooks TA, Howe MD, Vo S, Jia Z, Hegde P, Schacht N, Aldrich CC, Baughn AD.
2025. Mechanism of the Dual Action Self-Potentiating Antitubercular Drug
Morphazinamide. bioRxiv doi:10.1101/2024.10.08.617272.

46.   Bellerose MM, Baek SH, Huang CC, Moss CE, Koh E, Proulx M, Smith C, Baker RE, Lee
JS, Eum S, Shin SJ, Cho SN, Murray M, Sassetti CM. 2019. Common variants in the
glycerol kinase gene reduce tuberculosis drug efficacy. mBio in press.

47.   Jackson M, Raynaud C, Laneelle MA, Guilhot C, Laurent-Winter C, Ensergueix D, Gicquel
B, Daffe M. 1999. Inactivation of the antigen 85C gene profoundly affects the mycolate
content and alters the permeability of the Mycobacterium tuberculosis cell envelope. Mol
Microbiol 31:1573-87.

48.   Long JE, DeJesus M, Ward D, Baker RE, Ioerger T, Sassetti CM. 2015. Identifying
essential genes in Mycobacterium tuberculosis by global phenotypic profiling. Methods
Mol Biol 1279:79-95.

49.   Ochman H, Gerber AS, Hartl DL. 1988. Genetic applications of an inverse polymerase
chain reaction. Genetics 120:621-3.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Free CoA Pool in Minimal media + Propionate

A Propionate

Free CoA Pool in Minimal media + Vit B12

B Propionate + Vitamin B12

Free CoA pool in Minimal media + Pyruvate

C Pyruvate

Free CoA Pool in Minimal media + HC106 + MSU-39446

D Propionate

■ WTCDC   ■ Δ*phoPR*   ■ COMP::*phoPR*

Figure 6

Figure 7

Figure 8

Supplemental Table and Figures

Mycobacterium tuberculosis **growth arrest on propionate at acidic pH is suppressed by mutations in** *phoPR* **and pyrazinamide treatment**

Heather M. Murdoch[1,2], Shelby J. Dechow[1,2], Bassel J. Abdalla[1] and Robert B. Abramovitch[1]

[1]Department of Microbiology, Genetics & Immunology, Michigan State University, East Lansing, MI,48824

[2]Co-first authors contributed equally to this study.

<u>**Supplemental Table**</u>

Table S1: Transposon Mutant Propionate Selection Summary

<u>**Supplemental Figure Legends**</u>

Figure S1. Bacterial Growth and viability of WT, Δ*phoPR* mutant, and complemented strains grown on propionate and treated with MSU-43996 at neutral pH. A) bacterial growth of the three strains on propionate at pH 7.0, showing no significant differences in growth rate. B) bacterial growth of the three strains on propionate at pH 7.0, treated with MSU-43996 or the vehicle, showing no major differences in growth rates based on strain or treatment.

Figure S2. PhoPR restricts growth at acidic pH by diverting carbon from the methyl citrate cycle. A) CFUs from the corresponding experiment in Figure 5A, showing cell death in the ETZ treated icl2 and icl1/2 knockouts. **B)** CFUs of Vitamin B12 treated Mtb strains. **C)** CFUs for figure 5C showing enhanced growth of MSU-43996 treated Mtb. Note, that timepoints are different, but CFUs are from the same experiment. **D)** Vitamin B12 does not impact the growth of strains at neutral pH. A one-way ANOVA was used for the viability experiments, ***** <0.05, ****** <0.01, ******* <0.001, ******** <0.0001. Experiments were replicated at least twice with similar results.

Figure S3. Function of Icl and PhoPR at neutral pH on propionate. Growth of Mtb strains at pH 7.0 on propionate. ETZ did not impact growth in the *Icl1* or *Icl2* single mutants. The double icl1/2 mutant dies due to propionate toxicity. An unpaired t-test was used between individual groups for the growth curves, ***** <0.05, ****** <0.01, ******* <0.001, ******** <0.0001. Experiments were replicated at least twice with similar results.

Figure S4. Itaconic acid inhibits enhanced growth of the Δ*phoPR* mutant. Itaconic acid inhibits the ability of the Δ*phoPR* to grow on propionate at pH 5.7. An unpaired t-test was used between individual groups for the growth curves, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure S5. Free CoA pools on glycerol as a sole carbon source. During acid growth arrest on glycerol, all three strains maintain relatively low free CoA pools. Two-way ANOVA was used for this analysis, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure S6. Vitamin B12 inhibits enhanced growth of WT Mtb treated with PZA. The inhibition of enhanced growth is consistent with PZA inhibiting lipid synthesis to promote growth, as making another lipid synthesis pathway available abolishes enhanced growth. Multiple comparison unpaired t-test was used for the analysis between treatments and the control, and between strains receiving the same treatment, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure S7. Vitamin B12 inhibits enhanced growth of WT Mtb treated with PZA. CFU data for OD data presented in Figure S6.

Figure S8. Sensitivity of Mtb to PZA on Pyruvate as a sole carbon source at pH 5.7. PZA treatment does not result in enhanced growth with pyruvate as a sole carbon source.

Figure S9. PZA suppresses acid growth arrest on propionate in Mtb Erdman. To determine if the enhanced growth by PZA is conserved across strains, the PZA treatment experiment from Figure 7 which as conducted in CDC1551, was repeated in Mtb Erdman. Multiple comparison Unpaired t-tests were used for the growth assays, and one-way ANOVA was used for the viability assays, * <0.05, ** <0.01, *** <0.001, **** <0.0001. Experiments were replicated at least twice with similar results.

Figure S10. Sensitivity of Mtb to PZA treatment at pH 7.0. Control experiment for data in Figure 7, showing PZA is not active on Mtb grown in propionate at pH 7.0.

Table S1: Transposon Mutant Propionate Selection Summary

Mutant #	Approx AT or TA site/gene length (nt)	MT number	Rv Number	Gene Number	PDIM Mutation?	Fold Change (Day 14/Day0)
1.1	416/1458	MT0783	Rv0758	*phoR*	None	3.5
1.2	159/1458	MT0783	Rv0758	*phoR*	None	3.3
1.4	311/744	MT0782	Rv0757	*phoP*	None	1.5
2.1	278/1458	MT0783	Rv0758	*phoR*	*ppsE*	2.1
2.4	145/1458	MT0783	Rv0758	*phoR*	*ppsE*	1.9
3.2	145/1458	MT0783	Rv0758	*phoR*	*ppsE*	1.8
3.3	1124/1458	MT0783	Rv0758	*phoR*	*ppsE*	3.6
3.4	416/1458	MT0783	Rv0758	*phoR*	*ppsE*	1.8
4.1	378/1458	MT0783	Rv0758	*phoR*	*ppsE*	1.7
4.2	378/1458	MT0783	Rv0758	*phoR*	*ppsE*	3.5
4.3	416/1458	MT0783	Rv0758	*phoR*	*ppsE*	2.1
4.4	416/1458	MT0783	Rv0758	*phoR*	*ppsE*	2.2

Supplementary Figure 1

Supplementary Figure 3

Supplementary Figure 5

Supplementary Figure 9