FuChi: A cell cycle biosensor for investigating cell-cycle kinetics during avian development.

Zoe R. Sudderick[1,2], Tiernan Briggs[3], Shirooza Mubarak[4], Melinda Van Kerckvoorde[3], Ana R. Hernandez-Rodriguez[5], Sudeepta K. Panda[1,2,7], Jon Riddell;[1,2], Cameron Batho-Samblas[1,2], Lorna Taylor[1,2], Lynn McTeir[1,2], Dominique Meunier[1,2], Amy Findlay[1,2], Flossie Roberts[1,2], Anna Raper[1], Asako Sakaue-Sawano[6], Atsushi Miyawaki[6], Joe Rainger[1,2], Jeffrey J. Schoenebeck[1,2], Cornelis J. Weijer[4], Mike J. McGrew[1,2], Denis J. Headon[1,2]*, Megan G. Davey[1,2]*, Richard L. Mort[3]*, James D. Glover[1,2]*.

*Corresponding authors

1. The Roslin Institute & R(D)SVS, University of Edinburgh, Edinburgh, UK
2. Roslin Institute Chicken Embryology (RICE) group
3. Division of Biomedical and Life Sciences, Faculty of Health and Medicine, Lancaster University, Lancaster, UK.
4. School of Life Sciences, University of Dundee, Dundee, UK
5. Gurdon Institute, University of Cambridge, Cambridge, UK
6. Laboratory for Cell Function Dynamics, RIKEN Center for Brain Science, Wako, Saitama 351-0198, Japan.
7. Division of Physiology and Climatology, ICAR-Indian Veterinary Research Institute, Izatnagar, 243 122, U.P., India.

Abstract

The ability to monitor the proliferative status in live cells both *in vitro* and *in vivo* over time has revolutionised our understanding of development, growth and disease. This was first made possible by fluorescent ubiquitination-based cell cycle indicator (Fucci) technology, which distinguishes specific cell cycle phases through the reciprocal degradation of fluorescently tagged, truncated forms of human CDT1 and GMNN proteins. Fucci genetic systems have been successfully implemented in transgenic mice, zebrafish, and axolotls. To date, no viable, stably expressing Fucci line has been developed in an avian species. Although a range of continuously improving Fucci constructs have been developed in recent years, existing *in vivo* Fucci models remain limited because they rely on older reporter technology that fails to distinguish cells in S, G2, and M phases or to label cells in early G1. As a result, these models can be challenging to interpret and their utility for continuous cell tracking and precise analysis of cell-cycle dynamics is limited. Here, we introduce FuChi, a multicistronic Fucci-expressing chicken line incorporating a newly optimised reporter construct comprising of an mCerulean-tagged Histone H1.0 linker protein fused via a self-cleaving 2A peptide to the tandem Fucci(CA) cell cycle biosensor, with additional epitope tags included for detection in fixed tissues. We show that this system accurately discriminates and tracks cells in G1, S, G2, and M phases both *in vitro* and *in vivo*, enabling faithful visualisation of cell cycle status in intact tissues and organs. Using FuChi embryos, we mapped proliferation dynamics across developing tissues, analysed cell cycle states of migrating cells, and performed live imaging of early embryos. These latter experiments revealed that transition from S phase may be a key morphogenetic event during gastrulation as mesendoderm cells egress from the primitive streak to form embryonic structures including the prechordal plate. Pairing this advanced reporter with the intrinsic experimental advantages of the chicken embryo, positions FuChi as a premier *in vivo* system for studying cell-cycle kinetics in development, delivering clear technological improvements over current Fucci models. FuChi chickens provide a powerful new resource for studying embryonic development, organ growth, tissue homeostasis, disease processes, and infection responses.

Introduction

The ability to monitor the proliferative status of cells has transformed our view of development, growth, and disease. Cell cycle control diversifies as development proceeds[1] and is tightly coordinated within tissues, underpinning key aspects of morphogenesis and organogenesis such as kidney branching and digit formation[2]. In recent years, live imaging of cell cycle progression has become the gold standard for cell cycle analysis.

The E3 ubiquitin ligases APC[Cdh1], SCF[Skp2], and CUL4[Ddb1] regulate the cell cycle by targeting specific proteins for ubiquitin-mediated degradation. SCF[Skp2] not only acts as a substrate of APC[Cdh1] but also directly inhibits it, resulting in reciprocal oscillations in their activities and the levels of their target proteins. APC[Cdh1] is predominantly active in late mitosis and G1 phase, SCF[Skp2] functions during S and G2 phases, and CUL4[Ddb1] is predominantly active during S phase[3-6]. CDT1 and GMNN, which regulate DNA replication licensing, are substrates of SCF[Skp2]/CUL4[Ddb1] and APC[Cdh1], respectively, and therefore also exhibit oscillations in their protein levels[7,8]. These dynamics have been exploited to develop the Fucci (Fluorescent Ubiquitination-based Cell Cycle Indicator) systems[9-11].

The original Fucci(SA) probe pair included a fusion of monomeric Kusabira Orange (mKO2) with a truncated human CDT1 (amino acids 30–120), and a fusion of monomeric Azami Green (mAG) with the N-terminal amino acids (1-110) of human GMNN. The mKO2-hCDT1(30/120) probe accumulates in G1 phase, labelling cell nuclei red, and is degraded at the G1–S transition, while the mAG-hGem(1/110) probe accumulates during S, G2, and M phases, labelling cell nuclei green, and is rapidly degraded before cytokinesis[9].

An improved version of Fucci(SA) (sometimes referred to as Fucci2 or Fucci(SA)2) replaces the original fluorescent proteins with mCherry and mVenus, respectively, increasing signal separation and compatibility with other reporters[12]. The original probe set was expressed from separate constructs; incorporation of a 2A self-cleaving peptide enabled equimolar expression from a single promoter (termed Fucci2a or tandem-Fucci(SA)) and facilitated the development of Cre-inducible mouse models[11,13].

Fucci4 enables discrimination of all four cell cycle phases by combining four sensors: hCDT1(30-120) degraded by SCF[Skp2] marks G1, a PIP degron targeted by CUL4[Ddb1] labels S phase, GMNN(1–110) degraded by APC[Cdh1] marks S and G2, and an H1.0 linker histone fusion highlights chromatin during mitosis. Although powerful for *in vitro* imaging Fucci4 is not easily deployable in animal models as the sensors are expressed separately[14].

The bicistronic tandem-Fucci(CA) probe pair updates the CDT1 probe from Fucci(SA) by incorporating amino acids 1–100, removing the SCF[Skp2]-targeted CY motif and instead

including an N-terminal PIP box, which is targeted for degradation via CUL4[Ddb1], resulting in accumulation of the CDT1 probe from the S–G2 transition. This enables labelling of S-phase nuclei as green, G1-phase nuclei as red, and S and G2 phases nuclei as yellow with just two sensors[10].

Fucci(SA) has primarily been used in mouse and human cell lines[9-11,13-15] while tandem-Fucci(SA) has also been deployed in mouse models[11,14,15] and has been optimised for use in zebrafish, xenopus (zFucci)[16,17] and in axolotls[18] using species-specific homologues of Cdt1 and Geminin. However, none of these *in vivo* models label all four cell cycle phases, and detecting the fluorescent proteins in fixed tissues remains challenging.

Chickens are a valuable developmental model because their embryos develop externally, allowing real-time imaging and manipulation, and their early amniote morphology closely parallels that of humans[19,20]. Fucci systems have previously been introduced into cultured chicken fibroblasts and primordial germ cells, respectively[21,22], suggesting compatibility with avian species; however, no tractable *in vivo* germline model system has yet been developed.

Here, we describe a new Fucci-based biosensor, H1.0-Fucci(CA)2, designed to label all four cell cycle phases and incorporating epitope tags for detection in fixed tissues. Using the PiggyBac system, we integrated H1.0-Fucci(CA)2 into chicken fibroblasts and primordial germ cells, and demonstrate that it outperforms Fucci(SA)2 by faithfully discriminating between cells in G1, S, G2, and M phases. We further generated a stable germline line of H1.0-Fucci(CA)2-expressing transgenic chickens (FuChi), enabling, for the first time, *in vivo* studies of cell cycle progression during real time embryonic development in an avian model.

Materials and methods

H1.0-Fucci(CA)2 construction

To generate pCDNA5-CAG-H1.0Cerulean-FucciCA, a gene fragment encoding the *mCerulean* fluorescent protein fused to an HA epitope tag and a porcine teschovirus-1 2A peptide (P2A), flanked by 5′ MluI and 3′ HindIII restriction sites, was synthesised by GeneArt (Thermo Fisher). The *mCerulean* fragment was cloned into pZERO-1 (Thermo Fisher), and the coding sequence of the mouse *H1f0* gene (encoding the H1.0 linker histone) was PCR-amplified from mouse cDNA (Table 1) and cloned into the MluI site.

Subsequently, an *mCherry-hCDT(1/100)Cy(-)* gene fragment[23] - including a 5′ 6xHis tag and HindIII, and 3′ NsiI sites - was synthesised by GeneArt. An *mVenus-hGem(1/110)* fragment[23] was PCR-amplified to incorporate a 5′ Myc tag, NsiI, and 3′ AgeI sites (Table 1). These three components were sequentially cloned into a pCDNA5-Flp backbone modified to contain the CAG promoter in place of CMV[24].

To generate pB-H1.0Cerulean-Fucci(CA)2, a polylinker sequence (Table 1) was cloned into pB-H2BCerulean-Fucci2a[25] using MluI/PmeI, and the H1.0-Fucci(CA) construct was transferred as an MluI/AgeI fragment.

Transfection into PGCs

To generate H1.0-Fucci(CA)2 PGCs, 2 μg of pB-H1.0Cerulean-Fucci(CA)2 was co-transfected with 1 μg of piggyBac Hybase vector[26] into a previously derived line of male PGCs (FR2M, Hyline birds) using Lipofectamine 2000 (Life Technologies), and grown in FAOT medium[27]. Stably transduced cells were then selected with puromycin (Sigma Aldrich) and expanded prior to PGC injection.

Generation of FuChi embryos

H1.0-Fucci(CA)2 PGCs were micro-injected into HH15-16 surrogate iCaspase9 sterile hosts as described previously[28]. Briefly, 1 μl of 25 mM B/B compound (Takara Bio) was added to a 50 μl cell suspension (5000 PGCs/μl) before 1 μl of this suspension was injected into the dorsal aorta of host embryos within windowed eggs. Egg shells were sealed with medical Leukosilk tape (BSN Medical) and incubated until hatch. Founder F_0 FuChi males were mated with wild type Hyline females in pens to produce heterozygous F_1 FuChi eggs and offspring. Chicken experiments were conducted under UK Home Office licence PP9565661 and approved by the Roslin Institute Animal Welfare and Ethical Review Board Committee.

Eggs

The FuChi line is maintained in the National Avian Research Facility (NARF), Roslin Institute, UK. Eggs were incubated vertically in rocking or non-rocking incubators (Masalles) at 37.8°C. All post incubation day 14 embryos were treated in accordance with the ASPA 1986 schedule 1 culling methods. Fluorescent embryos identified using a Nightsea SFA-RB fluorescence adapter and/or a Zeiss Axiozoom V6 microscope.

Cell culture and transfections

DF1 cells were maintained in DMEM (Life Technologies) containing 10% FBS, 2% chicken serum (CS), 1% penicillin/streptomycin. Cells were transfected with either pB-H1.0Cerulean-Fucci(CA)2 or pB-H2BCerulean-Fucci2A[25], together with a vector expressing piggyBac Hybase using Lipofectamine LTX (Life Technologies) as per the manufacturer's instructions. The cell lines were termed H1.0-Fucci(CA)2-DF1 and H2B-Fucci(SA)2-DF1 in line with Sakaue-Sawano et al.[23] .

FuChi primary fibroblasts were derived from HH19 limb bud micromasses. HH19 embryos were dissected from the egg into DMEM and all limb buds removed. limb buds were placed in 10X Trypsin (Life Technologies) on ice for 30-40 minutes, followed by two washes with PBS and removal of the ectoderm with sharpened tungsten needles. Limb bud mesoderm was collected in 100 μl of micromass culture media (50:50 DMEM/Ham's F12, 10% FCS, 1% Antibiotic/Antimycotic, 1% L-Glutamine) and dissociated by trituration. 900 μl of cell culture medium was added and the samples were centrifuged at 300 rcf for 2 minutes. The cell pellet was resuspended in 10 μl of cell culture media per limb bud. 10 μl drops were added to individual wells of a 24 well plate and incubated for 30 minutes to 1 hour at 37°C. Wells were then flooded with 1 ml of micromass culture media

To derive FuChi PGCs, 1-2 μl of blood was extracted from HH16 F_1 FuChi embryos, transferred into FAOT medium and cells expanded as previously described[27].

Cell treatments

Prior to flow cytometry analysis, H2B-Fucci(SA)2-DF1, H1.0-Fucci(CA)2-DF1 cells or FuChi PGCs were grown in their regular medium conditions as described above. Cells were then treated for 16 hours in control (regular medium plus DMSO), base (regular culture medium minus FBS and CS for DF1, or removal of FAOT growth factors for PGCs) or Nocodazole treated (regular culture media containing 100 ng/ml Nocodazole (Sigma Aldrich) conditions. Following treatment, cells were dissociated from the flasks / PGCs removed, centrifuged, put through a 70 μm cell strainer, and stained with DAPI (Sigma Aldrich) for 30 minutes. Cells were analysed for DNA content and mCherry/mVenus fluorescence using a BD Fortessa X20.

Live cell imaging

Primary FuChi limb micromass cultures were grown on glass bottom culture plates in micromass culture medium and imaged for 72 hours. H2B-Fucci(SA)2-DF1 and H1.0-Fucci(CA)2-DF1 cultures were grown on glass bottom plates in cell culture media overnight then imaged for 48 hours. All live imaging was carried out on a Zeiss LSM880 confocal microscope, with z-stack images taken at 10-minute intervals. Cell tracking and generation of individual cell montages was carried out using ImageJ/Fiji and the Fucci Tools plugin[29].

Lightsheet imaging and analysis

FuChi eggs were obtained from the NARF and stored at 16ºC until the start of the experiments. The eggs were incubated for 1 hr to equilibrate to 37ºC, after which embryos were isolated on filter paper and mounted in a specially constructed imaging chamber and imaged on a dedicated lightsheet microscope as described previously[30]. To image the embryos at 10x magnification two slightly overlapping scans were performed. During the scanning the surface of the embryo was kept in focus using a dedicated surface tracking algorithm as previously described[30]. Each scan consisted of 2500-3000 successive slices taken at 1.84 µm intervals and imaged at 488 and 562 nm excitation. This procedure allowed us to collect whole embryo images at two wavelengths at 8-10 minute intervals.

To process the raw image data, the images were transformed into a rectangular coordinate system and the surface was detected using a surface finding algorithm. The top 30 z slices (14 µm) were projected onto a plane and the nuclei detected in each channel using a custom Matlab Laplacian of Gaussian (LoG) blob detector algorithm. Nuclei were classified into three classes, those detected exclusively in either the green or red channel and those detected in both, as determined by their co-localisation. For analysis of variation of hGMNN-mVenus (488 channel) and hCDT1-mCherry (562 channel) expression during the cell cycle *in vivo* selected nuclei were traced manually and their intensities in both channels measured. Individual traces were aligned using the sudden break down of hGMNN-mVenus and hCDT1-mCherry at mitosis. The boundary between the embryonic and extra embryonic area at the start of the experiment was derived from the Lagrangian repeller[31,32] and subsequently evolved for all successive timepoints using calculated tissue velocity[30].

Whole genome sequencing

Sequencing was carried out as in Oh et al.[33]. Briefly, E9 livers were snap frozen on dry ice and stored at -80°C. Genomic DNA was extracted from the frozen samples using an Monarch HMW DNA Extraction Kit (New England Biolabs) per manufacturer's instructions. Diluted gDNA was assessed using an Agilent TapeStation and quantified by a Qubit fluorometer.

gDNA from 5 samples was pooled. Library preparation and sequencing was carried out by Edinburgh Genomics. The SMRTbell library was sequenced on a Pacific Biosystems Revio platform (25M ZMW, HiFi mode) using a single SMRT cell. 58 gigabases were produced with an average fragment size of 19 kilobase pairs and Phred quality of Q30. Data was returned in bam and fastq file formats (ENA accession: PRJEB98847). To detect the genomic location(s) of transgene insertion, hi-fi reads in fastq.gz format were first aligned against the corresponding transgene sequence (using Minimap2 (v2.24-r1122)[34]. In total, 69 of 2,983,364 hi-fi reads were mapped. Transgene-mapped reads were retained by Samtools (v1.18 using htslib v1.18)[35]. The output file was converted from bam to fastq and the transgene-mapped reads were realigned using Minimap2 against the paternal haplotype of GRCg7w (white leghorn layer, GCF_016700215.2)[36].

Protein sequence comparison

Cdt1 and Gmnn protein sequences from various species were downloaded from NCBI and aligned in MegAlign Pro 17 (DNAstar) using the MAFFT algorithm.

PCR

Insertion sites in chromosomes 1 and 2 revealed by long read sequencing were confirmed by PCR (See Table 1 for primers) using Phusion polymerase (NEB), and the supplied HF or GC buffers. Chromosome 1 detection was carried out with HF buffer with the following conditions: 98°C/30s, (98°C/10 s, 60°C/30 s, 72°C/30 s) x33, 72°C/5 mins. Chromosome 2 detections were carried out with GC rich buffer with the following conditions: 98°C/30s, (98°C/10 s, 64°C/30 s, 72°C/30 s) x30, 72°C/5 mins. PCR samples were run on 2% agarose gels for assessment.

Cryo-embedding and paraffin processing

For cryo-processing, FuChi embryos were dissected from the egg, briefly washed in PBS, and fixed with 4% PFA or 10% NBF. Following fixation, embryos were washed in PBS then equilibrated in 15% sucrose / PBS at room temperature, then incubated in 15% sucrose / 7.5% gelatin / PBS at 37°C until fully equilibrated. Blocks of samples embedded in sucrose/gelatin blocks were flash frozen in isopentane on dry ice, and stored at -80°C. 10 μm cryosections were cut on a Leica CM1950 Cryostat. For imaging, cryosections were incubated in PBS at 37ºC for 15 minutes to remove gelatin, before mounting in Prolong Gold (Life Technologies) and imaged on a Zeiss LSM880 confocal microscope.

For paraffin embedding, fixed samples (10% NBF or 4% PFA) were washed in PBS, dehydrated through 70%, 90%, and 100% EtOH before clearing in 1:1 EtOH: Chloroform, and

a final wash in 100% Chloroform. Samples were incubated overnight in paraffin wax at 65°C prior to embedding. 6 μm sections for immunofluorescent analysis were cut using a Thermo Microtome.

Immunofluorescence and RNA in situ hybridisation chain reaction

Immunofluorescent detection of the epitope probes in the H1.0-Fucci(CA)2 biosensor was carried out on paraffin and cryo- sections of FuChi embryos using anti-6x-His Tag (Mouse (1:100), Thermo Fisher #MA1-21315) and anti-Myc-Tag (Rabbit (1:100), Cell Signaling Technologies #71D10) antibodies. For paraffin sections, after dewaxing antigen retrieval was performed with Tris-EDTA using an Antigen Retriever 2100 (Aptum Biologics). For cryosections, samples were equilibrated to room temperature and then incubated for 15 minutes in PBS at 37°C to remove gelatin. After these steps, paraffin and cryo- sections were treated similarly. Sections were permeabilised with PBS containing 0.5% Triton-X100, briefly washed in PBS containing 0.1% Tween 20 (PBST), and blocked in 5% goat serum / PBST for one hour at room temperature. Sections were then incubated overnight at 4°C in primary antibodies (anti-6x His Tag, anti-Myc Tag, anti-YAP1 (Rabbit (1:200) Proteintech #13584) diluted in blocking buffer. Samples were washed in PBST before incubation in fluorescent secondary antibodies (Goat anti-rabbit IgG Alexafluor 647 (1:500, Life Technologies # A-21245), Goat anti-mouse IgG Alexafluor 488 (1:500, Life Technologies, #A-11001) diluted in blocking buffer for one hour at room temperature. Samples were washed in PBST, counterstained with DAPI, mounted in Prolong Gold and imaged on a Zeiss LSM880 confocal microscope.

For wholemount immunofluorescence, FuChi embryos were dissected and fixed in 10% NBF. Samples were washed 3 x 30-60 mins in PBS containing 0.5% Triton X-100 (PBTx) with rocking, before blocking in 10% goat serum / PBTX for 2 hrs at room temperature with rocking. Samples were then incubated with anti-DAZL antibody (Rabbit (1:250), Abcam, #AB215718) diluted in blocking buffer for 20 hrs at 4°C with rocking. Following primary antibody incubation, samples were washed 5 x 30-60 mins in PBTx, before incubation with secondary antibody (Goat anti-rabbit IgG Alexafluor 647;1:200) for 20 hours at 4°C with rocking. Samples were then washed 5 x 30-60 mins in PBTx, counterstained with DAPI (1:2000 in PBTX) for 1 hour, mounted in Prolong Gold and imaged on a Zeiss LSM880 confocal microscope.

Hybridisation chain reaction (HCR; Molecular Instruments) was used to detect *SHH* expression. Briefly, FuChi embryos were dissected, fixed in 4% PFA, dehydrated into methanol and stored at -20°C overnight. HCR was then performed on these samples following the manufacturer's standard protocol using a *SHH* probe designed and ordered directly from

Molecular Instruments. After HCR, embryos were processed for cryosectioning and sections imaged using a Zeiss LSM880 confocal microscope.

Early embryo culture and imaging

Early embryo cultures were performed as in Chapman et al.[37] Briefly, embryos were collected on filter paper, washed in Ringer's solution to remove excess yolk, before culturing ventral side up on glass bottomed six well plate containing a thin layer of albumin / agar (0.3%) mix. Embryos were incubated in a 37°C imaging chamber, with wet tissue included in the surrounding space and/or unused wells on the plate to ensure humidity, and imaged on a Zeiss LSM880 confocal microscope.

Fluorescence quality assessment

E3 embryos were imaged *in ovo*, immediately upon removal, and at 4, 24, 48, 72, and 168 hrs. All images were taken with the same fluorescence settings on a Zeiss Axiozoom V16 microscope. Samples were fixed in either 4% PFA or10% NBF for 2, 4, or 24 hours, then stored in PBS or 70% EtOH.

Image based determination of cell cycle status in PGCs

Following flow cytometry of FuChi derived PGC lines under different conditions, a sample of PGCs from each line and treatment group were mounted on slides and imaged using a Zeiss LSM880 confocal microscope. Using the Fucci Tools plugin for ImageJ/Fiji[29], values of fluorescent intensity were obtained from cultured PGCs (3 independently derived lines; 3 culture conditions), *in vivo* PGCs in the germinal crescent (4 embryos), and from neighbouring somatic cells in embryo wholemounts.

Values for all cells were plotted as scatter graphs, using normalised values for mCherry and mVenus which were calculated by subtracting the population background value (mean of lowest 5 values) from the individual value and dividing this by the maximum population value mean of top 5 values). Using the percentages of PGCs in each cell cycle phase as indicated by flow cytometry analysis we determined boundaries of G1 (mCherry >0.1, mVenus <0.1), S (mCherry <0.1, mVenus >0.1), and G2/M (mCherry >0.1, mVenus >0.1). These boundaries were then applied to the values obtained from HH6 germinal crescent PGCs and surrounding somatic cells.

Table 1: Oligonucleotide sequences used for cloning and PCR.

Name	Oligonucleotide sequence 5' - 3'
Oligonucleotide sequences used for cloning*:	

H1.0_For	CGGCAC**ACGCGT**ACCGAGAACTCCACCTCCGCCCCGGC
H1.0_Rev	GCCCTG**GCGCGC**CTTCTTCTTGCTGGCCCTCTTGGCACTCGACTTGGC
mVenus_For	CGGCAC**ATGCAT**GAACAAAAACTCATCTCAGAAGAGGATCTGATGGTGAGCAAGGGCGAGGAGCTGTTCACCGG
hGem_Rev	GACGTC**ACCGGT**TTACAGCGCCTTTCTCCGTTTTTCTGCCACTTCCTTCCAATACTGAGAGG
Polylinker Fw	**ACGCGT**TCGCGA**ACCGGT**TTCGAAGTTTAAAC
Polylinker Rv	GTTTAAACTTCGAACCGGTTCGCGAACGCGT
Oligonucleotide sequences used for genotyping:	
Fucci internal Fw	GCAATTGGCCGCCACCATGAC
Fucci internal Rv	CTTGCTCACAGCGTAATCTGGAACATCGTATGG
Chromosome 1 Fw	CATAGTTAAATCAAACTCCCACAGG
Chromosome 1 Rv	TGTGCCAAAGTTGTTTCTGACTG
Chromosome 2 Fw	CAGACCGATAAAACACATGCGT
Chromosome 2 Rv	GAGCCACCCAGTTCTGAACA

*   Restriction endonuclease sites used for cloning are indicated in bold.

Results

Design of the H1.0-Fucci(CA)2 biosensor

To generate a Fucci-expressing chicken line that overcomes the limitations of previously described Fucci systems, we first aimed to design an optimal Fucci biosensor for chicken. Fucci(SA)2, the most commonly used Fucci pair, incorporates a truncated form of human CDT1(30–120) that undergoes degradation via the SCF[Skp2] pathway through its CY motif. While this system functions in mammalian models[11,38-40], the CY motif is not well conserved in non-mammalian species (Figure 1A) and does not function in zebrafish[41], suggesting that it may not function efficiently in chicken. By contrast, Fucci(CA)2 uses CDT1(1–110) with an alternative N-terminal PIP box, which is targeted for degradation via the CUL4[Ddb1] pathway and is well conserved across species (Figure 1A). The regulation of GMNN (Geminin), the counterpart in both Fucci pairs, is also highly conserved[41] (Figure 1A). This conservation, combined with the additional advantage of S-phase discrimination offered by the PIP box–based CDT1 sensor, suggested that the Fucci(CA)2 pair would be the best choice for avian species.

Neither Fucci(CA)2 nor Fucci(SA)2 discriminate between G2 and M phases and the existing animal models (which are based on Fucci(SA)2) are unable to track cells through early G1[10]. A tricistronic construct, H2B-Fucci(SA)2, has been described which enabled M-phase discrimination and tracking through early G1 by labelling DNA with an H2B-Cerulean fusion protein[25]. Similarly, the Fucci4 system, although composed of individual constructs and so not useful here, used a fusion with the H1.0 linker histone to achieve the same effect[14]. Additionally, toxicity associated with H2B fusions has also been reported *in vivo*[42]. Therefore, we selected the H1.0 linker histone fused to mCerulean (optimally spectrally separated from mVenus and mCherry) for use in our chick model. As a final optimisation step, we added HA, 6xHis and Myc tags to the H1.0-Cerulean, mCherry-CDT1(1/100)Cy(-) and mVenus-hGMNN(1/110) components, respectively, to aid in antibody detection in fixed and cleared tissues. Figure S1A shows our optimised tricistronic H1.0-Fucci(CA)2 biosensor.

Faithful discrimination of G1, S, G2 and M phases in chicken cells

We compared the utility of our new H1.0-Fucci(CA)2 biosensor against the previously described H2BFucci(SA)2 biosensor[25] by using the PiggyBac system to stably integrate each of the biosensors into immortalised chicken DF1 fibroblasts. We found that H1.0-Fucci(CA)2 accurately discriminated cell cycle stage, with similar degradation profiles to Fucci(CA)2 mammalian models[23]. This achieved a clear discrimination between G1, S and G2/M populations and a faithful response to cell cycle perturbations (Figures 1D-F, S1B,

Supplementary video 1). However, with H2BFucci(SA)2 we identified an impaired CDT1 degradation curve, resulting in persistence of mCherry beyond the G1/S transition (Figures 1G-I, S1B, Supplementary video 2), making cell cycle phase difficult to determine accurately. Both the H2B-Cerulean and H1.0-Cerulean fusions resulted in bright nuclear labelling throughout the cell cycle, including early G1, with sharp peaks in fluorescence intensity as the chromosomes condense in M-phase (Figures 1D, E, G & H). These results establish H1.0-Fucci(CA)2 as a superior cell cycle probe capable of functionally discriminating all cell cycle phases in chicken cells.

Figure 1. Faithful labelling of G1, S, G2 and M using H1.0-Fucci(CA)2 in chicken cells. A) Comparison of the PIP box (targeted by CUL4) and Cy motif (targeted by SCF) degradation domains of CDT1 and the destruction box (targeted by APC) of GMNN across vertebrate

species. While the destruction box of GMNN is well conserved in mouse, human and chicken, as are the core residues of the PIP box (Ψ represents any moderately hydrophobic amino acid (L, V, I, or M); Ω is an aromatic residue (Y or F)), the Cy motif of CDT1 is not well conserved across species. **B)** Comparison of the mCherry-hCDT1 fusion proteins used in the Fucci(SA)2 and Fucci(CA)2 systems. The mVenus-hGMNN construct is used in both systems. **C)** Schematic showing the expected fluorescence profiles of the two systems throughout the cell cycle. The Fucci(SA)2 and Fucci(CA)2 constructs contain nuclear H2B-mCerulean and H1.0-mCerulean sequences, respectively, which are present throughout the cell cycle. Note that double labeled cells are white, negative cells are grey. **D & G)** Example montages following an individual dividing cell and one of its daughters through two cell cycles (denoted by *) from live cell imaging of stably transduced chicken fibroblasts (DF1) expressing Fucci biosensors. **E & H)** Normalised fluorescence intensity (upper plots) throughout the cell cycle based on 20 cell division curves interpolated so that max time represents a full cell cycle. In Fucci(SA) cells, the degradation of CDT1 is delayed compared with the expected profile (arrows). Temporal normalised fluorescence intensity profiles (lower plots) of Fucci reporter systems based on live cell imaging data. N = 20 cells. **F & I)** Flow cytometric analysis of DF1 Fucci reporter cells in both control conditions and following treatment with nocodazole (NcZ) which leads to G2 arrest. Fucci(CA)2 shows more faithful correlation between Fucci biosensor fluorescence and DNA content as a proxy for cell cycle stage in comparison to Fucci(SA)2.

H1.0-Fucci(CA)2 expressing "FuChi" chickens

We used the piggyBac system[43] to generate stably integrated transgenic H1.0-Fucci(CA)2 cell-cycle biosensor expressing (FuChi – "**Fu**cci **Chi**cken") PGCs (Figure 2A). After transplantation of FuChi PGCs into sterile recipient embryos[28], founder F_0 males were bred with wild-type Hyline hens to generate heterozygote F_1 embryos for analysis (Figures 2A & 2B). FuChi chickens demonstrated fertility, viability and transgene transmission rates of 97%, 96% and 48%, respectively (Figure 2D). FuChi embryos demonstrated strong H1.0-Fucci(CA)2 transgene expression with no observed lethality across all developmental stages (Figures 2B & S1C), and could be successfully hatched (Figure S1D). Of note, hatched FuChi chicks are initially smaller than wild type counterparts but grow at comparative rates and are fertile. Mapping of the integration sites of the H1.0-Fucci(CA)2 transgene in FuChi embryos revealed that the majority of founders had undergone a single insertion event on chromosome 1 (NC_052573.1):139895885-139918586) or chromosome 2 (NC_052574.1):57700736-57701209) (Figure 2E). FuChi reporter expression was consistent between embryos regardless of insertion site.

To further investigate the operation of the H1.0-Fucci(CA)2 construct in our FuChi chickens we derived PGCs and limb micromass cultures from FuChi F_1 embryos for cytometry analysis and confocal imaging, respectively. The temporal fluorescent profiles of the H1.0-Fucci(CA)2 biosensor in micromass cells closely matched those of our H1.0-Fucci(CA)2-DF1 cells with clear G1, S, G2/M populations identifiable through Fucci(CA)2 biosensor abundance with labelling of nuclei by H1.0-Cerulean throughout the cell cycle and a strong H1.0-Cerulean peak evident during M-phase (Figures 2F & 1D; Supplementary video 3). Flow cytometry analysis of three independent PGC lines (two male, one female) confirmed the fidelity of the reporter in both control and experimental culture conditions in which cell cycle progression was perturbed (Figures 2G & S2E). Taken together, these results establish FuChi chickens as the first viable stably expressing avian cell cycle biosensor model for *in vivo* studies.

Figure 2. Generation and validation of FuChi reporter chicken. A) Schematic of FuChi generation strategy through primordial germ cell (PGC) injection into CIA (chemically induced ablation) sterile host embryos. **B)** H1.0-Fucci(CA)2 expression in FuChi F_1 embryos compared to wild type. **C)** Transgene integration sites in the FuChi chicken (pool of five embryos). **D)** Fertility, viability and carrier rate of FuChi embryos. **E)** Percentage of chromosome integration sites identified by sequencing present in FuChi embryos. 87 embryos were analysed at two

developmental stages (HH19; n = 32; HH35; n = 55). **F)** Individual cell montages from limb micromass cultures derived from FuChi embryos, together with subsequent temporal analysis of H1.0-Fucci(CA)2 biosensor abundance. **G)** Validation of H1.0-Fucci(CA)2 reporter system by flow analysis in PGCs derived from FuChi embryos in both control, growth factor removed, and nocodazole (Nz) treated conditions. The reporter profiles accurately reflect DNA content as determined by DAPI staining in both control and experimental conditions. Scale bars as annotated.

Monitoring cell cycle progression throughout chicken embryonic development

Having validated the functionality of H1.0-Fucci(CA)2 in cell lines derived from FuChi embryos we surveyed reporter activity during embryonic development (Figure 3). In Eyal Giyaldi & Kochav (EG & K)[44] stage X embryos, a large proportion of blastoderm cells were observed in G2/M phase, consistent with them entering diapause during lower storage temperature prior to incubation (Figure 3A)[45]. As gastrulation progresses, the primitive streak and presomitic mesoderm displayed a large proportion of cells in S, G2/M phase, whereas the developing mesendoderm and neural plate regions contain cells predominantly in G1 (Figure 3A).

Figure 3. H1.0-Fucci(CA)2 reporter expression in embryonic tissues. A) H1.0-Fucci(CA)2 biosensor expression in FuChi embryos during early development, visualised using either confocal or fluorescent stereo zoom microscopy. Known areas of high proliferation such as the blastoderm are predominantly double labelled with mCherry-hCDT1/mVenus-hGMNN (S, G2/M) while areas including the presumptive neural plate in HH4 embryos (highlighted in box) show a larger proportion of non-proliferative (G1) cells. FuChi embryos enable visualisation of circulating blood cells during vasculogenesis, the proliferative regions of the tail bud, and the

non-proliferative (G1) somites. Hn = Hensen's node. **B)** Images of embryonic tissues from FuChi embryos visualised using confocal or stereo zoom microscopy. Segregated proliferating and non-proliferating populations are present in the somites, neural tube, lens, and retinal epithelium. The stalling of cells into G1 during mesenchymal condensation is readily visible in structures such as the developing limb and feather condensates. **C)** Areas of regional proliferation are established in the developing intestine. Scale bars as annotated.

Cryosections of fixed FuChi retinal tissue, a tissue known to be highly proliferative during development[46], displayed a high proportion of cells in S, G2/M (Figure 3B). Static and live imaging of developing somites revealed that their epithelial cells undergo extensive proliferation, with a high proportion of S, and G2/M cells, compared to the mesenchyme, which is predominantly in G1 as they undergo condensation (Figure 3B & Supplementary video 4). Similarly, other sites of mesenchymal condensations such as the developing digits and feather buds show an abundance of G1-labeled cells (Figure 3B). Analysis of the developing intestine across several developmental stages revealed proliferative changes as the muscle layers developed, whilst also confirming the presence of proliferative epithelial cells in the base of the crypts at later stages consistent with a stem cell, and subsequent transit amplifying cell population, arising during late embryonic development[47,48] (Figure 3C). Taken together, these results confirm that the intensity profiles of the H1.0-Fucci(CA)2 biosensor in FuChi embryos recapitulate the known cell cycle profiles of those tissues.

Robust H1.0-Fucci(CA)2 stability in fixed tissues and antibody detection after paraffin embedding

Next, to assess the stability and usability of H1.0-Fucci(CA)2 in FuChi tissues we analysed persistence of the fluorescence signal following common methods of fixation and tissue preparation (Figure S2). H1.0-Fucci(CA)2 signal was stable following a range of fixation methods, compared to unfixed specimens. We observed only a slight decrease in the signal to noise ratio due to increased autofluorescence of the fixed samples (Figure S2).

While H1.0-Fucci(CA)2 signal can be directly observed in cryosections and fixed wholemount tissue, both specimen types can be used for immunofluorescence and *in situ* hybridisation chain reaction (HCR) to investigate overlap with gene/protein expression patterns of interest. For example, this approach can be used for identifying *SHH* expressing cells in the notochord and floor plate of HH20 embryos, or YAP1 in the developing spinal cord (Figures 4A & 4B). The inclusion of epitope tags in H1.0-Fucci(CA)2 enabled detection of antigens by immunofluorescence in formalin fixed paraffin embedded (FFPE) sections and cryosections with identical H1.0-Fucci(CA)2 profiles to fresh tissue (Figure 4C).

Figure 4. Additional downstream analysis methods for FuChi embryos include analysing cell cycle status in migrating cells *in vivo*. A) Hybridisation Chain Reaction (HCR) to detect *SHH* expression in HH20 FuChi embryos. Expression of the fluorescent reporters remains robust following HCR, with *SHH* expression localised to the notochord and floor plate of the neural tube. hGMNN-mVenus expression is notably fainter compared with hCDT1-mVenus and H1.0-mCerulean expression. **B)** Immunofluorescent detection of total YAP1 protein in a cryosection of the developing spinal cord in HH34 FuChi embryos. **C)** Examples of cryo-embedded and formalin fixed paraffin embedded (FFPE) sections of HH29 mesonephros with immunofluorescent detection for the epitope tags contained in the FuChi construct (His6-mCherry-hCDT1, Myc-mVenus-hGem). Fluorescent profiles are identical between the two section types. **D)** Expression of FuChi reporter in migrating PGCs, identified by wholemount immunofluorescence for the germ cell marker DAZL, as they move anteriorly towards the germinal crescent. **E)** Percentage of PGCs and neighbouring somatic cells in each cell cycle phase determined from confocal images (Between 55 and 172 of each cell type were analysed per embryo (n = 4 embryos). Scale bars as annotated.

To further explore the utility of FuChi embryos, we tested whether it was possible to assess the proliferation profile of native migrating cells using immunofluorescence on wholemount embryos (Figures 4D). In chickens, PGCs migrate anteriorly from the centre of the blastodisc, ahead of the forming embryo to populate the germinal crescent during early development. Here they enter the bloodstream and are translocated to the anterior vitelline veins where they exit the blood stream, migrate into the genital ridges before finally populating the gonads[49,50].

However, the proliferative status of PGCs during their initial migratory steps, prior to entering the bloodstream, is unknown. After establishing parameters (see methods for more details; Figure S3) for our imaging-based approach, we found that as PGCs migrate towards the germinal crescent the majority of cells reside in G1 (67.3%), compared to those in S (21.7%) and G2/M (11.0%) (Figure 4E). This differs from both neighbouring somatic cells (Figure 4E) and also from PGCs in culture (Figures 2G, S1D, S3), indicating that unlike the rapidly proliferating germ cells in the gonads, PGCs at these initial migratory stages are not undergoing frequent cell division.

Real time visualisation of cell cycle dynamics during gastrulation

A major attraction of Fucci systems is that they can be used to image cell cycle progression, not only in isolated live cells, but also in complex tissues and even whole embryos. Control of the cell cycle plays an important role in the embryonic development of all organisms, including chick. However, so far there is only limited information available on cell cycle changes during avian development. To show the potential of the H1.0-Fucci(CA)2 system for live imaging we used lightsheet microscopy to monitor cell cycle changes from the onset of development after egg laying (EGK XII) to the primitive streak stage (HH4) (Figure 5A, Supplementary video 5). Embryos were imaged at 10x magnification, allowing visualisation and detection of mVenus (hGMNN) and mCherry (hCDT1) in all nuclei of the surface layers of the embryo at 8-10 minute time intervals. To quantitatively assess changes in the distribution of cells in different cell cycle phases during the early stages of development we measured the total number of nuclei detected in G1, S, and G2/M based on hGMNN-mVenus and hCDT1-mVenus fluorescence as a function of time in both the embryonic and extra embryonic territories (Figure 5B). This revealed that at the onset of development most cells are in S-phase and progress into G2 followed by mitosis and G1 in a partially synchronised manner which persists in a further subset of cells during primitive streak formation (Figure 5B; same observation seen across four independent lightsheet experiments). Next, to characterise the temporal cell cycle dynamics *in vivo* we measured the kinetics of fluorescence changes in individual cells in the embryonic and extraembryonic regions at the onset of development (Figure 5C, Supplementary videos 6-8). This analysis revealed that the cell cycle is around 10 hrs in both territories (Figure 5D; representative of three independent repeats). The synchronised manner of cells progressing from S through G2 followed by mitosis to G1 from the onset of development was also apparent in the global analysis of cells in the specific regions analysed (Figure S4A).

Figure 5. Visualisation and assessment of cell cycle dynamics during early embryonic development. **A)** Epiblast surface views from a representative lighsheet experiment performed on a FuChi embryo at the start and after 11 hours of development. The boundary separating the embryonic (Em.) and extra embryonic regions (Ex.) is indicated by the white line, calculated as described in methods. **B)** Global cell counts of embryonic and extra embryonic regions of the FuChi embryo visualised by lightsheet microscopy across time in Figure 5A. All cells detected in the analysed regions were categorised based on their fluorescence profile at the time interval; hCDT1-mCherry positive (G1; magenta), hGMNN-mVenus positive (S; green), or positive in both channels (Late S, G2/M; black). **C)** Left panel: Representative image of epiblast surface view from light sheet timelapse imaging of a FuChi embryo with regions of interest in the embryonic and extra embryonic regions highlighted (yellow boxes). Right panels: Individual cells (blue rings) were tracked in the embryonic and extraembryonic regions. Images represent a single time point from a lightsheet timelapse video. **D)** Normalised fluorescence profiles of individual cells in embryonic and extra-embryonic cut out regions highlighted in (Figure 5C). Error bars represent standard deviation. **E)** 3D hypoblast view of a stage HH5 FuChi embryo at two successive time points 6 hrs apart. Four optical sections (i-iv) taken from anterior (Ant.) to posterior (Pos.) are also shown at positions indicated by white lines. Note the difference in cell cycle status of cells egressing from the streak (arrow) and prechordal plate. and anterior mesoderm cells already migrated away. Scale bars as annotated.

Finally, the epiblast view showed that during primitive streak formation many cells in the deeper layer of the embryo resided in G1, while migrating away from the streak (Figure S4B). To investigate this in more detail we performed hypoblast side up experiments (Figure 5D, Supplementary video 9), which allows better viewing of hypoblast cells and cells emerging from the streak after their ingression. Strikingly, these experiments (three independent repeats) revealed that mesendoderm cells emerging from the primitive streak are in S phase, transition through G2/M, and enter G1 during their migration away; this is particularly evident in cells forming the prechordal plate (Figure 5D, Supplementary video 9). These whole embryo live imaging results demonstrate that the H1.0-Fucci(CA)2 system allows tracking of cell cycle progression *in vivo* for prolonged periods of time. Importantly, it highlights the dynamic cell cycle changes that cells undergo during key morphological events such as gastrulation. The significance of these findings, including to what extent cell cycle progression influences cell fate, will be elaborated in more detail in future experiments on FuChi embryos.

Discussion

Here we describe the first avian cell cycle reporter model and demonstrate the functionality of the Fucci(CA)2 system *in vivo*. Fucci(CA)2 operates effectively in the chicken both *in vivo* and *in vitro*, displaying a fluorescent reporter profile consistent with that observed in mammalian systems[23]. A major advantage of this reporter is its ability to distinguish between S and G2/M phases - a key improvement over the original Fucci(SA)2 system. In addition, the inclusion of H1.0-mCerulean allows for continuous cell tracking throughout the cell cycle and enables specific identification of M-phase via labelling of mitotic chromatin. The incorporation of epitope tags further expands the system's utility by facilitating analysis in fixed tissues.

Although Fucci(SA)2 systems using human CDT1 fusion proteins have previously been used in chicken cells *in vitro* and through transplantation *in vivo*[21,22], our results suggest that in this system human CDT1 is not efficiently degraded during S phase, thereby reducing the overall fidelity of the Fucci(SA)2 reporter in chick. This likely arises from differences between the core residues of the CY motif in mammalian (Arg, Arg, Leu) and avian (Lys, Arg, Leu) CDT1, which are essential for SCF-mediated CDT1 degradation[3,51,52]. This is consistent with human CDT1 based-Fucci reporters requiring SCF degradation not being functional in zebrafish[41]. The Fucci(CA)2 system overcomes species differences in the CY motif by utilising the PIP box of CDT1, a feature conserved across metazoans[53]. As hGMNN degradation is well conserved and functional between species[41], our data suggest that the pre-existing human Fucci(CA)2

reporter system will function in other model organisms, without the need for the generation of new species-specific CDT1 variants.

The ability to analyse cell cycle status *in vivo* in both real time and in fixed sections enables many downstream applications. Here, we investigated the proliferative status of migrating PGCs during early embryonic development *in vivo*. Previous studies analysed chicken PGC cell cycle status either in culture[22,54] or from stages beyond E2.5[55,56], at which point the PGCs have entered the bloodstream. Our results reveal that at HH stage 6, as PGCs migrate anteriorly to the germinal crescent and prior to entering the bloodstream, the majority of cells are in G1 (~70%). A potential reduction in cell division before entering the vasculature is consistent with PGC number not rapidly increasing across these stages[57], as well as PGC cultures derived from EG & K stage X embryos showing reduced proliferation compared to those derived from circulating PGCs[54]. This suggests a trade-off between proliferation and active migration in PGCs at this stage, distinct from earlier passive embryo-wide movements or later circulation via the bloodstream[58]. In addition to further investigation of cell cycle events across germ cell development, FuChi embryos can be used interrogate the proliferative status of other migrating cells such as neural crest, especially given the known tractability and benefits of the chick embryo to study these cells[59,60].

The cell cycle duration observed in the embryonic and extra embryonic tissues here is faster than that measured in tissue culture and is in line with previous reports from tracking experiments[30]. The results show that most cells in diapause after egg laying are arrested in S phase of development. After re-entering development, cells complete S-phase, and enter G2, followed by mitosis resulting in the observed partial synchrony of cell divisions. The results also suggest the possibility that cells are in S phase while emerging from the primitive streak and are in G1 when they migrate away from the streak, which is particularly prominent for the cells going to form the pre chordal plate and lateral plate mesoderm. The significance of this and its link to differentiation may be elucidated by detailed cell tracking at higher magnification in future work.

Developing Fucci-expressing cell lines requires strategies that achieve stable integration of the construct at moderate expression levels with balanced stoichiometry of the fluorescent reporters. Previously, this was achieved using the Flp-In system[24,61]. In this study, we successfully applied PiggyBac transposon technology in the chick, consistent with recent results in human cells[25]. This approach achieves comparable integration without the need for prior genomic insertion of a Flp-In cassette or reliance on viral transduction. As such, the H1.0-Fucci(CA)2 PiggyBac construct described here should enable straightforward generation of Fucci reporter lines in a wide range of mammalian and avian cell types.

The chick embryo is a powerful model system that has been instrumental in advancing our understanding of vertebrate development and disease since the 17th century[62]. Over the past two decades, progress in chicken stem cell and transgenic technologies[27,28,63,64] has enabled the generation of several fluorescent reporter and gene knockout chicken lines, leading to important discoveries in embryonic development, human congenital disorders, and infectious diseases[33,65-69]. The growing repertoire of transgenic tools - now including the FuChi line - is helping to establish the chicken as a highly tractable alternative to more commonly used models, such as the mouse and zebrafish. Furthermore, the chorioallantoic membrane (CAM) of FuChi embryos provides a platform to investigate host tissue proliferation and dedifferentiation in response to human or animal xenografts (tumours or cell lines) and to evaluate potential therapies.

From a 3Rs (Replacement, Reduction and Refinement) perspective, the chick embryo offers clear ethical and logistical advantages. For example, in the UK, embryos up to embryonic day 14 are not classified as protected animals, and their use does not require the sacrifice of a pregnant female. This supports Reduction in the number of protected animals, as a single small transgenic flock can generate ~1,300 embryos annually[70]. The chick also contributes to Replacement of early-stage mammalian models and allows for Refinement through non-invasive *in ovo* imaging and precise manipulation during development.

Limitations

While the FuChi system offers comprehensive resolution of all major cell cycle phases (G1, S, G2, and M) and enables high-resolution live imaging of proliferation during early avian development, several limitations remain. First, although FuChi expression is strong and widespread in early embryos, its performance in later developmental stages has not yet been systematically evaluated. Differences in chromatin environment, epigenetic silencing, or protein turnover across tissues and timepoints could affect biosensor performance at later stages. Second, as previously reported, the Fucci(CA) probes rely on the CUL4^{DBB1}mediated degradation pathway; therefore, the intensity of the hCDT1(1/100)Cy(-) signal may be reduced in contexts where DNA damage is induced (e.g., by UV irradiation)[23]. Third, although Fucci(CA)2 distinguishes G1 from S, and S from G2/M, and the mCerulean-tagged histone H1.0 enables visualisation of mitotic chromatin, precise quantification of transitions (e.g., G2 to M) may still require complementary temporal data or additional markers, particularly in highly dynamic contexts. Finally, although the FuChi biosensor enables clear phase discrimination, analysing proliferation in rapidly morphing or migrating tissues (e.g., the primitive streak or neural crest streams) may require high-frame-rate imaging or integration

with lineage markers to avoid misinterpreting positional shifts as cell-cycle transitions. This will be enabled by application of the latest Ai-driven segmentation and tracking algorithms[71,72].

Conclusion

In conclusion, the FuChi transgenic line provides a powerful and versatile tool for real-time analysis of the cell cycle in the chicken embryo, both *in vivo* and *in vitro*, across multiple developmental stages. We show that the Fucci(CA) system functions robustly in the avian context, offering precise discrimination of all major cell cycle phases—including mitosis—while the addition of H1.0-mCerulean and epitope tags enhances both live imaging and fixed-tissue analysis, including in FFPE samples. The stability and accessibility of this reporter line make it a valuable resource for researchers worldwide. Collectively, these advances position FuChi embryos as a premier *in vivo* system for investigating developmental cell-cycle kinetics, with clear technological advantages over current mouse models. We anticipate that FuChi will advance studies in development, growth, tissue homeostasis, and disease. FuChi eggs are now available upon request from the National Avian Research Facility at the Roslin Institute, UK.

Acknowledgements

We would like to thank all of the staff belonging to the NARF and Bioimaging units of the Roslin Institute without whom this project would not have been possible. This research was funded by BBSRC (BBS/E/D/10002071; BBS/E/RL/230001C) and NC3Rs (NC/T002328/1). Edinburgh Genomics are supported with core funding from the Natural Environment Research Council (UKSBS PR18037). Figure schematics were created using BioRender.

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

During the preparation of this work the author(s) used ChatGPT in order to assist in grammar checks, and rephrasing of paragraphs for conciseness. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the published article.

Declaration of interests

The authors declare no competing interests.

References

1. Edgar, B. (1995). Diversification of Cell-Cycle Controls in Developing Embryos. Curr Opin Cell Biol *7*, 815-824. Doi 10.1016/0955-0674(95)80065-4.
2. Nogawa, H., Morita, K., and Cardoso, W.V. (1998). Bud formation precedes the appearance of differential cell proliferation during branching morphogenesis of mouse lung epithelium in vitro. Dev Dynam *213*, 228-235. Doi 10.1002/(Sici)1097-0177(199810)213:2<228::Aid-Aja8>3.0.Co;2-I.
3. Nishitani, H., Sugimoto, N., Roukos, V., Nakanishi, Y., Saijo, M., Obuse, C., Tsurimoto, T., Nakayama, K.I., Nakayama, K., Fujita, M., et al. (2006). Two E3 ubiquitin ligases, SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis. Embo Journal *25*, 1126-1136. 10.1038/sj.emboj.7601002.
4. Vodermaier, H.C. (2004). APC/C and SCF: Controlling each other and the cell cycle. Curr Biol *14*, R787-R796. 10.1016/j.cub.2004.09.020.
5. Wei, W., Ayad, N.G., Wan, Y., Zhang, G.J., Kirschner, M.W., and Kaelin, W.G. (2004). Degradation of the SCF component Skp2 in cell-cycle phase G1 by the anaphase-promoting complex. Nature *428*, 194-198. 10.1038/nature02381.
6. Benmaamar, R., and Pagano, M. (2005). Involvement of the SCF complex in the control of Cdh1 degradation in S-phase. Cell Cycle *4*, 1230-1232. DOI 10.4161/cc.4.9.2048.
7. Nishitani, H., Lygerou, Z., and Nishimoto, T. (2004). Proteolysis of DNA replication licensing factor Cdt1 in S-phase is performed independently of Geminin through its N-terminal region. J Biol Chem *279*, 30807-30816. 10.1074/jbc.M312644200.
8. Ang, X.L., and Harper, J.W. (2004). Interwoven ubiquitination oscillators and control of cell cycle transitions. Sci STKE *2004*, pe31. 10.1126/stke.2422004pe31.
9. Sakaue-Sawano, A., Kurokawa, H., Morimura, T., Hanyu, A., Hama, H., Osawa, H., Kashiwagi, S., Fukami, K., Miyata, T., Miyoshi, H., et al. (2008). Visualizing

spatiotemporal dynamics of multicellular cell-cycle progression. Cell *132*, 487-498. 10.1016/j.cell.2007.12.033.

10. Sakaue-Sawano, A., Yo, M., Komatsu, N., Hiratsuka, T., Kogure, T., Hoshida, T., Goshima, N., Matsuda, M., Miyoshi, H., and Miyawaki, A. (2017). Genetically Encoded Tools for Optical Dissection of the Mammalian Cell Cycle. Mol Cell *68*, 626-640 e625. 10.1016/j.molcel.2017.10.001.

11. Mort, R.L., Ford, M.J., Sakaue-Sawano, A., Lindstrom, N.O., Casadio, A., Douglas, A.T., Keighren, M.A., Hohenstein, P., Miyawaki, A., and Jackson, I.J. (2014). Fucci2a: a bicistronic cell cycle reporter that allows Cre mediated tissue specific expression in mice. Cell Cycle *13*, 2681-2696. 10.4161/15384101.2015.945381.

12. Sakaue-Sawano, A., Kobayashi, T., Ohtawa, K., and Miyawaki, A. (2011). Drug-induced cell cycle modulation leading to cell-cycle arrest, nuclear mis-segregation, or endoreplication. BMC Cell Biol *12*, 2. 10.1186/1471-2121-12-2.

13. Ford, M.J., Yeyati, P.L., Mali, G.R., Keighren, M.A., Waddell, S.H., Mjoseng, H.K., Douglas, A.T., Hall, E.A., Sakaue-Sawano, A., Miyawaki, A., et al. (2018). A Cell/Cilia Cycle Biosensor for Single-Cell Kinetics Reveals Persistence of Cilia after G1/S Transition Is a General Property in Cells and Mice. Dev Cell *47*, 509-523 e505. 10.1016/j.devcel.2018.10.027.

14. Bajar, B.T., Lam, A.J., Badiee, R.K., Oh, Y.H., Chu, J., Zhou, X.X., Kim, N., Kim, B.B., Chung, M., Yablonovitch, A.L., et al. (2016). Fluorescent indicators for simultaneous reporting of all four cell cycle phases. Nat Methods *13*, 993-996. 10.1038/nmeth.4045.

15. Ando, R., Sakaue-Sawano, A., Shoda, K., and Miyawaki, A. (2023). Two coral fluorescent proteins of distinct colors for sharp visualization of cell-cycle progression. Cell Struct Funct *48*, 135-144. 10.1247/csf.23028.

16. Pai, V.P., Lemire, J.M., Pare, J.F., Lin, G., Chen, Y., and Levin, M. (2015). Endogenous gradients of resting potential instructively pattern embryonic neural tissue via Notch signaling and regulation of proliferation. J Neurosci *35*, 4366-4385. 10.1523/JNEUROSCI.1877-14.2015.

17. Sugiyama, M., Sakaue-Sawano, A., Iimura, T., Fukami, K., Kitaguchi, T., Kawakami, K., Okamoto, H., Higashijima, S., and Miyawaki, A. (2009). Illuminating cell-cycle progression in the developing zebrafish embryo. Proc Natl Acad Sci U S A *106*, 20812-20817. 10.1073/pnas.0906464106.

18. Cura Costa, E., Otsuki, L., Rodrigo Albors, A., Tanaka, E.M., and Chara, O. (2021). Spatiotemporal control of cell cycle acceleration during axolotl spinal cord regeneration. Elife *10*. 10.7554/eLife.55665.

19. Davey, M.G., and Tickle, C. (2007). The chicken as a model for embryonic development. Cytogenetic and Genome Research *117*, 231-239. 10.1159/000103184.

20. Mok, G.F., Alrefaei, A.F., McColl, J., Grocott, T., and Münsterberg, A. Chicken as a Developmental Model. In Encyclopedia of Life Sciences, pp. 1-8. https://doi.org/10.1002/9780470015902.a0021543.

21. Esteves de Lima, J., Bonnin, M.A., Bourgeois, A., Parisi, A., Le Grand, F., and Duprez, D. (2014). Specific pattern of cell cycle during limb fetal myogenesis. Dev Biol *392*, 308-323. 10.1016/j.ydbio.2014.05.015.

22. Ecker, A., Lazar, B., Toth, R.I., Urban, M., Hoffmann, O.I., Fekete, Z., Barta, E., Uher, F., Matula, Z., Varkonyi, E., and Gocza, E. (2024). Creating a novel method for chicken primordial germ cell health monitoring using the fluorescent ubiquitination-based cell cycle indicator reporter system. Poult Sci *103*, 104144. 10.1016/j.psj.2024.104144.

23. Sakaue-Sawano, A., Yo, M., Komatsu, N., Hiratsuka, T., Kogure, T., Hoshida, T., Goshima, N., Matsuda, M., Miyoshi, H., and Miyawaki, A. (2017). Genetically Encoded Tools for Optical Dissection of the Mammalian Cell Cycle. Molecular Cell *68*, 626-640.e625. https://doi.org/10.1016/j.molcel.2017.10.001.

24. Ford, M.J., Yeyati, P.L., Mali, G.R., Keighren, M.A., Waddell, S.H., Mjoseng, H.K., Douglas, A.T., Hall, E.A., Sakaue-Sawano, A., Miyawaki, A., et al. (2018). A Cell/Cilia Cycle Biosensor for Single-Cell Kinetics Reveals Persistence of Cilia after G1/S

Transition Is a General Property in Cells and Mice. Developmental Cell *47*, 509-523.e505. https://doi.org/10.1016/j.devcel.2018.10.027.

25. Loftus, A.E.P., Romano, M.S., Phuong, A.N., McKinnel, B.J., Muir, M.T., Furqan, M., Dawson, J.C., Avalle, L., Douglas, A.T., Mort, R.L., et al. (2024). An ILK/STAT3 pathway controls glioblastoma stem cell plasticity. Developmental Cell *59*, 3197-3212.e3197. 10.1016/j.devcel.2024.09.003.

26. Yusa, K., Zhou, L., Li, M.A., Bradley, A., and Craig, N.L. (2011). A hyperactive piggyBac transposase for mammalian applications. Proceedings of the National Academy of Sciences *108*, 1531-1536. 10.1073/pnas.1008322108.

27. Whyte, J., Glover, J.D., Woodcock, M., Brzeszczynska, J., Taylor, L., Sherman, A., Kaiser, P., and McGrew, M.J. (2015). FGF, Insulin, and SMAD Signaling Cooperate for Avian Primordial Germ Cell Self-Renewal. Stem Cell Reports *5*, 1171-1182. 10.1016/j.stemcr.2015.10.008.

28. Ballantyne, M., Woodcock, M., Doddamani, D., Hu, T., Taylor, L., Hawken, R.J., and McGrew, M.J. (2021). Direct allele introgression into pure chicken breeds using Sire Dam Surrogate (SDS) mating. Nature Communications *12*, 659. 10.1038/s41467-020-20812-x.

29. Van Kerckvoorde, M., Ford, M.J., Yeyati, P.L., Mill, P., and Mort, R.L. (2021). Live Imaging and Analysis of Cilia and Cell Cycle Dynamics with the Arl13bCerulean-Fucci2a Biosensor and Fucci Tools. Methods Mol Biol *2329*, 291-309. 10.1007/978-1-0716-1538-6_21.

30. Rozbicki, E., Chuai, M., Karjalainen, A.I., Song, F., Sang, H.M., Martin, R., Knölker, H.J., MacDonald, M.P., and Weijer, C.J. (2015). Myosin-II-mediated cell shape changes and cell intercalation contribute to primitive streak formation. Nat Cell Biol *17*, 397-408. 10.1038/ncb3138.

31. Chuai, M., Serrano Nájera, G., Serra, M., Mahadevan, L., and Weijer, C.J. (2023). Reconstruction of distinct vertebrate gastrulation modes via modulation of key cell behaviors in the chick embryo. Science Advances *9*, eabn5429. doi:10.1126/sciadv.abn5429.

32. Serra, M., Streichan, S., Chuai, M., Weijer, C.J., and Mahadevan, L. (2020). Dynamic morphoskeletons in development. Proc Natl Acad Sci U S A *117*, 11444-11449. 10.1073/pnas.1908803117.

33. Oh, J.D.H., Freem, L., Saunders, D.D.Z., McTeir, L., Gilhooley, H., Jackson, M., Glover, J.D., Smith, J., Schoenebeck, J.J., Lettice, L.A., et al. (2024). Insights into digit evolution from a fate map study of the forearm using Chameleon, a new transgenic chicken line. Development *151*. 10.1242/dev.202340.

34. Li, H. (2021). New strategies to improve minimap2 alignment accuracy. Bioinformatics *37*, 4572-4574. 10.1093/bioinformatics/btab705.

35. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079. 10.1093/bioinformatics/btp352.

36. Smith, J., Alfieri, J.M., Anthony, N., Arensburger, P., Athrey, G.N., Balacco, J., Balic, A., Bardou, P., Barela, P., Bigot, Y., et al. (2022). Fourth Report on Chicken Genes and Chromosomes 2022. Cytogenet Genome Res *162*, 405-528. 10.1159/000529376.

37. Chapman, S.C., Collignon, J., Schoenwolf, G.C., and Lumsden, A. (2001). Improved method for chick whole-embryo culture using a filter paper carrier. Dev Dyn *220*, 284-289. 10.1002/1097-0177(20010301)220:3<284::Aid-dvdy1102>3.0.Co;2-5.

38. Sakaue-Sawano, A., Kurokawa, H., Morimura, T., Hanyu, A., Hama, H., Osawa, H., Kashiwagi, S., Fukami, K., Miyata, T., Miyoshi, H., et al. (2008). Visualizing Spatiotemporal Dynamics of Multicellular Cell-Cycle Progression. Cell *132*, 487-498. https://doi.org/10.1016/j.cell.2007.12.033.

39. Carlier, G., Maugein, A., Cordier, C., Pechberty, S., Garfa-Traoré, M., Martin, P., Scharfmann, R., and Albagli, O. (2014). Human fucci pancreatic Beta cell lines: new

tools to study Beta cell cycle and terminal differentiation. PLoS One *9*, e108202. 10.1371/journal.pone.0108202.

40. Pauklin, S., and Vallier, L. (2013). The cell-cycle state of stem cells determines cell fate propensity. Cell *155*, 135-147. 10.1016/j.cell.2013.08.031.

41. Sugiyama, M., Sakaue-Sawano, A., Iimura, T., Fukami, K., Kitaguchi, T., Kawakami, K., Okamoto, H., Higashijima, S.-i., and Miyawaki, A. (2009). Illuminating cell-cycle progression in the developing zebrafish embryo. Proceedings of the National Academy of Sciences *106*, 20812-20817. 10.1073/pnas.0906464106.

42. Verma, S., Moreno, I.Y., Gesteira, T.F., and Coulson-Thomas, V.J. (2024). Toxicity of nuclear-localized GFP in reporter mice. Sci Rep *14*, 24642. 10.1038/s41598-024-75741-2.

43. Zhao, S., Jiang, E., Chen, S., Gu, Y., Shangguan, A.J., Lv, T., Luo, L., and Yu, Z. (2016). PiggyBac transposon vectors: the tools of the human gene encoding. Transl Lung Cancer Res *5*, 120-125. 10.3978/j.issn.2218-6751.2016.01.05.

44. Eyal-Giladi, H., and Kochav, S. (1976). From cleavage to primitive streak formation: a complementary normal table and a new look at the first stages of the development of the chick. I. General morphology. Dev Biol *49*, 321-337.

45. Pokhrel, N., Genin, O., Sela-Donenfeld, D., and Cinnamon, Y. (2022). HREM, RNAseq and Cell Cycle Analyses Reveal the Role of the G2/M-Regulatory Protein, WEE1, on the Survivability of Chicken Embryos during Diapause. Biomedicines *10*. 10.3390/biomedicines10040779.

46. Rapaport, D.H., Wong, L.L., Wood, E.D., Yasumura, D., and LaVail, M.M. (2004). Timing and topography of cell genesis in the rat retina. J Comp Neurol *474*, 304-324. 10.1002/cne.20134.

47. Cloft, S.E., Uni, Z., and Wong, E.A. (2023). Profiling intestinal stem and proliferative cells in the small intestine of broiler chickens via in situ hybridization during the peri-hatch period. Poultry Science *102*, 102495. https://doi.org/10.1016/j.psj.2023.102495.

48. Kolev, H.M., and Kaestner, K.H. (2023). Mammalian Intestinal Development and Differentiation—The State of the Art. Cellular and Molecular Gastroenterology and Hepatology *16*, 809-821. https://doi.org/10.1016/j.jcmgh.2023.07.011.

49. Morimoto, M., and Saito, D. (2025). Avian primordial germ cell migration: History, mechanisms, applications, and unanswered questions. Protein Expression and Purification *235*, 106788. https://doi.org/10.1016/j.pep.2025.106788.

50. Nakamura, Y., Yamamoto, Y., Usui, F., Mushika, T., Ono, T., Setioko, A.R., Takeda, K., Nirasawa, K., Kagami, H., and Tagami, T. (2007). Migration and proliferation of primordial germ cells in the early chicken embryo. Poult Sci *86*, 2182-2193. 10.1093/ps/86.10.2182.

51. Takeda, D.Y., Parvin, J.D., and Dutta, A. (2005). Degradation of Cdt1 during S Phase Is Skp2-independent and Is Required for Efficient Progression of Mammalian Cells through S Phase*. Journal of Biological Chemistry *280*, 23416-23423. https://doi.org/10.1074/jbc.M501208200.

52. Liu, E., Li, X., Yan, F., Zhao, Q., and Wu, X. (2004). Cyclin-dependent Kinases Phosphorylate Human Cdt1 and Induce Its Degradation*. Journal of Biological Chemistry *279*, 17283-17288. https://doi.org/10.1074/jbc.C300549200.

53. Arias, E.E., and Walter, J.C. (2007). Strength in numbers: preventing rereplication via multiple mechanisms in eukaryotic cells. Genes Dev *21*, 497-518. 10.1101/gad.1508907.

54. Chen, D., Zhi, Y., Jia, X., Ji, N., Zhang, J., Liang, J., Ran, M., Liu, X., Xu, H., and Lu, Y. (2025). Characterization of primordial germ cells from EG&K stage X chicken embryos. Poultry Science *104*, 105308. https://doi.org/10.1016/j.psj.2025.105308.

55. Yang, S.Y., Lee, H.J., Lee, H.C., Hwang, Y.S., Park, Y.H., Ono, T., and Han, J.Y. (2018). The dynamic development of germ cells during chicken embryogenesis. Poultry Science *97*, 650-657. https://doi.org/10.3382/ps/pex316.

56.   Rengaraj, D., Cha, D.G., Park, K.J., Lee, K.Y., Woo, S.J., and Han, J.Y. (2022). Finer resolution analysis of transcriptional programming during the active migration of chicken primordial germ cells. Computational and Structural Biotechnology Journal *20*, 5911-5924. https://doi.org/10.1016/j.csbj.2022.10.034.

57.   Nakamura, Y., Yamamoto, Y., Usui, F., Mushika, T., Ono, T., Setioko, A.R., Takeda, K., Nirasawa, K., Kagami, H., and Tagami, T. (2007). Migration and Proliferation of Primordial Germ Cells in the Early Chicken Embryo. Poult Sci *86*, 2182-2193.

58.   Kang, K.S., Lee, H.C., Kim, H.J., Lee, H.G., Kim, Y.M., Lee, H.J., Park, Y.H., Yang, S.Y., Rengaraj, D., Park, T.S., and Han, J.Y. (2015). Spatial and temporal action of chicken primordial germ cells during initial migration. Reproduction *149*, 179-187. 10.1530/rep-14-0433.

59.   Khudyakov, J., and Bronner-Fraser, M. (2009). Comprehensive spatiotemporal analysis of early chick neural crest network genes. Dev Dyn *238*, 716-723. 10.1002/dvdy.21881.

60.   Selleck, M.A., and Bronner-Fraser, M. (1996). The genesis of avian neural crest cells: a classic embryonic induction. Proceedings of the National Academy of Sciences *93*, 9352-9357. 10.1073/pnas.93.18.9352.

61.   Mort, R.L., Keighren, M., Hay, L., and Jackson, I.J. (2014). Ex vivo culture of mouse embryonic skin and live-imaging of melanoblast migration. J Vis Exp, doi: 10.3791/51352. 10.3791/51352.

62.   Stern, C.D. (2005). The Chick: A Great Model System Becomes Even Greater. Developmental Cell *8*, 9-17. 10.1016/j.devcel.2004.11.018.

63.   Macdonald, J., Taylor, L., Sherman, A., Kawakami, K., Takahashi, Y., Sang, H.M., and McGrew, M.J. (2012). Efficient genetic modification and germ-line transmission of primordial germ cells using piggyBac and Tol2 transposons. Proceedings of the National Academy of Sciences. 10.1073/pnas.1118715109.

64.   van de Lavoir, M.-C., Diamond, J.H., Leighton, P.A., Mather-Love, C., Heyer, B.S., Bradshaw, R., Kerchner, A., Hooi, L.T., Gessaro, T.M., Swanberg, S.E., et al. (2006). Germline transmission of genetically modified primordial germ cells. Nature *441*, 766-769. 10.1038/nature04831.

65.   Doddamani, D., Carlson, D.F., McTeir, L., Taylor, L., Nandi, S., Davey, M.G., McGrew, M.J., and Glover, J.D. (2025). PRDM14 is essential for vertebrate gastrulation and safeguards avian germ cell identity. Developmental Biology *521*, 129-137. https://doi.org/10.1016/j.ydbio.2025.02.005.

66.   Ho Ching Chan, B., Hardy, H., Requena, T., Findlay, A., Ioannidis, J., Meunier, D., Toms, M., Moosajee, M., Raper, A., McGrew, M.J., and Rainger, J. (2025). A stable NTN1 fluorescent reporter chicken reveals cell specific molecular signatures during optic fissure closure. Sci Rep *15*, 10096. 10.1038/s41598-025-94589-8.

67.   Idoko-Akoh, A., Goldhill, D.H., Sheppard, C.M., Bialy, D., Quantrill, J.L., Sukhova, K., Brown, J.C., Richardson, S., Campbell, C., Taylor, L., et al. (2023). Creating resistance to avian influenza infection through genome editing of the ANP32 gene family. Nat Commun *14*, 6136. 10.1038/s41467-023-41476-3.

68.   McGrew, M.J., Sherman, A., Lillico, S.G., Ellard, F.M., Radcliffe, P.A., Gilhooley, H.J., Mitrophanous, K.A., Cambray, N., Wilson, V., and Sang, H. (2008). Localised axial progenitor cell populations in the avian tail bud are not committed to a posterior Hox identity. Development *135*, 2289-2299. 10.1242/dev.022020.

69.   Hagihara, Y., Okuzaki, Y., Matsubayashi, K., Kaneoka, H., Suzuki, T., Iijima, S., and Nishijima, K.I. (2020). Primordial germ cell-specific expression of eGFP in transgenic chickens. Genesis *58*, e23388. 10.1002/dvg.23388.

70.   McGrew, M.J., Holmes, T., and Davey, M.G. (2025). A scientific case for revisiting the embryonic chicken model in biomedical research. Developmental Biology *522*, 220-226. https://doi.org/10.1016/j.ydbio.2025.02.013.

71. Archit, A., Freckmann, L., Nair, S., Khalid, N., Hilt, P., Rajashekar, V., Freitag, M., Teuber, C., Spitzner, M., Tapia Contreras, C., et al. (2025). Segment Anything for Microscopy. Nature Methods 22, 579-591. 10.1038/s41592-024-02580-4.

72. Bragantini, J., Theodoro, I., Zhao, X., Huijben, T.A.P.M., Hirata-Miyasaki, E., VijayKumar, S., Balasubramanian, A., Lao, T., Agrawal, R., Xiao, S., et al. (2025). Ultrack: pushing the limits of cell tracking across biological scales. Nature Methods. 10.1038/s41592-025-02778-0.

Supplementary Figures

Supplementary Figure 1. Validation of Fucci reporter systems in chicken cells *in vitro* and *in vivo*. A) Design of PiggyBac (PB) Fucci reporter constructs. The Fucci(CA)2 reporter cassette uses a CAG promoter to drive expression of H1.0-mCerulean, hCDT1(1/100)Cy(-)-mCherry, and hGMNN(1/110)-mVenus fusion proteins, each containing a 5' epitope tag (HA, 6x His, Myc), and which are upstream of puromycin resistance gene. The Fucci(SA)2 reporter cassette expresses H2B-mCerulean, hCDT1(30/120)Cy(+)-mCherry and hGMNN(1/110)-mVenus fusion proteins, upstream of puromycin. ITR = Inverted terminal repeats. The presence or absence of the PIP domain and Cy motif of human CDT1 proteins are denoted in

parentheses. **B)** Comparison of Fucci(CA)2 vs Fucci(SA)2 reporter systems in chicken DF1 fibroblasts. In the absence of serum, an increased proportion of cells are observed in G1 and S phase through DNA content analysis using flow cytometry. This shift is accurately captured in the fluorescent profiles of Fucci(CA)2 but not Fucci(SA)2 cells. For comparison, control conditions are shown in Figures 1F and 1I. **C)** Table showing fertility, viability and transgenic rate of FuChi embryos across various developmental stages. **D)** FuChi chicks can be hatched, are viable, and reporter expression can be detected. **E)** Validation of H1.0-Fucci(CA)2 reporter system by flow analysis in two independently derived PGC lines derived from FuChi embryos in both control, growth factor removed, and nocodazole (Nz) treated conditions. The reporter profiles accurately reflect DNA content as determined by DAPI staining in both control and experimental conditions. Results from a third PGC line (#2) can be seen in Figure 2G.

Supplementary Figure 2. Reporter fluorescence quality post-fixation. A) Example images of FuChi or wildtype embryos which were fixed and/or stored in each medium condition as annotated and imaged at 0 and 72 hours. **B)** Fluorescent reporter levels at 0 and 72 hrs in FuChi and wild type embryos following fixation and/or storage. Fixation in 10% neutral buffered formalin (NBF) or 4 % paraformaldehyde (PFA) stabilises H1.0-Fucci(CA)2 reporter expression while reducing increase in autofluorescence levels observed without fixation. Error bars represent S.E.M. *P<0.05, **P<0.01, ***P<0.001; Two-Way ANOVA with pairwise comparisons. Scale bars as annotated.

Supplementary Figure 3. Parameterisation for image-based analysis of H1.0-Fucci(CA)2 reporter in FuChi cells *in vitro* and *in vivo*. A) Representative confocal images of PGCs cultured under different conditions used for analysis of cell cycle stage. The same experimental cells that were analysed by flow cytometry (Figures 2G and S1D) were used. **B)** Scatter plot of relative mCherry and mVenus values from image analysis. Gates for each cell cycle are marked. Orange, green and blue dots represent cells from three independently derived FuChi PGC lines. **C)** Quantification of cell cycle state in PGCs by image based and flow cytometry methodologies. Three independently derived FuChi PGC lines were analysed, with 100 PGCs analysed for image-based quantification. Nz. = Nocodazole. Scale bars as annotated.

Supplementary Figure 4. Cell cycle analysis in developing FuChi embryos using lighsheet microscopy. A) Cell counts of embryonic and extra embryonic selected regions of the FuChi embryo visualised by lightsheet microscopy across time in Figure 5C. All cells detected in the analysed regions were categorised based on their fluorescence profile at the time interval; hCDT1-mCherry positive (G1; magenta), hGMNN-mVenus positive (S; green), or positive in both channels (Late S, G2/M; black). These results are representative of three independent repeats. **B)** 3D epiblast view of a HH stage 5 FuChi embryo at two successive time points 6 hrs apart. Four optical sections (i-iv) taken from anterior (Ant.) to posterior (Pos.) are also shown at positions indicated by white lines. Note the difference in cell cycle status of cells egressing from the streak (arrow). Scale bars as annotated.

Supplementary video legends

Supplementary video 1: Imaging of DF1 cells expressing the H1.0-Fucci(CA)2 biosensor. These cells express H1.0-mCerulean (cyan), hCDT1(1/100)Cy(-)-mCherry (magenta), and hGMNN(1/110)-mVenus (green) fusion proteins. Images were captured every 10 minutes by confocal microscopy. Scale bar = 50 μm.

Supplementary video 2: Imaging of DF1 cells expressing the H2B-Fucci(SA)2 biosensor. These cells express H2B-mCerulean (cyan), hCDT1(30/120)Cy(+)-mCherry (magenta), and hGMNN(1/110)-mVenus (green) fusion proteins. Images were captured every 10 minutes by confocal microscopy. Scale bar = 50 μm

Supplementary video 3: Visualisation of H1.0-Fucci(CA)2 biosensor expression in limb micromass cell cultures derived from a FuChi embryo. hCDT1-mCherry (magenta), hGMNN-mVenus (green), and H1.0-mCerulean (cyan) show dynamic expression through cell division. Images were captured every 10 minutes by confocal microscopy. Scale bar = 50 μm

Supplementary video 4: Visualisation of somite formation in a FuChi embryo. Left panel: hCDT1-mCherry (magenta) and hGMNN-mVenus (green) expression. Right panel: H1.0-mCerulean (cyan) reporter expression. Images were captured every 10 minutes by confocal microscopy. Scale bar = 100 μm

Supplementary video 5: Lightsheet imaging of a developing FuChi embryo. Changes in cell cycle dynamics in the epiblast during development from stage EGK XIII to HH4. Time interval between images is -8 min and the scale bar represents 500 μm.

Supplementary video 6: The dynamic position of selected 330x330 μm regions in the embryonic and extra embryonic areas selected for analysis. The regions (yellow squares) were tracked over time using the velocity field calculated using particle image velocimetry. The time interval between images is 10 mins. Scale bar = 500 μm.

Supplementary video 7: Cutout of small epiblast region from the embryonic area of a developing FuChi embryo. Coloured circles indicate detection of hGMNN-mVenus (green) or hCDT1-mCherry (magenta) fluorescence. Manually tracked cells are outlined by blue squares. Scale bar = 50 μm.

Supplementary video 8: Cutout of small epiblast region from the extra embryonic tissue of a developing FuChi developing embryo. Coloured circles indicate detection of hGMNN-mVenus (green) or hCDT1-mCherry (magenta) fluorescence. Manually tracked cells are outlined by blue squares. Scale bar = 50 μm.

Supplementary video 9: Hypoblast side view of the hypoblast and emerging mesendoderm in a FuChi embryo developing from stage HH3 to stage HH5 with a depth of view of 60 µm. Time interval between images is 10 minutes. Scale bar = 500µm

A
1. Stably transduce PGCs with PiggyBac expression vector
2. Select and expand PGCs to inject into CIA host embryos
3. Raise founder males to sexual maturity and breed with WT hens
4. Generate heterozygote FuChi (F₁) embryos

H1.0-Fucci(CA)2 reporter → PGCs

FuChi Founder (F₀) Male X Wild Type Female

B

FuChi / WT — hCDT1
FuChi / WT — hGMNN
FuChi / WT — mCerulean
FuChi / WT — Brightfield
Scale bar: 1 mm

C

Chromosome 1
139,820 K – 139,980 K
SOX1 · 139,895,885 · LOC121107347

PB 5' ITR — CAG Promoter — H1.0 — mCerulean — mCherry — hCDT1 — mVenus — hGMNN — IRES2 — Puromycin — pA — PB 3' ITR

NRSN1 (+12) · 577,007,36 · LOC121109792
Chromosome 2
57,600 K – 57,820 K

D
FuChi embryos (n=1035)
Rate (%)
Fertility 87.8 · Viability 93.2 · Transgenic 48.5

E
FuChi transgene integration (n=87)
Insertion rate (%)
Chr. 1 83.9 · Chr. 2 2.3 · Other 13.8

F
FuChi limb micromass
hCDT1
hGMNN
hCDT1 / hGMNN
mCerulean

M — H1.0-mCerulean — G1 — hCDT1-mCherry — S — Geminin-mVenus — G2

Fluorescence index vs Time (hrs)
hCDT1 · hGMNN · mCerulean
n=20

G
FuChi PGCs

Control — mCherry vs mVenus
G1 (43.4%) · G2/M (32.5%) · S (22.7%)

No growth factors
G1 (57.2%) · G2/M (20.2%) · S (21.4%)

100 ng/ml Nz.
G1 (3.63%) · G2/M (80.2%) · S (15.3%)

Cell count vs DAPI
Control: G1 45.8% · S 15% · G2/M 39.2%
No growth factors: G1 62.8% · S 13.8% · G2/M 23.4%
100 ng/ml Nz.: G1 2.16% · S 10.7% · G2/M 86.9%

A Hybridisation Chain Reaction

Neural Tube · HH20 | Neural Tube · HH20

SHH | *SHH*
hCDT1 | mCerulean
hGMNN

B Immunofluorescence

Spinal Cord · HH34 | Spinal Cord · HH34

YAP | YAP
hCDT1 | mCerulean
hGMNN

C Cryosection | FFPE section

Myc-tag | Myc-tag
His6-tag | His6-tag

DAPI | DAPI

D Wholemount immunofluorescence

Embryo · HH6 | Embryo · HH6 | HH6

DAZL | DAZL | DAZL
hCDT1 | mCerulean | hCDT1
hGMNN | | hGMNN

HH6

DAZL
mCerulean

E Image based analysis of cell cycle phase *in vivo*

- G1
- S
- G2/M

Cell cycle status (%)

PGCs · Somatic cells

A — Lightsheet live imaging

hGMNN — hCDT1 — 0 hrs — Em. — Ex. — EG & K XII — 500 μm
hGMNN — hCDT1 — 11 hrs — 500 μm

B — Global cell analysis

Embryonic

Number of nuclei — Time (hrs) — S — G1 — Late S, G2/M

Extra embryonic

Number of nuclei — Time (hrs) — S — G1 — Late S, G2/M

C — Individual cell tracking

Ex. — Em. — 0 min — 500 μm
Embryonic — 0 min — 50 μm
Extra embryonic — 0 min — 50 μm

D — Individual cell tracking

Embryonic

Fluorescence index — Time (hrs) — M — hCDT1 — hGMNN — M

Extra embryonic

Fluorescence index — Time (hrs) — M — hCDT1 — hGMNN — M

E — Hypoblast side up lightsheet live imaging

hGMNN — hCDT1 — HH5 (0 hrs) — Pos. ←→ Ant. — 500 μm — i — ii — iii — iv
hGMNN — hCDT1 — HH5 (+6 hrs) — Pos. ←→ Ant. — 500 μm — i — ii — iii — iv

i — 250 μm
ii — 250 μm
iii — 250 μm
iv — 250 μm

i — 250 μm
ii — 250 μm
iii — 250 μm
iv — 250 μm